# Why is there so much variability in crop multi-model studies?

Wallach[a], Daniel; Palosuo[b,*], Taru; Mielenz[c], Henrike; Buis[d], Samuel; Thorburn[e], Peter; Asseng[f],

Senthold; Dumont[g], Benjamin; Ferrise[h], Roberto; Gayler[i], Sebastian; Ghahramani[j,1], Afshin; Harrison[k],

Matthew Tom; Hochman[e], Zvi; Hoogenboom[l], Gerrit; Huang[m], Mingxia; Jing[n], Qi; Justes[o], Eric;

Kersebaum[p,q,r], Kurt Christian; Launay[s], Marie; Lewan[t], Elisabet; Liu[k], Ke; Luo[u], Qunying; Mequanint[i], Fasil;

Nendel[p,q,v], Claas; Padovan[h], Gloria; Olesen[x], Jørgen Eivind; Pullens[x], Johannes Wilhelmus Maria; Qian[n],

Budong; Seserman[p], Diana-Maria; Shelia[l], Vakhtang; Souissi[y], Amir; Specka[p], Xenia; Wang[m], Jing; Weber[z],

Tobias, K.D.; Weihermüller[aa]; Lutz, Seidel, Sabine J.[a]

[a]Institute of Crop Science and Resource Conservation, University of Bonn, Germany

[b]Natural Resources Institute Finland (Luke), Helsinki, Finland

[c]Julius Kühn Institute (JKI) – Federal Research Centre for Cultivated Plants, Institute for Crop and Soil

Science , Braunschweig, Germany

[d]INRAE, UMR 1114 EMMAH, Avignon, France

[e]CSIRO Agriculture and Food, Brisbane, Queensland, Australia

[f]Technical University of Munich, Department of Life Science Engineering, Digital Agriculture, HEF World

Agricultural Systems Center, Freising, Germany

[g]Plant Sciences & TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech, University of Liege,

Gembloux, Belgium

[h]Department of Agriculture, Food, Environment and Forestry (DAGRI), University of Florence, Italy

[i]Institute of Soil Science and Land Evaluation, Biogeophysics, University of Hohenheim, Stuttgart,

Germany

---

[1] Present address: Department of Environment and Science and Innovation, Queensland Government, Toowoomba, QLD 4350, Australia.

[j]Centre for Sustainable Agricultural Systems, Institute for Life Sciences and the Environment, University

of Southern Queensland, Toowoomba, Queensland, Australia

[k]Tasmanian Institute of Agriculture, University of Tasmania, Launceston, Tasmania, Australia, 7248

[l]Agricultural and Biological Engineering Department, University of Florida, Gainesville, Florida, USA &

Global Food Systems Institute, University of Florida, Gainesville, Florida, USA

[m]College of Resources and Environmental Sciences, China Agricultural University, Beijing, China

[n]Ottawa Research and Development Centre, Agriculture and Agri-Food Canada, Ottawa, Canada

[o]CIRAD, Univ Montpellier, Persyst Department, Montpellier, France

[p]Leibniz Centre for Agricultural Landscape Research (ZALF), Müncheberg, Germany

[q]Global Change Research Institute CAS, Brno, Czech Republic

[r]Institute for Tropical Plant Production and Agricultural Systems Modelling, Georg-August University,

Göttingen, Germany

[s]INRAE, US 1116 AgroClim, Avignon, France

[t]Department of Soil and Environment, Swedish University of Agricultural Sciences (SLU), Uppsala,

Sweden

[u]Hillridge Technology Pty Ltd, Sydney, Australia

[v]Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany

[x]Department of Agroecology, Aarhus University, Tjele, Denmark

[y]Swift Current Research and Development Centre, Agriculture and Agri-Food Canada, Swift Current,

Saskatchewan, Canada

[z]Soil Science Section, Faculty of Organic Agricultural Sciences, University of Kassel, Germany (before [i])

[aa]Institute of Bio- and Geosciences - IBG-3, Agrosphere, Forschungszentrum Jülich GmbH, Jülich,

Germany

*Corresponding author. Email: taru.palosuo@luke.fi

**Author contribution:**

## Abstract

It has become common to compare crop model results in multi-model simulation experiments. In general, one observes a large variability in such studies, which reduces the confidence one can have in such models. It is important to understand the causes of this variability as a first step toward reducing it. For a given data set, the variability in a multi-model study can arise from uncertainty in structure or in parameter values for a given structure. Previous studies have made assumptions about the origin of parameter uncertainty, and then quantified its contribution, generally finding that parameter uncertainty is less important than structure uncertainty. However, those studies do not take account of the full parameter variability in multi-model studies. Here we propose estimating parameter uncertainty based on open-call multi-model ensembles where the same structure is used by more than one modeling group. The variability in such a case is due to the full variability of parameters among modeling groups. Then structure and parameter contributions can be estimated sing random effects analysis of variance. Based on three multi-model studies for simulating wheat phenology, it is found that the contribution of parameter uncertainty to total uncertainty is, on average, more than twice as large as the uncertainty from structure. A second estimate, based on a comparison of two different calibration approaches for multiple models leads to a very similar result. We conclude that improvement of crop models requires as much attention to parameters as to model structure.

## Keywords

Crop model, structure uncertainty, parameter uncertainty, multi-model studies

## 1. Introduction

Process-based crop models are widely used tools in agronomy, to help analyze past results and to explore scenarios related to alternative management options, effect of climate change, impact of new varieties etc. (Boote et al., 2010). Since the inception of the Agricultural Modeling Inter-comparison and Improvement Project (AgMIP, Rosenzweig et al., 2013), it has become common to organize multi-model simulation studies, where multiple modeling groups are provided with the same input data (i.e. weather, crop management, soil characteristics and initial conditions) and the same observational data for model calibration, and simulate the same response variables. Such studies have shown that the variability in simulated values is very substantial (Asseng et al., 2013; Durand et al., 2018; Salo et al., 2016; Sándor et al., 2020; Wallach et al., 2021a, 2021b). In several studies where multiple crop models were combined with multiple global circulation models (GCMs), it was found that the uncertainty contribution from crop models is as large or larger than that from the GCMs (Asseng et al., 2013; Müller et al., 2021; Wang et al., 2024, 2020). The variability in multi-model studies is probably the most pertinent measure of overall crop model uncertainty, since it reflects the current variability in simulated results among modeling groups. The large uncertainty clearly limits the confidence that one can have in the results of crop models, since it implies that results can vary widely depending on the specific model and modeling group providing the results. A reduction in uncertainty from crop models is necessary in order to increase confidence in the results of these models (Maiorano et al., 2017; Rötter et al., 2011).

Our case study here concerns prediction of wheat phenology. Crop phenology is important since it has a major effect on crop yield and is a major aspect of crop response to global warming (Ahmad et al., 2019; Fatima et al., 2020; Liu et al., 2022). Process-based models of phenology, usually embedded in crop or agroecosystem models, are often used to predict the effect of weather on phenology (Bindi et al., 2015; Muleke et al., 2022). Multi-model studies specifically of crop phenology have shown wide

variability in simulated results (Wallach et al., 2021b, 2021a). Thus the uncertainty in prediction of crop

phenology is of interest in its own right, as well as serving as an example for crop models more generally.

Since all modeling groups are given the same input and observation data in multi-model studies,

the variability between groups has two possible origins. The first is the fact that different groups use

different model structures. That is, there is uncertainty in crop model structure, which encompasses

both the processes included in the models and the specific forms of the equations used to model those

processes. The second source of variability is uncertainty in the parameter values. Different groups using

the same model structure can have different values for the parameters (Albanito et al., 2022;

Confalonieri et al., 2016), again leading to differences in simulated values. It is important to quantify the

contributions of structure and parameter uncertainty to overall uncertainty, as a first step in reducing

the overall uncertainty. In general, uncertainty is summarized as a variance value, so the question is the

relative contributions of structure and parameter variance to the overall variance of simulated values in

multi-model studies.

To proceed, it is helpful to have a clear definition of the parameter and structure uncertainties of

interest. We propose that an appropriate definition of model structure uncertainty is the range of

plausible crop model structures, where by "plausible" we mean a structure that would be deemed

acceptable by researchers knowledgeable in the field of crop modeling. The structures represented in

open-call multi-model studies can be considered a random sample of plausible structures; random

because all modeling groups were invited to participate, plausible because the structures are used by

researchers in the field.

Our definition of parameter uncertainty for a given model structure and data set is the range of plausible

parameter vectors for that structure and data. Here "plausible" means that research groups working

with that structure would consider those parameters, or the way in which the parameters are obtained,

reasonable. Note that parameter uncertainty is nested within structure uncertainty. That is, the

uncertainty in question is different for each model structure, since each structure has its own

parameterization. Overall parameter uncertainty is an average over the uncertainties for different

structures. Parameter uncertainty covers both the plausible range of parameter default values and the

plausible range of calibration procedures, including the choice of which parameters to fit to the data and

the procedures of doing the fitting. Each parameter vector represented in open-call multi-model studies

is plausible, since it is derived by a practicing research group. For a given model structure, each

parameter vector can be considered a random draw from among plausible vectors for that structure,

that is, all plausible parameter vectors have an equal chance of being represented in the multi-model

study.

The variability in calibration approach is a major source of parameter uncertainty. Calibration of

crop models and more generally process based models in other fields is difficult because of the large

number of parameters and the multiple types of observed variables. As a result, there is no general

consensus on the best calibration approach for crop modes (Ahuja and Ma, 2011). One question is how

to choose the subset of parameters to estimate by fitting to observed data. Various studies have

proposed sensitivity analysis approaches for choosing the parameters to estimate (e.g. Zadeh et al., 2022

for water quality modeling; Zhang et al., 2014 for rice phenology). Other studies have compared

different algorithms for searching for the best parameter values (e.g. Gao et al., 2020; Harrison et al.,

2019). The choice of objective function is also variable, and several choices are possible (e.g. Houska et

al., 2015 for ecological models). Wallach et al. (2021c) specifically studied the range of calibration

approaches used by different modeling groups in three multi-model studies. They found that some

groups used a frequentist approach, with various objective functions, and others a Bayesian approach,

with various assumptions about the likelihood function. There were also differences in the way to choose

the parameters to estimate, with some groups basing the choice on sensitivity analysis, others on expert

knowledge about the most important model parameters, sometimes combined with trial and error to

choose the parameter set that gives the best fit to the data. Even for modeling groups using the same

model structure, there were multiple differences in calibration approach, including different choices of

which parameters to fit to the data. The uncertainty in plausible parameter vectors includes all these

different choices in calibration practices. Albanito et al. (2022) also emphasized that different modeling

groups make different decisions about calibration

Several previous studies have specifically quantified the contributions of structure and

parameter uncertainty to total uncertainty for crop models or other process-based models. In these

studies, a single modeling group generates multiple parameter vectors for each of several models. Zhang

et al. (2017) considered five different models used to predict rice phenology. For each model, 50

parameter sets were generated, based on an assumed uncertainty range for each parameter. Analysis of

variance was used to estimate the separate contributions of structure and parameter uncertainty. Tao et

al. (2018) used a similar approach for an ensemble of seven models for simulating barley growth and

yield. Parameter ranges were based on expert opinion. Both studies identify a fixed subset of parameters

that are considered uncertain. These approaches are appropriate for estimating *a priori* uncertainty in a

fixed subset of parameter values. However, in multi-model studies, and in our definition of plausible

parameter values, models are calibrated using provided data. Calibration modifies the uncertainty in

parameter values compared to their *a priori* range. Furthermore, these studies ignore uncertainty in

parameters outside the subset of parameters considered. Thus, this approach does not provide a sample

of plausible parameter values according to our definition, and is therefore not appropriate for estimating

the contribution of parameter uncertainty to total uncertainty in multi-model studies.

A different approach assumes that parameter uncertainty derives from error in observations.

The variance of observation error is estimated and then used to estimate the distribution of the

parameter estimators. Thus Wallach et al ( 2017) estimated parameters of two models using the

standard regression technique of generalized least squares, which also provides estimates of the

variance-covariance matrix of the parameter estimators (Seber and Wild, 1989). Then samples of

parameter vectors were generated for each model, and analysis of variance was used to estimate

structure and parameter variance. Another approach explores the parameter space around the optimal

parameter values, and accepts parameter values that satisfy a goodness of fit test, which is a way of

relating parameter uncertainty to observation uncertainty. Migliavacca et al. (2012), looking at 12

budburst models, used this approach to generate 1000 parameter vectors that give fits to the data that

are statistically equivalent to the fit using the best parameter values for each model. However, while the

effect of observation error on parameter uncertainty is important, it is not relevant to uncertainty in

multi-model studies, which use a fixed data set. Thus the variability in parameter values generated in

these studies does not mimic the variability of plausible parameter values.

In a somewhat different context, Smallman et al. (2021) studied structure and parameter

uncertainty for five terrestrial ecosystem models. Bayesian calibration was used to determine parameter

distributions for each model. This approach evaluates parameter uncertainty as the result of *a priori*

uncertainty and observation error. Since as noted above observation error does not contribute to

variability in open-call multi-model ensembles, this approach again does not recreate the variability of

plausible parameter values, according to our definition.

Xiong et al. (2020) considered structure and parameter contributions to uncertainty in wheat

model simulations applied to two Chinese regions. Three models were considered. Parameter variability

was due to different levels of spatial detail (same parameters at all locations, different parameters for

each major agro-ecological zone, parameters fit to phenology data or parameters fit to both phenology

and yield data). In this case parameter uncertainty measures the variability between different specific

choices of the way to determine the parameter values. The approach is not aimed at reproducing the

variability in multi-model studies.

These previous approaches have sampled from specific types of parameter uncertainty, but

those samples do not represent the uncertainty, or not all the uncertainty, between different modeling

groups that could participate in open-call multi-model studies (i.e. in plausible parameter vectors). It

would be essentially impossible, in fact, to generate a sample of parameter values that takes into

account all the variability that exists between modeling groups.

The purpose of this paper is to propose and illustrate a novel approach to the estimation of

structure and parameter variances in open-call multi-model studies. Our hypothesis is that the most

realistic estimate is one based directly on the results of such studies. For our case studies, we revisit

three published open-call multi-model studies on predicting wheat phenology (Wallach et al., 2021b,

2021a), where some model structures were used by more than one modeling group. Given the results

from multiple different model structures, and from multiple groups using the same structure, we

estimate structure and parameter variances using random effects analysis of variance. As far as we

know, this is the first estimation of the two contributions to overall uncertainty that takes into account

the true variability between modeling groups, and is thus pertinent for explaining the uncertainty in

multi-model studies. The methodology employed here, namely using random effects analysis of variance

on data from multi-model studies, is applicable in general to process-based models.

In the particular cases treated here, we can obtain a second estimate of the variance in

simulated values due to parameter uncertainty. In the multi-model studies revisited here, each

participating group used their "usual" calibration approach. Subsequently an original calibration protocol

was developed, and most of the groups redid the calibration using the new protocol. Thus for multiple

groups we have results for two different calibration procedures, "usual" and "protocol", both of which

may be considered plausible The comparison between them can be used to estimate potential variability

in parameters due to calibration approach. These results can be compared with the results based on the

original multi-model studies.

## 2. Materials and Methods

2.1 Observational data

The three multi-model studies use three different data sets (**Table 1**). Two of the data sets are

from winter wheat variety trials in France. These two data sets have identical structures (same sites and

sowing dates) but concern two different wheat varieties, "Apache" and "Bermude". Both varieties are of

intermediate precocity, but Bermude has a longer cycle. The observed data from each environment

consist of dates of stem elongation (BBCH30 on the BBCH scale, Meier, 1997) and of the middle of

heading (BBCH55). The third data set is from a multi-site sowing date experiment on spring wheat variety

Janz in Australia. The original data were observations of growth stage at weekly intervals. The data

provided to modeling groups were the dates of each BBCH stage, from the first to the last observed

stage, derived from interpolation of the original data. All data sets were split into two subsets, one for

calibration and the other for evaluation. The calibration and evaluation data sets had neither site nor

year in common. Thus evaluation was a true test of how well models could predict out-of-sample

conditions.

| Data set | Number of calibration environments | Number of evaluation environments | Observed development stages* | Simulated development stages used for analysis* |
|---|---|---|---|---|
| France Variety Apache | 14 | 8 | BBCH30, BBCH55 | BBCH10, BBCH30, BBCH55 |
| France Variety Bermude | 14 | 8 | BBCH30, BBCH55 | BBCH10, BBCH30, BBCH55 |
| Australia | 24 | 18 | BBCH stage at | BBCH10, BBCH30, |

| | | | | weekly intervals | BBCH65, BBCH90 |
|---|---|---|---|---|---|
| Variety Janz | | | | | |

232       *The development stages, using the BBCH scale (Meier, 1997) are emergence (BBCH10), start of

233   stem elongation (BBCH30), middle of heading (BBCH55), full flowering (BBCH65), and maturity (BBCH90).

234                                     **Table 1. Data sets**

235           **The three data sets used in the multi-model ensemble studies analyzed here.**

## 2.2 Participating modeling groups and model structures

The  call for participants in these studies was published using  the mailing lists of AgMIP and of

several crop models. All modeling groups that volunteered (a modeling group is one or more individuals

who work together to run a model) were accepted. No specific attempt was made to encourage or

discourage participation by different groups that used the same structure. In the event, there were two

or three (depending on the simulation study) model structures used by more than one group (Table 2).

These are thus what we have termed "open-call" multi-model studies.

A total of 30 modeling groups using 23 different structures participated in one or more of the

simulation experiments. For the purposes of this study, each model with a unique name was considered

as a different model structure. A list of structures represented here, together with references that

describe the model equations, is given in Supplementary Table 1. In all cases, the phenology models

were embedded in general crop models, so the model structure here refers to the overall crop model.

Most but not all of the groups participated in all the studies.

In the original reports of these studies, the different modeling groups were identified by a code

(M1, M2 etc.) without identifying  the group or the model structure used (Wallach et al., 2021b, 2021a).

The same codes are used here. In addition, we assign here codes to the different model structures (S1,

S2, etc.). The same model structure can be associated with more than one modeling group.

2.3Ssimulation studies

The three multi-model simulation studies, corresponding to the three data sets, are summarized

in Table 2. In the first two studies, based on the two French data sets, each participating modeling group

implemented their usual calibration method to calibrate their model using the calibration data subsets

and then simulated days from sowing to development stages BBCH10, BBCH30, and BBCH55 for both the

calibration and evaluation environments (Wallach et al., 2021a) (Table 2). In the study based on the

Australian data set (Wallach et al., 2021b), each participating modeling group again used their usual

calibration method and then simulated days from sowing to development stages BBCH10, BBCH30,

BBCH65, and BBCH90 for the Australian environments. Neither the French nor the Australian data sets

had observations of BBCH10 (emergence), which was included in the variables to simulate in order to

have an example of a simulated stage without calibration data. Detailed information about the variability

in calibration procedures among the different modeling teams is presented in Wallach et al. (2021c).

| Simulation study | Data set (country, cultivar) | Calibration procedure | Number of modeling groups | Repeated structures* |
|---|---|---|---|---|
| multi-model study 1 | France, Apache | Usual | 27 | S2 (4), S4 (3), S18 (2) |
| multi-model study 2 | France, Bermude | Usual | 27 | S2 (4), S4 (3), S18 (2) |
| multi-model study 3 | Australia, Janz | Usual | 28 | S2 (3), S4 (3), S18 (2) |

265     *The model structures used by more than one modeling group, and in parentheses, the number of

266     groups that used  that structure.

267                                 **Table 2**

**268**   **The three open-call multi-model simulation studies analyzed here (Wallach et al., 2021a,**

**269**   **2021b).**

**270**   Following the three multi-model studies, a standardized calibration procedure (hereafter the

**271**   calibration "protocol") was proposed (Wallach et al., 2023). In a fourth simulation experiment, each of 19

**272**   participating groups used the protocol calibration to simulate for the same three data sets used in the

**273**   first studies. Any group that participated in the fourth study but that had not participated in the first

**274**   three studies first did the calibration using their usual method, before applying the protocol. Thus for

**275**   each data set and each participant, there were two parameter vectors, based on usual or protocol

**276**   calibration.

**277**   ## 2.4 Estimation of structure and parameter variance

**278**   ### Variance components approach

**279**   We make the usual statistical assumptions for random effects analysis of variance (Scheffé,

**280**   1959). We assume that for each of the open-call simulation experiments, the model structures are a

**281**   random sample from plausible structures. We assume that each parameter vector is a random draw

**282**   from plausible parameter vectors for the corresponding model structure. Finally, we assume

**283**   independence between the parameter estimation choices of different modeling groups.

**284**   The statistical model for a simulated value is then

**285**   $$y_{sp} = \mu + \alpha_s + \beta_{sp}$$

**286**   where $y_{sp}$ is the simulated value for model structure $s$ and the parameter vector $p$, $\mu$ is the overall

**287**   mean, $\alpha_s$ is the effect of model structure and $\beta_{sp}$ is the effect of parameter vector nested within the

**288**   model structure. The parameter effect is nested within structure, because different structures have

different parameterizations. There is no residual term because these are simulated values with no

measurement error. We treat $a_s$ and $\beta_{sp}$ as random effects, since we are interested in the variability in

the population not in the specific model structures or parameter vectors in the sample. The variances are

noted $\mathrm{var}(\alpha_s) = \sigma_{structure}^2$ for all $s$ and $\mathrm{var}(\beta_{sp}) = \sigma_{parameters}^2$ for all $sp$. The random terms are assumed

independent so that the total variance for one environment and one simulated variable is the sum of

model structure variance and parameter variance (Scheffé, 1959):

$$\sigma_{total}^2 = \sigma_{structure}^2 + \sigma_{parameters}^2 \tag{2}$$

To estimate parameter and model structure variance simultaneously , we use the lmer function

of the lme4 package (Bates et al., 2015) in R (R Core Team, 2017), which is designed to analyze linear

mixed effect models (see Supplementary for more details). An illustration of the data used in this

approach is given in Fig. 1. As that figure shows, three of the model structures (S2, S4, and S18) were

used by more than one modeling group. It is the variability between the different groups using the same

model structure that allows us to estimate parameter variance. The results for all groups contribute to

the estimation of structure variance. Note that the simplified approach to estimating parameter

uncertainty using these same data in Wallach et al. (2021a, 2021b) underestimates the contribution of

parameter uncertainty.

**Figure 1**

**Example of data used for estimating parameter and structure uncertainty in the "variance**

**components" approach. Each circle or star is a simulated value of days from sowing to mid-**

**heading (stage BBCH55) for variety Apache in one environment from simulation study 1.**

**All modeling groups used the same inputs. The x-axis shows the model structure. In most**

**cases, each structure was used by a single modeling group (open circles), but structures S2,**

**S4, and S18 were used respectively by 4, 3, and 2 groups (data shown as stars). It is the**

**variability between simulations from different groups using those model structures that is**

**the basis for estimating parameter uncertainty. The horizontal line is the average of all**

**simulated values.**

Comparison of usual and protocol calibration

A second estimation of parameter uncertainty is based on comparing the results between usual

and protocol calibration, for each group. For each environment, we calculate the variance of the two

calibration approaches, noted var$_i$ for modeling group $i$ (eq. 1), and then average over modeling groups

to obtain an estimated parameter variance for that environment (eq. 2).

321
$$\mathrm{var}_i = \left( y^{usual} - \bar{y} \right)^2 + \left( y^{protocol} - \bar{y} \right)^2 \ (1)$$

322
$$\hat{\sigma}^2_{parameters} = 1/R \sum_{i=1}^{R} \mathrm{var}_i \ (2)$$

where $\bar{y} = (y^{usual} + y^{protocol})/2$ and R is the number of modeling groups.

We do not have a rigorous method of estimating structure variance from these data. The

difficulty is that the assumptions underlying the random effects model described above are not satisfied;

the parameter vectors obtained by protocol calibration are not independent between modeling groups,

since all use the same protocol. Therefore, or each environment we use the estimate of structure

variance from the variance components approach above.

330                          **Figure 2**

**331    Example of simulated values where each group used two calibration approaches.**

**332    Simulations are of days from sowing to stem elongation (stage BBCH30) for variety**

**333    Bermude in one environmen. Each code (M1, etc.) refers to a specific modeling group (x-**

**334    axis). For each modeling group, there are two simulated values, one using parameters based**

**335    on usual calibration and the other using parameters based on protocol calibration. The**

**336    differences between the two values are used to estimate parameter variance. The horizontal**

**337    line shows the mean of simulations using usual calibration.**

## 339    3. Results

Initially, all calculations were done for the calibration and evaluation data subsets separately.

However, there were no systematic differences between the two (Fig. S1 and Fig. S2). All results here are

therefore based on the combined calibration and evaluation data. Table 3 and Table 4 show parameter

variance, and parameter variance as a fraction of total variance, averaged over all environments for the

French and Australian data sets, respectively. Note that the estimated contribution of parameter

variance to total variance is always much less for BBCH10, for which there were no observed values, than

for the other simulated development stages.

Table 5 shows results averaged over all environments of both the French and Australian data

sets, for both the variance-components approach and the approach based on comparing usual and

protocol calibration. Separate values are given for BBCH10 and for all other stages. For development

stages other than BBCH10, the estimated contribution of parameter uncertainty is large. It ranges from

24 to 93%, and is larger than 50% for 11 out of the 14 combinations of data set, development stage and

approach for estimating the contribution of parameter variance.

| | Apache | | | Bermude | | |
|---|---|---|---|---|---|---|
| Development stage | BBCH10 | BBCH30 | BBCH55 | BBCH10 | BBCH30 | BBCH55 |
| Ave number of modeling groups | 23 | 25 | 27 | 23 | 25 | 27 |
| Number of model structures | 17 | 19 | 21 | 17 | 19 | 21 |
| Parameter variance (days²) | 11.74 | 52.94 | 26.77 | 12.05 | 71.25 | 36.41 |
| Structure variance (days²) | 5.05 | 4.29 | 9.65 | 4.99 | 10.86 | 12.73 |
| Parameter variance/total | 0.55 | 0.93 | 0.81 | 0.56 | 0.92 | 0.8 |

**Table 3**

**Parameter and structure variance for French data sets estimated using the variance**

**components calculation. The variance components estimate is based on ensemble studies**

**where some model structures were used by multiple modeling groups. Results are averages**

**over environments. The ratio of parameter variance to total (last row) is an average over**

**the ratios for each environment.**

| Development stage | BBCH10 | BBCH30 | BBCH65 | BBCH90 |
|---|---|---|---|---|
| Average number of modeling groups | 22.95 | 23.93 | 27.64 | 26.05 |
| Average number of model structures | 17.95 | 19.93 | 22.64 | 21.05 |
| Parameter variance (days²) | 40.69 | 102.88 | 47.1 | 30.3 |
| Structure variance (days²) | 276.33 | 45.74 | 48.4 | 90.71 |
| Parameter variance / total | 0.12 | 0.66 | 0.49 | 0.24 |

**Table 4**

**Parameter and structure variance for the Australian data set estimated using the variance components calculation. The variance components estimate is based on ensemble studies where some model structures were used by multiple modeling groups. Results are averages over environments. The ratio of parameter variance to total (last row) is an average over the ratios for each environment.**

| Method for estimating parameter variance as fraction of total | BBCH10 | All other stages |
|---|---|---|
| Variance components | 0.41 | 0.69 |
| Usual compared to protocol calibration | 0.23 | 0.69 |

**Table 5**

**Fraction of total variance due to parameter variance based on two estimation procedures: variance components and comparison of usual and protocol calibration. Averages are over all environments of both the French and Australian data sets, either for the development stage BBCH10 or for all other development stages.**

## 4. Discussion

. The variability in parameters between different groups using the same model structure results from differences in fixed parameter values and in calibration approach. None of the data sets here

included time to emergence (BBCH10), so variability in simulating time to emergence likely depends only

on variability in default values. On the other hand, there are data for subsequent development stages, so

simulation of those stages will depend on the values of calibrated parameters. The parameter

uncertainty contribution to variability in days to emergence is always smaller (41% on average) than the

contribution to variability in days to other stages (69% on average). This shows that the importance of

parameter uncertainty depends on both the specific variable simulated and on the data available for

calibration. It seems that variability in simulated variables affected by calibration is larger than variability

in simulated variables which depend just on default parameter values.

When averaging over data sets and simulated variables, for variables other than time to

emergence, the resuts show that parameter uncertainty represents 69% of total uncertainty. The same

value is obtained using variance components or the comparison of usual and protocol calibration.

Structure uncertainty contributes 31%, so the contribution of parameter uncertainty is more than twice

as large as the contribution of structure uncertainty. Previous studies that have estimated structure and

parameter uncertainty contributions to overall uncertainty of plant or ecosystem models have generally

found that structure uncertainty is larger than parameter uncertainty (Tao et al., 2018; Xiong et al.,

2020), including studies that have specifically considered simulation of phenology (Migliavacca et al.,

2012; Wallach et al., 2017; Zhang et al., 2017). This is in fact not too surprising. Previous studies have

quantified the effect of some specific source of parameter uncertainty, such as observation error, or *a*

*priori* uncertainty, or choice of data to use. The study here on the other hand takes into account the full

variability in default values and calibration approach between different modeling groups who could

participate in open-call multi-model studies.

The present study has several limitations, which must be taken into account before generalizing

the results. First, this is a case study, so the specific results refer to the specific cases studied here.

However, the studies use three different data sets, with 22 or 42 different environments each. The

French and Australian data sets not only have quite different environmental conditions and plant

material (winter wheat in France and spring wheat in Australia), but also quite different measured

variables (two development stages for the French data set, weekly development stage for the Australian

data set). There is substantial variability in results for different environments and different simulated

variables, but the range of situations analyzed gives some confidence in the overall conclusion, that

parameter uncertainty is a major contribution to overall model uncertainty when simulating phenology.

A second limitation is that the study concerns only simulation of phenology. It remains to be seen

whether parameter uncertainty is also dominant when simulating crop growth and yield. Calibration in

that case is more complex, which may lead to larger calibration uncertainties, but also the simulations

involve more equations, which may lead to more structure uncertainty. In any case, it seems probable

that taking into account the full variability in plausible parameters will lead to a larger contribution of

parameter uncertainty to total uncertainty than previously thought, for all types of process-based

models. A third limitation is that the results rely on the variability between different groups that use the

same model structure, but only three model structures were used by more than one group, and at most

four groups used the same model structure. However, we also have a second estimate of the

contribution of parameter uncertainty, based on the comparison of usual and protocol calibration. This

second estimate is based on a sample of two different calibration approaches for all models. The average

contribution of parameter to total uncertainty using this second approach, again ignoring simulations of

time to emergence, was 69%, essentially the same as the value obtained using the variance components

analysis. This increases confidence in the variance component results.

The analysis of variance analysis done here can be easily applied to the results of any open-call

multi-model study where some model structures are used by more than a single modeling team. This

does not require any additional simulations, and the calculations can be easily done using existing

software.

## 5. Conclusions

The large variability in crop multi-model simulations reduces the confidence one can have in

such simulations, so it is important to understand the causes of this variability. A realistic estimate of

parameter uncertainty must take into account the actual variability in parameters between different

modeling groups. We illustrate a simple method of doing so, based on results of multi-model studies

associated with random effects analysis of variance. This methodology is easily applicable to any case

where the necessary multi-model results exist.

The parameter uncertainty taken into account here is different than in previous studies, where

specific sources of parameter uncertainty were examined. Unlike those previous studies, we find in the

case study here that parameter uncertainty makes the largest contribution to overall uncertainty,

substantially larger than structure uncertainty. The larger parameter uncertainty found in the present

study is unsurprising, since multiple sources of parameter uncertainty are included here, which suggests

that it may be generally true that parameter uncertainty makes a larger contribution to total uncertainty

than previously thought.

The importance of parameter uncertainty emphasizes the need for giving more attention to

parameter values and in particular to Improved calibration approaches and procedures. Improving

parameter values should be a major goal of the crop modeling community, on a par with work on

improving model equations.

## 6. Acknowledgements

This work was in part supported by the Collaborative Research Center 1253 CAMPOS (Project 7:

Stochastic Modelling Framework), funded by the German Research Foundation (DFG, Grant Agreement

SFB 1253/1 2017), the Academy of Finland through projects AICropPro (316172) and DivCSA (316215),

the BonaRes projects ''Soil3'' (BOMA 03037514) and "I4S" (031B053I) of the Federal Ministry of

Education and Research (BMBF), Germany, the Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation) under Germany's Excellence Strategy - EXC 2070 – 390732324 EXC (PhenoRob),

the project BiomassWeb of the GlobeE programme (Grant number: FKZ031A258B) funded by the Federal

Ministry of Education and Research (BMBF, Germany), the INRA ACCAF meta-programme, the German

Federal Ministry of Education and Research (BMBF) in the framework of the funding measure "Soil as a

Sustainable Resource for the Bioeconomy —- BonaRes", project "BonaRes (Module B): BonaRes Centre

for Soil Research, subproject B" (grant 031B0511B), the National Key Research and Development

Program of China (2017YFD0300205), the National Science Foundation for Distinguished Young Scholars

(31725020), the Priority Academic Program Development of Jiangsu Higher Education Institutions

(PAPD), the 111 project (B16026), and China Scholarship Council, the Agriculture and Agri-Food Canada's

Project J-002303 under the Interdepartmental Research Initiative in Agriculture, the DFG Research Unit

FOR 1695 'Agricultural Landscapes under Global Climate Change – Processes and Feedbacks on a

Regional Scale, the U.S. Department of Agriculture National Institute of Food and Agriculture (award no.

2015-68007-23133) and USDA/NIFA HATCH grant N. MCL02368, the National Key Research and

Development Program of China (2016YFD0300105), The Broadacre Agriculture Initiative, a research

partnership between University of Southern Queensland and the Queensland Department of Agriculture

and Fisheries, the Academy of Finland through project AI-CropPro (315896), the JPI FACCE MACSUR2

project, funded by the Italian Ministry for Agricultural, Food, and Forestry Policies (D.M. 24064/7303/15

of 6/Nov/2015), the SYSTEMIC funded by JPI HDHL, JPI-OCEANS and FACCE-JPI under ERA-NET

(n.696295), the SustEs project funded by the Ministry of Education, Youth and Sports of the Czech

Republic (CZ.02.1.01/0.0/0.0/16_019/000797). The time of MTH and KL was supported by the Grains

Research and Development Corporation (Project UOT1906-002RTX). The observed Australian phenology

data were jointly funded by CSIRO, the Grains Research and Development Corporation (GRDC) under the

"Adding Value to GRDC's National Variety Trial Network" project (CSA00027). The order in which the

donors are listed is arbitrary.

# 476     7. References

Ahmad, S., Abbas, G., Ahmed, M., Fatima, Z., Anjum, M.A., Rasul, G., Khan, M.A., Hoogenboom, G., 2019.

Climate warming and management impact on the change of phenology of the rice-wheat cropping

system in Punjab, Pakistan. F. Crop. Res. 230, 46–61. https://doi.org/10.1016/j.fcr.2018.10.008

Ahuja, L.R., Ma, L., 2011. A Synthesis of Current Parameterization Approaches and Needs for Further

Improvements, in: Methods of Introducing System Models into Agricultural Research, Advances in

Agricultural Systems Modeling. pp. 427–440.

https://doi.org/https://doi.org/10.2134/advagricsystmodel2.c15

Albanito, F., McBey, D., Harrison, M., Smith, P., Ehrhardt, F., Bhatia, A., Bellocchi, G., Brilli, L., Carozzi, M.,

Christie, K., Doltra, J., Dorich, C., Doro, L., Grace, P., Grant, B., Léonard, J., Liebig, M., Ludemann, C.,

Martin, R., Meier, E., Meyer, R., De Antoni Migliorati, M., Myrgiotis, V., Recous, S., Sándor, R.,

Snow, V., Soussana, J.-F., Smith, W.N., Fitton, N., 2022. How Modelers Model: the Overlooked

Social and Human Dimensions in Model Intercomparison Studies. Environ. Sci. Technol. 56, 13485–

13498. https://doi.org/10.1021/acs.est.2c02023

Asseng, S., Ewert, F., Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P.J.,

Rötter, R.P., Cammarano, D., Brisson, N., Basso, B., Martre, P., Aggarwal, P.K., Angulo, C., Bertuzzi,

P., Biernath, C., Challinor, A.J., Doltra, J., Gayler, S., Goldberg, R., Grant, R., Heng, L., Hooker, J.,

Hunt, L.A., Ingwersen, J., Izaurralde, R.C., Kersebaum, K.C., Müller, C., Naresh Kumar, S., Nendel, C.,

O'Leary, G., Olesen, J.E., Osborne, T.M., Palosuo, T., Priesack, E., Ripoche, D., Semenov, M.A.,

Shcherbak, I., Steduto, P., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Travasso, M.,

Waha, K., Wallach, D., White, J.W., Williams, J.R., Wolf, J., 2013. Uncertainty in simulating wheat

yields under climate change. Nat. Clim. Chang. 3, 827–832. https://doi.org/10.1038/nclimate1916

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using lme4. J.

Stat. Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Bindi, M., Palosuo, T., Trnka, M., Semenov, M., 2015. Modelling climate change impacts on crop

production for food security. Clim. Res. 65, 3–5. https://doi.org/10.3354/cr01342

Boote, K.J., Jones, J.W., Hoogenboom, G., White, J.W., 2010. The Role of Crop Systems Simulation in

Agriculture and Environment. Int. J. Agric. Environ. Inf. Syst. 1, 41–54.

Confalonieri, R., Orlando, F., Paleari, L., Stella, T., Gilardelli, C., Movedi, E., Pagani, V., Cappelli, G.,

Vertemara, A., Alberti, L., Alberti, P., Atanassiu, S., Bonaiti, M., Cappelletti, G., Ceruti, M.,

Confalonieri, A., Corgatelli, G., Corti, P., Dell'Oro, M., Ghidoni, A., Lamarta, A., Maghini, A.,

Mambretti, M., Manchia, A., Massoni, G., Mutti, P., Pariani, S., Pasini, D., Pesenti, A., Pizzamiglio, G.,

Ravasio, A., Rea, A., Santorsola, D., Serafini, G., Slavazza, M., Acutis, M., 2016. Uncertainty in crop

model predictions: What is the role of users? Environ. Model. Softw. 81, 165–173.

https://doi.org/10.1016/j.envsoft.2016.04.009

Durand, J.-L., Delusca, K., Boote, K., Lizaso, J., Manderscheid, R., Weigel, H.J., Ruane, A.C., Rosenzweig, C.,

Jones, J., Ahuja, L., Anapalli, S., Basso, B., Baron, C., Bertuzzi, P., Biernath, C., Deryng, D., Ewert, F.,

Gaiser, T., Gayler, S., Heinlein, F., Kersebaum, K.C., Kim, S.-H., Müller, C., Nendel, C., Olioso, A.,

Priesack, E., Villegas, J.R., Ripoche, D., Rötter, R.P., Seidel, S.I., Srivastava, A., Tao, F., Timlin, D.,

Twine, T., Wang, E., Webber, H., Zhao, Z., 2018. How accurately do maize crop models simulate the

interactions of atmospheric $CO_2$ concentration levels with limited water supply on water use and

yield? Eur. J. Agron. 100, 67–75. https://doi.org/10.1016/J.EJA.2017.01.002

Fatima, Z., Ahmed, M., Hussain, M., Abbas, G., Ul-Allah, S., Ahmad, S., Ahmed, N., Ali, M.A., Sarwar, G.,

Haque, E. ul, Iqbal, P., Hussain, S., 2020. The fingerprints of climate warming on cereal crops

phenology and adaptation options. Sci. Rep. 10, 18013. https://doi.org/10.1038/s41598-020-

74740-3

Gao, Y., Wallach, D., Liu, B., Dingkuhn, M., Boote, K.J., Singh, U., Asseng, S., Kahveci, T., He, J., Zhang, R.,

Confalonieri, R., Hoogenboom, G., 2020. Comparison of three calibration methods for modeling rice

phenology. Agric. For. Meteorol. 280, 107785. https://doi.org/10.1016/J.AGRFORMET.2019.107785

Harrison, M.T., Roggero, P.P., Zavattaro, L., 2019. Simple, efficient and robust techniques for automatic

multi-objective function parameterisation: Case studies of local and global optimisation using

APSIM. Environ. Model. Softw. 117, 109–133. https://doi.org/10.1016/j.envsoft.2019.03.010

Houska, T., Kraft, P., Chamorro-Chavez, A., Breuer, L., 2015. SPOTting Model Parameters Using a Ready-

Made Python Package. PLoS One 10, e0145180. https://doi.org/10.1371/journal.pone.0145180

Liu, Y., Bachofen, C., Wittwer, R., Silva Duarte, G., Sun, Q., Klaus, V.H., Buchmann, N., 2022. Using

PhenoCams to track crop phenology and explain the effects of different cropping systems on yield.

Agric. Syst. 195, 103306. https://doi.org/10.1016/J.AGSY.2021.103306

Maiorano, A., Martre, P., Asseng, S., Ewert, F., Müller, C., Rötter, R.P., Ruane, A.C., Semenov, M.A.,

Wallach, D., Wang, E., Alderman, P.D., Kassie, B.T., Biernath, C., Basso, B., Cammarano, D.,

Challinor, A.J., Doltra, J., Dumont, B., Rezaei, E.E., Gayler, S., Kersebaum, K.C., Kimball, B.A., Koehler,

A.-K., Liu, B., O'Leary, G.J., Olesen, J.E., Ottman, M.J., Priesack, E., Reynolds, M., Stratonovitch, P.,

Streck, T., Thorburn, P.J., Waha, K., Wall, G.W., White, J.W., Zhao, Z., Zhu, Y., 2017. Crop model

improvement reduces the uncertainty of the response to temperature of multi-model ensembles.

F. Crop. Res. 202. https://doi.org/10.1016/j.fcr.2016.05.001

Meier, U., 1997. Growth stages of mono- and dicotyledonous plants : BBCH-Monograph [WWW

Document]. URL

https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKEwjBo-

Ln4_r4AhVM3RoKHc-

TA5sQFnoECAcQAQ&url=https%3A%2F%2Fwww.politicheagricole.it%2Fflex%2FAppData%2FWebLi

ve%2FAgrometeo%2FMIEPFY800%2FBBCHengl2001.pdf&usg=AOvVaw1G_eUh_-qnVt91_qbj44d

(accessed 1.9.19).

Migliavacca, M., Sonnentag, O., Keenan, T.F., Cescatti, A., O'Keefe, J., Richardson, A.D., 2012. On the

uncertainty of phenological responses to climate change, and implications for a terrestrial

biosphere model. Biogeosciences 9, 2063–2083. https://doi.org/10.5194/bg-9-2063-2012

Muleke, A., Harrison, M.T., de Voil, P., Hunt, I., Liu, K., Yanotti, M., Eisner, R., 2022. Earlier crop flowering

caused by global warming alleviated by irrigation. Environ. Res. Lett. 17, 044032.

https://doi.org/10.1088/1748-9326/ac5a66

Müller, C., Franke, J., Jägermeyr, J., Ruane, A.C., Elliott, J., Moyer, E., Heinke, J., Falloon, P.D., Folberth, C.,

Francois, L., Hank, T., Izaurralde, R.C., Jacquemin, I., Liu, W., Olin, S., Pugh, T.A.M., Williams, K.,

Zabel, F., 2021. Exploring uncertainties in global crop yield projections in a large ensemble of crop

models and CMIP5 and CMIP6 climate scenarios. Environ. Res. Lett. 16, 034040.

https://doi.org/10.1088/1748-9326/abd8fc

R Core Team, 2017. A language and Environment for Statistical Computing.

Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P., Antle, J.M., Nelson, G.C.,

Porter, C., Janssen, S., Asseng, S., Basso, B., Ewert, F., Wallach, D., Baigorria, G., Winter, J.M., 2013.

The Agricultural Model Intercomparison and Improvement Project (AgMIP): Protocols and pilot

studies. Agric. For. Meteorol. 170. https://doi.org/10.1016/j.agrformet.2012.09.011

Rötter, R.P., Carter, T.R., Olesen, J.E., Porter, J.R., 2011. Crop–climate models need an overhaul. Nat.

Clim. Chang. 1, 175–177. https://doi.org/10.1038/nclimate1152

Salo, T.J., Palosuo, T., Kersebaum, K.C., Nendel, C., Angulo, C., Ewert, F., Bindi, M., Calanca, P., Klein, T.,

Moriondo, M., Ferrise, R., Olesen, J.E., Patil, R.H., Ruget, F., Takac, J., Hlavinka, P., Trnka, M., Rotter,

R.P., 2016. Comparing the performance of 11 crop simulation models in predicting yield response

to nitrogen fertilization. J. Agric. Sci. 154, 1218–1240. https://doi.org/10.1017/S0021859615001124

Sándor, R., Ehrhardt, F., Grace, P., Recous, S., Smith, P., Snow, V., Soussana, J.-F., Basso, B., Bhatia, A.,

Brilli, L., Doltra, J., Dorich, C.D., Doro, L., Fitton, N., Grant, B., Harrison, M.T., Kirschbaum, M.U.F.,

Klumpp, K., Laville, P., Léonard, J., Martin, R., Massad, R.-S., Moore, A., Myrgiotis, V., Pattey, E.,

Rolinski, S., Sharp, J., Skiba, U., Smith, W., Wu, L., Zhang, Q., Bellocchi, G., 2020. Ensemble

modelling of carbon fluxes in grasslands and croplands. F. Crop. Res. 252, 107791.

https://doi.org/10.1016/j.fcr.2020.107791

Scheffé, H., 1959. The analysis of variance. John Wiley & Sons, New York.

Seber, G.A.F., Wild, C.J., 1989. Nonlinear regression. Wiley , New York.

Smallman, T.L., Milodowski, D.T., Neto, E.S., Koren, G., Ometto, J., Williams, M., 2021. Parameter

uncertainty dominates C-cycle forecast  errors over most of Brazil for the 21st century. Earth Syst.

Dynam. 12, 1191–1237. https://doi.org/10.5194/esd-12-1191-2021

Tao, F., Rötter, R.P., Palosuo, T., Gregorio Hernández Díaz-Ambrona, C., Mínguez, M.I., Semenov, M.A.,

Kersebaum, K.C., Nendel, C., Specka, X., Hoffmann, H., Ewert, F., Dambreville, A., Martre, P.,

Rodríguez, L., Ruiz-Ramos, M., Gaiser, T., Höhn, J.G., Salo, T., Ferrise, R., Bindi, M., Cammarano, D.,

Schulman, A.H., 2018. Contribution of crop model structure, parameters and climate projections to

uncertainty in climate change impact assessments. Glob. Chang. Biol. 24, 1291–1307.

https://doi.org/10.1111/gcb.14019

Wallach, D., Nissanka, S.P., Karunaratne, A.S., Weerakoon, W.M.W., Thorburn, P.J., Boote, K.J., Jones,

J.W., 2017. Accounting for both parameter and model structure uncertainty in crop model

predictions of phenology: A case study on rice. Eur. J. Agron. 88.

https://doi.org/10.1016/j.eja.2016.05.013

Wallach, D., Palosuo, T., Thorburn, P., Gourdain, E., Asseng, S., Basso, B., Buis, S., Crout, N., Dibari, C.,

Dumont, B., Ferrise, R., Gaiser, T., Garcia, C., Gayler, S., Ghahramani, A., Hochman, Z., Hoek, S.,

Hoogenboom, G., Horan, H., Huang, M., Jabloun, M., Jing, Q., Justes, E., Kersebaum, K.C.,

Klosterhalfen, A., Launay, M., Luo, Q., Maestrini, B., Mielenz, H., Moriondo, M., Nariman Zadeh, H.,

Olesen, J.E., Poyda, A., Priesack, E., Pullens, J.W.M., Qian, B., Schütze, N., Shelia, V., Souissi, A.,

Specka, X., Srivastava, A.K., Stella, T., Streck, T., Trombi, G., Wallor, E., Wang, J., Weber, T.K.D.,

Weihermüller, L., de Wit, A., Wöhling, T., Xiao, L., Zhao, C., Zhu, Y., Seidel, S.J., 2021a. How well do

crop modeling groups predict wheat phenology, given calibration data from the target population?

Eur. J. Agron. 124, 126195. https://doi.org/https://doi.org/10.1016/j.eja.2020.126195

Wallach, D., Palosuo, T., Thorburn, P., Hochman, Z., Andrianasolo, F., Asseng, S., Basso, B., Buis, S., Crout,

N., Dumont, B., Ferrise, R., Gaiser, T., Gayler, S., Hiremath, S., Hoek, S., Horan, H., Hoogenboom, G.,

Huang, M., Jabloun, M., Jansson, P.-E., Jing, Q., Justes, E., Kersebaum, K.C., Launay, M., Lewan, E.,

Luo, Q., Maestrini, B., Moriondo, M., Olesen, J.E., Padovan, G., Poyda, A., Priesack, E., Pullens,

J.W.M., Qian, B., Schütze, N., Shelia, V., Souissi, A., Specka, X., Kumar Srivastava, A., Stella, T.,

Streck, T., Trombi, G., Wallor, E., Wang, J., Weber, T.K.D., Weihermüller, L., de Wit, A., Wöhling, T.,

Xiao, L., Zhao, C., Zhu, Y., Seidel, S.J., 2021b. Multi-model evaluation of phenology prediction for

wheat in Australia. Agric. For. Meteorol. 298–299, 108289.

https://doi.org/10.1016/J.AGRFORMET.2020.108289

Wallach, D., Palosuo, T., Thorburn, P., Hochman, Z., Gourdain, E., Andrianasolo, F., Asseng, S., Basso, B.,

Buis, S., Crout, N., Dibari, C., Dumont, B., Ferrise, R., Gaiser, T., Garcia, C., Gayler, S., Ghahramani,

A., Hiremath, S., Hoek, S., Horan, H., Hoogenboom, G., Huang, M., Jabloun, M., Jansson, P.-E., Jing,

Q., Justes, E., Kersebaum, K.C., Klosterhalfen, A., Launay, M., Lewan, E., Luo, Q., Maestrini, B.,

Mielenz, H., Moriondo, M., Nariman Zadeh, H., Padovan, G., Olesen, J.E., Poyda, A., Priesack, E.,

Pullens, J.W.M., Qian, B., Schütze, N., Shelia, V., Souissi, A., Specka, X., Srivastava, A.K., Stella, T.,

Streck, T., Trombi, G., Wallor, E., Wang, J., Weber, T.K.D., Weihermüller, L., de Wit, A., Wöhling, T.,

Xiao, L., Zhao, C., Zhu, Y., Seidel, S.J., 2021c. The chaos in calibrating crop models: Lessons learned

from a multi-model calibration exercise. Environ. Model. Softw. 145, 105206.

https://doi.org/10.1016/J.ENVSOFT.2021.105206

Wallach, D., Palosuo, T., Thorburn, P., Mielenz, H., Buis, S., Hochman, Z., Gourdain, E., Andrianasolo, F.,

Dumont, B., Ferrise, R., Gaiser, T., Garcia, C., Gayler, S., Harrison, M., Hiremath, S., Horan, H.,

Hoogenboom, G., Jansson, P.-E., Jing, Q., Justes, E., Kersebaum, K.-C., Launay, M., Lewan, E., Liu, K.,

Mequanint, F., Moriondo, M., Nendel, C., Padovan, G., Qian, B., Schütze, N., Seserman, D.-M.,

Shelia, V., Souissi, A., Specka, X., Srivastava, A.K., Trombi, G., Weber, T.K.D., Weihermüller, L.,

Wöhling, T., Seidel, S.J., 2023. Proposal and extensive test of a calibration protocol for crop

phenology models. Agron. Sustain. Dev. 43, 46. https://doi.org/10.1007/s13593-023-00900-0

Wang, B., Feng, P., Liu, D.L., O'Leary, G.J., Macadam, I., Waters, C., Asseng, S., Cowie, A., Jiang, T., Xiao,

D., Ruan, H., He, J., Yu, Q., 2020. Sources of uncertainty for wheat yield projections under future

climate are site-specific. Nat. Food 1, 720–728. https://doi.org/10.1038/s43016-020-00181-w

Wang, B., Jägermeyr, J., O'Leary, G.J., Wallach, D., Ruane, A.C., Feng, P., Li, L., Liu, D.L., Waters, C., Yu, Q.,

Asseng, S., Rosenzweig, C., 2024. Pathways to identify and reduce uncertainties in agricultural

climate impact assessments. Nat. Food 1–7. https://doi.org/10.1038/s43016-024-01014-w

Xiong, W., Asseng, S., Hoogenboom, G., Hernandez-Ochoa, I., Robertson, R., Sonder, K., Pequeno, D.,

Reynolds, M., Gerard, B., 2020. Different uncertainty distribution between high and low latitudes in

modelling warming impacts on wheat. Nat. Food 1, 63–69. https://doi.org/10.1038/s43016-019-

0004-2

Zadeh, F.K., Nossent, J., W., B., Bauwens, W., Van Griensven, A., 2022. A fast and effective

parameterization of water quality models. Environ. Model. Softw. 149, 105331.

https://doi.org/https://doi.org/10.1016/j.envsoft.2022.105331

Zhang, S., Tao, F., Shi, R., 2014. Modeling the rice phenology and production in China with SIMRIW:

sensitivity analysis and parameter estimation. Front. Earth Sci. 8, 505–511.

https://doi.org/10.1007/s11707-014-0468-1

Zhang, S., Tao, F., Zhang, Z., 2017. Uncertainty from model structure is larger than that from model

parameters in simulating rice phenology in China. Eur. J. Agron. 87, 30–39.

https://doi.org/10.1016/j.eja.2017.04.004