Reduced motoneuron excitability and common synaptic inputs of the aged human vastus lateralis

Yuxiao Guo[1], Eleanor J. Jones[1], Jakob Škarabot[2], Thomas B. Inns[1], Bethan E. Phillips[1], Philip J. Atherton[1], Mathew Piasecki[1]

1. *Centre of Metabolism, Ageing & Physiology (COMAP), MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research & National Institute for Health Research (NIHR) Nottingham Biomedical Research Centre, School of Medicine, University of Nottingham, Derby, UK*
2. *School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, UK*

Abstract

The age-related loss of neuromuscular function has been comprehensively documented with a disproportionate loss in the quadriceps compared to other muscle groups. Although muscle atrophy may partially account for strength decline, it is further influenced by alterations of the nervous system. The outputs of motoneurons are heavily dependent on their intrinsic excitability and the level of common synaptic inputs they receive. However, these have not yet been described in aged human quadriceps.

High density surface electromyography (HDsEMG) signals were collected from the vastus lateralis of ten young (mean±SD, 24 ± 6.2 y) and ten older (66 ± 7.7 y) men during a 20-s ramp contraction peaking at 20% of maximal voluntary isometric contraction. EMG signals were decomposed to identify individual motor unit discharges, and motoneuron persistent inward current (PIC) amplitude and intramuscular coherence were estimated.

Older participants produced significantly lower peak knee extensor torque (p<0.001) and poorer force control ability (p=0.002) than young. Older participants also had lower peak discharge rate (p=0.016), PIC amplitudes (p=0.001) and coherence estimates in the beta frequency band in both ascending (p=0.026) and descending phases (p=0.015) when compared to young.

These data provide new evidence that intrinsic motoneuron excitability and common synaptic input is lower in the vastus lateralis of older males when compared to young, and highlights the feasibility of this method in the clinically and functionally important

quadriceps, further underpinning age-related loss of force control occurring
independent of the loss of muscle mass.

Key Words: Ageing; motor unit; persistent inward current; common synaptic input;
vastus lateralis

Key Points

• The age-related loss of muscle mass is exceeded by the loss of function,
which is influenced by structural and functional alterations of the nervous
system.
• Motoneuronal persistent inward currents and common synaptic inputs play an
important role in the activation of motor units and subsequent force generation
and control ability.
• Here we show reduced estimates of persistent inward currents and lower
common synaptic inputs to older vastus lateralis, potentially contributing to
observed lower strength and poorer force tracking.
• These findings highlight decrements of the aged human motor system that act
independently of muscle atrophy in functionally relevant muscle groups which
should be considered in the application of interventions targeting aged human
muscle.

Introduction

Human ageing is characterised by a progressive reduction in muscle size and
muscle strength, resulting in an impairment of force generation, force control and
physical performance (Manini *et al.*, 2013; Wilkinson *et al.*, 2018; Pethick & Piasecki,
2022), all of which can negatively impact quality of life (Daley & Spinks, 2000; Smee
et al., 2012). The loss of muscle size is explicable by the combined effects of muscle
fibre atrophy and muscle fibre loss (Wilkinson et al. 2018), and although atrophy may
partially account for declines in muscle strength, it is further influenced by structural
and functional alterations of the nervous system.

The voluntary contraction of muscle fibres occurs via net excitatory signals to the
motor units (MUs); groups of muscle fibres innervated by a single motoneuron
(Heckman & Enoka, 2012). The central nervous system regulates muscle force via
two primary strategies; varying the number of MUs recruited and varying the
discharge rate of each motor neuron, both of which are susceptible to age-related
alterations. Substantial motoneuron loss accelerates in the sixth decade of life,
resulting in fewer MUs, and a compensatory process of MU remodelling further
results in larger MU innervation ratios (Hepple & Rice, 2016; Piasecki *et al.*, 2016).

Additionally, although this appears to be muscle specific, an age-related decrease in
MU discharge rate has also been documented at normalised contraction levels (Kirk
et al., 2021; Orssatto *et al.*, 2022) as well as being associated with circulating
hormone levels that are also affected by age (Guo *et al.*, 2022*b*).

Beyond releasing neurotransmitters into postsynaptic ligand-gated channels at the
spinal cord level via ionotropic mechanisms, activation of the motor nervous system
also relies on neuromodulation (Heckman *et al.*, 2005). This is mediated by
intracellular second messenger systems that modify the voltage- and ligand-gated
channels of neurons and alters several properties from their level of excitability to
their pattern of firing in response to a given input (Heckman *et al.*, 2008*b*; Nadim &
Bucher, 2014). This process is enhanced by monoamines, e.g. serotonin and
norepinephrine released from the brainstem nuclei by stimulating inward-flowing
persistent calcium and sodium currents (Heckman *et al.*, 200*5*). As long as the
membrane potential remains above the activation threshold, persistent inward
currents (PICs) tend to remain activated (Heckman *et al.*, 2005), amplifying the firing
behaviours of motoneurons to synaptic inputs by three to five fold (Binder & Powers,
2001) as well as generating long-lasting plateau potentials which results in self-
sustaining firing (Gorassini *et al.*, 2002). A reduction in PIC amplitude could result in
a reduction of force and an increase in perceived effort due to suppression of MU
discharge rates. Even though PIC amplitudes appear to decrease with age in upper
(Hassan *et al.*, 2021) and lower limb muscles (Orssatto *et al.*, 2021), PICs can be
mediated by local ionotropic inhibitory inputs that may change differently with ageing
between muscle groups, particularly those susceptible to functional losses such as
the vastus lateralis (VL).

Rather than controlling the discharge pattern of individual MUs, the central nervous
system controls the excitatory inputs to the MU pool, referred to as common synaptic
inputs (Farina *et al.*, 2014) which are defined as the proportion of net synaptic input
correlated between MUs. Force fluctuations are largely determined by the alterations
in low frequency components of common synaptic inputs to MUs, which is reflected
in concurrent fluctuations in MU discharge rates from the same MU pool (Farina &
Negro, 2015). Accordingly, a decrease in force steadiness associated with ageing is
a consequence of an increase in variance of common synaptic inputs (Castronovo *et*
al., 2018; Feeney *et al.*, 2018). However, the variability of common synaptic inputs in

quadriceps has only been explored in healthy young individuals (Avrillon *et al.*, 2020;
Rossato *et al.*, 2022), with no available data from older humans.

It is not possible to make intracellular recordings from human spinal motoneurons in
order to estimate PICs, so alternative methods are employed which distinguish
intrinsic excitability of the MU from its descending synaptic drive by using unit-wise
MU analysis technique (Hassan *et al.*, 2021). In view of the advances with which MU
discharging patterns can be measured, it provides an opportunity to investigate and
better understand the physical properties and the underlying mechanisms which may
impair motor control in older age. Therefore, the aim of this study was to explore the
differences in physical performance as well as the level of neuronal excitability and
common synaptic input in the clinically relevant VL between healthy young and older
males.

Methods
Participants and ethical approval
26 healthy male participants (11 young and 15 older) provided informed consent to
take part in the study, which was approved by the University of Nottingham Faculty
of Medicine and Healthy Science Research Ethics Committee (90-0820, 199-0221)
and conformed to the Declaration of Helsinki. In order to ensure a sufficient number
of MU pairs for PIC amplitude estimation, 20 participants were included; 10 young
(mean ± SD; age, 24 ± 6.2 y; body mass index, 24.5 ± 2.8 kg/m^2) and 10 older (age,
66 ± 7.7 y; body mass index, 25.5 ± 2.6 kg/m^2) males. All participants were recruited
through advertisements in the local community. Prior to enrolment, all participants
completed a comprehensive clinical examination and metabolic screening was
conducted at the School of Medicine, Royal Derby Hospital Centre. All recruited
participants were recreationally active and free from any metabolic disease, lower
limb musculoskeletal abnormalities, neurological disorder, cardiovascular disease
and uncontrolled hypertension.

Muscle cross sectional area

Ultrasound was used to measure the cross-sectional area (CSA) of the VL muscle of
the right leg, using an ultrasound probe (LA523 probe, B-mode, frequency range 26–
32 Hz, and MyLabTM50 scanner, Esaote, Genoa, Italy) at the anatomical mid-point
of the right thigh. Images were recorded with the participants laying in a supine

position and the mid-point obtained from the identification of the greater trochanter at
the top of the femur and the mid-point of the patella. Panoramic imaging (VPAN)
images were obtained in a medial to lateral fashion, beginning and ending the
capture of the image at the aponeurosis borders of the VL of the right leg. The
produced images were subsequently analysed using ImageJ software (National
Institute of Health, USA) through tracing around the contour of the aponeurosis via
polygon function to quantify the area of the VL. Each image was analysed three
times and the mean value of three images was accepted as CSA.

Muscle strength and force tracking accuracy

The participants were required to attend the lab at 0900 after an overnight fast and
abstain from strenuous exercise 24h prior to the testing sessions. Participants were
seated in a custom-built chair with hips and knees flexed at ~90 degree. The lower
leg was secured to a force dynamometer with noncompliant straps above the medial
malleolus. To minimize the movement of the upper trunk during testing, a seat belt
was fastened across the pelvis of the participant. After a standardised warm up
through several submaximal contractions (Guo *et al.*, 2022*a*), participants were
instructed to contract as hard and fast as they could – to the point where the holding
of the contraction is mechanically not possible. During the trial, participants were not
allowed to hold onto the side of the chair and were asked to cross their arms across
the chest. This was performed with real-time visual force feedback on a monitor
placed in front of the participant, with verbal encouragement to aid in producing
maximal effort. This was repeated two to three times further, giving 60 seconds of
rest between attempts standardised recovery. If the difference between the last 2
attempts was < 5%, the highest value recorded in newtons was taken and accepted
as the maximal isometric voluntary contraction force (MVC). Peak torque was
calculated by multiplying MVC by the lever arm length from the knee to the centre of
the ankle strap.

Following a short rest, participants performed four triangular shaped contractions
(10s up and 10s down) to 20% of their MVCs with 30s rests. All force targets were
set using OTBioLab software (OT Bioelecttronica, Turin, Italy). Ramp contractions to
20% MVC have been widely used for PICs estimation (details below) (Gorassini *et*
al., 2002; Hassan *et al.*, 2020; Afsharipour *et al.*, 2020). To assess force tracking
accuracy, all data were exported to Spike2 (version 9.11), where a virtual channel

was created by subtracting the performed path from the requested (target) path and

rectifying it. Area under the curve reflects the level of deviation from the target line

reflecting muscle force tracking accuracy, with higher values indicating greater

deviation.

High-density surface electromyography

A semi-disposable high-density surface electromyography (HDsEMG) array (64

electrodes, 13×5, 8mm, I.E.D., GR08MM1305, OT Bioelettronica, Inc., Turin, Italy)

was placed with the orientation of the muscle fibres (proximal to distal) over the

muscle belly of right vastus lateralis after skin preparation involving shaving, light

abrasion and cleansing with 70% ethanol. Electrodes were secured to the skin using

flexible tape. The adhesive grids were attached to the surface of the muscle by

disposable bi-adhesive foam layers (SpesMedica, Bettipaglia, Italy). The skin

electrode contact was facilitated by filling the cavities of the adhesive layers with

conductive paste (AC Cream, SpesMedica). A strap ground electrode (WS2,

OTBioelettronica, Turin, Italy) dampened with water was positioned around the ankle

of the right leg to optimise the signal quality. HDsEMG signals were acquired in a

monopolar configuration, amplified (x256) with filtering set at 10-500 Hz and digitally

converted at 2000 Hz by a 16-bit wireless amplifier (Sessantaquattro,

OTBioelettronica, Turin, Italy) and transferred to a PC for further offline analysis.

The recorded HDsEMG signals were converted into a MatLab file format with only

one contraction in a single data file and then analysed offline using DEMUSE

software in MatLab. Each participant performed four ramp contractions; however, a

single trial was selected for analysis based on the force tracking accuracy and the

quality of MU discharge profiles. Monopolar HDsEMG signals were band-pass

filtered at 20-500Hz and then decomposed into MU pulse trains by a convolutive

kernel compensation algorithm (Holobar & Zazula, 2007; Holobar *et al.*, 2014). After

the decomposition, a trained investigator manually inspected the MU spike trains and

edited the discharge patterns of each MU detected. Only MUs with a pulse-to-noise

ratio equal or greater than 30 dB were kept for further analysis.

Estimation of persistent inward current amplitude

The instantaneous firing rates of both MUs were calculated as the inverse of the

interspike intervals of each MU spike train and smoothed by fitting a fifth order

polynomial function. An estimate of the PIC amplitude can be derived from the well-
established unit-wise MU analysis technique (Hassan *et al.*, 2021). The lower
threshold unit to be recruited in the ramp contraction is commonly referred to as the
"control" unit; the "test" unit is a unit of higher threshold. Delta F (estimates of PIC
amplitudes) was calculated as the difference in "control" unit firing rate between the
onset and offset of a "test" unit (Gorassini *et al.*, 2002). This technique relies on
several assumptions: 1) test and control units share the common synaptic drive; 2)
PIC is activated before or at the recruitment; 3) firing rate of the control MU closely
reflects the depolarizing input to the parent motoneuron; 4) test and control unit
process the synaptic inputs in a similar way. The MU pairs for the delta F calculation
were selected based on the following criteria: 1) as a measure of common synaptic
modulation, rate-to-rate correlation r ≥ 0.7; 2) to avoid the high variability in delta F
calculation due to the initial acceleration phase of MU firings, test units were
recruited at least 1s after the control units; 3) to account for the possibility of control
unit saturation leading to an underestimation of delta F, pairs in which rate
modulation of the control unit fell within 0.5pps were removed from analysis
(Gorassini *et al.*, 2002; Udina *et al.*, 2010; Vandenberk & Kalmar, 2014; Hassan *et*
al., 2020). An average delta F was calculated when individual test units were paired
with multiple control units (Figure 1). Subsequently, a single delta F was determined
from each participant in VL by averaging all the delta F values obtained for all the
units from each participant. Peak discharge rates were measured as the highest
value in fifth order polynomials for each MU.

Figure 1. Example data showing delta frequency (∆F) calculation in vastus lateralis for paired-motor unit technique. Paired motor unit data from a young male is displayed in the left panels (mean ∆F=3.09) and from an older male in the right panels (mean ∆F=1.52). Test units are displayed in panel A and corresponding control units are in B, C and D.

Coherence Estimates

To estimate the level of common synaptic input, intramuscular coherence was calculated during ascending and descending phase of a ramp contraction separately (Figure 2), which represents a frequency domain correlation between cumulative spike trains (CSTs) of the identified motor units. The magnitude-squared coherence was calculated using the Welch's averaged periodogram with non-overlapping Hann windows of 1s at the different frequency bands: delta (0-5 Hz), alpha (5-12 Hz), beta (15-30 Hz) and piper (40-50 Hz) (Raethjen *et al.*, 2007; Laine & Valero-Cuevas, 2017). This analysis was repeated for 60 randomly chosen combinations of two equally sized CSTs calculated as the sum of firing times from three MUs randomly selected from the same set of MUs and then averaged. The Fisher Z-transform was

applied to the coherence estimates to obtain a normally distributed variable for
comparisons.

Figure 2. The group mean intramuscular coherence Z-Scores during ascending and
descending phase of a ramp contraction at 20% maximal isometric voluntary
contraction in young (blue) and older (yellow) participants. The magnitude-squared
coherence was calculated from all the combinations (a maximum of 60 random
permutations) of two equally sized cumulative spike trains, each composed by three
motor units.

Statistical analysis

Data management and analysis were performed using RStudio (Version 2022.07.1).
Descriptive data were generated for all variables. *An independent t-test* was
conducted to test whether any differences existed between young and older males in
torque, force tracking accuracy and coherence estimates. In order to preserve
variability within and across participants simultaneously to the greatest extent,
multilevel linear regression models were generated to compare peak discharge rate,
delta F, recruitment threshold and derecruitment threshold between groups using
lme4 package (Version 1.1-27.1). Recruitment threshold was included as a covariate
in the model to explore peak discharge rate and discharge rate range. In the
multilevel models, single MU was regarded as the first level; and individual

participant with clustered MUs was considered as the second level. To better show

the magnitude of the standardized mean difference between groups, the absolute

difference was also calculated by *Hedges' g effect size (ES). Simple linear*

regression models were used to explore the relationship between age, peak

discharge rate and delta F. For data visualization, individual participant means were

displayed in box-and-jitter plots. A *p* value <0.05 was considered significant.

Results

During the ramp contractions peaking at 20% of MVC, a total of 197 MUs were

identified from the right VL of 10 young males and 175 MUs from the right VL of 10

older males. For Delta F calculations, there was an average of 5 (SD: 3.68) pairs of

test and control units per person in the young and 6 (4.95) pairs in older group, with

the mean number of 4 (1.79) test units per person in young and 3 (1.97) in old.

There was no statistical difference in muscle cross sectional area (Y v O: 29.22 ±

4.88 v 23.59 ± 7.88 cm^2, ES = 0.82, p=0.071; Figure 3A), however the younger

group had greater muscle strength (279.60 ± 84.77 v 153.50 ± 38.06 Nm, ES = 1.84,

p<0.001; Figure 3B) when compared to the older group. There was a significant

difference in force tracking accuracy between groups, with the younger group

performing better (9.38 ± 3.88 v 17.54 ± 7.01 Ns, ES = -1.38, p=0.005; Figure 3D)

during a whole ramp contraction. When contraction phases were explored separately,

there was no group difference in the ascending phase (5.13 ± 3.12 v 7.32 ± 3.10 Ns,

ES = -0.67, p=0.132; Figure 3E), yet younger participants showed better force

control ability than the older participants in the descending phase (4.27 ± 1.03 v

10.26 ± 5.03 Ns, ES = -1.58, p=0.002; Figure 3F).

Figure 3. Scatter plots with means and SDs showing the differences in cross-sectional area (A) and torque (B) between young (blue) and older (yellow) males; during a single ramp contraction at 20% of maximal voluntary contraction (C), the differences in force tracking accuracy during whole contraction (D) as well as during ascending phase (E) and descending phase (F). Note: ***p<0.001, **p<0.010, *p<0.05.

The younger group had significantly higher peak discharge rates (10.64 ± 1.59 v 9.60 ± 1.34 pps, ES = 0.67, p=0.016; Figure 4A) and delta F values than the old (3.11 ± 0.77 v 1.86 ±0.65 pps, ES = 1.68, p=0.006; Figure 4B). In each group there were no significant relationships between delta F and peak discharge rate (Y: p=0.694; O: p=0.844) or between age and delta F, or between age and peak discharge rate (all p>0.05).

Figure 4. Box-and-jitter plots of the individual participant mean value and their corresponding rainclouds plots of peak discharge rate (A) and Delta F (B) in young (blue) and older (yellow) males. Note: ***p<0.001, **p<0.010, *p<0.05.

Younger participants had a greater mean discharge rate range in both ascending (3.57 ± 0.70 v 3.51 ± 0.81 pps, ES = 0.07, p=0.029) and descending (5.38 ± 1.22 v 4.88 ± 0.84 pps, ES = 0.46, p=0.022; Figure 5A) phases of the ramp contraction.

Figure 5. Box-and-jitter plots of the individual participant mean value and their corresponding rainclouds plots of discharge rate range (A) and recruitment threshold and derecruitment threshold (B) in young (blue) and older (yellow) males. Note: ***p<0.001, **p<0.010, *p<0.05.

There was a significant difference in MU recruitment threshold between young and older groups (Figure 5B). The mean recruitment threshold for old occurred at 6% MVC, significantly lower that the threshold for young, which was 10% MVC (p=0.001). During the descending phase of the contraction, significantly longer discharging of the activated MUs were observed in older group (p<0.001). The mean derecruitment threshold of the activated MUs in younger group occurred at 9.4% of MVC, with a mean derecuritment threshold at 5.6% of MVC in the older group.

There was no significant age-related difference in the COH Delta, Alpha or Piper

bands in either ascending or descending phase of the ramp contraction (all p>0.05;

Figure 6). Younger participants had greater coherence estimates in the beta

frequency band in both ascending (p=0.026) and descending phases (p=0.015)

when compared to older participants.

Figure 6. Individual participant means of intramuscular coherence Z-score at
different frequency bands during ascending and descending phases of a ramp
contraction at 20% maximal voluntary contraction in young (blue) and older (yellow)
males.

Discussion

Here we provide the first evidence for reduced neuronal excitability in the aged

human VL, a muscle with established susceptibility for age-related functional losses.

Our findings indicate that younger individuals have better force control ability during

a ramped contraction when compared to older individuals, and notably, this was

explained by an inferior tracking ability during the descending phase in the older

group. Older participants had lower peak discharge rate, smaller discharge rate

range, and lower mean recruitment and derecruitment thresholds compared to young.

We also show age-related differences in common synaptic input to the VL, with a
reduced commonality during both ascending and descending phases. These findings
highlight important aspects of impaired neural input to older muscle that should be
considered in the field of neuromuscular ageing.

The older participants in this cohort were healthy and living independently which may
partly explain the lack of a statistically significant (although an ES of 0.87) age-
related difference in muscle size, which is not consistent with the wider ageing
literature (Piaseck*i et al.*, 2018). However, the greater difference in muscle strength
is consistent with numerous previous studies (e.g. Maden-Wilkinson et al. 2013;
Piasecki et al. 2016) and highlights explanatory factors of strength loss beyond total
muscle size. Alongside the lower strength and MU discharge rate, reduced force
control ability in older participants was also observed during the descending phase of
a ramp contraction. Numerous factors associated with advancing age may plausibly
explain this poorer force control, including MU remodelling, reduced excitability of the
MUs, decreased muscle spindle sensitivity, less common synaptic inputs, and/or
impaired cutaneous afferents (Mynark & Koceja, 2001; Shaffer & Harrison, 2007).

Estimates of MU number in the VL are reduced in older age (Piasecki *et al.*, 2016*b*,
2019), and although direct human evidence is lacking, animal models show large
MUs are preferentially lost with ageing and denervated fibres are reinnervated by
early recruited MUs (Kadhiresan *et al.*, 1996), resulting in expansion of those earlier
recruited MUs such as those sampled in the current study (<20% MVC). These MUs
with a larger number of fibres, when activated, would have a greater influence on
force during their progressive re/derecruitment processes.

The poorer force control ability in the older group could partially be a result of
neuronal atrophy. Several studies have reported a decrease of grey matter and white
matter throughout the motor cortex with increasing age (Marner *et al.*, 2003; Salat *et
al.*, 2004; McGinnis *et al.*, 2011) and a strong correlation between cortical atrophy
and fine control capacity (Kennedy & Raz, 2005). A significant decrease in inhibition
has also been observed in aged muscles, as well as a greater level of cortical and
subcortical area activity during a targeted motor control task (Shinohar*a et al.*, 2003;
Marneweck *et al.*, 2011; Plow *et al.*, 2013). In the absence of the ability to inhibit
appropriate cortical areas during targeted motor tasks, MUs of older people may be
activated in an inappropriate manner relative to the task.

Proprioception is an essential aspect of functional muscle contraction, and again,
this is impaired in older age which can negatively impact balance (Ferlinc *et al.*,
2019). Both animal and human studies have revealed a decrease proprioception in
aged muscles with the evidence of declined muscle spindle numbers and
degeneration of the sensory nerve terminals, leading to the morphological
adaptations at a peripheral level and modulation of mechanoreceptor gain at a
central level (Winarakwong *et al.*, 2004; Macefield & Knellwolf, 2018). Through
inhibitory interneurons, inhibitory spinal circuits can affect descending inputs as well
as Ia afferents from muscle spindles to spinal motoneurons (Kido *et al.*, 2004). Older
participants exhibited a greater loss of Ia afferent feedback (Scaglion*i et al.*, 2003)
and an impaired ability to modulate Ia presynaptic inhibition (Baudry *et al.*, 2010).
Thus, compared to younger individuals, it is probable the older participants were less
able to sense force rate changes during the contraction.

Although not directly assessed, the points made thus far are supported by the
intramuscular coherence findings during ascending and descending phases.
Coherence in alpha band (5-12 Hz) is associated with Ia afferent feedback
(Mehrkanoon *et al.*, 2014) and when exploring the differences between ascending
and descending phases, a significant lower coherence estimate in alpha band was
observed in old only, corresponding to the poorer control ability. Beta band (15-30
Hz) differences between age groups in both ascending and descending phases may
reflect the reduced descending inputs from the motor cortex in older group (Walker
et al., 2019), suggesting that corticospinal tract function was compromised and there
may be neural remodelling process occurring for performance regulation in older
participants (Dodd *et al.*, 2017). Finally, age-related deterioration of cutaneous
afferent inputs may also contribute to reduced force control ability, which is reduced
in older adults (Peters *et al.*, 2016). Therefore, a decreased ability to regulate the
force may also be a result of the combination of reduced Ia afferent feedback from
muscle spindles and reduced cutaneous afferents.

MUs in older participants showed lower recruitment and derecruitment thresholds;
put simply, MUs in older individuals were activated at a lower force level when
compared to the younger participants and remained activated for longer. This may
be explained by the loss of higher-threshold MUs in older participants, leading to a
larger proportion of lower-threshold MUs in the MU pool (Piasecki et al. 2016). It is

widely accepted that in smaller muscles such as those in the hand, all MUs are
recruited by 50% MVC with forces above that level achieved via increases in
discharge rates (De Luca *et al.*, 1982). In larger muscles, recruitment continues up
until approximately 90% MVC and MU discharge rate facilitates force increases
beyond that (Masakado, 1994). However, MUs undergo remodelling during ageing,
and it is possible that the force regulation strategy is altered with a preferential
recruitment of MUs at lower relative contraction levels to compensate.

Compared to younger participants, we report a significantly reduced delta F in the VL
of older participants during a ramp contraction. Supportive of studies of other
muscles (Hassan *et al.*, 2021; Orssatto *et al.*, 2021), the current findings in VL
showed that the intrinsic synaptic excitation was lower in old, which is likely
attributable to the reduced activity of the monoaminergic system. Serotonin (5-HT)
and norepinephrine (NE) appears primarily to be involved in the modulation of motor
control and sensory input, partly influencing the magnitude of PIC amplitude (Lee &
Heckman, 1998). Ageing is associated with decreased levels of 5-HT receptors and
transporters in human brain (Karrer *et al.*, 2019; Deza-Araujo *et al.*, 2021) as well as
impaired NE synthesis and secretion in older participants (Shibata *et al.*, 2006),
leading to a reduced monoaminergic neurotransmission in healthy ageing.
Additionally, the magnitude of PICs may also be influenced by age-related
alterations in Na^+ and L-type Ca^{2+} ion channels, widely distributed on the dendrites
of motoneurons. The deregulation of Ca^{2+} has been reported to have an essential
role in the development of ageing and the neurodegeneration process via C. elegans
models (Alvarez *et al.*, 2020), which may also be apparent in human spinal
motoneurons. A high sensitivity of PIC amplitude to inhibitory input to motoneurons
has been reported (Heckmann et al. 2005) and as discussed above, inhibitory inputs
significantly decrease with advancing age at both central nervous system and spinal
cord levels. The reduced PICs in older participants here may be a result of increased
inhibitory input levels. Furthermore, the sex-based differences of PIC magnitude
recently observed (Jenz *et al.*, 2023) may be explained by hormonal differences
which influences 5-HT, that may similarly differ between young and older males.

Strengths/Limitations

The main strength of this study was this is the first study investigating the common
synaptic inputs and intrinsic excitability properties in aged VL between young and
older humans. However, there are several limitations. Firstly, the results reported
here were from males only which may not directly translate to females, as decreases
in estrogen levels may also contribute to the alterations in circulating serotonin and
its receptor densities in postmenopausal women (Gonzales & Carrillo, 1993).
Secondly, we selected the best contraction among four contractions, which
potentially ignored the age-related variability among and within participants. In
addition, we used 10s signal length for calculating coherence estimates, possibly
resulting in an underestimated effect of ageing on common synaptic inputs and
intrinsic properties.

Conclusion

The reduced muscle strength and control ability observed in older males is partially
attributable to impaired motoneuron excitability and reduced common synaptic input
which occur independently of reductions in muscle size. These findings have
important implications in the field of healthy human ageing and should be considered
when applying interventions to mitigate age-related functional decrements.

Data Availability Statement

The datasets generated and analysed during the current study are available from the
corresponding author upon reasonable request.

Competing interests

The authors have no competing interests to declare.

References

Afsharipour B, Manzur N, Duchcherer J, Fenrich KF, Thompson CK, Negro F,
Quinlan KA, Bennett DJ & Gorassini MA (2020). Estimation of self-sustained
activity produced by persistent inward currents using firing rate profiles of

multiple motor units in humans. *J Neurophysiol* **124,** 63–85.

Alvarez J, Alvarez-Illera P, García-Casas P, Fonteriz RI & Montero M (2020). The
Role of Ca(2+) Signaling in Aging and Neurodegeneration: Insights from
Caenorhabditis elegans Models. *Cells*; DOI: 10.3390/cells9010204.

Avrillon S, Del Vecchio A, Farina D, Pons JL, Vogel C, Umehara J & Hug F (2020).
Individual differences in the neural strategies to control the lateral and medial
head of the quadriceps during a mechanically constrained task. *J Appl Physiol*
130, 269–281.

Baudry S, Maerz AH & Enoka RM (2010). Presynaptic modulation of Ia afferents in
young and old adults when performing force and position control. *J
Neurophysiol* **103,** 623–631.

Binder MD & Powers RK (2001). Relationship between simulated common synaptic
input and discharge synchrony in cat spinal motoneurons. *J Neurophysiol* **86,**
2266–2275.

Castronovo AM, Mrachacz-Kersting N, Stevenson AJT, Holobar A, Enoka RM &
Farina D (2018). Decrease in force steadiness with aging is associated with
increased power of the common but not independent input to motor neurons. *J
Neurophysiol* **120,** 1616–1624.

Daley MJ & Spinks WL (2000). Exercise, Mobility and Aging. *Sport Med* **29,** 1–12.

Deza-Araujo YI, Baez-Lugo S, Vuilleumier P, Chocat A, Chételat G, Poisnel G &
Klimecki OM (2021). Whole blood serotonin levels in healthy elderly are
negatively associated with the functional activity of emotion-related brain regions.
Biol Psychol **160,** 108051.

Dodd KC, Nair VA & Prabhakaran V (2017). Role of the Contralesional vs.
Ipsilesional Hemisphere in Stroke Recovery. *Front Hum Neurosci* **11,** 469.

Farina D & Negro F (2015). Common synaptic input to motor neurons, motor unit
synchronization, and force control. *Exerc Sport Sci Rev* **43,** 23–33.

Farina D, Negro F & Dideriksen JL (2014). The effective neural drive to muscles is
the common synaptic input to motor neurons. *J Physiol* **592,** 3427–3441.

Feeney DF, Mani D & Enoka RM (2018). Variability in common synaptic input to
motor neurons modulates both force steadiness and pegboard time in young
and older adults. *J Physiol* **596,** 3793–3806.

Ferlinc A, Fabiani E, Velnar T & Gradisnik L (2019). The Importance and Role of
Proprioception in the Elderly: a Short Review. *Mater Sociomed* **31,** 219–221.

Gonzales GF & Carrillo C (1993). Blood serotonin levels in postmenopausal women:
Effects of age and serum oestradiol levels. *Maturitas* **17,** 23–29.

Gorassini M, Yang JF, Siu M & Bennett DJ (2002). Intrinsic Activation of Human
Motoneurons: Possible Contribution to Motor Unit Excitation. *J Neurophysiol* **87,**
1850–1858.

Guo Y, Jones EJ, Inns TB, Ely IA, Stashuk DW, Wilkinson DJ, Smith K, Piasecki J,
Phillips BE, Atherton PJ & Piasecki M (2022*a*). Neuromuscular recruitment

strategies of the vastus lateralis according to sex. *Acta Physiol* **235,** e13803.

Guo Y, Piasecki J, Swiecicka A, Ireland A, Phillips BE, Atherton PJ, Stashuk D, Rutter MK, McPhee JS & Piasecki M (2022*b*). Circulating testosterone and dehydroepiandrosterone are associated with individual motor unit features in untrained and highly active older men. *GeroScience* **44,** 1215–1228.

Hassan A, Thompson CK, Negro F, Cummings M, Powers RK, Heckman CJ, Dewald JPA & McPherson LM (2020). Impact of parameter selection on estimates of motoneuron excitability using paired motor unit analysis. *J Neural Eng* **17,** 16063.

Hassan AS, Fajardo ME, Cummings M, McPherson LM, Negro F, Dewald JPA, Heckman CJ & Pearcey GEP (2021). Estimates of persistent inward currents are reduced in upper limb motor units of older adults. *J Physiol* **599,** 4865–4882.

Heckman CJ & Enoka RM (2012). Motor unit. *Compr Physiol* **2,** 2629–2682.

Heckman CJ, Hyngstrom AS & Johnson MD (2008*a*). Active properties of motoneurone dendrites: diffuse descending neuromodulation, focused local inhibition. *J Physiol* **586,** 1225–1231.

Heckman CJ, Johnson M, Mottram C & Schuster J (2008*b*). Persistent inward currents in spinal motoneurons and their influence on human motoneuron firing patterns. *Neurosci a Rev J bringing Neurobiol Neurol psychiatry* **14,** 264–275.

Heckmann CJ, Gorassini MA & Bennett DJ (2005). Persistent inward currents in motoneuron dendrites: implications for motor output. *Muscle Nerve* **31,** 135–156.

Hepple RT & Rice CL (2016). Innervation and neuromuscular control in ageing skeletal muscle. *J Physiol* **594,** 1965–1978.

Holobar A, Minetto MA & Farina D (2014). Accurate identification of motor unit discharge patterns from high-density surface EMG and validation with a novel signal-based performance metric. *J Neural Eng* **11,** 16008.

Holobar A & Zazula D (2007). Multichannel Blind Source Separation Using Convolution Kernel Compensation. *IEEE Trans Signal Process* **55,** 4487–4496.

Jenz ST, Beauchamp JA, Gomes MM, Negro F, Heckman CJ & Pearcey GEP (2023). The effects of biological sex on estimates of persistent inward currents in the human lower limb. *bioRxiv*2022.10.09.511486.

Kadhiresan VA, Hassett CA & Faulkner JA (1996). Properties of single motor units in medial gastrocnemius muscles of adult and old rats. *J Physiol* **493 (Pt 2,** 543–552.

Karrer TM, McLaughlin CL, Guaglianone CP & Samanez-Larkin GR (2019). Reduced serotonin receptors and transporters in normal aging adults: a meta-analysis of PET and SPECT imaging studies. *Neurobiol Aging* **80,** 1–10.

Kennedy KM & Raz N (2005). Age, Sex and Regional Brain Volumes Predict Perceptual-Motor Skill Acquisition. *Cortex* **41,** 560–569.

Kido A, Tanaka N & Stein RB (2004). Spinal excitation and inhibition decrease as

humans age. *Can J Physiol Pharmacol* **82,** 238–248.

Kirk EA, Christie AD, Knight CA & Rice CL (2021). Motor unit firing rates during
constant isometric contraction: establishing and comparing an age-related
pattern among muscles. *J Appl Physiol* **130,** 1903–1914.

Laine CM & Valero-Cuevas FJ (2017). Intermuscular coherence reflects functional
coordination. *J Neurophysiol* **118,** 1775–1783.

Lee RH & Heckman CJ (1998). Bistability in spinal motoneurons in vivo: systematic
variations in persistent inward currents. *J Neurophysiol* **80,** 583–593.

De Luca CJ, LeFever RS, McCue MP & Xenakis AP (1982). Behaviour of human
motor units in different muscles during linearly varying contractions. *J Physiol*
329, 113–128.

Macefield VG & Knellwolf TP (2018). Functional properties of human muscle
spindles. *J Neurophysiol* **120,** 452–467.

Maden-Wilkinson TM, Degens H, Jones DA & McPhee JS (2013). Comparison of
MRI and DXA to measure muscle size and age-related atrophy in thigh muscles.
J Musculoskelet Neuronal Interact **13,** 320–328.

Manini TM, Hong SL & Clark BC (2013). Aging and muscle: a neuron's perspective.
Curr Opin Clin Nutr Metab Care. Available at: https://journals.lww.com/co-
clinicalnutrition/Fulltext/2013/01000/Aging_and_muscle__a_neuron_s_perspecti
ve.5.aspx.

Marner L, Nyengaard JR, Tang Y & Pakkenberg B (2003). Marked loss of myelinated
nerve fibers in the human brain with age. *J Comp Neurol* **462,** 144–152.

Marneweck M, Loftus A & Hammond G (2011). Short-interval intracortical inhibition
and manual dexterity in healthy aging. *Neurosci Res* **70,** 408–414.

Masakado Y (1994). Motor unit firing behavior in man. *Keio J Med* **43,** 137–142.

McGinnis SM, Brickhouse M, Pascual B & Dickerson BC (2011). Age-related
changes in the thickness of cortical zones in humans. *Brain Topogr* **24,** 279–291.

Mehrkanoon S, Breakspear M & Boonstra TW (2014). The reorganization of
corticomuscular coherence during a transition between sensorimotor states.
Neuroimage **100,** 692–702.

Mynark RG & Koceja DM (2001). Effects of Age on the Spinal Stretch Reflex. *J Appl*
Biomech **17,** 188–203.

Nadim F & Bucher D (2014). Neuromodulation of neurons and synapses. *Curr Opin*
Neurobiol **29,** 48–56.

Orssatto LBR, Borg DN, Blazevich AJ, Sakugawa RL, Shield AJ & Trajano GS
(2021). Intrinsic motoneuron excitability is reduced in soleus and tibialis anterior
of older adults. *GeroScience* **43,** 2719–2735.

Orssatto LBR, Borg DN, Pendrith L, Blazevich AJ, Shield AJ & Trajano GS (2022).
Do motoneuron discharge rates slow with aging? A systematic review and meta-
analysis. *Mech Ageing Dev* **203,** 111647.

Peters RM, McKeown MD, Carpenter MG & Inglis JT (2016). Losing touch: age-
related changes in plantar skin sensitivity, lower limb cutaneous reflex strength,
and postural stability in older adults. *J Neurophysiol* **116,** 1848–1858.

Pethick J & Piasecki M (2022). Alterations in Muscle Force Control With Aging: Is
There a Modulatory Effect of Lifelong Physical Activity?   . *Front Sport Act*
Living . Available at:
https://www.frontiersin.org/articles/10.3389/fspor.2022.817770.

Piasecki M, Ireland A, Jones DA & McPhee JS (2016*a*). Age-dependent motor unit
remodelling in human limb muscles. *Biogerontology* **17,** 485–496.

Piasecki M, Ireland A, Piasecki J, Degens H, Stashuk DW, Swiecicka A, Rutter MK,
Jones DA & McPhee JS (2019). Long-Term Endurance and Power Training May
Facilitate Motor Unit Size Expansion to Compensate for Declining Motor Unit
Numbers in Older Age   . *Front Physiol* . Available at:
https://www.frontiersin.org/articles/10.3389/fphys.2019.00449.

Piasecki M, Ireland A, Piasecki J, Stashuk DW, Swiecicka A, Rutter MK, Jones DA &
McPhee JS (2018). Failure to expand the motor unit size to compensate for
declining motor unit numbers distinguishes sarcopenic from non-sarcopenic
older men. *J Physiol* **596,** 1627–1637.

Piasecki M, Ireland A, Stashuk D, Hamilton-Wright A, Jones DA & McPhee JS
(2016*b*). Age-related neuromuscular changes affecting human vastus lateralis. *J*
Physiol **594,** 4525–4536.

Plow EB, Cunningham DA, Bonnett C, Gohar D, Bayram M, Wyant A, Varnerin N,
Mamone B, Siemionow V, Hou J, Machado A & Yue GH (2013).
Neurophysiological correlates of aging-related muscle weakness. *J*
Neurophysiol **110,** 2563–2573.

Raethjen J, Govindan RB, Kopper F, Muthuraman M & Deuschl G (2007). Cortical
involvement in the generation of essential tremor. *J Neurophysiol* **97,** 3219–
3228.

Rossato J, Tucker K, Avrillon S, Lacourpaille L, Holobar A & Hug F (2022). Less
common synaptic input between muscles from the same group allows for more
flexible coordination strategies during a fatiguing task. *J Neurophysiol* **127,** 421–
433.

Salat DH, Buckner RL, Snyder AZ, Greve DN, Desikan RSR, Busa E, Morris JC,
Dale AM & Fischl B (2004). Thinning of the Cerebral Cortex in Aging. *Cereb*
Cortex **14,** 721–730.

Scaglioni G, Narici M V, Maffiuletti NA, Pensini M & Martin A (2003). Effect of ageing
on the electrical and mechanical properties of human soleus motor units
activated by the H reflex and M wave. *J Physiol* **548,** 649–661.

Shaffer SW & Harrison AL (2007). Aging of the Somatosensory System: A
Translational Perspective. *Phys Ther* **87,** 193–207.

Shibata E, Sasaki M, Tohyama K, Kanbara Y, Otsuka K, Ehara S & Sakai A (2006).
Age-related Changes in Locus Ceruleus on Neuromelanin Magnetic Resonance
Imaging at 3 Tesla. *Magn Reson Med Sci* **5,** 197–200.

Shinohara M, Keenan KG & Enoka RM (2003). Contralateral activity in a
homologous hand muscle during voluntary contractions is greater in old adults. *J*
Appl Physiol **94,** 966–974.

Smee DJ, Anson JM, Waddington GS & Berry HL (2012). Association between
Physical Functionality and Falls Risk in Community-Living Older Adults ed.
Lauretani F. *Curr Gerontol Geriatr Res* **2012,** 864516.

Udina E, D'Amico J, Bergquist AJ & Gorassini MA (2010). Amphetamine increases
persistent inward currents in human motoneurons estimated from paired motor-
unit activity. *J Neurophysiol* **103,** 1295–1303.

Vandenberk MS & Kalmar JM (2014). An evaluation of paired motor unit estimates of
persistent inward current in human motoneurons. *J Neurophysiol* **111,** 1877–
1884.

Walker S, Avela J, Wikgren J, Meeusen R, Piitulainen H, Baker SN & Parviainen TM
(2019). Aging and Strength Training Influence Knee Extensor Intermuscular
Coherence During Low- and High-Force Isometric Contractions . *Front*
Physiol . Available at:
https://www.frontiersin.org/articles/10.3389/fphys.2018.01933.

Wilkinson DJ, Piasecki M & Atherton PJ (2018). The age-related loss of skeletal
muscle mass and function: Measurement and physiology of muscle fibre
atrophy and muscle fibre loss in humans. *Ageing Res Rev* **47,** 123–132.

Winarakwong L, Muramoto T, Soma K & Takano Y (2004). Age-related changes and
the possible adaptability of rat jaw muscle spindles: immunohistochemical and
fine structural studies. *Arch Histol Cytol* **67,** 227–240.

660