Title: Common synaptic inputs and persistent inward currents of vastus

lateralis motor units are reduced in older age

Yuxiao Guo[1], Eleanor J. Jones[1], Jakob Škarabot[2], Thomas B. Inns[1], Bethan E.

Phillips[1], Philip J. Atherton[1], Mathew Piasecki[1]

1. Centre of Metabolism, Ageing & Physiology (COMAP), MRC-Versus Arthritis

Centre for Musculoskeletal Ageing Research & National Institute for Health

Research (NIHR) Nottingham Biomedical Research Centre, School of

Medicine, University of Nottingham, Derby, UK

2. School of Sport, Exercise and Health Sciences, Loughborough University,

Loughborough, UK

Running title: Age-related reductions in PICs and coherence

Corresponding author details:

Mathew Piasecki (Mathew.piasecki@nottingham.ac.uk)

Centre Of Metabolism, Ageing & Physiology (COMAP)

Academic Unit of Injury, Inflammation & Recovery Sciences

School of Medicine, Faculty of Medicine & Health Sciences

University of Nottingham, Royal Derby Hospital Centre (Room 3011)

Derby, DE22 3DT, UK

Tel: +44 01332 724676

Abstract

Although muscle atrophy may partially account for age-related strength decline, it is

further influenced by alterations of neural input to muscle. Persistent inward currents

and the level of common synaptic inputs to motoneurons influence neuromuscular

function. However, these have not yet been described in aged human quadriceps.

High density surface electromyography (HDsEMG) signals were collected from the

vastus lateralis of 15 young (mean±SD, 23 ± 5 y) and 15 older (67 ± 9 y) men during

submaximal sustained and 20-s ramped contractions. HDsEMG signals were

decomposed to identify individual motor unit discharges, from which delta F and

intramuscular coherence were estimated.

Older participants produced significantly lower knee extensor torque (p<0.001) and

poorer force tracking ability (p<0.001) than young. Older participants also had lower

delta F (p=0.001) and coherence estimates in the alpha frequency band (p<0.001)

during ramp contractions when compared to young.

Persistent inward currents and common synaptic inputs are lower in the vastus

lateralis of older males when compared to young. These data highlight altered neural

input to the clinically and functionally important quadriceps, further underpinning age-

related loss of function which may occur independently of the loss of muscle mass.

Key Points

- The age-related loss of muscle mass is exceeded by the loss of function, which is influenced by structural and functional alterations of the nervous system.
- Motoneuronal persistent inward currents and common synaptic inputs play an important role in the activation of motor units and subsequent force generation and control ability.
- Here we show reduced estimates of persistent inward currents and lower common synaptic inputs to older vastus lateralis, potentially contributing to observed lower strength and poorer force tracking.
- These findings highlight decrements of the aged human motor system, accompanied by muscle atrophy in functionally relevant muscle groups, which should be considered in the application of interventions targeting aged human muscle.

Introduction

Human ageing is characterised by a progressive reduction in muscle size and muscle strength, resulting in an impairment of force generation, force control and physical performance [1–3], all of which can negatively impact quality of life [4,5]. The loss of muscle size is explicable by the combined effects of muscle fibre atrophy and muscle fibre loss [2], and although atrophy may partially account for declines in muscle strength, it is further influenced by structural and functional alterations of the nervous system.

The voluntary contraction of muscle fibres occurs via net excitatory signals to the motor units (MUs); groups of muscle fibres innervated by a single motoneuron [6]. The central nervous system controls muscle force via two primary strategies; varying the number of MUs recruited and varying the discharge rate of each motor neuron, both of which are susceptible to age-related alterations. Substantial motoneuron loss accelerates in the sixth decade of life, resulting in fewer MUs, and a compensatory process of MU remodelling further results in larger MU innervation ratios [7,8]. Additionally, although this appears to be muscle specific, an age-related decrease in MU discharge rate has also been documented at normalised contraction levels [9,10]

as well as being associated with circulating hormone levels that are also affected by
age [11].

In addition to direct neurotransmitter activation onto postsynaptic ligand-gated
channels at the spinal cord level via ionotropic mechanisms, activation of the motor
nervous system also relies on neuromodulation [12]. This is mediated by intracellular
second messenger systems that modify the voltage- and ligand-gated channels of
neurons and alters several properties from their level of excitability to their pattern of
firing in response to a given input [13,14]. This process is enhanced by monoamines,
e.g. serotonin and norepinephrine released from the brainstem nuclei by stimulating
inward-flowing persistent calcium and sodium currents [12], known as persistent
inward currents (PICs). As long as the membrane potential remains above the
activation threshold, these PICs tend to remain activated [12], combined with
neuromodulatory drive amplifying the firing behaviours of motoneurons to synaptic
inputs up to five fold [15] as well as generating long-lasting plateau potentials which
results in self-sustaining firing [16]. Even though PICs appear to decrease with age
in upper [17] and lower limb muscles [18], PICs can be mediated by local ionotropic
inhibitory inputs that may be muscle specific in the response to age [10,19,20],
particularly those highly susceptible to functional losses such as the vastus lateralis
(VL) [21].

Rather than controlling the discharge pattern of individual MUs, the central nervous
system controls the excitatory inputs to the MU pool, referred to as common synaptic
inputs [22] which are defined as the proportion of net synaptic input correlated
between MUs. Force fluctuations are largely determined by the alterations in low
frequency components of common synaptic inputs to MUs, which is reflected in
concurrent fluctuations in MU discharge rates from the same MU pool [23].
Accordingly, a decrease in force steadiness associated with ageing is a
consequence of an increase in variance of common synaptic inputs [24,25]. However,
the variability of common synaptic inputs in quadriceps has only been explored in
healthy young individuals [26,27], with no available data from older humans.

It is not possible to make intracellular recordings from human spinal motoneurons in
order to estimate PICs, so alternative methods are employed which distinguish
intrinsic excitability of the MU from its descending synaptic drive by using the unit-
wise MU analysis technique [17]. In view of the advances with which MU discharging

patterns can be measured, it provides an opportunity to investigate and better understand the physical properties and the underlying mechanisms which may impair motor control in older age. Therefore, the aim of this study was to explore the differences in physical performance as well as the magnitude of persistent inward currents and common synaptic inputs in the vastus lateralis between healthy young and older males. We hypothesized that older participants would exhibit greater physical decrements and lower levels of persistent inward currents and common synaptic inputs.

Methods

Participants and ethical approval

This study was approved by the University of Nottingham Faculty of Medicine and Healthy Science Research Ethics Committee (90-0820, 199-0221) and conformed to the Declaration of Helsinki.

15 healthy young males between the ages of 18-40 (mean ± SD; age: 23.1 ± 5.2 years; body mass index: 25.0 ± 2.5 kg/m^2) and 15 healthy older males between the ages of 55 to 85 (67.2 ± 8.9 years; 26.1± 2.3 kg/m^2) provided informed consent to take part in the study. All participants were recruited through advertisements in the local community. Prior to enrolment, all participants completed a comprehensive clinical examination and metabolic screening was conducted at the School of Medicine, Royal Derby Hospital Centre. All recruited participants were recreationally active and would be excluded if they display evidence of: BMI < 18.5 or > 35kg/m^2; are competitive in an athletic discipline at county level or above; musculoskeletal disorders; respiratory disease; neurological disorders; metabolic disease; active cardiovascular problems; active inflammatory bowel or renal disease; recent steroid treatment within 6 months or hormone replacement therapy; family history of early (< 55years) death from cardiovascular disease.

Anthropometry

Ultrasound was used to measure the cross-sectional area (CSA) of the VL muscle of the right leg, using an ultrasound probe (LA523 probe, B-mode, frequency range 26–32Hz, and MyLabTM50 scanner, Esaote, Genoa, Italy) at the anatomical mid-point of the right thigh. Images were recorded with the participants laying in a supine position and the mid-point obtained from the identification of the greater trochanter at the top

of the femur and the mid-point of the patella. Panoramic imaging (VPAN) images were obtained in a medial to lateral fashion, beginning and ending the capture of the image at the aponeurosis borders of the VL of the right leg. The produced images were subsequently analysed using ImageJ software (National Institute of Health, USA) through tracing around the contour of the aponeurosis via polygon function to quantify the area of the VL. Each image was analysed three times and the mean value of three images was accepted as CSA.

Experimental Protocol

All participants were required to attend the lab at 0900 after an overnight fast and abstain from strenuous exercise 24h prior to the testing sessions. Participants were seated in a custom-built chair with hips and knees flexed at ~90 degree. The lower leg was secured to a force dynamometer with noncompliant straps above the medial malleolus. To minimize the movement of the upper trunk during testing, a seat belt was fastened across the pelvis of the participant. After a standardised warm up through several submaximal contractions [28], participants were instructed to contract as hard and fast as they could. During the trial, participants were not allowed to hold onto the side of the chair and were asked to cross their arms across the chest. This was performed with real-time visual force feedback on a monitor placed in front of the participant, with verbal encouragement to aid in producing maximal effort. This was repeated two to three times further, giving 60 seconds of rest between attempts standardised recovery. If the difference between the last two attempts was < 5%, the highest value recorded in newtons was taken and accepted as the maximal isometric voluntary contraction force (MVC). Muscle torque was calculated by multiplying MVC by the lever arm length from the knee to the centre of the ankle strap.

Following a 60-second rest, participants were then asked to perform four submaximal voluntary isometric contractions at 25% of MVC (~12s) and four triangular shaped contractions (10s up and 10s down) peaking at 20% of MVC, with 30-second intervals between contractions. In each case, a target line was displayed on a screen and participants were instructed to follow the target as close as possible. Ramp contractions to 20% of MVC have been widely used for estimating persistent inward currents (details below) [16,29,30]. To quantify force steadiness during 25% of MVC, the coefficient of variation of the force (CoV) was calculated =

187 (SD/Mean)*100. To assess force tracking accuracy, all data were exported to Spike2

188 (version 9.11, CED Ltd., Cambridge, UK), where a virtual channel was created by

189 subtracting the performed path from the requested (target) path and rectifying it.

190 Area under the curve reflects the level of deviation from the target line reflecting

191 muscle force tracking accuracy, with higher values indicating greater deviation.

192 High-density surface electromyography

A semi-disposable high-density surface electromyography (HDsEMG) array (64

electrodes, 13×5, 8mm, I.E.D., GR08MM1305, OT Bioelettronica, Inc., Turin, Italy)

was placed with the orientation of the muscle fibres (proximal to distal) over the

muscle belly of right vastus lateralis after skin preparation involving shaving, light

abrasion and cleansing with 70% ethanol. Electrodes were secured to the skin using

flexible tape. The adhesive grids were attached to the surface of the muscle by

disposable bi-adhesive foam layers (SpesMedica, Bettipaglia, Italy). The skin

electrode contact was facilitated by filling the cavities of the adhesive layers with

conductive paste (AC Cream, SpesMedica). A strap ground electrode (WS2,

OTBioelettronica, Turin, Italy) dampened with water was positioned around the ankle

of the right leg to optimise the signal quality. HDsEMG signals were acquired in a

monopolar configuration, amplified (x256) with filtering set at 10-500 Hz and digitally

converted at 2000 Hz by a 16-bit wireless amplifier (Sessantaquattro,

OTBioelettronica, Turin, Italy) and transferred to a PC for further offline analysis.

HDsEMG signals were recorded in OTBioLab software (OT Bioelecttronica, Turin,

Italy).

The recorded HDsEMG signals were converted into a MatLab file comprising a

single contraction. Each participant performed four submaximal contractions and

ramp contractions; however, a single trial for each contraction protocol was selected

for analysis based on the on the smoothness of the force profile. Monopolar

HDsEMG signals were band-pass filtered at 20-500Hz and then decomposed offline

into MU pulse trains by a convolutive kernel compensation algorithm [31,32]. After

the decomposition, a trained investigator manually inspected the MU spike trains and

edited the discharge patterns of each MU detected. Only MUs with a pulse-to-noise

ratio equal or greater than 30 dB were kept for further analysis.

Mean discharge rate during the submaximal contraction at 25% of MVC was
obtained and discharge rate variability was reported as the coefficient of variation
(CoV) of the interspike interval (ISI) displayed as a percentage. Peak discharge rates
during a ramp contraction were measured as the highest value of the smoothed MU
discharge rate with a fifth order polynomial.

Estimation of persistent inward currents (PICs)

The instantaneous firing rates of both MUs were calculated as the inverse of the
interspike intervals of each MU spike train and smoothed by fitting a fifth order
polynomial function. An estimate of the PIC magnitude was derived using the unit-
wise MU analysis technique [17]. The lower threshold unit to be recruited in the ramp
contraction is commonly referred to as the "control" unit; the "test" unit is a unit of
higher threshold. A sufficient number of MUs were isolated for 10 young and 10 older
participants. Delta F, the contribution of persistent inward currents to self-sustained
firing, was calculated as the difference in "control" unit firing rate between the onset
and offset of a "test" unit [16]. This technique relies on several assumptions: 1) test
and control units share the common synaptic drive; 2) PIC is activated before or at
the recruitment; 3) firing rate of the control MU closely reflects the depolarizing input
to the parent motoneuron; 4) test and control unit process the synaptic inputs in a
similar way. The MU pairs for the delta F calculation were selected based on the
following criteria: 1) as a measure of common synaptic modulation, rate-to-rate
correlation r ≥ 0.7; 2) to avoid the high variability in delta F calculation due to the
initial acceleration phase of MU firings, test units were recruited at least 1s after the
control units; 3) to account for the possibility of control unit saturation leading to an
underestimation of delta F, pairs in which rate modulation of the control unit fell
within 0.5pps were removed from analysis [16,30,33,34]. An average delta F was
calculated when individual test units were paired with multiple control units (Figure 1).

Figure 1. Example data showing delta frequency (ΔF) calculation in vastus lateralis for paired-motor unit technique. Paired motor unit data from a young male is displayed in the left panels (mean ΔF=3.09) and from an older male in the right panels (mean ΔF=1.52). Test units are displayed in panel A and corresponding control units are in B, C and D.

Coherence Estimates

To estimate the level of common synaptic input, intramuscular coherence was calculated separately during sustained and ramp contractions (Figure 2), and represents a frequency domain correlation between cumulative spike trains (CSTs) of the identified motor units. The magnitude-squared coherence was calculated using the Welch's averaged periodogram with 50% overlapping Hann windows of 1s at the different frequency bands: delta (0-5 Hz), alpha (5-12 Hz), beta (15-30 Hz) and piper (40-50 Hz) [35,36]. This analysis was repeated for 60 randomly chosen combinations of two equally sized CSTs calculated as the sum of firing times from three MUs randomly selected from the same set of MUs and then averaged. The

261 Fisher Z-transform was applied to the coherence estimates to obtain a normally
262 distributed variable for comparisons.

Figure 2. The group mean intramuscular coherence Z-Scores during submaximal
contraction at 25% of maximal isometric voluntary contraction (MVC) and ramp
contraction at 20% of MVC in young (blue) and older (yellow) participants. The
magnitude-squared coherence was calculated from all the combinations (a maximum
of 60 random permutations) of two equally sized cumulative spike trains, each
composed of three motor units.

Statistical analysis

Data management and analysis were performed using RStudio (Version 2022.07.1).
Descriptive data were generated for all variables. *An independent t-test* was
conducted to test whether any differences existed between young and older males in
muscle cross-sectional area, torque, CoV-force, force tracking accuracy and
coherence estimates. In order to preserve variability within and across participants
simultaneously to the greatest extent, *multilevel linear regression models* were
generated to compare mean discharge rate, discharge rate variability, peak
discharge rate, delta F, recruitment threshold and derecruitment threshold between
groups using *lme4 package* (Version 1.1-27.1) [37]. In the multilevel models, single

MU was regarded as the first level; and individual participant with clustered MUs was
considered as the second level. For data visualization, individual participant means
are displayed in box-and-jitter plots. A *p* value <0.05 was considered statistically
significant.

Results

Young males had a larger muscle cross-sectional area (Y v O: 30.59 ± 5.87 v 23.34
± 7.03 cm^2, p=0.005; Figure 3A) and greater muscle torque (241.2 ± 62.69 v 143.3 ±
41.50 Nm, p<0.0001; Figure 3B) than older males. There was no difference in force
steadiness (CoV-Force) (2.18 ± 0.73 v 2.27 ± 0.65, p=0.735; Figure 3C) during
sustained contractions. However, there was a significant difference between groups
in force tracking accuracy of the ramp contraction, with the younger group
performing better than the old (9.41 ± 3.25 v 19.47 ± 8.36 Ns, p<0.001; Figure 3D).

*Figure 3. Cross-sectional area (A), torque (B), coefficient of variance of force at 25% maximal voluntary contraction (C) and force tracking accuracy (area under curve) during a single ramp contraction at 20% of maximal voluntary contraction (D) between young (blue) and older (yellow) males. Lines indicate group means and SD. ***p<0.001, **p<0.010, *p<0.05.*

During the submaximal contraction at 25% of MVC, a total of 246 MUs were identified from young males and 282 MUs from older males. There was no statistical difference in mean discharge rates (8.50 ± 2.08 v 7.98 ± 1.64, p=0.137, Figure 4A) or discharge rate variability (11.34 ± 6.01 v 10.22 ± 5.15, p=0.072, Figure 4B) between groups.

During the ramp contractions peaking at 20% of MVC, a total of 256 MUs were identified from young males and 262 MUs from older males. There was no statistical

difference in peak discharge rates (9.63 ± 2.20 v 9.35 ± 1.94 pps, p=0.154; Figure

4C). For Delta F calculations, there was an average of 5 (SD: 3.68) pairs of test and

control units per person in the young and 6 (4.95) pairs in older group, with the mean

number of 4 (1.79) test units per person in young and 3 (1.97) in old. Younger males

had significantly greater delta F values when compared to older participants (3.11 ±

0.77 v 1.86 ±0.65 pps, p=0.006; Figure 4D). There was no significant difference in

recruitment (10.60 ± 3.37 v 10.75 ± 3.55 %, p=0.604; Figure 4E) or derecruitment

thresholds (10.14 ± 3.40 v 10.18 ± 3.39 %, p=0.866; Figure 4F) of the identified MUs

between young and older groups.

Figure 4. Jitter plots of the individual participant mean values and box and raincloud plots of mean discharge rate (A), discharge rate variability (B), peak discharge rate (C), Delta F (D) and recruitment/decruitment threshold (E&F) in young (blue) and older (yellow) males.

In the sustained contraction, there was no significant age-related difference in coherence in Delta (0.49 ± 0.14 vs 0.54 ± 0.11, p=0.315), Alpha (0.24 ± 0.05 vs 0.23 ± 0.03, p=0.462) or Beta (0.22 ± 0.03 vs 0.21 ± 0.02, p=0.125) bands, however the

 young had higher coherence in the Piper band (0.23 ± 0.03 vs 0.21 ± 0.02, p=0.026)

(Figure 5A-D). In the ramped contraction, there were no age-related differences in

COH Delta (0.87 ± 0.08 vs 0.92 ± 0.12, p=0.182), Beta (0.19 ± 0.02 vs 0.18 ± 0.02,

p=0.110) or Piper (0.20 ± 0.04 vs 0.18 ± 0.02, p=0.156) bands however, young

participants had greater coherence estimates in Alpha band (0.21 ± 0.02 vs 0.18 ±

0.01, p<0.001) (Figure 5E-H).

Figure 5. Individual participant means of intramuscular coherence Z-score at
different frequency bands during a submaximal contraction at 25% maximal
voluntary contraction (MVC) and a ramp contraction at 20% MVC in young (blue)
and older (yellow) males.

Discussion

Here we provide the first evidence for reduced estimates of persistent inward

currents combined with reduced common synaptic inputs alongside poorer force

control ability in the aged human VL, a muscle with established susceptibility for age-

related functional losses. Though no significant age-related difference in MU

discharge rates was observed, our findings highlight important aspects of impaired

neural input to older muscle that likely contribute to functional decrements.

A smaller muscle size and a lower muscle strength were observed in older

participants, which is consistent with the wider ageing literature [38,39]. However, no

age-related difference in MU discharge rates was detected between groups in both

submaximal sustained contraction or ramp contraction, which is not aligned with our
previous findings in this muscle using needle EMG recordings [39]. This may reflect
differences in the population of MUs sampled, with needle EMG sampling from a
greater depth of muscle than surface-based HD EMG used here [40].

Age-related differences in force control ability were only observed during the ramp
contraction, highlighting the role of task complexity in age-related impairments [41].
Numerous factors associated with advancing age may also explain this poorer force
control in a more complex contraction, including MU remodelling, reduced excitability
of the MUs, decreased muscle spindle sensitivity, less common synaptic inputs,
and/or impaired cutaneous afferents [42,43]. Estimates of MU number in the VL are
reduced in older age [39,44], and although direct human evidence is lacking, animal
models show large MUs are preferentially lost with ageing and denervated fibres are
reinnervated by early recruited MUs [45], resulting in expansion of those earlier
recruited MUs [46] such as those sampled in the current study (< 20% MVC). These
MUs with a larger number of fibres, when activated, would have a greater influence
on force during their progressive re/derecruitment processes, and may partly explain
the age-related differences of force control observed here.

The progressive increase and decrease in force during the targeted isometric
ramped contraction used here is partly reliant on afferent neuromodulatory feedback
from the site of force application [47]. Several studies have reported a decrease of
grey matter and white matter throughout the motor cortex with increasing age [48–50]
and a strong correlation between cortical atrophy and fine control capacity [51]. A
significant decrease in inhibition has also been observed in aged muscles, as well as
a greater level of cortical and subcortical area activity during a targeted motor control
task [52–54], all of which may exert greater detrimental effects during a complex task
than simple (e.g. ramped vs sustained). Additionally, both animal and human studies
have revealed a decreased proprioception in aged muscles [55] with the evidence of
declined muscle spindle numbers and degeneration of the sensory nerve terminals,
leading to the morphological adaptations at a peripheral level and modulation of
mechanoreceptor gain at a central level [56,57]. Decreased inhibitory spinal circuits
with ageing has also been shown to have influence on Ia afferents from muscle
spindles to spinal motoneurons [58].

Although not directly assessed, the points made thus far are supported by the
intramuscular coherence findings during the ramp contraction. Coherence in alpha
band (5-12 Hz) is associated with Ia afferent feedback [59] and when exploring the
age-related differences, a significant lower coherence estimate in alpha band was
observed in old only during the ramp contraction, corresponding to the poorer force
tracking ability. This result supports evidence from previous studies, showing a
greater loss of Ia afferent feedback [60] and an impaired ability to modulate Ia
presynaptic inhibition [61] in older participants. Moreover, it has also been suggested
that age-related deterioration of cutaneous afferent inputs may also contribute to
reduced force control ability [62]. Therefore, a decreased ability to regulate the
progressive increase and decrease in force may also be a result of the combination
of reduced Ia afferent feedback from muscle spindles and reduced cutaneous
afferents.

The effect of ageing on MU recruitment thresholds is inconclusive, or more
accurately, it appears to be muscle specific. In a single study of the biceps and
triceps, MUs were recruited earlier in older triceps, but not the biceps [17]. This study
of VL found no age-related difference in mean MU recruitment or derecruitment
threshold, which was approximately 10% of MVC in both young and older
participants during contractions up to 20% of MVC; indicating age was probably not
the main predictor of MU recruitment/derecruitment threshold of VL.

Compared to younger participants, we report a significantly reduced delta F in the VL
of older participants during a ramp contraction, supportive of studies of other
muscles in the upper [17] and lower [18] limbs. The current findings in VL showed
that the estimates of PICs were lower in old, which is likely attributable to the
reduced activity of the monoaminergic system. Serotonin (5-HT) and norepinephrine
(NE) appears primarily to be involved in the modulation of motor control and sensory
input, partly influencing the magnitude of PICs [63]. Ageing is associated with
decreased levels of 5-HT receptors and transporters in the brain [64,65] as well as
impaired NE synthesis and secretion [66], leading to a reduced monoaminergic
neurotransmission. Additionally, the magnitude of PICs may also be influenced by
age-related alterations in Na^+ and L-type Ca^{2+} ion channels, widely distributed on the
dendrites of motoneurons. The dysregulation of Ca^{2+} has been reported to have an
essential role in the development of ageing and the neurodegeneration process via

C. elegans models [67], which may also be apparent in human spinal motoneurons.
A high sensitivity of PICs to inhibitory input to motoneurons has been reported [12]
and as discussed above, inhibitory inputs significantly decrease with advancing age
at the brain and spinal cord levels.

Strengths/Limitations

To our knowledge, this is the first study concurrently investigating the upper motor
neuron command (common synaptic inputs) and lower motor neuron excitability
(persistent inward currents) in young and older VL. Age-related impairments are
evident in each. However, there are several limitations. Firstly, the results reported
here were from males only which may not directly translate to females, as decreases
in estrogen levels may also contribute to the alterations in circulating serotonin and
its receptor densities in postmenopausal women [68]. Secondly, the MU data we
report are relevant to those recruited at lower contraction levels (< 25%) and reveal
little of those later recruited MUs. In addition, we used 12s and 20s signal length for
calculating coherence estimates during submaximal sustained contraction and ramp
contraction respectively, possibly resulting in an underestimated effect of ageing on
common synaptic inputs.

Conclusion

The reduced muscle strength and control ability observed in older males is partially
attributable to reduced estimates of persistent inward currents and common synaptic
input, which occur independently of reductions in MU discharge rates. These findings
have important implications in the field of healthy human ageing and should be
considered when applying interventions to mitigate age-related functional
decrements.

Data Availability Statement

The datasets generated and analysed during the current study are available from the
corresponding author upon reasonable request.

Competing interests

The authors have no competing interests to declare.

References

1. Manini TM, Hong SL, Clark BC. Aging and muscle: a neuron's perspective.
Curr. Opin. Clin. Nutr. Metab. Care [Internet] 2013;16. Available from:
https://journals.lww.com/co-
clinicalnutrition/Fulltext/2013/01000/Aging_and_muscle__a_neuron_s_perspec
tive.5.aspx

2. Wilkinson DJ, Piasecki M, Atherton PJ. The age-related loss of skeletal muscle
mass and function: Measurement and physiology of muscle fibre atrophy and
muscle fibre loss in humans. Ageing Res. Rev. 2018;47:123–32.

3. Pethick J, Piasecki M. Alterations in Muscle Force Control With Aging: Is There
a Modulatory Effect of Lifelong Physical Activity? [Internet]. Front. Sport. Act.
Living 2022;4. Available from:
https://www.frontiersin.org/articles/10.3389/fspor.2022.817770

4. Daley MJ, Spinks WL. Exercise, Mobility and Aging. Sport. Med. [Internet]
2000;29:1–12. Available from: https://doi.org/10.2165/00007256-200029010-
00001

5. Smee DJ, Anson JM, Waddington GS, Berry HL. Association between
Physical Functionality and Falls Risk in Community-Living Older Adults. Curr.
Gerontol. Geriatr. Res. [Internet] 2012;2012:864516. Available from:
https://doi.org/10.1155/2012/864516

6. Heckman CJ, Enoka RM. Motor unit. Compr. Physiol. 2012;2:2629–82.

7. Hepple RT, Rice CL. Innervation and neuromuscular control in ageing skeletal
muscle. J. Physiol. 2016;594:1965–78.

8. Piasecki M, Ireland A, Stashuk D, Hamilton-Wright A, Jones DA, McPhee JS.
Age-related neuromuscular changes affecting human vastus lateralis. J.
Physiol. 2016;594:4525–36.

9. Kirk EA, Christie AD, Knight CA, Rice CL. Motor unit firing rates during
constant isometric contraction: establishing and comparing an age-related
pattern among muscles. J. Appl. Physiol. [Internet] 2021;130:1903–14.
Available from: https://doi.org/10.1152/japplphysiol.01047.2020

10. Orssatto LBR, Borg DN, Pendrith L, Blazevich AJ, Shield AJ, Trajano GS. Do
motoneuron discharge rates slow with aging? A systematic review and meta-
analysis. Mech. Ageing Dev. [Internet] 2022;203:111647. Available from:
https://www.sciencedirect.com/science/article/pii/S004763742200029X

11. Guo Y, Piasecki J, Swiecicka A, Ireland A, Phillips BE, Atherton PJ, et al.
Circulating testosterone and dehydroepiandrosterone are associated with
individual motor unit features in untrained and highly active older men.

GeroScience 2022;44:1215–28.

12. Heckmann CJ, Gorassini MA, Bennett DJ. Persistent inward currents in
motoneuron dendrites: implications for motor output. Muscle Nerve
2005;31:135–56.

13. Heckman CJ, Johnson M, Mottram C, Schuster J. Persistent inward currents in
spinal motoneurons and their influence on human motoneuron firing patterns.
Neurosci. a Rev. J. bringing Neurobiol. Neurol. psychiatry 2008;14:264–75.

14. Nadim F, Bucher D. Neuromodulation of neurons and synapses. Curr. Opin.
Neurobiol. 2014;29:48–56.

15. Lee RH, Heckman CJ. Adjustable amplification of synaptic input in the
dendrites of spinal motoneurons in vivo. J. Neurosci. Off. J. Soc. Neurosci.
2000;20:6734–40.

16. Gorassini M, Yang JF, Siu M, Bennett DJ. Intrinsic Activation of Human
Motoneurons: Possible Contribution to Motor Unit Excitation. J. Neurophysiol.
[Internet] 2002;87:1850–8. Available from:
https://doi.org/10.1152/jn.00024.2001

17. Hassan AS, Fajardo ME, Cummings M, McPherson LM, Negro F, Dewald JPA,
et al. Estimates of persistent inward currents are reduced in upper limb motor
units of older adults. J. Physiol. 2021;599:4865–82.

18. Orssatto LBR, Borg DN, Blazevich AJ, Sakugawa RL, Shield AJ, Trajano GS.
Intrinsic motoneuron excitability is reduced in soleus and tibialis anterior of
older adults. GeroScience 2021;43:2719–35.

19. Yavuz UŞ, Negro F, Diedrichs R, Farina D. Reciprocal inhibition between
motor neurons of the tibialis anterior and triceps surae in humans. J.
Neurophysiol. 2018;119:1699–706.

20. Pearcey GEP, Khurram OU, Beauchamp JA, Negro F, Heckman CJ.
Antagonist tendon vibration dampens estimates of persistent inward currents
in motor units of the human lower limb. bioRxiv [Internet] 2022;Available from:
https://www.biorxiv.org/content/early/2022/08/04/2022.08.02.502526

21. Naruse M, Trappe S, Trappe TA. Human skeletal muscle-specific atrophy with
aging: a comprehensive review. J. Appl. Physiol. 2023;134:900–14.

22. Farina D, Negro F, Dideriksen JL. The effective neural drive to muscles is the
common synaptic input to motor neurons. J. Physiol. 2014;592:3427–41.

23. Farina D, Negro F. Common synaptic input to motor neurons, motor unit
synchronization, and force control. Exerc. Sport Sci. Rev. 2015;43:23–33.

24. Feeney DF, Mani D, Enoka RM. Variability in common synaptic input to motor
neurons modulates both force steadiness and pegboard time in young and
older adults. J. Physiol. 2018;596:3793–806.

25. Castronovo AM, Mrachacz-Kersting N, Stevenson AJT, Holobar A, Enoka RM,
Farina D. Decrease in force steadiness with aging is associated with increased
power of the common but not independent input to motor neurons. J.
Neurophysiol. [Internet] 2018;120:1616–24. Available from:

https://doi.org/10.1152/jn.00093.2018

26.   Avrillon S, Del Vecchio A, Farina D, Pons JL, Vogel C, Umehara J, et al.
Individual differences in the neural strategies to control the lateral and medial
head of the quadriceps during a mechanically constrained task. J. Appl.
Physiol. [Internet] 2020;130:269–81. Available from:
https://doi.org/10.1152/japplphysiol.00653.2020

27.   Rossato J, Tucker K, Avrillon S, Lacourpaille L, Holobar A, Hug F. Less
common synaptic input between muscles from the same group allows for more
flexible coordination strategies during a fatiguing task. J. Neurophysiol.
2022;127:421–33.

28.   Guo Y, Jones EJ, Inns TB, Ely IA, Stashuk DW, Wilkinson DJ, et al.
Neuromuscular recruitment strategies of the vastus lateralis according to sex.
Acta Physiol. [Internet] 2022;235:e13803. Available from:
https://doi.org/10.1111/apha.13803

29.   Afsharipour B, Manzur N, Duchcherer J, Fenrich KF, Thompson CK, Negro F,
et al. Estimation of self-sustained activity produced by persistent inward
currents using firing rate profiles of multiple motor units in humans. J.
Neurophysiol. 2020;124:63–85.

30.   Hassan A, Thompson CK, Negro F, Cummings M, Powers RK, Heckman CJ,
et al. Impact of parameter selection on estimates of motoneuron excitability
using paired motor unit analysis. J. Neural Eng. 2020;17:16063.

31.   Holobar A, Zazula D. Multichannel Blind Source Separation Using Convolution
Kernel Compensation. IEEE Trans. Signal Process. 2007;55:4487–96.

32.   Holobar A, Minetto MA, Farina D. Accurate identification of motor unit
discharge patterns from high-density surface EMG and validation with a novel
signal-based performance metric. J. Neural Eng. 2014;11:16008.

33.   Udina E, D'Amico J, Bergquist AJ, Gorassini MA. Amphetamine increases
persistent inward currents in human motoneurons estimated from paired
motor-unit activity. J. Neurophysiol. 2010;103:1295–303.

34.   Vandenberk MS, Kalmar JM. An evaluation of paired motor unit estimates of
persistent inward current in human motoneurons. J. Neurophysiol.
2014;111:1877–84.

35.   Raethjen J, Govindan RB, Kopper F, Muthuraman M, Deuschl G. Cortical
involvement in the generation of essential tremor. J. Neurophysiol.
2007;97:3219–28.

36.   Laine CM, Valero-Cuevas FJ. Intermuscular coherence reflects functional
coordination. J. Neurophysiol. [Internet] 2017;118:1775–83. Available from:
https://doi.org/10.1152/jn.00204.2017

37.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models
Using lme4. J. Stat. Softw. [Internet] 2015;67:1–48. Available from:
https://www.jstatsoft.org/index.php/jss/article/view/v067i01

38.   Maden-Wilkinson TM, Degens H, Jones DA, McPhee JS. Comparison of MRI
and DXA to measure muscle size and age-related atrophy in thigh muscles. J.

Musculoskelet. Neuronal Interact. 2013;13:320–8.

39. Piasecki M, Ireland A, Stashuk D, Hamilton-Wright A, Jones DA, McPhee JS.
Age-related neuromuscular changes affecting human vastus lateralis. J.
Physiol. 2016;594:4525–36.

40. Jones EJ, Piasecki J, Ireland A, Stashuk DW, Atherton PJ, Phillips BE, et al.
Lifelong exercise is associated with more homogeneous motor unit potential
features across deep and superficial areas of vastus lateralis. GeroScience
[Internet] 2021;43:1555–65. Available from: https://doi.org/10.1007/s11357-
021-00356-8

41. Bootsma JM, Caljouw SR, Veldman MP, Maurits NM, Rothwell JC, Hortobágyi
T. Neural Correlates of Motor Skill Learning Are Dependent on Both Age and
Task Difficulty   [Internet]. Front. Aging Neurosci. 2021;13. Available from:
https://www.frontiersin.org/articles/10.3389/fnagi.2021.643132

42. Mynark RG, Koceja DM. Effects of Age on the Spinal Stretch Reflex. J. Appl.
Biomech. 2001;17:188–203.

43. Shaffer SW, Harrison AL. Aging of the Somatosensory System: A Translational
Perspective. Phys. Ther. [Internet] 2007;87:193–207. Available from:
https://doi.org/10.2522/ptj.20060083

44. Piasecki M, Ireland A, Piasecki J, Degens H, Stashuk DW, Swiecicka A, et al.
Long-Term Endurance and Power Training May Facilitate Motor Unit Size
Expansion to Compensate for Declining Motor Unit Numbers in Older Age
[Internet]. Front. Physiol. 2019;10. Available from:
https://www.frontiersin.org/articles/10.3389/fphys.2019.00449

45. Kadhiresan VA, Hassett CA, Faulkner JA. Properties of single motor units in
medial gastrocnemius muscles of adult and old rats. J. Physiol. 1996;493 (Pt
2:543–52.

46. Jones EJ, Chiou SY, Atherton PJ, Phillips BE, Piasecki M. Ageing and
exercise-induced motor unit remodelling. J. Physiol. [Internet] 2022;600:1839–
49. Available from: https://doi.org/10.1113/JP281726

47. Koch SC, Acton D, Goulding M. Spinal Circuits for Touch, Pain, and Itch. Annu.
Rev. Physiol. [Internet] 2018;80:189–217. Available from:
https://doi.org/10.1146/annurev-physiol-022516-034303

48. Salat DH, Buckner RL, Snyder AZ, Greve DN, Desikan RSR, Busa E, et al.
Thinning of the Cerebral Cortex in Aging. Cereb. Cortex [Internet]
2004;14:721–30. Available from: https://doi.org/10.1093/cercor/bhh032

49. Marner L, Nyengaard JR, Tang Y, Pakkenberg B. Marked loss of myelinated
nerve fibers in the human brain with age. J. Comp. Neurol. [Internet]
2003;462:144–52. Available from: https://doi.org/10.1002/cne.10714

50. McGinnis SM, Brickhouse M, Pascual B, Dickerson BC. Age-related changes
in the thickness of cortical zones in humans. Brain Topogr.2011;24:279–91.

51. Kennedy KM, Raz N. Age, Sex and Regional Brain Volumes Predict
Perceptual-Motor Skill Acquisition. Cortex [Internet] 2005;41:560–9. Available
from: https://www.sciencedirect.com/science/article/pii/S0010945208701965

52. Shinohara M, Keenan KG, Enoka RM. Contralateral activity in a homologous
hand muscle during voluntary contractions is greater in old adults. J. Appl.
Physiol. [Internet] 2003;94:966–74. Available from:
https://doi.org/10.1152/japplphysiol.00836.2002

53. Marneweck M, Loftus A, Hammond G. Short-interval intracortical inhibition and
manual dexterity in healthy aging. Neurosci. Res. [Internet] 2011;70:408–14.
Available from:
https://www.sciencedirect.com/science/article/pii/S0168010211001222

54. Plow EB, Cunningham DA, Bonnett C, Gohar D, Bayram M, Wyant A, et al.
Neurophysiological correlates of aging-related muscle weakness. J.
Neurophysiol. 2013;110:2563–73.

55. Ferlinc A, Fabiani E, Velnar T, Gradisnik L. The Importance and Role of
Proprioception in the Elderly: a Short Review. Mater. Sociomed. 2019;31:219–
21.

56. Winarakwong L, Muramoto T, Soma K, Takano Y. Age-related changes and
the possible adaptability of rat jaw muscle spindles: immunohistochemical and
fine structural studies. Arch. Histol. Cytol. 2004;67:227–40.

57. Macefield VG, Knellwolf TP. Functional properties of human muscle spindles. J.
Neurophysiol. [Internet] 2018;120:452–67. Available from:
https://doi.org/10.1152/jn.00071.2018

58. Kido A, Tanaka N, Stein RB. Spinal excitation and inhibition decrease as
humans age. Can. J. Physiol. Pharmacol. 2004;82:238–48.

59. Mehrkanoon S, Breakspear M, Boonstra TW. The reorganization of
corticomuscular coherence during a transition between sensorimotor states.
Neuroimage 2014;100:692–702.

60. Scaglioni G, Narici M V, Maffiuletti NA, Pensini M, Martin A. Effect of ageing on
the electrical and mechanical properties of human soleus motor units
activated by the H reflex and M wave. J. Physiol. 2003;548:649–61.

61. Baudry S, Maerz AH, Enoka RM. Presynaptic modulation of Ia afferents in
young and old adults when performing force and position control. J.
Neurophysiol. 2010;103:623–31.

62. Peters RM, McKeown MD, Carpenter MG, Inglis JT. Losing touch: age-related
changes in plantar skin sensitivity, lower limb cutaneous reflex strength, and
postural stability in older adults. J. Neurophysiol. 2016;116:1848–58.

63. Lee RH, Heckman CJ. Bistability in spinal motoneurons in vivo: systematic
variations in persistent inward currents. J. Neurophysiol. 1998;80:583–93.

64. Karrer TM, McLaughlin CL, Guaglianone CP, Samanez-Larkin GR. Reduced
serotonin receptors and transporters in normal aging adults: a meta-analysis of
PET and SPECT imaging studies. Neurobiol. Aging [Internet] 2019;80:1–10.
Available from:
https://www.sciencedirect.com/science/article/pii/S019745801930106X

65. Deza-Araujo YI, Baez-Lugo S, Vuilleumier P, Chocat A, Chételat G, Poisnel G,
et al. Whole blood serotonin levels in healthy elderly are negatively associated

with the functional activity of emotion-related brain regions. Biol. Psychol.
[Internet] 2021;160:108051. Available from:
https://www.sciencedirect.com/science/article/pii/S0301051121000429

66. Shibata E, Sasaki M, Tohyama K, Kanbara Y, Otsuka K, Ehara S, et al. Age-
related Changes in Locus Ceruleus on Neuromelanin Magnetic Resonance
Imaging at 3 Tesla. Magn. Reson. Med. Sci. 2006;5:197–200.

67. Alvarez J, Alvarez-Illera P, García-Casas P, Fonteriz RI, Montero M. The Role
of Ca(2+) Signaling in Aging and Neurodegeneration: Insights from
Caenorhabditis elegans Models. Cells 2020;9.

68. Gonzales GF, Carrillo C. Blood serotonin levels in postmenopausal women:
Effects of age and serum oestradiol levels. Maturitas [Internet] 1993;17:23–9.
Available from:
https://www.sciencedirect.com/science/article/pii/0378512293901207

670