Bumblebee mouthparts exhibit poor acuity for the detection of pesticides in

nectar

Rachel H. Parkinson[1], Jennifer Scott[1], Anna L. Dorling[1], Hannah Jones[2], Martha

Haslam[1], Alex E. McDermott-Roberts[1], Geraldine A. Wright[1]

[1]Department of Biology, the University of Oxford, Oxford, UK; [2]Department of Life

Sciences, Imperial College, London

Author contributions: G.A.W, J.S, and R.H.P designed research. J.S, A.L.D, M.H, and

A.E.MD-R collected behavioural data. R.H.P and H.J. collected electrophysiology data,

R.H.P performed statistical analyses and prepared visualizations. R.H.P, J.S, A.L.D.,

and G.A.W. wrote the manuscript.

Abstract (max 200 words)

Bees are important pollinators of agricultural crops, but their populations are at risk

when pesticides are used. One of the largest risks bees face is poisoning of floral nectar

and pollen by insecticides. Studies of bee detection of neonicotinoids have reported

contradictory evidence about whether bees can taste these pesticides in sucrose

solutions and hence avoid them. Here, we use an assay for the detection of food

aversion combined with single-sensillum electrophysiology to test whether the

mouthparts of the buff-tailed bumblebee (*Bombus terrestris*) detect the presence of

pesticides in a solution that mimicked the nectar of oilseed rape (*Brassica napus*). Bees

did not avoid consuming solutions containing concentrations of imidacloprid,

thiamethoxam, clothianidin, or sulfoxaflor spanning six orders of magnitude, even when

these solutions contained lethal doses. Only extremely high concentrations of the

pesticides altered spiking in gustatory neurons through a slight reduction in firing rate or

change in the rate of adaptation. These data provide strong evidence that bumblebees

cannot detect or avoid field-relevant concentrations of pesticides using information from

their mouthparts. As bees rarely contact floral nectar with other body parts, we predict

that they are at high risk of unwittingly consuming pesticides in the nectar of pesticide-
treated crops.

Keywords: Neonicotinoids, sulfoxaflor, bumblebees, oil seed rape, taste perception

Main text

Insect pollination is a critical ecosystem service. The most economically valuable
pollinators are bees (Calderone, 2012; Garibaldi et al., 2014; Huang and An, 2018),
especially domesticated bees that secure crop yield and quality (Meehan et al., 2011).
In agricultural ecosystems, however, insecticides are widely used to protect crop yields
(Klein et al., 2007). Using insecticides impacts the health and survival of bees,
contributing to the recent population declines of all bee species (Goulson et al., 2015;
Potts et al., 2016; van der Sluijs et al., 2013; Vanbergen et al., 2013). Hundreds of
studies have shown that exposure to sublethal concentrations of neonicotinoid
pesticides (i.e. imidacloprid, IMD, thiamethoxam, TMX, and clothianidin, CLO) in food
impairs foraging behaviour, homing and orientation behaviour, and olfactory learning
and memory (e.g., Gill et al., 2012; Henry et al., 2012; Parkinson et al., 2022a;
Schneider et al., 2012a; Williamson et al., 2014; Wright et al., 2015). Most studies
administer pesticides in sugar solutions; bees consume the solution, and then a change
in behaviour or performance is recorded. If bees could detect and avoid pesticides in
food, however, these substances could be used to defend crops against pests without
risk to bees. Only a few have explicitly tested whether bees can detect and avoid
neonicotinoid pesticides in nectar (Arce et al., 2018; Kessler et al., 2015; Muth et al.,
2020), but the conclusions of these studies are contradictory.

We previously found that when bees are given a choice between sucrose solution and
sucrose containing field-relevant doses of IMD and TMX, bees choose the neonicotinoid
solution (Kessler et al., 2015). This could indicate that bees find neonicotinoid pesticides
phagostimulatory. However, we explicitly tested if bees could detect IMD, TMX, or CLO
in water and in sucrose solution by recording from the gustatory receptor neurons
(GRNs) of the galeal sensilla of the bees' mouthparts (Kessler et al., 2015). We failed to

find a change in the rate of GRN spiking that indicated detection and deduced that
these compounds were not detected by bees using their sense of taste. This
conclusion, however, has been challenged by a study in free-flying *B. terrestris* foragers
(Arce et al., 2018). They also found that free-flying bumblebees preferred specific
concentrations of solutions laced with TMX (Arce et al., 2018). In this study, the location
of the solutions was randomized, forcing a choice at the point of feeding. These authors
concluded that the only way bees could solve this problem was if they used their sense
of taste to identify the preferred solution, arguing that further work needed to be done to
identify the mechanism for sensation. Subsequently, however, another study in a
different species of bumblebees (*B. impatiens*) using methods similar to ours reported
that bees are neither attracted nor deterred from consuming sugar solutions containing
neonicotinoids, consistent with the idea that they cannot detect the pesticides (Muth et
al., 2020). Thus, whether or not bumblebees can detect neonicotinoids in food using
their sense of taste remains unresolved.

Insects detect potential toxins using GRNs located in contact chemosensilla on their
mouthparts, antennae, and tarsi using two mechanisms: active spiking of GRNs
expressing receptors tuned to bitter substances (Weiss et al., 2011) or via the
inactivation of sugar-sensing GRNs (French et al., 2015). In our previous work (Kessler
et al., 2015), stimulation of the mouthpart galeal sensilla with neonicotinoids did not
elicit spikes in putative bitter sensing GRNs. We also did not observe reduction in the
rate of firing of the GRNs in the sensilla. However, we have since developed more
advanced methods of spike sorting and data analysis that made it possible to discover
that bumblebees have highly specialized mechanisms for encoding sugars. In
bumblebee galeal sensilla, stimulation with sugars produces coherent spiking in 2
GRNs which burst in response to stimulation with high-value saccharide compounds
(Miriyala et al., 2018; Parkinson et al., 2022b). We discovered that bumblebees also
highly value the monosaccharide, fructose, and have a GRN tuned to detect it
(Parkinson et al., 2022b). It is possible that changes to the burst pattern of firing could
indicate that the presence of a compound such as a neonicotinoid in food. Furthermore,
the type of sugar used to stimulate the sensilla could also influence whether pesticides

were detected in food (Parkinson et al., 2022b). We also know that the labial palps, a
section of the mouthparts critical to toxin detection in other insect species (Chapman
and Sword, 1993), could house additional GRNs sensitive to bitter compounds like
pesticides. Our previous work used sucrose to stimulate the galeal sensilla and did not
analyse how pesticides might alter the burst structure of the galeal GRN response to
sugars or test the labial palps. Clearly, deeper investigations could determine whether
the mouthparts GRNs possess novel gustatory mechanisms for pesticide detection and
provide critical information about the risk to bees of consuming the nectar of pesticide-
defended crops in the field.

Here, we use a combination of sensitive behavioural assays and electrophysiology to
test in detail whether bumblebee mouthparts have mechanisms to detect pesticides in
nectar. Feeding assays using freely moving foragers make it possible to assess if bees
detect and avoid potential toxins in food (Ma et al., 2016). To definitively test whether
bumblebees detect neonicotinoids on their mouthparts, we quantified the structure of
feeding when the mouthparts of freely-moving bumblebees (*B. terrestris)* were
stimulated with nectar-like solutions containing pesticides over a 2 min period (Ma et al.,
2016). We specifically tested a mixture of primarily fructose and glucose that mimicked
the nectar of oilseed rape (*Brassica napus*) with field-relevant concentrations of
neonicotinoids. In addition, we comprehensively tested these solutions on the gustatory
receptor neurons in the A-type sensilla on the galea and the labial palps of the bees'
mouthparts using electrophysiology to determine whether sensilla in multiple locations
are able to detect neonicotinoids (IMD, CLO, and TMX) and the sulfoxamine pesticide,
sulfoxaflor (SFX). The burst-spiking of the galeal GRNs was analysed for changes to
structure and compared to the GRN response to the bitter compound, quinine (QUI),
which has been reported previously to be a feeding deterrent to bees (Ma et al., 2016;
Wright et al., 2010).

Methods

Bumblebee colonies

Bumblebee colonies (*Bombus terrestris audax,* Biobest, Westerlo, Belgium*)* were maintained at laboratory conditions (22-27°C and 35-40% RH) at the University of Oxford and fed *ad libitum* with the proprietary sugar syrup provided with the colonies (BioGluc®, Biobest, Westerlo, Belgium) containing fructose (37.5%), glucose (34.5%), sucrose (25%), maltose (2%), oligosaccharides (1%), and the preservatives potassium sorbate (E202) 0.15% and citric acid (E330) 0.06% (Wäckers et al., 2017). Bumblebees were provided with freeze-dried honeybee collected pollen (approx. 10 g) (Agralan Growers, Wiltshire, UK) three times/week.

In total, 12 colonies were used for the taste assays and 6 colonies were used for the electrophysiological study. Bees were randomly assigned to a treatment and colonies were counterbalanced across treatments to control for intercolony variation.

Solutions

An artificial oilseed rape nectar sugar solution was created containing 1.04 M glucose (D-(+)-Glucose, Sigma Aldrich), 0.746 M fructose (D-(-)-Fructose, Sigma Aldrich), and 0.007 M sucrose (Sigma Aldrich) (Carruthers et al., 2017), hereafter referred to as 'OSR'.

Stock solutions (1 mM) were made for each pesticide in OSR solution. No solvents are needed at these concentrations as all pesticides used are readily soluble in water at concentrations far exceeding 1 mM. Concentrations of imidacloprid (IMD), thiamethoxam (TMX), and clothianidin (CLO) in floral nectar range from 1.3 to 64 nM (Wood and Goulson, 2017), and sulfoxaflor (SFX) can reach concentrations as high as 71.4 µM (U.S. Environmental Protection Agency (EPA), 2014). In order to assess field-relevant concentrations of each chemical, a concentration series of each of IMD, TMX, CLO (Sigma Aldrich) and SFX (Chem Service) was made via tenfold serial dilution

using OSR to produce nectar containing the pesticides at 1000, 100, 10, and 1, and 0

nM (pure OSR). Additionally, 0.1 mM of each chemical dissolved in OSR was assessed

alongside 0.1 mM and 1.0 mM QUI ((-)-Quinine hydrochloride dihydrate, Sigma Aldrich)

in OSR, which functioned as positive controls for the effects of a bitter tastant. A

concentration of 0.1 mM was chosen for the pesticides as this encompasses the higher

range of SFX found in nectar, and allowed us to see if extremely high concentrations of

IMD, TMX and CLO could be detected by bees.

Aversive taste assay

Bumblebee workers were collected directly from the colony under red light and placed

individually into plastic holding vials with ventilation holes drilled into the lid. To minimise

inclusion of nest bees (i.e. bees that never or rarely forage), only workers with a thorax

width > 4.0 mm were used in these experiments (Goulson et al., 2002). In order to

motivate the bees to feed during the assay, bees were deprived of food between 3-6 h.

Bees were held in individual plastic holding vials at laboratory conditions in total

darkness throughout the starvation period. Following the starvation period, individual

bees were transferred into a 15 ml falcon tube, modified such that the tip of the tube

was removed and a small (10 mm x 20 mm) steel mesh was affixed contiguous with the

resulting hole. Bees could extend their proboscis through this hole to reach the feeding

solutions, whilst the mesh served as a surface for the bees to grip onto to maintain their

position during feeding. The tube was then affixed to a polystyrene holder with dental

wax (Ma et al., 2016).

A 60 mm long, 100 µL glass capillary tube was filled with a test solution and scanned

onto a computer at 600 dpi to produce an image from which the start volume could be

measured. The capillary tube was then connected to a silicone tube which was in turn

connected to a 1.0 ml syringe via a female connector, as in (Ma et al., 2016). This

syringe functioned as a pipette bulb to maintain the feeding solution at the tip of the

microcapillary. The capillary was held in place in the apparatus via a 1.0 ml modified

syringe affixed to a micromanipulator. A DinoLite digital microscope camera (Model

AM4815ZT, DinoLite, The Netherlands) was positioned 20 cm above the feeding site where the bees accessed the test solutions. This was connected to a computer and set to record using DinoLite Digital Microscope software (DinoLite, The Netherlands) at a frame rate of 640 X 480 at 25X magnification. All feeding behaviour was recorded for later analysis.

Once a bee was positioned in the apparatus, they were given 3 min to habituate to their environment before the assay began. Following this, bees were encouraged to extend their proboscis by touching the bee's terminal antennomers with a droplet (~3.5 µL) of 0.5 M sucrose (Sigma Aldrich) dissolved in deionised water. Once the bee extended their proboscis, the droplet was presented to the bees' mouthparts for them to consume. Each bee was given up to 5 min to extend their proboscis and consume the droplet. Bees that did not do so were removed from the experiment. When the bee finished consuming the droplet, the microcapillary tube containing the test solution was presented to the bees extended mouthparts. The test phase began once the bee's proboscis contacted the test solution, and the bee was given 2 min to consume the solution. After feeding, microcapillaries were re-scanned to measure the end volume.

All experimenters were blind to the experimental treatments. All solutions were tested in a randomised order and treatments were counterbalanced over time to eliminate any effects of starvation time on feeding behaviour.

Measuring solution consumption and feeding behaviour

ImageJ (Schneider et al., 2012b) was used to produce measurements of the length of solution in each microcapillary before and after the assay. The reference scale was set to 60.0 mm. Image files were zoomed in to 400% and the length of the solution inside the microcapillary was measured meniscus to meniscus. The length of test solution consumed by each bee was calculated as the difference between the measured length of the liquid inside the microcapillary tube before and after the test phase. These lengths were then converted to volumes using the formula:

$(100\ \mu L \times X\ \text{mm}) / 60.0\ \text{mm}$

Where 100 µL is the maximum volume of the capillary tube, 60.0 mm is the length of the

capillary tube, and X mm is the amount of solution consumed as a measured length

within the capillary.

A feeding bout is defined as a period in which contact is made between the proboscis

and test solution that is not separated by an absence of contacts for 5 s or more. The

number of feeding bouts and their duration can be used to evaluate the

phagostimulatory or deterrent activity of the compounds tested (Ma et al., 2016). To

identify bouts of feeding behaviour, videos were played back at 50 % speed and periods

of contact between the bees' mouthparts and the solution were scored using the Noldus

Observer (Noldus, the Netherlands).

Electrophysiology

We performed extracellular tip recordings on individual taste sensilla on the bumblebee

mouthparts to assess the peripheral taste perception of sucrose, a 10% dilution of

oilseed rape nectar (10% OSR), and the addition of pesticides or QUI to 10% OSR. We

recorded from 'A-type' sensilla from the galea (Gal), which are known to produce

bursting spike trains in response to sugars (Miriyala et al., 2018) and the third and fourth

segments of the labial palps (LP_{III} and LP_{IV}, respectively, Figure 1 A). Bumblebees were

prepared for electrophysiology as described previously (Miriyala et al., 2018; Parkinson

et al., 2022b). Sensilla were stimulated for 5 s with a borosilicate (Clark capillary glass

30-066, GC150TF-10) recording electrode (20 µm tip diameter, made with a Narishige

PC-10 electrode puller) containing the test solution. Signals were acquired with a pre-

amplifier (TasteProbe; Syntech, Germany), amplified (AC amplifier 1800, A-M Systems,

USA), digitized at 30 kHz (DT9803 Data Translation) and stored using DataView

(version 11.5). Pesticides (IMD, CLO, TMX, SFX) were tested in a background of 10%

OSR. Pesticides were tested using a concentration gradient (1, 10, 100, and 1000 nM)

and also at 0.1 mM. Additionally, QUI was tested in 10% OSR at concentrations of 0.1

and 1.0 mM. A minimum inter-stimulus interval of 3 min was used to prevent sensory

adaptation. We obtained recordings from n=12 bee per compound and tested 2-4

sensilla per mouthpart location (6-12 sensilla per bee). To ensure the pesticides or QUI

were not imposing a toxic effect on the taste sensilla, we included recordings at the

beginning and end of each stimulation series with 10% OSR (SI Figure 1).

Spike detection

Spikes from GRNs on the galea were extracted as described previously (Parkinson et

al., 2022b). Briefly, after band-pass filtering (300-2500 Hz, second order butterworth

filter) and normalization, spikes were detected using the peakfinder function with

adequate thresholds manually set for each recording. Interspike intervals were used to

detect the end of burst positions (EOB). Using this method, we detected three GRNs

within the galeal electrophysiological recordings, as seen previously (Parkinson et al.,

2022b). GRNs of the labial palps were non-bursting. A custom-written routine in Matlab

(The MathWorks) was used to differentiate spiking units (i.e. GRNs) within these

recordings. Spiking units were detected as described previously, and visual inspection

of the data suggested that two GRNs were present in these recordings. The spike

waveforms for each recording were compared to differentiate the spikes from the

putative units. Waveforms were aligned from the peak +/- 2 ms, and the Matlab singular

value decomposition (SVD) function was used to obtain the principal components of the

spike waveforms. Waveforms were then projected onto the space spanned by the first

two principal components, and K-means clustering was used to assign the waveforms

to units for each recording. Using this method, two (or in some cases just one) GRNs

were obtained from each recording, with the unit labelled GRN1 represented by larger

waveform amplitudes and higher spike frequencies in response to sugars. GRN

responses from individual animals were averaged across sensilla recorded in each

location (Gal, LP_{III}, LP_{IV}).

Statistical analyses and data presentation

We performed all further analyses in R version 4.2.1 (Team, 2016). Average firing rates

of the 7 GRN types (Gal GRNs 1-3, LP_{III} GRNs 1-2, and LP_{IV} GRNs 1-2) across

mouthpart locations were compared. Firing rates of a given GRN type were averaged

across sensilla from a single animal, and log transformed (log_{10}(average_rate + 1)) to fit

a normal distribution. We compared electrophysiological responses across stimuli using

linear mixed effects models (LME, lmerTest toolbox) (Kuznetsova et al., 2017) with

Gaussian   distributions   and   "bee   ID"   as   the   random   effect

(avg_rate~stimulus*GRN+(1|BeeID)). Models with non-significant interaction terms were

re-run without the term (avg_rate~stimulus+GRN+(1|BeeID)). *Post hoc* analyses were

performed using estimated marginal means (emmeans toolbox, (Lenth et al., 2018))

with Tukey's adjustment for family-wise error rates. Significant effects were denoted

with asterisks (*, <0.05; ** <0.001; ***, <0.0001).

The temporal patterns of GRN activity over time was also assessed by extracting spike

rates for each GRN within 0.1 s bins from 0.1 to 1.1 s after stimulation. Firing rates

within each bin were averaged by animal for a given GRN type. As the labial palp GRNs

did not display a bursting pattern, we averaged the firing rate within each bin across all

four labial palp GRNs. The pattern of bursting of galeal GRNs (GRNs 1 and 2) was

assessed using two measures: the burst length (number of GRN1 spikes per burst) and

burst rate. Firing rate histograms were constructed using these measurements over 1 s

of   stimulation,   and   stimuli   were   compared   with   LME   models

(temporal_parameter~stimulus+bin+(1|BeeID)). Significant differences in the slopes

between stimuli were denoted on plots with letters. To assess differences in the

complete temporal code of all 7 GRNs across mouthparts, we aligned the binned spike

times in series for each GRN. For each animal and stimulus, there was a vector of 70

bins (ten 0.1s bin per GRN type, aligned for 7 GRNs). To reduce the variability in

responses between animals and balance the scale of responses across GRNs, we

normalized the responses across GRN replicates within each animal by dividing by the

average responses of each GRN to 10% OSR. The normalized responses of single

GRNs were then averaged across replicates for each animal. We used t-distributed

stochastic neighbour embedding (t-SNE) to reduce the dimensionality of the temporal

data to two axes for each dataset (OSR vs sucrose, QUI, and pesticides). Results from

t-SNE were then clustered to reveal grouping of responses by stimulus. Optimal cluster

count was predicted using Monte Carlo reference-based consensus clustering (M3C,

(John et al., 2020)), an unsupervised learning technique that aggregates across k-
means clustering runs. Optimal clustering (if k>1) was included on t-SNE plots.

Feeding behaviours were compared using t-tests, ANOVA with Tukey's *post hoc*
multiple comparisons (HSD), or LME for concentration gradient comparisons of
pesticide mixtures with "colony" as a random effect. Behavioural data that was not
normally distributed was log transformed. For all boxplots in the paper, the median is
indicated by a line, bounds of the box mark the 1st and 3rd quartile, and whiskers
extend to 1.5 times the interquartile range. Individual data points are drawn.

Results

GRNs encode differences in sugar solution composition

Responses of bumblebee galeal GRNs to individual sugar compounds have previously
been described (Miriyala et al., 2018; Parkinson et al., 2022b). Here, we tested whether
a mixture of fructose, glucose, and negligible amounts of sucrose that mimics OSR
nectar altered the GRN spiking responses compared to sucrose alone. We also
expanded our investigation to include responses form GRNs in the labial palps (Figure 1
A). We found that GRNs in each location had specific patterns of activation that
characterized their responses to sugar stimulation (Figure 1 B). Stimulation of galeal
sensilla with sucrose or with OSR nectar produced bursts of spikes resulting from the
co-activation of two GRNs (Gal1 and Gal2), as we found for individual sugars previously
(Miriyala et al., 2018; Parkinson et al., 2022b, Figure 1 B). Each labial palp sensillum
(LP_{III} and LP_{IV}) contained two sugar-sensitive GRNs (LP_{III} 1 and 2, LP_{IV} 1 and 2), but
these GRNs exhibited tonic firing patterns in response to stimulation instead of bursts of
spikes (Figure 1 B). The spiking rates averaged over 1 s stimulation of Gal1 and the
labial palp GRNs were unchanged towards the two stimuli, but the average rate of Gal2
and Gal3 when stimulated with sucrose was lower than the rate for stimulation with 10%
OSR (Figure 1 C, LME, stimulus*GRN $F_{6,1607}$ = 9.16, p < 0.0001). (Note: GRN3 was
active only when stimulated with 10% OSR, but not by an equimolar sucrose solution,
Figure 1 C).

We compared the time-varying responses of GRNs using binned spikes (0.1 s bins over

1 s stimulation). We found that the rate of adaptation, measured by the change in the

interspike interval of labial palp GRNs were not significantly different for 10% OSR or

sucrose (Figure 1 D, stim: $F_{1,461} = 3.11$, $p > 0.05$; time, $F_{1,450} = 586$, $p < 0.0001$; SI

Figure 1 B). However, galeal GRN burst structure (Gal1 and Gal2) differed between

stimuli (Figure 1 E, stim*GRN: $F_{1,938} = 13.8$, $p = 0.0002$, time: $F_{1,938} = 1.42$, $p > 0.05$).

OSR elicited a 20% higher burst rate (i.e., Gal2 rate, SI Figure 1 C) and shorter bursts

(i.e., fewer Gal1 spikes per burst, SI Figure 1 D), although the rate of adaptation of Gal1

did not differ as a function of stimulus (Figure 1 E). To understand if the population of

neurons we recorded from contained information about stimulus identity, we combined

the temporal responses binned at 100 ms intervals over 1 s of recording of all 7 GRNs

across mouthparts using a clustering algorithm (t-SNE and k-means clustering). The

algorithm predicted two statistically separate stimulation groups, one for OSR and one

for sucrose (Figure 1 F).

In spite of the differences observed in GRN responses, the total volume of the 10%

OSR and equimolar sucrose solutions (1.79 M, Figure 1 G) consumed by bumblebees

was not significantly different in the free-feeding assay (Figure 1 H, t-test, t = -1.44, df =

28, $p = 0.49$). We also did not observe a difference in the duration of the first feeding

bout (Figure 1 I, t-test, t = -0.50, df = 27, $p = 0.081$), or the total time the bees spent in

contact with the solutions over 2 min (Figure 1 J, t-test, t = -0.837, df = 27, $p = 0.38$). We

also tested whether the concentration of these sugar solutions influenced behaviour

using a concentration gradient of OSR and sucrose solutions at 2.0, 1.5, 1.0, 0.5 M and

water. A greater volume of sucrose was consumed for the 1 M and 1.5 M

concentrations than for the OSR solution, but there was no difference at any other

concentration (SI Figure 2). First bout and total contact duration were not significantly

different.

Labial palp GRNs may enable bitter detection

QUI is routinely used to test an animal's sensitivity to bitter substances. Here, we used

a concentration series of QUI in 10% OSR solution to test how mouthparts GRNs

respond to QUI, so that we could use this as a positive control for our tests with
pesticides. A 1 mM QUI concentration suppressed all galeal and labial palp GRNs,
while 0.1 mM QUI suppressed the activity of $LP_{III}2$ and $LP_{IV}2$ alone (Figure 2 A, LME,
stimulus*GRN $F_{12,1581} = 11.3$, $p < 0.0001$, SI Figure 3 A-C). The rate of adaptation,
however, of the labial palp GRNs was faster when they were stimulated with the 0.1 mM
QUI solution (Figure 2 B, stim*time: $F_{1,450} = 13.8$, $p = 0.0002$). Galeal GRN burst
structure was affected by the addition of QUI (Figure 2 C, stim*GRN: $F_{2,1116} = 308$, $p <$
0.0001, time: $F_{1,1116} = 0.062$, $p = 0.803$; SI Figure 3 B-C). Gal1 adapted faster when
stimulated with 0.1 mM QUI compared to QUI, although there was no effect on Gal2
(Figure 2 C). Clustering of all GRN responses predicted two groups, with the responses
to 1 mM QUI separating from the responses to 10% OSR and 0.1 mM QUI (Figure 2 D).
OSR and 0.1 mM QUI were not significantly different.
365

The detection threshold observed in the behaviour experiments was lower than
predicted by the t-SNE clustering but consistent with our measurements of adaptation in
labial palp neurons. Bumblebees consumed significantly less OSR solution when it
contained 0.1 mM QUI, while 1 mM QUI was completely deterrent (Figure 2 E, LME,
concentration: $F_{2,35} = 49.1$, $p < 0.0001$). However, the structure of feeding towards 1
mM QUI was similar to the results found for the cluster analysis of the GRNs: bees
tested with 1 mM QUI exhibited much shorter average first feeding bout duration (Figure
2 F, LME, concentration: $F_{2,40} = 18.2$, $p < 0.0001$) and shorter average cumulative bout
duration (Figure 2 G, LME, concentration: $F_{2,36} = 61.5$, $p < 0.0001$) than those fed with
OSR or 0.1 mM QUI solution.
376

Extremely high concentrations of pesticides in OSR nectar reduce the rate of
spiking of GRNs
In our previous work (Kessler et al. (2015), field-relevent concentrations of neonicotinoid
pesticides were not detected by galeal GRNs. Here, we used OSR instead of sucrose
and included sulfoxaflor. We also extended this experiment to test the labial palps. As
before, we found no significant difference in galeal GRN responses between the control
(OSR) solution and all of the solutions containing pesticides at field-relevant

concentrations (SI Figure 4). No significant differences were detected in the responses
of the labial palp neurons to the pesticide solutions (SI Figure 4).

To test whether all concentrations of pesticides go undetected by bees, we stimulated
GRNs with 0.1 mM of each pesticide mixed with 10% OSR. Such concentrations are 10-
100k fold greater than has been reported for neonicotinoids from the nectar and pollen
of agricultural crops and are at least 1k fold greater than the LD_{50} values for
bumblebees (Cresswell et al., 2014; Mundy-Heisz et al., 2022; Siviter et al., 2022).
Sulfoxaflor, however, is present at much higher concentrations in the field (e.g., 71.4
µM, U.S. Environmental Protection Agency (EPA), 2014)). Stimulation with extremely
high concentrations of TMX reduced the average firing rates of LP_{III} GRN2 and LP_{IV}
GRN2; none of the other pesticides affected the average GRN firing rates (Figure 3 A,
LME, stimulus*GRN: $F_{24,2139} = 1.65$, $p = 0.024$.)

We also examined how the temporal response of galeal and labial palp neurons was
influenced by stimulation with extreme concentrations of the pesticides. Labial palp
GRNs adapted faster when IMD or TMX were added to 10% OSR (Figure 3 B, stim:
$F_{1,821} = 49.9$, $p < 0.0001$; time, $F_{1,808} = 571$, $p < 0.0001$), and firing rates of labial palp
GRNs over 1 s were significantly reduced (SI Figure 5 A). Gal1 and Gal2 spiking and
adaptation rates were affected by CLO and IMD (Figure 3 C, stim*GRN: $F_{4,1653} = 3.47$, p
$= 0.0078$; time, $F_{1,1653} = 2.33$, $p = 0.127$). Galeal GRN bursting rate was slightly reduced
when stimulated with OSR mixed with CLO or IMD (SI Figure 5 B), while the burst
length was not affected by the addition of pesticides (SI Figure 5 C). We used the same
clustering algorithm that was able to differentiate the responses to sucrose and OSR.
Clustering using all of the data for each of the sensillum types failed to separate the
pesticide responses from responses to the control (10% OSR, Figure 3 D).

Bumblebees readily consume pesticides in nectar
We assessed whether the pesticides were aversive in the range of values that included
field-realistic concentrations (0 - 1000 nM) and one of the extreme values (0.1mM)
added to a 10% OSR solution. Field-relevant concentrations of the pesticides in OSR

nectar did not deter the bees from feeding, as the presence of these compounds did not
significantly alter the amount of food consumed (LME, concentration: $F_{4,297}$ = 0.209, p =
0.933), first bout duration (LME, concentration: $F_{4,290}$ = 0.648, p = 0.629) or cumulative
bout duration (LME, concentration: $F_{4,288}$ = 0.310, p = 0.871) over the 2 min assay
period (Figure 4 A-C). Importantly, we also tested feeding behaviour using an extreme
concentration (0.1 mM) of the pesticides in OSR. Even at this very high concentration,
we did not observe a significant difference in the amount of food consumed (AOV: $F_{4,78}$
= 0.975, p = 0.426), first bout duration (AOV, $F_{4,74}$ = 0.403, p = 0.806), or cumulative
bout duration (AOV, $F_{4,74}$ = 1.22, p = 0.309) over 2 min (Figure 4 D-F).
424

Discussion and conclusions

Using an assay developed to study avoidance behaviour towards bitter compounds in
bumblebees (Ma et al., 2016), we show that bumblebees do not avoid drinking OSR
nectar solutions even when very high concentrations of common pesticides are present.
Our electrophysiology assay also showed that the only concentrations that produced a
measurable reduction in the spiking of sugar-sensing GRNs were extremely high
concentrations of these pesticides. These small changes in the rate of spiking did not
translate into features that the clustering algorithm could use to differentiate pesticide
laced stimuli from the control OSR solution, however. Likewise, bumblebees were not
deterred from drinking the OSR solutions with high pesticide concentrations. From these
data, we conclude that the buff-tailed bumblebee's mouthparts do not have mechanisms
for the detection and avoidance of common cholinergic pesticides in nectar, as
previously reported by Kessler et al. (2015).
438

In most insects, bitter, non-nutrient compounds are detected by specialized GRNs
which spike in response to stimulation (Weiss et al., 2011). Bitter compounds can also
be detected through the suppression of the sugar-sensing GRNs (French et al., 2015).
In the present experiments, we were mainly testing for the reduction of GRN spiking
caused by the suppression of sugar-sensing GRNs, as all of the 'bitter' compounds we
tested were presented to the sensilla in OSR nectar (a mixture of primarily fructose and

glucose). The known bitter compound, QUI, clearly suppressed the activity of sugar-

sensing GRNs in both labial palps and the galea, but it didn't measurably alter the burst

pattern of firing of the galeal GRNs. Importantly, our new data were able to confirm that

field-relevant concentrations of pesticides did not significantly alter the pattern or rate of

spiking of galeal or labial palp GRNs towards a nectar solution.

We used a clustering algorithm that integrated input from all the sensilla we recorded

from over a 1 s time interval with every stimulus to identify whether population-level

input made it possible to differentiate the stimuli. Although we were able to measure

small but significant changes in the responses of both labial palp and galeal neurons

caused by the high concentrations (100 µM) of IMD, SFX, and TMX, this effect was not

sufficient for the clustering algorithm to form classes of distinct stimuli for the OSR-

pesticide mixtures. This is similar to what we observed for 0.1 mM QUI; though a small

change in firing was detected, the classification algorithm did not differentiate it from

OSR. Our electrophysiological recordings are the first to report GRN responses from the

bumblebee's labial palps. The labial palps in other insects such as locusts are used to

detect both phagostimulants and deterrant compounds (Blaney and Chapman, 1970),

so we might have rationally expected to observe labial palp GRNs that spike in

response to QUI and cholinergic pesticides, but we did not.

It is interesting to note that in the case of all bitter compounds tested, the output of the

clustering algorithm matched the behaviour of the bees, as the bees themselves did not

reject any of the pesticide solutions or the 0.1 mM QUI solution. This could indicate that

that small differences detected at the sensory periphery do not translate into a sufficient

signal across the whole population of gustatory neurons to produce an aversion to foods

containing potential toxins.   However, it is worth noting that the OSR background

solutions in both cases were different: in the behavioural experiments the concentration

of OSR was an average value of that naturally found in nectar. Our physiology

experiments were performed using a 10% dilution of this solution. Different solutions

were used because it was not possible to record from neurons with such a concentrated

sugar solution, and in behavioural experiments, most bumblebees do not find 10% OSR
sufficiently phagostimulatory to feed.

Our new data also indicate that gustatory coding is more than the presence or absence
of spikes arising from GRNs tuned to detect particular tastants. In bumblebees, it is
clear that the pattern of spiking itself conveys information which is a feature that enables
the brain to distinguish gustatory stimuli (Parkinson et al., 2022b). This supports the
idea that encoding of gustatory stimuli is accomplished by the whole population of
GRNs on a given body part (e.g. the proboscis) as was observed in multichannel
recordings from the adult hawkmoth maxillary nerve (Reiter et al. 2015). Thus, activity
across the mouthparts GRN population that includes temporal input over time is likely to
be used by the brain to differentiate taste stimuli and to facilitate the valuation of food
quality. Our new data testing the difference between a nectar stimulus composed of a
mixture of sugars (OSR) and sucrose clearly illustrates this. The changes in the rate of
bursting and spiking in the galeal and labial palps were sufficient for the clustering
algorithm to classify sucrose and OSR as separate stimuli. Although our behavioural
assay did not show a difference, we would not expect it to, as it is an assay developed
to study food rejection behaviour by bees, not differences in phagostimulation. We
predict that these two solutions are completely distinguishable by bumblebees; this has
exciting implications for the bee's ability to distinguish naturally occurring nectar of
different flowering plant species. Future studies will be necessary to confirm this.

Previously, Arce et al. (2018) reported that *B. terrestris* were capable of selectively
foraging on 30% w/v sucrose solutions containing thiamethoxam. They challenged our
previous conclusion that honeybee and bumblebee preferences for solutions containing
TMX were accomplished in the absence of taste cues in our choice assays (Kessler et
al. 2015). There are important differences in the experiments of Arce et al. (2018) and
those we present now. They used gravity feeders in an enclosed arena in which bees
could consume solutions *ad libitum*, switching the feeders between training and testing
phases to ensure bees could not predict where the TMX laced food was. The feeders
did not have visual or olfactory cues, so they deduced that the bees were capable of

tasting the presence of TMX in the solution. However, gravity feeders do not simulate
how bees feed on real flowers. Such feeders have large openings in which several bees
can feed simultaneously; in this setting, bees can contact the solution with all body
parts, including the tarsi and antennae. It is possible that the sensilla on the antennae or
tarsi could detect TMX, but unlikely. In honeybees, bitter detection in these locations is
limited and is only observed when the sensilla are contacted with concentrations of
compounds like QUI that are >1 mM (de Brito Sanchez et al., 2005; de Brito Sanchez et
al., 2014). Field-relevant concentrations of the pesticides we tested are between 3-6
orders of magnitude lower in concentration (see Kessler et al. 2015 for review); such
dilute concentrations of a bitter compound present in a highly concentrated sugar
solution would be difficult for most organisms to detect. The post-ingesitve mechanisms
that produce bee preferences for foods containing neonicotinoids remain to be
discovered.

Floral morphology such as corolla length has often co-evolved with particular groups of
pollinators with longer mouthparts; a long corolla protects nectar when visitors are
efficient and reliable pollinators. In many flowers, nectar is hidden deep within the
corolla such that the only contact that bumblebees have with it is via the distal end of
the proboscis (also the location where most of the sensilla are). For example, in OSR
flowers, nectaries are located at a depth of 5-8 mm (Cresswell et al., 2001) and the
nectar cannot be contacted otherwise unless the bee bites through the petals. In
addition, bees learn to handle flowers quickly to improve their foraging efficiency
(Heinrich, 1976) and even extend the proboscis in anticipation of entering the corolla
prior to landing. Thus, they are unlikely to contact nectar using gustatory sensilla on
their antennae or tarsi while foraging. For this reason, it is likely that nectar palatability is
largely determined by the sensilla on the proboscis. Based on our current results and
those of previous studies (Sculfort et al., 2021; Tiedeken et al., 2014), we predict that
bumblebees have a limited ability to detect naturally-occurring concentrations of toxic or
bitter compounds in nectar. Future studies that survey a greater variety of compounds
will be necessary to confirm this.

Acknowledgements

We would like to thank Priya Chakrabarti-Basu for the pilot studies that led to this work.

Funding

This research was supported by a BBSRC grant (BB/S000402/1) to GA Wright, and an NSERC PDF (PDF-546125-2020) and Royal Society Newton International Fellowship (NIF/R1/19368) to RH Parkinson.

Figure legends

Figure 1: Electrophysiological and behavioural responses to sucrose and oilseed rape (OSR) nectar.

A: Diagram of the bumblebee's mouthparts from which tip-recordings were made, including the galea, and segments III and IV of the labial palps. Tip-recordings were made from the longer 'A-type' sensilla.

B: Filtered electrophysiological recordings from galeal (Gal), labial palp segment III (LP_{III}), and labial palp segment IV (LP_{IV}) sensilla. Spikes from GRN 1 at each location are labeled with blue circles, GRN 2 spikes in green, and GRN 3 spikes in magenta (only two GRNs present in labial palp recordings).

C: Average firing rates of GRNs from sensilla on the galea (Gal), segment III of the labial palps (LP_{III}), and segment IV (LP_{IV}) over 1 s of stimulation with 10% OSR or an equimolar (0.173 M) sucrose solution (SUC). Mean and standard deviation illustrated with black bars, data shown with coloured points. Asterisks represent significant differences between SUC and OSR.

D: Linear regression of inter-spike intervals per 0.1 s bin of labial palp GRNs versus time. Slope of regression is the adaptation rate. Shading is standard error. No significant difference between the adaptation rate when stimulated with OSR versus sucrose.

E: Firing rates of Gal1 versus Gal2 over 1 s stimulation with 10% OSR and SUC. Points represent mean rate in each 100 ms bin across all trials. A black marker highlights the

first bin (i.e., time = 100 ms). *Post hoc* comparisons showed that Gal2 firing rates were
significantly different between stimuli.

F: t-SNE of all GRN responses for each animal following stimulation with OSR (gray) or
SUC (orange) and k-means clusters (k=2, predicted by Monte Carlo reference-based
consensus clustering) in gray shading.

G: Image of an untethered bumblebee in a holding tube feeding from a capillary in the
2-min feeding assay.

H: Total volume consumed of 1.79 M SUC or 100% OSR of freely moving bumblebees
during 2 minutes.

I: The duration of the first feeding bout duration of bumblebees feeding on SUC or OSR.

J: The cumulative duration of feeding bouts within a 2 min period.

Figure 2: Quinine shuts down GRN and behavioural responses to sugars.

A: Average firing rates of 7 GRNs from the galea (Gal), and labial palps (LP$_{III}$, LP$_{IV}$) over
a 1 s stimulations with 10% OSR (OSR), 10% OSR plus 0.1 mM quinine (QUI 0.1), and
10% OSR plus 1 mM quinine (QUI 1). Mean and standard deviation illustrated with
black bars, data shown with coloured points (n = 12). Asterisks represent significant
differences between stimulus and OSR.

B: Linear regressions of inter-spike intervals per 0.1s bin of labial palp GRNs versus
time illustrating the adaptation rate, with SEM in grey shading. The addition of 0.1 mM
QUI to OSR significantly affected labial palp adaptation. 1 mM QUI not shown as there
was insufficient spiking to calculate adaptation.

C: Firing rates of Gal1 versus Gal2 over 1 s stimulation. Points represent mean rate in
each 100 ms bin across all trials. A black marker highlights the first bin (i.e., time = 100
ms). *Post hoc* comparisons showed that only Gal1 (not Gal2) firing rates were
significantly different at 0.1 mM QUI, while both were affected by 1.0 mM QUI.

D: t-SNE of all GRN responses for each animal following stimulation with 10% OSR,
QUI 0.1 or QUI 1, and k-means clusters (k=2, predicted by Monte Carlo reference-
based consensus clustering) in gray shading.

E: Volume consumed by freely-moving bumblebees of 100% OSR, OSR plus 0.1 mM
QUI, or OSR plus 1mM QUI over a 2 min period (n = 15 bees/group). Asterisks denote
results of Dunnett's test.

F: First bout duration when feeding on OSR, QUI 0.1 or QUI 1 (n = 15 bees/group).
Asterisks denote results of Dunnett's test.

G: Cumulative bout duration over a 2 min period of bumblebees feeding on OSR, QUI
0.1 or QUI 1 (n = 15 bees/group). Asterisks denote results of Dunnett's test.

Figure 3: Pesticides affect the temporal code of bumblebee GRNs.

A: Average firing rates of 7 GRNs from the galea (Gal) and labial palps (LP$_{III}$, LP$_{IV}$) over
a 1 s stimulation with 10% OSR (OSR), or OSR plus 0.1 mM clothianidin (CLO),
imidacloprid (IMD), sulfoxaflor (SFX), or thiamethoxam (TMX). Mean and standard
deviation illustrated with black bars, and data shown with coloured points. Asterisks
represent significant differences between stimulus and OSR.

B: Linear regressions of inter-spike intervals per 0.1s bin of labial palp GRNs versus
time illustrating the adaptation rate, with SEM in grey shading.

C: Firing rates of Gal1 versus Gal2 over 1 s stimulation. Points represent mean rate in
each 100 ms bin across all trials. A black marker highlights the first bin (i.e., time = 100
ms). *Post hoc* comparisons showed that only Gal1 firing rates were significantly different
for CLO and IMD versus OSR, and Gal2 firing rates were affected by IMD only.

D: t-SNE of all GRN responses for each animal following stimulation with OSR, CLO,
IMD, SFX, or TMX. Monte Carlo reference-based consensus clustering predicted an
optimal cluster count of k=1 on the t-SNE.

Figure 4: Bumblebees do not avoid consuming pesticides in nectar at field
relevant concentrations

A: The total volume consumed by freely moving bumblebees during 2 min of 100% OSR
solutions containing increasing concentrations (nM) of CLO, IMD, TMX and SFX (n = 15
bees per group).

B: First feeding bout duration of OSR mixtures with increasing concentrations (nM) of
pesticides (CLO, IMD, SFX, TMX, n = 15 bees per group).

C: Cumulative feeding duration during 2 min of OSR mixtures with increasing

concentrations (nM) of pesticides (n = 15 bees per group).

D: Total volume consumed of OSR or OSR plus 0.1 mM CLO, IMD, TMX or SFX over 2

min by freely moving bumblebees (n = 15 bees per group).

E: First bout duration of bumblebees feeding on OSR and OSR mixed with 0.1 mM

pesticides (n = 15 bees per group).

F: Cumulative bout duration over 2 min by bees feeding on OSR and OSR mixed with

0.1 mM pesticides (n = 15 bees per group).

References

Arce, A. N., Ramos Rodrigues, A., Yu, J., Colgan, T. J., Wurm, Y. and Gill, R. J. (2018).

Foraging bumblebees acquire a preference for neonicotinoid-treated food with prolonged

exposure. *Proceedings of the Royal Society of London B: Biological Sciences* **285**, 8–11.

Blaney, W. M. and Chapman, R. F. (1970). The functions of the maxillary palps of acrididae

(Orthoptera). *Entomol Exp Appl* **13**, 363–376.

Calderone, N. W. (2012). Insect Pollinated Crops , Insect Pollinators and US Agriculture:

Trend Analysis of Aggregate Data for the Period 1992 – 2009. *PLoS One* **7**, 24–28.

Carruthers, J. M., Cook, S. M., Wright, G. A., Osborne, J. L., Clark, S. J., Swain, J. L. and

Haughton, A. J. (2017). Oilseed rape (Brassica napus) as a resource for farmland insect

pollinators: quantifying floral traits in conventional varieties and breeding systems. *GCB*

Bioenergy **9**, 1370–1379.

Chapman, R. F. and Sword, G. (1993). The importance of palpation in food selection by a

polyphagous grasshopper (Orthoptera: Acrididae). *J Insect Behav* **6**, 79–91.

Cresswell, J. E., Hagen, C. and Woolnough, J. M. (2001). Attributes of individual flowers of

Brassica napus L. are affected by defoliation but not by intraspecific competition. *Ann Bot*

88, 111–117.

Cresswell, J. E., Robert, F. X. L., Florance, H. and Smirnoff, N. (2014). Clearance of

ingested neonicotinoid pesticide (imidacloprid) in honey bees (Apis mellifera) and

bumblebees (Bombus terrestris). *Pest Manag Sci* **70**, 332–337.

de Brito Sanchez, M. G., Giurfa, M., de Paula Mota, T. R. and Gauthier, M. (2005).

Electrophysiological and behavioural characterization of gustatory responses to antennal

"bitter" taste in honeybees. *European Journal of Neuroscience* **22**, 3161–3170.

de Brito Sanchez, M. G., Lorenzo, E., Su, S., Liu, F., Zhan, Y. and Giurfa, M. (2014). The

tarsal taste of honey bees: behavioral and electrophysiological analyses. *Front Behav*

Neurosci **8**, 1–16.

French, A. S., Sellier, M.-J., Ali Agha, M., Guigue, A., Chabaud, M.-A., Reeb, P. D., Mitra,

A., Grau, Y., Soustelle, L. and Marion-Poll, F. (2015). Dual Mechanism for Bitter

Avoidance in Drosophila. *Journal of Neuroscience* **35**, 3990–4004.

Garibaldi, L. A., Steffan-Dewenter, I., Winfree, R., Aizen, M. A., Bommarco, R.,

Cunningham, S. A., Kremen, C. and Carvalheiro, L. G. (2014). Wild Pollinators

Enhance Fruit Set of Crops Regardless of Honey Bee Abundance. *Science (1979)* **339**,

1608–1611.

Gill, R. J., Ramos-Rodriguez, O. and Raine, N. E. (2012). Combined pesticide exposure

severely affects individual- and colony-level traits in bees. *Nature* **491**, 105–8.

Goulson, D., Peat, J., Stout, J. C., Tucker, J., Darvill, B., Derwent, L. C. and Hughes, W. O.

H. (2002). Can alloethism in workers of the bumblebee, bombus terrestris, be explained in

terms of foraging efficiency? *Anim Behav* **64**, 123–130.

Goulson, D., Nicholls, E., Botías, C. and Rotheray, E. L. (2015). Bee declines driven by

combined Stress from parasites, pesticides, and lack of flowers. *Science (1979)* **347**,.

Heinrich, B. (1976). The Foraging Specializations of Individual Bumblebees. *Ecol Monogr* **46**,

105–128.

Henry, M., Béguin, M., Requier, F., Rollin, O., Odoux, J., Aupinel, P., Aptel, J.,

Tchamitchian, S. and Decourtye, A. (2012). A common pesticide decreases foraging

success and survival in honey bees. *Science (1979)* **336**, 348–350.

Huang, J. X. and An, J. D. (2018). Species diversity, pollination application and strategy for

conservation of the bumblebees of China. *Biodiversity Science* **26**, 486–497.

John, C. R., Watson, D., Russ, D., Goldmann, K., Ehrenstein, M., Pitzalis, C., Lewis, M.

and Barnes, M. (2020). M3C: Monte Carlo reference-based consensus clustering. *Sci Rep*

10, 1–14.

Kessler, S. C., Tiedeken, E. J., Simcock, K. L., Derveau, S., Mitchell, J., Softley, S., Stout, J.
C. and Wright, G. a. (2015). Bees prefer foods containing neonicotinoid pesticides. *Nature*
521, 74–76.

Klein, A. M., Vaissière, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A.,
Kremen, C. and Tscharntke, T. (2007). Importance of pollinators in changing landscapes
for world crops. *Proceedings of the Royal Society B: Biological Sciences* **274**, 303–313.

Kuznetsova, A., Brockhoff, P. B. and Christensen, R. H. B. (2017). lmerTest Package: Tests
in Linear Mixed Effects Models . *J Stat Softw* **82**,.

Lenth, R., Singmann, H., Love, J., Buerkner, P. and Herve, M. (2018). emmeans: Estimated
Marginal Means, aka Least-Squares Means. *R package version*.

Ma, C., Kessler, S., Simpson, A. and Wright, G. (2016). A Novel Behavioral Assay to
Investigate Gustatory Responses of Individual, Freely-moving Bumble Bees. *Journal of*
Visualized Experiments **113**, e54233.

Meehan, T. D., Werling, B. P., Landis, D. A. and Gratton, C. (2011). Agricultural landscape
simplification and insecticide use in the Midwestern United States. *Proceedings of the*
National Academy of Sciences **108**, 11500–11505.

Miriyala, A., Kessler, S., Rind, F. C. and Wright, G. A. (2018). Burst Firing in Bee Gustatory
Neurons Prevents Adaptation. *Current Biology* **28**, 1585-1594.e3.

Mundy-Heisz, K. A., Prosser, R. S. and Raine, N. E. (2022). Acute oral toxicity and risks of
four classes of systemic insecticide to the Common Eastern Bumblebee (Bombus
impatiens). *Chemosphere* **295**, 133771.

Muth, F., Gaxiola, R. and Leonard, A. S. (2020). No evidence for neonicotinoid preferences in
the bumblebee Bombus impatiens. *R Soc Open Sci* **7**,.

Parkinson, R. H., Fecher, C. and Gray, J. R. (2022a). Chronic exposure to insecticides impairs
honeybee optomotor behaviour. *Frontiers in Insect Science* **2**, 1–14.

Parkinson, R. H., Kessler, S. C., Scott, J., Simpson, A., Bu, J., Al-Esawy, M., Mahdi, A.,
Miriyala, A. and Wright, G. A. (2022b). Temporal responses of bumblebee gustatory
neurons to sugars. *iScience* **25**,.

Potts, S. G., Imperatriz-Fonseca, V., Ngo, H. T., Aizen, M. A., Biesmeijer, J. C., Breeze, T.
D., Dicks, L. v., Garibaldi, L. A., Hill, R., Settele, J., et al. (2016). Safeguarding
pollinators and their values to human well-being. *Nature* **540**, 220–229.

Schneider, C. W., Tautz, J., Grünewald, B. and Fuchs, S. (2012a). RFID tracking of sublethal
effects of two neonicotinoid insecticides on the foraging behavior of Apis mellifera. *PLoS*
One **7**, 1–9.

Schneider, C. A., Rasband, W. S. and Eliceiri, Kevin, W. (2012b). NIH Image to ImageJ: 25
years of Image Analysis. *Nat Methods* **9**, 671–675.

Sculfort, O., Gérard, M., Gekière, A., Nonclercq, D., Gerbaux, P., Duez, P. and
Vanderplanck, M. (2021). Specialized Metabolites in Floral Resources: Effects and
Detection in Buff-Tailed Bumblebees. *Front Ecol Evol* **9**,.

Siviter, H., Matthews, A. J. and Brown, M. J. F. (2022). A Combined LD50 for
Agrochemicals and Pathogens in Bumblebees (Bombus terrestris [Hymenoptera: Apidae]).
Environ Entomol **51**, 378–384.

Team, R. C. (2016). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Tiedeken, E. J., Stout, J. C., Stevenson, P. C. and Wright, G. A. (2014). Bumblebees are not
deterred by ecologically relevant concentrations of nectar toxins. *Journal of Experimental*
Biology **217**, 1620–1625.

U.S. Environmental Protection Agency (EPA) (2014). Ecological Risk Assessment for the
Registration Review of Sulfoxaflor. 79.

van der Sluijs, J. P., Simon-Delso, N., Goulson, D., Maxim, L., Bonmatin, J. M. and
Belzunces, L. P. (2013). Neonicotinoids, bee disorders and the sustainability of pollinator
services. *Curr Opin Environ Sustain* **5**, 293–305.

Vanbergen, A. J., Garratt, M. P., Vanbergen, A. J., Baude, M., Biesmeijer, J. C., Britton,
N. F., Brown, M. J. F., Brown, M., Bryden, J., Budge, G. E., et al. (2013). Threats to an
ecosystem service: Pressures on pollinators. *Front Ecol Environ* **11**, 251–259.

Wäckers, F. L., Alberola, J. S., Garcia-Marí, F. and Pekas, A. (2017). Attract and distract:
Manipulation of a food-mediated protective mutualism enhances natural pest control. *Agric*
Ecosyst Environ **246**, 168–174.

Weiss, L. A., Dahanukar, A., Kwon, J. Y., Banerjee, D. and Carlson, J. R. (2011). The
molecular and cellular basis of bitter taste in Drosophila. *Neuron* **69**, 258–272.

Williamson, S. M., Willis, S. J. and Wright, G. A. (2014). Exposure to neonicotinoids
influences the motor function of adult worker honeybees. *Ecotoxicology* **23**, 1409–1418.

Wood, T. J. and Goulson, D. (2017). The environmental risks of neonicotinoid pesticides: a

review of the evidence post 2013. *Environmental Science and Pollution Research* **24**,

17285–17325.

Wright, G. A., Mustard, J. A., Simcock, N. K., Ross-Taylor, A. A. R., McNicholas, L. D.,

Popescu, A. and Marion-Poll, F. (2010). Parallel reinforcement pathways for conditioned

food aversions in the honeybee. *Current Biology* **20**, 2234–2240.

Wright, G. A., Softley, S. and Earnshaw, H. (2015). Low doses of neonicotinoid pesticides in

food rewards impair short-term olfactory memory in foraging-age honeybees. *Sci Rep* **5**, 1–

7.

Figure 1

Figure 2

Figure 3

Figure 4