Mouthparts of the bumblebee (*Bombus terrestris)* exhibit poor acuity for the
detection of pesticides in nectar

Rachel H. Parkinson[1], Jennifer Scott[1], Anna L. Dorling[1], Hannah Jones[2], Martha Haslam[1],
Alex E. McDermott-Roberts[1], Geraldine A. Wright[1]

[1]Department of Biology, the University of Oxford, Oxford, UK; [2]Department of Life
Sciences, Imperial College, London

Author contributions: G.A.W, J.S, and R.H.P designed research. J.S, A.L.D, M.H, and
A.E.MD-R collected behavioural data. R.H.P and H.J. collected electrophysiology data,
R.H.P performed statistical analyses and prepared visualizations. R.H.P, J.S, A.L.D., and
G.A.W. wrote the manuscript.

Abstract (max 200 words)

Bees are important pollinators of agricultural crops, but their populations are at risk when
pesticides are used. One of the largest risks bees face is poisoning of floral nectar and
pollen by insecticides. Studies of bee detection of neonicotinoids have reported
contradictory evidence about whether bees can taste these pesticides in sucrose
solutions and hence avoid them. Here, we use an assay for the detection of food aversion
combined with single-sensillum electrophysiology to test whether the mouthparts of the
buff-tailed bumblebee (*Bombus terrestris*) detect the presence of pesticides in a solution
that mimicked the nectar of oilseed rape (*Brassica napus*). Bees did not avoid consuming
solutions containing concentrations of imidacloprid, thiamethoxam, clothianidin, or
sulfoxaflor spanning six orders of magnitude, even when these solutions contained lethal
doses. Only extremely high concentrations of the pesticides altered spiking in gustatory
neurons through a slight reduction in firing rate or change in the rate of adaptation. These
data provide strong evidence that bumblebees cannot detect or avoid field-relevant
concentrations of pesticides using information from their mouthparts. As bees rarely
contact floral nectar with other body parts, we predict that they are at high risk of
unwittingly consuming pesticides in the nectar of pesticide-treated crops.

Keywords: Neonicotinoids, sulfoxaflor, bumblebees, oil seed rape, taste perception

Main text

Insect pollination is a critical ecosystem service. The most economically valuable pollinators are bees (Calderone, 2012; Garibaldi et al., 2014; Huang and An, 2018), especially domesticated bees that secure crop yield and quality (Meehan et al., 2011). In agricultural ecosystems, however, insecticides are widely used to protect crop yields (Klein et al., 2007). Using insecticides impacts the health and survival of bees, contributing to the recent population declines of all bee species (Goulson et al., 2015; Potts et al., 2016; Van der Sluijs et al., 2013; Vanbergen et al., 2013). Hundreds of studies have shown that exposure to sublethal concentrations of neonicotinoid pesticides (i.e. imidacloprid, IMD, thiamethoxam, TMX, and clothianidin, CLO) in food impairs foraging behaviour, homing and orientation behaviour, and olfactory learning and memory (e.g., Gill et al., 2012; Henry et al., 2012; Parkinson et al., 2022a; Schneider et al., 2012a; Williamson et al., 2014; Wright et al., 2015). Most studies administer pesticides in sugar solutions; bees consume the solution, and then a change in behaviour or performance is recorded. If bees could detect and avoid pesticides in food, however, these substances could be used to defend crops against pests without risk to bees. Only a few have explicitly tested whether bees can taste and avoid neonicotinoid pesticides in nectar (Arce et al., 2018; Kessler et al., 2015; Muth et al., 2020), but the conclusions of these studies are contradictory. In addition, other compounds that affect cholinergic signalling in the insect nervous system, such as sulfoxaflor, have been proposed as alternatives to neonicotinoids, but whether or not these compounds can be detected by the bee's sense of taste has not previously been tested.

Insects detect non-volatile compounds (i.e. tastants) through activation of GRNs in contact chemosensilla on mouthparts, antennae, and tarsi which express receptor proteins that bind with specific types of ligands (e.g. sugars, non-nutrient 'bitter' compounds, salts, etc.) (Wright, 2016). When a taste receptor binds with its ligand, this sets off a signal transduction cascade that results in depolarization of the neuron (Dethier, 1976). For example, several types of receptors tuned to bitter substances are expressed

in a specialized subset of GRNs (Weiss et al., 2011). When these GRNs spike in

response to stimulation, feeding reflexes are inhibited (French et al., 2015). Bitter

compound detection is also accomplished by the inactivation of sugar-sensing GRNs

(French et al., 2015). For compounds which are potentially toxic, simultaneous activation

of GRNs that directly inhibit feeding reflexes and silencing of GRNs that activate feeding

make it possible to form a rapid response in reaction to contact (Wright, 2016). Thus, we

expected that if bees could detect neonicotinoids as bitter compounds, we would see

activation of bitter-sensing GRNs and/or inactivation of sugar-sensing GRNs. Previously,

we observed that stimulation of the mouthpart's galeal sensilla with neonicotinoids in

water did not elicit spikes from any GRNs, nor did we see a reduction in the rate of firing

of sugar-sensing GRNs in the sensilla when stimulated with a mixture of sucrose and

neonicotinoid compound (Kessler et al., 2015).

In a two choice assay, we unexpectedly found that when bees were given a choice

between sucrose solution and sucrose containing field-relevant doses of IMD and TMX

over a 24 h period, the bees choose the neonicotinoid solution (Kessler et al., 2015). This

could indicate that bees find neonicotinoid pesticides phagostimulatory, like sugars.

However, we also failed to find evidence that neonicotinoid pesticides in water elicited

spikes in any galeal GRNs including the nutrient/sugar sensing neurons. For this reason,

we concluded that bees could not taste neonicotinoids in nectar or on their own in water.

This conclusion, however, has been challenged by a study in free-flying *B. terrestris*

foragers (Arce et al., 2018). They found that free-flying bumblebees preferred specific

concentrations of solutions laced with TMX, but only after a period of at least 10 days of

feeding on the solutions (Arce et al., 2018). In this study, the location of the solutions was

randomized, forcing a choice at the point of feeding. These authors concluded that the

only way bees could solve this problem was if they used their sense of taste to identify

the preferred solution, arguing that further work needed to be done to identify the

mechanism for sensation. Subsequently, however, another study in a different species of

bumblebees (*B. impatiens*) using methods similar to ours reported that bees are neither

attracted nor deterred from consuming sugar solutions containing neonicotinoids,

consistent with the idea that they cannot detect the pesticides (Muth et al., 2020). Thus,

whether or not bumblebees can taste neonicotinoids in food remains unresolved.

Bees have highly specialized mechanisms for encoding sugars which are different to

those reported for any other insect species. For example, in bumblebee galeal sensilla,

stimulation with sugars produces coherent spiking in 2 GRNs which burst in response to

stimulation with high-value saccharide compounds (Miriyala et al., 2018; Parkinson et al.,

2022b). We recently discovered that bumblebees also highly value the monosaccharide,

fructose, and have a GRN tuned to detect it (Parkinson et al., 2022b). It is possible that

changes to the burst pattern of firing could indicate that the presence of a compound such

as a neonicotinoid in food. Furthermore, the type of sugar used to stimulate the sensilla

could influence whether bees could taste neonicotinoids (Parkinson et al., 2022b). We

also know that the labial palps are critical to toxin detection in other insect species

(Chapman and Sword, 1993), and could house additional GRNs sensitive to bitter

compounds like pesticides. Our previous work on neonicotinoid taste used sucrose to

stimulate the galeal sensilla and did not analyse how pesticides might alter the burst

structure of the galeal GRN response to sugars or test the labial palps. Clearly, deeper

investigations could determine whether the mouthparts GRNs possess novel gustatory

mechanisms for pesticide detection and provide critical information about the risk to bees

of consuming the nectar of pesticide-defended crops in the field. They would also

elucidate novel mechanisms for the detection of bitter compounds.

Here, we use a combination of sensitive behavioural assays and electrophysiology to test

in detail whether bumblebee mouthparts have mechanisms to detect pesticides in nectar.

Feeding assays using freely moving foragers make it possible to assess if bees detect

and avoid potential toxins in food (Ma et al., 2016). To definitively test whether

bumblebees detect neonicotinoids on their mouthparts, we quantified the structure of

feeding when the mouthparts of freely-moving bumblebees (*B. terrestris)* were stimulated

with nectar-like solutions containing pesticides over a 2 min period (Ma et al., 2016). We

specifically tested a mixture of primarily fructose and glucose that mimicked the nectar of

oilseed rape (*Brassica napus*) with field-relevant concentrations of neonicotinoids. In

addition, we comprehensively tested these solutions on the gustatory receptor neurons

in the A-type sensilla on the galea and the labial palps of the bees' mouthparts using
electrophysiology to determine whether sensilla in multiple locations are able to detect
neonicotinoids (IMD, CLO, and TMX) and the previously untested sulfoxamine pesticide,
sulfoxaflor (SFX). The burst-spiking of the galeal GRNs was analysed for changes to
structure and compared to the GRN response to the bitter compound, quinine (QUI),
which has been reported previously to be a feeding deterrent to bees (Ma et al., 2016;
Wright et al., 2010).

Methods

Bumblebee colonies

Bumblebee colonies (*Bombus terrestris audax,* Biobest, Westerlo, Belgium*)* were
maintained at laboratory conditions (22-27°C and 35-40% RH) at the University of Oxford
and fed *ad libitum* with the proprietary sugar syrup provided with the colonies (BioGluc®,
Biobest, Westerlo, Belgium) containing fructose (37.5%), glucose (34.5%), sucrose
(25%), maltose (2%), oligosaccharides (1%), and the preservatives potassium sorbate
(E202) 0.15% and citric acid (E330) 0.06% (Wäckers et al., 2017). Bumblebees were
provided with freeze-dried honeybee collected pollen (approx. 10 g) (Agralan Growers,
Wiltshire, UK) three times/week.

In total, 12 colonies were used for the taste assays and 6 colonies were used for the
electrophysiological study. Bees were randomly assigned to a treatment and colonies
were counterbalanced across treatments to control for intercolony variation.

Solutions

An artificial oilseed rape nectar sugar solution was created containing 1.04 M glucose (D-
(+)-Glucose, Sigma Aldrich), 0.746 M fructose (D-(-)-Fructose, Sigma Aldrich), and 0.007
M sucrose (Sigma Aldrich) (Carruthers et al., 2017), hereafter referred to as 'OSR'.

Stock solutions (1 mM) were made for each pesticide in OSR solution. No solvents are

needed at these concentrations as all pesticides used are readily soluble in water at

concentrations far exceeding 1 mM. Concentrations of imidacloprid (IMD), thiamethoxam

(TMX), and clothianidin (CLO) in floral nectar range from 1.3 to 64 nM (Wood and

Goulson, 2017), and sulfoxaflor (SFX) can reach concentrations as high as 71.4 µM (U.S.

Environmental Protection Agency (EPA), 2014). In order to assess field-relevant

concentrations of each chemical, a concentration series of each of IMD, TMX, CLO

(Sigma Aldrich) and SFX (Chem Service) was made via tenfold serial dilution using OSR

to produce nectar containing the pesticides at 1000, 100, 10, and 1, and 0 nM (pure OSR).

Additionally, 0.1 mM of each chemical dissolved in OSR was assessed alongside 0.1 mM

and 1.0 mM QUI ((-)-Quinine hydrochloride dihydrate, Sigma Aldrich) in OSR, which

functioned as positive controls for the effects of a bitter tastant. A concentration of 0.1

mM was chosen for the pesticides as this encompasses the higher range of SFX found

in nectar, and allowed us to see if extremely high concentrations of IMD, TMX and CLO

could be detected by bees.

Aversive taste assay

Bumblebee workers were collected directly from the colony under red light and placed

individually into plastic holding vials with ventilation holes drilled into the lid. To minimise

inclusion of nest bees (i.e. bees that never or rarely forage), only workers with a thorax

width > 4.0 mm were used in these experiments (Goulson et al., 2002). In order to

motivate the bees to feed during the assay, bees were deprived of food between 3-6 h.

Bees were held in individual plastic holding vials at laboratory conditions in total darkness

throughout the starvation period. Following the starvation period, individual bees were

transferred into a 15 ml falcon tube, modified such that the tip of the tube was removed

and a small (10 mm x 20 mm) steel mesh was affixed contiguous with the resulting hole.

Bees could extend their proboscis through this hole to reach the feeding solutions, whilst

the mesh served as a surface for the bees to grip onto to maintain their position during

feeding. The tube was then affixed to a polystyrene holder with dental wax (Ma et al.,

2016).

A 60 mm long, 100 μL glass capillary tube was filled with a test solution and scanned onto
a computer at 600 dpi to produce an image from which the start volume could be
measured. The capillary tube was then connected to a silicone tube which was in turn
connected to a 1.0 ml syringe via a female connector, as in (Ma et al., 2016). This syringe
functioned as a pipette bulb to maintain the feeding solution at the tip of the microcapillary.
The capillary was held in place in the apparatus via a 1.0 ml modified syringe affixed to a
micromanipulator. A DinoLite digital microscope camera (Model AM4815ZT, DinoLite,
The Netherlands) was positioned 20 cm above the feeding site where the bees accessed
the test solutions. This was connected to a computer and set to record using DinoLite
Digital Microscope software (DinoLite, The Netherlands) at a frame rate of 640 X 480 at
25X magnification. All feeding behaviour was recorded for later analysis.

Once a bee was positioned in the apparatus, they were given 3 min to habituate to their
environment before the assay began. Following this, bees were encouraged to extend
their proboscis by touching the bee's terminal antennomers with a droplet (~3.5 μL) of 0.5
M sucrose (Sigma Aldrich) dissolved in deionised water. Once the bee extended their
proboscis, the droplet was presented to the bees' mouthparts for them to consume. Each
bee was given up to 5 min to extend their proboscis and consume the droplet. Bees that
did not do so were removed from the experiment. When the bee finished consuming the
droplet, the microcapillary tube containing the test solution was presented to the bees
extended mouthparts. The test phase began once the bee's proboscis contacted the test
solution, and the bee was given 2 min to consume the solution. After feeding,
microcapillaries were re-scanned to measure the end volume.

All experimenters were blind to the experimental treatments. All solutions were tested in
a randomised order and treatments were counterbalanced over time to eliminate any
effects of starvation time on feeding behaviour.

Measuring solution consumption and feeding behaviour

ImageJ (Schneider et al., 2012b) was used to produce measurements of the length of
solution in each microcapillary before and after the assay. The reference scale was set
to 60.0 mm. Image files were zoomed in to 400% and the length of the solution inside the
microcapillary was measured meniscus to meniscus. The length of test solution
consumed by each bee was calculated as the difference between the measured length
of the liquid inside the microcapillary tube before and after the test phase. These lengths
were then converted to volumes using the formula:

$$(100 \ \mu L \times X \ mm) \ / \ 60.0 \ mm$$

Where 100 µL is the maximum volume of the capillary tube, 60.0 mm is the length of the
capillary tube, and X mm is the amount of solution consumed as a measured length within
the capillary.

A feeding bout is defined as a period in which contact is made between the proboscis and
test solution that is not separated by an absence of contacts for 5 s or more (Ma et al.,
2016). The number of feeding bouts and their duration can be used to evaluate the
phagostimulatory or deterrent activity of the compounds tested (Ma et al., 2016). To
identify bouts of feeding behaviour, videos were played back at 50 % speed and periods
of contact between the bees' mouthparts and the solution were scored using the Noldus
Observer (Noldus, the Netherlands). The duration of the first bout and the cumulative bout
duration (over the 2 minute period) were compared between stimuli.

Electrophysiology

We performed extracellular tip recordings on individual taste sensilla on the bumblebee
mouthparts to assess the peripheral taste perception of sucrose, a 10% dilution of oilseed
rape nectar (10% OSR), and the addition of pesticides or QUI to 10% OSR. We recorded
from 'A-type' sensilla from the galea (Gal), which are known to produce bursting spike
trains in response to sugars (Miriyala et al., 2018) and the third and fourth segments of
the labial palps (LP_{III} and LP_{IV}, respectively, Figure 1 A). Bumblebees were prepared for
electrophysiology as described previously (Miriyala et al., 2018; Parkinson et al., 2022b).

Briefly, bees were cold-anesthetized and harnessed with the proboscis extended. The
mouthpart nerves were severed by making an incision at the base of the mouthparts to
prevent movement. The galea and labial palps were oriented for access to the sensilla
using dental wax and wire pins. Sensilla were stimulated for 5 s with a borosilicate (Clark
capillary glass 30-066, GC150TF-10) recording electrode (20 µm tip diameter, made with
a Narishige PC-10 electrode puller) filled with the test solution. Signals were acquired
with a pre-amplifier (TasteProbe; Syntech, Germany), amplified (AC amplifier 1800, A-M
Systems, USA), digitized at 30 kHz (DT9803 Data Translation) and stored using
DataView (version 11.5). Pesticides (IMD, CLO, TMX, SFX) were tested in a background
of 10% OSR. Pesticides were tested using a concentration gradient (1, 10, 100, and 1000
nM) and also at 0.1 mM. Additionally, QUI was tested in 10% OSR at concentrations of
0.1 and 1.0 mM. A minimum inter-stimulus interval of 3 min was used to prevent sensory
adaptation. We obtained recordings from n=12 bee per compound and tested 2-4 sensilla
per mouthpart location (6-12 sensilla per bee). To ensure the pesticides or QUI were not
imposing a toxic effect on the taste sensilla, we included recordings at the beginning and
end of each stimulation series with 10% OSR (SI Figure 1).

Spike detection

Spikes from GRNs on the galea were extracted as described previously (Parkinson et al.,
2022b). Briefly, after band-pass filtering (300-2500 Hz, second order butterworth filter)
and normalization, spikes were detected using the peakfinder function with adequate
thresholds manually set for each recording. Interspike intervals were used to detect the
end of burst positions (EOB). Using this method, we detected three GRNs within the
galeal electrophysiological recordings, as seen previously (Parkinson et al., 2022b).
GRNs of the labial palps were non-bursting, and visual inspection of the data suggested
that two GRNs were present in these recordings. The spike waveforms for each recording
were compared to differentiate the spikes from the putative units. Waveforms were
aligned from the peak +/- 2 ms, and the Matlab singular value decomposition (SVD)
function was used to obtain the principal components of the spike waveforms. Waveforms
were then projected onto the space spanned by the first two principal components, and
k-means clustering was used to assign the waveforms to units for each recording. Using

this method, two (or in some cases just one) GRNs were obtained from each recording,

with the unit labelled GRN1 represented by larger waveform amplitudes and higher spike

frequencies in response to sugars.

Statistical analyses and data presentation

We performed all further analyses in R version 4.2.1 (Team, 2016). Average firing rates

of the 7 GRN types (Gal GRNs 1-3, LP_{III} GRNs 1-2, and LP_{IV} GRNs 1-2) across mouthpart

locations were compared. Firing rates of a given GRN type were averaged across sensilla

from a single animal, and log transformed (log_{10}(average_rate + 1)) to fit a normal

distribution. We compared electrophysiological responses across stimuli using linear

mixed effects models (LME, lmerTest package) (Kuznetsova et al., 2017) with Gaussian

distributions and "bee ID" as the random effect (avg_rate~stimulus*GRN+(1|BeeID)).

Models with non-significant interaction terms were re-run without the term

(avg_rate~stimulus+GRN+(1|BeeID)). *Post hoc* analyses were performed using

estimated marginal means (emmeans package, (Lenth et al., 2018)) with Tukey's

adjustment for family-wise error rates. Significant effects were denoted with asterisks (*,

<0.05; ** <0.001; ***, <0.0001).

We also assessed the temporal patterns of GRN activity over time. Time series spike data

were averaged by animal for a given GRN type, and firing rates were calculated in 0.1 s

bins from 0.1 to 1.1 s. Because the labial palp GRNs did not display a bursting pattern,

we averaged the firing rate within each bin across all four labial palp GRNs. We assessed

the bursting pattern of galeal GRNs (Gal 1 and 2) was assessed using two measures: the

burst length (number of Gal1 spikes per burst) and burst rate. We constructed firing rate

histograms (firing rate versus time) using these measurements over 1 s of stimulation,

and compared stimuli with LME models (temporal_parameter~stimulus+bin+(1|BeeID))

implemented in the lmerTest package (Kuznetsova et al., 2017). We denoted significant

differences in the resulting slopes between stimuli on plots with letters, assessed with the

estimated marginal means (emmeans) package (Lenth et al., 2018) with Tukey's

adjustment for family-wise error rates. To assess differences in the complete temporal

code of all 7 GRNs across mouthparts, we aligned the binned spike times in series for

each GRN. For each animal and stimulus, there was a vector of 70 bins (ten 0.1s bin per

GRN type, aligned for 7 GRNs). To reduce the variability in responses between animals

and balance the scale of responses across GRNs, we normalized the responses across

GRN replicates within each animal by dividing by the average responses of each GRN to

10% OSR. The normalized responses of single GRNs were then averaged across

replicates for each animal. We applied t-distributed stochastic neighbor embedding (t-

SNE) to reduce the dimensionality of the temporal data into two components for each

dataset (OSR vs sucrose, QUI, and pesticides), using Euclidean distance as the distance

metric. The results from the t-SNE were subsequently clustered to reveal the grouping of

responses by stimulus. To predict the optimal cluster count, we employed Monte Carlo

reference-based consensus clustering (M3C, (John et al., 2020)), an unsupervised

learning technique that aggregates across multiple k-means clustering runs. If the optimal

number of clusters (k) was greater than one, we included clustering on the t-SNE plots.

Measurements of feeding behaviour (volume consumed, first bout, and cumulative bout)

were compared using t-tests LME, with "colony" as a random effect:

(feeding_measurement~stimulus+ (1|ColonyID)), or

(feeding_measurement~stimulus+concentration+(1|ColonyID)) (lmerTest package)

(Kuznetsova et al., 2017), with emmeans *post hoc* tests (emmeans package) (Lenth et

al., 2018) and Tukey's adjustment for family-wise error rates. The non-significant

interaction term (stimulus*concentration) was removed for all pesticide concentration

gradient models due to increased model fit (using AIC and BIC). Behavioural data that

was not normally distributed was log transformed. For all boxplots, the median is indicated

by a line, bounds of the box mark the 1st and 3rd quartile, and whiskers extend to 1.5

times the interquartile range. Where data points are drawn, these represent the

responses of individual GRNs.

Results

Sugar solution composition is encoded by GRNs

Responses of bumblebee galeal GRNs to individual sugar compounds have previously

been described (Miriyala et al., 2018; Parkinson et al., 2022b). Here, we tested whether

a mixture of fructose, glucose, and negligible amounts of sucrose that mimics OSR nectar

altered the GRN spiking responses compared to sucrose alone. We also expanded our

investigation to include responses form GRNs in the labial palps (Figure 1 A). We found

that GRNs in each location had specific patterns of activation that characterized their

responses to sugar stimulation (Figure 1 B). Stimulation of galeal sensilla with sucrose or

with OSR nectar produced bursts of spikes resulting from the co-activation of two GRNs

(Gal1 and Gal2), as we found for individual sugars previously (Miriyala et al., 2018;

Parkinson et al., 2022b, Figure 1 B). Each labial palp sensillum (LP$_{III}$ and LP$_{IV}$) contained

two sugar-sensitive GRNs (LP$_{III}$ 1 and 2, LP$_{IV}$ 1 and 2), but these GRNs exhibited tonic

firing patterns in response to stimulation instead of bursts of spikes (Figure 1 B). The

spiking rates averaged over 1 s stimulation of Gal1 and the labial palp GRNs were

unchanged towards the two stimuli, but the average rate of Gal2 and Gal3 when

stimulated with sucrose was lower than the rate for stimulation with 10% OSR (Figure 1

C, LME, stimulus*GRN $F_{6,1607}$ = 9.16, p < 0.0001). (Note: Gal3 was active only when

stimulated with 10% OSR, but not by an equimolar sucrose solution, Figure 1 C).

We compared the time-varying responses of GRNs using binned spikes (0.1 s bins over

1 s stimulation) averaged by animal across sensilla. The coefficient of variation of spike

rates across sensilla for each animal for a given stimulus ranged from 0.023 to 2.44 with

a median of 0.35 and IQR of 0.47. We found that the rate of adaptation, measured by the

change in the interspike interval of labial palp GRNs were not significantly different for

10% OSR or sucrose (Figure 1 D, stim: $F_{1,461}$ = 3.11, p = 0.079; time, $F_{1,450}$ = 586, p <

0.0001; SI Figure 1 B). However, galeal GRN burst structure (Gal1 and Gal2) differed

between stimuli (Figure 1 E, stim*GRN: $F_{1,938}$ = 13.8, p = 0.0002, time: $F_{1,938}$ = 1.42, p =

0.23). OSR elicited a 20% higher burst rate (i.e., Gal2 rate, SI Figure 1 C) and shorter

bursts (i.e., fewer Gal1 spikes per burst, SI Figure 1 D), although the rate of adaptation

of Gal1 did not differ as a function of stimulus (Figure 1 E). To understand if the population

of neurons we recorded from contained information about stimulus identity, we combined

the temporal responses binned at 100 ms intervals over 1 s of recording of all 7 GRNs

across mouthparts using a clustering algorithm (t-SNE and k-means clustering). The

algorithm predicted two statistically separate stimulation groups, one for OSR and one for
sucrose (Figure 1 F).

In spite of the differences observed in GRN responses, the total volume of the 10% OSR
and equimolar sucrose solutions (1.79 M, Figure 1 G) consumed by bumblebees was not
significantly different in the free-feeding assay (Figure 1 H, LME, $F_{1,28}$ = 2.09, p = 0.16).
We also did not observe a difference in the duration of the first feeding bout (Figure 1 I,
LME, $F_{1,26}$ = 0.176, p = 0.678), or the total time the bees spent in contact with the solutions
over 2 min (Figure 1 J, LME, LME, $F_{1,26}$ = 0.555, p = 0.463). We also tested whether the
concentration of these sugar solutions influenced behaviour using a concentration
gradient of OSR and sucrose solutions at 2.0, 1.5, 1.0, 0.5 M and water. A greater volume
of sucrose was consumed for the 1 M and 1.5 M concentrations than for the OSR solution,
but there was no difference at any other concentration (SI Figure 2). First bout and total
contact duration were not significantly different.

Labial palp GRNs may enable bitter detection
QUI is routinely used to test an animal's sensitivity to bitter substances. Here, we tested
a concentration series of QUI in 10% OSR solution to test how GRNs on the labial palps
and galea respond to QUI, for use as a positive control for our tests with pesticides. A 1
mM QUI concentration suppressed all galeal and labial palp GRNs, while 0.1 mM QUI
suppressed the activity of LP$_{III}$2 and LP$_{IV}$2 alone (Figure 2 A, LME, stimulus*GRN $F_{12,1581}$
= 11.3, p < 0.0001, SI Figure 3 A-C). The rate of adaptation, however, of the labial palp
GRNs was faster when they were stimulated with the 0.1 mM QUI solution (Figure 2 B,
stim*time: $F_{1,450}$ = 13.8, p = 0.0002). Galeal GRN burst structure was affected by the
addition of QUI (Figure 2 C, stim*GRN: $F_{2,1116}$ = 308, p < 0.0001, time: $F_{1,1116}$ = 0.062, p
= 0.803; SI Figure 3 B-C). Gal1 adapted faster when stimulated with 0.1 mM QUI
compared to QUI, although there was no effect on Gal2 (Figure 2 C). Clustering of all
GRN responses predicted two groups, with the responses to 1 mM QUI separating from
the responses to 10% OSR and 0.1 mM QUI (Figure 2 D). OSR and 0.1 mM QUI were
not significantly different.

The detection threshold observed in the behaviour experiments was lower than predicted
by the t-SNE clustering but consistent with our measurements of adaptation in labial palp
neurons. Bumblebees consumed significantly less OSR solution when it contained 0.1
mM QUI, while 1 mM QUI was completely deterrent (Figure 2 E, LME, concentration: $F_{2,35}$
= 49.1, p < 0.0001). However, the structure of feeding towards 1 mM QUI was similar to
the results found for the cluster analysis of the GRNs: bees tested with 1 mM QUI
exhibited much shorter average first feeding bout duration (Figure 2 F, LME,
concentration: $F_{2,40}$ = 18.2, p < 0.0001) and shorter average cumulative bout duration
(Figure 2 G, LME, concentration: $F_{2,36}$ = 61.5, p < 0.0001) than those fed with OSR or 0.1
mM QUI solution.
395

Extremely high concentrations of pesticides in OSR nectar reduce the rate of
spiking of GRNs
In our previous work (Kessler et al. (2015), field-relevent concentrations of neonicotinoid
pesticides were not detected by galeal GRNs. Here, we used OSR instead of sucrose
and included sulfoxaflor. We also extended this experiment to test the labial palps. As
before, we found no significant difference in galeal GRN responses between the control
(OSR) solution and all of the solutions containing pesticides at field-relevant
concentrations (SI Figure 4). No significant differences were detected in the responses of
the labial palp neurons to the pesticide solutions (SI Figure 4).
405

To test whether all concentrations of pesticides go undetected by bees, we stimulated
GRNs with 0.1 mM of each pesticide mixed with 10% OSR. Such concentrations are 10-
100k fold greater than has been reported for neonicotinoids from the nectar and pollen of
agricultural crops and are at least 1k fold greater than the LD_{50} values for bumblebees
(Cresswell et al., 2014; Mundy-Heisz et al., 2022; Siviter et al., 2022). Sulfoxaflor,
however, is present at much higher concentrations in the field (e.g., 71.4 µM, U.S.
Environmental Protection Agency (EPA), 2014)). Stimulation with extremely high
concentrations of TMX reduced the average firing rates of $LP_{III}2$ and $LP_{IV}2$; none of the
other pesticides affected the average GRN firing rates (Figure 3 A, LME, stimulus*GRN:
$F_{24,2139}$ = 1.65, p = 0.024.)

We also examined how the temporal response of galeal and labial palp neurons was

influenced by stimulation with extreme concentrations of the pesticides. Labial palp GRNs

adapted faster when IMD or TMX were added to 10% OSR (Figure 3 B, stim: $F_{1,821}$ =

49.9, $p < 0.0001$; time, $F_{1,808}$ = 571, $p < 0.0001$), and firing rates of labial palp GRNs over

1 s were significantly reduced (SI Figure 5 A). Gal1 and Gal2 spiking and adaptation rates

were affected by CLO and IMD (Figure 3 C, stim*GRN: $F_{4,1653}$ = 3.47, $p = 0.0078$; time,

$F_{1,1653}$ = 2.33, $p = 0.127$). Galeal GRN bursting rate was slightly reduced when stimulated

with OSR mixed with CLO or IMD (SI Figure 5 B), while the burst length was not affected

by the addition of pesticides (SI Figure 5 C). We used the same clustering algorithm that

was able to differentiate the responses to sucrose and OSR. Clustering using all of the

data for each of the sensillum types failed to separate the pesticide responses from

responses to the control (10% OSR, Figure 3 D).

Bumblebees readily consume pesticides in nectar

We assessed whether the pesticides were aversive in the range of values that included

field-realistic concentrations (0 - 1000 nM) and one of the extreme values (0.1mM) added

to a 10% OSR solution. Field-relevant concentrations of the pesticides in OSR nectar did

not deter the bees from feeding, as the presence of these compounds did not significantly

alter the amount of food consumed (LME, concentration: $F_{4,297}$ = 0.209, $p = 0.933$), first

bout duration (LME, concentration: $F_{4,290}$ = 0.648, $p = 0.629$) or cumulative bout duration

(LME, concentration: $F_{4,288}$ = 0.310, $p = 0.871$) over the 2 min assay period (Figure 4 A-

C). Importantly, we also tested feeding behaviour using an extreme concentration (0.1

mM) of the pesticides in OSR. Even at this very high concentration, we did not observe a

significant difference in the amount of food consumed (LME: $F_{4,73}$ = 0.872, $p = 0.485$),

first bout duration (LME, $F_{4,74}$ = 0.392, $p = 0.814$), or cumulative bout duration (LME, $F_{4,74}$

= 1.06, $p = 0.382$) over 2 min (Figure 4 D-F).

Discussion and conclusions

For the first time, we report how populations of GRNs on the bee's mouthparts respond to stimulation with putatively bitter compounds. We verified that bees do not have mechanisms on their mouthparts that enable them to taste neonicotinoids or sulfoxaflor when these compounds are present in nectar. Neonicotinoids and sulfoxaflor did not change the bursting of sugar-sensing neurons in the galea, nor did they elicit or inhibit spikes in labial palp neurons. Strikingly, we show that bumblebees do not avoid drinking OSR nectar solutions even when very high concentrations of common pesticides are present. Very high concentrations of these pesticides could produce a small but measurable reduction in the spiking of sugar-sensing GRNs, but these compounds never elicited spikes on their own. What's more, these small changes in the rate of spiking did not translate into features that the clustering algorithm could use to differentiate pesticide laced stimuli from the control OSR solution. From these data, we conclude that the buff-tailed bumblebee's mouthparts do not have mechanisms for the detection and avoidance of common cholinergic pesticides in nectar.

In the present experiments, we expected that if neonicotinoids and sulfoxaflor could be tasted by bees, they would exhibit the same responses in GRNs produced by other non-nutrient compounds (i.e. bitter responses). For this reason, we were mainly testing for the reduction of GRN spiking caused by the suppression of sugar-sensing GRNs, as all of the 'bitter' compounds we tested were presented to the sensilla in OSR nectar (a mixture of primarily fructose and glucose). The known bitter compound, QUI, clearly suppressed the activity of sugar-sensing GRNs in both labial palps and the galea, but it did not measurably alter the burst pattern of firing of the galeal GRNs. Importantly, our new data were able to confirm that field-relevant concentrations of pesticides did not significantly alter the pattern or rate of spiking of galeal or labial palp GRNs towards a nectar solution.

We used a clustering algorithm that integrated input from all the sensilla we recorded from over a 1 s time interval with every stimulus to identify whether population-level input made it possible to differentiate the stimuli. Although we were able to measure small but significant reduction in the responses of both labial palp and galeal neurons caused by

the high concentrations (100 μM) of IMD, SFX, and TMX, this effect was not sufficient for

the clustering algorithm to form classes of distinct stimuli for the OSR-pesticide mixtures.

This is similar to what we observed for 0.1 mM QUI; though a small change in firing was

detected, the clustering algorithm did not differentiate it from OSR. It is worth noting that

the OSR background solutions in both cases were different: in the behavioural

experiments the concentration of OSR was an average value of that naturally found in

nectar. Our physiology experiments were performed using a 10% dilution of this solution.

Different solutions were used because it was not possible to record from neurons with

such a concentrated sugar solution, and in behavioural experiments, most bumblebees

do not find 10% OSR sufficiently phagostimulatory to feed.

Our electrophysiological recordings are the first to report GRN responses from the

bumblebee's labial palps. The labial palps in other insects such as locusts are used to

detect both phagostimulants and deterrant compounds (Blaney and Chapman, 1970), so

we might have rationally expected to observe labial palp GRNs that spike in response to

QUI and cholinergic pesticides, but we did not. This could indicate that on the mouthparts,

bitter compounds are only detected through the suppression of sugar-sensing GRNs (de

Brito Sanchez et al., 2005).

It is interesting to note that in the case of all bitter compounds tested, the output of the

clustering algorithm matched the behaviour of the bees, as the bees did not reject any of

the pesticide solutions or the 0.1 mM QUI solution after the first feeding bout. This could

indicate that that small differences detected at the sensory periphery do not translate into

a sufficient signal across the whole population of gustatory neurons to produce an

aversion to foods containing potential toxins. Bumblebees did drink less overall of the 0.1

mM QUI solution compared to OSR, however the amount of time they spent in contact

with the feeding solution was the same. This suggests that, while 0.1 mM QUI is not

immediately aversive to bumblebees (i.e., by taste), they can integrate negative post-

ingestive feedback within a 2 min feeding period. Compounds reach the hemolymph of

honeybees within 30 s of ingestion (Simcock et al., 2018), and quinine causes malaise-

like symptoms (Ayestaran et al., 2010; Hurst et al., 2014), so it is feasible that the

bumblebees in our behavioural assay factored in post-ingestive feedback while consuming 0.1 mM QUI. Unfortunately, when fed high concentrations of pesticides, post-ingestive feedback did not prevent bumblebees from consuming potentially lethal doses. For example, the LD50 for TMX in *Bombus terrestris* is approximately 6 ng/bee (Siviter et al., 2022), and bumblebees in our assay consumed a median amount of 3.5 ng of TMX when dissolved at 100 uM in OSR.

Our data also indicate that gustatory coding is more than the presence or absence of spikes arising from GRNs tuned to detect particular tastants. In bumblebees, it is clear that the pattern of spiking itself conveys information which is a feature that enables the brain to distinguish gustatory stimuli (Parkinson et al., 2022b). This supports the idea that encoding of gustatory stimuli is accomplished by the whole population of GRNs on a given body part (e.g. the proboscis) as was observed in multichannel recordings from the adult hawkmoth maxillary nerve (Reiter et al. 2015). Thus, activity across the mouthparts GRN population that includes temporal input over time is likely to be used by the brain to differentiate taste stimuli and to facilitate the valuation of food quality. Our new data testing the difference between a nectar stimulus composed of a mixture of sugars (OSR) and sucrose clearly illustrates this. The changes in the rate of bursting and spiking in the galeal and labial palps were sufficient for the clustering algorithm to classify sucrose and OSR as separate stimuli. Although our behavioural assay did not show a difference, we would not expect it to, as it is an assay developed to study food rejection behaviour by bees, not differences in phagostimulation.

Previously, Arce et al. (2018) reported that *B. terrestris* were capable of selectively foraging on 30% w/v sucrose solutions containing TMX. This finding confirmed our previous data that honeybees and bumblebees prefer solutions containing TMX (Kessler et al. 2015). Arce et al (2018) did not observe a preference for or against the solutions containing the neonicotinoids during the initial days of their experiments. We expect that if taste was the mechanism for the preference, the bees would have shown a preference for the TMX solution within the first foraging bout, similar to the time frame of our behavioural assay in this study (2 min). Instead, the authors found that the preference for

the TMX solution developed over time. Similarly, in our previous work, we found that

individual bees developed a preference for solutions containing TMX if they were confined

to feed on a choice between sucrose and sucrose laced with pesticide over longer time

period (i.e., 24 h, Kessler et al., 2015). We concluded that the preference was caused by

the pharmacological action of TMX on cholinergic circuits involved in encoding reward in

the insect brain (Barnstedt et al., 2016), as we found no electrophysiological evidence to

support that bees could taste neonicotinoids (Kessler et al., 2015). This conclusion is

consistent with the expectation that a preference resulting from post-ingestive

reinforcement caused by amplification of the brain's response to sugars would be

expected to develop following long-term, repeated exposure to a pharmacological

compound (Palmer et al., 2013; Wright et al., 2013). If TMX caused bees to associate a

higher 'reward value' with food found in a particular location, they would learn to return to

it in preference over other solutions. Although Arce et al. (2018) switched the feeders

between training and testing phases, this may not have been sufficient to rule out the

possibility that bees could learn the location of the food within a single feeding bout.

Arce et al. (2018) conducted their experiments using gravity feeders in an enclosed arena,

allowing bees to consume solutions *ad libitum*. Since the feeders did not have visual or

olfactory cues, the authors inferred that the bees were capable of tasting the TMX in the

solution. However, gravity feeders do not simulate how bees feed on real flowers. These

feeders have large openings that allow multiple bees to feed simultaneously, enabling

them to contact the solution with all body parts, including the tarsi and antennae. While it

is possible that the sensilla on the antennae or tarsi could detect TMX, it seems unlikely

given that honeybees have limited bitter detection in these locations, only observed when

the sensilla are contacted with concentrations of compounds like QUI that are >1 mM (De

Brito Sanchez et al., 2005; de Brito Sanchez et al., 2014). The field-relevant

concentrations of the pesticides we tested are between 3-6 orders of magnitude lower in

concentration (see Kessler et al. 2015 for review); such dilute concentrations of a bitter

compound present in a highly concentrated sugar solution would be difficult for most

organisms to detect.

Floral morphology such as corolla length has often co-evolved with particular groups of
pollinators with longer mouthparts; a long corolla protects nectar when visitors are
efficient and reliable pollinators. In many flowers, nectar is hidden deep within the corolla
such that the only contact that bumblebees have with it is via the distal end of the
proboscis (also the location where most of the sensilla are). For example, in OSR flowers,
nectaries are located at a depth of 5-8 mm (Cresswell et al., 2001) and the nectar cannot
be contacted otherwise unless the bee bites through the petals. In addition, bees learn to
handle flowers quickly to improve their foraging efficiency (Heinrich, 1976) and even
extend the proboscis in anticipation of entering the corolla prior to landing. Thus, they are
unlikely to contact nectar using gustatory sensilla on their antennae or tarsi while foraging.
For this reason, it is likely that nectar palatability is largely determined by the sensilla on
the proboscis.

Acknowledgements
We would like to thank Priya Chakrabarti-Basu for the pilot studies that led to this work.

Funding
This research was supported by a BBSRC grant (BB/S000402/1) to GA Wright, and an
NSERC PDF (PDF-546125-2020) and Royal Society Newton International Fellowship
(NIF/R1/19368) to RH Parkinson.

Figure legends

Figure 1: Electrophysiological and behavioural responses to sucrose and oilseed
rape (OSR) nectar.
A: Diagram of the bumblebee's mouthparts from which tip-recordings were made,
including the galea, and segments III and IV of the labial palps. Tip-recordings were made
from the longer 'A-type' sensilla.
B: Filtered electrophysiological recordings from galeal (Gal), labial palp segment III (LP_{III}),
and labial palp segment IV (LP_{IV}) sensilla. Spikes from GRN 1 at each location are labeled

with blue circles, GRN 2 spikes in green, and GRN 3 spikes in magenta (only two GRNs

present in labial palp recordings).

C: Average firing rates of GRNs from sensilla on the galea (Gal), segment III of the labial

palps (LP$_{III}$), and segment IV (LP$_{IV}$) over 1 s of stimulation with 10% OSR or an equimolar

(0.173 M) sucrose solution (SUC, n=315 sensilla from 37 bees). Mean and standard

deviation illustrated with black bars, data shown with coloured points. Asterisks represent

significant differences between SUC and OSR (***, p < 0.0001).

D: Linear regression of inter-spike intervals per 0.1 s bin of labial palp GRNs versus time.

Slope of regression is the adaptation rate. Shading is standard error. No significant

difference between the adaptation rate when stimulated with OSR versus SUC.

E: Firing rates of Gal1 versus Gal2 over 1 s stimulation with 10% OSR and SUC. Points

represent mean rate in each 100 ms bin across all trials. A black marker highlights the

first bin (i.e., time = 100 ms). *Post hoc* comparisons showed that Gal2 firing rates were

significantly different between stimuli.

F: t-SNE of all GRN responses for each animal following stimulation with OSR (gray) or

SUC (orange) and k-means clusters (k=2, predicted by Monte Carlo reference-based

consensus clustering) in gray shading.

G: Image of an untethered bumblebee in a holding tube feeding from a capillary in the 2-

min feeding assay.

H: Total volume consumed of 1.79 M SUC or 100% OSR of freely moving bumblebees

during 2 minutes (n = 15 bees per group).

I: The duration of the first feeding bout duration of bumblebees feeding on SUC or OSR.

J: The cumulative duration of feeding bouts within a 2 min period.

Figure 2: Quinine shuts down GRN and behavioural responses to sugars.

A: Average firing rates of 7 GRNs from the galea (Gal), and labial palps (LP$_{III}$, LP$_{IV}$) over

a 1 s stimulations with 10% OSR (OSR), 10% OSR plus 0.1 mM quinine (QUI 0.1), and

10% OSR plus 1 mM quinine (QUI 1). Mean and standard deviation illustrated with black

bars, data shown with coloured points (n = 325 sensilla from 37 bees). Asterisks represent

significant differences between stimulus and OSR (*, <0.05; ** <0.001; ***, <0.0001).

B: Linear regressions of inter-spike intervals per 0.1s bin of labial palp GRNs versus time illustrating the adaptation rate, with SEM in grey shading. The addition of 0.1 mM QUI to OSR significantly affected labial palp adaptation. 1 mM QUI not shown as there was insufficient spiking to calculate adaptation.

C: Firing rates of Gal1 versus Gal2 over 1 s stimulation. Points represent mean rate in each 100 ms bin across all trials. A black marker highlights the first bin (i.e., time = 100 ms). *Post hoc* comparisons showed that only Gal1 (not Gal2) firing rates were significantly different at 0.1 mM QUI, while both were affected by 1.0 mM QUI.

D: t-SNE of all GRN responses for each animal following stimulation with 10% OSR, QUI 0.1 or QUI 1, and k-means clusters (k=2, predicted by Monte Carlo reference-based consensus clustering) in gray shading.

E: Volume consumed by freely-moving bumblebees of 100% OSR, OSR plus 0.1 mM QUI, or OSR plus 1mM QUI over a 2 min period (n = 15 bees/group). Asterisks denote results of Dunnett's test (***, $p < 0.0001$).

F: First bout duration when feeding on OSR, QUI 0.1 or QUI 1 (n = 15 bees/group). Asterisks denote results of Dunnett's test (***, $p < 0.0001$).

G: Cumulative bout duration over a 2 min period of bumblebees feeding on OSR, QUI 0.1 or QUI 1 (n = 15 bees/group). Asterisks denote results of Dunnett's test (***, $p < 0.0001$).

Figure 3: Pesticides affect the temporal code of bumblebee GRNs.

A: Average firing rates of 7 GRNs from the galea (Gal) and labial palps (LP_{III}, LP_{IV}) over a 1 s stimulation with 10% OSR (OSR), or OSR plus 0.1 mM clothianidin (CLO), imidacloprid (IMD), sulfoxaflor (SFX), or thiamethoxam (TMX, n = 331 sensilla from 37 bees). Mean and standard deviation illustrated with black bars, and data shown with coloured points. Asterisks represent significant differences between stimulus and OSR (*, <0.05; ** <0.001; ***, <0.0001).

B: Linear regressions of inter-spike intervals per 0.1s bin of labial palp GRNs versus time illustrating the adaptation rate, with SEM in grey shading. Asterisks represent significant differences between stimulus and OSR (*** $p < 0.0001$).

C: Firing rates of Gal1 versus Gal2 over 1 s stimulation. Points represent mean rate in each 100 ms bin across all trials. A black marker highlights the first bin (i.e., time = 100

ms). *Post hoc* comparisons showed that only Gal1 firing rates were significantly different

for CLO and IMD versus OSR, and Gal2 firing rates were affected by IMD only.

D: t-SNE of all GRN responses for each animal following stimulation with OSR, CLO,

IMD, SFX, or TMX. Monte Carlo reference-based consensus clustering predicted an

optimal cluster count of k=1 on the t-SNE.

Figure 4: Bumblebees do not avoid consuming pesticides in nectar at field relevant

concentrations

A: The total volume consumed by freely moving bumblebees during 2 min of 100% OSR

solutions containing increasing concentrations (nM) of CLO, IMD, TMX and SFX (n = 15

bees per group).

B: First feeding bout duration of OSR mixtures with increasing concentrations (nM) of

pesticides (CLO, IMD, SFX, TMX, n = 15 bees per group).

C: Cumulative feeding duration during 2 min of OSR mixtures with increasing

concentrations (nM) of pesticides (n = 15 bees per group).

D: Total volume consumed of OSR or OSR plus 0.1 mM CLO, IMD, TMX or SFX over 2

min by freely moving bumblebees (n = 15 bees per group).

E: First bout duration of bumblebees feeding on OSR and OSR mixed with 0.1 mM

pesticides (n = 15 bees per group).

F: Cumulative bout duration over 2 min by bees feeding on OSR and OSR mixed with 0.1

mM pesticides (n = 15 bees per group).

References

Arce, A. N., Ramos Rodrigues, A., Yu, J., Colgan, T. J., Wurm, Y. and Gill, R. J. (2018).
Foraging bumblebees acquire a preference for neonicotinoid-treated food with prolonged
exposure. *Proceedings of the Royal Society of London B: Biological Sciences* **285**, 8–11.
Ayestaran, A., Giurfa, M. and de Brito Sanchez, M. G. (2010). Toxic but drank: Gustatory
aversive compounds induce post-ingestional malaise in harnessed honeybees. *PLoS One* **5**,.
Barnstedt, O., Owald, D., Felsenberg, J., Brain, R., Moszynski, J. P., Talbot, C. B., Perrat,
P. N. and Waddell, S. (2016). Memory-Relevant Mushroom Body Output Synapses Are
Cholinergic. *Neuron* **89**, 1237–1247.
Blaney, W. M. and Chapman, R. F. (1970). The functions of the maxillary palps of acrididae
(Orthoptera). *Entomol Exp Appl* **13**, 363–376.

Calderone, N. W. (2012). Insect Pollinated Crops , Insect Pollinators and US Agriculture :
Trend Analysis of Aggregate Data for the Period 1992 – 2009. *PLoS One* **7**, 24–28.
Carruthers, J. M., Cook, S. M., Wright, G. A., Osborne, J. L., Clark, S. J., Swain, J. L. and
Haughton, A. J. (2017). Oilseed rape (Brassica napus) as a resource for farmland insect
pollinators: quantifying floral traits in conventional varieties and breeding systems. *GCB*
Bioenergy **9**, 1370–1379.
Chapman, R. F. and Sword, G. (1993). The importance of palpation in food selection by a
polyphagous grasshopper (Orthoptera: Acrididae). *J Insect Behav* **6**, 79–91.
Cresswell, J. E., Hagen, C. and Woolnough, J. M. (2001). Attributes of individual flowers of
Brassica napus L. are affected by defoliation but not by intraspecific competition. *Ann Bot*
88, 111–117.
Cresswell, J. E., Robert, F. X. L., Florance, H. and Smirnoff, N. (2014). Clearance of
ingested neonicotinoid pesticide (imidacloprid) in honey bees (Apis mellifera) and
bumblebees (Bombus terrestris). *Pest Manag Sci* **70**, 332–337.
De Brito Sanchez, M. G., Giurfa, M., De Paula Mota, T. R. and Gauthier, M. (2005).
Electrophysiological and behavioural characterization of gustatory responses to antennal
"bitter" taste in honeybees. *European Journal of Neuroscience* **22**, 3161–3170.
de Brito Sanchez, M. G., Lorenzo, E., Su, S., Liu, F., Zhan, Y. and Giurfa, M. (2014). The
tarsal taste of honey bees: behavioral and electrophysiological analyses. *Front Behav*
Neurosci **8**, 1–16.
Dethier, V. G. (1976). *The Hungry Fly: A physiological study of the behavior associated with*
feeding. Harvard University Press.
French, A. S., Sellier, M.-J., Ali Agha, M., Guigue, A., Chabaud, M.-A., Reeb, P. D., Mitra,
A., Grau, Y., Soustelle, L. and Marion-Poll, F. (2015). Dual Mechanism for Bitter
Avoidance in Drosophila. *Journal of Neuroscience* **35**, 3990–4004.
Garibaldi, L. A., Steffan-Dewenter, I., Winfree, R., Aizen, M. A., Bommarco, R.,
Cunningham, S. A., Kremen, C. and Carvalheiro, L. G. (2014). Wild Pollinators
Enhance Fruit Set of Crops Regardless of Honey Bee Abundance. *Science (1979)* **339**,
1608–1611.
Gill, R. J., Ramos-Rodriguez, O. and Raine, N. E. (2012). Combined pesticide exposure
severely affects individual- and colony-level traits in bees. *Nature* **491**, 105–8.
Goulson, D., Peat, J., Stout, J. C., Tucker, J., Darvill, B., Derwent, L. C. and Hughes, W. O.
H. (2002). Can alloethism in workers of the bumblebee, bombus terrestris, be explained in
terms of foraging efficiency? *Anim Behav* **64**, 123–130.
Goulson, D., Nicholls, E., Botías, C. and Rotheray, E. L. (2015). Bee declines driven by
combined Stress from parasites, pesticides, and lack of flowers. *Science (1979)* **347**,.
Heinrich, B. (1976). The Foraging Specializations of Individual Bumblebees. *Ecol Monogr* **46**,
105–128.
Henry, M., Béguin, M., Requier, F., Rollin, O., Odoux, J., Aupinel, P., Aptel, J.,
Tchamitchian, S. and Decourtye, A. (2012). A common pesticide decreases foraging
success and survival in honey bees. *Science (1979)* **336**, 348–350.
Huang, J. X. and An, J. D. (2018). Species diversity, pollination application and strategy for
conservation of the bumblebees of China. *Biodiversity Science* **26**, 486–497.
Hurst, V., Stevenson, P. C. and Wright, G. A. (2014). Toxins induce 'malaise' behaviour in
the honeybee (Apis mellifera). *J Comp Physiol A Neuroethol Sens Neural Behav Physiol*
200, 881–890.

John, C. R., Watson, D., Russ, D., Goldmann, K., Ehrenstein, M., Pitzalis, C., Lewis, M. and Barnes, M. (2020). M3C: Monte Carlo reference-based consensus clustering. *Sci Rep* **10**, 1–14.

Kessler, S. C., Tiedeken, E. J., Simcock, K. L., Derveau, S., Mitchell, J., Softley, S., Stout, J. C. and Wright, G. a. (2015). Bees prefer foods containing neonicotinoid pesticides. *Nature* **521**, 74–76.

Klein, A. M., Vaissière, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C. and Tscharntke, T. (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences* **274**, 303–313.

Kuznetsova, A., Brockhoff, P. B. and Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models . *J Stat Softw* **82**,.

Lenth, R., Singmann, H., Love, J., Buerkner, P. and Herve, M. (2018). emmeans: Estimated Marginal Means, aka Least-Squares Means. *R package version*.

Ma, C., Kessler, S., Simpson, A. and Wright, G. (2016). A Novel Behavioral Assay to Investigate Gustatory Responses of Individual, Freely-moving Bumble Bees. *Journal of Visualized Experiments* **113**, e54233.

Meehan, T. D., Werling, B. P., Landis, D. A. and Gratton, C. (2011). Agricultural landscape simplification and insecticide use in the Midwestern United States. *Proceedings of the National Academy of Sciences* **108**, 11500–11505.

Miriyala, A., Kessler, S., Rind, F. C. and Wright, G. A. (2018). Burst Firing in Bee Gustatory Neurons Prevents Adaptation. *Current Biology* **28**, 1585-1594.e3.

Mundy-Heisz, K. A., Prosser, R. S. and Raine, N. E. (2022). Acute oral toxicity and risks of four classes of systemic insecticide to the Common Eastern Bumblebee (Bombus impatiens). *Chemosphere* **295**, 133771.

Muth, F., Gaxiola, R. and Leonard, A. S. (2020). No evidence for neonicotinoid preferences in the bumblebee Bombus impatiens. *R Soc Open Sci* **7**,.

Palmer, M. J., Moffat, C., Saranzewa, N., Harvey, J., Wright, G. a and Connolly, C. N. (2013). Cholinergic pesticides cause mushroom body neuronal inactivation in honeybees. *Nat Commun* **4**, 1634.

Parkinson, R. H., Fecher, C. and Gray, J. R. (2022a). Chronic exposure to insecticides impairs honeybee optomotor behaviour. *Frontiers in Insect Science* **2**, 1–14.

Parkinson, R. H., Kessler, S. C., Scott, J., Simpson, A., Bu, J., Al-Esawy, M., Mahdi, A., Miriyala, A. and Wright, G. A. (2022b). Temporal responses of bumblebee gustatory neurons to sugars. *iScience* **25**,.

Potts, S. G., Imperatriz-Fonseca, V., Ngo, H. T., Aizen, M. A., Biesmeijer, J. C., Breeze, T. D., Dicks, L. V., Garibaldi, L. A., Hill, R., Settele, J., et al. (2016). Safeguarding pollinators and their values to human well-being. *Nature* **540**, 220–229.

Schneider, C. W., Tautz, J., Grünewald, B. and Fuchs, S. (2012a). RFID tracking of sublethal effects of two neonicotinoid insecticides on the foraging behavior of Apis mellifera. *PLoS One* **7**, 1–9.

Schneider, C. A., Rasband, W. S. and Eliceiri, Kevin, W. (2012b). NIH Image to ImageJ: 25 years of Image Analysis. *Nat Methods* **9**, 671–675.

Simcock, N. K., Gray, H., Bouchebti, S. and Wright, G. A. (2018). Appetitive olfactory learning and memory in the honeybee depend on sugar reward identity. *J Insect Physiol* **106**, 71–77.

Siviter, H., Matthews, A. J. and Brown, M. J. F. (2022). A Combined LD50 for
Agrochemicals and Pathogens in Bumblebees (Bombus terrestris [Hymenoptera: Apidae]).
Environ Entomol **51**, 378–384.
Team, R. C. (2016). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.
U.S. Environmental Protection Agency (EPA) (2014). Ecological Risk Assessment for the
Registration Review of Sulfoxaflor. 79.
Van der Sluijs, J. P., Simon-Delso, N., Goulson, D., Maxim, L., Bonmatin, J. M. and
Belzunces, L. P. (2013). Neonicotinoids, bee disorders and the sustainability of pollinator
services. *Curr Opin Environ Sustain* **5**, 293–305.
Vanbergen, A. J., Garratt, M. P., Vanbergen, A. J., Baude, M., Biesmeijer, J. C., Britton,
N. F., Brown, M. J. F., Brown, M., Bryden, J., Budge, G. E., et al. (2013). Threats to an
ecosystem service: Pressures on pollinators. *Front Ecol Environ* **11**, 251–259.
Wäckers, F. L., Alberola, J. S., Garcia-Marí, F. and Pekas, A. (2017). Attract and distract:
Manipulation of a food-mediated protective mutualism enhances natural pest control. *Agric*
Ecosyst Environ **246**, 168–174.
Weiss, L. A., Dahanukar, A., Kwon, J. Y., Banerjee, D. and Carlson, J. R. (2011). The
molecular and cellular basis of bitter taste in Drosophila. *Neuron* **69**, 258–272.
Williamson, S. M., Willis, S. J. and Wright, G. A. (2014). Exposure to neonicotinoids
influences the motor function of adult worker honeybees. *Ecotoxicology* **23**, 1409–1418.
Wood, T. J. and Goulson, D. (2017). The environmental risks of neonicotinoid pesticides: a
review of the evidence post 2013. *Environmental Science and Pollution Research* **24**,
17285–17325.
Wright, G. A. (2016). To feed or not to feed: circuits involved in the control of feeding in
insects. *Curr Opin Neurobiol* **41**, 87–91.
Wright, G. A., Mustard, J. A., Simcock, N. K., Ross-Taylor, A. A. R., McNicholas, L. D.,
Popescu, A. and Marion-Poll, F. (2010). Parallel reinforcement pathways for conditioned
food aversions in the honeybee. *Current Biology* **20**, 2234–2240.
Wright, G. A., Baker, D. D., Palmer, M. J., Stabler, D., Mustard, J. A., Power, E. F.,
Borland, A. M. and Stevenson, P. C. (2013). Caffeine in floral nectar enhances a
pollinator's memory of reward. *Science (1979)* **339**, 1202–1204.
Wright, G. A., Softley, S. and Earnshaw, H. (2015). Low doses of neonicotinoid pesticides in
food rewards impair short-term olfactory memory in foraging-age honeybees. *Sci Rep* **5**, 1–
7.
812

Figure 1

Figure 2

Figure 3

Figure 4