1 **Loss of RdDM-independent CHH methylation is sufficient for disruption of**
2 **chromosome-specific rRNA gene silencing**
3

4 Anastasia McKinlay[1,†], Ramya Enganti[1,†], Navinchandra V. Puppala[2,†], Gargi P.
5 Saradadevi[2,†], Todd Blevins[3] , and Gireesha Mohannath[2*]
6
7 [1]Department of Biology and Department of Molecular and Cellular Biochemistry, Indiana
8 University, Bloomington, IN, United States.
9
10 [2]Department of Biological Sciences, Birla Institute of Technology and Science-Pilani,
11 Hyderabad campus, Hyderabad, Telangana, India.
12
13 [3]Institut de Biologie Moléculaire des Plantes, CNRS, Université de Strasbourg, Strasbourg,
14 France.
15
16 † These authors contributed equally to this work
17
18 Corresponding author:
19 Email: gireesha@hyderabad.bits-pilani.ac.in

Summary
Arabidopsis thaliana has two nucleolus organizer regions (NORs), consisting of long tandem
arrays of 45S rRNA genes on chromosomes 2 (NOR2) and 4 (NOR4). The rRNA gene
subtypes mapping to NOR2 are mostly silenced during development, whereas those
mapping to NOR4 are active. By analyzing DNA methylation-deficient Arabidopsis mutants,
we found that the degree of disruption of NOR2 rRNA gene silencing correlates consistently
with the loss of rDNA cytosine methylation in the CHH sequence context, rather than with

methylation losses in CG or CHG contexts. Comparison of Arabidopsis and tomato NOR
sequences revealed that rRNA gene promoter cytosines mostly occur in the CHH context,
followed by fewer sites in the CHG context. By contrast, almost no CG dinucleotides occur in
plant rRNA gene promoters. This prevalence of CHH sites in plant rDNA promoters suggests
a mechanism to explain the stronger disruption of NOR2 gene silencing in CHH methylation
mutants lacking CHROMOMETHYLASE 2 (CMT2) or DECREASE IN DNA METHYLATION
1 (DDM1) functions as compared to other DNA methylation-deficient mutant lines. Notably,
CMT2 mediates CHH methylation independently of RNA-directed DNA methylation factors
but with a requirement for the chromatin remodeler DDM1. Furthermore, we demonstrate
that NOR2 genes are hypermethylated and exist in more condensed chromatin than NOR4
genes. Collectively, our data define a critical role for CHH methylation in plant rRNA gene
silencing, which is mostly mediated by CG methylation in mammals.
53 **Key words:**
54 Ribosomal RNA (rRNA), rDNA promoter, Nucleolus organizer region (NOR), Gene silencing,
55 DNA cytosine methylation, chromatin
56
Significance Statement

In plants, DNA cytosine methylation occurs in three sequence contexts (CG, CHG, and
CHH), but it was not clear how these distinct contexts contribute to chromosome-specific
large-scale rRNA gene silencing. Using rRNA gene expression, methylation assays, and
bioinformatic analysis of rDNA promoters and methylation patterns, this study shows that
CHH methylation, rather than CG or CHG methylation, plays a critical role in plant rRNA
gene silencing, which is mostly mediated by CG methylation in mammals.
64

Introduction
In eukaryotes, hundreds to thousands of 45S ribosomal RNA (rRNA) genes are arranged as
tandem repeats at chromosomal loci known as nucleolus organizer regions (NORs) (Gerbi,
1986, Long and Dawid, 1980, Viktorovskaya and Schneider, 2015). RNA polymerase I (Pol I)
transcribes rRNA genes to produce 45S pre-rRNA transcripts that are then processed into
18S, 5.8S, and 25-28S rRNAs that comprise the catalytic core of ribosomes, which

synthesize proteins in cells (Saradadevi et al., 2020, Turowski and Tollervey, 2015, Viktorovskaya and Schneider, 2015). Although eukaryotic genomes have many rRNA genes, not all of them are transcribed all the time. Rather, the number of genes transcribed varies during development, dictated by the developmental stage-specific need for ribosomes and protein synthesis (Grummt, 2013, Hannan et al., 2013, Moss et al., 2007). rRNA 'dosage control' involves selective gene silencing rather than selective gene activation (Chen and Pikaard, 1997, Pikaard et al., 2023). In *Arabidopsis thaliana*, rRNA genes are clustered within two NORs located at the northern tips of chromosomes 2 (*NOR2,* which spans ~ 5.9 Mb) and 4 (*NOR4,* which spans ~ 3.9 Mb) (Copenhaver et al., 1995, Copenhaver and Pikaard, 1996b, Copenhaver and Pikaard, 1996a, Fultz et al., 2023). Recent findings in *A. thaliana* demonstrated that the selective rRNA gene silencing is determined by their chromosomal location, not rRNA gene sequence variation (Chandrasekhara et al., 2016, Mohannath et al., 2016). During development, rRNA genes located on *NOR2* are selectively silenced while those on *NOR4* remain active (Chandrasekhara et al., 2016, Mohannath et al., 2016).

Transcriptional gene silencing in eukaryotes often involves cytosine methylation and chromatin condensation (Rhee et al., 2002, Zhang et al., 2006, Zilberman et al., 2007). However, the relative role of cytosine methylation in all three sequence contexts (CG, CHG, and CHH, where H=A, C, or T) in selectively silencing *NOR2* rRNA genes is unknown. In plants, cytosine methylation is deposited by several DNA methyltransferases belonging to different families. METHYLTRANSFERASE 1 (MET1) and CHROMOMETHYLASE 3 (CMT3) mediate virtually all Arabidopsis CG and CHG methylation, respectively (Cokus et al., 2008, Lister et al., 2008, Zemach et al., 2013). CHH methylation is directed by small RNAs and mediated by DOMAINS REARRANGED METHYLTRANSFERASE 2 (DRM2) via the RNA-directed DNA methylation (RdDM) pathway, but also by CHROMOMETHYLASE 2 (CMT2) independently of RdDM (Zemach et al., 2013, Law and Jacobsen, 2010, Rymen et al., 2020, Sasaki et al., 2019). *CMT2* and *CMT3* belong to a group of plant-specific chromomethylases (Bewick et al., 2017). *cmt2-3* mutation is known to reduce most of the CHH methylation (Zemach et al., 2013). In addition to the abovementioned DNA methyltransferases, DECREASE IN DNA METHYLATION 1 (*DDM1*), a SWI2/SNF2 family chromatin remodeler, acts as a master regulator of DNA methylation (Brzeski and Jerzmanowski, 2003, Lee et al., 2023, Osakabe et al., 2021). Loss of DDM1 results in genome-wide demethylation at 70% of methylated cytosines in all three sequence contexts (CG, CHG, and CHH) (Jeddeloh et al., 1999, Vongs et al., 1993, Zemach et al., 2013).

An earlier study investigated the effects of *drm2*, *cmt3,* and *met1* mutants on rRNA gene silencing (Pontvianne et al., 2013). These mutants predominantly show loss of CG or

CHG methylation with minimal to modest loss of CHH methylation (Cokus et al., 2008, Zemach et al., 2013). In this study, we analysed *ddm1* and *cmt2* mutants, which display significant losses in CHH methylation when compared to *drm2*, *cmt3*, and *met1* mutants. We studied these five DNA methylation-deficient mutants to compare the relative effects of differential loss of CG, CHG, and CHH methylation on the disruption of chromosome-specific *NOR2* rRNA gene silencing. In this mutant panel, we show a correlation between the relative loss in *NOR2* rRNA gene silencing and the disruption of cytosine methylation at CHH sites, but not at CG or CHG sites within rDNA loci.

By analyzing the 45S rRNA gene promoters using the recently assembled and published NOR sequences of Arabidopsis (a total of ~950 45S rRNA genes) (Fultz et al., 2023) and tomato (a total of ~1500 45S rRNA genes) (Shirasawa and Ariizumi, 2023), we found that 45S rRNA gene promoter cytosines are mostly in the CHH context, distantly followed by the CHG context and with almost no cytosines in the CG context, providing a putative mechanistic basis for higher *NOR2* derepression observed in *cmt2* and *ddm1* mutants that display higher losses of CHH methylation than other mutants. Lastly, using multiple methylation and chromatin condensation assays, we show that *NOR2* rRNA genes are hypermethylated and exist in more condensed chromatin than *NOR4* genes. In contrast to the key role of CG methylation in mammalian rRNA gene regulation, our study demonstrates a critical role for CHH methylation mediated by CMT2 in plant rRNA gene regulation.

Results

Selective *NOR2* rRNA gene silencing is disrupted in DNA methylation mutants

The rRNA genes in *A. thaliana* are organized into two tandem arrays on NORs, one each on chromosomes 2 (*NOR2*), and 4 (*NOR4*) (Figure 1). In the reference strain Col-0, *NOR2* is 5.5 Mb and *NOR4* is 3.9 Mb, with ~550 and ~390 tandem repeats of 45S rRNA genes, respectively (Fultz et al., 2023) (Figure 1). The 18S, 5.8S, and 25S rRNA sequences are identical among 45S rRNA genes, but variation exists in the internal transcribed spacer (ITS) and the external transcribed spacer (ETS) regions. Four distinct variants, named VAR1 through VAR4 had previously been defined based on the length polymorphism in the 3' ETS region (Pontvianne et al., 2010) (Figure 1B-C). Of these, VAR1, VAR2, and a subset of VAR3 (VAR3a) lack HindIII sites in the 3' ETS region, but the remaining variants (VAR3b,c and VAR4) have a HindIII site (Chandrasekhara et al., 2016, Mohannath et al., 2016) (Fig.1C). During development, the rRNA gene variants that map primarily to *NOR2* (VAR1, VAR3a) are selectively silenced as a consequence of dosage control. In contrast, the

variants that map primarily to *NOR4* (VAR2, VAR3c, and VAR4) remain active (Figure 1D)(Chandrasekhara et al., 2016).

To determine the relative impact of loss of CG, CHG, and CHH methylation on selective *NOR2* rRNA gene silencing, we analysed rRNA variant expression in the mutants defective for DNA methylation, which include *ddm1*, *cmt2*, *met1*, *cmt3*, and *drm2* mutants. In these mutants and the WT-Col-0 strain, we performed RT-PCR analysis using RNA isolated from 4-week-old leaf tissues. VAR1 type rRNA genes constitute ~50% of total 45S rRNA genes, and most of them map to *NOR2*, and undergo selective silencing during development (Chandrasekhara et al., 2016, Pontvianne et al., 2010). Our results showed that among the DNA methylation mutants, VAR1 expression levels were found to be the highest in *cmt2 and ddm1,* followed by *met1* mutants (Figure 2)*.* In contrast, the *cmt3* and *drm2* mutants have displayed low levels of VAR1 expression, close to that of wild-type Col-0 plants (Figure 2).

Using published whole-genome bisulfite sequencing data (Stroud et al., 2013), we generated heatmaps of cytosine methylation for a 45S rRNA reference gene sequence (Figure S2) using Methylpy (Schultz et al., 2015) (Figure 3). The rDNA cytosine methylation heatmaps show the highest loss of CHH methylation in *cmt2*, followed by *ddm1* mutants (Figure 3). Indeed these mutants also show higher *NOR2* gene silencing disruption than the other mutants (Figure 2). Next, *met1* mutants showed loss of CHH methylation and *NOR2* gene silencing disruption, which in degree are lower compared to *cmt2* and *ddm1* mutants (Figs 2 &3). Lastly, among all, *cmt3* and *drm1drm2* mutants showed the least CHH methylation loss at rDNA loci (Figure 3) and the least *NOR2* gene silencing disruption (Figure 2). Interestingly, the minimal loss of rDNA CHH methylation in *drm1drm2,* known RdDM mutants, indicate that CHH methylation at rDNA loci is mostly mediated by RdDM-independent *CMT2*, providing one more basis for the highest *NOR2* gene silencing disruption observed in *cmt2* mutants (Figs. 2&3).

Next, the highest loss of CG methylation at rDNA loci was found in *ddm1* and *met1* mutants and the highest loss of CHG methylation was found in *ddm1* and *cmt3* mutants (Figure 3). Based on the data from Figures 2 and 3, we conclude that the degree of disruption of *NOR2* rRNA gene silencing consistently correlates with the levels of CHH methylation loss, rather than the levels of CG or CHG methylation loss, at rDNA loci. Our data depicting varying loss of CG, CHG, and CHH methylation at rDNA loci in *met1*, *cmt3*, and *drm2* mutants are consistent with findings from a previous study (Cokus et al., 2008). However, *ddm1* and *cmt2* mutants were not previously analyzed for context-specific cytosine methylation loss at rDNA loci or for their effects on disruption of selective *NOR2* rRNA gene silencing. Studying these mutants turned out to be key in establishing a critical role for CHH methylation in the selective *NOR2* rRNA gene silencing.

Next, we wondered why when compared to CHH methylation loss, CG or CHG

methylation loss does not significantly impact *NOR2* rRNA gene silencing despite the *NOR2*

genes showing the highest levels of CG methylation, followed by CHG methylation (Fultz et

al., 2023, Mo et al., 2023). To address this question, we first analyzed the distribution of

cytosines in CG, CHG, and CHH sequence contexts in the rRNA gene promoter regions of

the recently assembled and published NOR sequences in *A. thaliana* (Col-0 strain),

constituting a total of ~ 950 copies of 45S rRNA genes (Doelling and Pikaard, 1995, Fultz et

al., 2023). In Arabidopsis, -55 to +6 region with respect to the transcription start site (+1) in

the 45S rRNA genes constitutes the gene promoter region (Doelling et al., 1993, Doelling

and Pikaard, 1995). () First, we found that the gene promoter has eight cytosines but

strikingly, none of them exist in CG context. Six (75%) exist in CHH context and the

remaining two (25%) exist in CHG context (Figure 4). Next, we found that the gene promoter

region is highly conserved across most of the *A. thaliana* 45S rRNA genes (Fultz et al.,

2023). Of 894 total 45S rRNA gene repeats combinedly present on *NOR2* and *NOR4*, 859

repeats (96%) showed 100% homology to the gene promoter sequence (-55 to +6) (Figure

4B, 5[th] column). The remaining repeats showed one to few mismatches. Further, we

extended our analysis to -300 bp to include the upstream promoter region (Pontvianne et al.,

2013), which had showed only a small positive effect on transcription of 45S rRNA genes

(Doelling et al., 1993, Doelling and Pikaard, 1995), and these upstream sequences are not

conserved among 45S rRNA genes (Fultz et al., 2023). Among the 51 cytosines found in this

region, again, predominant occurrence of cytosines was found in the CHH context (41

cytosines; ~80%), distantly followed by the CHG (6 cytosines; ~12%) and the CG (4

cytosines; ~8%) contexts (Figure S3-A).

Similarly, we analysed gene promoter regions in the recently assembled tomato

NOR2 constituting ~1500 45S rRNA genes (Shirasawa and Ariizumi, 2023) and the

previously published tomato 45S rRNA gene promoter sequence (Doelling and Pikaard,

1995). First, we found that the gene promoter has 12 cytosines, of which 10 (~83%) exist in

the CHH context, 2 (~17%) the CHG context, and none in the CG context (Figure 4)

(Doelling and Pikaard, 1995, Shirasawa and Ariizumi, 2023). We also found the gene

promoter sequences to be highly conserved across the tomato *NOR2*. Of 1500 total 45S

rRNA gene repeats present on *NOR*2, 1473 repeats (98%) showed 100% homology to the

gene promoter sequence (-55 to +6) (Figure 4B, 5[th] column) (Shirasawa and Ariizumi, 2023)

and the remaining repeats showed one to few mismatches. The results also appear to be

similar in other plant species (Figure 4) (Doelling and Pikaard, 1995). Collectively, the rDNA

promoter analyses across different plant species provide a putative mechanistic basis for

maximum disruption of *NOR2* rRNA gene silencing observed in the *cmt2* and *ddm1* mutants

that displayed the highest loss of CHH methylation. Overall, these results define a novel critical role for CHH methylation in 45S rRNA gene regulation in plants.

In contrast, when we analyzed known human rDNA promoter sequences (Zhou et al., 2016), we found that out of 65 cytosines, 26 are in CG (~40%; the highest among the three), 19 in CHG (~29%), and 20 in CHH (~31%) contexts (Figure S3.B). These results are consistent with the mammalian rRNA gene regulation, which is almost entirely dictated by CG methylation (Bernstein et al., 2007, Priyadarshini et al., 2023, Zhou et al., 2016). These results indicate the presence of evolutionarily divergent mechanisms of rRNA gene regulation between plants and mammals.

NOR2 rRNA genes are hypermethylated while NOR4 rRNA genes are hypomethylated in Col-0

We carried out methylation analysis of rRNA variants using HpaII-digested genomic DNA (gDNA) of Col-0 and *met1* mutants (Figure 5). Because the rRNA variant-defining 3' ETS region lacks HpaII sites, HpaII digestion of gDNA does not impact PCR amplification of this region. Therefore, we chose to separate HpaII-digested and HpaII-undigested fractions of gDNA by resolving them on an agarose gel. The fragments that remained at the top of the gel (>20 kb) constituted the HpaII-undigested fractions while those resolved at 0.6-1.0 kb range constituted the HpaII-digested fractions (Figure 5A, left panel). Both the digested and the undigested fractions were gel-eluted, followed by PCR amplification of rRNA variants. Through this analysis, we were able to identify the rRNA variants that were enriched in HpaII-digested or undigested fractions. The PCR data revealed VAR2 genes from *NOR4* to be the most predominant in Col-0 among the HpaII-digested products, indicating relative hypomethylation of *NOR4* genes compared to *NOR2*-localized VAR1 genes (Figure 5A, lanes 1 and 2, right panel). However, in *met1* mutants, used as a control, all rRNA variants were amplified proportionate to their composition due to loss of cytosine methylation in all of them (Figure 5A, lanes 3 and 4, right panel,).

In another experiment, we subjected gDNA to digestion by McrBC, an enzyme that preferentially digests DNA if its recognition site ($R^mC(N_{40-3000})R^mC$) consists of methylated cytosines. McrBC digestion of gDNA followed by PCR amplification of rRNA variants revealed that VAR1 was digested the most and VAR2 the least (Figure 5B), indicating that *NOR2* genes are hypermethylated in contrast to *NOR4* genes, consistent with recent findings in Arabidopsis which employed sequencing-based methylation analysis of rRNA variants (Fultz et al., 2023, Mo et al., 2023).

NOR2 rRNA genes are hypermethylated while NOR4 rRNA genes are hypomethylated in the introgression line Col$^{Sf\text{-}NOR4}$

In a previous study, we showed that an introgression line named Col$^{Sf\text{-}NOR4}$ consists of *NOR2* from Col-0 (VAR1 and VAR3 genes) and *NOR4* from Sf-2 ecotype (Chandrasekhara et al., 2016) (Figure 6A). Sf-2 has only VAR1 type of rRNA genes, but they can be differentiated from VAR1 genes of Col-0 by HindIII digestion. Sf-2 VAR1 genes have a HindIII site in their 3' ETS region, while Col-0 VAR1 genes lack it (Chandrasekhara et al., 2016). In Col$^{Sf\text{-}NOR4}$, Sf-2 *NOR4* genes are active though they are similar in sequence to the silenced Col-0 *NOR2* (VAR1) genes (Chandrasekhara et al., 2016). However, Col-0 *NOR2* genes are still selectively silenced in Col$^{Sf\text{-}NOR4}$ (Chandrasekhara et al., 2016). Therefore, we analyzed this introgression line to see if VAR1 type of genes residing in *NOR2* and *NOR4* of Col$^{Sf\text{-}NOR4}$ are differentially methylated or not, using similar methylation assays, as described above. In one experiment, gDNA of Col$^{Sf\text{-}NOR4}$ was subjected to HpaII digestion, followed by gel elutions of the undigested and the digested fractions. The gel-eluted DNA samples were then subjected to PCR amplification of rRNA variants followed by HindIII digestion. In control samples (no HpaII), HindIII digestion of PCR-amplified rRNA variants resulted in the expected products, which include digested Sf2-VAR1 genes, undigested Col-0 VAR1 genes, and partially digested Col-0 VAR3 genes (Figure 6B, lanes 1 and 2, top right gel image, and Figure 6C) (Chandrasekhara et al., 2016). However, HindIII digestion of the PCR products obtained from the gel-purified HpaII-digested fraction as the template, revealed that Col-0 VAR1 genes predominantly constitute the HpaII-undigested fraction (Figure 6B, lanes 3 and 4, top right gel image, and Figure 6C). On the other hand, we observed a significant increase in the Sf-2 VAR1 genes in the HpaII-digested fraction (Figure 6B, lanes 3 and 4, bottom right gel image, and Figure 6C) compared to the undigested fraction (Figure 6B, lanes 3 and 4, top right gel image, and Figure 6C). These data indicate hypermethylation of Col-0 *NOR2* genes and hypomethylation of Sf-2 *NOR4* genes, despite the presence of the VAR1 type of genes on both the NORs in the Col$^{Sf\text{-}NOR4}$ introgression line.

In another experiment, gDNA of Col$^{Sf\text{-}NOR4}$ was first digested with McrBC, followed by PCR amplification of rRNA variants. These PCR products were then digested with HindIII and resolved using regular gel electrophoresis. HindIII-digested control PCR products (no McrBC digestion) behaved as expected based on the presence or the absence of HindIII sites (Figure 6D, E, lanes 1 and 2). However, the McrBC digestion patterns of the rRNA variants (Figure 6D, E) were the opposite of the experimental results involving HpaII digestion (compare to Figure 6B, C). Predominantly, the Sf-2 VAR1 genes constituted the McrBC-undigested fraction, while the Col-0 VAR1 genes constituted the McrBC-digested fraction (Figure 6D, E, lanes 3 and 4). We also observed complete digestion of the Col-0

VAR3 (from *NOR2*) genes by McrBC (Figure 6D, lane 3). These data again demonstrate that

NOR4 genes are hypomethylated and *NOR2* genes are hypermethylated in Col^Sf-NOR4.

In Figures 4 and 5, resistance to HpaII digestion indicates the presence of CG and

CHG methylation while digestion by McrBC indicates the prevalence of CG, CHG, and CHH

methylation. Based on these data we conclude that *NOR2* genes are hypermethylated than

NOR4 genes, in all cytosine contexts.

NOR2 rRNA genes exist in more condensed chromatin than NOR4 rRNA genes

To investigate the chromatin accessibility of *NOR2* and *NOR4* genes, we carried out Tn5

transposase-mediated transposition using nuclei isolated from 3-week-old Col-0 plants. Tn5

is a prokaryotic hyperactive transposase that has been developed as a tool to insert short

DNA sequences into open chromatin regions (Adey et al., 2010, Goryshin and Reznikoff,

1998, Buenrostro et al., 2013). A pair of 19 bp IS50 repeats that serve as 'outside ends'

(OEs) sequences are sufficient for recognition by Tn5 transposase and transposition of OEs-

containing sequences into open chromatin (Ahmed, 2009). We designed short DNA

sequences consisting of OEs for Tn5 recognition and used them in a native chromatin

transposition assay. Purified Col-0 nuclei were incubated with or without 5 pmol oligos

containing Tn5 recognition sites for transposition, followed by DNA extraction from the nuclei

and PCR analysis of rRNA variants. The PCR data did not reveal differences between *NOR2*

and *NOR4* genes. We then tried an indirect approach to focus on the chromatin accessibility

state of the promoter regions. In this approach, the DNA samples isolated from Tn5

transposase-treated nuclei were subjected to two rounds of PCR amplification. The first

round of PCR (PCR1) amplifies ~2.8 kb region that spans the 3' ETS region, IGS, and the

promoter of the following 45S rRNA gene repeat (Figure 7 A-B). Due to the lack of variant-

specific sequence variation in the promoter region, we exploited the proximity of the rRNA

variant-defining 3' ETS region of one rRNA gene to the promoter of the following rRNA gene.

The occurrence of rRNA genes as tandem repeats (Figure 1A) at NORs permitted this

approach. The idea is that, if promoters of VAR1 genes (primarily present on *NOR2*) are

more condensed than the promoters of VAR2 genes (primarily present on *NOR4*), then the

2.8 kb PCR products should be enriched for VAR1 genes. Because the 2.8 kb PCR products

obtained during PCR1 (using primers 5'allvar and p34) cannot discriminate between *NOR2*

and *NOR4* genes, they were resolved on a 1% agarose gel followed by gel-purification of 2.8

kb amplicons. These amplicons were then subjected to the second round of PCR

amplification (PCR2) with primers that either amplified only VAR1 and VAR2 genes (Figure

7C) or all four rRNA variants (Figure 7D). The results showed severe depletion of *NOR4*-

specific VAR2 genes in the reactions with 5 pmol of Tn5 oligos (Figure 7C, D), indicating

depletion of *NOR4*-specific fragments in the PCR1 amplicon pool. This depletion is presumably due to higher transposition rates in the promoter regions of *NOR4*-specific rRNA genes compared to the *NOR2* genes. Collectively, these data indicate that *NOR4* genes are predominantly present in a more accessible open chromatin state than *NOR2* genes, consistent with recent findings from a chromatin condensation study in Arabidopsis (Mo et al., 2023).

It is still not clear why *NOR2* rRNA genes selectively undergo hypermethylation and chromatin condensation. Methylation maps of centromere-proximal regions of NORs show that ~ 75 kb of the *NOR2*-adjoining region, composed of transposable elements and transposon remnants, is heavily methylated in all cytosine contexts compared to the *NOR4*-adjoining region (Figure S4), which has a protein-coding gene just 3 kb away (Lewis and Pikaard, 2001) and do not contain any stretch of transposons or their remnants thereafter. In plants, transposons are silenced through DNA methylation-mediated mechanisms. Could such silencing of the *NOR2*-adjoining transposon-rich region spread to *NOR2* rRNA genes, resulting in their selective silencing? The question remains to be answered.

Discussion

To know the relative role of cytosine methylation in each sequence context (CG, CHG, and CHH) in *NOR2* rRNA gene silencing, we analyzed various DNA methylation-deficient mutants of Arabidopsis that display loss of CG, CHG, and CHH methylation to varying degrees. In Arabidopsis, CHH methylation is mainly determined by RdDM-dependent DRM2 and RdDM-independent CMT2 (Rhee et al., 2002, Stroud et al., 2013, Zemach et al., 2013). Between these two, CMT2 appears to be responsible for most of the CHH methylation at rDNA loci (Figure 3). Interestingly, cytosines in plant rRNA gene promoters, including those of Arabidopsis predominantly occur in the CHH context followed by a few in the CHG context and almost none in the CG context (Figure 4 and Figure S3-A). These data potentially explain why *cmt2* mutants displayed the highest *NOR2* gene silencing disruption, despite showing minimal loss of CG or CHG methylation (Figs. 2&3). Therefore, we propose to rule out a significant role for CG or CHG methylation in rRNA gene silencing.

CMT2 and CMT3, methylate DNA through the dual binding of H3K9 di-methylation (H3K9me2)-containing nucleosomes via their bromo adjacent homology (BAH) and Chromo domains (Du et al., 2012). Interestingly, it has been found that H3K9 methyltransferases are required for rRNA variant-specific silencing in *A. thaliana* (Pontvianne et al., 2012). Therefore, it is likely that nucleosomes in *NOR2* rRNA gene promoters undergo H3K9 dimethylation followed by CHG and CHH methylation, resulting in heterochromatinization and silencing. However, the loss of CHH methylation at rDNA loci has more impact on *NOR2*

derepression than the loss of CHG methylation presumably due to the predominant occurrence of cytosines in the CHH context than the CHG context in the rRNA gene promoters (Figs. 2-4).

Next, because DDM1 mediates CHH methylation by CMT2 (Zemach et al., 2013), it is reasonable to find levels of CHH methylation loss and *NOR2* gene silencing in *ddm1* mutants comparable to that of *cmt2* mutants (Figs. 2&3). Recent findings have shown that DDM1 deposits histone variant H2A.W, which marks plant heterochromatin, onto potentially mobile transposable elements (TEs) resulting in their transcriptional gene silencing (Osakabe et al., 2021). It remains to be determined if a similar mechanism involving DDM1-mediated histone H2A.W deposition is involved in *NOR2* rRNA gene silencing.

Due to apparent lack of a significant role of CG or CHG methylation in rRNA gene silencing (Figs. 2&3), we surmise that the *NOR2* gene silencing disruption observed in *ddm1* and *met1* mutants (Figure 2) is primarily due to their loss of CHH methylation at rDNA loci than due to their loss of CG or CHG methylation (Figure 3). Overall, the data establish a novel critical role for CMT2-mediated RdDM-independent CHH methylation in plant rRNA gene regulation, as shown in our proposed model in Figure 8.

It is interesting to note that knocking down DNA methyltransferases MET1 or CMT3, responsible for maintaining methylation in CG and CHG contexts respectively, did not affect nucleolar dominance in tetraploid species *Arabidopsis suecica,* in which rRNA genes from one of the progenitor species *A. thaliana* are silenced during development, while those from the other progenitor species *A. arenosa* remain active (Costa-Nunes et al., 2010). In contrast, knocking down DRM2 DNA methyltransferase (responsible for most of the RdDM-dependent CHH methylation) disrupted nucleolar dominance via release of silencing of *A. thaliana* rRNA genes (Costa-Nunes et al., 2010). Therefore, we predict that the loss of CMT2, responsible for most of the RdDM-independent CHH methylation, would severely impact nucleolar dominance in *A. suecica*, but this prediction needs to be tested.

In a similar study in plant species *Tragopogon porrifolius*, the methylation dynamics of cytosines in CWG sequence contexts (where W= A or T) has been suggested to be most responsible for regulating silent and active rDNA states (Matyášek et al., 2024). It would be interesting to know if similar conservation and enrichment of cytosines in the CWG context among the rDNA gene promoters exist in *Tragopogon porrifolius*. Such a cytosine enrichment would indicate the possibility of involvement of different plant-specific chromomethyltransferases (CMTs) in different plant species. Nevertheless, several findings in multiple plant species collectively underscore the importance of non-CG methylation in plant rRNA gene silencing.

In contrast to plant species, CG methylation is known to play a prominent role in mammalian rRNA gene regulation. For example, human rDNA promoters are mostly methylated in the CG context and their dysregulation has been linked to several types of cancers (Priyadarshini et al., 2023, Bacalini et al., 2014, Ghoshal et al., 2004, Karahan et al., 2015, Raval et al., 2012, Zhou et al., 2016). Moreover, cytosines occur predominantly in the CG context in the human rRNA gene promoters (Figure S3-B). CHG and CHH methylation are largely absent in mammals, but they occur commonly in plant genomes, carried out predominantly by plant-specific chromomethylases (Bernstein et al., 2007, Bewick et al., 2017, Henderson and Jacobsen, 2007). Further, it is to be noted that, spacer promoters in animal models (e.g., frog species *Xenopus laevis*) significantly contribute to the enhancement of rRNA gene transcription (De Winter and Moss, 1986). In contrast, spacer promoters have minimal contribution, if any, to rRNA gene transcription, in plant species (e.g. *Arabidopsis thaliana*) (Doelling et al., 1993). These findings indicate the evolution of divergent rRNA gene regulation mechanisms in plants and animals.

The silent *NOR2* genes are hypermethylated in all cytosine contexts (CG, CHG, and CHH) and exist in more condensed chromatin state than the active *NOR4* rRNA genes (Figs. 5-7) (Fultz et al., 2023, Mo et al., 2023). However, despite the striking similarities in rRNA gene sequences between *NOR2* and *NOR4*, why rRNA genes on *NOR2* are selectively hypermethylated and silenced? The question remains to be answered.

Materials and Methods

Plant material and growth

A. thaliana ecotype Col-0 (stock no. CS22681), and mutants *met1-7* (stock no. SALK_076522), *cmt2-3* (stock no. SALK_012874C), and *cmt3-11t* (stock no. SALK_148381), were obtained from the Arabidopsis Biological Resource Center (ABRC; http://abrc.osu.edu). *drm2-2b* (stock no. SAIL_70_E12) and Col[Sf-NOR4] (Chandrasekhara et al., 2016, Lee and Amasino, 1995) were provided by Andrzej Wierzbicki and Scott Michaels, respectively, and *ddm1-2* was provided by Eric Richards. Plants were grown in soil in a plant growth room under a 16-h light/8-h dark photoperiod cycle with 50–55% humidity and temperature of 23–25°C.

RNA isolation

Four-week-old leaf tissue of individual plants was frozen in liquid nitrogen and ground into a fine powder. RNeasy Plant Mini Kit (Qiagen) was used to extract total RNA from ground powder. Total RNA was then treated with Turbo DNA-free kit (Invitrogen) to eliminate contaminating DNA.

PCR and RT-PCR assays

For PCR of genomic DNA to amplify all rRNA variants, PCR amplification was conducted using 22-30 cycles (depending on the source of DNA) of 30 sec at 95°C, 30 sec at 55°C–60°C, and 60 sec at 72°C. For RT–PCR, the reverse transcription reaction was performed using 1 µg of total RNA using SuperScript III reverse transcriptase (Invitrogen). One microliter (~100 ng) of reverse transcription product was then used for PCR amplification of rRNA variable regions or actin gene control using 30 cycles of 30 sec at 94°C, 30 sec at 55°C, and 60 sec at 72°C. In all the PCR or RT-PCR reactions, primers 5'allvar and 3'allvar were used to amplify rRNA variable regions (all four variant types). PCR and RT–PCR products were resolved by electrophoresis on 2%–2.5% agarose gels in TBE buffer. Sequences of primers used in this study are given in Figure S1.

Generation of DNA methylation heat maps

Whole genome bisulfite sequencing data (GSE39901) published by Stroud et al. (2013)(Stroud et al., 2013), was used to assess the methylation state of rDNA in various methylation mutants. The sequencing data were analysed using Methylpy (Schultz et al., 2015). The reads were mapped to a converted 45S rDNA reference sequence allowing a single mismatch. The methylation state of all cytosines in CG, CHG, and CHH contexts was then analyzed and represented as a fraction of methylated cytosines over the total cytosines in each context. Replicates for a given genotype were merged into a single file.

DNA methylation map of the NOR2 and NOR4-adjacent regions

DNA methylation profiles of the *NOR2* and *NOR4*-adjacent regions on *A. thaliana* Col-0 chromosomes 2 and 4 were compared using whole genome bisulfite sequencing data published by Stroud et al. (2013) (GEO accession# GSM980986). In short, the WT_rep2_CG.wig.gz, WT_rep2_CHG.wig.gz and WT_rep2_CHH.wig.gz data tracks available in Wiggle format at https://www.ncbi.nlm.nih.gov/geo/ were used to display each 150 kb, centromere-proximal NOR-adjacent region in a local JBrowse instance with TAIR10 gene annotations and TE annotations of Buisine et al. (2008)

DNA isolation from leaf tissue and agarose gel pieces (gel elution).

Genomic DNA (gDNA) was isolated from the leaf tissue of 4-week-old plants by using Illustra Nucleon Phytopure extraction kit reagents (GE Healthcare). This gDNA was used for HindIII CAPS assay (Figure 1C), PCR of rRNA variants (Figure 2, topmost panel), methylation analysis of gDNA (Figure 5A left panel, Figure 5B, and Figure 6B, left panel). In some cases, following gDNA PCR or digestion by methylation-sensitive enzymes, the products were

resolved on an agarose gel and the DNA fragments were gel-purified using QIAquick gel
extraction kit (Qiagen). This gel-purified DNA was subsequently used as a template for
downstream PCR amplification of rRNA variants (Figure 5B right panel, Figure 7C) or HindIII
CAPS assay (Figure 6B right panel). DNA from the EZ-Tn5 transposase-treated nuclei was
isolated using the cetyl trimethyl ammonium bromide (CTAB) method.

Nuclei isolation

Nuclei were isolated using 10 g tissue from 3-week-old Col-0 plant seedlings as described
previously (Gaudino and Pikaard, 1997) with a few modifications. Briefly, 3-week-old plants
were harvested and washed in ice-cold water. Then the plants were submerged in cold
diethyl ether for 5 min and washed twice in ice-cold water. Homogenization of plant material
was performed in 100 mL of nuclei isolation buffer (1M sucrose, 50 mM Tris-HCl pH 7.2, 5
mM $MgCl_2$, 5 mM KCl, 10 mM 2-mercaptoethanol, 0.2 mM PMSF) with a motorized
homogenizer (Fisher Scientific Powergen 700). Homogenate was filtered through 1 layer of
Miracloth and 1 layer of nylon mesh (pore size of 50 μm). After centrifugation of the filtered
homogenate at 14,000 g for 15 min at 4°C the pellet was resuspended in 10 ml of nuclei
isolation buffer and prepared for loading into discontinuous Percoll gradient (15%, 30%, 45%,
and 60%) by adding one volume of "100% Percoll solution" (34.23 g sucrose, 5 mL 1 M Tris-
HCl pH 7.2, 0.5 mL 1 M $MgCl_2$, 0.5 mL 1 M KCl, 34 μL 2-mercaptoethanol, 100 μL 2 mM
PMSF and Percoll to 100 mL) to 19 volumes of nuclei. Gradients were centrifuged at 2,000
rpm for 10 min, then 8,000 rpm for 20 min. Nuclei were harvested from 30%/45% and
45%/60% boundaries and pooled. Nuclei were diluted with 5 volumes of nuclei isolation
buffer, mixed by inversion, and collected by centrifugation at 1,500 g for 10 min at 4°C. A
second wash of nuclei was performed with 25 volumes of nuclei isolation buffer. The final
wash was done with 50 mL of Nuclease digestion buffer (50 mM Tris-HCl, pH 8.0, 0.3 M
sucrose, 5 mM $MgCl2$, 1.5 mM NaCl, 0.1 mM $CaCl2$, 5 mM 2-mercaptoethanol). The
resulting pellet was resuspended in 2 mL of Nuclease digestion buffer.

Transposase-accessible chromatin assay

The assay was carried out based on the methodology described by Buenrostro et al.
(Buenrostro et al., 2013). Oligos (5'-Ez-Tn5 and 3'-Ez-Tn5) were annealed by heating a
mixture of equal volumes of 100 mM oligos at 95 °C for 5 min, followed by turning the
heating off and incubating for 10 min. Transposition reactions were carried out using 2 μL of
purified nuclei, 20 μL of EZ-Tn5 transposase with varying amounts of the oligos 5'-Ez-Tn5
and 3'-Ez-Tn5, according to the manufacturer's instructions (Epicentre). Incubation was

performed at 37°C for 90 min. DNA was isolated using the cetyl trimethyl ammonium
bromide (CTAB) method. PCR1 and PCR2 were carried out as described above. The
sequences of primers and oligos used in this study are listed in Figure S1.
502
503

Acknowledgements
This work is supported by:
Birla Institute of Science and Technology (BITS) Pilani, Hyderabad campus through a
research grant to G.M. (BITS/GAU/ACRG/2019/H0576) and an Institute Fellowship to G.P.S.
Science and Engineering Research Board (SERB), Government of India through Ramanujan
Fellowship (SB-S2-RJN-062-2017) and CRG (CRG/2020/002855) research grants to GM,
and internship to N.V.P.
Department of Biotechnology through research grant (BT/PR38410/GET/119/310/2020) to
GM.
A.M. and R.E. were supported by funds to Craig S. Pikaard. as an investigator of the Howard
Hughes Medical Institute.
T.B. was supported by the ANR (ANR-17-CE20-0004), and by the Interdisciplinary Thematic
Institute IMCBio (ITI 2021-2028 program), including funds from IdEx University of Strasbourg
(ANR-10-IDEX-0002), SFRI-STRAT'US (ANR 20-SFRI-0012) and EUR IMCBio (ANR-17-
EURE-0023) in the framework of the French Investments for the Future Program.
519
520 We thank the members of the Pikaard lab and the Mohannath lab for their critique of the
521 manuscript.

523 **Competing interests**
524 The authors declare no competing interests.
525

526 **Author contributions**
527 A.M., R.E., N.V.P, G.P.S., and G.M. planned and designed the research; A.M., R.E., N.V.P.,
528 G.P.S., T.B., and G.M. performed the research; A.M., R.E., N.V.P, G.P.S., and G.M.
529 analysed the data; and A.M., R.E., N.V.P, G.P.S., and G.M. wrote the paper.
530

Data availability
For this article we analyzed the publicly available/published data, and the details are as
follows: The whole genome bisulfite sequencing data of *Arabidopsis thaliana* (WT and
mutants) were published by Stroud et al. (2013) (GEO accession# GSM980986). NOR-

adjacent methylation data are available at https://www.ncbi.nlm.nih.gov/geo/. The fully
assembled NOR sequences of *Arabidopsis thaliana* (strain Col-0) were published by Fultz et
al. (2023) and the sequences are available at GenBank as accessions OR453402
(https://ncbi.nlm.nih.gov/nuccore/OR453402)   for   *NOR2*   and   OR453401
(https://ncbi.nlm.nih.gov/nuccore/OR453401)   for   *NOR4*.   Assembled   whole   genome
sequences of tomato (*Solanum lycopersicum*, cultivar Micro-Tom) have been published on a
preprint server Shirasawa and Ariizumi (2023) (https://doi.org/10.1101/2023.10.26.564283)
and the sequences are available at DDBJ (accession numbers AP028935–1 AP028946) and
KaTomicsDB (https://www.kazusa.or.jp/ tomato).
544

545   **Supplementary figure legends**

546   **Figure S1:** Primers/oligos used in this study.
547

548   **Figure S2:** The 45S rRNA gene reference sequence used to generate methylation heat
549   maps.
550

551   **Figure S3:** Occurrence of cytosines in the promoter regions of 45S rRNA genes.
552   Diagram depicting relative distribution of cytosines in CG, CHG, and CHH contexts in the
553   promoter regions of 45S rRNA genes in (A) Arabidopsis thaliana, and in (B) humans.
554

Figure S4: Centromere-proximal NOR2-adjoining region is hyper methylated compared to
the corresponding NOR4-adjoining region. Map of cytosine methylation in all sequence
contexts (CG, CHG and CHH) obtained for the wild-type plants (Col-0) using deep bisulfite
sequencing data. Methylation data for about 150 Kb region right next to the NORs on the
centromere-proximal side has been depicted. Height of the bars for each cytosine residue
indicates depth of the methylation at that residue. Annotated loci in the 150 Kb region are
shown above the methylation data.
562

References
ADEY, A., MORRISON, H. G., ASAN, XUN, X., KITZMAN, J. O., TURNER, E. H., STACKHOUSE, B.,
MACKENZIE, A. P., CARUCCIO, N. C., ZHANG, X. & SHENDURE, J. 2010. Rapid, low-input, low-
bias construction of shotgun fragment libraries by high-density in vitro transposition.
Genome Biol, 11**,** R119.
AHMED, A. 2009. Alternative mechanisms for tn5 transposition. *PLoS Genet,* 5**,** e1000619.

BACALINI, M. G., PACILLI, A., GIULIANI, C., PENZO, M., TRERÉ, D., PIRAZZINI, C., SALVIOLI, S., FRANCESCHI, C., MONTANARO, L. & GARAGNANI, P. 2014. The nucleolar size is associated to the methylation status of ribosomal DNA in breast carcinomas. *BMC Cancer,* 14**,** 361.

BERNSTEIN, B. E., MEISSNER, A. & LANDER, E. S. 2007. The mammalian epigenome. *Cell,* 128**,** 669-81.

BEWICK, A. J., NIEDERHUTH, C. E., JI, L., ROHR, N. A., GRIFFIN, P. T., LEEBENS-MACK, J. & SCHMITZ, R. J. 2017. The evolution of CHROMOMETHYLASES and gene body DNA methylation in plants. *Genome Biol,* 18**,** 65.

BRZESKI, J. & JERZMANOWSKI, A. 2003. Deficient in DNA methylation 1 (DDM1) defines a novel family of chromatin-remodeling factors. *J Biol Chem,* 278**,** 823-8.

BUENROSTRO, J. D., GIRESI, P. G., ZABA, L. C., CHANG, H. Y. & GREENLEAF, W. J. 2013. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat Methods,* 10**,** 1213-8.

BUISINE, N., QUESNEVILLE, H. & COLOT, V. 2008. Improved detection and annotation of transposable elements in sequenced genomes using multiple reference sequence sets. *Genomics,* 91**,** 467-75.

CHANDRASEKHARA, C., MOHANNATH, G., BLEVINS, T., PONTVIANNE, F. & PIKAARD, C. S. 2016. Chromosome-specific NOR inactivation explains selective rRNA gene silencing and dosage control in Arabidopsis. *Genes Dev,* 30**,** 177-90.

CHEN, Z. J. & PIKAARD, C. S. 1997. Epigenetic silencing of RNA polymerase I transcription: a role for DNA methylation and histone modification in nucleolar dominance. *Genes Dev,* 11**,** 2124-36.

COKUS, S. J., FENG, S., ZHANG, X., CHEN, Z., MERRIMAN, B., HAUDENSCHILD, C. D., PRADHAN, S., NELSON, S. F., PELLEGRINI, M. & JACOBSEN, S. E. 2008. Shotgun bisulphite sequencing of the Arabidopsis genome reveals DNA methylation patterning. *Nature,* 452**,** 215-9.

COPENHAVER, G. P., DOELLING, J. H., GENS, S. & PIKAARD, C. S. 1995. Use of RFLPs larger than 100 kbp to map the position and internal organization of the nucleolus organizer region on chromosome 2 in Arabidopsis thaliana. *Plant J,* 7**,** 273-86.

COPENHAVER, G. P. & PIKAARD, C. S. 1996a. RFLP and physical mapping with an rDNA-specific endonuclease reveals that nucleolus organizer regions of Arabidopsis thaliana adjoin the telomeres on chromosomes 2 and 4. *Plant J,* 9**,** 259-72.

COPENHAVER, G. P. & PIKAARD, C. S. 1996b. Two-dimensional RFLP analyses reveal megabase-sized clusters of rRNA gene variants in Arabidopsis thaliana, suggesting local spreading of variants as the mode for gene homogenization during concerted evolution. *Plant J,* 9**,** 273-82.

COSTA-NUNES, P., PONTES, O., PREUSS, S. B. & PIKAARD, C. S. 2010. Extra views on RNA-dependent DNA methylation and MBD6-dependent heterochromatin formation in nucleolar dominance. *Nucleus,* 1**,** 254-9.

DE WINTER, R. F. & MOSS, T. 1986. Spacer promoters are essential for efficient enhancement of X. laevis ribosomal transcription. *Cell,* 44**,** 313-8.

DOELLING, J. H., GAUDINO, R. J. & PIKAARD, C. S. 1993. Functional analysis of Arabidopsis thaliana rRNA gene and spacer promoters in vivo and by transient expression. *Proc Natl Acad Sci U S A,* 90**,** 7528-32.

DOELLING, J. H. & PIKAARD, C. S. 1995. The minimal ribosomal RNA gene promoter of Arabidopsis thaliana includes a critical element at the transcription initiation site. *Plant J,* 8**,** 683-92.

DU, J., ZHONG, X., BERNATAVICHUTE, Y. V., STROUD, H., FENG, S., CARO, E., VASHISHT, A. A., TERRAGNI, J., CHIN, H. G., TU, A., HETZEL, J., WOHLSCHLEGEL, J. A., PRADHAN, S., PATEL, D. J. & JACOBSEN, S. E. 2012. Dual binding of chromomethylase domains to H3K9me2-containing nucleosomes directs DNA methylation in plants. *Cell,* 151**,** 167-80.

FULTZ, D., MCKINLAY, A., ENGANTI, R. & PIKAARD, C. S. 2023. Sequence and epigenetic landscapes of active and silent nucleolus organizer regions in Arabidopsis. *Sci Adv,* 9**,** eadj4509.

GAUDINO, R. J. & PIKAARD, C. S. 1997. Cytokinin induction of RNA polymerase I transcription in Arabidopsis thaliana. *J Biol Chem,* 272**,** 6799-804.

GERBI, S. A. 1986. The evolution of eukaryotic ribosomal DNA. *Biosystems,* 19**,** 247-58.
GHOSHAL, K., MAJUMDER, S., DATTA, J., MOTIWALA, T., BAI, S., SHARMA, S. M., FRANKEL, W. &
JACOB, S. T. 2004. Role of human ribosomal RNA (rRNA) promoter methylation and of
methyl-CpG-binding protein MBD2 in the suppression of rRNA gene expression. *J Biol Chem,*
279**,** 6783-93.
GORYSHIN, I. Y. & REZNIKOFF, W. S. 1998. Tn5 in vitro transposition. *J Biol Chem,* 273**,** 7367-74.
GRUMMT, I. 2013. The nucleolus—guardian of cellular homeostasis and genome integrity.
Chromosoma, 122**,** 487-97.
HANNAN, K. M., SANIJ, E., ROTHBLUM, L. I., HANNAN, R. D. & PEARSON, R. B. 2013. Dysregulation of
RNA polymerase I transcription during disease. *Biochim Biophys Acta,* 1829**,** 342-60.
HENDERSON, I. R. & JACOBSEN, S. E. 2007. Epigenetic inheritance in plants. *Nature,* 447**,** 418-24.
JEDDELOH, J. A., STOKES, T. L. & RICHARDS, E. J. 1999. Maintenance of genomic methylation requires
a SWI2/SNF2-like protein. *Nat Genet,* 22**,** 94-7.
KARAHAN, G., SAYAR, N., GOZUM, G., BOZKURT, B., KONU, O. & YULUG, I. G. 2015. Relative
expression of rRNA transcripts and 45S rDNA promoter methylation status are dysregulated
in tumors in comparison with matched-normal tissues in breast cancer. *Oncol Rep,* 33**,** 3131-
45.
LAW, J. A. & JACOBSEN, S. E. 2010. Establishing, maintaining and modifying DNA methylation
patterns in plants and animals. *Nat Rev Genet,* 11**,** 204-20.
LEE, I. & AMASINO, R. M. 1995. Effect of Vernalization, Photoperiod, and Light Quality on the
Flowering Phenotype of Arabidopsis Plants Containing the FRIGIDA Gene. *Plant Physiol,* 108**,**
157-162.
LEE, S. C., ADAMS, D. W., IPSARO, J. J., CAHN, J., LYNN, J., KIM, H. S., BERUBE, B., MAJOR, V.,
CALARCO, J. P., LEBLANC, C., BHATTACHARJEE, S., RAMU, U., GRIMANELLI, D., JACOB, Y.,
VOIGT, P., JOSHUA-TOR, L. & MARTIENSSEN, R. A. 2023. Chromatin remodeling of histone H3
variants by DDM1 underlies epigenetic inheritance of DNA methylation. *Cell,* 186**,** 4100-
4116.e15.
LEWIS, M. S. & PIKAARD, C. S. 2001. Restricted chromosomal silencing in nucleolar dominance. *Proc*
Natl Acad Sci U S A, 98**,** 14536-40.
LISTER, R., O'MALLEY, R. C., TONTI-FILIPPINI, J., GREGORY, B. D., BERRY, C. C., MILLAR, A. H. & ECKER,
J. R. 2008. Highly integrated single-base resolution maps of the epigenome in Arabidopsis.
Cell, 133**,** 523-36.
LONG, E. O. & DAWID, I. B. 1980. Repeated genes in eukaryotes. *Annu Rev Biochem,* 49**,** 727-64.
MATYÁŠEK, R., KALFUSOVÁ, R., KUDEROVÁ, A., ŘEHŮŘKOVÁ, K., SOCHOROVÁ, J. & KOVAŘÍK, A. 2024.
Transcriptional Silencing of 35S rDNA in Tragopogon porrifolius Correlates with Cytosine
Methylation in Sequence-Specific Manner. *Int J Mol Sci,* 25.
MO, W., SHU, Y., LIU, B., LONG, Y., LI, T., CAO, X., DENG, X. & ZHAI, J. 2023. Single-molecule targeted
accessibility and methylation sequencing of centromeres, telomeres and rDNAs in
Arabidopsis. *Nat Plants,* 9**,** 1439-1450.
MOHANNATH, G., PONTVIANNE, F. & PIKAARD, C. S. 2016. Selective nucleolus organizer inactivation
in Arabidopsis is a chromosome position-effect phenomenon. *Proc Natl Acad Sci U S A,* 113**,**
13426-13431.
MOSS, T., LANGLOIS, F., GAGNON-KUGLER, T. & STEFANOVSKY, V. 2007. A housekeeper with power
of attorney: the rRNA genes in ribosome biogenesis. *Cell Mol Life Sci,* 64**,** 29-49.
OSAKABE, A., JAMGE, B., AXELSSON, E., MONTGOMERY, S. A., AKIMCHEVA, S., KUEHN, A. L.,
PISUPATI, R., LORKOVIĆ, Z. J., YELAGANDULA, R., KAKUTANI, T. & BERGER, F. 2021. The
chromatin remodeler DDM1 prevents transposon mobility through deposition of histone
variant H2A.W. *Nat Cell Biol,* 23**,** 391-400.
PIKAARD, C. S., CHANDRASEKHARA, C., MCKINLAY, A., ENGANTI, R. & FULTZ, D. 2023. Reaching for
the off switch in nucleolar dominance. *Plant J,* 115**,** 1185-1192.

PONTVIANNE, F., ABOU-ELLAIL, M., DOUET, J., COMELLA, P., MATIA, I., CHANDRASEKHARA, C.,
DEBURES, A., BLEVINS, T., COOKE, R., MEDINA, F. J., TOURMENTE, S., PIKAARD, C. S. & SAEZ-
VASQUEZ, J. 2010. Nucleolin is required for DNA methylation state and the expression of
rRNA gene variants in Arabidopsis thaliana. *PLoS Genet,* 6**,** e1001225.
PONTVIANNE, F., BLEVINS, T., CHANDRASEKHARA, C., FENG, W., STROUD, H., JACOBSEN, S. E.,
MICHAELS, S. D. & PIKAARD, C. S. 2012. Histone methyltransferases regulating rRNA gene
dose and dosage control in Arabidopsis. *Genes Dev,* 26**,** 945-57.
PONTVIANNE, F., BLEVINS, T., CHANDRASEKHARA, C., MOZGOVA, I., HASSEL, C., PONTES, O. M.,
TUCKER, S., MOKROS, P., MUCHOVA, V., FAJKUS, J. & PIKAARD, C. S. 2013. Subnuclear
partitioning of rRNA genes between the nucleolus and nucleoplasm reflects alternative
epiallelic states. *Genes Dev,* 27**,** 1545-50.
PRIYADARSHINI, N., VENKATARAMA PUPPALA, N., PEROTH JAYAPRAKASH, J., KHANDELIA, P.,
SHARMA, V. & MOHANNATH, G. 2023. Ribosomal RNA gene downregulation in head and
neck squamous carcinoma cells correlates with rDNA promoter hypermethylation. *Gene***,**
147793.
RAVAL, A., SRIDHAR, K. J., PATEL, S., TURNBULL, B. B., GREENBERG, P. L. & MITCHELL, B. S. 2012.
Reduced rRNA expression and increased rDNA promoter methylation in CD34+ cells of
patients with myelodysplastic syndromes. *Blood,* 120**,** 4812-8.
RHEE, I., BACHMAN, K. E., PARK, B. H., JAIR, K. W., YEN, R. W., SCHUEBEL, K. E., CUI, H., FEINBERG, A.
P., LENGAUER, C., KINZLER, K. W., BAYLIN, S. B. & VOGELSTEIN, B. 2002. DNMT1 and
DNMT3b cooperate to silence genes in human cancer cells. *Nature,* 416**,** 552-6.
RYMEN, B., FERRAFIAT, L. & BLEVINS, T. 2020. Non-coding RNA polymerases that silence
transposable elements and reprogram gene expression in plants. *Transcription,* 11**,** 172-191.
SARADADEVI, G. P., PRIYADARSHINI, N., BERA, A. & MOHANNATH, G. 2020. Together we are on
together we are off -a conserved rule for ribosomal RNA (rRNA) gene regulation? *Journal of*
Plant Biochemistry and Biotechnology, 29**,** 743-753.
SASAKI, E., KAWAKATSU, T., ECKER, J. R. & NORDBORG, M. 2019. Common alleles of CMT2 and
NRPE1 are major determinants of CHH methylation variation in Arabidopsis thaliana. *PLoS*
Genet, 15**,** e1008492.
SCHULTZ, M. D., HE, Y., WHITAKER, J. W., HARIHARAN, M., MUKAMEL, E. A., LEUNG, D., RAJAGOPAL,
N., NERY, J. R., URICH, M. A., CHEN, H., LIN, S., LIN, Y., JUNG, I., SCHMITT, A. D., SELVARAJ, S.,
REN, B., SEJNOWSKI, T. J., WANG, W. & ECKER, J. R. 2015. Human body epigenome maps
reveal noncanonical DNA methylation variation. *Nature,* 523**,** 212-6.
SHIRASAWA, K. & ARIIZUMI, T. 2023. Near-complete genome assembly of tomato (Solanum
lycopersicum) cultivar Micro-Tom. *bioRxiv***,** 2023.10.26.564283.
STROUD, H., GREENBERG, M. V., FENG, S., BERNATAVICHUTE, Y. V. & JACOBSEN, S. E. 2013.
Comprehensive analysis of silencing mutants reveals complex regulation of the Arabidopsis
methylome. *Cell,* 152**,** 352-64.
TUROWSKI, T. W. & TOLLERVEY, D. 2015. Cotranscriptional events in eukaryotic ribosome synthesis.
Wiley Interdiscip Rev RNA, 6**,** 129-39.
VIKTOROVSKAYA, O. V. & SCHNEIDER, D. A. 2015. Functional divergence of eukaryotic RNA
polymerases: unique properties of RNA polymerase I suit its cellular role. *Gene,* 556**,** 19-26.
VONGS, A., KAKUTANI, T., MARTIENSSEN, R. A. & RICHARDS, E. J. 1993. Arabidopsis thaliana DNA
methylation mutants. *Science,* 260**,** 1926-8.
ZEMACH, A., KIM, M. Y., HSIEH, P. H., COLEMAN-DERR, D., ESHED-WILLIAMS, L., THAO, K., HARMER,
S. L. & ZILBERMAN, D. 2013. The Arabidopsis nucleosome remodeler DDM1 allows DNA
methyltransferases to access H1-containing heterochromatin. *Cell,* 153**,** 193-205.
ZHANG, X., YAZAKI, J., SUNDARESAN, A., COKUS, S., CHAN, S. W., CHEN, H., HENDERSON, I. R., SHINN,
P., PELLEGRINI, M., JACOBSEN, S. E. & ECKER, J. R. 2006. Genome-wide high-resolution
mapping and functional analysis of DNA methylation in arabidopsis. *Cell,* 126**,** 1189-201.

ZHOU, H., WANG, Y., LV, Q., ZHANG, J., WANG, Q., GAO, F., HOU, H., ZHANG, H., ZHANG, W. & LI, L.
2016. Overexpression of Ribosomal RNA in the Development of Human Cervical Cancer Is
Associated with rDNA Promoter Hypomethylation. *PLoS One,* 11**,** e0163340.
ZILBERMAN, D., GEHRING, M., TRAN, R. K., BALLINGER, T. & HENIKOFF, S. 2007. Genome-wide
analysis of Arabidopsis thaliana DNA methylation uncovers an interdependence between
methylation and transcription. *Nat Genet,* 39**,** 61-9.

Figure Legends

Figure 1: Ribosomal RNA (rRNA) gene organization, subtype variation, localization, and
their relative arrangement. (A) The diagram shows a generic tandem arrangement of 45S
rRNA genes on a NOR. (B) The diagram shows the 3' ETS region of 45S rRNA genes that
exhibit length and sequence polymorphisms among the rRNA variants. (C) The gel image
shows HindIII CAPS assay of rRNA variants and the diagram at the bottom shows the
position of HindIII restriction site relative to the binding sites of primers that amplify all four
rRNA variants. (D) The diagram shows known locations of the rRNA variants on *NOR2* and
NOR4 in the Col-0 ecotype of *A. thaliana*.

Figure 2: Disruption of *NOR2* gene silencing to varying degrees in DNA methylation mutants.
The gel images show DNA-PCR data for rRNA gene subtypes and RT-PCR expression data
for rRNA gene subtypes and actin. The gel images were spliced to remove the second lane.

Figure 3: Methylation state of rDNA in CG, CHG, and CHH contexts. Whole genome
bisulfite sequencing data from various DNA methylation mutants were analyzed using
Methylpy, and the overall cytosine methylation state of rDNA is shown as heatmaps in all
three cytosine contexts. The Y-axis represents a single rDNA unit starting with a promoter
and ending with 3' ETS. Each line on the heatmap represents the methylation state of the
cytosines in a given methylation context in 50 nucleotide bins. CG is shown in red, CHG in
green, and CHH in blue. Grey indicates that no methylation sites were present in that bin for
a given context.
TSS= Transcription Start Site.

Figure 4: Cytosines in the promoters of 45S rRNA genes occur mostly in CHH, distantly
followed by CHG, and almost none in CG context.

(A) Relative distribution of cytosines among different cytosine contexts (CG, CHG, and CHH) in the minimal rDNA promoter regions of different plant species. (B) For *A. thaliana* and tomato, publicly available assembled NOR sequences were analyzed to quantify the number of 45S rRNA gene repeats which show 100% conserved gene promoter sequence (-55 to +6).

NA-Entire NOR sequences not available.

Figure 5*: NOR2* genes are hypermethylated and *NOR4* genes are hypomethylated. (A) Gel images on the left show resolution of HpaII- or buffer-treated Col-0 and *met1-7* DNA samples, and the gel images on the right show PCR amplification of rRNA gene subtypes. For these PCR reactions, gel-eluted DNA samples representing the HpaII-undigested ($\geqslant$ 20 kb) and the HpaII-digested (0.6-1.0 kb) fractions were used. (B) The gel images show rRNA gene subtype PCR products obtained from McrBC-digested or no enzyme-treated Col-0 DNA.

Figure 6: Col-0 *NOR2* genes are hypermethylated and Sf-2 *NOR4* genes are hypomethylated in the Col[Sf-NOR4] introgression line. (A) Diagram showing rRNA subtype composition of the Arabidopsis introgression line called Col[Sf-NOR4] in which Sf-2 *NOR4* and adjacent sequences were introgressed into the Col-0 genetic background. (B) The agarose gel images on the left show resolution of HpaII- or buffer-treated Col[Sf-NOR4] gDNA samples, and the gel images on the right show PCR amplification of rRNA variants with or without HindIII digestion. For these PCR reactions, gel-eluted DNA samples representing the HpaII-undigested ($\geqslant$ 20 kb) and the HpaII-digested (0.6-1.0 kb) fractions were used. (C) Summary of results shown in (B). (D) The gel images show PCR amplification of rRNA variants obtained from McrBC- or buffer-treated Col[Sf-NOR4] gDNA with or without HindIII digestion. (E) Summary of results shown in (D).

Figure 7: *NOR2* genes exist in a more condensed chromatin than *NOR4* genes. (A) Tn5 Transposase-accessible chromatin assay (ATAC) was performed on nuclei isolated from wild-type Col-0 plants. (B) DNA isolated from transposase-treated nuclei was used in PCR1 to amplify ~2.8kb rRNA gene promoter-proximal upstream regions involving the 3' ETS of the preceding gene and IGS. (C) The PCR1-amplified 2.8 kb fragments were gel-eluted followed by PCR2, products of which are visualized on a 2.8% agarose gel stained with ethidium bromide. Amplification of VAR1 and VAR2 genes from 3' ETS variable region from genomic DNA (gDNA) as a control is shown in the first lane. (D) The PCR1-amplified 2.8 kb fragments were gel-eluted followed by PCR3, products of which are visualized on a 2.8%

791 agarose gel stained with ethidium bromide. Here, PCR3 involves amplification of all rRNA
792 variants. Amplification of all rRNA variants from 3' ETS variable region from genomic DNA
793 (gDNA) as a control is shown in the first lane.
794
Figure 8: Model depicting a role for CMT2-dependent CHH methylation in rRNA gene
silencing in Arabidopsis. In Wild-type, NOR2 rRNA genes are methylated in all cytosine
contexts. Loss of RdDM-dependent mCHH does not affect rRNA gene silencing at NOR2.
However, loss of CMT2-mediated mCHH results in disruption of NOR2 rRNA gene silencing
and likely alleviates chromatin condensation at these loci. Collectively, these findings
suggest that CMT2-mediated CHH methylation plays a critical role in rRNA gene silencing.
801

Figure 1. Ribosomal RNA (rRNA) gene organization, subtype variation, localization and their relative arrangement
(A) The diagram shows a generic tandem arrangement of 45S rRNA genes on a NOR. (B) The diagram shows the 3' ETS region of 45S rRNA genes that exhibit length and sequence polymorphisms among the rRNA variants. (C) The gel image shows HindIII CAPS assay of rRNA variants and the diagram at the bottom shows the position of HindIII restriction site relative to the binding sites of primers that amplify all four rRNA variants. (D) The diagram shows known locations of the rRNA variants on *NOR2* and *NOR4* in the Col-0 ecotype of *A. thaliana*.

rRNA subtype expression in DNA methylation mutants

Figure 2. Disruption of *NOR2* gene silencing to varying degrees in DNA methylation mutants.

The gel images show DNA-PCR data for rRNA gene subtypes and RT-PCR expression data for rRNA gene subtypes and actin. The gel images were spliced to remove the second lane.

Figure 3: Methylation state of rDNA in CG, CHG, and CHH contexts. Whole genome bisulfite sequencing data from various DNA methylation mutants were analyzed using Methylpy, and the overall cytosine methylation state of rDNA is shown as heatmaps in all three cytosine contexts. The Y-axis represents a single rDNA unit starting with a promoter and ending with 3' ETS. Each line on the heatmap represents the methylation state of the cytosines in a given methylation context in 50 nucleotide bins. CG is shown in red, CHG in green, and CHH in blue. Grey indicates that no methylation sites were present in that bin for a given context. TSS= Transcription Start Site

Figure 4. Cytosines in the promoters of 45S rRNA genes occur mostly in CHH, distantly followed by CHG, and almost none in CG context.
(A) Relative distribution of cytosines among different cytosine contexts (CG, CHG, and CHH) in the minimal rDNA promoter regions of different plant species. (B) For *A. thaliana* and tomato, publicly available assembled NOR sequences were analyzed to quantify the number of 45S rRNA gene repeats which show 100% conserved gene promoter sequence (-55 to +6).
NA-Entire NOR sequences not available.

(A) PCR analysis of rRNA gene subtypes upon HpaII digestion of genomic DNA

PCR analysis of rRNA gene variants

(B) PCR analysis of rRNA gene variants upon McrBc digestion of genomic DNA

Figure 5. *NOR2* genes are hypermethylated and *NOR4* genes are hypomethylated.

(A) Gel images on the left show resolution of HpaII- or buffer-treated Col-0 and *met1-7* DNA samples and the gel images on the right show PCR amplification of rRNA gene subtypes. For these PCR reactions, gel-eluted DNA samples representing the HpaII-undigested (≥ 20 kb) and the HpaII-digested (0.6-1.0 kb) fractions were used. (B) The gel image shows rRNA gene subtype PCR products obtained from McrBc-digested or no enzyme-treated Col-0 DNA.

(A) Col^{Sf-NOR4} line has Col-0 *NOR2* and Sf-2 *NOR4*

(B) HindIII CAPS analysis of rRNA gene subtypes amplified from gel-eluted fractions of HpaII-treated genomic DNA

(C)

rRNA variant composition of PCR amplicons		Position of HindIII in PCR amplicons	Relative DNA methylation status	Lane 1 (no enzyme control)	Lane 2 (HindIII after PCR)	Lane 3 (HpaII before PCR & No HindIII)	Lane 4 (HpaII before PCR & HindIII after PCR)
VAR1	Sf-2 VAR1 (471 bp)	F ⊢ HindIII ⊣ R 90 bp — 381 bp	hypomethylated	uncut	cut	cut by HpaII	cut by HpaII & HindIII
	Colo-0 VAR1 (471 bp)	F ⊢ R	hypermethylated	uncut	uncut	uncut	uncut by HpaII or HindIII
VAR3	Col-0 VAR3a (348 bp)	F ⊢ R	hypermethylated	uncut	uncut	uncut	uncut by HpaII or HindIII
	Col-0 VAR3b (348 bp)	F ⊢ HindIII ⊣ R 90 bp — 258 bp	hypermethylated	uncut	cut	uncut	uncut by HpaII but cut by HindIII

F-forward primer; R-reverse primer

(D) HindIII CAPS analysis of rRNA gene subtypes amplified from McrBc-treated genomic DNA

(E)

rRNA variant composition of PCR amplicons		Position of HindIII in PCR amplicons	Relative DNA methylation status	Lane 1 (no enzyme control)	Lane 2 (HindIII after PCR)	Lane 3 (McrBc before PCR & No HindIII)	Lane 4 (HpaII before PCR & HindIII after PCR)
VAR1	Sf-2 VAR1 (471 bp)	F — HindIII — R / 90 bp — 381 bp	hypomethylated	uncut	cut	uncut	uncut by McrBc & cut by HindIII
	Colo-0 VAR1 (471 bp)	F — R	hypermethylated	uncut	uncut	cut by McrBc	cut by McrBc & uncut by HindIII
VAR3	Col-0 VAR3a (348 bp)	F — R	hypermethylated	uncut	uncut	cut by McrBc	cut by McrBc & uncut by HindIII
	Col-0 VAR3b (348 bp)	F — HindIII — R / 90 bp — 258 bp	hypermethylated	uncut	cut	cut by McrBc	cut by McrBc & HindIII

F-forward primer; R-reverse primer

Figure 6. Col-0 *NOR2* genes are hypermethylated and Sf-2 *NOR4* genes are hypomethylated in the Col^sf-NOR4 introgression line.
(A) Diagram showing rRNA subtype composition of the Arabidopsis introgression line called Col^sf-NOR4 in which Sf-2 *NOR4* and adjacent sequences were introgressed into the Col-0 genetic background. (B) The agarose gel images on the left show resolution of HpaII- or buffer-treated Col^sf-NOR4 gDNA samples, and the gel images on the right show PCR amplification of rRNA variants with or without HindIII digestion. For these PCR reactions, gel-eluted DNA samples representing the HpaII-undigested (≥ 20 kb) and the HpaII-digested (0.6-1.0 kb) fractions were used. (C) Summary of results shown in (B). (D) The Gel image shows PCR amplification of rRNA variants obtained from McrBc- or buffer-treated Col^sf-NOR4 gDNA with or without HindIII digestion. (E) Summary of results shown in (D).

Figure 7. *NOR2* genes exist in a more condensed chromatin than *NOR4* genes.

(A) Tn5 Transposase-accessible chromatin assay (ATAC) was performed on nuclei isolated from wild-type Col-0 plants. (B) DNA isolated from the transposase-treated nuclei was used in PCR1 to amplify ~2.8 kb of rRNA gene promoter-proximal upstream regions involving the variable 3'ETS of the preceding gene and IGS. (C) The PCR1-amplified 2.8 kb fragments were gel-eluted followed by PCR2, products of which are visualized on a 1% agarose gel stained with ethidium bromide. Amplification of VAR1 and VAR2 genes from 3' ETS variable region from genomic DNA (gDNA) as a control is shown in the first lane. (D) The PCR1-amplified 2.8 kb gments were gel-eluted followed by PCR3, products of which are visualized on a 2.8% agarose gel stained with ethidium bromide. Here, PCR3 involves amplification of all rRNA variants. Amplification of all rRNA variants from 3' ETS variable region from genomic DNA (gDNA) as a control is shown in the first lane.

Figure 8. Model depicting a role for CMT2-dependent CHH methylation in rRNA gene silencing in Arabidopsis. In Wild-type, *NOR2* rRNA genes are methylated in all cytosine contexts. Loss of RdDM-dependent mCHH does not affect rRNA gene silencing at *NOR2.* However, loss of CMT2-mediated mCHH results in disruption of *NOR2* rRNA gene silencing and likely alleviates chromatin condensation at these loci. Collectively, these findings suggest that CMT2-mediated CHH methylation plays a critical role in rRNA gene silencing.