Multimegabase-scale DNA hypermethylation and condensed chromatin correlate with chromosome-specific rRNA gene silencing in Arabidopsis

Gireesha Mohannath[1,*], Anastasia McKinlay[2,3,†], Navinchandra Venkata Puppala[1,,†], Ramya Enganti[2,†], Gargi Prasad Saradadevi[1,†], Todd Blevins[4] and Craig S. Pikaard[2,3,]

[1]Department of Biological Sciences, Birla Institute of Technology and Science-Pilani, Hyderabad campus, Hyderabad, Telangana, 500078, India.

[2]Department of Biology and Department of Molecular and Cellular Biochemistry, Indiana University, Bloomington, IN, 47405, United States.

[3]Howard Hughes Medical Institute, Indiana University, Bloomington, IN, 47405, United States.

[4]Institut de Biologie Moléculaire des Plantes, CNRS, Université de Strasbourg, F-67084 Strasbourg, France.

* To whom correspondence should be addressed.

Tel: +91-40-66303697

    +1-812-855-1874

Fax: +91-40-66303998

Email: gireesha@hyderabad.bits-pilani.ac.in

† Joint second authors

ABSTRACT

In eukaryotes, hundreds of ribosomal RNA (rRNA) genes are clustered at chromosomal loci called nucleolus organizer regions (NORs). *Arabidopsis thaliana* has two NORs, one on chromosome 2 (NOR2) and the other on chromosome 4 (NOR4). Each NOR consists of ~ 400 rRNA gene copies. We recently showed that rRNA gene subtypes that map to the NOR2 are silenced during development, whereas those that map to NOR4 are active. In several DNA methylation mutants of Arabidopsis, we show release of the NOR2 silencing to varying degrees. Significantly, the highest release of NOR2 silencing correlates with a maximum loss of cytosine methylation in the CHH context followed by CG, but independent of RNA-directed DNA methylation (RdDM). Next, we show that NOR2 is relatively hypermethylated and NOR4 is hypomethylated using methylation analysis of genomic DNA carried out with different types of methylation-sensitive restriction enzymes. We demonstrate similar differential methylation status between NOR2 and NOR4 in genotypes carrying NORs from multiple ecotypes. Lastly, using Tn5 transposon-mediated transposition assay, we show that NOR2 chromatin is in more condensed state than NOR4.

INTRODUCTION

In eukaryotes, hundreds to thousands of 45S ribosomal RNA (rRNA) genes are arranged as tandem repeats at chromosomal loci known as nucleolus organizer regions (NORs) (Fig. 1A). RNA polymerase I (Pol I) transcribes rRNA genes to produce 45S pre-rRNA transcripts that are then processed into 18S, 5.8S, and 25-28S rRNAs that comprise the catalytic core of ribosomes, which synthesize proteins in cells. (1-3). Although eukaryotic genomes have many rRNA genes, not all of them are transcribed all the time. Rather, the number of genes transcribed varies during development, dictated by the developmental stage-specific need for ribosomes and protein synthesis (4-6). The rRNA gene regulation in eukaryotes has been found to be a form of 'dosage control' that involves selective gene silencing, not selective gene activation (7). Our recent findings in *Arabidopsis thaliana* (ecotype Col-0) revealed that such selective silencing is determined by chromosomal location and not by the rRNA gene sequence variation (8,9).

Ribosomal RNA genes in *A. thaliana* are organized into two NORs; one NOR on chromosome 2 (NOR2) and the other NOR on chromosome 4 (NOR4). Each NOR consists of ~ 400 rRNA gene copies. The 18S, 5.8S, and 25S rRNA sequences are identical among 45S rRNA genes, but some variation exists in the internal transcribed spacer (ITS) and the external transcribed spacer (ETS) regions. Four distinct variants had been defined based on the variation in the 3'ETS region named VAR1 through 4 (Fig. 1B-C) (10). Of these, VAR1, VAR2, and a subset of VAR3 (VAR3a) lack a HindIII site in the 3'ETS region, but the remaining variants (VAR3b,c and VAR4) have a HindIII site (8,9) (Fig.1C).

Earlier, we had shown that the rRNA gene variants that map to NOR2 (VAR1, VAR3a) are selectively silenced as a consequence of dosage control during development. In contrast, the variants that map to NOR4 (VAR2, VAR3b,c, and VAR4) remain active (Fig. 1D)(8). Silencing of many genes in eukaryotes involves cytosine methylation and chromatin condensation. First, we sought to know the specific role of cytosine methylation in all three contexts (CG, CHG and CHH) in selectively silencing NOR2. To address this question, we analyzed various DNA methylation mutants of *A. thaliana*, including mutants of DECREASE IN DNA METHYLATION 1 (*ddm1-2*), METHYLTRANSFERASE 1 (*met1-7*), DOMAINS REARRANGED METHYLASE 2 (*drm2*), CHROMOMETHYLASE (*cmt2-3*) and CMT3 (*cmt3*). DDM1 is a SWI2/SNF2 family chromatin remodeler which acts as a master regulator of DNA methylation (11). It promotes DNA methyltransferases' access to the H1 histone-containing heterochromatin to enable silencing of transposable elements in cooperation with RNA-directed DNA Methylation (RdDM) pathway (12). Loss of DDM1 function resulted in genome-wide demethylation of 70% of all cytosines (13,14). In plants, MET1 is involved in cytosine

methylation in CG context, Chromomethylases (CMT) in CHG context and DRM in CHH context (15). However, another study on CMT2 and DDM1 revealed that most of the cytosine methylation in CHH context is facilitated by DDM1 and mediated by CMT2, but independent of RNA-directed DNA methylation (RdDM). Using RT-PCR analysis, we show that the highest release of NOR2 silencing correlates with a maximum loss of cytosine methylation in the CHH context, followed by CG, but independent of RNA-directed DNA methylation (RdDM).

Further, to determine if DNA methylation and chromatin condensation are involved in the selective silencing of NOR2, we analyzed the cytosine methylation and chromatin status of NOR2 and NOR4 regions using multiple molecular approaches. First, using methylation-sensitive restriction digestion analyses, we show hypermethylation of silent rRNA genes at NOR2 and hypomethylation of the active rRNA genes at NOR4. We demonstrate similar results in different Arabidopsis genotypes containing NORs from multiple ecotypes. Lastly, using Tn5 transposase-mediated transposition assay of native chromatin (16), we show that active NOR4 exists as open chromatin while the silenced NOR2 exists as condensed chromatin.

MATERIAL AND METHODS

Plant material and growth

The *A. thaliana* ecotype Col-0 (stock no. CS22681), and mutants *met1-7* (stock no. SALK_076522), *cmt2-3* (stock no. SALK_012874C), *cmt3*, and *hda6-7* (stock no. CS66154) were obtained from the Arabidopsis Biological Resource Center (ABRC; http://abrc.osu.edu). *drm2-2b* (stock no. SAIL_70_E12) and Col$^{Sf-NOR4}$ were provided by Andrejez Wierzbicki and Scott Michaels, respectively, and *ddm1-2* and *met1-1* were provided by Eric Richards. All plants were grown under standard growth conditions for Arabidopsis in a plant growth chamber.

RNA isolation

Four-week-old leaf tissue of individual plants was frozen in liquid nitrogen and ground into a fine powder. RNeasy plant minikit (Qiagen) was used to extract total RNA from ground powder. Total RNA was then treated with Turbo DNA-free kit (Invitrogen) to eliminate contaminating DNA.

PCR and RT-PCR assays

For PCR of genomic DNA to amplify all rRNA variants, PCR amplification was conducted using 22-30 cycles (depending on the source of DNA) of 30 sec at 95°C, 30 sec at 55°C–60°C, and 60 sec at 72°C. For RT–PCR, the reverse transcription reaction was performed using 1 µg of total RNA using SuperScript III reverse transcriptase (Invitrogen). One microliter (~100 ng) of reverse transcription product was then used for PCR amplification of rRNA variable regions or an actin control using 30 cycles of 30 sec at 94°C, 30 sec at 55°C, and 60 sec at 72°C. In all the PCR or RT-PCR reactions, primers 5'allvar and 3'allvar were used to amplify rRNA variable regions (all the 4 variant types). PCR and RT–PCR products were resolved by electrophoresis on 2%–2.5% agarose gels in TBE buffer. Sequences of primers used in this study are given in Figure S1.

Generation of methylation heat maps

Whole genome bisulfite sequencing data (GSE39901) published by Stroud et al. (2013)(17), was used to assess the methylation state of rDNA in various methylation mutants. The sequencing data were analyzed using Methylpy (18). The reads were mapped to a converted 45S rDNA reference sequence allowing a single mismatch. The methylation state of all cytosines in CG, CHH and CHG contexts was then analyzed and represented as a fraction of methylated cytosines over the total cytosines in each context. Replicates for a given genotype were merged into a single file.

DNA isolation from leaf tissue and agarose gel pieces (gel elution).

Four-week-old individual plant leaf tissue was incubated for 15 min at 99°C in extraction buffer (200 mM Tris-HCl at pH 8.0, 250 mM NaCl, 25 mM EDTA, 0.5% SDS). Genomic DNA from the supernatant was recovered by ethanol precipitation and resuspended in 60 µL of Tris-HCl (pH 8.0). In some cases, DNA was isolated from leaf tissue of 4-wk-old plants by using Illustra Nucleon Phytopure extraction kit reagents (GE Healthcare). DNA was isolated from agarose gel pieces using QIAquick gel extraction kit (Qiagen).

Nuclei isolation

Nuclei were isolated from 3-week-old Col-0 plant seedlings as described previously (22) with a few modifications. Briefly, 3-week-old plants were harvested and washed in ice-cold water. Then plants were submerged in cold diethyl ether for 5 min and washed twice in ice-cold water. Homogenization of plant material was performed in 100 mL of nuclei isolation buffer

(1M sucrose, 50 mM Tris-HCl pH 7.2, 5 mM MgCl₂, 5 mM KCl, 10 mM 2-mercaptoethnol, 0.2

mM PMSF) with a motorized homogenizer (Fisher Scientific Powergen 700). Homogenate

was filtered through 1 layer of Miracloth and 1 layer of nylon mesh (pore size of 50 μm).

After centrifugation of the filtered homogenate at 14,000 g for 15 min at 4°C the pellet was

resuspended in 10 ml of nuclei isolation buffer and prepared for loading into discontinuous

Percoll gradient (15%, 30%, 45%, and 60%) by adding one volume of "100% Percoll

solution" (34.23 g sucrose, 5 mL 1 M Tris-HCl pH 7.2, 0.5 mL 1 M MgCl₂, 0.5 mL 1 M KCl,

34 μL 2-mercaptoethnol, 100 μL 2 mM PMSF and Percoll to 100 mL) to 19 volumes of nuclei.

Gradients were centrifuged at 2,000 rpm for 10 min, then 8,000 rpm for 20 min. Nuclei were

harvested from 30%/45% and 45%/60% boundaries and pooled. Nuclei were diluted with 5

volumes of nuclei isolation buffer, mixed by inversion and collected by centrifugation at 1,500

g for 10 min at 4°C. A second wash of nuclei was performed with 25 volumes of nuclei

isolation buffer. Final wash was done with 50 mL of Nuclease digestion buffer (50 mM Tris-

HCl, pH 8.0, 0.3 M sucrose, 5 mM MgCl2, 1.5 mM NaCl, 0.1 mM CaCl2, 5 mM beta-

mercaptoethanol). The resulting pellet was resuspended in 2 mL of Nuclease digestion

buffer.

Transposase-accessible chromatin assay

The assay was carried out based on the methodology described by Buenrostro et al. (16).

Oligos (5'-Ez-Tn5 and 3'-Ez-Tn5) were annealed by heating a mixture of equal volumes of

100 mM oligos at 95 °C for 5 min, followed by letting the mixture cool down gradually by

turning the heating off followed by another incubation at RT for 10 min. Transposition

reactions were carried out using 2 μL of purified nuclei, 20 μL of EZ-Tn5 transposase with

varying amount of the oligos 5'-Ez-Tn5 and 3'-Ez-Tn5, according to the manufacturer

instructions (Epicentre). Incubation was performed at 37°C for 90 min. DNA was isolated

using cetyl trimethylammonium bromide (CTAB) method. PCR1 and PCR2 were carried out

as described above. For the sequences of the primers and oligos used in this study, refer to

the table (Fig. S1).

RESULTS

Selective NOR2 silencing is released in DNA methylation mutants

To determine the relative impact of loss of cytosine methylation in three different contexts

(CG, CHG and CHH) on the release of selective NOR2 silencing, we analysed rRNA variant

expression in several DNA methylation mutants. We performed RT-PCR analysis using RNA
isolated from 4-week-old leaf tissues of various mutants. VAR1 type of rRNA genes
constitute ~50% of total 45S rRNA genes, and all of them map to NOR2, which undergoes
selective silencing during development (8,10). Therefore, VAR1 expression levels are used
as a measure of the release of NOR2 silencing. Among the DNA methylation mutants, VAR1
expression levels was found to be the highest among *cmt2-3* and *ddm1-2* mutants followed
by *met1-7* (Fig. 2 and Fig. S2)*. We confirmed high VAR1 expression in these mutants by
analysing multiple independent mutant plants using RT-PCR analysis (Fig. S2). However,
the *cmt3* and *drm2* (a member of RdDM pathway) mutants displayed low levels of VAR1
expression, if any (Fig. 2). The *ddm 1-2* mutation is known to reduce cytosine methylation in
all three contexts (CG, CHG and CHH) while *cmt2-3* is known to reduce most of cytosine
methylation in the CHH context, independent of RdDM (12). Using published whole-genome
bisulfite sequencing data (17), we generated heatmaps of cytosine methylation for a 45S
rRNA reference gene sequence (Fig. S3) using Methylpy (18) for multiple DNA methylation
mutants and wild-type Col-0 (Fig. 3). The cytosine methylation heatmaps confirm loss of
most of the cytosine methylation in CHH context in *cmt2-3* and *ddm1-2* mutants (Fig. 3), in
which release of NOR2 silencing was observed to a higher degree (Fig. 2 and Fig. S2). The
hda6-7 mutant was included as a positive control which has already been known to release
NOR2 silencing to a very high degree (8,23). Similarly, heatmaps confirm the loss of most of
the cytosine methylation in CG context in *ddm1-2* and *met1-7* mutants (Fig. 3). Overall, the
patterns of cytosine methylation loss for the rDNA loci align with what has been known of
these mutants regarding their effect on genome-wide methylation status. Nevertheless,
generating these heatmaps for the rDNA loci was important as most of the earlier genome-
wide studies excluded rRNA genes due to their repetitive nature.

NOR2 is hypermethylated while NOR4 is hypomethylated in Col-0

First, we carried out methylation analysis of rRNA variants using HpaII-digested genomic
DNA (DNA isolated by grinding leaf tissue) (Fig. 4). Because rRNA variants-defining 3' ETS
region lacks HpaII site, HpaII digestion will not have any impact on amplification of rRNA
variants. Therefore, we chose to separate HpaII-digested and HpaII-undigested fractions of
genomic DNA by resolving them on an agarose gel. The fragments that remained at the top
of the gel (>20 Kb) constituted the HpaII-undigested fractions while those resolved at 0.6-1.0
Kb range constituted the HpaII-digested fractions (Fig. 4A, left panel). Both the digested and
the undigested fractions were gel-eluted, followed by PCR amplification of rRNA variants.
Through this analysis, we were able to identify the rRNA variants that were enriched in
HpaII-digested or undigested fraction. The PCR data again revealed VAR2 genes from

NOR4 to be the most predominant variant among the HpaII-digested products, indicating relative hypomethylation of NOR4 compared to NOR2 (Fig. 4A, right panel).

In another experiment, we subjected gDNA to digestion by McrBC, an enzyme which preferentially digests methylated DNA (cytosine methylation). Because rRNA variants-defining 3' ETS region has multiple McrBc sites ($R^mC(N_{40-3000})R^mC$), McrBC digestion will impact PCR amplification of rRNA variants. Therefore gel-elution is not required in this case. McrBC digestion of gDNA followed by PCR amplification of rRNA variants revealed that VAR1 was digested the most and VAR2 the least (Fig. 4B), again indicating hypermethylation of NOR2 and hypomethylation of NOR4. HpaII digestion indicates cytosine methylation levels in the CG context only, but McrBc digestion indicates cytosine methylation levels in all three contexts (CG, CHG, and CHH). Methylation maps of centromere-proximal regions of NORs show that ~ 75 Kb of the NOR2-adjoining region is heavily methylated compared to the NOR4-adjoining region (Fig. S4), which has a protein-coding gene just 3 Kb away (24). It is possible that the heavily methylated NOR2-adjoining region might act as a source for spreading methylation-mediated silencing into the NOR2 region, but this hypothesis remains to be tested.

NOR2 is hypermethylated while NOR4 is hypomethylated in Col[Sf-NOR4]

An introgression line named Col[Sf-NOR4] consists of NOR2 from Col-0 (VAR1 and VAR3 genes) and NOR4 from another ecotype called Sf-2 (VAR1 genes) (8) (Fig. 5A). Sf-2 consists of only VAR1 type of genes, but they can be differentiated from VAR1 genes of Col-0 by HindIII digestion. Sf-2 VAR1 genes have a HindIII site in their 3'ETS region, while Col-0 VAR1 genes lack it (8). We also showed that in Col[Sf-NOR4], Sf-2 NOR4 is active and Col-0 NOR2 is selectively silenced (8). The methylation status of rRNA variants was determined using similar experiments as described above. In one experiment, genomic DNA (DNA isolated by grinding leaf tissue) of Col[Sf-NOR4] was subjected to HpaII digestion, followed by gel elutions of the undigested and the digested fractions. The gel-eluted DNA samples were then subjected to PCR amplification of rRNA variants followed by HindIII digestion. In control samples (no HpaII), HindIII digestion of PCR-amplified rRNA variants resulted in the expected products, which include digested Sf2-VAR1 genes, undigested Col-0 VAR1 genes, and partially digested Col-0 VAR3 genes (Fig. 5B, lanes 1 and 2, top right gel image) (8). However, HindIII digestion of the PCR products obtained from the gel-purified HpaII-digested fraction as template, revealed that Col-0 VAR1 genes predominantly constitute the HpaII-undigested fraction (Fig. 5B, lanes 3 and 4, top right gel image). We also observed a significant increase in the Sf-2 VAR1 genes in the HpaII-digested fraction (Fig. 5B, lanes 3 and 4, bottom right

gel image) compared to the undigested fraction (Fig. 5B, lanes 3 and 4, top right gel image). These data indicate hypermethylation of Col-0 NOR2 and hypomethylation of Sf-2 NOR4, despite the presence of the VAR1 genes on both the NORs in the Col$^{Sf-NOR4}$ introgression line.

In another experiment, gDNA (DNA isolated by grinding leaf tissue) of Col$^{Sf-NOR4}$ was first digested with McrBc, followed by PCR amplification of rRNA variants. These PCR products were then digested with HindIII and resolved using regular gel electrophoresis. HindIII-digested control PCR products (no McrBc digestion) behaved as expected (Fig. 5C, lanes 1 and 2). However, the McrBC digestion patterns of the rRNA variants (Fig. 5C) turned out to be exactly the opposite of the experimental results involving HpaII digestion (compare to Fig. 5B). Predominantly, the Sf-2 VAR1 genes constituted the McrBc-undigested fraction, while the Col-0 VAR1 genes constituted the McrBc-digested fraction (Fig. 5C, lanes 3 and 4). We also observed complete digestion of the Col-0 VAR3 genes by McrBC (Fig. 5C, lane 3). These data again indicate that NOR4 is hypomethylated and NOR2 is hypermethylated in the Col$^{Sf-NOR4}$ introgression line despite the presence of VAR1 type of genes on both the NORs.

NOR2 exists as condensed chromatin while NOR4 exists as open chromatin

Next, to determine the chromatin condensation status of NOR2 and NOR4, we carried out Tn5 transposase-mediated transposition using nuclei isolated from 3-week-old Col-0 plants. Tn5 is a prokaryotic hyperactive transposase that has been developed as a tool to insert short DNA sequences into open chromatin regions (16,25,26) where in 'outside ends' (OEs) sequences are sufficient for Tn5 recognition for transposition and gDNA nicking (27). We designed short DNA sequences consisting of OEs for Tn5 recognition and used them in a transposition assay. Purified Col-0 nuclei were incubated with either no or 5 pmol oligos containing Tn5 recognition sites for transposition, followed by DNA extraction from the nuclei and PCR analysis of rRNA variants. The PCR data did not reveal differences between NOR2 and NOR4 (data not shown). Then, we tried an indirect approach to focus on the chromatin accessibility state of the promoter regions. In this approach, the DNA samples isolated from nuclei subjected to Tn5 transposition were subjected to two rounds of PCR amplification. The first round of PCR (PCR1) amplifies ~2.8 Kb region that spans 3'ETS region, IGS and the promoter of the following gene repeat (Fig. 6B). Due to lack of variant-specific sequence variation in the promoter region, we exploited the proximity of the rRNA variant-defining 3'ETS region of one rRNA gene to the promoter of the following rRNA gene. The occurrence of rRNA genes as tandem repeats (Fig. 1A) at NORs permitted us to explore this approach.

The idea is that, if promoters of VAR1 genes (NOR2) are more condensed than the
promoters of VAR2 genes, then the 2.8 Kb PCR products should be enriched for VAR1
genes. Because the 2.8 Kb PCR products obtained during PCR1 (using primers 5'allvar and
p34) can't discriminate between NOR2 and NOR4, they were resolved on a 1% agarose gel
followed gel-purification of 2.8 Kb amplicons. These amplicons were then subjected to the
second round of PCR amplification (PCR2) with primers that either amplify only VAR1 and
VAR2 genes (Fig. 6C) or primers that amplify all rRNA variants (Fig. 6D). The results
showed severe depletion of NOR4-specific VAR2 genes in the reactions with 5 pmol of Tn5
oligos (Fig. 6C&D), indicating depletion of NOR4-specific fragments in the PCR1 amplicon
pool. This depletion could be presumably due to higher transposition rates in the promoter
regions of NOR4-specific rRNA variants compared to the NOR2 variants. Collectively, these
data indicate that NOR4 is predominantly in a more accessible open chromatin state and
NOR2 in a more condensed state.

DISCUSSION

Our earlier findings revealed that during development, NOR2 is silenced and NOR4 remains
transcriptionally active (8,9). In eukaryotes, often gene silencing involves DNA methylation
and chromatin condensation. First, we analysed several DNA methylation mutants to know
their effect on the release of NOR2 silencing (Fig. 2 and Fig. S2). In plants, cytosine
methylation occurs in all three contexts (CG, CHG and CHH). MET1 is primarily involved in
cytosine methylation in CG context whereas CHROMOMETHYLASES (CMT) are mainly
involved in cytosine methylation in CHG and CHH context, and DRM in cytosine methylation
in CHH context (12,15). DDM1, a SWI2/SNF2 family chromatin remodeler, is required for
cytosine methylation in all three contexts (CG, CHG, and CHH).

Although the release of NOR2 silencing was observed in most of the cytosine
methylation mutants studied, it was found to be the highest among *cmt2-3* and *ddm1-2,*
followed *met1-7* mutants (Fig. 2). The levels of release of NOR2 silencing in *ddm1-2* can be
comparable to that of *cmt2-3* if you look at multiple *cmt2-3* plants (Fig. S2). Notably, *ddm1-2*
and *cmt2-3* mutants also exhibit the highest loss of CHH methylation (Fig. 3)*.* Interestingly,
an earlier study in Arabidopsis has shown that most asymmetric cytosine methylation is
mediated by CMT2 and DDM1, independent of the RNA-directed DNA methylation (RdDM)
pathway (12). This could also explain why the release of NOR2 silencing is the least in *drm2*
mutants compared to the other mutants (Fig. 2). Note that DRM2 is a methyltransferase
primarily involved in RdDM. Although *ddm1-2* loses most of cytosine methylation in all

contexts (CG, CHG, CHH) (13,14) (Fig. 3), why the levels of release of NOR2 silencing is not greater than *cmt2-3,* which significantly affects cytosine methylation only in CHH context? One possible reason is that loss of CG and/or CHG methylation could dysregulate the expression of some of the positive regulators of NOR2 silencing, limiting the release of NOR silencing to some extent. Therefore, the effect of loss of cytosine methylation in all cytosine contexts in *ddm1-2* could be redundant than cumulative. Overall, the loss of cytosine methylation in the CHH context appears to have the maximum effect on the release of NOR2 silencing.

Next, to determine if DNA methylation and chromatin condensation are involved in selectively silencing NOR2, we thoroughly analysed DNA methylation and chromatin condensation status of NOR2 and NOR4 using various approaches in multiple genotypes carrying NORs from different ecotypes. To analyze DNA methylation status of NOR2 and NOR4 regions, we carried out rRNA variant analysis upon digestion of genomic DNA by different types of methylation-sensitive restriction enzymes (HpaII and McrBC) (Fig. 4). The HpaII does not digest DNA if its recognition site (CCGG) has cytosine methylation in the CG context. In contrast, McrBc preferentially digests DNA if its recognition site ($R^mC(N_{40-3000})R^mC$) consists of methylated cytosines. From these experiments involving two methylation-sensitive restriction enzymes with contrasting mechanisms of detecting cytosine methylation, we again showed that NOR2 is hypermethylated and NOR4 is hypomethylated (Fig. 4). HpaII digestion reveals the methylation status of cytosines only in the CG context, but McrBc digestion reveals the methylation status of cytosines in all three contexts (CG, CHG, CHH). Therefore, hypermethylation of cytosine on NOR2 could likely have occurred in all three contexts.

We conducted similar experiments using an introgression line named Col-[Sf-NOR4] which consists of NOR2 from Col-0 (VAR1 and VAR3 genes) and NOR4 from another ecotype called Sf-2 (8) (Fig. 5). Interestingly, Sf-2 consists of only VAR1 type of genes, but they can be differentiated from VAR1 genes of Col-0 by HindIII digestion. We have earlier shown that in Col[Sf-NOR4], Sf-2 NOR4 is active and Col-0 NOR2 is selectively silenced (8). Results from these experiments involving digestions with methylation-sensitive HpaII and McrBc, again showed hypermethylation of NOR2 and hypomethylation of NOR4 (Fig. 5). Based on these data, we conclude that NOR2 is hypermethylated compared to NOR4, and DNA methylation is involved in selectively silencing NOR2.

Further, we examined the chromatin condensation status of NOR2 and NOR4 using a Tn5 transposase-mediated transposition assay. Tn5 is a prokaryotic hyperactive

transposase that can insert short DNA sequences into open chromatin regions (16,25,26)
where in 'outside ends' (OEs) sequences are sufficient for Tn5 recognition for transposition
(27). The underlying principle is that the higher the frequency of insertion, the more open the
chromatin will be. PCR analysis of DNA obtained from Tn5-treated nuclei did not reveal any
difference in chromatin condensation states of NOR2 and NOR4. However, an indirect
approach involving two rounds of PCR reactions, as described above, revealed that NOR2 is
more condensed than NOR4 (Fig. 6). Well, one possibility for the observed result is that
there is not much difference between chromatin states at/near 3'ETS of NOR2 and NOR4,
but most of the differences could relatively be more at the promoter regions.

In conclusion, hypermethylation and chromatin condensation of NOR2 appears to be
primary contributing factor for selective NOR2 silencing. However, despite striking
similarities in rRNA gene sequences of NOR2 and NOR4, why does NOR2 undergo
selective hypermethylation, condensation and silencing? Interestingly, the immediate
centromere-proximal region of NOR2 is composed of transposable elements and transposon
remnants that stretch for ~ 60 kb (Fig. S4). The corresponding centromere-proximal regions
of NOR4 do not contain any stretch of transposons or their remnants (Fig. S4). In plants,
transposons are silenced through methylation-mediated mechanisms. Could such silencing
of the NOR2-adjacent transposon-rich region spread to NOR2, resulting in its selective
silencing? The question remains to be answered.

DATA AVAILABILITY

388 Whole genome bisulfite sequencing data published by Stroud et al. (2013)(17) with a GEO
389 accession number GSE39901 can be found in NCBI database
390 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE39901).

ACKNOWLEDGEMENT

393 We thank the members of the Pikaard lab and the Mohannath lab for their critique of the
394 manuscript.

FUNDING

396 G.M. is thankful to the Birla Institute of Science and Technology (BITS) Pilani, Hyderabad
397 campus for the research grant (BITS/GAU/ACRG/2019/H0576) to G.M. and an Institute

Fellowship to G.P.S., and to the Science and Engineering Research Board (SERB), Government of India, for the Ramanujan Fellowship Research Grant (SB-S2-RJN-062-2017) and CRG grant (CRG/2020/002855). C.S.P. is thankful to research funding from Howard Hughes Medical Institute, Indiana University (USA), National Institute of Health, and Gordon and Betty Moore Foundation.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. Saradadevi, G.P., Priyadarshini, N., Bera, A. and Mohannath, G. (2020) Together we are on together we are off -a conserved rule for ribosomal RNA (rRNA) gene regulation? *Journal of Plant Biochemistry and Biotechnology*, **29**, 743-753.

2. Turowski, T.W. and Tollervey, D. (2015) Cotranscriptional events in eukaryotic ribosome synthesis. *Wiley interdisciplinary reviews. RNA*, **6**, 129-139.

3. Viktorovskaya, O.V. and Schneider, D.A. (2015) Functional divergence of eukaryotic RNA polymerases: unique properties of RNA polymerase I suit its cellular role. *Gene*, **556**, 19-26.

4. Grummt, I. (2013) The nucleolus—guardian of cellular homeostasis and genome integrity. *Chromosoma*, **122**, 487-497.

5. Hannan, K.M., Sanij, E., Rothblum, L.I., Hannan, R.D. and Pearson, R.B. (2013) Dysregulation of RNA polymerase I transcription during disease. *Biochimica et biophysica acta*, **1829**, 342-360.

6. Moss, T., Langlois, F., Gagnon-Kugler, T. and Stefanovsky, V. (2007) A housekeeper with power of attorney: the rRNA genes in ribosome biogenesis. *Cell Mol Life Sci*, **64**, 29-49.

7. Chen, Z.J. and Pikaard, C.S. (1997) Epigenetic silencing of RNA polymerase I transcription: a role for DNA methylation and histone modification in nucleolar dominance. *Genes Dev*, **11**, 2124-2136.

8. Chandrasekhara, C., Mohannath, G., Blevins, T., Pontvianne, F. and Pikaard, C.S. (2016) Chromosome-specific NOR inactivation explains selective rRNA gene silencing and dosage control in Arabidopsis. *Genes Dev*, **30**, 177-190.

9. Mohannath, G., Pontvianne, F. and Pikaard, C.S. (2016) Selective nucleolus organizer inactivation in Arabidopsis is a chromosome position-effect phenomenon. *Proc Natl Acad Sci U S A*, **113**, 13426-13431.

10. Pontvianne, F., Abou-Ellail, M., Douet, J., Comella, P., Matia, I., Chandrasekhara, C., Debures, A., Blevins, T., Cooke, R., Medina, F.J. *et al.* (2010) Nucleolin is required for DNA methylation state and the expression of rRNA gene variants in Arabidopsis thaliana. *PLoS genetics*, **6**, e1001225.

11. Brzeski, J. and Jerzmanowski, A. (2003) Deficient in DNA methylation 1 (DDM1) defines a novel family of chromatin-remodeling factors. *The Journal of biological chemistry*, **278**, 823-828.

12. Zemach, A., Kim, M.Y., Hsieh, P.H., Coleman-Derr, D., Eshed-Williams, L., Thao, K., Harmer, S.L. and Zilberman, D. (2013) The Arabidopsis nucleosome remodeler DDM1 allows DNA methyltransferases to access H1-containing heterochromatin. *Cell*, **153**, 193-205.

13. Jeddeloh, J.A., Stokes, T.L. and Richards, E.J. (1999) Maintenance of genomic methylation requires a SWI2/SNF2-like protein. *Nature genetics*, **22**, 94-97.

14. Vongs, A., Kakutani, T., Martienssen, R.A. and Richards, E.J. (1993) Arabidopsis thaliana DNA
methylation mutants. *Science*, **260**, 1926-1928.
15. Law, J.A. and Jacobsen, S.E. (2010) Establishing, maintaining and modifying DNA methylation
patterns in plants and animals. *Nature reviews. Genetics*, **11**, 204-220.
16. Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y. and Greenleaf, W.J. (2013) Transposition
of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-
binding proteins and nucleosome position. *Nature methods*, **10**, 1213-1218.
17. Stroud, H., Greenberg, M.V., Feng, S., Bernatavichute, Y.V. and Jacobsen, S.E. (2013)
Comprehensive analysis of silencing mutants reveals complex regulation of the Arabidopsis
methylome. *Cell*, **152**, 352-364.
18. Schultz, M.D., He, Y., Whitaker, J.W., Hariharan, M., Mukamel, E.A., Leung, D., Rajagopal, N.,
Nery, J.R., Urich, M.A., Chen, H. *et al.* (2015) Human body epigenome maps reveal
noncanonical DNA methylation variation. *Nature*, **523**, 212-216.
19. Mohannath, G. and Pikaard, C.S. (2016) Analysis of rRNA Gene Methylation in Arabidopsis
thaliana by CHEF-Conventional 2D Gel Electrophoresis. *Methods in molecular biology (Clifton,
N.J.)*, **1455**, 183-202.
20. Chu, G., Vollrath, D. and Davis, R.W. (1986) Separation of large DNA molecules by contour-
clamped homogeneous electric fields. *Science*, **234**, 1582-1585.
21. Copenhaver, G.P., Doelling, J.H., Gens, S. and Pikaard, C.S. (1995) Use of RFLPs larger than
100 kbp to map the position and internal organization of the nucleolus organizer region on
chromosome 2 in Arabidopsis thaliana. *Plant J*, **7**, 273-286.
22. Gaudino, R.J. and Pikaard, C.S. (1997) Cytokinin induction of RNA polymerase I transcription
in Arabidopsis thaliana. *The Journal of biological chemistry*, **272**, 6799-6804.
23. Blevins, T., Pontvianne, F., Cocklin, R., Podicheti, R., Chandrasekhara, C., Yerneni, S., Braun,
C., Lee, B., Rusch, D., Mockaitis, K. *et al.* (2014) A two-step process for epigenetic inheritance
in Arabidopsis. *Molecular cell*, **54**, 30-42.
24. Lewis, M.S. and Pikaard, C.S. (2001) Restricted chromosomal silencing in nucleolar
dominance. *Proc Natl Acad Sci U S A*, **98**, 14536-14540.
25. Adey, A., Morrison, H.G., Asan, Xun, X., Kitzman, J.O., Turner, E.H., Stackhouse, B.,
MacKenzie, A.P., Caruccio, N.C., Zhang, X. *et al.* (2010) Rapid, low-input, low-bias
construction of shotgun fragment libraries by high-density in vitro transposition. *Genome
biology*, **11**, R119.
26. Goryshin, I.Y. and Reznikoff, W.S. (1998) Tn5 in vitro transposition. *The Journal of biological
chemistry*, **273**, 7367-7374.
27. Ahmed, A. (2009) Alternative mechanisms for tn5 transposition. *PLoS genetics*, **5**, e1000619.

(A) rRNA gene repeat arrangement at *A. thaliana* NORs

NOR

Spacer Repeats
Promoter
5'ETS
ITS1
ITS2
3'ETS
Variable Region

Intergenic Spacer (IGS)
45S pre-rRNA precursor

(B) 3'ETS variation defining rRNA variants

3' ETS variable region

+1

25S

PCR Primers

(C) HindIII digestion pattern of 3'ETS region of rRNA variants

Col-0
− +

VAR1 — Uncut VAR1 (471bp)
VAR2 — Uncut VAR2 (389bp)
VAR3 — Uncut VAR3a (348bp)

Cut VAR3b,c (258bp)

Cut VAR3b,c (90bp)

primer → HindIII ← primer
90bp   258bp

(D) Locations of silenced versus active rRNA gene subtypes

VAR1, VAR 3a,b* (Interspersed)

Chromosome 2

TEL2N ← NOR2(~4 Mb)

TEL4N ← NOR4 (~4 Mb)

Chromosome 4

VAR 3b*,c*, VAR2 & VAR4*

Figure 1. Ribosomal RNA (rRNA) gene organization, subtype variation, localization and their relative arrangement

rRNA subtype expression in DNA methylation mutants

Figure 2. Release of NOR2 silencing to varying degrees in DNA methylation mutants.

The gel images show DNA-PCR data for rRNA gene subtypes and RT-PCR expression data for rRNA gene subtypes and actin.

Figure 3: Methylation state of rDNA in CG, CHG and CHH contexts. Methylation state of rDNA in CG, CHG and CHH contexts. Whole genome bisulfite sequencing data from various mutants was analyzed using Methylpy and the overall cytosine methylation state of the rDNA is shown as heatmaps in all three contexts. The Y-axis is a single rDNA unit starting at the promoter and ending in the 3'ETS. Each line on the heatmap represents the methylation state of cytosines in a given methylation context in 50 nucleotide bins. CG is shown in red, CHG in green and CHH in blue. Grey indicates that no methylation sites were present in that bin for a given context.

(A) PCR analysis of rRNA gene subtypes upon HpaII digestion of genomic DNA

PCR analysis of rRNA gene variants

Figure 4. NOR2 is hypermethylated and NOR4 is hypomethylated

(A) Gel images on the left show resolution of HpaII- or buffer-treated Col-0 DNA samples and the gel images on the right show PCR amplification of rRNA gene subtypes. For these PCR reactions, gel-eluted DNA samples representing the HpaII-undigested (≥ 20 kb) and the HpaII-digested (0.6-1.0 kb) fractions were used. (B) The gel image shows rRNA gene subtype PCR products obtained from McrBc-digested or no enzyme-treated Col-0 DNA.

Figure 5. Col-0 NOR2 is hypermethylated and Sf-2 NOR4 is hypomethylated in the Col^sf-NOR4 introgression line.
(A) Diagram showing rRNA subtype composition of the Arabidopsis introgression line called Col^Sf-NOR4 in which Sf-2 NOR4 and adjacent sequences were introgressed into the Col-0 genetic background. (B) The agarose gel images on the left show resolution of HpaII- or buffer-treated Col^sf-NOR4 gDNA samples, and the gel images on the right show PCR amplification of rRNA variants with or without HindIII digestion. For these PCR reactions, gel-eluted DNA samples representing the HpaII-undigested (≥ 20 kb) and the HpaII-digested (0.6-1.0 kb) fractions were used. (C) The Gel image shows PCR amplification of rRNA variants obtained from McrBc- or buffer-treated Col^sf-NOR4 gDNA with or without HindIII digestion.

Figure 6. NOR2 exists as condensed chromatin while NOR4 exists as open chromatin.

(A) Tn5 Transposase-accessible chromatin assay (ATAC) was performed on nuclei isolated from wild-type Col-0 plants. (B) DNA isolated from the transposase-treated nuclei was used in PCR1 to amplify ~2.8 Kb of rRNA gene promoter-proximal upstream regions involving the variable 3'ETS of the preceding gene and IGS. (C) The PCR1-amplified 2.8 Kb fragments were gel-eluted followed by PCR2, products of which are visualized on a 2.8% agarose gel stained with ethidium bromide. Amplification of VAR1 and VAR2 genes from 3'ETS variable region from genomic DNA (gDNA) as a control is shown in the first lane. (D) The PCR1-amplified 2.8 Kb fragments were gel-eluted followed by PCR3, products of which are visualized on a 2.8% agarose gel stained with ethidium bromide. Here, PCR3 involves amplification of all rRNA variants. Amplification of all rRNA variants from 3'ETS variable region from genomic DNA (gDNA) as a control is shown in the first lane.