Title:

Gene panel selection for targeted spatial transcriptomics

Yida Zhang[1,2], Viktor Petukhov[1], Evan Biederstedt[1], Richard Que[3], Kun Zhang[3,4],

Peter V. Kharchenko[1,4,*]

[1]Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA.

[2]Department of Neurobiology, Duke University, Durham, NC, USA.

[3]Department of Bioengineering, University of California San Diego, La Jolla, CA, USA.

[4]San Diego Institute of Science, Altos Labs, San Diego, CA, USA.

[*]Corresponding author: peter_kharchenko@hms.harvard.edu

Abstract

Targeted spatial transcriptomics hold particular promise in analysis of complex

tissues. Most such methods, however, measure only a limited panel of transcripts,

which need to be selected in advance to inform on the cell types or processes being

studied. A limitation of existing gene selection methods is that they rely on scRNA-

seq data, ignoring platform effects between technologies. Here we describe gpsFISH,

a computational method to perform gene selection through optimizing detection of

known cell types. By modeling and adjusting for platform effects, gpsFISH

outperforms other methods. Furthermore, gpsFISH can incorporate cell type

hierarchies and custom gene preferences to accommodate diverse design

requirements.

28 **Background**

The building block of complex tissues is the diverse range of cell types [1–4].

Knowing the identity and spatial location of cells from different cell types is the key

for understanding how they communicate with each other to carry out specific

functions and how diseases emerge when this complex network of interactions goes

awry [5–11]. Single-cell RNA sequencing (scRNA-seq) provides a powerful tool to

study the identity of cell types and cell states [12–17]. However, the spatial

information is lost due to cell disassociation during library preparation. Recent

advances in spatial transcriptomics technologies have overcome this limitation by

providing ways to quantify gene expression while keeping the spatial information of

cells, leading to more comprehensive and detailed understanding of diseases and

normal functions [18–23].

Based on the number of transcripts that can be probed, spatial transcriptomics

technologies can be broadly categorized as (1) targeted, measuring a limited panel

of transcripts; and (2) untargeted, capturing all transcripts from the transcriptome.

Targeted spatial transcriptomics include in situ hybridization (ISH)-based [24–28]

and most in situ sequencing (ISS)-based methods [29–32]. Untargeted spatial

transcriptomics include next-generation sequencing (NGS)-based methods [33–39].

Compared to untargeted spatial transcriptomics, targeted spatial transcriptomics

can achieve high sensitivity and subcellular resolution. However, their targeted

nature requires a panel of genes (from a few hundred to thousand) to be selected in

advance to recognize cell types or processes relevant to the tissue being studied.

Gene selection methods are used to design gene panels. They can be classified into

two major categories based on their gene selection objectives. One category with an

imputation-based objective aims to select genes based on their ability to capture as

much of transcriptional variation in the scRNA-seq data as possible. Examples range

from simply selecting highly-variable genes to more advanced methods like L1000

[40], geneBasis [41], and SCMER [42]. Specifically, L1000 identified the optimal set

of 'landmark' transcripts that construct a reduced representations of the

transcriptome. geneBasis finds genes that can yield a k-nearest neighbor graph that

is similar to the "true" graph constructed using the entire transcriptome. SCMER

aims to select genes that preserve the manifold of scRNA-seq data. Another category

of gene selection method with a classification-based objective selects genes given

their ability to reconstruct cell classifications or relationships. Examples range from

selecting differentially expressed genes (DEGs) to more advanced methods like

scGeneFit [43], RankCorr [44], and NS-Forest [45]. scGeneFit selects marker genes

that jointly optimize cell type recovery using a label-aware compressive

classification method. RankCorr is a rank-based one-vs-all feature selection method

that selects marker genes for each cell type based on a sparsity parameter that

controls the number of marker genes selected per cell type. NS-Forest is a machine

learning-based marker gene selection algorithm that uses the nonlinear attributes of

random forest feature selection and a binary expression scoring approach to select

the minimal combination of marker genes that captures the cell type identity in

scRNA-seq data. All these methods can be used to design gene panels for targeted

spatial transcriptomics technologies.

A key limitation of current gene selection methods is that they select genes purely

based on scRNA-seq data without considering potential differences between scRNA-

seq and the target spatial transcriptomics technologies. Such platform effects

include systematic differences in capture efficiency of genes between platforms

caused by technology-dependent factors, including detection technique and library

preparation chemistry. Platform effects have been previously noted when

comparing gene expression measurements from single-cell and single-nucleus RNA-

seq on the same biological sample [46]. Platform effects also exist between scRNA-

seq and spatial transcriptomics technologies [47–49], posing a challenge when

transferring cell type information from scRNA-seq to spatial transcriptomics

technologies. When selecting gene panels using scRNA-seq data, such platform-

specific distortions can lead to reduced performance of selected gene panels in the

resulting spatial measurements.

Besides platform effects, there are other complications involved in gene panel

selection. First, current classification-based gene selection methods [43–45] treat

cell types as equally distinct. However, cell types are organized in a hierarchical

manner with cell subpopulations belonging to the same broad cell type more similar

to each other than subpopulations belonging to different broad cell types [50–56].

Depending on the biological questions and capabilities of the assays, a gene

selection method could optimize for fidelity at lower cell type resolution, or place

increased emphasis on certain subgroups of cell types. More generally, this is not

only useful for selecting genes that inform on cell types but can also be extended to

selecting genes for other biological entities with a hierarchical structure, e.g., gene

ontology and pathways [57,58]. Second, both imputation-based and classification-

based gene selection methods select genes solely based on a pre-defined objective

function. However, in practice of gene panel design for targeted spatial

transcriptomics, there can be other criteria contributing to the gene selection.

Examples range from technical factors, such as ability to design probes for targeting

certain transcripts, to biological factors such as preferences for certain pathways or

marker genes commonly used in the literature. A framework that takes such

orthogonal preferences into consideration can be helpful in practice.

To address these challenges, we developed gpsFISH, a classification-based gene

selection method that models and adjusts for the platform effects between scRNA-

seq and targeted spatial transcriptomics technologies, yielding more informative

gene panels and better cell type classification compared to previously published

classification-based gene selection methods. In addition, gpsFISH provides options

to account for cell type hierarchy and gene-specific custom preferences during gene

panel design, offering flexible and finer control of cell type granularity and gene

selection for different biological questions.

Results

Platform effects between scRNA-seq and targeted spatial transcriptomics

Even molecule counting assays carry inherent detection biases, posing challenges

for joint analysis of multiple assays, such as scRNA-seq and spatially-resolved

counts [47–49]. Indeed, we observed a systematic difference of transcript detection

rate across platforms (**Fig. 1A-D**), which distorts the resulting transcriptional

profile estimates. Consequently, a panel of genes selected based on scRNA-seq that

works well on distinguishing cell types may not achieve similar level of performance

when measured by targeted spatial transcriptomics.

To address this challenge, we estimate the level of gene expression distortion in

targeted spatial transcriptomics data relative to scRNA-seq and from the same

tissue using a Bayesian model (**Fig. S1, Methods**). Bayesian inference estimates the

posterior distribution of distortion magnitudes, which will be used to predict the

potential distortion levels for genes that have not yet been observed in a given

assay. Specifically, we assume platform effects are on a per gene basis. γ_i and c_i

represent gene specific multiplicative and additive platform effect for each gene i,

respectively. These distortion parameters are assumed to follow two normal

distributions with μ_γ, μ_c as mean and σ_γ, σ_c as standard deviation, respectively. The

posterior distribution of σ_γ and σ_c can be considered as a generalized description of

the magnitude of multiplicative and additive platform effects. We can use them to

sample the magnitudes of gene specific multiplicative and additive platform

distortions for unobserved genes. The model is fitted for a given pair of scRNA-seq

and targeted spatial transcriptomics platforms to account for the differences

between them.

To check the extent to which the model is able to capture platform biases, we used

three paired scRNA-seq and targeted spatial transcriptomics datasets: scRNA-seq

and MERFISH data from mouse hypothalamic preoptic region [24] (Moffit dataset),

scRNA-seq and osmFISH data from mouse cortex [26,59] (Codeluppi dataset), and

scRNA-seq and DARTFISH data from healthy human kidney [60] (Zhang dataset)

(**Methods, Table S1**). Fitting a model for each pair of datasets, we then performed

posterior predictive check, i.e., we simulated spatial transcriptomics measurements

from scRNA-seq data using the fitted Bayesian model (**Methods**). Comparisons of

the distribution of simulated and observed spatial transcriptomics measurements

demonstrated that the Bayesian model can accurately recapitulate the platform

effects from different pairs of technologies (**Fig. 1E, Fig. S2A-C**). The posterior

distributions of σ_γ and σ_c (**Fig. 1F-G**) on the three datasets showed distinct levels of

additive and multiplicative platform effects, indicating the need to account for

platform-specific properties during gene panel selection.

Gene panel selection using genetic algorithm

To take platform distortions into account during selection of the gene panels, we use

the platform specific Bayesian model to simulate spatial transcriptomics

measurements with distortions (**Methods**). The gene panels are optimized for their

ability to recover cell type labels from such simulated spatial measurements, rather

than the original scRNA-seq measurements. Such an approach is intended to

provide a more accurate estimation of panel performance in a real spatially-

resolved measurement. Instead of selecting top-performing genes, gpsFISH

optimizes the entire gene panel in its combined ability to recover cell type labels. To

optimize within this combinatorial gene space, gpsFISH uses genetic algorithm

optimizer [61,62] (**Fig. 2, Methods**).

Within each iteration of optimization, multiple cross validations of classification are

performed for each proposed gene panel. To avoid biasing towards a specific

realization of spatial transcriptomics distortions, gpsFISH performs the platform

simulations separately in each cross validation. As a result, gene panels that are

more robust to unexpected platform distortions will be favored. This gene panel

selection framework ensures the evaluation is reflective of the gene panel's real

classification performance when measured by specific targeted spatial

transcriptomics technologies.

We first tested gpsFISH on the mouse hypothalamic scRNA-seq data (Moffitt

dataset) with simulated platform effect by optimizing a 200 gene panel to

distinguish "level 1" cell type annotation, which includes 12 broadly defined cell

types (**Fig 3A**). Most of the cells are correctly classified, yielding an overall accuracy

of 0.983 and high area under the receiver-operator curve (AUC) across all cell types

(**Fig 3B, C**). The optimized gene panel selected with considering platform effect was

also more successful in separating the 12 cell types on the resulting UMAP

embedding compared to the gene panel selected without considering platform effect

(**Fig. 3D, E**).

To quantify performance of different methods we simulated spatial transcriptomics

data from scRNA-seq, separating training and test sets (**Methods**). Simulations

were performed both with and without distortions in order to evaluate how taking

platform effects into account impacts gene panel performance. We also compared

gpsFISH with two previously published classification-based gene selection methods:

RankCorr and scGenefit. Both methods rely on the scRNA-seq expression profiles

without considering platform effects. RankCorr is a rank-based one-vs-all feature

selection method that selects marker genes for each cell type given a sparsity

parameter, which controls the number of marker genes selected per cell type. We

tuned this parameter to make sure the panels generated using RankCorr have the

same size (200 genes). scGenefit selects gene markers that jointly optimize cell label

recovery using label-aware compressive classification methods. As control, we

provided a naïve way to simulate spatial transcriptomics measurements without

platform effects (**Methods**). In addition, we also generated a panel of randomly

selected genes as baseline.

The objective of gpsFISH optimization is to achieve high quality cell type

classification on the spatial transcriptomics data. This entails two tasks: (1)

selecting a good gene panel, and (2) using the gene panel for accurate cell type

classification. In practice, while the design of an initial gene panel may rely on the

scRNA-seq data, optimization of subsequent panels can take advantage of the probe-

specific distortions that have already been observed in earlier measurements.

Similarly, as more and more spatial transcriptomics data are generated, when

classifying cell types in a newly generated spatially-resolved measurement, it is

likely that some partial annotations may already be available for that platform

either on the current or previously acquired datasets. Regardless of the cell type

granularity of partial annotation, it contains gene-specific platform effect

information of genes in the spatial transcriptomics data, which can be estimated

using our Bayesian model to improve cell type classification. Following this logic, we

used two benchmark strategies, which evaluate the impact of platform effects on the

two tasks (**Methods**). Both strategies share the same general framework in which a

classifier is trained on the training data with gene expression profiles for all cell

types, and then applied onto the testing data for cell type classification. The

difference is how the two strategies incorporate partial annotation into the training

data when available. Specifically, for the first strategy, evaluation with platform

effect re-estimation (**Methods**), platform effects are estimated from the partial

annotation data and incorporated into the training data for all gene selection

methods. Since under this strategy the gene panels from all methods are evaluated

in the same manner, it is useful in evaluating the impact of platform effect on the

first task, i.e., selecting a good gene panel. In contrast, under the second evaluation

strategy, evaluation without platform effect re-estimation (**Methods**), only gpsFISH

panels are evaluated with platform effect estimation as described above (**Methods**),

illustrating the impact of platform effects on both tasks.

Evaluation with platform effect re-estimation on the Moffit dataset using naïve

Bayes as the classifier shows gpsFISH outperforms the control with naïve simulation

and other gene selection methods (**Fig. 4A**), indicating that taking platform effects

into consideration leads to more informative gene panels. Similar results were

observed for the Zhang and Codeluppi dataset (**Fig. S3A and S3C**) and using random

forest as classifier (**Fig. S4A-C**). From the normalized confusion matrix of the gene

panel selected by gpsFISH with hierarchical tree on the left showing the relationship

between cell types (**Fig. 4C**) we can see that most of the misclassifications are

within the complex subpopulations of inhibitory and excitatory neurons.

A larger performance improvement of gpsFISH over other gene selection methods is

observed using evaluation without platform effect re-estimation, especially when

the level of partial annotation is low (**Fig. 4B, Fig. S3B and S3D, Fig. S4D-F**),

indicating that considering platform effects can lead to more accurate cell type

classification.

Overall, the comparison results show that gpsFISH outperforms other gene selection

methods and considering platform effects can result in more informative gene

panels and better cell type classification.

Redundancy in gene space across independent gene panel optimizations

enables incorporation of customized preferences

Independent panel optimizations performed multiple times (10) for each of the

three datasets showed high level of redundancy in the gene space (**Fig. 5A**).

Specifically, despite similar levels of overall performance, the overlap between

independently optimized 200-gene panels was around 85, 65, and 35 genes, and

more than 20%, 30%, and 45% of the genes showed up in only one of the 10

optimized gene panels for the Zhang, Moffit, and Codeluppi datasets, respectively

(**Fig. S5A-C**). We observed similar level of redundancy even when the optimization

was performed for a more granular "level 2" cell type annotations (46, 87, and 47

cell types for Zhang, Moffit, and Codeluppi dataset, **Fig. S5D**). The ability to achieve

similar level of performance with different gene sets suggests that the panels can be

further optimized to accommodate secondary criteria, such as inclusion of pre-

selected genes, emphasis on genes with specific features or from specific pathways,

etc.

gpsFISH allows to incorporate secondary preferences during gene panel

optimization, by specifying custom gene weights. To illustrate how panel

redundancy can be used to incorporate secondary preferences with little impact on

the classification performance, we evaluated the ability to increase the number of

technical probes per gene. Specifically, many ISH-based assays, including DARTFISH,

can include multiple different probes to enhance detection of any given transcript.

The number of probes that can be designed to target each gene is determined by

gene-specific factors like gene length. Genes with more probes are preferred, as they

can be used to improve robustness and sensitivity of detection. To generate a gene

panel with high number of potential probes, we used the predicted number of

probes for each gene in the DARTFISH data (Zhang dataset) (**Methods**) as gene

weight during gene panel selection. Of note, we capped the probe count at 15 to

avoid bias towards a small portion of genes with extremely high number of probes

(**Fig. S6A**). This also agrees with the fact that sensitivity will saturate when we have

enough probes for a gene.

Following this approach, we performed 10 optimizations with and without probe

count gene weights on the Zhang dataset using "level 1" cell type annotations. As

expected, the optimizations with gene weight had slightly lower accuracy (**Fig. 5B**)

but achieved a significantly higher number of total probes (**Fig. 5C**). This

demonstrates that the redundancy of gene spaces allows one to incorporate

additional customized constraints/preferences based on orthogonal information to

design gene panels with preferred features without sacrificing the overall cell type

classification performance.

Hierarchical gene selection based on cell type hierarchy

Cell types are organized in a hierarchical manner with broad cell types divided into

more detailed subpopulations. This hierarchical relationship can be considered

when evaluating cell classification errors. For example, failure to distinguish two

closely related subtypes, such as Th1 and Th17, of cells is likely to be considered

less severe than mis-annotation of a Th cell into a different major cell type such as B

cells.

In addition to the default "flat" cell type evaluation, gpsFISH therefore, implements a

hierarchical classification option (**Fig. 6A, Methods**), in which correct classifications

or misclassification between different cell types will receive different credit/penalty

specified by a weighted penalty matrix according to cell type hierarchy. Using this

hierarchical classification framework, gpsFISH provides flexibility to customize

optimization based on desired level of cell type granularity.

To evaluate the effect of the hierarchical classification for gene selection, we

performed hierarchical gene selection at level 2 cell annotation of all three datasets.

Under a hierarchical penalty scheme, misclassifications of cells between different

level 1 categories incur a fixed penalty, whereas misclassifications within the same

level 1 category were given partial credit, proportional to the expression similarity

between the called and true subtypes (**Methods, Fig. 6B**). To quantify to what

extent this hierarchical classification framework reduces misclassifications across

broad cell types at level 1, we calculated the percentage of across broad cell type

mistakes over all mistakes (**Methods**). We observed that the optimized gene panels

using hierarchical classification tend to make significantly fewer misclassifications

across broad cell types at level 1 compared to flat classification (**Fig. 6C-E**),

indicating that cell type granularity can be controlled through the hierarchical

classification framework.

Discussion and conclusions

Accurate cell type classification is crucial for understanding the spatial relationship

of cells in complex tissues. We implemented gpsFISH, a method for gene panel

design of targeted spatial transcriptomics. By accounting for platform effects

between scRNA-seq and targeted spatial transcriptomics technologies, gpsFISH is

able to find more robust and informative gene panels and achieve better cell type

classification.

Different technology has different patterns of platform effects. Specifically, we

decomposed platform effects into two components: multiplicative and additive

platform effects. While the multiplicative effect has been considered in

deconvolution contexts (e.g., RCTD [47]), neither type of platform-specific

distortions have been considered by other gene selection methods. Among other

things, the additive platform effect enables gpsFISH to describe situations where

specific genes show no expression in scRNA-seq data, but is detected in spatial

transcriptomics data (dots forming the vertical line in **Fig. 1A** and **1B**). This

observation is common for osmFISH (Codeluppi dataset) and MERFISH (Moffit

dataset), and cannot be modelled using only multiplicative platform effect.

Comparing the three targeted spatial transcriptomics platforms, we found highest

levels of additive platform effects in DARTFISH, followed by osmFISH and MERFISH

(**Fig. 1G**). More specifically, DARTFISH had the lowest μ_c, indicating the highest level

of signal reduction compared to MERFISH and osmFISH (**Fig. S2E**). Signal reduction

increases the possibility of good marker genes from scRNA-seq losing cell type

specificity in spatial transcriptomics data (dots forming the horizontal line in **Fig.**

1C), which is a main scenario where platform effects affect gene panel selection.

Higher level of signal reduction for DARTFISH agrees with our result that the

performance improvement of gpsFISH over other gene selection methods is the

largest in the Zhang dataset compared to the other two datasets, indicating the

necessity to account for additive platform effects, especially for targeted spatial

transcriptomics technologies with higher level of signal reduction.

In addition to additive platform effect, multiplicative platform effect also contributes

to the systematic difference of transcripts detection rate across technologies, posing

a challenge when transferring cell type information from scRNA-seq to spatial

transcriptomics technologies. Comparison of three targeted spatial transcriptomics

platforms shows osmFISH has the highest level of multiplicative platform effect,

followed by MERFISH and then DARTFISH (**Fig. 1F**). Higher level of multiplicative

platform effect leads to poorer cell type classification when there is no or low level

of partial annotation compared to high level of partial annotation (**Fig. 4A and 4B**,

Fig. S3 and S4), especially for evaluation without platform effect re-estimation due

to distorted expression profiles between scRNA-seq and targeted spatial

transcriptomics technologies. For evaluation with platform effect re-estimation, low

level of partial annotation provided limited statistical power to accurately estimate

gene specific platform effects, thus not able to increase the classification

performance. This reduced performance is gone when we have more than one cell

type included in the partial annotation, indicating partial annotation of a few cell

types is enough to enhance cell type classification if multiplicative platform effects

are accounted for.

Redundancy across independent optimizations allows incorporation of customized

preferences into gene selection. However, gene weight needs to be carefully

specified to ensure no sacrifice on overall gene panel performance. For the result in

Fig. 5B and **5C**, we capped the number of probes for each gene at 15. For cutoffs

lower than 15, gene weight difference between genes are small, leading to gene

panels with similar performance but also similar total number of probes. However,

for cutoffs higher than 15, the optimization will bias towards a small group of genes

with high probe count, resulting in local minimum during optimization (**Fig. S6B-C**).

This does achieve panels with significantly higher total number of probes, but the

classification accuracy is dropped. This emphasizes the need to test different ways

for gene weight specification in order to get the expected result without sacrificing

performance.

Similarly, in our test of hierarchical gene selection, we specified the weighted

penalty matrix directly from cell type hierarchy. Although we reduced

misclassifications across broad cell types, the overall accuracy is slightly lower than

flat classification (**Fig. S7**). This shows that partial credit of misclassifications needs

to be given carefully, especially when there are many similar subpopulations within

the same broad cell type like in the Moffit dataset. In real usage, it is suggested to

prune the weighted penalty matrix constructed from the cell type hierarchy to

remove unnecessary partial credit. Gene panel selection using flat classification can

be run first to help adjust the weighted penalty matrix constructed using cell type

hierarchy. In addition, the hierarchical classification provides a generic framework

to fine tune emphasis of classification on certain cell types. Here we showed its

usage to incorporate cell type hierarchy, but it is not restricted to cell type

hierarchy. Customized weighted penalty matrix can be constructed using other

information that provides preferences towards different classifications.

A major goal of spatial transcriptomics is to understand the spatial distribution of

cell types and their corresponding cellular environment. gpsFISH facilitates this by

selecting more informative and robust gene panels and providing ways for better

cell type annotation. We also provide options to account for various custom

preferences. As more targeted spatial transcriptomics data are generated, we expect

that gpsFISH can facilitate the study of cellular organization of complex tissues

under different biological contexts.

Methods

Datasets

In our study, we used three datasets that have both scRNA-seq and targeted spatial

transcriptomics data from the same tissue. Information regarding the three datasets

is summarized in **Table S1**. Further processing details are discussed below.

Moffit dataset

scRNA-seq data was downloaded from Gene Expression Omnibus (GEO) [63] under

accession code GSE113576. MERFISH data was downloaded from Dryad [64]. Of

note, the MERFISH data from Dryad is normalized and batch corrected. We undid

the volume normalization and batch correction to get the original data.

In the scRNA-seq data, we first filtered out cells annotated as "Ambiguous" and

"Unstable". We then used information in the supplementary Table 1 of the original

study to assign cell types. "Cell class (determined from clustering of all cells)"

column was used as level 1 cell type annotation. "Neuronal cluster (determined

from clustering of inhibitory or excitatory neurons)" and "Non-neuronal cluster

(determined from clustering of all cells)" were used as level 2 cell annotation.

Normalization was performed as described in the original study.

Only MERFISH data from naïve mice was used (to match scRNA-seq data). In

addition, we also filtered out cells annotated as "Ambiguous" and "Unstable". Fos

gene and five blank genes were filtered out. 135 genes imaged in the combinatorial

smFISH imaging were kept. Following the naming of cell types in Fig. 3D of the

original study, we modified the cell type annotation of MERFISH data to make it

consistent with the scRNA-seq data. Specifically, at level 1 cell type annotation, cells

annotated as "Endothelial 1", "Endothelial 2", "Endothelial 3" were merged into

"Endothelial". "Astrocyte" was changed to "Astrocytes". "OD Immature 1" and "OD

Immature 2" were changed to "Immature_oligodendrocyte". "OD Mature 1", "OD

Mature 2", "OD Mature 3", and "OD Mature 4" were changed to

"Mature_oligodendrocyte". "Pericytes" was changed to "Mural". At cell type level 2,

"Endothelial 1", "Endothelial 2", and "Endothelial 3" were changed to

"Endothelial_1", "Endothelial_2", and "Endothelial_3", respectively. "Ependymal" was

changed to "Ependymal_1". "OD Immature 1" and "OD Immature 2" were changed to

"Immature_oligodendrocyte_1" and "Immature_oligodendrocyte_2", respectively.

"OD Mature 1", "OD Mature 2", "OD Mature 3", and "OD Mature 4" were changed to

"Mature_oligodendrocyte_1", "Mature_oligodendrocyte_2",

"Mature_oligodendrocyte_3", and "Mature_oligodendrocyte_4", respectively.

After the processing above, additional filters were applied on the raw and

normalized scRNA-seq data before gene panel selection. Genes with maximum cell

type average expression lower than 1 were filtered out. In addition, long non-coding

RNAs were also removed. As a result, 2886 and 5100 genes were used for gene

panel selection at level 1 and 2, respectively. For platform effects estimation, the

subset of the raw scRNA-seq and MERFISH data with cells from overlapping cell

types were used.

Codeluppi dataset

scRNA-seq data was downloaded from GEO under accession code GSE60361.

Annotation data was downloaded from [65]. osmFISH and corresponding

annotation data was downloaded from [66].

For scRNA-seq data, cell labels in row 9 of the annotation were used as level 1 cell

type annotation, and row 11 were used as level 2 cell type annotation. However, the

level 1 cell type annotation is too broad (only 5 major cell types). Therefore, we

regenerated level 1 cell type annotation by merging similar cell types at level 2

following descriptions from the original study. Specifically, in generating data for

gene panel selection at level 1, "S1PyrDL", "S1PyrL23", "S1PyrL4", "S1PyrL5",

"S1PyrL5a", "S1PyrL6", S1PyrL6b", "ClauPyr" were merged into "S1_Excitatory".

"CA1Pyr1", "CA1Pyr2", "CA1PyrInt", "CA2Pyr2", "SubPyr" were merged into

"Hippocampus_Excitatory". 16 subclasses of interneurons ("Int1" to "Int16") were

merged into "Interneuron". "Astro1" and "Astro2" were merged into "Astrocyte".

"Mgl1" and "Mgl2" were merged into "Microglia". "Pvm1" and "Pvm2" were merged

into "Pvm". Six subpopulations of oligodendrocytes ("Oligo1" to "Oligo6") were

merged into "Oligodendorcyte". "Vend1" and "Vend2" were merged into

"Endothelial". To make cell type labels consistent between scRNA-seq and osmFISH,

"Peric" was changed to "Pericyte". "Choroid" was changed to "Ventricle". "Epend"

was changed to "Ependymal".

To generate the data for platform effect estimation, cell type labels were modified

slightly differently to reflect the correspondence between cell types in scRNA-seq

and osmFISH as shown in Fig. 2C and Fig. 2D of the original study. Specifically, three

CA1 subclasses ("CA1Pyr1", "CA1Pyr2", "CA1PyrInt") were merged into

"Hippocampus_Excitatory". 16 subclasses of interneurons ("Int1" to "Int16") were

merged into "Interneuron". Two subclasses of microglia ("Mgl1" and "Mgl2") were

merged into "Microglia". Two subclasses of perivascular macrophages ("Pvm1" and

"Pvm2") were merged into "Pvm". Subclasses of S1 pyramidal cells were also

merged: "S1PyrL4" and "S1PyrL5a" were merged into "S1_Excitatory_L45a",

"S1PyrL5" and "S1PyrL6b" were merged into "S1_Excitatory_L56b". In addition, to

make the cell type labels consistent between scRNA-seq and osmFISH, we changed

"Astro1" and "Astro2" to "Astrocyte1" and "Astrocyte2", respectively. We changed

"Oligo6" to "Oligo_Mature", "Oligo5" to "Oligo_MF", "Oligo1", to "Oligo_COP", "Vend1"

to "Endothelial1", "Vend2" to "Endothelial2", "Peric" to "Pericyte", "Choroid" to

"Ventricle", "Epend" to "Ependymal", "S1PyrL23" to "S1_Excitatory_L23", and

"S1PyrL6" to "S1_Excitatory_L6". Cell types with fewer than 50 cells were removed.

For osmFISH data, we first filtered out invalid cells based on the "Valid" column of

the annotation data. Then, similar to scRNA-seq data, we modified cell type labels

according to Fig. 2C and Fig.2D in the original study, which shows correspondence

between cell types in scRNA-seq and osmFISH. Specifically, "Astrocyte Gfap" was

changed to "Astrocyte1". "Astrocyte Mfge8" was changed to "Astrocyte2".

"Hippocampus" was changed to "Hippocampus_Excitatory". "pyramidal L4" was

changed to "S1_Excitatory_L45a". "Pyramidal L5" was changed to

"S1_Excitatory_L56b". "Pyramidal L6" was changed to "S1_Excitatory_L6".

"Perivascular Macrophages" was changed to "Pvm". "Oligodendrocyte COP" was

changed to "Oligo_COP". ""Oligodendrocyte Mature"" was changed to

"Oligo_Mature". "Oligodendrocyte MF" was changed to "Oligo_MF". "Endothelial 1"

was changed to "Endothelial1", and "Endothelial" was changed to "Endothelial2".

"Pericytes" was changed to "Pericyte". "Vascular Smooth Muscle" was changed to

"Vsmc", "C. Plexus" was changed to "Ventricle". "Pyramidal L2-3" and "Pyramidal L2-

3 L5" were merged into "S1_Excitatory_L23". "Inhibitory Cnr1", "Inhibitory CP",

"Inhibitory Crhbp", "Inhibitory IC", "Inhibitory Kcnip2", "Inhibitory Pthlh", and

"Inhibitory Vip" were merged into "Interneuron".

scRNA-seq data was normalized using the count_normalize function in the scran

package. Similar to the Moffit dataset, the raw and normalized scRNA-seq were

further filtered before gene panel selection using the same filters.

6123 and 9052 genes were used for gene panel selection at level 1 and 2,

respectively. For platform effect estimation, the subset of the raw scRNA-seq and

osmFISH data with cells from overlapping cell types were used.

Zhang dataset

Raw and normalized scRNA-seq data from kidney were obtained from [60]. They

were further filtered before gene panel selection using the same filters. 2920 and

3796 genes were used for gene panel selection at level 1 and 2, respectively.

The DARTFISH data is unpublished. It can be found in Zenodo [67]. We annotated

the cells in the DARTFISH data manually using curated marker genes (**Table S2**) at

subclass level (third column). For platform effect estimation, the subset of the raw

scRNA-seq and DARTFISH data with cells from overlapping cell types were used.

Platform effects estimation using a Bayesian model

We assume the observed number of molecules y_{ij} in the spatial transcriptomics

data for gene i in cell j follows a zero-inflated negative bimonial (ZINB) distribution

with:

$$y_{ij} \sim ZINB(\mu_{ij}, \theta_{ij}, \pi) \qquad (1)$$

537 where π is the zero inflation parameter which is assumed to be constant across

538 genes and cells. μ_{ij} is the mean parameter determined by a global intercept α, true

539 expression level of gene i in cell j denoted as λ_{ij}, and the cell depth (total number of

540 molecules) of cell j from spatial transcriptomics data as CD_j^{SP}:

$$\ln(\mu_{ij}) = \alpha + \ln(\lambda_{ij}) + \ln(CD_j^{SP}) \qquad (2)$$

544 To account for platform effects, we assume the true expression level λ_{ij} is a random

545 variable defined by:

547
$$\text{logit}(\lambda_{ij}) = \gamma_i \times \sqrt{x_{ij}} + c_i \qquad (3)$$

549 where γ_i is a gene specific coefficient representing multiplicative platform effects,

550 and c_i is a gene specific intercept representing additive platform effects. x_{ij}

551 represents the relative expression of gene i in cell j calculated from scRNA-seq data:

553
$$x_{ij} = \frac{c_{ij}}{\sum_{i=1}^{N} c_{ij}}$$

555 where c_{ij} is the number of count for gene i in cell j from the scRNA-seq data, and N

556 is the totol number of genes. When fitting the Bayesian model, in order to match

557 measurement between scRNA-seq data and targeted spatial transcriptomics data,

558 we used cell type average relative expression to replace individual cell level relative

559 expression:

561
$$x_{ij,j \subset k} = x_{ik} = \frac{\sum_{j=1}^{M_k} c_{ij}}{\sum_{i=1}^{N} \sum_{j=1}^{M_k} c_{ij}}$$

563 where M_k is the number of cells in cell type k.

565 For the dispersion parameter θ_{ij} of the ZINB distribution, we assume it is also

566 dependent on λ_{ij}:

568
$$\ln(\theta_{ij}) = \beta + \lambda_{ij} \qquad (4)$$

570 where β is the intercept.

572 A Beta prior distribution is assumed for π. For α, β, and c_i, we assume they follow

573 normal distribution. γ_i is assumed to follow a log-normal distribution:

575
$$\pi \sim \text{Beta}(1, 1)$$

576
$$\alpha \sim \text{Normal}(0, \sigma_\alpha)$$

577
$$\beta \sim \text{Normal}(0, \sigma_\beta)$$

578
$$c_i \sim \text{Normal}(\mu_c, \sigma_c)$$

579
$$\gamma_i \sim \text{LogNormal}(\mu_\gamma, \sigma_\gamma)$$

581 where the hyperparameters are assumed to follow Cauchy and half Cauchy

582 distribution:

584
$$\mu_c, \mu_\gamma \sim \text{Cauchy}(0, 5)$$

585
$$\sigma_\alpha, \sigma_\beta, \sigma_c, \sigma_\gamma \sim \text{HalfCauchy}(0, 5)$$

scRNA-seq and targeted spatial transcriptomics data from overlapping genes and

overlapping cell types were used as input. Additional filters were applied on the

MERFISH data to reduce the totol number of cells for more efficient estimation.

Specifically, cells with cell depth lower than 100 were filtered out. Cell types with

fewer than 1000 cells were filtered out. Then we subsampled each cell type to keep

at most 1000 cells for each cell type. Variational inference in Stan was used for

model fitting.

Simulation of spatial transcriptomics measurements from scRNA-seq data

with platform effects

We used fitted Bayesian models to simulate spatial transcriptomics measurements

from scRNA-seq data. Specifically, $\alpha, \beta, \pi, \mu_c, \sigma_c, \mu_\gamma, \sigma_\gamma$ were randomly sampled from

their estimated posterior distribution. c_i, and γ_i were randomly sampled from their

corresponding normal and log normal distribution for each new gene that is not

observed in the data used to fit the Bayesian model. If a gene is already seen during

fitting the Bayesian model, we can either use the empirical c_i, and γ_i estimated

during model fitting (used in this study) or randomly sample them from the

corresponding normal and log norml distribution. CD_j^{SP} was randomly sampled

from empirical cell depth distribution from observed targeted spatial

transcriptomics data. x_{ij} was calculated from scRNA-seq data. It can be cell type

average as we used in model fitting or calculated within each individual cell. In our

study, the latter was used when simulating spatial transcriptomics measurements to

maintain the cell level heterogenity in scRNA-seq data. Finally, the generated values

were plugged into equations (1), (2), (3), and (4) to generate spatial transcriptomics

measurements.

Simulation of spatial transcriptomics measurements from scRNA-seq data

without platform effects (naïve simulation)

We provided a naïve way to simulate spatial transcriptomics measurements without

platform effects as control. During the simulation without platform effects, cell

depth of simulated spatial transcriptomics cell were randomly sampled from the

empirical cell depth distribution of observed targeted spatial transcriptomics data.

Of note, the empirical cell depth distribution was adjusted proportionally based on

the ratio between relative expression of new genes for simulation and relative

expression of overlapping genes used in fitting the Bayesian model. After having the

simulated cell depth for each cell, the number of molecules for each gene within

each cell was sampled from a multinomial distribution with size equal to the

simulated cell depth and probability equal to each gene's relative expression in that

cell. At the end, genes were randomly selected given the probe failure rate. Then,

simulated molecule count of selected genes were set to 0 to reflect probe failure.

Genetic algorithm for gene panel selection

We used genetic algorithm as the framework for gene panel selection. Each

individual in a population is one candidate gene panel. We set the gene panel size to

200 genes. Each population contains 200 individuals.

The first step of genetic algorithm is to initialize a population of candidate gene

panels. The genes can be either randomly selected from all candidate genes or

selected based on their differential expression between cell types. In this study, we

took a hybrid approach. 95% of the 200 gene panels were initiated randomly from

all candidate genes to maintain population diversity. The rest 5% were initialized

using DEGs for each cell type. DE analysis was performed using Pagoda2. Genes

with AUC greater than 0.7 were considered significant.

The second step is to evaluate the fitness of each candidate gene panel in the

population. Here we define fitness as the average classification accuracy over 5

cross validations. Classification was performed on simulated spatial transcriptomics

measurements from scRNA-seq data. Cell type annotation from scRNA-seq data was

used as ground truth. The accuracy was calculated based on the original confusion

matrix for flat classification and weighted confusion matrix for hierarchical

classification. We provided two classifiers, random forest and naïve Bayes. In this

study we used naïve Bayes due to its fast speed and relatively similar level of

accuracy compared to random forest. To improve the efficiency, scRNA-seq data was

subsampled to reduce the number of cells for large cell types and resampled to

increase the number of cells for small cell types. Specifically, for level 1 cell type

annotation, cell type size was capped at 1500 cells. The lower bound was set as

1000 cells for Moffit dataset and 500 for Zhang and Codeluppi dataset. For level 2

cell type annotation, 250 and 500 were used as the cell type size range for Moffit

dataset. The range for Zhang and Codeluppi dataset was 300 and 900.

The third step is selection and mutation. The selection strategy we used is

tournaments. Specifically, randomly selected candidate gene panels face each other

1 vs. 1. The one with a higher fitness value was used as parent. In addition,

candidate gene panels with higher fitness values were more likely to be selected in

the tournaments. After having the parent gene panels, uniform crossover was

performed to generate the offspring gene panels. Duplicated genes after uniform

crossover were replaced by randomly sampled genes in the parent candidate gene

panels but not in the offspring gene panel. Mutation was then performed to maintain

gene diversity and prevent premature convergence. We set the mutation rate to 1%.

When gene weight was provided, genes with higher weight were (1) more likely to

be selected during crossover, (2) less likely to be mutated if it is already in the

population, (3) and more likely to be introduced into the population through

mutation if it is not in the current population.

Finally, the same process was repeated for the offspring population. The candidate

gene panel with the highest fitness value for one iteration was considered as the

optimal gene panel. If the iteration after it has a candidate gene panel with higher

fitness value, the optimal panel will be replaced by this new candidate gene panel.

Otherwise, the optimal panel will stay the same. The iterative process will end either

when it reaches a given number of iterations, or the accuracy doesn't improve more

than a threshold for a given number of iterations. In our study, we ran all the

optimizations for at least 500 iterations to ensure convergence although in all cases

the optimization converged a lot earlier.

If a list of pre-selected genes, e.g., canonical marker genes based on previous

knowledge, is provided, genes in the list will be included in each candidate gene

panel as well as the final optimal gene panel.

Hierarchical classification using cell type hierarchy

During genetic algorithm optimization, a weighted penalty matrix can be provided

to assign partial credit or extra penalty to classification between certain cell types.

The weighted penalty matrix is a square matrix with each row and each column

representing one cell type. For each value p_{ij} ($i \neq j$) in the weighted penalty matrix,

if $p_{ij} > 1$, an extra penalty is given to misclassifying cells from cell type j to cell type

i. If $p_{ij} < 1$, a partial credit is given to misclassifying cells from cell type j to cell type

i. $p_{ij} = 1$ means no penalty or partial credit. In hierarchical classification, the

weighted penalty matrix was incorporated to the confusion matrix by element-wise

multiplication to provide a weighted confusion matrix. The accuracy of the weighted

confusion matrix was used to evaluate the fitness of candidate gene panels.

Essentially, the weighted penalty matrix can be constructed arbitrarily by user's

preference. In this study, we constructed the weighted penalty matrix from cell type

hierarchy. First, pairwise distance between cell types was calculated. Specifically,

average expression profile of each cell type was calculated using normalized count

by taking average expression of all cells in each cell type. Top 1000 genes with

highest standard deviation were used to calculate pairwise Pearson correlation

coefficient. One minus the pairwise Pearson correlation coefficient was used as

pairwise distance between cell types. Second, the pairwise distance matrix was

normalized by the largest distance so the values range from 0 to 1. Third, the

pairwise distance matrix was then adjusted based on cell type hierarchy.

Specifically, a level of cell type annotation was selected as reference. For cell types

below the reference level that are from the same cell type at the reference level, the

pairwise distance (between 0 and 1) between them was kept unchanged to reflect

partial credit to wrong classifications among them. For cell types below the

reference level that are from different cell types at the reference level, the pairwise

distance between them was changed to a user defined value where 1 means no extra

penalty and greater than 1 means extra penalty. In this study, we used 1 for no extra

penalty and level 1 cell type annotation was used as reference. Finally, the diagonal

value was changed to 1 to reflect no extra credit to correct classifications. This

weighted penalty matrix was used for hierarchical classification in our study.

Calculating the percentage of across broad cell type mistakes over all mistakes

We performed 5 optimizations with flat classification and hierarchical classification

for all three datasets, respectively. Average confusion matrix over 5 optimizations

for each data was calculated. After that, for each cell type, we counted the total

number of misclassifications and among all the misclassifications, what proportion

of them misclassifies cells to cell types at level 2 that don't belong to the same cell

type at level 1.

Gene panel selection using RankCorr and scGeneFit

The same scRNA-seq data from the three datasets after filtering were used as input.

For RankCorr, raw scRNA-seq data before normalization was used as suggested. The

lamb parameter was tuned to make sure the output marker gene list has 200 genes.

For scGeneFit, normalized scRNA-seq data was used by following the examples on

its GitHub page. Panel size was set to 200.

Evaluation of optimized gene panel

To evaluate optimized gene panels, we first simulated spatial transcriptomics

measurements with platform effects based on the gene panel's expression profile in

scRNA-seq data. Then this simulated spatial transcriptomics data was split into

training and testing data. The training data contains cells from a subset of cell types

whose cell type labels are known. This was used as the partial annotation of the

simulated spatial transcriptomics data. The testing data contains cells from all cell

types (excluding cells in the training data), which is considered as part of the

simulated spatial transcriptomics data that hasn't been annotated yet. We varied the

number of cell types in the training data from zero to all the cell types to reflect

different levels of partial annotation. When there was no partial annotation, scRNA-

seq data was used as the final training data for classifier training. When there was

partial annotation, information in the partial annotation was included in the final

training data. After that, a classifier (naïve Bayes or random forest) was trained

using the final training data and applied on the testing data for cell type

classification evaluation. Since the testing data was simulated from scRNA-seq data,

the cell type labels in scRNA-seq data were used as ground truth. Classification

accuracy was used as the metric to evaluate a gene panel. At each level of partial

annotation, we repeated the same calculation 10 times. To separately evaluate the

impact of platform effects on gene panel selection and cell type classification, within

the same framework described here, we designed two different strategies to

evaluate a gene panel by varying whether platform effect distortions that can be

learned from partial annotation examples are used to produce more realistic

training data for cell type classification.

Evaluation with platform effect re-estimation

This evaluation strategy was designed to focus on the performance of the optimized

gene panels, and not on the differences in the cell type classification (evaluation)

stage. In this strategy, partial annotation was first used to estimate gene specific

platform effects using the Bayesian model. We then used these estimated gene

specific platform effects to simulate an updated spatial transcriptomics training

data, which will be combined with the partially annotated spatial transcriptomics

data and then used for training cell type classifiers for all the methods being

evaluated. Only cell types not already available in the partially annotated spatial

transcriptomics data were simulated. When partial annotation was available for 5 or

fewer cell types, the final training data combined the partially annotated and

simulated spatial transcriptomics training data with scRNA-seq data. When more

than 5 cell types were available, training was performed on the partially annotated

and simulated spatial transcriptomics training data only. The final training data and

testing data were normalized by the total number of transcripts within each cell and

scaled by 10000. It was then log transformed after adding 1 pseudocount. This

normalized training and testing data were used for classifier training and testing.

Evaluation without platform effect re-estimation

In this evaluation strategy, only gpsFISH is able to make use of the platform effects

information in the partial annotation (as described above). All the other methods

used the partial annotation according to their own method design. Specifically, for

the control which used naïve simulation during gene panel selection, the empirical

cell depth distribution of the complete testing data was used to simulate a spatial

transcriptomics training data without platform effect. This simulated spatial

transcriptomics training data was used in the same way as described above to get

the final training data. For RankCorr, scGeneFit, and the random panel, since the

gene selection was solely based on scRNA-seq data, cells in the partial annotation

were directly combined with the scRNA-seq data of cell types not already available

in the partial annotation. The combined data were used as the final training data.

Same normalization was performed on the final training data and testing data

before classifier training and testing.

Calculate the number of probes for each gene for the DARTFISH data

During the generation of the DARTFISH data, ppDesigner [68] was used to calculate

the number of probes that can be designed to target each gene.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Scripts to generate data and to perform the above analysis are available in Zenodo

[67].

gpsFISH's open-source code is maintained and documented on Github [69] and is

publicly available under the MIT license.

Pre-fitted Bayesian models based on the Zhang, Moffit, and Codeluppi dataset

respectively are deposited in Zenodo [70].

Competing interests

P.V.K. serves on the Scientific Advisory Board to Celsius Therapeutics Inc. and

Biomage Inc. P.V.K. is an employee of Altos Labs.

Funding

P.V.K. and Y.Z. were supported by 5U54HL145608 grant from NIH.

Authors' contributions

P.V.K. and Y.Z. formulated the study and the overall approach. Y.Z. developed the

detailed algorithms and performed the analysis with advice from P.V.K. and V.P. Y.Z.

implemented the gpsFISH package with help from E.B. R.Q. and K.Z. generated the

DARTFISH dataset. Y.Z. and P.V.K. drafted the manuscript.

Acknowledgements

We thank Hirak Sarkar and Teng Gao for useful discussions on the method

development. We also thank Chienju Chen and Kian Kalhor for useful discussions on

the DARTFISH technology and data processing. We are also grateful to Jeffrey

Moffitt who provided help with obtaining the original MERFISH data before

normalization and batch correction. P.V.K. serves on the Scientific Advisory Board to

Celsius Therapeutics Inc. and Biomage Inc. P.V.K. is an employee of Altos Labs.

Supplementary Information

Additional file 1: Table S1.xlsx

Information of the Moffit, Codeluppi, and Zhang dataset

Additional file 2: Table S2.xlsx

Curated marker genes for the Zhang dataset

Reference

1. Arendt D. The evolution of cell types in animals: emerging principles from
molecular studies. Nat Rev Genet. 2008;9:868–82.

2. Elmentaite R, Domínguez Conde C, Yang L, Teichmann SA. Single-cell atlases:
shared and tissue-specific cell types across human organs. Nat Rev Genet.
2022;23:395–410.

3. Lindeboom RGH, Regev A, Teichmann SA. Towards a Human Cell Atlas: Taking
Notes from the Past. Trends in Genetics. 2021;37:625–30.

4. Zeng H. What is a cell type and how to define it? Cell. 2022;185:2739–55.

5. Kölsch Y, Hahn J, Sappington A, Stemmer M, Fernandes AM, Helmbrecht TO, et al.
Molecular classification of zebrafish retinal ganglion cells links genes to cell types to
behavior. Neuron. 2021;109:645-662.e9.

6. Osterhout JA, Kapoor V, Eichhorn SW, Vaughn E, Moore JD, Liu D, et al. A preoptic
neuronal population controls fever and appetite during sickness. Nature.
2022;606:937–44.

7. Xu S, Yang H, Menon V, Lemire AL, Wang L, Henry FE, et al. Behavioral state
coding by molecularly defined paraventricular hypothalamic cell type ensembles.
Science. 2020;370:eabb2494.

8. Elmentaite R, Kumasaka N, Roberts K, Fleming A, Dann E, King HW, et al. Cells of
the human intestinal tract mapped across space and time. Nature. 2021;597:250–5.

9. Armingol E, Officer A, Harismendy O, Lewis NE. Deciphering cell–cell interactions
and communication from gene expression. Nat Rev Genet. 2021;22:71–88.

10. Chen W-T, Lu A, Craessaerts K, Pavie B, Sala Frigerio C, Corthout N, et al. Spatial
Transcriptomics and In Situ Sequencing to Study Alzheimer's Disease. Cell.
2020;182:976-991.e19.

11. Hwang WL, Jagadeesh KA, Guo JA, Hoffman HI, Yadollahpour P, Reeves JW, et al.
Single-nucleus and spatial transcriptome profiling of pancreatic cancer identifies
multicellular dynamics associated with neoadjuvant treatment. Nature Genetics
[Internet]. 2022; Available from: https://doi.org/10.1038/s41588-022-01134-8

12. Darmanis S, Sloan SA, Zhang Y, Enge M, Caneda C, Shuer LM, et al. A survey of
human brain transcriptome diversity at the single cell level. Proc Natl Acad Sci USA.
2015;112:7285–90.

13. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly
Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter
Droplets. Cell. 2015;161:1202–14.

14. Tasic B, Menon V, Nguyen TN, Kim TK, Jarsky T, Yao Z, et al. Adult mouse cortical
cell taxonomy revealed by single cell transcriptomics. Nat Neurosci. 2016;19:335–
46.

15. Ner-Gaon H, Melchior A, Golan N, Ben-Haim Y, Shay T. JingleBells: A Repository
of Immune-Related Single-Cell RNA–Sequencing Datasets. JI. 2017;198:3375–9.

16. Trapnell C. Defining cell types and states with single-cell genomics. Genome Res.
2015;25:1491–8.

17. Regev A, Teichmann SA, Lander ES, Amit I, Benoist C, Birney E, et al. The Human
Cell Atlas. eLife. 2017;6:e27041.

18. Marx V. Method of the Year: spatially resolved transcriptomics. Nat Methods.
2021;18:9–14.

19. Chen R, Blosser TR, Djekidel MN, Hao J, Bhattacherjee A, Chen W, et al. Decoding
molecular and cellular heterogeneity of mouse nucleus accumbens. Nat Neurosci.
2021;24:1757–71.

20. Fernandez J. Molecular atlas of the adult mouse brain. SCIENCE ADVANCES.
2020;14.

21. Rao A, Barkley D, França GS, Yanai I. Exploring tissue architecture using spatial
transcriptomics. Nature. 2021;596:211–20.

22. Zhang M, Eichhorn SW, Zingg B, Yao Z, Cotter K, Zeng H, et al. Spatially resolved
cell atlas of the mouse primary motor cortex by MERFISH. Nature. 2021;598:137–
43.

23. Wang Y, Eddison M, Fleishman G, Weigert M, Xu S, Wang T, et al. EASI-FISH for
thick tissue defines lateral hypothalamus spatio-molecular organization. Cell.
2021;184:6361-6377.e24.

24. Moffitt JR, Bambah-Mukku D, Eichhorn SW, Vaughn E, Shekhar K, Perez JD, et al.
Molecular, spatial, and functional single-cell profiling of the hypothalamic preoptic
region. Science. 2018;362:eaau5324.

25. Chen KH, Boettiger AN, Moffitt JR, Wang S, Zhuang X. Spatially resolved, highly
multiplexed RNA profiling in single cells. Science. 2015;348:aaa6090–aaa6090.

26. Codeluppi S, Borm LE, Zeisel A, La Manno G, van Lunteren JA, Svensson CI, et al.
Spatial organization of the somatosensory cortex revealed by osmFISH. Nat
Methods. 2018;15:932–5.

27. Shah S, Lubeck E, Zhou W, Cai L. In Situ Transcription Profiling of Single Cells
Reveals Spatial Organization of Cells in the Mouse Hippocampus. Neuron.
2016;92:342–57.

28. Cai M, Zhang K. Spatial mapping of single cells in human cerebral cortex using
DARTFISH: A highly multiplexed method for in situ quantification of targeted RNA
transcripts. eScholarship, University of California; 2019.

29. Wang X, Allen WE, Wright MA, Sylwestrak EL, Samusik N, Vesuna S, et al. Three-
dimensional intact-tissue sequencing of single-cell transcriptional states. Science.
2018;361:eaat5691.

30. Chen X, Sun Y-C, Church GM, Lee JH, Zador AM. Efficient in situ barcode
sequencing using padlock probe-based BaristaSeq. Nucleic Acids Research.
2018;46:e22–e22.

31. Chen X, Sun Y-C, Zhan H, Kebschull JM, Fischer S, Matho K, et al. High-
Throughput Mapping of Long-Range Neuronal Projection Using In Situ Sequencing.
Cell. 2019;179:772-786.e19.

32. Gyllborg D, Langseth CM, Qian X, Choi E, Salas SM, Hilscher MM, et al.
Hybridization-based *in situ* sequencing (HybISS) for spatially resolved
transcriptomics in human and mouse brain tissue. Nucleic Acids Research.
2020;48:e112–e112.

33. Chen A, Liao S, Cheng M, Ma K, Wu L, Lai Y, et al. Spatiotemporal transcriptomic
atlas of mouse organogenesis using DNA nanoball-patterned arrays. Cell.
2022;185:1777-1792.e21.

34. Ståhl PL, Salmén F, Vickovic S, Lundmark A, Navarro JF, Magnusson J, et al.
Visualization and analysis of gene expression in tissue sections by spatial
transcriptomics. Science. 2016;353:78–82.

35. Rodriques SG, Stickels RR, Goeva A, Martin CA, Murray E, Vanderburg CR, et al.
Slide-seq: A scalable technology for measuring genome-wide expression at high
spatial resolution. 2019;6.

36. Vickovic S, Eraslan G, Salmén F, Klughammer J, Stenbeck L, Schapiro D, et al.
High-definition spatial transcriptomics for in situ tissue profiling. Nat Methods.
2019;16:987–90.

37. Liu Y, Yang M, Deng Y, Su G, Enninful A, Guo CC, et al. High-Spatial-Resolution
Multi-Omics Sequencing via Deterministic Barcoding in Tissue. Cell. 2020;183:1665-
1681.e18.

38. Cho C-S, Xi J, Si Y, Park S-R, Hsu J-E, Kim M, et al. Microscopic examination of
spatial transcriptome using Seq-Scope. Cell. 2021;184:3559-3572.e22.

39. Fu X, Sun L, Chen JY, Dong R, Lin Y, Palmiter RD, et al. Continuous Polony Gels for
Tissue Mapping with High Resolution and RNA Capture Efficiency. bioRxiv.
2021;2021.03.17.435795.

40. Subramanian A, Narayan R, Corsello SM, Peck DD, Natoli TE, Lu X, et al. A Next
Generation Connectivity Map: L1000 Platform and the First 1,000,000 Profiles. Cell.
2017;171:1437-1452.e17.

41. Missarova A, Jain J, Butler A, Ghazanfar S, Stuart T, Brusko M, et al. geneBasis: an
iterative approach for unsupervised selection of targeted gene panels from scRNA-
seq. Genome Biology. 2021;22:333.

42. Liang S, Mohanty V, Dou J, Miao Q, Huang Y, Müftüoğlu M, et al. Single-cell
manifold-preserving feature selection for detecting rare cell populations. Nat
Comput Sci. 2021;1:374–84.

43. Dumitrascu B, Villar S, Mixon DG, Engelhardt BE. Optimal marker gene selection
for cell type discrimination in single cell analyses. Nat Commun. 2021;12:1186.

44. Vargo AHS, Gilbert AC. A rank-based marker selection method for high
throughput scRNA-seq data. BMC Bioinformatics. 2020;21:477.

45. Aevermann BD, Zhang Y, Novotny M, Keshk M, Bakken TE, Miller JA, et al. A
machine learning method for the discovery of minimum marker gene combinations
for cell-type identification from single-cell RNA sequencing. Genome Res.
2021;gr.275569.121.

46. Bakken TE, Hodge RD, Miller JA, Yao Z, Nguyen TN, Aevermann B, et al. Single-
nucleus and single-cell transcriptomes compared in matched cortical cell types.
Soriano E, editor. PLoS ONE. 2018;13:e0209648.

47. Cable DM, Murray E, Zou LS, Goeva A, Macosko EZ, Chen F, et al. Robust
decomposition of cell type mixtures in spatial transcriptomics. Nat Biotechnol.
2022;40:517–26.

48. Okochi Y, Sakaguchi S, Nakae K, Kondo T, Naoki H. Model-based prediction of
spatial gene expression via generative linear mapping. Nat Commun. 2021;12:3731.

49. Andersson A, Bergenstråhle J, Asp M, Bergenstråhle L, Jurek A, Fernández
Navarro J, et al. Single-cell and spatial transcriptomics enables probabilistic
inference of cell type topography. Commun Biol. 2020;3:565.

50. the FANTOM Consortium, Liang C, Forrest ARR, Wagner GP. The statistical
geometry of transcriptome divergence in cell-type evolution and cancer. Nat
Commun. 2015;6:6066.

51. Pliner HA, Shendure J, Trapnell C. Supervised classification enables rapid
annotation of cell atlases. Nat Methods. 2019;16:983–6.

52. Tasic B. Single cell transcriptomics in neuroscience: cell classification and
beyond. Current Opinion in Neurobiology. 2018;50:242–9.

53. Zeng H, Sanes JR. Neuronal cell-type classification: challenges, opportunities and
the path forward. Nat Rev Neurosci. 2017;18:530–46.

54. Yuste R, Hawrylycz M, Aalling N, Aguilar-Valles A, Arendt D, Armañanzas R, et al.
A community-based transcriptomics classification and nomenclature of neocortical
cell types. Nat Neurosci. 2020;23:1456–68.

55. Bard J, Rhee SY, Ashburner M. An ontology for cell types. Genome Biology.
2005;5.

56. Bakken T, Cowell L, Aevermann BD, Novotny M, Hodge R, Miller JA, et al. Cell
type discovery and representation in the era of high-content single cell phenotyping.
BMC Bioinformatics. 2017;18:559.

57. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene
Ontology: tool for the unification of biology. Nat Genet. 2000;25:25–9.

58. The Gene Ontology Consortium, Carbon S, Douglass E, Good BM, Unni DR, Harris
NL, et al. The Gene Ontology resource: enriching a GOld mine. Nucleic Acids
Research. 2021;49:D325–34.

59. Zeisel A, Muñoz-Manchado AB, Codeluppi S, Lönnerberg P, La Manno G, Juréus A,
et al. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-
seq. Science. American Association for the Advancement of Science; 2015;347:1138–
42.

60. Lake BB, Menon R, Winfree S, Hu Q, Ferreira RM, Kalhor K, et al. An atlas of
healthy and injured cell states and niches in the human kidney. bioRxiv.
2021;2021.07.28.454201.

61. Goldberg DE. Genetic Algorithms in Search, Optimization and Machine Learning.
1st ed. USA: Addison-Wesley Longman Publishing Co., Inc.; 1989.

62. Holland JH. Adaptation in Natural and Artificial Systems: An Introductory
Analysis with Applications to Biology, Control, and Artificial Intelligence [Internet].
The MIT Press; 1992 [cited 2022 Jul 30]. Available from:
https://doi.org/10.7551/mitpress/1090.001.0001

63. Gene Expression Omnibus. www.ncbi.nlm.nih.gov/geo

64. Dryad. https://datadryad.org/stash/dataset/doi:10.5061/dryad.8t8s248

65. Zeisel A, Muñoz-Manchado AB, Codeluppi S, Lönnerberg P, La Manno G, Juréus A,
et al. Single-cell analysis of mouse cortex. http://linnarssonlab.org/cortex/

66. Codeluppi S, Borm LE, Zeisel A, La Manno G, van Lunteren JA, Svensson CI, et al.
osmFISH: Spatial organization of the somatosensory cortex revealed by cyclic
smFISH. http://linnarssonlab.org/osmFISH/

67. Zhang Y, Petukhov V, Biederstedt E, Que R, Zhang K, Kharchenko PV. gpsFISH
analysis code and data (Zenodo link). 2023;
https://doi.org/10.5281/zenodo.7613712

68. ppDesigner: Algorithm to design Padlock Probes. http://genome-
tech.ucsd.edu/public/Gen2_BSPP/ppDesigner/ppDesigner.php

69. Zhang Y, Petukhov V, Biederstedt E, Que R, Zhang K, Kharchenko PV. gpsFISH R
package. GitHub. 2023; https://github.com/kharchenkolab/gpsFISH

70. Zhang Y, Petukhov V, Biederstedt E, Que R, Zhang K, Kharchenko PV. Pre-fitted
Bayesian models (Zenodo link). 2023; https://doi.org/10.5281/zenodo.6946054

1022

1023

1024

1025

Fig. 1

$y_{ij} \sim ZINB(\mu_{ij}, \theta_{ij}, \pi)$
$\ln(\mu_{ij}) = \alpha + \ln(\lambda_{ij}) + \ln(cell\ depth_j)$
$\text{logit}(\lambda_{ij}) = \gamma_i \times \sqrt{x_{ij}} + c_i$

Fig. 2

Fig. 3

A Evaluation with platform effect re-estimation

B Evaluation without platform effect re-estimation

Accuracy

Number of cell types in the partially annotated data

Method
gpsFISH
Naive simulation
Random
RankCorr
scGeneFit

Classifier:
Naive Bayes

C

Cell type

Non-neuronal

Excitatory

Inhibitory

Distance

Cell type

Count
30
20
10
0

Fig. 4

Fig. 5

Fig. 6

$$y_{ij} \sim \text{ZINB}(\mu_{ij}, \theta_{ij}, \pi)$$
$$\ln(\mu_{ij}) = \alpha + \ln(\lambda_{ij}) + \ln(CD_j)$$
$$\ln(\theta_{ij}) = \beta + \lambda_{ij}$$
$$\text{logit}(\lambda_{ij}) = \gamma_i \times \sqrt{x_{ij}} + c_i$$

$$\alpha \sim \mathcal{N}(0, \sigma_\alpha)$$
$$\beta \sim \mathcal{N}(0, \sigma_\beta)$$
$$\gamma_i \sim \mathcal{LN}(\mu_\gamma, \sigma_\gamma)$$
$$c_i \sim \mathcal{N}(\mu_c, \sigma_c)$$
$$\sigma_\alpha, \sigma_\beta, \sigma_\gamma, \sigma_c \sim \text{HalfCauchy}(0, 5)$$
$$\mu_\gamma, \mu_c \sim \text{Cauchy}(0, 5)$$
$$\pi \sim \mathcal{B}(1, 1)$$

y_{ij}: number of molecules in gene i and cell j from spatial data
x_{ij}: relative expression of gene i in cell j from scRNA-seq data
CD_j: total number of molecules in cell j from spatial data
λ_{ij}: corrected relative expression of gene i in cell j
γ_i and c_i: platform effect magnitude
$\mu_\gamma, \sigma_\gamma, \mu_c, \sigma_c$: platform effect hyperparameters
μ_{ij}: mean expression of gene i in cell j
θ_{ij}: dispersion of gene i in cell j
π: zero inflation parameter

Fig. S1

Fig. S2

Fig. S3

Fig. S4

Fig. S5

A Mean:
0.660 (hierarchical)
vs.
0.689 (flat)

B Mean:
0.700 (hierarchical)
vs.
0.716 (flat)

C Mean:
0.702 (hierarchical)
vs.
0.711 (flat)

Fig. S7

Figure 1 Platform effect between scRNA-seq and targeted spatial transcriptomics technologies.
A-C:
Scatter plot showing the log transformed relative expression of genes measured by both scRNA-seq and targeted spatial transcriptomics across three datasets, Moffit (**A**), Codeluppi (**B**), and Zhang (**C**), respectively. A small value is added to avoid negative infinity after log transformation. Each dot represents the relative expression of one gene in one cell type. Denominator for relative expression calculation is from all genes measured by both technologies. Color indicates density of dots. Dots should fall on the diagonal when there is no platform effect.
D:
Density plot of Deming regression coefficient for each dataset. Deming regression is fitted for each gene using relative expression measured by scRNA-seq and spatial transcriptomics data with intercept fixed to 0.
E:
Posterior predictive check of the Bayesian models fitted using each of the three datasets. QQ plot showing the distribution of simulated vs. observed spatial transcriptomics measurements.
F-G:
Density plot showing the estimated posterior distribution of σ_γ (**F**) and σ_c (**G**).

Figure 2: Schematic overview of gpsFISH.
Upper left, an scRNA-seq dataset with cell type annotation is used as input. Bottom, a genetic algorithm framework is used for gene panel selection. Platform effects are accounted for using a Bayesian model. Cell type hierarchy can also be incorporated. Upper right, output includes optimized gene panel with classification statistics.

Figure 3: Gene panel selection using gpsFISH.
A:
UMAP of cells based on the mouse hypothalamic scRNA-seq data from Moffit dataset at level 1 cell type annotation.
B:
Normalized confusion matrix of the optimized gene panel for Moffit dataset at level 1 cell type annotation.
C:
AUC for each cell type of the same gene panel.
D-E:
UMAP of cells based on simulated spatial transcriptomics measurements with platform effect of the optimized gene panel selected with (**D**) and without (**E**)considering platform effect at level 1 cell type annotation.

Figure 4: Comparison between gpsFISH and other gene selection methods.
A-B:
Box plot showing classification accuracy distribution of gene panels selected by 5 gene panel selection methods at different levels of partial annotation. The result is

based on the Moffit dataset using evaluation with (**A**) and without (**B**) platform effect re-estimation. Naïve Bayes is used as classifier.
C:
Normalized confusion matrix of the optimized gene panel for Moffit dataset at level 2 cell type annotation with dendrogram showing the cell type hierarchy. Diagonal values of the confusion matrix are removed for better visualization of misclassifications.

Figure 5: Redundancy in gene space across independent gene panel optimizations enables incorporation of customized preferences.
A:
Distribution of overlap of independent gene panels across 10 optimizations within each platform at level 1 cell type annotation.
B:
Accuracy of optimized gene panels without vs. with gene weight across 10 optimizations.
C:
Total number of probes of optimized gene panels without vs. with gene weight across 10 optimizations.

Figure 6: Gene panel selection with cell type hierarchy.
A:
Schematic of hierarchical gene selection using cell type hierarchy. A weighted penalty matrix is constructed using cell type hierarchy information quantified by pairwise distance between cell types. Additional penalty can be specified according to the cell type hierarchy. The weighted penalty matrix is then multiplied element-wise with the original confusion matrix to get the weighted confusion matrix for fitness evaluation.
B:
Original (left) vs. weighted (right) confusion matrix of the same optimized gene panel from Moffit dataset at level 2 cell type annotation with dendrogram showing the cell type hierarchy. Diagonal values of the confusion matrix are removed for better visualization of misclassifications.
C-E:
Percentage of across broad cell type (level 1) misclassifications over all misclassifications for flat vs. hierarchical classification on the Moffit (**C**), Codeluppi (**D**), and Zhang (**E**) dataset. Each dot represents one cell type with dots representing the same cell type connected. Wilcoxon paired test is performed between the percentages from flat vs. hierarchical classification and the p value is shown.

Figure S1: Schematic of the Bayesian model for platform effect estimation.
Circles in gray represent observed variables. Circles in green correspond to platform effect related variables to be estimated.

Figure S2: Bayesian model captures platform effect between scRNA-seq and targeted spatial transcriptomics technologies.

A-C:
Scatter plot showing the log transformed relative expression of genes measured by scRNA-seq vs. simulated spatial transcriptomics data using fitted Bayesian model for Moffit (**A**), Codeluppi (**B**), and Zhang (**C**), respectively.
D-E:
Density plot showing the estimated posterior distribution of μ_γ (**D**) and μ_c. (**E**).

Figure S3: Comparison between gpsFISH and other gene selection methods on the Zhang and Codeluppi dataset.
Box plot showing classification accuracy distribution of gene panels selected by 5 gene panel selection methods at different levels of partial annotation. (**A**) Zhang dataset using evaluation with platform effect re-estimation. (**B**) Zhang dataset using evaluation without platform effect re-estimation. (**C**) Codeluppi dataset using evaluation with platform effect re-estimation. (**D**) Codeluppi dataset using evaluation without platform effect re-estimation. Naïve Bayes is used as classifier.

Figure S4: Comparison between gpsFISH and other gene selection methods using random forest as classifier.
Box plot showing classification accuracy distribution of gene panels selected by 5 gene panel selection methods at different levels of partial annotation for the three datasets using evaluation with (**A-C**) and without (**D-E**) platform effect re-estimation. Random forest is used as classifier.

Figure S5: High redundancy across optimizations using gpsFISH.
A-C:
Bar plot showing among all the genes selected in 10 optimizations, the percentage of them that are included in 1 to 10 optimized panels for Moffit (**A**), Codeluppi (**B**), and Zhang (**C**) dataset at level 1 cell type annotation.
D:
Distribution of overlap of independent gene panels across 10 optimizations within each platform at level 2 cell type annotation.

Figure S6: Weighted gene panel selection based on probe count per gene.
A:
Distribution of probe count per gene for the Zhang dataset.
B-C:
Distribution of accuracy (**B**) and total number of probes (**C**) of optimized gene panels from optimization without and with gene weight. Optimization without gene weight is performed 10 times. Optimization with gene weight is performed 6 times, each time with a different probe count cutoff (no cutoff, 5, 10, 15, 20, 30).

Figure S7: Accuracy of optimized gene panels using flat vs. hierarchical gene selection.
A-C:

Distribution of accuracy of optimized gene panels using flat vs. hierarchical gene selection for Moffit (**A**), Codeluppi (**B**), and Zhang (**C**), respectively. Both flat and hierarchical gene selection are performed 5 times.

Information of the Moffit, Codeluppi, and Zhang dataset

Dataset	Moffit	Codeluppi	Zhang
Single-cell RNA sequencing platform	10X Genomics Chromium v2 & Illumina NextSeq500	Illumina HiSeq 2000	10X Genomics Chromium v3 & Illumina NovaSeq
Spatially resolved transcriptomics platform	MERFISH	osmFISH	DARTFISH
Number of cell types in scRNA-seq at level 1 for gene panel selection	12	12	16
Number of cells in scRNA-seq at level 1 for gene panel selection	30370	2816	64693
Number of cell types in scRNA-seq at level 2 for gene panel selection	87	47	46
Number of cells in scRNA-seq at level 2 for gene panel selection	30370	2816	64693
Number of overlapping cell types at level 1 for platform effect estimation	9	11	7
Number of cells in scRNA-seq at level 1 for platform effect estimation	29760	2139	43261
Number of cells in spatial transcriptomics data at level 1 for platform effect estimation	417026	3127	1341

Curated marker genes for the Zhang dataset

Subclass (Full Name)	Subclass Level 3	Subclass Level 1	Class	Substructure	Positive Markers (Curated)	Negative Markers	Degenerative State Upregulated	Degenerative State Downregulated	Adaptive State	Cycling State
Podocyte	POD	POD	epithelial cells	glomerulus	NPHS1, NPHS2, PTPRQ, CLIC5, NTNG1		CDKN1C, SPOCK2	PTPRQ		
Parietal Epithelial Cell	PEC	PEC	epithelial cells	glomerulus	CLDN1, VCAM1, CFH					
Proximal Tubule Cell		PT	epithelial cells	proximal tubules	LRP2, CUBN, AQP1		CST3, HAVCR1, CLU, APOE, S100A6, B2M		ITGB8, CDH6, DCDC2, VCAM1, DLGAP1, HAVCR1, PLSCR1	MKI67, TOP2A
Proximal Tubule Epithelial Cell Segment 1	PT-S1	PT	epithelial cells	proximal tubules	SLC5A12, SLC22A6,SLC22A8, SLC5A2					
Proximal Tubule Epithelial Cell Segment 2	PT-S2	PT	epithelial cells	proximal tubules	SLC34A1, SLC22A7					
Proximal Tubule Epithelial Cell Segment 3	PT-S3	PT	epithelial cells	proximal tubules	SLC5A11, MOGAT1, SLC22A7, SLC22A24, SLC7A13					
Thin Limb Cell	TL	TL	epithelial cells	intermediate tubules	CRYAB, TACSTD2, AKR1B1					
Descending Thin Limb Cell Type 2	DTL2	DTL	epithelial cells	intermediate tubules	AQP1, UNC5D	CLDN10				
Descending Thin Limb Cell Type 1	DTL1	DTL	epithelial cells	intermediate tubules	ADGRL3, ID1	CLDN10, AQP1				
Descending Thin Limb Cell Type 3	DTL3	DTL	epithelial cells	intermediate tubules	CLDN1, SH3GL3, SLC14A2, SMOC2	CLDN10, AQP2				
Ascending Thin Limb Cell	ATL	ATL	epithelial cells	intermediate tubules	CLDN1, SH3GL3, CLDN10, PROX1					
Thick Ascending Limb Cell		TAL	epithelial cells	Distal tubules	SLC12A1, CASR, UMOD, EGF			UMOD, EGF	ITGB6, PROM1, CCL2, PLSCR1, DCDC2	
Medullary Thick Ascending Limb Cell	M-TAL	TAL	epithelial cells	Distal tubules	PROX1					
Cortical Thick Ascending Limb Cell	C-TAL	TAL	epithelial cells	Distal tubules						
Macula Densa Cell	MD	TAL	epithelial cells	Distal tubules	NOS1, ROBO2	UMOD				
Distal Convoluted Tubule Cell		DCT	epithelial cells	Distal tubules	Distal tubules					
Distal Convoluted Tubule Cell Type 1	DCT1	DCT	epithelial cells	Distal tubules						
Distal Convoluted Tubule Cell Type 2	DCT2	DCT	epithelial cells	Distal tubules	SLC8A1					
Connecting Tubule		CNT	epithelial cells	Collecting tubules	SLC8A1, HSD11B2, CALB1					
Connecting Tubule Cell	CNT	CNT	epithelial cells	Collecting tubules						
Connecting Tubule Principal Cell	CNT-PC	CNT	epithelial cells	Collecting tubules	SCNN1G, SCNN1B					
Principal Cell		PC	epithelial cells	Collecting tubules	AQP2, AQP3					
Cortical Collecting Duct Principal Cell	CCD-PC	PC	epithelial cells	Collecting tubules	SCNN1G, SCNN1B					
Outer Medullary Collecting Duct Principal Cell	OMCD-PC	PC	epithelial cells	Collecting tubules	SCNN1G, SCNN1B					
Inner Medullary Collecting Duct Cell	IMCD	PC	epithelial cells	Collecting tubules	SLC14A2, HS3ST5					
Papillary Epithelial Cells	PapE	PC	epithelial cells	Collecting tubules	TP63, KRT5					
Intercalated Cell		IC	epithelial cells	Collecting tubules	ATP6V0D2					
Cortical Collecting Duct Intercalated Cell Type A	CCD-IC-A	IC	epithelial cells	Collecting tubules	SLC26A7, SLC4A1					
Connecting Tubule Intercalated Cell Type A	CNT-IC-A	IC	epithelial cells	Collecting tubules	SLC26A7, SLC4A1, SLC8A1					
Outer Medullary Collecting Duct Intercalated Cell Type A	OMCD-IC-A	IC	epithelial cells	Collecting tubules	SLC26A7, SLC4A1, KIT					
Intercalated Beta Cell	IC-B	IC	epithelial cells	Collecting tubules	SLC26A4, SLC4A9					
Endothelial Cell		EC	endothelial cells	vessels	PECAM1, CD34					
Glomerular Capillary Endothelial Cell	EC-GC	EC	endothelial cells	glomerulus	EMCN, HECW2, PLAT, EHD3					
Afferent / Efferent Arteriole Endothelial Cell	EC-AEA	EC	endothelial cells	vessels	BTNL9, PALMD, TM4SF1, SERPINE2, AQP1					
Descending Vasa Recta Endothelial Cell	EC-DVR	EC	endothelial cells	vessels	BTNL9, PALMD, TM4SF1, SERPINE2, AQP1, SLC14A1					
Peritubular Capilary Endothelial Cell	EC-PTC	EC	endothelial cells	vessels	DNASE1L3, PLVAP					
Ascending Vasa Recta Endothelial Cell	EC-AVR	EC	endothelial cells	vessels	DNASE1L3, PLVAP, TLL1	PALMD, BTNL9, SLC14A1				
Lymphatic Cell	EC-LYM	EC	endothelial cells	vessels	PROX1, MMRN1					
Vascular Smooth Muscle Cell and Pericyte		VSM/P	stroma cells	interstitium	PDGFRB, NOTCH3		TAGLN, ACTA2			
Mesangial Cell	MC	VSM/P	stroma cells	glomerulus	POSTN, PIEZO2, ITGA8					
Renin-positive Juxtaglomerular Granular Cell	REN	VSM/P	stroma cells	interstitium	REN					
Vascular Smooth Muscle Cell	VSMC	VSM/P	stroma cells	interstitium	MYH11, MCAM					
Vascular Smooth Muscle Cell / Pericyte	VSMC/P	VSM/P	stroma cells	interstitium						
Fibroblast		FIB	stroma cells	interstitium	C7, DCN, COL1A1, PDGFRA				FLRT2, FGF14, IGF1	
Fibroblast	FIB	FIB	stroma cells	stroma cells	MEG3, LAMA2					
Medullary Fibroblast	M-FIB	FIB	stroma cells	interstitium	SYT1, TNC					
Myofibroblast	MyoF	FIB	stroma cells	interstitium	FAP, ACTA2, TAGLN, POSTN, GLI2, COL5A1					
Immune Cells		IMM	immune cells	interstitium	PTPRC					
B Cell	B	IMM	immune cells	interstitium	MS4A1, BANK1					
Plasma Cell	PL	IMM	immune cells	interstitium	IGKC, MZB1					
T Cell	T	IMM	immune cells	interstitium	CD3E, CD4					
Natural Killer T Cell	NKT	IMM	immune cells	intrrstitium	NKG7, GNLY, CD96, RUNX3					
Mast Cell	MAST	IMM	immune cells	interstitium	MS4A2, CPA3, KIT	IL3RA				
M2 Macrophage	MAC-M2	IMM	immune cells	interstitium	CD163, F13A1, MRC1, CD14					
Classical Dendritic Cell	cDC	IMM	immune cells	interstitium	ITGAX, FLT3	CD14				
Plasmacytoid Dendritic Cell	pDC	IMM	immune cells	interstitium	IL3RA, FLT3	CD14				
Non-Classical Monocyte	ncMON	IMM	immune cells	interstitium	FCN1, HLA-DRA, FCGR3A					
Neutrophil	NC	IMM	immune cells	interstitium	S100A8, S100A9, IFITM2, FCGR3B					
Schwann Cell / Neural	SC/NEU	NEU	neural like cells	interstitium	CDH19, NRXN1, PLP1, S100B					

Table S2