1   Neanderthal Introgression Shaped Human Circadian Traits
2

3   Keila Velazquez-Arcelay[1], Laura L. Colbran[2], Evonne McArthur[3], Colin Brand[4,5], Justin
4   Siemann[1], Douglas McMahon[1], and John A. Capra[4,5*]
5
6   [1] Department of Biological Sciences, Vanderbilt University
7   [2] Department of Genetics, Perelman School of Medicine, University of Pennsylvania
8   [3] Vanderbilt University School of Medicine
9   [4] Department of Epidemiology and Biostatistics, University of California, San Francisco
10 [5] Bakar Computational Health Sciences Institute, University of California, San Francisco
11
[*] Corresponding author: tony@capralab.org

ABSTRACT
Introduction: When the ancestors of modern Eurasians migrated out of Africa and interbred with
Eurasian archaic hominins, namely Neanderthals and Denisovans, DNA of archaic ancestry
integrated into the genomes of anatomically modern humans. This process potentially
accelerated adaptation to Eurasian environmental factors, including reduced ultra-violet radiation
and an increased variation in seasonal dynamics. However, whether these groups differed
substantially in circadian biology, and whether archaic introgression adaptively contributed to
human chronotypes remains unknown.
Results: Here we traced the evolution of chronotype based on genomes from archaic hominin
and present-day humans. First, we inferred differences in circadian gene sequences, splicing, and
regulation between archaic hominins and modern humans. We identified 28 circadian genes
containing variants likely to alter splicing in archaics (e.g., *CLOCK*, *PER2*, *RORB*, *RORC*), and
16 circadian genes likely divergently regulated between present-day humans and archaic
hominins, including *RORA*. These differences suggest the potential for introgression to modify
circadian gene expression. Testing this hypothesis, we found that introgressed variants are
enriched among eQTLs for circadian genes. Supporting the functional relevance of these
regulatory effects, we found that many introgressed alleles have strong associations with
chronotype. Strikingly, the strongest introgressed effects on chronotype increase morningness,
which is consistent with adaptations to high latitude in other species. Finally, we identified 26
circadian loci with evidence of adaptive introgression, including *PER2* and *MYBBP1A*.
Conclusions: These findings identify differences in circadian gene regulation between modern
humans and archaic hominins and support the contribution of introgression via coordinated
effects on variation in human chronotype.
37
38
39 **Keywords:** circadian biology, chronotype, Neanderthals, adaptive introgression

40
41

SIGNIFICANCE STATEMENT

Interbreeding between modern humans and Neanderthals created the potential for adaptive introgression as humans moved into new environments that had been populated by Neanderthals for hundreds of thousands of years. Here we discover substantial lineage-specific genetic differences in circadian genes and their regulatory elements between humans and Neanderthals. We then show that introgressed archaic alleles are enriched for effects on circadian gene regulation and consistently increase propensity for morningness in modern Europeans. These results substantially expand our understanding of how the genomes of humans and our closest relatives responded to living in environments with different light/dark cycles, and they demonstrate a coordinated contribution of archaic admixture to modern human chronotype in a direction that is consistent with adaptation to higher latitudes.

INTRODUCTION

All anatomically modern humans (AMH) trace their origin to the African continent around 300 ka (Hublin et al. 2017; Stringer 2016), where environmental factors shaped many of their biological features. Approximately seventy-thousand years ago (Bae, Douka, and Petraglia 2017), the ancestors of modern Eurasian AMH began to migrate out of Africa, where they were exposed to diverse new environments. In Eurasia, the novel environmental factors included greater seasonal variation in temperature and photoperiod.

Changes in the pattern and level of light exposure has biological and behavioral consequences in organisms. For example, *D. melanogaster* that are native to Europe harbor a polymorphism in *timeless*, a key gene in the light response of the circadian system, that follows a latitudinal cline in allele frequency (Sandrelli et al. 2007; Tauber et al. 2007). This polymorphism is an insertion in the 5' region of the gene, resulting in two initiation codons and as a result two isoforms of the protein: L-TIM with prevalence in southern Europe and S-TIM with prevalence in northern Europe. The isoforms have different levels of affinity to cryptochrome (CRY), creating a change in photosensitivity and altering the length of the period. Another example is found in pacific salmon. The Chinook salmon (*Oncorhynchus tshawytscha*) populations show a latitudinal-cline in the frequency and length of repeat motifs in the gene *OtsClock1b* (O'Malley, Ford, and Hard 2010; O'Malley and Banks 2008). Thus, this locus is under selection associated with latitude and photoperiod. The evolution of circadian adaptation to diverse environments has been widely studied in insects, plants (Q. Zhang et al. 2008; Michael et al. 2003), and fishes, but it is understudied in humans. Adaptive processes could have helped to align human biology and chronotype to new natural conditions.

Previous studies in humans found a correlation between latitude and chronotype (morningness vs. eveningness) variation (Leocadio-Miguel et al. 2017; Lowden et al. 2018; Randler and Rahafar 2017) and a latitudinal-cline in circadian allele frequencies (Putilov et al. 2019; Putilov, Dorokhov, and Poluektov 2018; Dorokhov et al. 2018), highlighting the contribution of the environment to circadian behavior and circadian biology. Many human health effects are linked to the misalignment of chronotype (Knutson and von Schantz 2018), including cancer, obesity (Gan et al. 2018; Gyarmati et al. 2016; Papantoniou et al. 2016; 2017; Shi et al. 2020; Yousef et al. 2020), and diabetes (Gan et al. 2015; Larcher et al. 2015; 2016). There is also evidence of a correlation between evening chronotype and mood disorders, most notably seasonal affective disorder (SAD), depression, and worsening of bipolar disorder episodes (Srinivasan et al. 2006; Taylor and Hasler 2018; Kivelä, Papadopoulos, and Antypa 2018). Thus, we hypothesize that the differences in geography and environment encountered by early AMH populations moving into higher latitudes created potential for circadian misalignment and health risk.

Although AMHs arrived in Eurasia ~70 ka, other hominins (e.g. Neanderthals and Denisovans) lived there for more than 400 ka (Meyer et al. 2016; 2014; Arnold et al. 2014). These archaic hominins diverged from AMHs around 700 ka (Gómez-Robles 2019; Nielsen et al. 2017; Meyer et al. 2012; Prüfer et al. 2014; 2017; Mafessoni et al. 2020), and as a result, the ancestors of AMHs and archaic hominins evolved under different environmental conditions. While there was substantial variation in the latitudinal ranges of each group, the Eurasian hominins largely lived at consistently higher latitudes and, thus, were exposed to higher amplitude seasonal variation in photoperiods. Given the influence of environmental cues on

circadian biology, we hypothesized that these separate evolutionary histories produced
differences in circadian traits adapted to the distinct environments.
When AMH migrated into Eurasia, they interbred with the archaic hominins that were
native to the continent, initially with Neanderthals (Green et al. 2010; Villanea and Schraiber
2019) around 60 ka (Sankararaman et al. 2012; Skoglund and Mathieson 2018) and later with
Denisovans (Jacobs et al. 2019). Due to this, a substantial fraction (>40%) of the archaic
variation remains in present-day Eurasians (Skov et al. 2020; Vernot and Akey 2014), although
each human individual carries only ~2% DNA of archaic ancestry (Vernot et al. 2016; Prüfer et
al. 2017). Most of the archaic ancestry in AMH was subject to strong negative selection, but
some of these introgressed alleles remaining in AMH populations show evidence of adaptation
(Gower et al. 2021; Racimo et al. 2015). Archaic alleles have been associated with differences in
hemoglobin levels at higher altitude in Tibetans, immune resistance to new pathogens, levels of
skin pigmentation, and fat composition (Dannemann and Kelso 2017; Racimo et al. 2015; 2017;
Racimo, Marnetto, and Huerta-Sánchez 2017; Huerta-Sánchez et al. 2014; McArthur, Rinker,
and Capra 2021). Introgressed alleles are also associated with chronotype (Dannemann and
Kelso 2017), suggesting a potential adaptive pressure stemming from migration to higher
latitudes. Summarizing effects across introgressed alleles, chronotype was also moderately
enriched for heritability in introgressed regions (McArthur, Rinker, and Capra 2021).
Motivated by the potential for a role of archaic introgression in AMH circadian variation,
we explore two related questions: 1) Can comparative genomic analysis identify differences in
AMH and archaic hominin circadian biology?, and 2) Do introgressed archaic alleles influence
human circadian biology? Understanding the ancient history and evolution of chronotypes in
humans will shed light on human adaptation to high latitudes and provide context for the genetic
basis for the modern misalignment caused by the development of technology and night
shiftwork.
124
125
126
127 **RESULTS**
128
129 **Did archaic hominins and modern humans diverge in circadian biology?**
130 Following divergence ~700,000 years ago (Gómez-Robles 2019; Nielsen et al. 2017), archaic
131 hominins and AMH were geographically isolated, resulting in the accumulation of lineage-
132 specific genetic variation (Figure 1). In the next several sections, we evaluate the evidence for
133 divergence in circadian biology between archaic hominin and modern human genomes.
134
Identifying archaic-hominin-specific circadian gene variation
With the sequencing of several genomes of archaic hominins, we now have a growing, but
incomplete, catalog of genetic differences specific to modern and archaic lineages. We evaluated
archaic-specific variants (Kuhlwilm and Boeckx 2019) for their ability to influence proteins,
splicing, and regulation of 246 circadian genes (Methods). The circadian genes were identified
by a combination of literature search, expert knowledge, and existing annotations (Table S1;
Figure S1; Methods).
We identified 1,136 archaic-specific variants in circadian genes, promoters, and
candidate distal cis-regulatory elements (cCREs). The circadian genes with the most archaic-

specific variants are *CLDN4*, *NAMPT*, *LRPPRC*, *ATF4*, and *AHCY* (125, 112, 110, 104, 102
respectively) (Table S2).

146
147

148
Figure 1. Did the sharing of functionally diverged alleles from archaic hominins influence
human circadian biology? A) Anatomically modern humans and archaic hominins evolved
separately at different latitudes for hundreds of thousands of years. The ancestors of modern
Eurasian humans left Africa approximately 70 thousand years ago and admixed with archaics,
likely in southwestern Asia. The shaded purple range represents the approximate Neanderthal
range. The purple dot represents the location of the sequenced Denisovan individual in the Altai
Mountains; the full range of Denisovans is currently unknown. Silhouettes from phylopic.org. **B)**
After the split between the human and archaic lineages, each group accumulated variation and
evolved in their respective environments for approximately 700 thousand years. We first test for
evidence for divergent circadian evolution during this time. Humans acquired introgressed alleles
from Neanderthals and from Denisovans around 60 and 45 thousand years ago, respectively.
These alleles experienced strong selective pressures; however, ~40% of the genome retains
archaic ancestry in some modern populations. The second question we explore is whether
introgression made adaptive contributions to human circadian biology. **C)** The core circadian
clock machinery is composed of a dimer between the CLOCK and ARNTL (BMAL1)
transcription factors, which bind to E-box enhancer elements and activate the expression of the

Period (*PER1/2/3*) and Cryptochrome (*CRY1/2*) genes. PERs and CRYs form heterodimers that
inhibit the positive drive of CLOCK-BMAL1 on E-boxes, inhibiting their own transcription in a
negative feedback loop. CLOCK-BMAL1 also drive the expression of *NR1D1/2* (Nuclear
Receptor Subfamily 1 Group D Member 1 and 2), *RORA/B* (RAR Related Orphan Receptor A
and B), *DBP* (D-Box Binding PAR BZIP Transcription Factor), and other clock-controlled genes
(CCG). ROR and REV-ERB are transcriptional regulators of BMAL1. CK1 binds to the
PER/CRY heterodimer, phosphorylating PER and regulating its degradation. Similarly, FBXL3
marks CRY for degradation.
173
174
Fixed human- and archaic-specific variants are enriched in circadian genes and associated
regulatory elements
After the archaic and AMH lineages diverged, each group accumulated genetic variation specific
to each line. Variants fixed in each lineage are likely to be enriched in genomic regions that
influence traits that experienced positive selection. We tested whether human- and archaic-
specific fixed variants are enriched compared to other sites with archaic variants in circadian
genes, their promoters, and in annotated regulatory elements within 1 Mb (Figure 2). We found
that human- and archaic-specific fixed variants are enriched in circadian genes (Fisher's exact
test; human: OR=1.84, P=7.06e-12; archaic: OR=1.13, P=0.023) and distal regulatory elements
(Fisher's exact test; human: OR=1.25, P=8.39e-4; archaic: OR=1.16, P=6.15e-5). Although
promoter regions have a similar enrichment pattern as that in gene and regulatory regions, the p-
values are high (Fisher's exact test; human: OR=1.21, P=0.65; archaic: OR=1.09, P=0.63). This
is likely due to the small number of such variants in promoters. These results suggest that both
groups had a greater divergence in genomic regions related to circadian biology than expected.
189

190
Figure 2. **Human- and archaic-specific fixed variants are enriched in circadian regulatory,**
promoter, and gene regions. Human-specific fixed variants are significantly enriched in
circadian regulatory elements (Fisher's exact: OR=1.25, P=8.39e-4) and gene regions (Fisher's
exact: OR=1.84, P=7.06e-12) compared to variants that are not fixed. Promoters show a similar
enrichment, but the higher p-value is the result of the small number of variants (Fisher's exact

test: OR=1.21, P=0.65). Likewise, archaic-specific variants are enriched in circadian regulatory
regions (Fisher's exact: OR=1.16, P=6.15e-5) and gene regions (Fisher's exact: OR=1.13,
P=0.023), with the promoters showing a similar trend (Fisher's exact test: OR=1.09, P=0.63).
The number in parentheses gives the number of variants observed in each type.
200
201
Several core circadian genes have evidence of alternative splicing between humans and
archaic hominins
We find only two archaic-specific coding variants in circadian genes: one missense and one
synonymous. The missense variant (hg19: chr17_46923411_A_G) is in the gene *CALCOCO2*,
calcium-binding and coiled-coil domain-containing protein 2. SIFT, PolyPhen, and CADD all
predict that the variant does not have damaging effects. The second variant (hg19:
chr7_119914770_G_T) is in the gene *KCND2*, which encodes a component of a voltage-gated
potassium channels, but it is synonymous and the variant effect predictors suggest it is tolerated.
To explore potential splicing differences in circadian genes between humans and
archaics, we applied SpliceAI to predict whether any sequence differences between modern
humans and archaics are likely to modify splicing patterns. Four archaic individuals were
included in this analysis (the Altai, the Vindija, the Chagyrskaya Neanderthals, and the Altai
Denisovan). We found that 28 genes contained at least one archaic-specific variant predicted to
result in alternative splicing in archaics. These included several of the core clock genes *CLOCK*,
PER2, *RORB*, *RORC*, and *FBXL13* (Figure 3A,C; Table S3). For example, the variant
chr2:239187088-239187089 in the 1st intron of *PER2* is predicted to result in a longer 5' UTR.
The splice-altering variants were largely specific to the two different archaic linages (Figure 3A),
with 13 specific to the Denisovan, eight shared among the three Neanderthals, and only one
shared among all four archaic individuals.
221
Circadian gene regulatory divergence between humans and archaic hominins
Given the enrichment of variants in regulatory regions of circadian genes, we sought to explore
the potential for differences in circadian gene regulation between humans and archaics with
causes beyond single lineage-specific variants. We leveraged an approach we recently developed
for predicting gene regulatory differences between modern and archaic individuals from
combinations of genetic variants (Colbran et al. 2019). The approach uses PrediXcan, an elastic
net regression method, to impute gene transcript levels in specific tissues from genetic variation.
Previous work demonstrated that this approach has a modest decrease in performance when
applied to Neanderthals, but that it can accurately applied between humans and Neanderthals for
thousands of genes. Here, we quantify differences in predicted regulation of the 246 circadian
genes between 2,504 humans in the 1000 Genomes Project (1KGP) and three archaic hominins
(the Altai Neanderthal, the Vindija Neanderthal, and the Altai Denisovan). The predicted
regulation values are normalized to the distribution in the training set from the Genotype Tissue
Expression Atlas (GTEx).
236
237

238
Figure 3. Many circadian genes have evidence of alternative splicing and divergent
regulation between modern and archaic hominins. A) The sharing across archaic individuals
of the 28 predicted archaic-specific splice-altering variants (SAV) in circadian genes. Most were
specific to either the Denisovan or Neanderthal lineage (Table S3). **B)** The sharing of predicted
divergently regulated (DR) gene/tissue pairs across three archaic individuals. (Predictions were
not available for the Chagyrskaya Neanderthal.) Seventeen divergently regulated gene/tissue
pairs were present in all three archaics (representing 16 unique genes). Additionally, 7
gene/tissue DR pairs are shared between the Altai Neanderthal and the Denisovan individual.
One pair is shared between the Vindija Neanderthal and the Denisovan (Table S4). **C)** The
proportion of circadian genes containing archaic splice-altering variants predicted by SpliceAI
(SAV; 11.4%) or divergently regulated circadian genes predicted by PrediXcan (DR; 6.5%).
Thus, 17.9% of the circadian genes are predicted to contain differences to AMH via these
mechanisms.

252
253
We first analyzed gene regulation predictions in the core circadian clock genes. Archaic gene
regulation was at the extremes of the human distribution for many core clock genes including
PER2, *CRY1*, *NPAS2*, *RORA, NR1D1* (Figure 4; Figure S2). For example, the regulation of
PER2 in the two Neanderthals is lower than 2,491 of the 2,504 (99.48%) modern humans
considered. The Denisovan has a predicted *PER2* regulation that is lower than 2,410 (96.25%).
Expanding to all circadian genes and requiring archaic regulation to be more extreme than all
humans (Methods), we identified 24 circadian genes across 23 tissues with strong divergent
regulation between humans and at least one archaic hominin (Figure 3B; Table S4). For example,
all three archaic individuals' regulation values for *RORA*, a core clock gene, are lower than for
any of the 2,504 modern humans. We found that 16 of these genes (Figure S3; Table S4),
including *RORA*, *MYBBP1A*, and *TIMELESS*, were divergently regulated in all archaic
individuals. This represents 6.5% of all the circadian genes (Figure 3C). Surprisingly, the two
Neanderthals only shared one DR gene not found in the Denisovan, while the Altai Neanderthal
and Denisovan shared seven not found in Vindija (Figure 3B). The Altai and Vindija
Neanderthals represent deeply diverging lineages, and this result suggests that they may have
experienced different patterns of divergence in the regulation of their circadian genes.
Given these differences in circadian gene regulation between humans and archaics, we
tested whether circadian genes are more likely to be divergently regulated than other gene sets.
Each archaic individual shows nominal enrichment for divergent regulation of circadian genes,

but given the small sample size, the P-values are moderate (Permutation test; Altai: OR=1.21,
P=0.19, Vindija: OR=1.05, P=0.43, Denisovan: OR=1.20, P=0.24). The enrichment was stronger
(~1.2x) in the Altai Neanderthal and Denisovan individual.
276
277

278
Figure 4. Many circadian genes are divergently regulated between modern humans and
archaic hominins. Comparison of the imputed regulation of core circadian genes between 2504
humans in 1000 Genomes Phase 3 (gray bars) and three archaic individuals (vertical lines). For
each core circadian gene, the tissue with the lowest average P-value for archaic difference from
humans is plotted. Archaic gene regulation is at the extremes of the human distribution for
several core genes: *CRY1*, *PER2*, *NPAS2, NR1D1 RORA*. See Figure S2 for all core clock genes
and Figure S3 for all divergently regulated circadian genes.
286
287
288　**Did introgressed archaic variants influence modern human circadian biology?**
289　The previous sections demonstrate lineage-specific variation in many genes essential to the
290　function of the core circadian clock and related pathways. Given this evidence of functional
291　differences between archaic hominins and AMH in these systems, we next evaluated the
292　influence of archaic introgression on AMH circadian biology.
293
Introgressed variants are enriched in circadian gene eQTL
Given the differences between archaic and modern sequences of circadian genes and their
regulatory elements, we investigated whether Neanderthal introgression contributed functional
circadian variants to modern Eurasian populations. We considered a set of 863,539 variants with

evidence being introgressed from archaic hominins to AMH (Browning et al. 2018). These
variants were identified using the S* algorithm, which searches for regions containing a high
density of alleles in common with Neanderthals and not present or at very low frequency in
Africans.
We identified 3,857 introgressed variants associated with the regulation of circadian
genes in modern non-Africans; i.e., they are eQTL for circadian genes in at least one of the
tissues in GTEx (Table S5). The genes *PTPRJ*, *HTR1B*, *NR1D2*, *CLOCK*, and *ATOH7* had the
most eQTL (304, 273, 262, 256, and 252 respectively). The tissues artery tibial, cells cultured
fibroblasts, skin sun exposed lower leg, adipose subcutaneous, and thyroid had the most eQTL
(1361, 962, 876, 871, 716, respectively); though, these patterns are driven in large part by
different power between tissues. Notably, several of these circadian genes (e.g., *NR1D2* and
CLOCK) with introgressed eQTL were also found to be divergently regulated in our previous
analyses. This indicates that some of the archaic-derived variants that drove divergent regulation
were retained after introgression and continue to influence circadian regulation in modern
humans.
Introgressed variants are significantly more likely to be eQTL for circadian genes than
expected by chance (Figure 5A; Fisher's exact test: OR=1.45, P=9.71e-101). Examining these
associations in each tissue, we found that introgressed eQTL showed significant enrichment for
circadian genes in most tissues (34 of 49; Figure 5B; Table S6) and trended this way in all but
five. Again, the differences between tissues likely influenced by differential power. These results
suggest that the circadian pressures were widespread across tissues. Given the previously
observed depletion for introgressed variants in regulatory elements and eQTL (Petr et al. 2019;
Rinker et al. 2020; Telis, Aguilar, and Harris 2020), this enrichment for circadian genes among
introgressed eQTL is surprising and suggests that the archaic circadian alleles could have been
beneficial after introgression.
323
324

325
Figure 5. Circadian genes are enriched for introgressed eQTL. A) Archaic introgressed
variants are more likely to be eQTL for circadian genes in GTEx than for non-circadian genes
(Fisher's exact test: OR=1.45, P=9.71e-101). Purple represents the set of introgressed variants;
blue represents the set of circadian variants; 3,857 are introgressed eQTL in circadian genes. The
overlaps are not to scale. **B)** The enrichment for circadian genes among the targets of
introgressed eQTLs in each GTEx tissue. Introgressed eQTL in most tissues show significant
enrichment for circadian genes (Fisher's exact test; Table S6). Kidney cortex did not have any
circadian introgressed eQTLs and thus is not shown. Numbers inside the parenthesis indicate the
count of variants in each tissue. Gray bars indicate lack of statistical significance; light blue bars
indicated nominal significance (p <= 0.05); and dark blue bars indicate significance at the 0.05
level after Bonferroni multiple testing correction (p <= 0.00102).

337
338
Introgressed variants predominantly increase propensity for morningness
After observing that circadian gene expression is influenced by archaic variants, we evaluated
whether these effects are likely to result in a change in organism-level phenotype. To do this, we
evaluated evidence that introgressed variants influence chronotype. The heritability of
chronotype has been estimated in a range from 12 to 38% (Lane et al. 2016; Jones et al. 2019;
2016). Previous studies have identified individual introgressed loci associated with chronotype
(Dannemann and Kelso 2017). We recently found modest enrichment for heritability of
chronotype (morning/evening person phenotype in a GWAS of the UK Biobank) among
introgressed variants genome-wide using stratified LD score regression (heritability enrichment:
1.58, P=0.25) and a suggestive bias toward increasing morningness (McArthur, Rinker, and
Capra 2021).

To test comprehensively for a directional effect, we calculated the cumulative fraction of
introgressed loci associated with chronotype in the UK Biobank that increase morningness (after
clumping based on LD at R^2=0.9 in CEU). Each of the nine introgressed loci most strongly
associated with chronotype increase propensity for morningness (Figure 6; Table S7; Table S8; 9
loci, P<=1.35e-13). As the strength of the association with morningness decreases, the bias
begins to decrease, but the effect is maintained well past the genome-wide significance threshold
(P<5e-8). When focusing the analysis on introgressed variants in proximity (<1 Mb) to circadian
genes, the pattern becomes even stronger. The bias toward morningness remains above 80% at
the genome-wide significance threshold. This suggests that introgressed variants act in a
consistent direction on chronotype, especially when they influence circadian genes.
360
361

362
Figure 6. Introgressed variants associate with increased morningness. The cumulative
fraction of introgressed loci significantly associated with the morning vs. evening person trait in
the UK Biobank that increase morningness (y-axis) at a given p-value threshold (x-axis).
Introgressed loci associated with chronotype are strongly biased towards increasing
morningness, and this effect is strongest at the most strongly associated loci. Introgressed
variants nearby (<1 Mb) circadian genes (blue) are even more strongly biased towards increasing
morningness than introgressed variants overall (gray). Each dot (triangle) represents an
associated locus; variants were clumped by LD for each set (R^2=0.9 CEU).
371
372
Circadian rhythms are involved in a wide variety of biological systems. To explore other
phenotypes potentially influenced by the introgressed circadian variants, we evaluated evidence
for pleiotropic associations. First, we retrieved all the genome-wide associations reported for
introgressed variants in the Open Targets Genetics (https://genetics.opentargets.org) database,
which combines GWAS data from the GWAS Catalog, UK Biobank, and several other sources.
Introgressed circadian variants are associated with traits from a diverse range of categories

(Table S9). Associations with blood related traits are by far the most common; however, this is
likely because they have more power in the UK Biobank. Overall, circadian introgressed variants
are significantly more likely to have at least one trait association than introgressed variants not in
the circadian set (Fisher's exact test: OR=1.25, P=7.03e-25) (Figure S4A). The circadian variants
also associate with significantly more traits per variant than the non-circadian set (Mann-
Whitney U: P=9.93e-14) (Figure S4B; Table S10). These results suggest effects for introgressed
circadian variants beyond chronotype.
386
Evidence for adaptive introgression at circadian loci
The gene flow from Eurasian archaic hominins into AMH played a role in adaptations to some of
the new environmental conditions encountered outside of Africa (Racimo et al. 2015). The above
analyses demonstrate the effects of introgressed variants on circadian gene regulation and
chronotype. To explore whether these circadian regions show evidence of adaptive introgression,
we considered two sets of introgressed regions predicted to have contributed to AMH adaptation
by two recent machine learning algorithms: *genomatnn* (Gower et al. 2021) and *MaLAdapt* (X.
Zhang et al. 2023). We intersected the circadian introgressed variants with the adaptive
introgression regions from each method.
We identified 26 regions in or near circadian genes with evidence of adaptive
introgression—seven from genomatnn and 19 from MaLAdapt (including 53 and 116 SNPs,
respectively; Table S11). One of these adapive introgression regions overlaps the upstream
region of the gene *PER2*, a core clock gene involved in the generation of circadian rhythms
through feedback loops. This gene is also predicted to have lower gene regulation in archaics
hominins than most humans. We also found several introgressed eQTL for *PER2*, though they do
not overlap the predicted adaptive introgression region. At the phenotype level, *PER2*
introgressed variants have been associated with having an effect on morning/evening preference
in humans in the UK Biobank. The Vindija Neanderthal also carries a lineage-specific variant in
this gene that has splice-altering effects; this suggests that *PER2* may have experienced multiple
functional changes in different modern and archaic lineages, with potential adaptive effects.
Another region with evidence of adaptive introgression overlaps the circadian variants in
the gene *MYBBP1A*. In mouse, MYBBP1A acts as co-repressor of PER2 by binding to its
promoter in combination with Cryptochrome-1 (CRY1). *MYBBP1A* is predicted to be
divergently regulated between AMH and archaic hominins, with decreased expression in the
archaics, and it also contains introgressed variants that are circadian eQTLs.
412
413
414
415 **DISCUSSION**
416
The Eurasian environments where Neanderthals and Denisovans lived for several hundred
thousand years are located at higher latitudes with more variable photoperiods than the landscape
where AMH evolved before leaving Africa. Evaluating genetic variation that arose separately in
each of the archaic and AMH lineages after their split ~700 MYA, we identified lineage-specific
genetic variation in circadian genes, their promoters, and flanking distal regulatory elements. We
found that both archaic- and human-specific variants are observed more often than expected in
each class of functional region. This result suggests that, while each group evolved separately
during hundreds of thousands of years in divergent environments, both experienced pressure on

circadian related variation. Leveraging sequence-based machine learning methods, we identified
many archaic-specific variants likely to influence circadian gene splicing and regulation. For
example, core clock genes (*CLOCK*, *PER2*, *RORB*, *RORC*, and *FBXL13*) have archaic variants
predicted to cause alternative splicing compared to AMH. Several core genes were also predicted
in archaics to be at the extremes of human gene regulation, including *PER2*, *CRY1*, *NPAS2*,
RORA, NR1D1. Surprisingly, the Altai Neanderthal shared more divergent regulation in the
circadian genes with the Denisovan individual than the Vindija Neanderthals. The two
Neanderthals represent populations that were quite distantly diverged with substantially different
histories and geographical ranges. The Denisovan and Altai Neanderthal also come from the
same region in Siberia, while the Vindija Neanderthal came from a region in Croatia with
slightly lower latitude.
Introgression introduced variation that first appeared in the archaic hominin lineage into
Eurasian AMH. While most of this genetic variation experienced strong negative selection in
AMH, a smaller portion is thought to have provided adaptive benefits in the new environments
(Racimo et al. 2015). Given the divergence in many circadian genes' regulation, we explored the
landscape of introgression on circadian genes. We first looked at introgressed circadian variants
that are likely to influence gene regulation in AMH. Variants in this set are observed more often
than expected, suggesting the importance of maintaining circadian variation in the population.
We then evaluated the association of these introgressed variants with variation in
circadian phenotypes of Eurasians. We previously reported a modest enrichment among
introgressed variants for heritability of the morning/evening person phenotype. Here we further
discovered a consistent directional effect of the introgressed circadian variants on chronotype.
The strongest associated variants increase the probability of being a morning person in
Eurasians.
While it is not immediately clear why increased morningness would be beneficial at
higher latitudes, considering this directional effect in the context of clock gene regulation and the
challenge of adaptation to higher latitudes suggests an answer. In present day humans, behavioral
morningness is correlated with shortened period of the circadian molecular clockworks in
individuals. This earlier alignment of sleep/wake with external timing cues is a consequence of a
quickened pace of the circadian gene network (Brown et al. 2008). Therefore, the morningness
directionality of introgressed circadian variants may indicate selection toward shortened
circadian period in the archaic populations living at high latitudes. Supporting this interpretation,
shortened circadian periods are required for synchronization to the extended summer
photoperiods of high latitudes in *Drosophila*, and selection for shorter periods has resulted in
latitudinal clines of decreasing period with increasing latitude, as well as earlier alignment of
behavioral rhythms (Hut et al. 2013). In addition, *Drosophila* populations exhibit decreased
amplitude of behavioral rhythms at higher latitudes which is also thought to aid in
synchronization to long photoperiods (Hut et al. 2013).
Our finding that introgressed circadian variants generally decrease gene regulation of
circadian genes suggests that they could lead to lower amplitude clock gene oscillations.
However, when assayed in present day humans there is not a strong correlation between the
overall expression level of *NR1D1* and the transcriptional amplitudes of other clock genes within
individuals (Brown et al. 2008), and quantitative modeling of the mammalian circadian
clockworks suggests that stable clock gene rhythms can result across a wide range of absolute
levels of gene expression as long as the stoichiometric ratios of key positive and negative clock
genes are reasonably conserved (Kim and Forger 2012). Interestingly, lower transcriptional

amplitude of *NR1D1* does confer greater sensitivity of the present-day human clockworks to
resetting stimuli, a potentially adaptive characteristic for high latitudes (Brown et al. 2008).
473   Thus, given the studies of latitudinal clines and adaptation from *Drosophila* and the
474 nascent understanding of clock gene contributions to behavioral phenotypes in present day
475 humans, the directional effects of introgressed circadian gene variants toward early chronotype
476 and decreased gene regulation we observed can be viewed as potentially adaptive. More
477 complex chronotype phenotyping and mechanistic studies of the variants of interest are needed
478 to fully understand these observations.
479   Finally, to explore evidence for positive selection on introgressed variants in AMH, we
480 analyzed results from two recent methods for detecting adaptive introgression: *genomatnn* and
481 *MaLAdapt*. Both methods identified circadian loci as candidates for adaptive introgression.
482 However, we note that the predictions of these two methods have only modest overlap with one
483 another, underscoring the difficulty of identifying adaptive introgression.
484   Several limitations must be considered when interpreting our results. First, multiple sets
of predictions of introgressed variants are available. While these sets are qualitatively similar in
their genomic attributes, additional introgressed circadian variants could be discovered by
including other lines of evidence for introgression. Second, it is challenging to quantify the
complexity of traits with a large behavioral component (like chronotype) and infer their variation
from genomic information alone. Nevertheless, we believe our approach of focusing on
molecular aspects (splicing, gene regulation) of genomic loci with relevance to circadian
biology, in parallel to GWAS-based associations, lends additional support to the divergence in
chronotype between archaic hominins and modern humans. Third, we also note that circadian
rhythms contribute to many biological systems, so the variants in these genes tend to be
associated with a variety of phenotypes. Thus, there is also the potential that selection acted on
other phenotypes influenced by circadian variation than those related directly to chronotype.
Fourth, recent adaptive evolution is challenging to identify, and this is especially challenging for
introgressed loci. Nonetheless, we find several circadian loci with evidence of adaptive
introgression from each of the scans we considered. Finally, given the many environmental
factors that differed between African and non-African environments, it is difficult to definitively
determine whether selection on a particular locus was the result of variation in light levels vs.
other related factors, such as temperature. Nonetheless, given the observed modern associations
with chronotype for many of these variants, we believe it is the most plausible target.
503   In conclusion, studying how humans evolved in the face of changing environmental
pressures is necessary to understanding variation in present-day phenotypes and the potential
tradeoffs that influence propensity to different diseases in modern environments (Benton et al.
2021). Here, we show that genomic regions involved in circadian biology exhibited substantial
functional divergence between separate hominin populations. Furthermore, we show that
introgressed variants contribute to variation in AMH circadian phenotypes today in ways that are
consistent with an adaptive benefit.
510
511
512
513 **METHODS**
514
515 **Circadian gene selection**

Circadian biology is a complex system due to its high importance in the functioning of biological timing in diverse biological systems. For that reason, determining which genes are crucial for selection to environment response related to light exposure is not a straight forward process. To address this issue, we look at different sources of genome annotation databases and searched for genes and variants associated with circadian related phenotypes. We considered all human protein-coding genes in the Gene Ontology database annotated with the GO:0007623 ("circadian rhythm") term or terms annotated with relationship "is_a", "part_of", "occurs_in", or "regulates" circadian rhythm. We also considered genes containing experimental or orthologous evidence of circadian function in the Circadian Gene Database (CGDB), the GWAS Catalog genes containing "chronotype" or "circadian rhythm" associated variants, and a curated set of genes available in WikiPathways [https://www.wikipathways.org/index.php/Pathway:WP3594, https://doi.org/10.1093/nar/gkaa1024]. The final set of circadian genes was curated by Dr. Douglas McMahon.

To select the candidate circadian genes with the highest confidence, we defined a hierarchy system where genes annotated by McMahon or annotated in 3 out of 4 other sources receive a "High" level of confidence. Genes with evidence from 2 out of 4 of the sources are assigned a "Medium" level of confidence. Genes annotated as circadian only in 1 out of 4 sources are assigned to Low confidence and not considered in our circadian gene set. We then defined our set of circadian variants from the 1000 Genomes Project using the official list of circadian genes. The variants are included in analysis of coding, non-coding, regulatory, eQTL, human-specific, archaic-specific, and introgressed variants.

Definition of lineage-specific variants

To identify candidate variants that are specific to the human and the archaic lineages, we used a set of variants published by Kuhlwilm and Boeckx (Kuhlwilm and Boeckx 2019) (https://doi.org/10.1038/s41598-019-44877-x). The variants were extracted from the high-coverage genotypes of three archaics: a 122,000-year-old Neanderthal from the Altai Mountains (52x coverage), a 52,000-year-old Neanderthal from Vindija in Croatia (30x coverage), and a 72,000-year-old Denisovan from the Altai Mountains (30x coverage). The total variant sites retrieved from this set is 4,437,803. We used a filtering method similar to the one proposed in this publication. That is, a human-specific is defined as a position where all the humans carry the derived allele and all the archaics carry the ancestral allele. An archaic-specific is defined as a position where all the archaics carry the derived allele and the human allele frequency is not greater than 0.00001 (1e-5). These filters resulted in 9,424 human specific and 33,184 archaic-specific variants.

Enrichment of lineage-specific variants among functional regions of the genome

We intersected the sets of lineage-specific variants with several sets of annotated functional genomic regions. Inside circadian gene regions (Gencode v29), we found 156 human-specific variants and 341 archaic-specific variants. In circadian promoter regions, we found 6 human-specific variants and 19 archaic-specific variants. Promoters were defined as regions 5 kb up- to 1 kb downstream from a transcription start site. In distal regulatory elements, we found 247 human-specific variants and 807 archaic-specific variants. For this last set, we considered candidate cis-regulatory elements (cCREs) published by ENCODE (Moore et al. 2020) within 1 Mb of the circadian genes.

To compute whether lineage-specific variants are more abundant than expected in circadian genes, we applied a Fisher's exact test to the sets of human- and archaic-specific variants in regulatory, promoter, and gene regions. Human and archaic-specific variants are significantly enriched in both regulatory (Human: OR=1.25, P=8.39e-4; Archaic: OR=1.16, P=6.15e-5) and gene (Human: OR=1.84, P=7.06e-12; Archaic: OR=1.13, P=0.023) regions. The enrichment observed in the promoters of both lineages is not supported by a significant p-value (Human: OR=1.21, P=0.65; Archaic: OR=1.09, P=0.63).

Genes containing archaic variants with evidence of alternative splicing

We used a set of archaic variants annotated with the splice altering probabilities to identify circadian genes that may be differentially spliced between archaic hominins and AMH (Brand, Colbran, and Capra 2022). We considered variants from four archaic individuals: the Altai, Chagyrskaya, and Vindija Neanderthals and the Altai Denisovan. These archaic variants were annotated using SpliceAI (Jaganathan et al. 2019) and we considered any variant with a maximum delta, or splice altering probability, > 0.2. We identified 36 archaic-specific splice altering variants, defined as those variants absent from 1KGP, among 28 circadian genes. Next, we tested for enrichment among this gene set using an empirical null approach (Brand, Colbran, and Capra 2022; McArthur et al. 2022). We shuffled the maximum deltas among 1,607,350 variants 10,000 times and counted the number of circadian genes with a splice altering variant each iteration. Enrichment was calculated as the number of observed genes (N = 28) divided by the mean gene count among 10,000 shuffles. In addition to all genes with archaic-specific variants, we considered six other subsets among these variants: 1) genes with variants private to the Altai Neanderthal, 2) genes with variants private to the Chagyrskaya Neanderthal, 3) genes with variants private to the Altai Denisovan, 4) genes with variants private to all Neanderthals, 5) genes with variants shared among all archaic individuals, and 6) genes with variants private to the Vindija Neanderthal. Finally, we considered a subset of splice altering variants that were identified as tag SNPs by Vernot et al. (Vernot et al. 2016).

PrediXcan

To understand the difference in circadian biology between present-day humans and archaic hominins, we analyzed predictions on gene regulation. We considered the results from PrediXcan gene regulation predictions across 44 tissues from the PredictDB Data Repository (http://predictdb.org/). The models were trained on GTEx V6 using variants identified in 2,504 present-day humans in the 1000 Genomes Project (1KGP) phase 3 within 1 Mb of each circadian gene. The original analysis includes predictions for 17,748 genes for which the models explained a significant amount of variance in gene expression in each tissue (FDR < 0.05). The prediction models were also applied to the Altai and Vindija Neanderthals and the Denisovan. The resulting predictions are normalized values of the distribution observed in GTEx individuals used to train the original prediction models. Each prediction contains an empirical P-value which was calculated for each gene and tissue pair to define genes that are divergently regulated between archaic hominins and humans. The P-value is obtained by calculating the proportion of humans from the 1KGP that have predictions more extreme compared to the human median than the archaic individual. Significantly DR genes are defined as those where the archaic prediction falls outside the distribution of humans in the 1KGP predictions.

We tested whether the circadian genes in our set are more likely to be DR compared to an empirical null distribution from random gene sets of the same size. We account for the fact that

some genes are modeled in more tissues than others by matching the distribution of tissues in
which each gene could be modeled in the random sets to our set. Among 1,467 DR genes in the
Altai Neanderthal we find 23 DR circadian genes out of the total 236 genes in the circadian set.
We iterate through the permutation analysis 1,000,000 times and find an enrichment of 1.21
(P=0.19). A similar analysis is done in the Vindija Neanderthal (1,536 total DR, 21 circadian
DR, enrichment of 1.05, P=0.43) and the Denisovan individual (1,214 total DR, 19 circadian DR,
enrichment of 1.20, P=0.24). In this study, we define a set of DR genes as the intersection
between DR genes in all three archaics, resulting in a set of 16 genes.
615

Enrichment of introgressed variants in eQTL
We performed an enrichment analysis using Pearson's chi-squared test to evaluate if there is
overrepresentation of introgressed alleles in our set of circadian variants using the GTEx dataset.
We did a liftOver of the GTEx v8 dataset from hg38 to hg19. The original hg38 set contains
4,631,659 eQTLs across 49 tissues. After the LiftOver, 4,608,446 eQTLs remained, with the rest
not mapping. We used the archaic introgressed variants dataset from Browning 2018. The set
contains 863,539 variants that are introgressed in humans originating in archaic hominins. We
performed an intersection between the set of genes containing evidence for eQTLs and our set of
246 circadian genes to retrieve a subset of variant sites with evidence of being eQTL in circadian
genes. The resulting subset contained 97,441 circadian eQTLs in 49 tissues and 239 genes. We
further intersected the introgressed variants and the set of eQTL, resulting in 128,138
introgressed eQTLs. The final set of eQTLs that are circadian and also introgressed is 3,857.
628

Direction of effect of chronotype associations
To explore the effect of archaic introgression in circadian dreams on human chronotype, we
quantified the direction of effect of variants associated to a Morning/Evening person trait in a
GWAS analysis of the UK Biobank (http://www.nealelab.is/uk-biobank/). The variants were LD
clumped using PLINK v1.9 (R2 0.9). We generated cumulative proportion values on the beta
values assigned to each associated variant on an ascending order of P-values.
635

Detection of pleiotropy in the set of introgressed circadian variants
To understand the extent of different phenotypes associated with the introgressed circadian
variants, we first extracted genome-wide associations from Open Targets Genetics
(https://genetics.opentargets.org/) for each of the variants with evidence of introgression
(Browning et al. 2018). Only the variants with significant p-values were analyzed. The p-value
threshold was set at the genome-wide significance level (P=5e-8). We split the variants in two
sets: introgressed circadian and introgressed non-circadian. Many of these variants are not
associated with any phenotype. We performed a Fisher's exact test to analyze which of the two
sets contains a higher ratio of SNPs with at least one association versus SNPs with no
association. The result showed that the circadian set had a significantly higher ratio (OR=1.36,
P=5e-29). Then we calculated the total of unique traits associated with each of the variants, given
that the SNP has at least one association. We used a Mann-Whitney U test to understand which
set is represented by a higher level of traits per SNP. The circadian set was slightly more
pleiotropic, and the result is supported by a significant p-value (P=5.4e-3).
650

651 **Identifying introgressed circadian variants with evidence of adaptive introgression**

We sought out to identify circadian variants that contain evidence of adaptive introgression (AI). To achieve this, we collected two sets of genomic regions that were measured for their likelihood to be under AI by two machine learning methods: *genomatnn* and *MaLAdapt*. *genomatnn* is a convolutional neural network trained to identify adaptive introgression based on simulations (Gower et al. 2021). *MaLAdapt* is a machine learning algorithm trained to find adaptive introgression based on simulations using an extra-trees classifier (ETC) (X. Zhang et al. 2023). Following the thresholds used in each paper, a region is considered to be under AI if the prediction value assigned to it meets a threshold of 0.5 or 0.9, respectively. Because the two methods are different, we performed two separate analyses instead of integrating them. To find the variants that fall into AI regions, we intersected the set of introgressed circadian SNPs with the genomatnn and the MaLAdapt regions individually. The set of introgressed circadian variants contains variants inside circadian genes, in circadian promoter regions (5 kb up- and 1 kb downstream of the TSS), and variants with regulatory function (cCREs) flanking circadian genes by 1 Mb. We found 53 introgressed circadian variants in 7 genes defined as AI by genomatnn, and 116 variants in 19 genes defined as AI by MaLAdapt.

DATA AVAILABILITY
The data underlying this article are available in the article and in its online supplementary material.

DECLARATION OF INTERESTS
The authors declare that they have no competing interests.

ACKNOWLEDGMENTS
We thank members of the Capra Lab for helpful comments on this work. This work was conducted in part using the resources of the Advanced Computing Center for Research and Education at Vanderbilt University, Nashville, TN. This work was supported by the National Institutes of Health [R35GM127087 to JAC, R01GM117650 to DM, F30HG011200 to EM, T32GM080178 to Vanderbilt University (EM), and T32HG009495 to the University of Pennsylvania (LLC)].

AUTHOR CONTRIBUTIONS
Conceptualization: KV, JAC; Methodology: KV, LC, EM, CB, JS, DM, JAC; Investigation: KV, LC, EM, CB, JAC; Writing – Original Draft: KV, JAC; Writing – Review & Editing: KV, LC, EM, CB, DR, DM, JAC; Funding Acquisition: JAC; Resources: JAC; Supervision: JAC.

REFERENCES

Arnold, Lee J., Martina Demuro, Josep M. Parés, Juan Luis Arsuaga, Arantza Aranburu, José
      María Bermúdez de Castro, and Eudald Carbonell. 2014. "Luminescence Dating and
      Palaeomagnetic Age Constraint on Hominins from Sima de Los Huesos, Atapuerca, Spain."
      Journal of Human Evolution 67 (1): 85–107. https://doi.org/10.1016/j.jhevol.2013.12.001.

Bae, Christopher J., Katerina Douka, and Michael D. Petraglia. 2017. "On the Origin of Modern
Humans: Asian Perspectives." *Science* 358 (6368). https://doi.org/10.1126/science.aai9067.
Benton, Mary Lauren, Abin Abraham, Abigail L. LaBella, Patrick Abbot, Antonis Rokas, and
John A. Capra. 2021. "The Influence of Evolutionary History on Human Health and
Disease." *Nature Reviews Genetics*. Nature Research. https://doi.org/10.1038/s41576-020-
00305-9.
Brand, Colin M, Laura L Colbran, and John A Capra. 2022. "Resurrecting the Alternative
Splicing Landscape of Archaic Hominins Using Machine Learning." *BioRxiv*.
Brown, Steven A., Dieter Kunz, Amelie Dumas, Pål O. Westermark, Katja Vanselow, Amely
Tilmann-Wahnschaffe, Hanspeter Herzel, and Achim Kramer. 2008. "Molecular Insights
into Human Daily Behavior." *Proceedings of the National Academy of Sciences of the
United States of America* 105 (5): 1602–7. https://doi.org/10.1073/pnas.0707772105.
Browning, Sharon R., Brian L. Browning, Ying Zhou, Serena Tucci, and Joshua M. Akey. 2018.
"Analysis of Human Sequence Data Reveals Two Pulses of Archaic Denisovan
Admixture." *Cell* 173 (1): 53–61. https://doi.org/https://doi.org/10.1016/j.cell.2018.02.031.
Colbran, Laura L., Eric R. Gamazon, Dan Zhou, Patrick Evans, Nancy J. Cox, and John A.
Capra. 2019. "Inferred Divergent Gene Regulation in Archaic Hominins Reveals Potential
Phenotypic Differences." *Nature Ecology & Evolution* 3 (November): 1598–1606.
https://doi.org/10.1038/s41559-019-0996-x.
Dannemann, Michael, and Janet Kelso. 2017. "The Contribution of Neanderthals to Phenotypic
Variation in Modern Humans." *American Journal of Human Genetics* 101 (4): 578–589.
https://doi.org/10.1016/j.ajhg.2017.09.010.
Dorokhov, Vladimir B, Alexandra N Puchkova, Anton O Taranov, A Petr, Tatiana V Tupitsina,
Igor D Ivanov, Valentin A Vavilin, et al. 2018. "An Hour in the Morning Is Worth Two in
the Evening: Association of Morning Component of Morningness – Eveningness with
Single Nucleotide Polymorphisms in Circadian Clock Genes." *Biological Rhythm Research*
49 (4): 622–42. https://doi.org/10.1080/09291016.2017.1390823.
Gan, Yong, Liqing Li, Liangwen Zhang, Shijiao Yan, Chao Gao, Sai Hu, Yan Qiao, Sha Tang,
Chao Wang, and Zuxun Lu. 2018. "Association between Shift Work and Risk of Prostate
Cancer: A Systematic Review and Meta-Analysis of Observational Studies."
Carcinogenesis. https://doi.org/10.1093/carcin/bgx129.
Gan, Yong, Chen Yang, Xinyue Tong, Huilian Sun, Yingjie Cong, Xiaoxu Yin, Liqing Li, et al.
2015. "Shift Work and Diabetes Mellitus: A Meta-Analysis of Observational Studies."
Occupational and Environmental Medicine. https://doi.org/10.1136/oemed-2014-102150.
Gómez-Robles, Aida. 2019. "Dental Evolutionary Rates and Its Implications for the
Neanderthal–Modern Human Divergence." *Science Advances* 5 (5): eaaw1268.
https://doi.org/10.1126/sciadv.aaw1268.
Gower, Graham, Pablo Iáñez Picazo, Matteo Fumagalli, and Fernando Racimo. 2021. "Detecting
Adaptive Introgression in Human Evolution Using Convolutional Neural Networks." *ELife*
10 (May). https://doi.org/10.7554/eLife.64669.
Green, Richard E, Johannes Krause, Adrian W Briggs, Tomislav Maricic, Udo Stenzel, Martin
Kircher, Nick Patterson, et al. 2010. "A Draft Sequence of the Neandertal Genome."
Science 328 (5979): 710–22. https://doi.org/10.1126/science.1188021.
Gyarmati, Georgina, Michelle C. Turner, Gemma Castaño-Vinyals, Ana Espinosa, Kyriaki
Papantoniou, Juan Alguacil, Laura Costas, et al. 2016. "Night Shift Work and Stomach
Cancer Risk in the MCC-Spain Study." *Occupational and Environmental Medicine*.

https://doi.org/10.1136/oemed-2016-103597.

Hublin, Jean Jacques, Abdelouahed Ben-Ncer, Shara E. Bailey, Sarah E. Freidline, Simon
Neubauer, Matthew M. Skinner, Inga Bergmann, et al. 2017. "New Fossils from Jebel
Irhoud, Morocco and the Pan-African Origin of Homo Sapiens." *Nature*.
https://doi.org/10.1038/nature22336.

Huerta-Sánchez, Emilia, Xin Jin, Asan, Zhuoma Bianba, Benjamin M. Peter, Nicolas
Vinckenbosch, Yu Liang, et al. 2014. "Altitude Adaptation in Tibetans Caused by
Introgression of Denisovan-like DNA." *Nature*. https://doi.org/10.1038/nature13408.

Hut, Roelof A., Silvia Paolucci, Roi Dor, Charalambos P. Kyriacou, and Serge Daan. 2013.
"Latitudinal Clines: An Evolutionary View on Biological Rhythms." *Proceedings of the*
Royal Society B: Biological Sciences 280 (1765). https://doi.org/10.1098/rspb.2013.0433.

Jacobs, Guy S., Georgi Hudjashov, Lauri Saag, Pradiptajati Kusuma, Chelzie C. Darusallam,
Daniel J. Lawson, Mayukh Mondal, et al. 2019. "Multiple Deeply Divergent Denisovan
Ancestries in Papuans." *Cell* 177 (April). https://doi.org/10.1016/j.cell.2019.02.035.

Jaganathan, Kishore, Sofia Kyriazopoulou Panagiotopoulou, Jeremy F. McRae, Siavash Fazel
Darbandi, David Knowles, Yang I. Li, Jack A. Kosmicki, et al. 2019. "Predicting Splicing
from Primary Sequence with Deep Learning." *Cell* 176 (3): 535-548.e24.
https://doi.org/10.1016/j.cell.2018.12.015.

Jones, Samuel E., Jacqueline M. Lane, Andrew R. Wood, Vincent T. van Hees, Jessica Tyrrell,
Robin N. Beaumont, Aaron R. Jeffries, et al. 2019. "Genome-Wide Association Analyses of
Chronotype in 697,828 Individuals Provides Insights into Circadian Rhythms." *Nature*
Communications 10 (343). https://doi.org/10.1038/s41467-018-08259-7.

Jones, Samuel E., Jessica Tyrrell, Andrew R. Wood, Robin N. Beaumont, Katherine S. Ruth,
Marcus A. Tuke, Hanieh Yaghootkar, et al. 2016. "Genome-Wide Association Analyses in
128,266 Individuals Identifies New Morningness and Sleep Duration Loci." *PLoS Genetics*
12 (6): e1006125. https://doi.org/10.1371/journal.pgen.1006125.

Kim, Jae Kyoung, and Daniel B. Forger. 2012. "A Mechanism for Robust Circadian
Timekeeping via Stoichiometric Balance." *Molecular Systems Biology* 8 (1).
https://doi.org/10.1038/msb.2012.62.

Kivelä, Liia, Marinos Rodolfos Papadopoulos, and Niki Antypa. 2018. "Chronotype and
Psychiatric Disorders." *Current Sleep Medicine Reports* 4 (2): 94–103.
https://doi.org/10.1007/s40675-018-0113-8.

Knutson, Kristen L., and Malcolm von Schantz. 2018. "Associations between Chronotype,
Morbidity and Mortality in the UK Biobank Cohort." *Chronobiology International* 35 (8):
1045–53. https://doi.org/10.1080/07420528.2018.1454458.

Kuhlwilm, Martin, and Cedric Boeckx. 2019. "A Catalog of Single Nucleotide Changes
Distinguishing Modern Humans from Archaic Hominins." *Nature Scientific Reports* 9
(8463). https://doi.org/10.1038/s41598-019-44877-x.

Lane, Jacqueline M., Irma Vlasac, Simon G. Anderson, Simon D. Kyle, William G. Dixon,
David A. Bechtold, Shubhroz Gill, et al. 2016. "Genome-Wide Association Analysis
Identifies Novel Loci for Chronotype in 100,420 Individuals from the UK Biobank." *Nature*
Communications 7 (1): 1–10. https://doi.org/10.1038/ncomms10889.

Larcher, Sandra, P. Y. Benhamou, J. L. Pépin, and A. L. Borel. 2015. "Sleep Habits and
Diabetes." *Diabetes and Metabolism*. https://doi.org/10.1016/j.diabet.2014.12.004.

Larcher, Sandra, Anne Sophie Gauchez, Sandrine Lablanche, Jean Louis Pépin, Pierre Yves
Benhamou, and Anne Laure Borel. 2016. "Impact of Sleep Behavior on Glycemic Control

in Type 1 Diabetes: The Role of Social Jetlag." *European Journal of Endocrinology*. https://doi.org/10.1530/EJE-16-0188.

Leocadio-Miguel, Mario André, Fernando Mazzili Louzada, Leandro Lourenção Duarte, Roberta Peixoto Areas, Marilene Alam, Marcelo Ventura Freire, John Fontenele-araujo, Luiz Menna-barreto, and Mario Pedrazzoli. 2017. "Latitudinal Cline of Chronotype." *Scientific Reports* 7 (5437): 2–7. https://doi.org/10.1038/s41598-017-05797-w.

Lowden, Arne, Nelson Lemos, Bruno Gonçalves, Gülçin Öztürk, Fernando Louzada, Mario Pedrazzoli, and Claudia Moreno. 2018. "Delayed Sleep in Winter Related to Natural Daylight Exposure among Arctic Day Workers." *Clocks & Sleep* 1 (1): 105–16. https://doi.org/10.3390/clockssleep1010010.

Mafessoni, Fabrizio, Steffi Grote, Cesare De Filippo, Viviane Slon, Kseniya A. Kolobova, Bence Viola, Sergey V. Markin, et al. 2020. "A High-Coverage Neandertal Genome from Chagyrskaya Cave." *Proceedings of the National Academy of Sciences of the United States of America* 117 (26). https://doi.org/10.1073/pnas.2004944117.

McArthur, Evonne, David C. Rinker, and John A. Capra. 2021. "Quantifying the Contribution of Neanderthal Introgression to the Heritability of Complex Traits." *Nature Communications* 12 (1): 1–14. https://doi.org/10.1038/s41467-021-24582-y.

McArthur, Evonne, David C. Rinker, Erin N. Gilbertson, Geoff Fudenberg, Maureen Pittman, Kathleen Keough, Katherine S. Pollard, and John A. Capra. 2022. "Reconstructing the 3D Genome Organization of Neanderthals Reveals That Chromatin Folding Shaped Phenotypic and Sequence Divergence." *BioRxiv*. https://www.biorxiv.org/content/10.1101/2022.02.07.479462v1%0Ahttps://www.biorxiv.org/content/10.1101/2022.02.07.479462v1.abstract.

Meyer, Matthias, Juan Luis Arsuaga, Cesare De Filippo, Sarah Nagel, Ayinuer Aximu-Petri, Birgit Nickel, Ignacio Martínez, et al. 2016. "Nuclear DNA Sequences from the Middle Pleistocene Sima de Los Huesos Hominins." *Nature* 531 (7595). https://doi.org/10.1038/nature17405.

Meyer, Matthias, Qiaomei Fu, Ayinuer Aximu-Petri, Isabelle Glocke, Birgit Nickel, Juan Luis Arsuaga, Ignacio Martínez, et al. 2014. "A Mitochondrial Genome Sequence of a Hominin from Sima de Los Huesos." *Nature* 505 (7483). https://doi.org/10.1038/nature12788.

Meyer, Matthias, Martin Kircher, Marie Theres Gansauge, Heng Li, Fernando Racimo, Swapan Mallick, Joshua G. Schraiber, et al. 2012. "A High-Coverage Genome Sequence from an Archaic Denisovan Individual." *Science*. https://doi.org/10.1126/science.1224344.

Michael, Todd P., Patrice A. Salomé, Hannah J. Yu, Taylor R. Spencer, Emily L. Sharp, Mark A. McPeek, José M. Alonso, Joseph R. Ecker, and C. Robertson McClung. 2003. "Enhanced Fitness Conferred by Naturally Occurring Variation in the Circadian Clock." *Science* 302 (5647): 1049–53. https://doi.org/10.1126/science.1082971.

Moore, Jill E., Michael J. Purcaro, Henry E. Pratt, Charles B. Epstein, Noam Shoresh, Jessika Adrian, Trupti Kawli, et al. 2020. *Expanded Encyclopaedias of DNA Elements in the Human and Mouse Genomes. Nature*. Vol. 583. https://pubmed.ncbi.nlm.nih.gov/32728249/.

Nielsen, Rasmus, Joshua M. Akey, Mattias Jakobsson, Jonathan K. Pritchard, Sarah Tishkoff, and Eske Willerslev. 2017. "Tracing the Peopling of the World through Genomics." *Nature* 541 (7637): 302–10. https://doi.org/10.1038/nature21347.

O'Malley, Kathleen G., and Michael A. Banks. 2008. "A Latitudinal Cline in the Chinook Salmon (Oncorhynchus Tshawytscha) Clock Gene: Evidence for Selection on PolyQ

Length Variants." *Proceedings of the Royal Society B: Biological Sciences* 275 (1653): 2813–21. https://doi.org/10.1098/rspb.2008.0524.

O'Malley, Kathleen G., Michael J. Ford, and Jeffrey J. Hard. 2010. "Clock Polymorphism in Pacific Salmon: Evidence for Variable Selection along a Latitudinal Gradient." In *Proceedings of the Royal Society B: Biological Sciences*, 277:3703–14. Royal Society. https://doi.org/10.1098/rspb.2010.0762.

Papantoniou, Kyriaki, Gemma Castaño-Vinyals, Ana Espinosa, Nuria Aragonés, Beatriz Pérez-Gómez, Eva Ardanaz, Jone Miren Altzibar, et al. 2016. "Breast Cancer Risk and Night Shift Work in a Case–Control Study in a Spanish Population." *European Journal of Epidemiology*. https://doi.org/10.1007/s10654-015-0073-y.

Papantoniou, Kyriaki, Gemma Castaño-Vinyals, Ana Espinosa, Michelle C. Turner, Maria Henar Alonso-Aguado, Vicente Martin, Nuria Aragonés, et al. 2017. "Shift Work and Colorectal Cancer Risk in the MCC-Spain Case–Control Study." *Scandinavian Journal of Work, Environment and Health*. https://doi.org/10.5271/sjweh.3626.

Petr, Martin, Svante Pääbo, Janet Kelso, and Benjamin Vernot. 2019. "Limits of Long-Term Selection against Neandertal Introgression." *Proceedings of the National Academy of Sciences* 116 (5): 1639–1644. https://doi.org/10.1073/pnas.1814338116.

Prüfer, Kay, Cesare De Filippo, Steffi Grote, Fabrizio Mafessoni, Petra Korlević, Mateja Hajdinjak, Benjamin Vernot, et al. 2017. "A High-Coverage Neandertal Genome from Vindija Cave in Croatia." *Science* 358: 655–658. https://doi.org/10.1126/science.aao1887.

Prüfer, Kay, Fernando Racimo, Nick Patterson, Flora Jay, Sriram Sankararaman, Susanna Sawyer, Anja Heinze, et al. 2014. "The Complete Genome Sequence of a Neanderthal from the Altai Mountains." *Nature* 505 (7481): 43–49. https://doi.org/10.1038/nature12886.

Putilov, Arcady A., Vladimir B. Dorokhov, and Michael G. Poluektov. 2018. "How Have Our Clocks Evolved? Adaptive and Demographic History of the out-of-African Dispersal Told by Polymorphic Loci in Circadian Genes." *Chronobiology International* 35 (4): 511–532. https://doi.org/10.1080/07420528.2017.1417314.

Putilov, Arcady A., Vladimir B. Dorokhov, Alexandra N. Puchkova, Gleb N. Arsenyev, and Dmitry S. Sveshnikov. 2019. "Genetic-Based Signatures of the Latitudinal Differences in Chronotype." *Biological Rhythm Research* 50 (2): 255–71. https://doi.org/10.1080/09291016.2018.1465249.

Racimo, Fernando, David Gokhman, Matteo Fumagalli, Amy Ko, Torben Hansen, Ida Moltke, Anders Albrechtsen, Liran Carmel, Emilia Huerta-Sanchez, and Rasmus Nielsen. 2017. "Archaic Adaptive Introgression in TBX15/WARS2." *Molecular Biology and Evolution* 34 (3): 509–24. https://doi.org/10.1093/molbev/msw283.

Racimo, Fernando, Davide Marnetto, and Emilia Huerta-Sánchez. 2017. "Signatures of Archaic Adaptive Introgression in Present-Day Human Populations." *Molecular Biology and Evolution* 34 (2): 296–317. https://doi.org/10.1093/molbev/msw216.

Racimo, Fernando, Sriram Sankararaman, Rasmus Nielsen, and Emilia Huerta-Sánchez. 2015. "Evidence for Archaic Adaptive Introgression in Humans." *Nature Reviews Genetics* 16 (6): 359–71. https://doi.org/10.1038/nrg3936.

Randler, Christoph, and Arash Rahafar. 2017. "Latitude Affects Morningness- Eveningness: Evidence for the Environment Hypothesis Based on a Systematic Review." *Nature Publishing Group* 7 (39976): 1–6. https://doi.org/10.1038/srep39976.

Rinker, David C., Corinne N. Simonti, Evonne McArthur, Douglas Shaw, Emily Hodges, and John A. Capra. 2020. "Neanderthal Introgression Reintroduced Functional Ancestral Alleles

Lost in Eurasian Populations." *Nature Ecology and Evolution* 4 (10): 1332–41.
https://doi.org/10.1038/s41559-020-1261-z.
Sandrelli, Federica, Eran Tauber, Mirko Pegoraro, Gabriella Mazzotta, Paola Cisotto, Johannes
Landskron, Ralf Stanewsky, et al. 2007. "A Molecular Basis for Natural Selection at the
Timeless Locus in Drosophila Melanogaster." *Science* 316 (5833): 1898–1900.
https://doi.org/10.1126/science.1138426.
Sankararaman, Sriram, Nick Patterson, Heng Li, Svante Pääbo, and David Reich. 2012. "The
Date of Interbreeding between Neandertals and Modern Humans." *PLoS Genetics* 8
(e1002947). https://doi.org/10.1371/journal.pgen.1002947.
Shi, Yan, Li Liu, Tsuyoshi Hamada, Jonathan A. Nowak, Marios Giannakis, Yanan Ma,
Mingyang Song, et al. 2020. "Night-Shift Work Duration and Risk of Colorectal Cancer
According to IRS1 and IRS2 Expression." *Cancer Epidemiology Biomarkers and*
Prevention. https://doi.org/10.1158/1055-9965.EPI-19-0325.
Skoglund, Pontus, and Iain Mathieson. 2018. "Ancient Genomics of Modern Humans: The First
Decade." *Annual Review of Genomics and Human Genetics*. Annual Reviews Inc.
https://doi.org/10.1146/annurev-genom-083117-021749.
Skov, Laurits, Moisès Coll Macià, Garðar Sveinbjörnsson, Fabrizio Mafessoni, Elise A. Lucotte,
Margret S. Einarsdóttir, Hakon Jonsson, et al. 2020. "The Nature of Neanderthal
Introgression Revealed by 27,566 Icelandic Genomes." *Nature* 582 (7810): 78–83.
https://doi.org/10.1038/s41586-020-2225-9.
Srinivasan, Venkataramanujan, Marcel Smits, Warren Spence, Alan Lowe, Leonid Kayumov,
Seithikurippu Pandi-Perumal, Barbara Parry, and Daniel Cardinali. 2006. "Melatonin in
Mood Disorders." *World Journal of Biological Psychiatry*.
https://doi.org/10.1080/15622970600571822.
Stringer, Chris. 2016. "The Origin and Evolution of Homo Sapiens." *Philosophical Transactions*
of the Royal Society B: Biological Sciences 371 (1698): 20150237.
https://doi.org/10.1098/rstb.2015.0237.
Tauber, Eran, Mauro Zordan, Federica Sandrelli, Mirko Pegoraro, Nicolò Osterwalder, Carlo
Breda, Andrea Daga, et al. 2007. "Natural Selection Favors a Newly Derived Timeless
Allele in Drosophila Melanogaster." *Science* 316 (5833): 1895–98.
https://doi.org/10.1126/science.1138412.
Taylor, Briana J., and Brant P. Hasler. 2018. "Chronotype and Mental Health: Recent
Advances." *Current Psychiatry Reports* 20 (8). https://doi.org/10.1007/s11920-018-0925-8.
Telis, Natalie, Robin Aguilar, and Kelley Harris. 2020. "Selection against Archaic Hominin
Genetic Variation in Regulatory Regions." *Nature Ecology and Evolution* 4 (11): 1558–66.
https://doi.org/10.1038/s41559-020-01284-0.
Vernot, Benjamin, and Joshua M. Akey. 2014. "Resurrecting Surviving Neandertal Lineages
from Modern Human Genomes." *Science* 343 (6174): 1017–21.
https://doi.org/10.1126/science.1245938.
Vernot, Benjamin, Serena Tucci, Janet Kelso, Joshua G. Schraiber, Aaron B. Wolf, Rachel M.
Gittelman, Michael Dannemann, et al. 2016. "Excavating Neandertal and Denisovan DNA
from the Genomes of Melanesian Individuals." *Science* 352 (6282): 235–39.
https://doi.org/10.1126/science.aad9416.
Villanea, Fernando A., and Joshua G. Schraiber. 2019. "Multiple Episodes of Interbreeding
between Neanderthal and Modern Humans." *Nature Ecology and Evolution* 3 (1): 39–44.
https://doi.org/10.1038/s41559-018-0735-8.

Yousef, Einas, Noha Mitwally, Noha Noufal, and Muhammad Ramzan Tahir. 2020. "Shift Work
and Risk of Skin Cancer: A Systematic Review and Meta-Analysis." *Scientific Reports* 10
(1): 1–11. https://doi.org/10.1038/s41598-020-59035-x.
Zhang, Qingzhu, Hongyu Li, Rui Li, Ruibo Hu, Chengming Fan, Fulu Chen, Zonghua Wang, Xu
Liu, Yongfu Fu, and Chentao Lin. 2008. "Association of the Circadian Rhythmic
Expression of GmCRY1a with a Latitudinal Cline in Photoperiodic Flowering of Soybean."
Proceedings of the National Academy of Sciences of the United States of America 105 (52):
21028–33. https://doi.org/10.1073/pnas.0810585105.
Zhang, Xinjun, Bernard Kim, Armaan Singh, Sriram Sankararaman, Arun Durvasula, and Kirk E
Lohmueller. 2023. "MaLAdapt Reveals Novel Targets of Adaptive Introgression From
Neanderthals and Denisovans in Worldwide Human Populations." *Molecular Biology and*
Evolution 40 (1): msad001.
940