1   Archaic Introgression Shaped Human Circadian Traits
2

3   Keila Velazquez-Arcelay[1], Laura L. Colbran[2], Evonne McArthur[3], Colin Brand[4,5], David
4   Rinker[1], Justin Siemann[1], Douglas McMahon[1], and John A. Capra[4,5*]
5
6   [1] Department of Biological Sciences, Vanderbilt University
7   [2] Department of Genetics, Perelman School of Medicine, University of Pennsylvania
8   [3] Vanderbilt University School of Medicine
9   [4] Department of Epidemiology and Biostatistics, University of California, San Francisco
10   [5] Bakar Computational Health Sciences Institute, University of California, San Francisco
11
[*] Corresponding author: tony@capralab.org

ABSTRACT

Introduction: When the ancestors of modern Eurasians migrated out of Africa and interbred with
Eurasian archaic hominins, namely Neanderthals and Denisovans, DNA of archaic ancestry
integrated into the genomes of anatomically modern humans. This process potentially
accelerated adaptation to Eurasian environmental factors, including reduced ultra-violet radiation
and increased variation in seasonal dynamics. However, whether these groups differed
substantially in circadian biology, and whether archaic introgression adaptively contributed to
human chronotypes remains unknown.
Results: Here we traced the evolution of chronotype based on genomes from archaic hominins
and present-day humans. First, we inferred differences in circadian gene sequences, splicing, and
regulation between archaic hominins and modern humans. We identified 28 circadian genes
containing variants with potential to alter splicing in archaics (e.g., *CLOCK*, *PER2*, *RORB*,
RORC), and 16 circadian genes likely divergently regulated between present-day humans and
archaic hominins, including *RORA*. These differences suggest the potential for introgression to
modify circadian gene expression. Testing this hypothesis, we found that introgressed variants
are enriched among eQTLs for circadian genes. Supporting the functional relevance of these
regulatory effects, we found that many introgressed alleles have associations with chronotype.
Strikingly, the strongest introgressed effects on chronotype increase morningness, consistent
with adaptations to high latitude in other species. Finally, we identified several circadian loci
with evidence of adaptive introgression or latitudinal clines in allele frequency.
Conclusions: These findings identify differences in circadian gene regulation between modern
humans and archaic hominins and support the contribution of introgression via coordinated
effects on variation in human chronotype.

37
38
39   **Keywords:** circadian biology; chronotype; Neanderthals; adaptive introgression; gene
40   expression; adaptive evolution
41
42

43   **SIGNIFICANCE STATEMENT**
44
Interbreeding between humans and Neanderthals created the potential for adaptive introgression
as humans moved into environments that had been populated by Neanderthals for hundreds of
thousands of years. Here we discover lineage-specific genetic differences in circadian genes and
their regulatory elements between humans and Neanderthals. We show that introgressed alleles
are enriched for effects on circadian gene regulation, consistently increase propensity for
morningness in Europeans, and show evidence of adaptive introgression or associations between
latitude and frequency. These results expand our understanding of how the genomes of humans
and our closest relatives responded to environments with different light/dark cycles, and
demonstrate a coordinated contribution of admixture to human chronotype in a direction that is
consistent with adaptation to higher latitudes.

INTRODUCTION

All anatomically modern humans (AMH) trace their origin to the African continent around 300 thousand years ago (ka) (Stringer, 2016; Hublin *et al.*, 2017), where environmental factors shaped many of their biological features. Approximately seventy-thousand years ago (Bae, Douka, and Petraglia 2017), the ancestors of modern Eurasian AMH began to migrate out of Africa, where they were exposed to diverse new environments. In Eurasia, the novel environmental factors included greater seasonal variation in temperature and photoperiod.

Changes in the pattern and level of light exposure have biological and behavioral consequences in organisms. For example, *D. melanogaster* that are native to Europe harbor a polymorphism in *timeless*, a key gene in the light response of the circadian system, that follows a latitudinal cline in allele frequency (Sandrelli et al. 2007; Tauber et al. 2007). The ancestral haplotype produces a short TIM (S-TIM) protein that is sensitive to degradation by light because of its strong affinity to cryptochromes (CRY), photoreceptor proteins involved in the entrainment of the circadian clock. An insertion of a G nucleotide in the 5' coding region of the gene originated approximately 10 kya in Europe and created a start codon that produces a new long TIM isoform (L-TIM). The L-TIM variant has a lower affinity to CRY, creating a change in photosensitivity and altering the length of the period. L-TIM flies are at a higher frequency in southern Europe, while S-TIM flies are more prevalent in northern Europe. Another example is found in pacific salmon. Chinook salmon (*Oncorhynchus tshawytscha*) populations show a latitudinal cline in the frequency and length of repeat motifs in the gene *OtsClock1b*, strongly suggesting that this locus is under selection associated with latitude and photoperiod (O'Malley, Ford, and Hard 2010; O'Malley and Banks 2008). The evolution of circadian adaptation to diverse environments has also been widely studied in insects, plants (Michael *et al.*, 2003; Zhang *et al.*, 2008), and fishes, but it is understudied in humans. Adaptive processes could have helped to align human biology and chronotype to new natural conditions.

Previous studies in humans found a correlation between latitude and chronotype (morningness vs. eveningness) variation (Leocadio-Miguel et al. 2017; Lowden et al. 2018; Randler and Rahafar 2017) and a latitudinal cline in some circadian allele frequencies (Dorokhov *et al.*, 2018; Putilov, Dorokhov and Poluektov, 2018; Putilov *et al.*, 2019), highlighting the contribution of the environment to behavior and circadian biology. Many human health effects are linked to the misalignment of chronotype (Knutson and von Schantz 2018), including cancer, obesity (Gyarmati *et al.*, 2016; Papantoniou *et al.*, 2016, 2017; Gan *et al.*, 2018; Shi *et al.*, 2020; Yousef *et al.*, 2020), and diabetes (Gan *et al.*, 2015; Larcher *et al.*, 2015, 2016). There is also evidence of a correlation between evening chronotype and mood disorders, most notably seasonal affective disorder (SAD), depression, and worsening of bipolar disorder episodes (Srinivasan *et al.*, 2006; Kivelä, Papadopoulos and Antypa, 2018; Taylor and Hasler, 2018). Thus, we hypothesize that the differences in geography and environment encountered by early AMH populations moving into higher latitudes created potential for circadian misalignment and health risk.

Although AMHs arrived in Eurasia ~70 ka, other hominins (e.g., Neanderthals and Denisovans) lived there for more than 400 ka (Arnold *et al.*, 2014; Meyer *et al.*, 2014, 2016). These archaic hominins diverged from AMHs around 700 ka (Meyer *et al.*, 2012; Prüfer *et al.*, 2014, 2017; Nielsen *et al.*, 2017; Gómez-Robles, 2019; Mafessoni *et al.*, 2020), and as a result, the ancestors of AMHs and archaic hominins evolved under different environmental conditions. While there was substantial variation in the latitudinal ranges of each group, the Eurasian

hominins largely lived at consistently higher latitudes and, thus, were exposed to higher
amplitude seasonal variation in photoperiods. Given the influence of environmental cues on
circadian biology, we hypothesized that these separate evolutionary histories produced
differences in circadian traits adapted to the distinct environments.
When AMH migrated into Eurasia, they interbred with the archaic hominins that were
native to the continent, initially with Neanderthals (Green et al. 2010; Villanea and Schraiber
2019) around 60 ka (Sankararaman et al. 2012; Skoglund and Mathieson 2018) and later with
Denisovans (Jacobs et al. 2019). Due to this, a substantial fraction (>40%) of the archaic
variation remains in present-day Eurasians (Skov et al. 2020; Vernot and Akey 2014), although
each human individual carries only ~2% DNA of archaic ancestry (Vernot *et al.*, 2016; Prüfer *et*
al., 2017). Most of the archaic ancestry in AMH was subject to strong negative selection, but
some of these introgressed alleles remaining in AMH populations show evidence of adaptation
(Racimo *et al.*, 2015; Gower *et al.*, 2021). For example, archaic alleles have been associated with
differences in hemoglobin levels at higher altitude in Tibetans, immune resistance to new
pathogens, levels of skin pigmentation, and fat composition (Huerta-Sánchez *et al.*, 2014;
Racimo *et al.*, 2015, 2017; Dannemann and Kelso, 2017; Racimo, Marnetto and Huerta-Sánchez,
2017; McArthur, Rinker and Capra, 2021). Previous work also suggests that introgressed alleles
could have adaptively influenced human chronotype. First, a phenome-wide association study
(PheWAS) in the UK Biobank found loci near *ASB1* and *EXOC6* with introgressed variants that
significantly associated with self-reported sleeping patterns (Dannemann and Kelso, 2017). One
of these alleles showed a significant association between frequency and latitude. Second,
summarizing effects genome-wide, introgressed alleles are also moderately enriched for
heritability of chronotype compared to non-introgressed alleles (McArthur, Rinker and Capra,
2021). These results suggest a potential role for introgressed alleles in adaptation to pressures
stemming from migration to higher latitudes.
Motivated by the potential for a role of archaic introgression in AMH circadian variation,
we explore two related questions: 1) Can comparative genomic analysis identify differences in
AMH and archaic hominin circadian biology?, and 2) Do introgressed archaic alleles influence
human circadian biology? Understanding the ancient history and evolution of chronotypes in
humans will shed light on human adaptation to high latitudes and provide context for the genetic
basis for the modern misalignment caused by the development of technology and night
shiftwork.
133
134
135
136 **RESULTS**
137
Did archaic hominins and modern humans diverge in circadian biology?
Following divergence ~700,000 years ago (ka) (Nielsen *et al.*, 2017; Gómez-Robles, 2019),
archaic hominins and AMH were geographically isolated, resulting in the accumulation of
lineage-specific genetic variation and phenotypes (Figure 1). In the next several sections, we
evaluate the genomic evidence for divergence in circadian biology between archaic hominin and
modern human genomes.
144
Identifying archaic-hominin-specific circadian gene variation

With the sequencing of several genomes of archaic hominins, we now have a growing, but
incomplete, catalog of genetic differences specific to modern and archaic lineages. Following
recent work (Kuhlwilm and Boeckx, 2019), we defined archaic-specific variants as genomic
positions where archaic hominins (Altai Neanderthal, Vindija Neanderthal, and Denisovan) all
have the derived allele while in humans the derived allele is absent or present at such an
extremely low frequency in the 1000 Genome Project (<0.00001) that it is likely an independent
occurrence. We defined human-specific variants as positions where all individuals in the 1000
Genomes Project carry the derived allele and all the archaics carry the ancestral allele.
We evaluated archaic-specific variants for their ability to influence proteins, splicing, and
regulation of 246 circadian genes (Methods). The circadian genes were identified by a
combination of literature search, expert knowledge, and existing annotations (Table S1; Figure
S1; Methods). The core circadian clock machinery is composed of a dimer between the CLOCK
and ARNTL (BMAL1) transcription factors, which binds to E-box enhancer elements and
activates the expression of the Period (*PER1/2/3*) and Cryptochrome (*CRY1/2*) genes (Figure
1C). PERs and CRYs form heterodimers that inhibit the positive drive of the CLOCK-BMAL1
dimer on E-boxes, inhibiting their own transcription in a negative feedback loop. CLOCK-
BMAL1 also drives the expression of many other clock-controlled genes (CCG), including
NR1D1/2 (Nuclear Receptor Subfamily 1 Group D Member 1 and 2), *RORA/B* (RAR Related
Orphan Receptor A and B), and *DBP* (D-Box Binding PAR BZIP Transcription Factor). ROR
and REV-ERB are transcriptional regulators of BMAL1. CK1 binds to the PER/CRY
heterodimer, phosphorylating PER and regulating its degradation. Similarly, FBXL3 marks CRY
for degradation. Beyond the core clock genes, we included other upstream and downstream
genes that are involved in maintenance and response of the clock (Table S1; Figure S1).
We identified 1,136 archaic-specific variants in circadian genes, promoters, and
candidate distal cis-regulatory elements (cCREs). The circadian genes with the most archaic-
specific variants are *CLDN4*, *NAMPT*, *LRPPRC*, *ATF4*, and *AHCY* (125, 112, 110, 104, 102
respectively) (Table S2).
173
174

175
Figure 1. Did the sharing of functionally diverged alleles from archaic hominins influence
human circadian biology? A) Anatomically modern humans and archaic hominins evolved
separately at different latitudes for hundreds of thousands of years. The ancestors of modern
Eurasian humans left Africa approximately 70 thousand years ago (ka) and admixed with
archaics, likely in southwestern Asia. The shaded purple range represents the approximate
Neanderthal range. The purple dot represents the location of the sequenced Denisovan individual
in the Altai Mountains; the full range of Denisovans is currently unknown. Silhouettes from
phylopic.org. **B)** After the split between the human and archaic lineages, each group accumulated
variation and evolved in their respective environments for approximately 700 ka. We first test for
evidence for divergent circadian evolution during this time. Humans acquired introgressed alleles
from Neanderthals and from Denisovans around 60 and 45 ka, respectively. These alleles
experienced strong selective pressures; however, ~40% of the genome retains archaic ancestry in
some modern populations. The second question we explore is whether introgression made
contributions to human circadian biology. **C)** The core circadian clock machinery is composed of
several transcription factors (ovals) that dimerize and interact with E-box enhancer elements and
each other to create a negative feedback loop. We defined a set of 246 circadian genes through a
combination of literature search, expert knowledge, and existing annotations (Table S1; Figure
S1; Methods). Lines with arrows represent activation, and lines with bars represent suppression.
194
195

Fixed human- and archaic-specific variants are enriched in circadian genes and associated
regulatory elements
After the archaic and AMH lineages diverged, each group accumulated genetic variation specific
to each group. Variants fixed in each lineage are likely to be enriched in genomic regions that
influence traits that experienced positive selection. We tested whether human- and archaic-
specific fixed variants are enriched compared to other variants in circadian genes, their
promoters, and in annotated candidate cis-regulatory elements within 1 megabase (Mb) (Figure
2). We found that human- and archaic-specific fixed variants are enriched in circadian genes
(Fisher's exact test; human: OR=1.84, P=7.06e-12; archaic: OR=1.13, P=0.023) and distal
regulatory elements (Fisher's exact test; human: OR=1.25, P=8.39e-4; archaic: OR=1.16,
P=6.15e-5) compared to variants derived on each lineage, but not fixed. Promoter regions have a
similar enrichment pattern as that in gene and regulatory regions, but the p-values are high
(Fisher's exact test; human: OR=1.21, P=0.65; archaic: OR=1.09, P=0.63). This is likely due to
the small number of such variants in promoters. These results suggest that both groups had a
greater divergence in genomic regions related to circadian biology than expected.
211

Enrichment

Human specific	1.25 (247)	1.21 (6)	1.84 (156)
Archaic specific	1.16 (807)	1.09 (19)	1.13 (341)

Regulatory Elements   Promoters   Genes

Circadian genes

212
Figure 2. **Human- and archaic-specific fixed variants are enriched in circadian regulatory,**
promoter, and gene regions. Human-specific fixed variants are significantly enriched compared
to variants that are not fixed in circadian regulatory elements (Fisher's exact: OR=1.25, P=8.39e-
4) and gene regions (Fisher's exact: OR=1.84, P=7.06e-12). Promoters show a similar
enrichment, but the higher p-value is the result of the small number of variants (Fisher's exact
test: OR=1.21, P=0.65). Likewise, archaic-specific variants are enriched in circadian regulatory
regions (Fisher's exact: OR=1.16, P=6.15e-5) and gene regions (Fisher's exact: OR=1.13,
P=0.023), with the promoters showing a similar trend (Fisher's exact test: OR=1.09, P=0.63).
The numbers in parentheses give the counts of fixed variants observed in each type of element.
Regulatory elements were defined based on the ENCODE candidate cis-regulatory elements.
223
224
Several core circadian genes have evidence of alternative splicing between humans and
archaic hominins

We find only two archaic-specific coding variants in circadian genes: one missense and one
synonymous. The missense variant (hg19: chr17_46923411_A_G) is in the gene *CALCOCO2*,
calcium-binding and coiled-coil domain-containing protein 2. SIFT, PolyPhen, and CADD all
predict that the variant does not have damaging effects. The second variant (hg19:
chr7_119914770_G_T) is in the gene *KCND2*, which encodes a component of a voltage-gated
potassium channel that contributes to the regulation of the circadian rhythm of action potential
firing, but it is synonymous and the variant effect predictors suggest it is tolerated.
To explore potential splicing differences in circadian genes between humans and
archaics, we applied SpliceAI to predict whether any sequence differences between modern
humans and archaics are likely to modify splicing patterns. Four archaic individuals were
included in this analysis (the Altai, the Vindija, the Chagyrskaya Neanderthals, and the Altai
Denisovan) (Meyer *et al.*, 2012; Prüfer *et al.*, 2014, 2017; Mafessoni *et al.*, 2020). We found that
28 genes contained at least one archaic-specific variant predicted to result in alternative splicing
in archaics. These included several of the core clock genes *CLOCK*, *PER2*, *RORB*, *RORC*, and
FBXL13 (Figure 3A,C; Table S3). For example, the variant chr2:239187088-239187089 in the
1st intron of *PER2* is predicted to result in a longer 5' UTR. The splice-altering variants were
largely specific to the two different archaic linages (Figure 3A), with 13 specific to the
Denisovan, 8 shared among the three Neanderthals, and only one shared among all four archaic
individuals.
246
Circadian gene regulatory divergence between humans and archaic hominins
Given the enrichment of variants in regulatory regions of circadian genes, we sought to explore
the potential for differences in circadian gene regulation between humans and archaics with
causes beyond single lineage-specific variants. We leveraged an approach we recently developed
for predicting gene regulatory differences between modern and archaic individuals from
combinations of genetic variants (Colbran *et al.*, 2019). The approach uses PrediXcan, an elastic
net regression method, to impute gene transcript levels in specific tissues from genetic variation.
Previous work demonstrated that this approach has a modest decrease in performance when
applied to Neanderthals, but that it can accurately applied between humans and Neanderthals for
thousands of genes. Here, we quantify differences in predicted regulation of the 246 circadian
genes between 2,504 humans in the 1000 Genomes Project (1000 Genomes Project Consortium,
2010) and the archaic hominins. The predicted regulation values are normalized to the
distribution in the training set from the Genotype Tissue Expression Atlas (GTEx).
260
261

262
Figure 3. Many circadian genes have evidence of alternative splicing and divergent
regulation between modern and archaic hominins. A) The distribution of the 28 predicted
archaic-specific splice-altering variants (SAV) in circadian genes across archaic individuals.
Most are specific to either the Denisovan or Neanderthal lineage (Table S3). **B)** The sharing of
predicted divergently regulated (DR) gene/tissue pairs across three archaic individuals.
(Predictions were not available for the Chagyrskaya Neanderthal.) Seventeen divergently
regulated gene/tissue pairs were present in all three archaics (representing 16 unique genes).
Additionally, 7 gene/tissue DR pairs are shared between the Altai Neanderthal and the
Denisovan individual. One pair is shared between the Vindija Neanderthal and the Denisovan
(Table S4). **C)** The proportion of circadian genes containing archaic splice-altering variants
predicted by SpliceAI (SAV; 11.4%) or divergently regulated circadian genes predicted by
PrediXcan (DR; 6.5%). Thus, 17.9% of the circadian genes are predicted to contain differences
to AMH via these mechanisms.
276
277
We first analyzed gene regulation predictions in the core circadian clock genes. Archaic gene
regulation was at the extremes of the human distribution for many core clock genes including
PER2, *CRY1*, *NPAS2*, *RORA, NR1D1* (Figure 4; Figure S2). For example, the regulation of
PER2 in the Altai and Vindija Neanderthals is lower than 2,491 of the 2,504 (99.48%) modern
humans considered. The Denisovan has a predicted *PER2* regulation that is lower than 2,410
(96.25%). Expanding to all circadian genes and requiring archaic regulation to be more extreme
than all humans (Methods), we identified 24 circadian genes across 23 tissues with strong
divergent regulation between humans and at least one archaic hominin (Figure 3B; Table S4).
For example, all archaic regulation values for *RORA*, a core clock gene, are lower than for any of
the 2,504 modern humans. We found that 16 of these genes (Figure S3; Table S4), including
RORA, *MYBBP1A*, and *TIMELESS*, were divergently regulated in all archaic individuals. This
represents 6.5% of all the circadian genes (Figure 3C). Surprisingly, the two Neanderthals only
shared one DR gene not found in the Denisovan, while the Altai Neanderthal and Denisovan
shared seven not found in Vindija (Figure 3B). The Altai and Vindija Neanderthals represent
deeply diverging lineages, and this result suggests that they may have experienced different
patterns of divergence in the regulation of their circadian genes.
Given these differences in circadian gene regulation between humans and archaics, we
tested whether circadian genes are more likely to be divergently regulated than other gene sets.
Each archaic individual shows nominal enrichment for divergent regulation of circadian genes,

and the enrichment was stronger (~1.2x) in the Altai Neanderthal and Denisovan individual.
However, given the small sample size, the P-values are moderate (Permutation test; Altai:
OR=1.21, P=0.19, Vindija: OR=1.05, P=0.43, Denisovan: OR=1.20, P=0.24).
300
301

302
Figure 4. Many circadian genes are divergently regulated between modern humans and
archaic hominins. Comparison of the imputed regulation of core circadian genes between 2504
humans in 1000 Genomes Phase 3 (gray bars) and three archaic individuals (vertical lines). For
each core circadian gene, the tissue with the lowest average P-value for archaic difference from
humans is plotted. Archaic gene regulation is at the extremes of the human distribution for
several core genes: *CRY1*, *PER2*, *NPAS2*, *NR1D1 RORA*. See Figure S2 for all core clock genes
and Figure S3 for all divergently regulated circadian genes.
310
311
312 **Did introgressed archaic variants influence modern human circadian biology?**
313 The previous sections demonstrate lineage-specific genetic variation in many genes and
314 regulatory elements essential to the function of the core circadian clock and related pathways.
315 Given this evidence of functional differences between archaic hominins and AMH in these
316 systems, we next evaluated the influence of archaic introgression on AMH circadian biology.
317
Introgressed variants are enriched in circadian gene eQTL
Given the differences between archaic and modern sequences of circadian genes and their
regulatory elements, we investigated whether Neanderthal introgression contributed functional
circadian variants to modern Eurasian populations. We considered a set of 863,539 variants with

evidence of being introgressed from archaic hominins to AMH (Browning *et al.*, 2018). These
variants were identified using the Sprime algorithm, which searches for regions containing a high
density of alleles in common with Neanderthals and not present or at very low frequency in
Africans. Since many approaches have been developed to identify introgressed variants, we also
considered two stricter sets: 47,055 variants that were supported by all of six different
introgression maps (Sankararaman *et al.*, 2014; Vernot *et al.*, 2016; Browning *et al.*, 2018;
Steinrücken *et al.*, 2018; Skov *et al.*, 2020; Schaefer, Shapiro and Green, 2021) and 755,653
variants that were supported by Sprime and at least one other introgression map. As described
below, our main results replicated on both of these stricter sets.
We first tested whether the presence of introgressed variants across modern individuals
associated with the expression levels of any circadian genes, i.e., whether the introgressed
variants are expression quantitative trait loci (eQTL). We identified 3,857 introgressed variants
associated with the regulation of circadian genes in modern non-Africans (Table S5). The genes
PTPRJ, *HTR1B*, *NR1D2*, *CLOCK*, and *ATOH7* had the most eQTL (304, 273, 262, 256, and 252
respectively). We found introgressed circadian eQTL for genes expressed in all tissues in GTEx,
except kidney cortex. Notably, several of these circadian genes (e.g., *NR1D2* and *CLOCK*) with
introgressed eQTL were also found to be divergently regulated in our comparison of modern and
archaic gene regulation. This indicates that some of the archaic-derived variants that contributed
to divergent regulation were retained after introgression and continue to influence circadian
regulation in modern humans.
Introgressed variants are significantly more likely to be eQTL for circadian genes than
expected by chance from comparison to all eQTL (Figure 5A; Fisher's exact test: OR=1.45,
P=9.71e-101). The stricter set of introgressed variants identified by Browning et al. plus at least
one other introgression map had similar levels of eQTL enrichment for circadian genes
(OR=1.47; P=2.4e-103). The highest confidence set of introgressed variants that were identified
by all six maps considered had even stronger enrichment (OR=1.68; P=6.5e-23).
Most core circadian genes are expressed broadly across tissues; the fraction expressed in
each GTEx tissue ranges from 57% (whole blood) to 83% in testis, and an average of 72%
(Table S6). As a result, we anticipated that the enrichment of introgressed variants among eQTL
for circadian genes would hold across tissues. Examining the associations in each tissue, we
found that introgressed eQTL showed significant enrichment for circadian genes in most tissues
(34 of 49; Figure 5B; Table S7) and trended this way in all but five. Given that tissues in GTEx
have substantial differences in sample size and cellular heterogeneity, statistical power to detect
enrichment differs. We anticipate that this is the main driver of differences in enrichment across
tissues.
These results suggest that circadian pressures were widespread across tissues. Given the
previously observed depletion for introgressed variants in regulatory elements and eQTL (Petr *et*
al., 2019; Rinker *et al.*, 2020; Telis, Aguilar and Harris, 2020), this enrichment for circadian
genes among introgressed eQTL is surprising and suggests that the archaic circadian alleles
could have been beneficial after introgression.
362
363

364
Figure 5. Circadian genes are enriched for introgressed eQTL. A) Archaic introgressed
variants are more likely to be eQTL for circadian genes in GTEx than for non-circadian genes
(Fisher's exact test: OR=1.45, P=9.71e-101). Purple represents the set of introgressed variants,
and blue represents the set of circadian variants. 3,857 are introgressed eQTL in circadian genes.
Gray represents the universe of all GTEx eQTLs lifted over to hg19. The overlaps are not to
scale. **B)** The enrichment for circadian genes among the targets of introgressed eQTLs in each
GTEx tissue. Introgressed eQTL in most tissues show significant enrichment for circadian genes
(Fisher's exact test; Table S7). Kidney cortex did not have any circadian introgressed eQTLs and
thus is not shown. Numbers inside the parenthesis indicate the count of variants in each tissue.
Gray bars indicate lack of statistical significance; light blue bars indicate nominal significance (p
<= 0.05); and dark blue bars indicate significance at the 0.05 level after Bonferroni multiple
testing correction (p <= 0.00102).

377
378
Introgressed variants predominantly increase propensity for morningness
After observing that circadian gene expression is influenced by archaic variants, we evaluated
whether these effects are likely to result in a change in organism-level phenotype. To do this, we
evaluated evidence that introgressed variants influence chronotype. The heritability of
chronotype has been estimated in a range from 12 to 38% (Jones *et al.*, 2016, 2019; Lane *et al.*,
2016), and previous studies have identified two introgressed loci associated with sleep patterns
(Dannemann and Kelso, 2017; Putilov *et al.*, 2019). We recently found modest enrichment for
heritability of chronotype (morning/evening person phenotype in a GWAS of the UK Biobank)
among introgressed variants genome-wide using stratified LD score regression (heritability
enrichment: 1.58, P=0.25) (McArthur, Rinker and Capra, 2021). This analysis also suggested
that introgressed variants were more likely to increase morningness.

To test for this proposed directional effect, we calculated the cumulative fraction of
introgressed loci associated with chronotype in the UK Biobank that increase morningness (after
collapsing based on LD at $R^2>0.5$ in EUR). The introgressed loci most strongly associated with
chronotype increase propensity for morningness (Figure 6; Table S8; Table S9). As the strength
of the association with morningness decreases, the bias begins to decrease, but the effect is
maintained well past the genome-wide significance threshold (P<5e-8). When focusing the
analysis on introgressed variants in proximity (<1 Mb) to circadian genes, the pattern becomes
even stronger. The bias toward morningness remains above 80% at the genome-wide
significance threshold. This result also held when limiting to introgressed variants found in
Browning plus one or all other introgression maps considered (Figure S4). This suggests that
introgressed variants act in a consistent direction on chronotype, especially when they influence
circadian genes.

404
Figure 6. Introgressed variants associate with increased morningness. The cumulative
fraction of introgressed loci significantly associated with the morning vs. evening person trait in
the UK Biobank that increase morningness (y-axis) at a given p-value threshold (x-axis).
Introgressed loci associated with chronotype are biased towards increasing morningness, and this
effect is greatest at the most strongly associated loci. Introgressed variants nearby (<1 Mb)
circadian genes (blue) are even more strongly biased towards increasing morningness than
introgressed variants overall (gray). Each dot (triangle) represents an associated locus; variants
were clumped by LD for each set ($R^2>0.5$ in EUR).

Circadian rhythms are involved in a wide variety of biological systems. To explore other
phenotypes potentially influenced by the introgressed circadian variants, we evaluated evidence
for pleiotropic associations. First, we retrieved all the genome-wide associations reported for
introgressed variants in the Open Targets Genetics (https://genetics.opentargets.org) database,
which combines GWAS data from the GWAS Catalog, UK Biobank, and several other sources.

Introgressed circadian variants are associated with traits from a diverse range of categories
(Table S10). Associations with blood related traits are by far the most common; however, this is
likely because they have more power in the UK Biobank. Overall, circadian introgressed variants
are significantly more likely to have at least one trait association than introgressed variants not in
the circadian set (Fisher's exact test: OR=1.25, P=7.03e-25) (Figure S5A). The circadian variants
also associate with significantly more traits per variant than the non-circadian set (Mann-
Whitney U: P=9.93e-14) (Figure S5B; Table S11). These results suggest effects for introgressed
circadian variants beyond chronotype.
428
Evidence for adaptive introgression at circadian loci
The gene flow from Eurasian archaic hominins into AMH contributed to adaptations to some of
the new environmental conditions encountered outside of Africa (Racimo *et al.*, 2015). The
above analyses demonstrate the effects of introgressed variants on circadian gene regulation and
chronotype. To explore whether these circadian regions show evidence of adaptive introgression,
we considered three sets of introgressed regions predicted to have contributed to AMH
adaptation: one from an outlier approach based on allele frequency statistics (Racimo, Marnetto
and Huerta-Sánchez, 2017) and two from recent machine learning algorithms: *genomatnn*
(Gower *et al.*, 2021) and *MaLAdapt* (Zhang *et al.*, 2023). We intersected the circadian
introgressed variants with the adaptive introgression regions from each method.
We identified 47 circadian genes with evidence of adaptive introgression at a nearby
variant from at least one of the methods (Table S12). No region was supported by all three
methods; however, six were shared between Racimo and *MaLAdapt* and three were shared by
Racimo and *genomatnn*. The relatively small overlap between these sets underscores the
challenges of identifying adaptive introgression. Nonetheless, these represent promising
candidate regions for further exploration of the effects of introgressed variants on specific
aspects of circadian biology. For example, an introgressed haplotype on chr10 tagged by
rs76647913 was identified by both *MaLAdapt* and Racimo. This introgressed haploype is an
eQTL for the nearby *ATOH7* gene in many GTEx tissues. *ATOH7* is a circadian gene that is
involved in retinal ganglion cell development, and mice with this gene knocked out are unable to
entrain their circadian clock based on light stimuli (Brzezinski *et al.*, 2005).
450
Latitudinal clines for introgressed circadian loci
Motivated by the previous discovery of an introgressed haplotype on chr2 that is associated with
chronotype and increases in frequency with latitude (Dannemann and Kelso, 2017; Putilov *et al.*,
2019), we also tested each introgressed circadian variant for a correlation between allele
frequency and latitude in modern non-African populations from the 1000 Genomes Project.
The strongest association between latitude and frequency was a large chromosome 2
haplotype that contains the previously discovered introgressed SNP (rs75804782, R=0.85)
associated with chronotype. This haplotype is present in all non-African populations, and
rs61332075 showed the strongest latitudinal cline (R=0.87). The second strongest consisted of a
smaller haplotype of introgressed variants a few kb upstream of the previous haplotype (tagged
by rs35333999 and rs960783) that overlaps the core circadian gene *PER2*. These variants have a
correlation between latitude and frequency of ~0.68 They are also in moderate LD (R^2 of ~0.35
in EUR) with an additional introgressed variant (rs62194932) that has a similar latitudinal cline
of 0.70 (Figure S6; Table S13). These variants are each in very low LD with the previously
discovered haplotype (R^2 of ~0.01) and are each supported by multiple introgression maps.

Moreover, these introgressed variants are absent in all EAS populations, absent or at very low
frequency in SAS (<3%), and at higher frequency in EUR populations (~13%).
The EUR-specific introgressed variant rs35333999 causes a missense change in the PER2
protein (V903I) that overlaps a predicted interaction interface with PPARG. PER2 controls lipid
metabolism by directly repressing PPARG's proadipogenic activity (Grimaldi *et al.*, 2010). The
rs62194932 variant is an eQTL of *HES6* in the blood in the eQTLGen cohort (Võsa *et al.*, 2021).
HES6 encodes a protein that contributes to circadian regulation of LDLR and cholesterol
homeostasis (Lee *et al.*, 2012).
Thus, this genomic region, that includes circadian genes and introgressed variants
associated with chronotype, has population-specific structure and at least two distinct sets of
introgressed variants with latitudinal clines and functional links to lipid metabolism. *PER2* is
also predicted to have lower gene regulation in archaic hominins than most humans (Figure 4).
and the Vindija Neanderthal carries a lineage-specific variant in this gene that has splice-altering
effects. These results together suggests that *PER2* may have experienced multiple functional
changes in different modern and archaic lineages, with potential adaptive effects mediated by
introgression.
We did not discover any other significant associations between latitude and frequency for
other introgressed circadian loci. The rapid migration and geographic turnover of populations in
recent human history is likely to obscure many latitude-dependent evolutionary signatures, so we
did not anticipate many circadian loci would have a strong signal.
486
487

488 **DISCUSSION**
489
The Eurasian environments where Neanderthals and Denisovans lived for several hundred
thousand years are located at higher latitudes with more variable photoperiods than the landscape
where AMH evolved before leaving Africa. Evaluating genetic variation that arose separately in
each of the archaic and AMH lineages after their split ~700 MYA, we identified lineage-specific
genetic variation in circadian genes, their promoters, and flanking distal regulatory elements. We
found that both archaic- and human-specific variants are observed more often than expected in
each class of functional region. This result suggests that, while each group evolved separately
during hundreds of thousands of years in divergent environments, both experienced pressure on
circadian related variation. Leveraging sequence-based machine learning methods, we identified
many archaic-specific variants likely to influence circadian gene splicing and regulation. For
example, core clock genes (*CLOCK*, *PER2*, *RORB*, *RORC*, and *FBXL13*) have archaic variants
predicted to cause alternative splicing compared to AMH. Several core genes were also predicted
in archaics to be at the extremes of human gene regulation, including *PER2*, *CRY1*, *NPAS2*,
RORA, NR1D1. Surprisingly, the Altai Neanderthal shared more divergent regulation in the
circadian genes with the Denisovan individual than the Vindija Neanderthals. The two
Neanderthals represent populations that were quite distantly diverged with substantially different
histories and geographical ranges. The Denisovan and Altai Neanderthal also come from the
same region in Siberia, while the Vindija Neanderthal came from a region in Croatia with
slightly lower latitude.
Introgression introduced variation that first appeared in the archaic hominin lineage into
Eurasian AMH. While most of this genetic variation experienced strong negative selection in
AMH, a smaller portion is thought to have provided adaptive benefits in the new environments

(Racimo *et al.*, 2015). Given the divergence in many circadian genes' regulation, we explored
the landscape of introgression on circadian genes. We first looked at introgressed circadian
variants that are likely to influence gene regulation in AMH. Variants in this set are observed
more often than expected, suggesting the importance of maintaining circadian variation in the
population. We also verified that these results held over variants identified by different methods
for calling archaic introgression.
We then evaluated the association of these introgressed variants with variation in
circadian phenotypes of Eurasians. We previously reported a modest enrichment among
introgressed variants for heritability of the morning/evening person phenotype (McArthur,
Rinker and Capra, 2021). Here, we further discovered a consistent directional effect of the
introgressed circadian variants on chronotype. The strongest associated variants increase the
probability of being a morning person in Eurasians.
While it is not immediately clear why increased morningness would be beneficial at
higher latitudes, considering this directional effect in the context of clock gene regulation and the
challenge of adaptation to higher latitudes suggests an answer. In present day humans, behavioral
morningness is correlated with shortened period of the circadian molecular clockworks in
individuals. This earlier alignment of sleep/wake with external timing cues is a consequence of a
quickened pace of the circadian gene network (Brown *et al.*, 2008). Therefore, the morningness
directionality of introgressed circadian variants may indicate selection toward shortened
circadian period in the archaic populations living at high latitudes. Supporting this interpretation,
shortened circadian periods are required for synchronization to the extended summer
photoperiods of high latitudes in *Drosophila*, and selection for shorter periods has resulted in
latitudinal clines of decreasing period with increasing latitude, as well as earlier alignment of
behavioral rhythms (Hut *et al.*, 2013). In addition, *Drosophila* populations exhibit decreased
amplitude of behavioral rhythms at higher latitudes which is also thought to aid in
synchronization to long photoperiods (Hut *et al.*, 2013).
Our finding that introgressed circadian variants generally decrease gene regulation of
circadian genes suggests that they could lead to lower amplitude clock gene oscillations.
However, when assayed in present day humans there is not a strong correlation between the
overall expression level of *NR1D1* and the transcriptional amplitudes of other clock genes within
individuals (Brown *et al.*, 2008), and quantitative modeling of the mammalian circadian
clockworks suggests that stable clock gene rhythms can result across a wide range of absolute
levels of gene expression as long as the stoichiometric ratios of key positive and negative clock
genes are reasonably conserved (Kim and Forger, 2012). Interestingly, lower transcriptional
amplitude of *NR1D1* does confer greater sensitivity of the present-day human clockworks to
resetting stimuli, a potentially adaptive characteristic for high latitudes (Brown *et al.*, 2008).
Thus, given the studies of latitudinal clines and adaptation from *Drosophila* and the
nascent understanding of clock gene contributions to behavioral phenotypes in present day
humans, the directional effects of introgressed circadian gene variants toward early chronotype
and decreased gene regulation we observed can be viewed as potentially adaptive. More
complex chronotype phenotyping and mechanistic studies of the variants of interest are needed
to fully understand these observations.
Finally, to explore evidence for positive selection on introgressed variants in AMH, we
analyzed results from three recent methods for detecting adaptive introgression. All methods
identified circadian loci as candidates for adaptive introgression. However, we note that the
predictions of these methods have only modest overlap with one another, underscoring the

difficulty of identifying adaptive introgression. Nonetheless, many of these loci, especially those
supported by both Racimo and *MaLAdapt*, are good candidates for adaptive introgression given
their functional associations with circadian genes
Several limitations must be considered when interpreting our results. First, it is
challenging to quantify the complexity of traits with a large behavioral component (like
chronotype) and infer their variation from genomic information alone. Nevertheless, we believe
our approach of focusing on molecular aspects (splicing, gene regulation) of genomic loci with
relevance to circadian biology, in parallel to GWAS-based associations, lends additional support
to the divergence in chronotype between archaic hominins and modern humans. Second, we also
note that circadian rhythms contribute to many biological systems, so the variants in these genes
tend to be associated with a variety of phenotypes. Thus, there is also the potential that selection
acted on other phenotypes influenced by circadian variation than those related directly to
chronotype. Third, given the complexity of circadian biology, there is no gold standard set of
circadian genes. We focus on the core clock genes and a broader set of expert-curated genes
relevant to circadian systems, but it is certainly possible that other genes with circadian effects
are not considered. Fourth, recent adaptive evolution is challenging to identify, and this is
especially challenging for introgressed loci. Nonetheless, we find several circadian loci with
evidence of adaptive introgression from more than one method. Finally, given the many
environmental factors that differed between African and non-African environments, it is difficult
to definitively determine whether selection on a particular locus was the result of variation in
light levels vs. other related factors, such as temperature. Nonetheless, given the observed
modern associations with chronotype for many of these variants, we believe it is a plausible
target.
In conclusion, studying how humans evolved in the face of changing environmental
pressures is necessary to understanding variation in present-day phenotypes and the potential
tradeoffs that influence propensity to different diseases in modern environments (Benton *et al.*,
2021). Here, we show that genomic regions involved in circadian biology exhibited substantial
functional divergence between separate hominin populations. Furthermore, we show that
introgressed variants contribute to variation in AMH circadian phenotypes today in ways that are
consistent with an adaptive benefit.
588
589
590
591 **METHODS**
592
Circadian gene selection
Circadian biology is a complex system due to its high importance in the functioning of biological
timing in diverse biological systems. For that reason, determining which genes are crucial for
selection to environment response related to light exposure is not a straight forward process. To
address this issue, we look at different sources of genome annotation databases and searched for
genes and variants associated with circadian related phenotypes. We considered all human
protein-coding genes in the Gene Ontology database annotated with the GO:0007623 ("circadian
rhythm") term or terms annotated with relationship "is_a", "part_of", "occurs_in", or "regulates"
circadian rhythm. We also considered genes containing experimental or orthologous evidence of
circadian function in the Circadian Gene Database (CGDB), the GWAS Catalog genes
containing "chronotype" or "circadian rhythm" associated variants, and a curated set of genes

available in WikiPathways [https://www.wikipathways.org/index.php/Pathway:WP3594,
https://doi.org/10.1093/nar/gkaa1024]. The final set of circadian genes was curated by Dr.
Douglas McMahon.
607   To select the candidate circadian genes with the highest confidence, we defined a
hierarchy system where genes annotated by McMahon or annotated in 3 out of 4 other sources
receive a "High" level of confidence. Genes with evidence from 2 out of 4 of the sources are
assigned a "Medium" level of confidence. Genes annotated as circadian only in 1 out of 4
sources are assigned to Low confidence and not considered in our circadian gene set. We then
defined our set of circadian variants from the 1000 Genomes Project using the official list of
circadian genes. The variants are included in analysis of coding, non-coding, regulatory, eQTL,
human-specific, archaic-specific, and introgressed variants.
615
Definition of lineage-specific variants
To identify candidate variants that are specific to the human and the archaic lineages, we used a
set of variants published by Kuhlwilm and Boeckx (Kuhlwilm and Boeckx, 2019)
(https://doi.org/10.1038/s41598-019-44877-x). The variants were extracted from the high-
coverage genomes of three archaics: a 122,000-year-old Neanderthal from the Altai Mountains
(52x coverage), a 52,000-year-old Neanderthal from Vindija in Croatia (30x coverage), and a
72,000-year-old Denisovan from the Altai Mountains (30x coverage). The variants were called in
the context of the human genome hg19/GRCh37 reference. The total number of variant sites
after applying filters for high coverage sites and genotype quality is 4,437,803. A human-specific
variant is defined as a position where all the humans in the 1000 Genomes Project carry the
derived allele and all the archaics carry the ancestral allele. An archaic-specific is defined as a
position where all the archaics carry the derived allele and the derived allele is absent or
extremely rare ($<= 0.00001$) across all human populations. Note that introgressed archaic alleles
are not included in the "archaic-specific" set. These criteria resulted in 9,424 human specific and
33,184 archaic-specific variants.
631
Enrichment of lineage-specific variants among functional regions of the genome
We intersected the sets of lineage-specific variants with several sets of annotated functional
genomic regions. Inside circadian gene regions (Gencode v29), we found 156 human-specific
variants and 341 archaic-specific variants. In circadian promoter regions, we found 6 human-
specific variants and 19 archaic-specific variants. Promoters were defined as regions 5 kb up- to
1 kb downstream from a transcription start site. In distal regulatory elements, we found 247
human-specific variants and 807 archaic-specific variants. For this last set, we considered
candidate cis-regulatory elements (cCREs) published by ENCODE (Moore *et al.*, 2020) within 1
Mb of the circadian genes.
641   To compute whether lineage-specific variants are more abundant than expected in
circadian genes, we applied a Fisher's exact test to the sets of human- and archaic-specific
variants in regulatory, promoter, and gene regions. Human and archaic-specific variants are
significantly enriched in both regulatory (Human: OR=1.25, P=8.39e-4; Archaic: OR=1.16,
P=6.15e-5) and gene (Human: OR=1.84, P=7.06e-12; Archaic: OR=1.13, P=0.023) regions. The
enrichment observed in the promoters of both lineages is not supported by a significant p-value
(Human: OR=1.21, P=0.65; Archaic: OR=1.09, P=0.63).
648
Genes containing archaic variants with evidence of alternative splicing

We used a set of archaic variants annotated with the splice altering probabilities to identify
circadian genes that may be differentially spliced between archaic hominins and AMH (Brand,
Colbran and Capra, 2023). We considered variants from four archaic individuals: the Altai,
Chagyrskaya, and Vindija Neanderthals and the Altai Denisovan. These archaic variants were
annotated using SpliceAI (Jaganathan *et al.*, 2019) and we considered any variant with a
maximum delta, or splice altering probability, > 0.2. We identified 36 archaic-specific splice
altering variants, defined as those variants absent from 1000 Genomes Project, among 28
circadian genes. Next, we tested for enrichment among this gene set using an empirical null
approach (McArthur *et al.*, 2022; Brand, Colbran and Capra, 2023). We shuffled the maximum
deltas among 1,607,350 variants 10,000 times and counted the number of circadian genes with a
splice altering variant each iteration. Enrichment was calculated as the number of observed genes
(N = 28) divided by the mean gene count among 10,000 shuffles. In addition to all genes with
archaic-specific variants, we considered six other subsets among these variants: 1) genes with
variants private to the Altai Neanderthal, 2) genes with variants private to the Chagyrskaya
Neanderthal, 3) genes with variants private to the Altai Denisovan, 4) genes with variants private
to all Neanderthals, 5) genes with variants shared among all archaic individuals, and 6) genes
with variants private to the Vindija Neanderthal. Finally, we considered a subset of splice
altering variants that were identified as tag SNPs by Vernot et al. (Vernot *et al.*, 2016).
668
PrediXcan
To understand the difference in circadian biology between present-day humans and archaic
hominins, we analyzed predictions on gene regulation. We considered the results from
PrediXcan gene regulation predictions across 44 tissues from the PredictDB Data Repository
(http://predictdb.org/). The models were trained on GTEx V6 using variants identified in 2,504
present-day humans in the 1000 Genomes Project phase 3 within 1 Mb of each circadian gene.
The original analysis includes predictions for 17,748 genes for which the models explained a
significant amount of variance in gene expression in each tissue (FDR < 0.05). The prediction
models were also applied to the Altai and Vindija Neanderthals and the Denisovan. The resulting
predictions are normalized values of the distribution observed in GTEx individuals used to train
the original prediction models. Each prediction contains an empirical P-value which was
calculated for each gene and tissue pair to define genes that are divergently regulated between
archaic hominins and humans. The P-value is obtained by calculating the proportion of humans
from the 1000 Genomes Project that have predictions more extreme compared to the human
median than the archaic individual. Significantly DR genes are defined as those where the
archaic prediction falls outside the distribution of humans in the 1000 Genomes Project
predictions.
We tested whether the circadian genes in our set are more likely to be DR compared to an
empirical null distribution from random gene sets of the same size. We account for the fact that
some genes are modeled in more tissues than others by matching the distribution of tissues in
which each gene could be modeled in the random sets to our set. Among 1,467 DR genes in the
Altai Neanderthal we find 23 DR circadian genes out of the total 236 genes in the circadian set.
We iterate through the permutation analysis 1,000,000 times and find an enrichment of 1.21
(P=0.19). A similar analysis is done in the Vindija Neanderthal (1,536 total DR, 21 circadian
DR, enrichment of 1.05, P=0.43) and the Denisovan individual (1,214 total DR, 19 circadian DR,
enrichment of 1.20, P=0.24). In this study, we define a set of DR genes as the intersection
between DR genes in all three archaics, resulting in a set of 16 genes.

696
Enrichment of introgressed variants in eQTL
We performed an enrichment analysis using Pearson's chi-squared test to evaluate if there is overrepresentation of introgressed alleles in our set of circadian variants using the GTEx dataset. We did a liftOver of the GTEx v8 dataset from hg38 to hg19. The original hg38 set contains 4,631,659 eQTLs across 49 tissues. After the LiftOver, 4,608,446 eQTLs remained, with the rest not mapping. We used the archaic introgressed variants dataset from Browning 2018. The set contains 863,539 variants that are introgressed in humans originating in archaic hominins. We performed an intersection between the set of genes containing evidence for eQTLs and our set of 246 circadian genes to retrieve a subset of variant sites with evidence of being eQTL in circadian genes. The resulting subset contained 97,441 circadian eQTLs in 49 tissues and 239 genes. We further intersected the introgressed variants and the set of eQTL, resulting in 128,138 introgressed eQTLs. The final set of eQTLs that are circadian and also introgressed is 3,857.

Direction of effect of chronotype associations
To explore the effect of archaic introgression in circadian dreams on human chronotype, we quantified the direction of effect of variants associated to a Morning/Evening person trait in a GWAS analysis of the UK Biobank (http://www.nealelab.is/uk-biobank/). The variants were LD clumped using PLINK v1.9 (R^2>0.5). We generated cumulative proportion values on the beta values assigned to each associated variant on an ascending order of P-values.

Detection of latitudinal clines in chronotype associations
To evaluate latitudinal clines in chronotype-associated variants, we assigned a latitude to each of the Eurasian 1000 Genomes Project populations. The latitude of diaspora populations was set to their ancestral country (GIH Gandhinagar in Gujarat: 23.223, STU Sri Jayawardenepura Kotte: 6.916667, ITU Amaravati in Andhra Pradesh: 16.5131, CEU: 52.372778). CEU was assigned a latitude in Amsterdam, following an analysis that shows that this group is more closely related to Dutch individuals (Lao *et al.*, 2008). We then used the LDlink API to retrieve allele frequencies for each introgressed morningness variant in Eurasian individuals (Machiela and Chanock, 2015). Variants that follow a latitudinal cline were identified using linear regression statistics requiring correlation coefficient (R >= 0.65) and P-value (P <= 0.5).

Detection of pleiotropy in the set of introgressed circadian variants
To understand the extent of different phenotypes associated with the introgressed circadian variants, we first extracted genome-wide associations from Open Targets Genetics (https://genetics.opentargets.org/) for each of the variants with evidence of introgression (Browning *et al.*, 2018). Only the variants with significant p-values were analyzed. The p-value threshold was set at the genome-wide significance level (P=5e-8). We split the variants in two sets: introgressed circadian and introgressed non-circadian. Many of these variants are not associated with any phenotype. We performed a Fisher's exact test to analyze which of the two sets contains a higher ratio of SNPs with at least one association versus SNPs with no association. The result showed that the circadian set had a significantly higher ratio (OR=1.36, P=5e-29). Then we calculated the total of unique traits associated with each of the variants, given that the SNP has at least one association. We used a Mann-Whitney U test to understand which set is represented by a higher level of traits per SNP. The circadian set was slightly more pleiotropic, and the result is supported by a significant p-value (P=5.4e-3).

Identifying introgressed circadian variants with evidence of adaptive introgression

We sought to identify circadian variants that contain evidence of adaptive introgression (AI). To achieve this, we collected AI predictions from a method that applied various summary statistics on 1000 Genomes Project data (Racimo, Marnetto and Huerta-Sánchez, 2017) and two sets of genomic regions that were measured for their likelihood to be under AI by two machine learning methods: *genomatnn* and *MaLAdapt*. *genomatnn* is a convolutional neural network trained to identify adaptive introgression based on simulations (Gower *et al.*, 2021). *MaLAdapt* is a machine learning algorithm trained to find adaptive introgression based on simulations using an extra-trees classifier (ETC) (Zhang *et al.*, 2023). Following the thresholds used in each paper, a region is considered to be under AI if the prediction value assigned to it meets a threshold of 0.5 or 0.9, respectively. To find the variants of interest that fall into AI regions, we intersected the set of introgressed circadian SNPs with the Racimo et al. 2015, *genomatnn* and the *MaLAdapt* regions individually. The set of introgressed circadian variants contains variants inside circadian genes, in circadian promoter regions (5 kb up- and 1 kb downstream of the TSS), and variants with regulatory function (cCREs) flanking circadian genes by 1 Mb.

DATA AVAILABILITY
The data underlying this article are available in the article and in its online supplementary material.

DECLARATION OF INTERESTS
The authors declare that they have no competing interests.

ACKNOWLEDGMENTS
We thank members of the Capra Lab for helpful comments on this work. This work was conducted in part using the resources of the Advanced Computing Center for Research and Education at Vanderbilt University, Nashville, TN. This work was supported by the National Institutes of Health [R35GM127087 to JAC, R01GM117650 to DM, F30HG011200 to EM, T32GM080178 to Vanderbilt University (EM), and T32HG009495 to the University of Pennsylvania (LLC)].

AUTHOR CONTRIBUTIONS
Conceptualization: KV, JAC; Methodology: KV, LC, EM, CB, DR, JS, DM, JAC; Investigation: KV, LC, EM, CB, JAC; Writing – Original Draft: KV, JAC; Writing – Review & Editing: KV, LC, EM, CB, DR, DM, JAC; Funding Acquisition: JAC; Resources: JAC; Supervision: JAC.

REFERENCES
1000 Genomes Project Consortium (2010) 'A map of human genome variation from population-scale sequencing', *Nature*, 467(7319), p. 1061. Available at: https://doi.org/10.1038/nature09534.

Arnold, L.J. *et al.* (2014) 'Luminescence dating and palaeomagnetic age constraint on hominins
from Sima de los Huesos, Atapuerca, Spain', *Journal of Human Evolution*, 67(1), pp. 85–107.
Available at: https://doi.org/10.1016/j.jhevol.2013.12.001.
Bae, C.J., Douka, K. and Petraglia, M.D. (2017) 'On the origin of modern humans: Asian
perspectives', *Science*, 358(6368), p. eaai9067. Available at:
https://doi.org/10.1126/science.aai9067.
Benton, M.L. *et al.* (2021) 'The influence of evolutionary history on human health and disease',
Nature Reviews Genetics. Nature Research, pp. 269–283. Available at:
https://doi.org/10.1038/s41576-020-00305-9.
Brand, C.M., Colbran, L.L. and Capra, J.A. (2023) 'Resurrecting the alternative splicing
landscape of archaic hominins using machine learning', *Nature Ecology and Evolution*, 7(6), pp.
939–953. Available at: https://doi.org/10.1038/s41559-023-02053-5.
Brown, S.A. *et al.* (2008) 'Molecular insights into human daily behavior', *Proceedings of the
National Academy of Sciences of the United States of America*, 105(5), pp. 1602–1607. Available
at: https://doi.org/10.1073/pnas.0707772105.
Browning, S.R. *et al.* (2018) 'Analysis of Human Sequence Data Reveals Two Pulses of Archaic
Denisovan Admixture', *Cell*, 173(1), pp. 53–61. Available at:
https://doi.org/https://doi.org/10.1016/j.cell.2018.02.031.
Brzezinski, J.A. *et al.* (2005) 'Loss of circadian photoentrainment and abnormal retinal
electrophysiology in Math5 mutant mice', *Investigative Ophthalmology and Visual Science*,
46(7), pp. 2540–2551. Available at: https://doi.org/10.1167/iovs.04-1123.
Colbran, L.L. *et al.* (2019) 'Inferred divergent gene regulation in archaic hominins reveals
potential phenotypic differences', *Nature Ecology & Evolution*, 3(November), pp. 1598–1606.
Available at: https://doi.org/10.1038/s41559-019-0996-x.
Dannemann, M. and Kelso, J. (2017) 'The Contribution of Neanderthals to Phenotypic Variation
in Modern Humans', *American Journal of Human Genetics*, 101(4), pp. 578–589. Available at:
https://doi.org/10.1016/j.ajhg.2017.09.010.
Dorokhov, V.B. *et al.* (2018) 'An hour in the morning is worth two in the evening: association of
morning component of morningness – eveningness with single nucleotide polymorphisms in
circadian clock genes', *Biological Rhythm Research*, 49(4), pp. 622–642. Available at:
https://doi.org/10.1080/09291016.2017.1390823.
Gan, Y. *et al.* (2015) 'Shift work and diabetes mellitus: A meta-analysis of observational
studies', *Occupational and Environmental Medicine*, pp. 72–78. Available at:
https://doi.org/10.1136/oemed-2014-102150.
Gan, Y. *et al.* (2018) 'Association between shift work and risk of prostate cancer: A systematic
review and meta-analysis of observational studies', *Carcinogenesis*, pp. 87–97. Available at:
https://doi.org/10.1093/carcin/bgx129.
Gómez-Robles, A. (2019) 'Dental evolutionary rates and its implications for the Neanderthal–
modern human divergence', *Science Advances*, 5(5), p. eaaw1268. Available at:
https://doi.org/10.1126/sciadv.aaw1268.
Gower, G. *et al.* (2021) 'Detecting adaptive introgression in human evolution using
convolutional neural networks', *eLife*, 10, p. e64669. Available at:
https://doi.org/10.7554/eLife.64669.
Green, R.E. *et al.* (2010) 'A Draft Sequence of the Neandertal Genome', *Science*, 328(5979), pp.
710–722. Available at: https://doi.org/10.1126/science.1188021.

Grimaldi, B. *et al.* (2010) 'PER2 controls lipid metabolism by direct regulation of PPARγ', *Cell Metabolism*, 12(5), pp. 509–520. Available at: https://doi.org/10.1016/j.cmet.2010.10.005.

Gyarmati, G. *et al.* (2016) 'Night shift work and stomach cancer risk in the MCC-Spain study', *Occupational and Environmental Medicine*, 73(8), pp. 520–527. Available at: https://doi.org/10.1136/oemed-2016-103597.

Hublin, J.J. *et al.* (2017) 'New fossils from Jebel Irhoud, Morocco and the pan-African origin of Homo sapiens', *Nature*, 546(7657), pp. 289–292. Available at: https://doi.org/10.1038/nature22336.

Huerta-Sánchez, E. *et al.* (2014) 'Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA', *Nature*, 512(7513), pp. 194–197. Available at: https://doi.org/10.1038/nature13408.

Hut, R.A. *et al.* (2013) 'Latitudinal clines: An evolutionary view on biological rhythms', *Proceedings of the Royal Society B: Biological Sciences*, 280(1765). Available at: https://doi.org/10.1098/rspb.2013.0433.

Jacobs, G.S. *et al.* (2019) 'Multiple Deeply Divergent Denisovan Ancestries in Papuans', *Cell*, 177. Available at: https://doi.org/10.1016/j.cell.2019.02.035.

Jaganathan, K. *et al.* (2019) 'Predicting Splicing from Primary Sequence with Deep Learning', *Cell*, 176(3), pp. 535-548.e24. Available at: https://doi.org/10.1016/j.cell.2018.12.015.

Jones, S.E. *et al.* (2016) 'Genome-Wide Association Analyses in 128,266 Individuals Identifies New Morningness and Sleep Duration Loci', *PLoS Genetics*, 12(6), p. e1006125. Available at: https://doi.org/10.1371/journal.pgen.1006125.

Jones, S.E. *et al.* (2019) 'Genome-wide association analyses of chronotype in 697,828 individuals provides insights into circadian rhythms', *Nature Communications*, 10(1), p. 343. Available at: https://doi.org/10.1038/s41467-018-08259-7.

Kim, J.K. and Forger, D.B. (2012) 'A mechanism for robust circadian timekeeping via stoichiometric balance', *Molecular Systems Biology*, 8(1). Available at: https://doi.org/10.1038/msb.2012.62.

Kivelä, L., Papadopoulos, M.R. and Antypa, N. (2018) 'Chronotype and Psychiatric Disorders', *Current Sleep Medicine Reports*, 4(2), pp. 94–103. Available at: https://doi.org/10.1007/s40675-018-0113-8.

Knutson, K.L. and von Schantz, M. (2018) 'Associations between chronotype, morbidity and mortality in the UK Biobank cohort', *Chronobiology International*, 35(8), pp. 1045–1053. Available at: https://doi.org/10.1080/07420528.2018.1454458.

Kuhlwilm, M. and Boeckx, C. (2019) 'A catalog of single nucleotide changes distinguishing modern humans from archaic hominins', *Nature Scientific Reports*, 9(8463). Available at: https://doi.org/10.1038/s41598-019-44877-x.

Lane, J.M. *et al.* (2016) 'Genome-wide association analysis identifies novel loci for chronotype in 100,420 individuals from the UK Biobank', *Nature Communications*, 7(1), pp. 1–10. Available at: https://doi.org/10.1038/ncomms10889.

Lao, O. *et al.* (2008) 'Correlation between Genetic and Geographic Structure in Europe', *Current Biology*, 18(16), pp. 1241–1248. Available at: https://doi.org/10.1016/j.cub.2008.07.049.

Larcher, S. *et al.* (2015) 'Sleep habits and diabetes', *Diabetes and Metabolism*, pp. 263–271. Available at: https://doi.org/10.1016/j.diabet.2014.12.004.

Larcher, S. *et al.* (2016) 'Impact of sleep behavior on glycemic control in type 1 diabetes: The role of social jetlag', *European Journal of Endocrinology*, 175(5), pp. 411–419. Available at: https://doi.org/10.1530/EJE-16-0188.

Lee, Y.J. *et al.* (2012) 'Circadian regulation of low density lipoprotein receptor promoter activity
by CLOCK/BMAL1, Hes1 and Hes6', *Experimental and Molecular Medicine*, 44(11), pp. 642–
652. Available at: https://doi.org/10.3858/emm.2012.44.11.073.
Leocadio-Miguel, M.A. *et al.* (2017) 'Latitudinal cline of chronotype', *Scientific Reports*,
7(5437), pp. 2–7. Available at: https://doi.org/10.1038/s41598-017-05797-w.
Lowden, A. *et al.* (2018) 'Delayed Sleep in Winter Related to Natural Daylight Exposure among
Arctic Day Workers', *Clocks & Sleep*, 1(1), pp. 105–116. Available at:
https://doi.org/10.3390/clockssleep1010010.
Machiela, M.J. and Chanock, S.J. (2015) 'LDlink: A web-based application for exploring
population-specific haplotype structure and linking correlated alleles of possible functional
variants', *Bioinformatics*, 31(21), pp. 3555–3557. Available at:
https://doi.org/10.1093/bioinformatics/btv402.
Mafessoni, F. *et al.* (2020) 'A high-coverage Neandertal genome from Chagyrskaya Cave',
Proceedings of the National Academy of Sciences of the United States of America, 117(26), pp.
15132–15136. Available at: https://doi.org/10.1073/pnas.2004944117.
McArthur, E. *et al.* (2022) 'Reconstructing the 3D genome organization of Neanderthals reveals
that chromatin folding shaped phenotypic and sequence divergence', *bioRxiv* [Preprint].
McArthur, E., Rinker, D.C. and Capra, J.A. (2021) 'Quantifying the contribution of Neanderthal
introgression to the heritability of complex traits', *Nature Communications*, 12(1), pp. 1–14.
Available at: https://doi.org/10.1038/s41467-021-24582-y.
Meyer, M. *et al.* (2012) 'A high-coverage genome sequence from an archaic Denisovan
individual', *Science*, 338(6104), pp. 222–226. Available at:
https://doi.org/10.1126/science.1224344.
Meyer, M. *et al.* (2014) 'A mitochondrial genome sequence of a hominin from Sima de los
Huesos', *Nature*, 505(7483). Available at: https://doi.org/10.1038/nature12788.
Meyer, M. *et al.* (2016) 'Nuclear DNA sequences from the Middle Pleistocene Sima de los
Huesos hominins', *Nature*, 531(7595), pp. 504–507. Available at:
https://doi.org/10.1038/nature17405.
Michael, T.P. *et al.* (2003) 'Enhanced Fitness Conferred by Naturally Occurring Variation in the
Circadian Clock', *Science*, 302(5647), pp. 1049–1053. Available at:
https://doi.org/10.1126/science.1082971.
Moore, J.E. *et al.* (2020) 'Expanded encyclopaedias of DNA elements in the human and mouse
genomes', *Nature*, 583(7818), pp. 699–710. Available at: https://doi.org/10.1038/s41586-020-
2493-4.
Nielsen, R. *et al.* (2017) 'Tracing the peopling of the world through genomics', *Nature*,
541(7637), pp. 302–310. Available at: https://doi.org/10.1038/nature21347.
O'Malley, K.G. and Banks, M.A. (2008) 'A latitudinal cline in the Chinook salmon
(Oncorhynchus tshawytscha) Clock gene: Evidence for selection on PolyQ length variants',
Proceedings of the Royal Society B: Biological Sciences, 275(1653), pp. 2813–2821. Available
at: https://doi.org/10.1098/rspb.2008.0524.
O'Malley, K.G., Ford, M.J. and Hard, J.J. (2010) 'Clock polymorphism in Pacific salmon:
Evidence for variable selection along a latitudinal gradient', in *Proceedings of the Royal Society
B: Biological Sciences*. Royal Society, pp. 3703–3714. Available at:
https://doi.org/10.1098/rspb.2010.0762.

Papantoniou, K. *et al.* (2016) 'Breast cancer risk and night shift work in a case–control study in a
Spanish population', *European Journal of Epidemiology*, 31(9), pp. 867–878. Available at:
https://doi.org/10.1007/s10654-015-0073-y.
Papantoniou, K. *et al.* (2017) 'Shift work and colorectal cancer risk in the MCC-Spain case–
control study', *Scandinavian Journal of Work, Environment and Health*, 43(3), pp. 250–259.
Available at: https://doi.org/10.5271/sjweh.3626.
Petr, M. *et al.* (2019) 'Limits of long-term selection against Neandertal introgression',
Proceedings of the National Academy of Sciences, 116(5), pp. 1639–1644. Available at:
https://doi.org/10.1073/pnas.1814338116.
Prüfer, K. *et al.* (2014) 'The complete genome sequence of a Neanderthal from the Altai
Mountains', *Nature*, 505(7481), pp. 43–49. Available at: https://doi.org/10.1038/nature12886.
Prüfer, K. *et al.* (2017) 'A high-coverage Neandertal genome from Vindija Cave in Croatia',
Science, 358(6363), pp. 655–658. Available at: https://doi.org/10.1126/science.aao1887.
Putilov, A.A. *et al.* (2019) 'Genetic-based signatures of the latitudinal differences in
chronotype', *Biological Rhythm Research*, 50(2), pp. 255–271. Available at:
https://doi.org/10.1080/09291016.2018.1465249.
Putilov, A.A., Dorokhov, V.B. and Poluektov, M.G. (2018) 'How have our clocks evolved?
Adaptive and demographic history of the out-of-African dispersal told by polymorphic loci in
circadian genes', *Chronobiology International*, 35(4), pp. 511–532. Available at:
https://doi.org/10.1080/07420528.2017.1417314.
Racimo, F. *et al.* (2015) 'Evidence for archaic adaptive introgression in humans', *Nature*
Reviews Genetics, 16(6), pp. 359–371. Available at: https://doi.org/10.1038/nrg3936.
Racimo, F. *et al.* (2017) 'Archaic adaptive introgression in TBX15/WARS2', *Molecular Biology*
and Evolution, 34(3), pp. 509–524. Available at: https://doi.org/10.1093/molbev/msw283.
Racimo, F., Marnetto, D. and Huerta-Sánchez, E. (2017) 'Signatures of archaic adaptive
introgression in present-day human populations', *Molecular Biology and Evolution*, 34(2), pp.
296–317. Available at: https://doi.org/10.1093/molbev/msw216.
Randler, C. and Rahafar, A. (2017) 'Latitude affects Morningness- Eveningness: evidence for the
environment hypothesis based on a systematic review', *Nature Publishing Group*, 7(39976), pp.
1–6. Available at: https://doi.org/10.1038/srep39976.
Rinker, D.C. *et al.* (2020) 'Neanderthal introgression reintroduced functional ancestral alleles
lost in Eurasian populations', *Nature Ecology and Evolution*, 4(10), pp. 1332–1341. Available at:
https://doi.org/10.1038/s41559-020-1261-z.
Sandrelli, F. *et al.* (2007) 'A molecular basis for natural selection at the timeless locus in
Drosophila melanogaster', *Science*, 316(5833), pp. 1898–1900. Available at:
https://doi.org/10.1126/science.1138426.
Sankararaman, S. *et al.* (2012) 'The date of interbreeding between Neandertals and modern
humans', *PLoS Genetics*, 8(10), p. e1002947. Available at:
https://doi.org/10.1371/journal.pgen.1002947.
Sankararaman, S. *et al.* (2014) 'The genomic landscape of Neanderthal ancestry in present-day
humans', *Nature*, 507(7492), pp. 354–359. Available at: https://doi.org/10.1038/nature12961.
Schaefer, N.K., Shapiro, B. and Green, R.E. (2021) 'An ancestral recombination graph of
human, Neanderthal, and Denisovan genomes', *Science advances*, 7(29), pp. 776–792. Available
at: https://doi.org/10.1126/sciadv.abc077.

Shi, Y. *et al.* (2020) 'Night-shift work duration and risk of colorectal cancer according to IRS1
and IRS2 expression', *Cancer Epidemiology Biomarkers and Prevention*, 29(1), pp. 133–140.
Available at: https://doi.org/10.1158/1055-9965.EPI-19-0325.
Skoglund, P. and Mathieson, I. (2018) 'Ancient genomics of modern humans: The first decade',
Annual Review of Genomics and Human Genetics. Annual Reviews Inc., pp. 381–404. Available
at: https://doi.org/10.1146/annurev-genom-083117-021749.
Skov, L. *et al.* (2020) 'The nature of Neanderthal introgression revealed by 27,566 Icelandic
genomes', *Nature*, 582(7810), pp. 78–83. Available at: https://doi.org/10.1038/s41586-020-
2225-9.
Srinivasan, V. *et al.* (2006) 'Melatonin in mood disorders', *World Journal of Biological*
Psychiatry, pp. 138–151. Available at: https://doi.org/10.1080/15622970600571822.
Steinrücken, M. *et al.* (2018) 'Model-based detection and analysis of introgressed Neanderthal
ancestry in modern humans', *Molecular Ecology*, 27(19), pp. 3873–3888. Available at:
https://doi.org/10.1111/mec.14565.
Stringer, C. (2016) 'The origin and evolution of Homo sapiens', *Philosophical Transactions of*
the Royal Society B: Biological Sciences, 371(1698), p. 20150237. Available at:
https://doi.org/10.1098/rstb.2015.0237.
Tauber, E. *et al.* (2007) 'Natural selection favors a newly derived timeless allele in Drosophila
melanogaster', *Science*, 316(5833), pp. 1895–1898. Available at:
https://doi.org/10.1126/science.1138412.
Taylor, B.J. and Hasler, B.P. (2018) 'Chronotype and Mental Health: Recent Advances', *Current*
Psychiatry Reports, 20(8). Available at: https://doi.org/10.1007/s11920-018-0925-8.
Telis, N., Aguilar, R. and Harris, K. (2020) 'Selection against archaic hominin genetic variation
in regulatory regions', *Nature Ecology and Evolution*, 4(11), pp. 1558–1566. Available at:
https://doi.org/10.1038/s41559-020-01284-0.
Vernot, B. *et al.* (2016) 'Excavating Neandertal and Denisovan DNA from the genomes of
Melanesian individuals', *Science*, 352(6282), pp. 235–239. Available at:
https://doi.org/10.1126/science.aad9416.
Vernot, B. and Akey, J.M. (2014) 'Resurrecting Surviving Neandertal Lineages from Modern
Human Genomes', *Science*, 343(6174), pp. 1017–1021. Available at:
https://doi.org/10.1126/science.1245938.
Villanea, F.A. and Schraiber, J.G. (2019) 'Multiple episodes of interbreeding between
Neanderthal and modern humans', *Nature Ecology and Evolution*, 3(1), pp. 39–44. Available at:
https://doi.org/10.1038/s41559-018-0735-8.
Võsa, U. *et al.* (2021) 'Large-scale cis- and trans-eQTL analyses identify thousands of genetic
loci and polygenic scores that regulate blood gene expression', *Nature Genetics*, 53(9), pp.
1300–1310. Available at: https://doi.org/10.1038/s41588-021-00913-z.
Yousef, E. *et al.* (2020) 'Shift work and risk of skin cancer: A systematic review and meta-
analysis', *Scientific Reports*, 10(1), pp. 1–11. Available at: https://doi.org/10.1038/s41598-020-
59035-x.
Zhang, Q. *et al.* (2008) 'Association of the circadian rhythmic expression of GmCRY1a with a
latitudinal cline in photoperiodic flowering of soybean', *Proceedings of the National Academy of*
Sciences of the United States of America, 105(52), pp. 21028–21033. Available at:
https://doi.org/10.1073/pnas.0810585105.

Zhang, X. *et al.* (2023) 'MaLAdapt Reveals Novel Targets of Adaptive Introgression From
Neanderthals and Denisovans in Worldwide Human Populations', *Molecular Biology and*
Evolution, 40(1), p. msad001. Available at: https://doi.org/10.1093/molbev/msad001.
1014