Too disturbed or too hot: summer temperatures more than hikers disturbance affect circadian activity of females in northern chamois

Thel Lucie[1,2], Garel Mathieu[3], Marchand Pascal[4], Bourgoin Gilles[5,6], Loison Anne[1]

Authors details

[1] Laboratoire d'Ecologie Alpine, UMR 5553 (CNRS, Université Savoie Mont--Blanc), Savoie Technolac, 73376 Le Bourget-du-Lac, France.

[2] Laboratoire Évolution et Diversité Biologique, UMR 5174 (CNRS, Université Paul Sabatier), 118 route de Narbonne, 31062 Toulouse, France.

[3] Office Français de la Biodiversité, Direction de la Recherche et de l'Appui Scientifique, Service Anthropisation et Fonctionnement des Ecosystèmes Terrestres, Gières, France.

[4] Office Français de la Biodiversité, Direction de la Recherche et de l'Appui Scientifique, Service Anthropisation et Fonctionnement des Ecosystèmes Terrestres, Juvignac, France.

[5] Univ Lyon, VetAgro Sup, Campus Vétérinaire de Lyon, 1 avenue Bourgelat, F-69280 Marcy l'Etoile, France.

[6] Univ Lyon, Université Lyon 1, CNRS, UMR 5558, Laboratoire de Biométrie et Biologie Evolutive (LBBE), F-69622 Villeurbanne Cedex, France.

Correspondence author: Thel Lucie, lucie.mc.thel@gmail.com.

Abstract

Though non-lethal, recreational activities often result in a spatial and/or temporal shift of activity in a large range of species because they are perceived as a risk. With the concurrent development of outdoors activities and the increase of temperatures due to climate change in mountainous ecosystems, mountain species face an increasing pressure in terms of managing their activity pattern to both limit risk exposure and reduce thermal stress. In this context, we investigated how female northern chamois adjust their summer circadian activity to spatio-temporal variations of both temperature and hikers disturbance. We found that chamois behaviour was more affected by high temperatures than by hikers disturbance. During the hottest days, they shifted their activity peak earlier in the morning (50 minutes earlier; and to a lesser extent, later in the evening), and were more active at night (+ 21.5% *approx.*) than during daytime (- 10.10% *approx.*). Their global daily activity was higher (+ 4.54% *approx.*) during the hottest than coldest days owing to a significant increase of activity level during the activity peaks. Conversely, hikers disturbance had almost no effect on activity peaks of chamois most likely due to a temporal mismatch with hikers arrival and departure. Spatial and temporal disturbance had also a weak effect on chamois activity level, except when considered together and during the hottest days (*i.e.* the most disturbed conditions) exacerbating the temporal report of activity at night those days. Low tolerance of chamois to hikers presence might have led to the pattern reported with a similar response whatever the intensity of encounter risk. These results might also suggest that behavioural changes buffering animals from high temperature (relocation into refuge areas) and/or from hikers presence (moving away temporarily from trails) allow them to marginally modify their activity pattern. In the context of ongoing socio-environmental changes, these findings stress the importance of conserving habitats providing both shelter against summer heat and human disturbance to mitigate their potential detrimental consequences.

48   **Keywords**: circadian rhythm, ecotourism, human disturbance, mountain ecosystem,

49   nocturnality, recreational activities, *Rupicapra rupicapra rupicapra*

Introduction

A noticeable development and diversification of outdoor activities has been ongoing in the

twenty-first century, especially in Western countries (Wheaton, 2010). While mountain

ecosystems have been preserved until recently because of their low accessibility, they now

face an increase of unaffiliated activities such as hiking, cross country skiing, trail running, or

climbing (Chanteloup, Perrin-Malterre, Duparc, & Loison, 2016). Those recreational

activities are non-lethal, but humans can be consistently perceived as predators by wildlife

(Byers, 1997; Frid & Dill, 2002), leading to a suite of physiological and behavioural

responses (Arlettaz *et al.,* 2007; Marchand *et al.,* 2014; Larson, Reed, Merenlender, &

Crooks, 2016). The consequences of repeated reactions to increasing disturbance level can be

detrimental on several fitness-related components (Phillips & Alldredge, 2000; French,

González-Suárez, Young, Durham, & Gerber, 2011). This could be particularly true in

mountain ecosystems, where the diversification of human activities challenges animals all

year long, both during the period of energy acquisition (*i.e.* spring to autumn) and loss (*i.e.*

winter). This increased pressure on the animals is accompanied by environmental changes,

including climate change, that are particularly strong in the Alps (Pepin *et al.,* 2015).

Accordingly, temperatures are increasing rapidly in all seasons, with consequences such as

earlier springs and plant cycles, more frequent summer heatwaves and droughts, and

decreased duration of snow cover (associated with an augmented risk of frost for plants;

Gobiet *et al.,* 2014). Mountain animals are therefore facing concomitantly two major changes

(increased disturbance level and climate change), with immediate (*e.g.* avoidance of heat and

disturbance) and long-term (*e.g.* through food quality or changes in habitat use) consequences

on their behaviour, their physiology and their energetic balance (food intake *vs.* energy

expenditure). Indeed, energy loss can be severe and highly detrimental for high-altitude

organisms undergoing harsh climatic conditions in winter and benefiting from only short

periods with mild weather to store enough energy to survive all year round (Lindstedt &

Boyce, 1985).

Ongoing climate change and expansion of human activities in mountain ecosystems

are also ultimately connected. An extended summer season or more frequent summer

heatwaves in valleys may, for instance, exacerbate human attendance in mountains (Scott,

2006). Although predicting tourism attendance under scenarios of climate change is probably

not straightforward (Beniston, 2003; Scott, 2006), this context raises questions on animal

responses to those interrelated human-induced environmental changes and on their potential

cumulative (even multiplicative) consequences for populations in terms of both immediate

and compensatory responses. One critical challenge is to understand whether behavioural

mechanisms used by animals to buffer them against increasing temperatures may also provide

a benefit against human attendance or *vice versa*.

Wildlife is constantly subject to the foraging/risk avoidance trade-off (Lima &

Bednekoff, 1999). Animals navigate and take decisions about where to be and what to do in

multiple layers of the landscape described as their "foodscape" (the distribution of edible

food resource in a landscape), their landscape of energy (the energy to be spent to move from

one place to another in a landscape), their thermal landscape and their landscape of fear (the

distribution of the risk perception in space; Laundre, Hernández, & Altendorf, 2001; Searle,

Hobbs, & Gordon, 2007). Individuals respond to the different values of these variables in

terms of movements, habitat use and selection, and time devoted to different activities

(foraging, resting, vigilance or social/sexual interactions). When facing a disturbance

perceived as a risk, animals can modify their activities (*e.g.* from foraging to vigilance;

Benoist, Garel, Cugnasse, & Blanchard, 2013; Xu, Gong, & Wang, 2021), and move away

temporarily or even definitively (Rogala *et al.,* 2011; Courbin *et al.,* 2022). Another way to

face disturbance is to modify their activity pattern by delaying their activities, such as

foraging in disturbed places, from high- to low-risk periods (Bonnot *et al.,* 2013). The benefit

of such responses to disturbances have been explained within the "risk allocation hypothesis"

framework (Lima & Bednekoff, 1999). Indeed, the costs of lost opportunities to forage during

risky periods could be dampened if animals transfer part of their diurnal activity to night-time

when human activities usually stop (*e.g.* Marchand *et al.,* 2014, Bonnot *et al.,* 2020) or adjust

their activity hour by hour (Li *et al.,* 2009) when disturbed.

However, as night can be significantly shorter than day (*e.g.* in summer), the transfer

of the diurnal activities to night-time does not necessarily allow to fully compensate for the

loss of foraging opportunities during day and may have consequences, for instance, on mass

gain (Bourgoin *et al.*, 2011). More generally, activity patterns are precisely regulated (such as

the critical foraging/rumination balance in ruminants, Hamel & Côté, 2008), and any

modification of those patterns can negatively impact individual's fitness, *e.g.* by increasing

encounter risk with predators (Lesmerises *et al.,* 2018), by leading to mismatches in

interactions with conspecifics (Greives *et al.,* 2015), or by increasing risk of diseases

infection (Takahashi, Hong, Ko, & McDearmon, 2008). At a coarse time scale, activity

patterns are seasonally structured, driven by photoperiod, environmental conditions and

species life-cycle (Bourgoin *et al.*, 2008). At a fine pace of time, such as a day, they depend

on internal drivers (Saper, Lu, Chou, & Gooley, 2005), fine scale weather conditions (*e.g.*

wind speed, Bourgoin *et al.*, 2011) or activity patterns of other species (*e.g.* insect harassment

disrupt activity budget in reindeer *Rangifer tarandus tarandus*, Hagemoen & Reimers, 2002).

At both coarse and fine time scales, temperature is among the most important regulators of

activity patterns (*e.g.* in Alpine ibex *Capra ibex*, Semenzato *et al.,* 2021). Temperature

influences space use (in arctic ground squirrels *Spermophilus parryii*, Long, Martin, &

Barnes, 2005; in Mediterranean mouflon *Ovis gmelini musimon x ovis* sp., Marchand *et al.,*

2015) and activity budget (in Japanese monkeys *Macaca fuscata*, Watanuki & Nakayama,

1993), with animals having to trade off foraging and thermal cover to limit thermoregulatory

costs while keeping their energy balance (van Beest & Milner, 2013). For example, female

Mediterranean mouflons decrease the length of their activity boots, shift their activity peaks

(earlier in the morning and later in the evening) and increase their nocturnal activity during

the hottest summer days (Bourgoin *et al.*, 2008; Bourgoin *et al.*, 2011). These results

highlight the importance of investigating full daily cycles to get a comprehensive picture of

how animals may adjust their behaviour in the face of human-induced environmental changes

(Marchand *et al.,* 2014; Marchand *et al.,* 2015).

Here, we relied on the GPS monitoring of 62 females of northern chamois *Rupicapra*

rupicapra rupicapra to investigate at a fine scale their behavioural responses to temporal and

spatial stressors in a dynamic landscape of fear (Palmer *et al.,* 2022). More specifically, we

assessed the day-to-day variation in both temperature and hikers attendance for each

individual and also computed an individual-specific index of spatial disturbance combining

data on the attendance of each trail section by hikers, and the density of hiking trails in each

chamois summer home range. This allowed us to investigate individual behavioural

responses, using a diversity of activity metrics, to spatio-temporal variations in both

temperature and human disturbance, and their relative contributions to changes in summer

circadian activity patterns. We expected female chamois to adjust their circadian activity

pattern by avoiding being active during the hottest hours during the hottest days (**H1**) and

during the most disturbed days (**H2**), with a compensation of lost activity opportunities

during daytime at night. However, as individuals were exposed to variable levels of

disturbance depending on the location of their summer home range, we also expected more

marked responses in chamois which were the most exposed to hiker disturbance (**H3**) as most

of them show low tolerance to hikers presence (Courbin *et al.,* 2022).

Materials and methods

Study site and species

The study took place in the National Game and Hunting Reserve of Les Bauges massif

(NGHRB), in the French Alps (45°40' N, 6°13' E; 900 to 2200 m a.s.l.; 5,200 ha). The

Bauges massif is made of limestone, with cold (mean annual temperature: 9.6°C, snow cover

from November to April above 1,600 m, Duparc *et al.,* 2013) and wet (mean cumulative

annual rainfall: 1,600 mm, Duparc *et al.,* 2013) climatic conditions (see Supporting

information 1 for details about summer temperatures). Fifty-six percent of the area is covered

by forests, dominated by beech *Fagus sylvatica* and fir *Abies alba*, 36% by grasslands, and

8% by rocks (Lopez, 2001). This heterogeneous environment leads to a huge diversity of

habitats and landscapes that attracts numerous tourists every year (> 37,000, most of them

during summer). All alpine grasslands are crossed by a network of trails where nearly all

hikers are concentrated, so that this network is a major component of the landscape of fear

perceived by chamois (Courbin *et al.,* 2022). Other recreational activities (*e.g.* paragliding)

occur in the study site during summer, but far less frequently than hiking.

      Most adult female chamois annually give birth to one young in late May. During

summer, they form groups with other females, kids and yearlings, that share and keep the

same home range from year to year (Loison, Jullien, & Menaut, 1999). Females play a major

role in the demography of the population (Gaillard, Festa-Bianchet, Yoccoz, Loison, &

Toigo, 2000). Chamois had no natural predators in the study area during the study period,

except golden eagles *Aquila chrysaetos* and red foxes *Vulpes vulpes* that may occasionally

predate newborns and injured animals. However, the presence of some non-resident wolves

Canis lupus has been noticed in recent years. Hunting (performed by small groups of three to

four hunters) occurs in the NGHRB from the beginning of September to the end of February

(Courbin *et al.,* 2022), with chamois being the main target species (~ 100 chamois harvested

per year on average, *i.e.* ~ 70% of the total number of ungulates harvested), in addition to

Mediterranean mouflon (~ 20%), roe deer *Capreolus capreolus* (~ 4%), wild boar *Sus scrofa*

(~ 4%) and red deer *Cervus elaphus* (~ 2%; see Courbin *et al.,* 2022 for details).

Data collection

We focused on summer, *i.e.* between 15th June and 7th September, 2004-2020, when the

hunting season had not started yet so that chamois were exclusively exposed to hikers'

disturbance. This period is critical for female chamois because it follows the birth season and

it is characterised by high energy requirements for lactation and storage of fat reserves for

winter survival (Clutton-Brock, Albon, & Guinness, 1989), so that any disturbance during

this period could be particularly detrimental. We used a subset of 62 adult female chamois >

2 years old trapped during summer using falling nets baited with salt licks located in four

alpine grasslands near two trails highly used by hikers (Armenaz and Dent des Portes). Those

females were equipped with GPS collars Lotek 3300S (Lotek Engineering Inc., Carp,

Ontario, Canada). All captures, handling and sampling were conducted according to the

appropriate national laws for animal welfare, following the ethical conditions detailed in the

specific accreditations delivered by the Préfecture de Paris (prefectorial decree n°2009–014)

in agreement with the French environmental code (Art. R421-15 to 421–31 and R422-92 to

422–94-1).

Chamois locations were recorded daily and only chamois with locations during at

least 2/3 of the study period (*i.e.* 56 days, because deployment/recovery of GPS collars

occurred during summer), for which we could reliably estimate the summer home range (95%

kernel contour using the *adehabitatHR* package; Worton, 1989; Calenge, 2006) were

included in the study. Animal motion was recorded every 5 minutes by two perpendicular

captive-balls in the GPS collars monitoring neck movements (0: no movement, 255: highly

mobile) and the proportion of time the animal was head down. We discriminated each

chamois as "active" or "inactive" for every 5-minutes boot using the discriminant model

developed by Bourgoin and colleagues (2011; see also Darmon *et al.,* 2014). We used a total

of 4,543 complete days of activity records from the 62 previously selected chamois (*n* = 137

chamois-year, 1,308,384 5-minutes activity measurements in total). The activity of each

chamois was monitored during 18 to 132 days in total, corresponding to 1 to 49 days (*median*

= 36, *SD* = 12) during a single summer period for a given chamois.

We assessed daily characteristics (see below) from 03:00 UTC of day *d* to 03:00 UTC

of day *d+1*. This allowed us to explore potential compensation processes during the night

following the day of interest. We selected the starting hour in accordance with the beginning

of the chamois activity peak during summer mornings (Darmon *et al.,* 2014). We used ERA5

reanalysis combined with a micro-climatic model (Kearney & Porter, 2017; Maclean,

Mosedale, & Bennie, 2019; Kearney, Gillingham, Bramer, Duffy, & Maclean, 2020) to

retrieve the mean daily temperatures within alpine pastures used by chamois during summer

months instead of the closest weather station located in the valley. We explored the effect of

temperature on female chamois activity pattern through a categorical variable with two levels

defined according to the quantiles of the mean daily temperature distribution. Days with

mean temperature ≤ 25% quantile (*i.e.* 10.7°C) were considered as **cold days** and days with

mean temperature > 75% quantile (*i.e.* 15.5°C) were considered as **hot days**. Cold days

experienced by chamois ranged between 3.4°C and 10.7°C (*median* = 8.9°C, *SD* = 1.6°C),

while hot days ranged between 15.5°C and 21.0°C (*median* = 17.1°C, *SD* = 1.2°C, see also

Supporting information 1).

Temporal disturbance associated with hiking practice was defined as a categorical

variable with two levels. Week-ends and public holidays (*i.e.* vacation days such as

Ascension Day) were considered as **highly disturbed days**, whereas days between Monday

and Friday (*i.e.* working days, including weekdays during school holidays) were considered

as **weakly disturbed days**. To check for the reliability of these two categories in the field, we

compared the daily attendance of hikers (number of hikers recorded on the trail) between

03:00 UTC and 23:00 UTC and the hour of first arrival for both temporal disturbance levels

using an eco-counter (PYRO sensor Eco-Compteur, www.eco-compteur.com; 90 cm above

ground) located on Armenaz trail (45°37'18.60" N, 6°13'2.53" E), on a subset of the study

period (2016 and 2017 summers). Our study period encompassed 193 theoretically highly

disturbed days and 442 weakly disturbed days. Hikers were almost two times more on the

trails during highly (*median* = 73 hikers, *SD* = 53 hikers) than during weakly (*median* = 40

hikers, *SD* = 34 hikers) disturbed days (*Mann-Whitney test statistic* = 2,565, *p-value* = 0.005).

Hikers also tended to arrive one hour earlier during highly (*median* = 07:00 UTC, *SD* = 2

hours) compared to weakly (*median* = 08:00 UTC, *SD* = 3 hours) disturbed days although

this difference was not statistically significant (*Mann-Whitney test statistic* = 1,447, *p-value* =

0.071).

Potential spatial exposure to hikers disturbance was evaluated within each female

chamois home range using frequentation rates of the trails extracted from Strava Global

Heatmap (Strava, 2022; www.strava.com, download date: 2022/03/09, data from the two

previous years, see also Courbin *et al.,* 2022 for a similar approach on this study site). Strava

heatmap provides a value ranging from 0 for no attendance to 255 for the highest attendance

for each pixel of the map and is a good proxy of relative human frequentation (Supporting

information 2, see also Thorsen *et al.,* 2022). The lowest value recorded for a trail pixel in

our study site was 51, so we defined frequentation classes according to the quantiles of the

[51; 255] distribution (*i.e.* covering trail pixels exclusively): 1 = [51; 119] (between 0% and

33% quantile, low frequentation); 2 =]119; 187] (between 33% and 66% quantiles,

intermediate frequentation); 3 =]187; 255] (between 66% and 100% quantiles, high

frequentation). We counted the number of trail pixels of each class in a given home range,

and multiplied this number by the corresponding class value (*i.e.* 1, 2 or 3) to give a different

weight to each trail pixel according to its frequentation level. We then summed these values

for a given home range and normalised the total by dividing it by the home range surface to

obtain our spatial disturbance index. We used the median spatial disturbance index (*i.e.*

0.00577) as a threshold to discriminate between **highly exposed chamois** and **weakly**

exposed chamois.

As our study period ran over almost three months, day length varied widely and is

known to influence chamois activity (Fattorini *et al.,* 2019). We evaluated day length as the

time elapsed between dawn and dusk using the *crepuscule* function from the *maptools*

package (Bivand & Lewin-Koh, 2022; reference angle of the sun below the horizon = 6°, *i.e.*

civil dawn/dusk).

Statistical analyses: predicting activity patterns and calculating activity metrics

To account for the effect of day length and inter-individual variations on circadian activity

patterns (see Supporting information 3), we first modelled the daily activity of chamois as a

series of values of either 0 (no activity) or 1 (highly active) for each 5-minutes boot. We used

a generalised additive mixed model with a binomial distribution of errors (*mgcv* package,

Wood, 2011; Wood, Goude, & Shaw 2015; Wood, 2017). We included three fixed effects in

the model: 1) *time of the day* (*i.e.* exact time every 5 minutes) and 2) *day length* (and their

interaction) as continuous variables with a cyclic cubic regression splines as smoothing basis;

3) a synthesis 8-levels categorical variable representing all possible combinations between

temperature classes, levels of temporal disturbance and of spatial disturbance, in interaction

with the *time of the day*. The levels were ordered, with the reference level corresponding to a

chamois living in a weakly exposed area during a cold and weakly disturbed day. We defined

the unique annual id-year ID of each chamois as a random effect. We set the maximum

number of knots for the smoothing functions to 24. We also fitted the null model, which was

similar to the model described above but excluded the *temperature-disturbance* variable, for

comparison purposes.

We used the predictions of the model to investigate the effect of temperature and

hikers disturbance on several activity metrics derived from this model (described below). We

controlled for the effect of *day length* by fixing it to the median of the complete dataset (*i.e.*

963 minutes) when predicting chamois activity for each level of the *temperature-disturbance*

combination. We derived the *diurnal*, *nocturnal* and *total activity* as the area under the

activity curve for each level of the *temperature-disturbance* variable. We defined daytime as

the period elapsed between two hours after dawn and two hours before dusk, and night-time

as the period elapsed between two hours after dusk and two hours before dawn (see Bonnot *et*

al., 2020 for a similar approach). We assessed the *timing of the morning* and *evening activity*

peak (time when the highest activity level occurs in the morning and the evening

respectively). We calculated confidence intervals around these values using bootstraps

method (Supporting information 4). We then calculated the deviation from the reference level

by subtracting the value for the reference level to the value for the level of interest (*i.e.* each

level of the *temperature-disturbance* variable) for each activity metric. The percentage of

variation between two situations was calculated as *(value$_{end}$ - value$_{strat}$)/value$_{start}$*100*. We

performed all analyses using the R software (R Core Development Team 2022).

Results

Model adjustment

The full generalised additive mixed model (*AIC* = 1,606,766, *explained deviance* = 11.4%, *adjusted R2* = 0.149) was better than the null model (including *time of the day* and *day length* effects and excluding the *temperature-disturbance* variable; *AIC* = 1,619,884, *explained deviance* = 10.7%, *adjusted R2* = 0.140) to predict activity pattern (Fig. 1). All smooth terms (fixed and random effects) were significant (*p-values* < 0.001).

Temperature

Whatever the level of spatial exposure and temporal disturbance, chamois always significantly increased their total activity during hot days compared to cold days of about 0.5 unit (area under curve, see Table 1, see also Fig. 2.a). This change corresponds to a significant increase of *approx.* 0.2 unit (more than 21% variation) in the nocturnal activity and decrease of *approx.* 0.6 unit (more than 10% variation) in the diurnal activity, from cold to hot days (Table 1, Fig. 2.b and 2.c). Chamois also significantly advanced their morning peak, *i.e.* by slightly less than 1h from cold to hot days (*Timing*$_{morn. peak cold}$ ranging between 05:40 [05:15; 05:40] UTC and 05:55 [05:30; 05:55] UTC, *Timingm*$_{orn. peak hot}$ ranging between 04:45 [04:40; 05:10] UTC and 04:55 [04:45; 05:10] UTC), but they only slightly delayed their evening peak (difference not significant for chamois living in a weakly exposed area during weakly disturbed days, Table 2, Fig. 3). Both morning and evening peaks were more intense (*i.e.* higher activity level) of about 14% during hot compared to cold days (Fig. 1). Chamois maintained a certain amount of activity in the middle of the day during cold days

(*i.e.* slightly above 40%), whereas the activity level at the same hours was lower and more

variable during hot days (Fig. 1).

Hikers disturbance

All the summer home ranges used by the studied chamois were crossed by at least one hiking

trail, more or less used by hikers. Our spatial disturbance index ranged between 0.00317 and

0.02912 (*median* = 0.00577, *SD* = 0.00433), with only seven chamois above 0.01000

(Supporting information 1). Chamois living in highly exposed areas always had a

significantly higher total activity than those living in weakly exposed areas of *approx.* 0.1

unit (almost 1% variation; Table 1, Fig. 2.a), except during cold highly disturbed days where

they were similar. However, most exposed chamois significantly decreased their diurnal

activity during hot days by *approx.* 0.1 unit (more than 2% variation) as compared to weakly

exposed ones and both did not extensively modify their nocturnal activity (Table 1, Fig. 2.b

and 2.c). Similarly, chamois did not significantly modify the timing of their activity peaks

according to the level of hikers exposure of their summer home range (consider large

confidence intervals in Table 2 and Fig. 3).

Chamois significantly increased their total activity level of *approx.* 0.1 unit (1%

variation) during highly disturbed days compared to weakly disturbed days, except for

chamois living in highly exposed areas during cold days, where it is the opposite (Table 1,

Fig. 2.a). In terms of nocturnal/diurnal activity, patterns were less clear regarding the effect of

temporal disturbance. Chamois living in highly disturbed areas tended to slightly reduce their

diurnal activity during highly disturbed compared to weakly disturbed days by less than 0.1

unit (less than 1% variation). To the contrary, chamois living in weakly disturbed areas

increased their diurnal activity by *approx.* 0.1 unit (more than 1% variation). The amount of

nocturnal activity did not change extensively according to temporal disturbance (Fig. 2.b and

2.c). Moreover, the timing of morning and evening activity peaks was not affected by the

level of temporal disturbance (consider large confidence intervals in Table 2 and Fig. 3).

Discussion

General results

Our objective was to determine and compare the relative contribution of summer

temperatures and hikers disturbance on the activity pattern of a wild ungulate, the northern

chamois. In agreement with **H1**, chamois adjusted their circadian activity to temperature by

shifting their activity peak earlier in the morning (and to a lesser extent, later in the evening),

and by being more active at night and less active during daytime during hot days.

Surprisingly, their total daily activity was higher during hot than cold days owing to a

significant increase of activity level during the activity peaks. Furthermore, even if this

response was less pronounced, chamois living in areas highly exposed to hikers also adjusted

their activity pattern, particularly during hot days, as expected from **H3**. When living in

disturbed areas and during disturbed days, they slightly decreased their activity during

daytime and increased it during the following night, confirming **H2**. Although they also

slightly delayed the morning activity peak during cold days, hikers disturbance had very little

effect on timing of activity peaks, most likely due to a temporal mismatch with hiker arrival

and departure (*e.g.* more than 2 hours after the morning peak). We also revealed

modifications of the activity levels in chamois living in areas weakly exposed to hikers

disturbance.

Additional effect of potential versus actual exposure to hikers disturbance

Even during weakly disturbed days (*i.e.* week days), chamois highly exposed to hikers disturbance, due to the presence of highly used trails in their home range, generally demonstrated stronger responses than those living in weakly exposed areas, *i.e.* home ranges where trails were less used and/or less represented. For instance, during hot days, chamois living in highly exposed areas decreased their diurnal activity compared to chamois living in weakly exposed areas, either during highly or weakly disturbed days. During highly disturbed days (*i.e.* week-ends), chamois living in highly disturbed areas decreased their activity even more, demonstrating a cumulative effect of potential exposure to disturbance and actual disturbance itself, directly illustrating the effect of the landscape of fear (Laundre, Hernández, & Altendorf, 2001). But contrary to our expectations, most of the activity metrics explored also varied for chamois living in weakly exposed areas according to the level of temporal disturbance they experienced. Samia and colleagues showed that weakly exposed populations among birds, mammals and lizards can be less tolerant to human disturbance because they are less used to it (Samia, Nakagawa, Nomura, Rangel, & Blumstein, 2015). In our population, most of the individuals (85%) move temporarily away from trails every day when hikers are present (Courbin *et al.,* 2022). This low tolerance might explain the weak difference between most/less disturbed chamois either through a similar response to human presence whatever the encounter risk (human being almost present everyday) or through the relocation of their activity in habitats farther away from the main source of disturbance (*i.e.* trails). Cumulative effects of exposure to human disturbance and actual disturbance are far more important than they seem, particularly because they can induce a delayed response instead of a direct, easily observable one (Chassagneux *et al.,* 2020).

Interaction between human disturbance and temperature

Though chamois seem to be less sensitive to high temperatures than other wild ungulate species (see Darmon *et al.,* 2014 for a comparison between chamois and mouflon), their faecal glucocorticoid metabolites concentration increases during drought conditions in summer (Anderwald, Campell Andri, & Palme 2021), as well as the time they spend in thermal refuges (Malagnino *et al.*, unpublished). This may explain that chamois responded much less strongly to human disturbance than to temperature. Moreover, in our study site, hiker attendance was significantly higher when temperature was high (42 ± 55 SD hikers during cold days, 61 ± 37 SD hikers during hot days; *Mann-Whitney test statistic* = 322.5, *p-value* = 0.003). Hence, the response to human disturbance could be lower than expected as the behavioural changes to buffer from high temperatures may also prevent chamois from hikers disturbance, as both factors are correlated. Animals have to face several sources of disturbance (*e.g.* human disturbance, predation, high temperatures) and they eventually cannot indefinitely increase their behavioural response as disturbance factors accumulate (Benoist, Garel, Cugnasse, & Blanchard, 2013).

Recent studies provided evidence of a non-negligeable nocturnal activity peak in chamois (Carnevali, Lovari, Monaco, & Mori, 2016). Here, we also highlighted that nocturnal activity significantly increased while diurnal activity decreased during hot days, confirming the influence of high temperatures and human disturbance on chamois activity. Similarly to numerous other species, chamois decreased their activity when exposed to high temperatures and/or hikers disturbance and attempted to compensate for the activity loss when it is absent (*e.g.* later in the day or at night, Gaynor, Hojnowski, Carter, & Brashares, 2018). However, an increase in nocturnal activity could expose individuals to other threats, such as nocturnal predators (Bonnot *et al.,* 2020). This is particularly true in our study site with the progressive return of wolves recorded since the end of the study period. Large

herbivores may face the difficult challenge of avoiding being exposed to high temperature in

the middle of the day while limiting their exposure to human disturbance during the rest of

the day, but without shifting completely their activity at night because of predation risk.

Chamois can spatially adjust to dampen the impact of hiker disturbance (Duparc,

2016; Courbin *et al.,* 2022). These diel migrations (Courbin *et al.,* 2022) allow chamois to

maintain a certain level of activity during disturbed periods. Unfortunately, our data provides

no clue about the nature of the behaviours displayed (*i.e.* foraging, resting, moving or social

behaviours). But contrary to high temperatures for which any kind of behaviour is costly and

should thus be reduced, leading to a lower activity level, some behaviours are favoured in the

context of human disturbance and could produce inconsistent activity patterns. For instance,

an escape movement is a response to risk which leads to an increase in the activity level,

while resting in a refuge area can also be a response to risk (through avoidance) but leads to a

lower activity level.

Mountain animals face increasing temperatures through the ongoing climate warming,

and modifications of their landscape of fear through the development of recreational activities

and the return of predators in these landscapes. Both constraints can lead either to similar

responses (*e.g.* rest in refuge areas), or to opposite reactions (*e.g.* reduce activity to avoid

thermal stress or escape to limit predation risk), which complicates their behavioural response

(Courbin *et al.,* 2022). That is why identifying the effect of human disturbance on wildlife

remains a difficult task (Tablado & Jenni, 2017), and an integrated approach is needed to take

into account all the covariables affecting chamois activity, such as temperature and predation.

Such an approach will be possible in our study site in the forthcoming years (consider

Courbin *et al.,* 2022 and Malagnino *et al.*, unpublished together with this study).

Management implications

As ecosystem engineers, wild ungulates play a major role in mountain landscapes, by their movements (*e.g.* seed dispersal via zoochory, Albert *et al.,* 2015) and grazing (*e.g.* plant selection via grazing, Austrheim & Eriksson, 2001). Most of the populations of large mammals living in the Alps and surrounding massifs have to face several kinds of disturbance, ranging from hiking to cable car ebb and flow, while constrained by increasing temperature during summer due to climate change (Brivio *et al.,* 2016). As tourism development is often a pillar of the local economy, the right balance must be found between natural environment and tourism management (Reynolds & Braithwaite, 2001). Human impact on chamois could become an issue given that it may decrease the size of their suitable habitats (Peksa & Ciach, 2015), generate a necessary reorganisation of their circadian activity patterns, and reduce their ability to satisfactorily cope with thermal stress, particularly in the context of increasing temperatures. The interaction between those factors could eventually jeopardise their demography if particular attention is not paid in conserving habitats providing both shelter against summer heat and human disturbance (Frid & Dill, 2002; Chirichella, Stephens, Mason, & Apollonio, 2021). Our study contributes to filling the knowledge gap on the cumulative effects of weather and human disturbance on chamois circadian activity in this context of ongoing socio-environmental changes (Corlatti *et al.,* 2022).

Authors' contributions

LT, MG, PM, AL, GB conceived the ideas and designed methodology; MG, PM and AL organised data collection; LT analysed the data; LT led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Acknowledgments

This study was funded by Agence Nationale de la Recherche (French National Research Agency) (ANR) Grant Mov-It No.16-CE02-0010 coordinated by Anne Loison, Centre National de la Recherche Scientifique (Centre National de la Recherche Scientifique) (CNRS). The authors thank all participants for helping with the capture and marking of animals, for the recovery of GPS collars, and for collecting GPS data on human activities: volunteers and professionals from the Office Français de la Biodiversité (formerly Office National de la Chasse et de la Faune Sauvage), the Office National des Forêts, the Groupement d'Intérêt Cynégétique des Bauges, the Parc Naturel Régional du massif des Bauges, the Observatoire Grande Faune et Habitats, and the laboratory Environnements Dynamiques Territoires Montagnes (EDYTEM). This work was performed using the computing facilities of the IFB Cloud.

Conflict of Interest

The authors declared they have no conflict of interest.

Data availability statement

Data will be available after acceptance of the manuscript.

References

Albert, A., Auffret, A. G., Cosyns, E., Cousins, S. A. O., D'hondt, B., Eichberg, C., Eycott, A. E., Heinken, T., Hoffmann, M., Jaroszewicz, B., & others. (2015). Seed dispersal by ungulates as an ecological filter: A trait-based meta-analysis. Oikos, 124(9), 1109–1120.

Anderwald, P., Campell Andri, S., & Palme, R. (2021). Reflections of ecological differences? Stress responses of sympatric Alpine chamois and red deer to weather, forage quality, and human disturbance. Ecology and Evolution, 11(22), 15740–15753. https://doi.org/10.1002/ece3.8235

Arlettaz, R., Patthey, P., Baltic, M., Leu, T., Schaub, M., Palme, R., & Jenni-Eiermann, S. (2007). Spreading free-riding snow sports represent a novel serious threat for wildlife. Proceedings of the Royal Society B: Biological Sciences, 274(1614), 1219–1224.

Austrheim, G., & Eriksson, O. (2001). Plant species diversity and grazing in the Scandinavian mountains-patterns and processes at different spatial scales. Ecography, 24(6), 683–695.

Beniston, M. (2003). Climatic change in mountain regions: a review of possible impacts. Climate Variability and Change in High Elevation Regions: Past, Present & Future, 5–31.

Benoist, S., Garel, M., Cugnasse, J.-M., & Blanchard, P. (2013). Human disturbances, habitat characteristics and social environment generate sex-specific responses in vigilance of Mediterranean mouflon. PLoS One, 8(12), e82960.

Bivand, R., & Lewin-Koh, N. (2022). maptools: Tools for Handling Spatial Objects. https://cran.r-project.org/package=maptools

Bonnot, N. C., Couriot, O., Berger, A., Cagnacci, F., Ciuti, S., De Groeve, J. E., Gehr, B., Heurich, M., Kjellander, P., Kröschel, M., & others. (2020). Fear of the dark?

511  Contrasting impacts of humans versus lynx on diel activity of roe deer across Europe.

512  Journal of Animal Ecology, 89(1), 132–145. https://doi.org/10.1111/1365-2656.13161

513 Bonnot, N., Morellet, N., Verheyden, H., Cargnelutti, B., Lourtet, B., Klein, F., & Hewison,

514  A. J. (2013). Habitat use under predation risk: hunting, roads and human dwellings

515  influence the spatial behaviour of roe deer. European Journal of Wildlife Research,

516  59(2), 185–193.

517 Bourgoin, G, Garel, M., Van Moorter, B., Dubray, D., Maillard, D., Marty, E., & Gaillard, J.-

518  M. (2008). Determinants of seasonal variation in activity patterns of mouflon. Canadian

519  Journal of Zoology, 86(12), 1410–1418.

520 Bourgoin, G., Garel, M., Blanchard, P., Dubray, D., Maillard, D., & Gaillard, J.-M. (2011).

521  Daily responses of mouflon (Ovis gmelini musimon x Ovis sp.) activity to summer

522  climatic conditions. Canadian Journal of Zoology, 89(9), 765–773.

523  https://doi.org/10.1139/Z11-046

524 Brivio, F., Bertolucci, C., Tettamanti, F., Filli, F., Apollonio, M., & Grignolio, S. (2016). The

525  weather dictates the rhythms: Alpine chamois activity is well adapted to ecological

526  conditions. Behavioral Ecology and Sociobiology, 70(8), 1291–1304.

527 Byers, J. A. (1997). American pronghorn: social adaptations and the ghosts of predators past.

528  University of Chicago Press.

529 Calenge, C. (2006). The package adehabitat for the R software: tool for the analysis of space

530  and habitat use by animals. Ecological Modelling, 197, 1035.

531 Carnevali, L., Lovari, S., Monaco, A., & Mori, E. (2016). Nocturnal activity of a diurnal

532  species, the northern chamois, in a predator-free Alpine area. Behavioural Processes,

533  126, 101–107.

534 Chanteloup, L., Perrin-Malterre, C., Duparc, A., & Loison, A. (2016). Quels points de vue sur

535  les espaces partagés entre humains et animaux sauvages? Espaces et Sociétés, 1, 33–47.

Chassagneux, A., Calenge, C., Marchand, P., Richard, E., Guillaumat, E., Baubet, E., & Saïd,

S. (2020). Should I stay or should I go? Determinants of immediate and delayed

movement responses of female red deer (Cervus elaphus) to drive hunts. Plos One,

15(3), e0228865.

Chirichella, R., Stephens, P. A., Mason, T. H. E., & Apollonio, M. (2021). Contrasting

effects of climate change on Alpine chamois. The Journal of Wildlife Management,

85(1), 109–120.

Clutton-Brock, T. H., Albon, S. D., & Guinness, F. E. (1989). Fitness costs of gestation and

lactation in wild mammals. Nature, 337(6204), 260–262.

https://doi.org/10.1038/337260a0

Corlatti, L., Iacolina, L., Safner, T., Apollonio, M., Buzan, E., Ferretti, F., Hammer, S. E.,

Herrero, J., Rossi, L., Serrano, E., & others. (2022). Past, present and future of chamois

science. Wildlife Biology, e01025. https://doi.org/10.1002/wlb3.01025

Courbin, N., Garel, M., Marchand, P., Duparc, A., Debeffe, L., Börger, L., & Loison, A.

(2022). Interacting lethal and nonlethal human activities shape complex risk tolerance

behaviors in a mountain herbivore. Ecological Applications, e2640.

https://doi.org/10.1002/eap.2640

Darmon, G., Bourgoin, G., Marchand, P., Garel, M., Dubray, D., Jullien, J.-M., & Loison, A.

(2014). Do ecologically close species shift their daily activities when in sympatry? A

test on chamois in the presence of mouflon. Biological Journal of the Linnean Society,

111(3), 621–626. https://doi.org/10.1111/bij.12228

Duparc, A. (2016). Dynamiques spatio-temporelles des ressources alimentaires et des

activités humaines: impacts sur la sélection d'habitat d'un grand herbivore de

montagne. Université Grenoble Alpes.

Duparc, A., Redjadj, C., Viard-Crétat, F., Lavorel, S., Austrheim, G., & Loison, A. (2013).

Co-variation between plant above-ground biomass and phenology in sub-alpine

grasslands. Applied Vegetation Science, 16(2), 305–316.

Fattorini, N., Brunetti, C., Baruzzi, C., Chiatante, G., Lovari, S., & Ferretti, F. (2019).

Temporal variation in foraging activity and grouping patterns in a mountain-dwelling

herbivore: Environmental and endogenous drivers. Behavioural Processes, 167,

103909.

French, S. S., González-Suárez, M., Young, J. K., Durham, S., & Gerber, L. R. (2011).

Human disturbance influences reproductive success and growth rate in California sea

lions (Zalophus californianus). PloS One, 6(3), e17686.

https://doi.org/10.1371/journal.pone.0017686

Frid, A., & Dill, L. (2002). Human-caused disturbance stimuli as a form of predation risk.

Conservation Ecology, 6(1). https://doi.org/10.5751/es-00404-060111

Gaillard, J.-M., Festa-Bianchet, M., Yoccoz, N. G., Loison, A., & Toigo, C. (2000).

Temporal variation in fitness components and population dynamics of large herbivores.

Annual Review of Ecology and Systematics, 31(1), 367–393.

https://doi.org/10.1146/annurev.ecolsys.31.1.367

Gaynor, K. M., Hojnowski, C. E., Carter, N. H., & Brashares, J. S. (2018). The influence of

human disturbance on wildlife nocturnality. Science, 360(6394), 1232–1235.

https://doi.org/10.1126/science.aar7121

Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., & Stoffel, M. (2014). 21st

century climate change in the European Alps - A review. Science of the Total

Environment, 493, 1138–1151.

Greives, T. J., Kingma, S. A., Kranstauber, B., Mortega, K., Wikelski, M., van Oers, K.,

Mateman, A. C., Ferguson, G. A., Beltrami, G., & Hau, M. (2015). Costs of sleeping in:

circadian rhythms influence cuckoldry risk in a songbird. Functional Ecology, 29(10),

1300–1307.

Hagemoen, R. I. M., & Reimers, E. (2002). Reindeer summer activity pattern in relation to

weather and insect harassment. Journal of Animal Ecology, 71(5), 883–892.

Hamel, S., & Côté, S. D. (2008). Trade-offs in activity budget in an alpine ungulate:

contrasting lactating and nonlactating females. Animal Behaviour, 75(1), 217–227.

Kearney, M. R., Gillingham, P. K., Bramer, I., Duffy, J. P., & Maclean, I. M. D. (2020). A

method for computing hourly, historical, terrain-corrected microclimate anywhere on

Earth. Methods in Ecology and Evolution, 11(1), 38–43.

Kearney, M. R., & Porter, W. P. (2017). NicheMapR - an R package for biophysical

modelling: the microclimate model. Ecography, 40(5), 664–674.

Larson, C. L., Reed, S. E., Merenlender, A. M., & Crooks, K. R. (2016). Effects of recreation

on animals revealed as widespread through a global systematic review. PloS One,

11(12), e0167259.

Laundré, J. W., Hernández, L., & Altendorf, K. B. (2001). Wolves, elk, and bison:

reestablishing the "landscape of fear" in Yellowstone National Park, USA. Canadian

Journal of Zoology, 79(8), 1401–1409.

Lesmerises, F., Déry, F., Johnson, C. J., & St-Laurent, M.-H. (2018). Spatiotemporal

response of mountain caribou to the intensity of backcountry skiing. Biological

Conservation, 217, 149–156.

Li, C., Jiang, Z., Feng, Z., Yang, X., Yang, J., & Chen, L. (2009). Effects of highway traffic

on diurnal activity of the critically endangered Przewalski gazelle. Wildlife Research,

36(5), 379–385.

Lima, S. L., & Bednekoff, P. A. (1999). Temporal variation in danger drives antipredator

behavior: the predation risk allocation hypothesis. The American Naturalist, 153(6),

649–659.

Lindstedt, S. L., & Boyce, M. S. (1985). Seasonality, fasting endurance, and body size in

mammals. The American Naturalist, 125(6), 873–878. https://doi.org/10.1086/284385

Loison, A., Jullien, J.-M., & Menaut, P. (1999). Subpopulation structure and dispersal in two

populations of chamois. Journal of Mammalogy, 80(2), 620–632.

Long, R. A., Martin, T. J., & Barnes, B. M. (2005). Body temperature and activity patterns in

free-living arctic ground squirrels. Journal of Mammalogy, 86(2), 314–322.

Lopez, J. F. (2001). Inventaire systématique de la flore et de la végétation du Massif des

Bauges.

Maclean, I. M. D., Mosedale, J. R., & Bennie, J. J. (2019). Microclima: An r package for

modelling meso-and microclimate. Methods in Ecology and Evolution, 10(2), 280–290.

Marchand, P., Garel, M., Bourgoin, G., Dubray, D., Maillard, D., & Loison, A. (2014).

Impacts of tourism and hunting on a large herbivores spatio-temporal behavior in and

around a French protected area. Biological Conservation, 177, 1–11.

Marchand, P., Garel, M., Bourgoin, G., Dubray, D., Maillard, D., & Loison, A. (2015). Sex-

specific adjustments in habitat selection contribute to buffer mouflon against summer

conditions. Behavioral Ecology, 26(2), 472–482.

Pęksa, Ł., & Ciach, M. (2015). Negative effects of mass tourism on high mountain fauna: the

case of the Tatra chamois Rupicapra rupicapra tatrica. Oryx, 49(3), 500–505.

Palmer M. S., Gaynor K. M., Becker J. A., Abraham J. O., Mumma M. A., & Pringle R. M.

(2022). Dynamic landscapes of fear: understanding spatiotemporal risk. Trends in

Ecology & Evolution. https://doi.org/10.1016/j.tree.2022.06.007

Pepin, N., Bradley, R. S., Diaz, H. F., Baraër, M., Caceres, E. B., Forsythe, N., Fowler, H.,

Greenwood, G., Hashmi, M. Z., Liu, X. D., & others. (2015). Elevation-dependent

warming in mountain regions of the world. Nature Climate Change, 5(5), 424–430.

Phillips, G. E., & Alldredge, A. W. (2000). Reproductive success of elk following

disturbance by humans during calving season. The Journal of Wildlife Management,

521–530.

Reynolds, P. C., & Braithwaite, D. (2001). Towards a conceptual framework for wildlife

tourism. Tourism Management, 22(1), 31–42.

Rogala, J. K., Hebblewhite, M., Whittington, J., White, C. A., Coleshill, J., & Musiani, M.

(2011). Human activity differentially redistributes large mammals in the Canadian

Rockies National Parks. Ecology and Society, 16(3).

Samia, D. S. M., Nakagawa, S., Nomura, F., Rangel, T. F., & Blumstein, D. T. (2015).

Increased tolerance to humans among disturbed wildlife. Nature Communications, 6(1),

1–8.

Saper, C. B., Lu, J., Chou, T. C., & Gooley, J. (2005). The hypothalamic integrator for

circadian rhythms. Trends in Neurosciences, 28(3), 152–157.

Scott, D. (2006). Global environmental change and mountain tourism. Tourism and Global

Environmental Change, 54–75.

Searle, K. R., Hobbs, N. T., & Gordon, I. J. (2007). It's the" foodscape", not the landscape:

using foraging behavior to make functional assessments of landscape condition. Israel

Journal of Ecology and Evolution, 53(3–4), 297–316.

Semenzato, P., Cagnacci, F., Ossi, F., Eccel, E., Morellet, N., Hewison, A. J. M., Sturaro, E.,

& Ramanzin, M. (2021). Behavioural heat-stress compensation in a cold-adapted

ungulate: Forage-mediated responses to warming Alpine summers. Ecology Letters,

24(8), 1556–1568.

Strava. 2022. Strava Global Heatmap. https://www.strava.com/heatmap.

Tablado, Z., & Jenni, L. (2017). Determinants of uncertainty in wildlife responses to human

disturbance. Biological Reviews, 92(1), 216–233.

Takahashi, J. S., Hong, H.-K., Ko, C. H., & McDearmon, E. L. (2008). The genetics of

mammalian circadian order and disorder: implications for physiology and disease.

Nature Reviews Genetics, 9(10), 764–775.

Thorsen, N. H., Bischof, R., Mattisson, J., Hofmeester, T. R., Linnell, J. D. C., & Odden, J.

(2022). Smartphone app reveals that lynx avoid human recreationists on local scale, but

not home range scale. Scientific Reports, 12(1), 1–13.

van Beest, F. M., & Milner, J. M. (2013). Behavioural responses to thermal conditions affect

seasonal mass change in a heat-sensitive northern ungulate. PloS One, 8(6), e65972.

Watanuki, Y., & Nakayama, Y. (1993). Age difference in activity pattern of Japanese

monkeys: effects of temperature, snow, and diet. Primates, 34(4), 419–430.

Wheaton, B. (2010). Introducing the consumption and representation of lifestyle sports. Sport

in Society, 13(7–8), 1057–1081.

Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood

estimation of semiparametric generalized linear models. Journal of the Royal Statistical

Society (B), 73(1), 3–36.

Wood, S. N. (2017). Generalized Additive Models: An Introduction with R (2nd ed.).

Chapman and Hall/CRC.

Wood, S. N, Goude, Y., & Shaw, S. (2015). Generalized additive models for large data sets.

Journal of the Royal Statistical Society: Series C (Applied Statistics), 64(1), 139–155.

Worton, B. J. (1989). Kernel methods for estimating the utilization distribution in home-

range studies. Ecology, 70(1), 164–168.

Xu, W., Gong, Y., & Wang, H. (2021). Alert time reflects the negative impacts of human

disturbance on an endangered bird species in Changbai Mountain, China. Global

Ecology and Conservation, 28, e01709.

Tables and Figures

Table 1: Total, diurnal and nocturnal Area Under Curve (unitless) estimated from predicted circadian activity patterns (median value of the bootstraps distribution, $n = 1001$) of northern chamois of the National Game and Hunting Reserve of Les Bauges massif, in summer (June to September, 2004-2020), according to temperature and human disturbance. We defined "diurnal" as the period elapsed between two hours after dawn and two hours before dusk, and "nocturnal" as the period elapsed between two hours after dusk and two hours before dawn (see text for details). Values between brackets correspond to 2.5% and 97.5% quantiles (95% confidence intervals) estimated with bootstraps.

Disturbance-Temperature variable levels			Total activity	Diurnal activity	Nocturnal activity
Potential exposure	Hikers disturbance	Temperature			
Low	Low	Low	10.89 [10.72; 11.06]	5.89 [5.77; 6.01]	1.09 [1.08; 1.09]
Low	High	Low	11.03 [10.86; 11.19]	5.98 [5.87; 6.11]	1.10 [1.08; 1.10]
Low	Low	High	11.41 [11.23; 11.54]	5.39 [5.28; 5.51]	1.34 [1.31; 1.34]
Low	High	High	11.53 [11.35; 11.66]	5.46 [5.36; 5.59]	1.31 [1.28; 1.31]
High	Low	Low	11.07 [10.90; 11.24]	6.03 [5.91; 6.15]	1.11 [1.09; 1.11]
High	High	Low	11.03 [10.86; 11.20]	5.99 [5.87; 6.12]	1.12 [1.10; 1.12]
High	Low	High	11.50 [11.32; 11.63]	5.33 [5.22; 5.45]	1.34 [1.31; 1.34]
High	High	High	11.58 [11.40; 11.72]	5.30 [5.19; 5.42]	1.38 [1.35; 1.38]

Table 2: Timing of the morning and evening peaks (UTC) estimated from predicted circadian

activity patterns (median value of the bootstraps distribution, n = 1001) of northern chamois

of the National Game and Hunting Reserve of Les Bauges massif, in summer (June to

September, 2004-2020), according to temperature and human disturbance. Values between

brackets correspond to 2.5% and 97.5% quantiles (95% confidence intervals) estimated with

bootstraps.

Disturbance-Temperature variable levels			Morning peak timing	Evening peak timing
Potential exposure	Hikers disturbance	Tempera-ture		
Low	Low	Low	05:40 [05:10; 05:45]	19:10 [19:10; 19:15]
Low	High	Low	05:50 [05:25; 05:55]	19:05 [19:05; 19:10]
Low	Low	High	04:50 [04:40; 05:05]	19:10 [19:10; 19:15]
Low	High	High	04:55 [04:45; 05:05]	19:15 [19:15; 19:20]
High	Low	Low	05:40 [05:15; 05:40]	19:10 [19:05; 19:10]
High	High	Low	05:55 [05:30; 05:55]	19:05 [19:05; 19:10]
High	Low	High	04:45 [04:40; 05:10]	19:15 [19:15; 19:15]
High	High	High	04:55 [04:45; 05:10]	19:20 [19:20; 19:20]

<mark>Figure 1</mark>: predicted circadian activity patterns (activity ranges between 0 for no activity, to 1

for the highest level of activity) of female northern chamois in the National Game and

Hunting Reserve of Les Bauges massif, in summer (15th June to 7th September, 2004-2020),

according to temperature (blue: cold day; red: hot day) and human disturbance: a) chamois

living in a weakly exposed area during a weakly disturbed day; b) chamois living in a weakly

exposed area during a highly disturbed day; c) chamois living in a highly exposed area during

a weakly disturbed day; d) chamois living in a highly exposed area during a highly disturbed

day. Solid lines represent predicted values from the model, shaded areas represent 95%

confidence intervals. Time of the day represents hours in Coordinated Universal Time (UTC).

Figure 2: Area Under Curve (AUC) differential (unitless) between the reference situation (*i.e.*

chamois living in a weakly exposed area during a cold weakly disturbed day) and the

situation of interest, a combination of different levels of temperature (blue: cold day; red: hot

day), spatial (dot: low exposure; triangle: high exposure) and hikers attendance (low = week

days; high = week-ends): a) total, b) diurnal and c) nocturnal activity. AUC are calculated

from predicted circadian activity patterns of northern chamois of the National Game and

Hunting Reserve of Les Bauges massif, in summer (June to September, 2004-2020; see Fig.

1). The central point represents the median, and the vertical bars represent the 2.5% and

97.5% quantiles of the bootstraps distribution (*n* = 1001). Note the change of scale between

Fig. 2.a and Fig. 2.b and 2.c.

Figure 3: Activity Peak Timing (APT) differential (in minutes) between the reference

situation (*i.e.* chamois living in a weakly exposed area during a cold weakly disturbed day)

and the situation of interest, a combination of different levels of temperature (blue: cold day;

red: hot day), spatial (dot: low exposure; triangle: high exposure) and temporal human

disturbance: a) morning and b) evening peak. APT are calculated from predicted circadian

activity patterns of northern chamois of National Game and Hunting Reserve of Les Bauges

massif, in summer (June to September, 2004-2020). The central point represents the median,

and the vertical bars represent the 2.5% and 97.5% quantiles of the bootstraps distribution (*n*

= 1001). The points have been shifted on both sides to improve readability.