Too hot or too disturbed? Temperatures more than hikers affect circadian activity of females in northern chamois

Thel Lucie[1,2], Garel Mathieu[3], Marchand Pascal[4], Bourgoin Gilles[5,6], Loison Anne[1]

Authors details

[1] Laboratoire d'Écologie Alpine, UMR 5553 (CNRS, Université Savoie Mont--Blanc), Savoie Technolac, 73376 Le Bourget-du-Lac, France.

[2] Laboratoire Évolution et Diversité Biologique, UMR 5174 (CNRS, Université Paul Sabatier), 118 route de Narbonne, 31062 Toulouse, France.

[3] Office Français de la Biodiversité, Direction de la Recherche et de l'Appui Scientifique, Service Anthropisation et Fonctionnement des Écosystèmes Terrestres, Gières, France.

[4] Office Français de la Biodiversité, Direction de la Recherche et de l'Appui Scientifique, Service Anthropisation et Fonctionnement des Écosystèmes Terrestres, Juvignac, France.

[5] Univ Lyon, VetAgro Sup, Campus Vétérinaire de Lyon, 1 avenue Bourgelat, F-69280 Marcy l'Etoile, France.

[6] Univ Lyon, Université Lyon 1, CNRS, UMR 5558, Laboratoire de Biométrie et Biologie Évolutive (LBBE), F-69622 Villeurbanne Cedex, France.

Correspondence author: Thel Lucie, lucie.mc.thel@gmail.com.

Too hot or too disturbed? Temperatures more than hikers

affect circadian activity of females in northern chamois

Abstract

Though non-lethal, recreational activities often result in a spatial and/or temporal shift of

activity in a large range of species because they are perceived as a risk. With the concurrent

development of outdoors activities and the increase of temperatures due to climate change in

mountainous ecosystems, mountain species face an increasing pressure in terms of managing

their activity pattern to both limit risk exposure and reduce thermal discomfort. In this

context, we investigated how female northern chamois adjust their summer circadian activity

to spatio-temporal variations of both temperature and hikers' disturbance. Chamois behaviour

was more affected by high temperatures than by hikers' disturbance. During the hottest days

(*median ± SD* = 17.1±1.2°C *vs.* 8.8±1.7°C during the coldest days), they shifted their activity

peak earlier in the morning (- 17 minutes), and were more active at night (+ 22%) than during

daytime (- 10%). Their global daily activity was slightly higher during the hottest days than

during the coldest days (+ 5%). Conversely, hikers' spatial and temporal disturbance had only

weak effects on activity of chamois, most likely due to a temporal mismatch with hikers,

except during the hottest conditions, when the duration of the morning activity peak

decreased (- 46 minutes) and the duration of the evening peak increased (+ 18 minutes) with

the highest compared to the lowest disturbance. The overall low effect of hikers' disturbance

on chamois activity pattern suggested that behavioural changes buffering animals from high

temperature (relocation into thermal refuges) and from hikers' presence (moving away from

trails) allow them to just marginally modify their activity pattern in the presence of hikers. In

the context of ongoing socio-environmental changes, these findings stress the importance of

conserving habitats providing thermal refuges against summer heat and preventing them from

human disturbance to mitigate their potential detrimental consequences.

Keywords: behavioural thermoregulation, circadian rhythm, climate warming, ecotourism,

human disturbance, mountain ecosystem, nocturnality, recreational activities, *Rupicapra*

rupicapra rupicapra

Introduction

A noticeable development and diversification of outdoor activities has been ongoing in the

twenty-first century, especially in Western countries (Wheaton, 2010). While mountain

ecosystems have been preserved until recently because of their low accessibility, they now

face an increase of unaffiliated activities such as hiking, cross country skiing, trail running, or

climbing. Those recreational activities are non-lethal, but humans can be consistently

perceived as predators by wildlife (Frid & Dill, 2002), leading to a suite of negative

physiological and behavioural responses (Arlettaz *et al.,* 2007; Marchand *et al.,* 2014;

Larson, Reed, Merenlender, & Crooks, 2016). The consequences of repeated reactions to

disturbance can be detrimental on several fitness-related components (Phillips & Alldredge,

2000; French, González-Suárez, Young, Durham, & Gerber, 2011). This is particularly true

in mountain ecosystems, where the diversification of human activities challenges animals all

year long, both during the period of energy acquisition from spring to autumn, and of energy

loss during winter (Lindstedt & Boyce, 1985). This increased pressure on the animals is

accompanied by environmental modifications, including climate change, that are particularly

strong in the Alps (Pepin *et al.,* 2015). Accordingly, temperatures are rising rapidly in all

seasons, with consequences such as earlier springs and plant cycles, more frequent summer

heatwaves and droughts, and decreased duration of snow cover associated with an augmented

risk of frost for plants (*e.g.,* Gobiet *et al.,* 2014). Mountain animals are therefore facing

concomitantly two major changes, namely an increased disturbance level and climate change.

To reduce their exposure to human disturbance, animals can adjust their behaviour in

terms of movements, habitat use and selection, and time devoted to their different activities

(foraging, resting, vigilance and social interactions). When facing a disturbance perceived as

a risk, animals can modify their activities (*e.g.,* from foraging to vigilance; Benoist, Garel,

Cugnasse, & Blanchard, 2013; Xu, Gong, & Wang, 2021), and move away temporarily or

even definitively (Rogala *et al.,* 2011). They can also modify their activity pattern by

delaying activities, such as foraging in disturbed places, from high- to low- risk periods

(Bonnot *et al.,* 2013). The benefits of such responses have been explained within the "risk

allocation hypothesis" framework (Lima & Bednekoff, 1999): the costs of lost opportunities

to forage during risky periods could be dampened if animals transfer part of their diurnal

activity to night-time when human activities stop for instance (*e.g.,* Marchand *et al.,* 2014;

Bonnot *et al.,* 2020). However, activity patterns are regulated, such as the critical

foraging/rumination succession in ruminants (Hamel & Côté, 2008). Any forced modification

of those patterns can negatively impact the fitness of individuals *via* an increased encounter

risk with predators (Lesmerises *et al.,* 2018) or mismatches in interactions with conspecifics

(Greives *et al.,* 2015) for instance.

At the same time, animals need to deal with climatic constraints, and behavioural

thermoregulation is among the most important mechanisms to avoid thermal stress. Indeed,

temperature influences space use (in arctic ground squirrels *Spermophilus parryii*, Long,

Martin, & Barnes, 2005; in Mediterranean mouflon *Ovis gmelini musimon* x *Ovis* sp.,

Marchand *et al.,* 2015) and activity budget (in Japanese monkeys *Macaca fuscata*, Watanuki

& Nakayama, 1993), with animals having to trade off foraging and thermal cover to limit

thermoregulatory costs while keeping their energy balance (van Beest & Milner, 2013). In the

same way as to avoid human disturbance, animals can shift their activity peaks earlier in the morning and later in the evening, and increase their nocturnal activity during the hottest summer days (Bourgoin *et al.*, 2008; Bourgoin *et al.*, 2011). However, in temperate ecosystems, night can be significantly shorter than day in summer, and riskier when nocturnal predators are present. Thus, the transfer of the diurnal activity to night-time does not necessarily allow to fully compensate for the loss of foraging opportunities during day and may have consequences on mass gain (Garel, Loison, Gaillard, Cugnasse & Maillard, 2004) or predation risk (Gehr *et al.*, 2017) for instance. These results highlight the importance of investigating full daily cycles to get a comprehensive picture of how animals may adjust their behaviour in the face of human-induced environmental changes (Marchand *et al.,* 2014; Marchand *et al.,* 2015).

Moreover, ongoing climate change and expansion of human activities in mountain ecosystems are also ultimately connected. An extended summer season or more frequent summer heatwaves in valleys may, for instance, exacerbate human attendance in mountains. Although predicting tourism attendance under scenarios of climate change is probably not straightforward (Beniston, 2003), this context raises questions on animal responses to those interrelated human-induced environmental changes and on their potential cumulative or even multiplicative consequences for populations in terms of both immediate and compensatory responses (Schmeller *et al.*, 2022). One critical challenge is therefore to understand whether the behavioural mechanisms used by animals to buffer against increasing temperatures may also provide an advantage against human attendance or *vice versa*.

Here, we relied on the GPS monitoring of 62 females of northern chamois *Rupicapra rupicapra rupicapra* to investigate at a fine scale their behavioural responses to temporal and spatial stressors during summer, through the study of their activity pattern. More specifically, we assessed the day-to-day variation in temperature and hikers' attendance for each chamois.

We also computed an individual-specific index of spatial disturbance, combining data on the

density of hiking trails in each chamois summer home range and the attendance of each trail

section by hikers. A diversity of activity metrics allowed us to investigate the relative

contribution of spatio-temporal variations in both temperature and human disturbance to

changes in circadian activity patterns of female chamois in summer. We expected the

chamois to adjust their circadian activity pattern by avoiding being active during the hottest

hours during the hottest days (**H1**) and during the most disturbed days (**H2**), with a

compensation of activity loss during daytime at night. However, as individuals were exposed

to variable levels of disturbance depending on the location of their summer home range, we

also expected more marked responses in chamois that were the most exposed to hikers'

disturbance (**H3**), as most of them show low tolerance to hikers' presence in this population

(Courbin *et al.,* 2022).

Materials and methods

Study site and species

The study took place in the National Game and Hunting Reserve of Les Bauges massif

(NGHRB), a protected area in the French Alps (45°40' N, 6°13' E; 900 to 2200 m a.s.l.;

5,200 ha). The Bauges massif is made of limestone, with cold (mean annual temperature:

9.6°C, snow cover from November to April above 1,600 m, Duparc *et al.,* 2013) and wet

(mean cumulative annual rainfall: 1,600 mm, Duparc *et al.,* 2013) climatic conditions (see

Supporting Information 1 for details about summer temperatures). Fifty-six percent of the

area is covered by forests (treeline around 1,500 m a.s.l. in the study site), dominated by

beech *Fagus sylvatica* and fir *Abies alba*, 36% by grasslands, and 8% by rocks (Lopez,

2001). This heterogeneous environment leads to a huge diversity of habitats and landscapes

that attracts numerous tourists every year (> 37,000, most of them during summer; 1.1

million and 0.5 million overnight stays in summer and winter respectively in 2014, Cloud

Parc des Bauges, 2014). All alpine grasslands are crossed by a network of trails where nearly

all hikers are concentrated due to the NGHRB restrictions, exposing chamois to a very

localised disturbance (Courbin *et al.,* 2022). Other recreational activities (*e.g.,* paragliding)

occur in the study site during summer, but far less frequently than hiking.

Most adult female chamois annually give birth to one young in late May. During

summer, they form groups with other females, kids and yearlings, that share and keep the

same home range from year to year (Loison, Jullien, & Menaut, 1999). Females play a major

role in the demography of the population, particularly through fecundity and survival of

young females (Gaillard, Festa-Bianchet, Yoccoz, Loison, & Toigo, 2000). Chamois have no

natural predators in the study area during the study period, except golden eagles *Aquila*

chrysaetos and red foxes *Vulpes vulpes* that may occasionally predate newborn and injured

animals. However, the presence of some non-resident wolves *Canis lupus* has been noticed in

recent years (Loupfrance.fr, 2021). Hunting performed by small groups of three to four

hunters occurs in the NGHRB from the beginning of September to the end of February

(Courbin *et al.,* 2022), with chamois being the main target species (~100 chamois harvested

per year on average, *i.e.,* <12% of the chamois population in the area and ~70% of the total

number of ungulates harvested), in addition to Mediterranean mouflon (~20%), roe deer

Capreolus capreolus (~4%), wild boar *Sus scrofa* (~4%) and red deer *Cervus elaphus* (~2%;

see Courbin *et al.,* 2022 for details).

Data collection

We focused on summers from 2004 to 2020, between the 15th of June and the 7th of

September. This period allows to encompass days falling in and out from the French summer

holidays which starts in early July, but excludes the hunting season which ranges from the

first weekend of September to the end of February, so that chamois were exclusively exposed

to hikers' disturbance. The study period is critical for female chamois because it follows the

birth season and it is characterised by high energy requirements for lactation and storage of

fat reserves for winter survival (Clutton-Brock, Albon, & Guinness, 1989; Richard, Toïgo,

Appolinaire, Loison & Garel 2017), so that any disturbance during this period could be

particularly detrimental.

We trapped 62 adult female chamois >2 years old during summer using falling nets

baited with salt licks located in four alpine grasslands near two trails highly used by hikers

(Armenaz and Dent des Portes). Although captures occurred during the birth and weaning

period, females generally avoided drop nets (positioned on flat areas) when juveniles were

particularly vulnerable (leading to the use by females of very steep slopes as an anti-predator

strategy). During the weaning period, chamois form nursery groups and kids are not always

closely associated with their mother, so that temporary separation between them, longer than

those resulting from capture/handling, also naturally occurred in this species (Ruckstuhl &

Ingold, 1998). The drop nets were triggered manually by an observer located a few dozen

meters from the nets, preventing non-target captures, and chamois were processed

immediately after the capture by several experienced animal handlers coordinated by the

Office Français de la Biodiversité. As soon as an animal was handled, it was restrained with

the eyes covered to reduce stress and legs attached to avoid injuries. An average of three

individuals were captured at the same time and this number was adjusted according to the

number of handlers to limit the waiting time for each chamois. All efforts were made to

minimise the handling time (approximately 10 to 15 minutes per individual) to release

animals on site as soon as possible. Juveniles and mothers were released at the same time to

prevent them from separation. As part of a long-term Capture-Marking-Recapture

monitoring, recaptures were not avoided, although unmarked individuals were targeted when

present. Since 2003, 916 individuals have been (re)captured, causing 5 deaths (*i.e.,* 0.55% of the individuals handled, generally euthanised using recommended ethical/legal ways after important injuries). All captures, handling and sampling were conducted according to the appropriate national laws for animal welfare, following the ethical conditions detailed in the specific accreditations delivered by the Préfecture de Paris (prefectorial decree n°2009–014) in agreement with the French environmental code (Art. R421-15 to 421–31 and R422-92 to 422–94-1).

Females were equipped with GPS collars Lotek 3300S (Lotek Engineering Inc., Carp, Ontario, Canada). The GPS collars used weighed <450 g and were deployed only on individuals >20 kg judged to be fully healthy so that it never exceeded 2.5% of the body mass of the equipped animal (average weight of females fitted with GPS collar = 30 kg). To allow for possible seasonal variation in neck circumference, we left a gap of ~3 cm between the GPS collar and the animal's neck while making sure that the collar would not rotate. We retrieved the collar using a remote drop-off system triggered manually at the end of the monitoring of the individuals, nearly one year after deployment. From repeated observations of GPS-collared individuals throughout the year as part of long-term Capture-Mark-Recapture monitoring of this population, we have no indication that GPS collars cause changes in behaviour compared to other non-equipped individuals. Similarly, we did not detect any adverse effects linked to the wearing of the collars in terms of injury or reproductive activity.

As deployment/recovery of GPS collars occurred during summer, chamois locations were not necessarily recorded daily during all the study period. Only chamois (n = 62) for which we could reliably estimate the summer home range (95% kernel contour using the *adehabitatHR* package; Worton, 1989; Calenge, 2006) were included in the study, *i.e.,* chamois with locations available during at least 2/3 of their monitoring period, corresponding

to 56 days in total. Animal motion was recorded every 5 minutes by two perpendicular

captive-balls in the GPS collars monitoring neck movements (0: no movement, 255: highly

mobile) and the proportion of time the animal was head down. We discriminated each

chamois as "active" (*i.e.,* feeding, standing, travelling, or doing other activities such as

interactions or scratching) or "inactive" (*i.e.,* sleeping, ruminating or resting) for every 5-

minutes boot using the discriminant model developed by Bourgoin *et al.* (2011; see also

Darmon *et al.,* 2014). From these animals, we obtained a total of 4,464 complete days of

activity records in summer (*n* = 131 chamois-year, 1,285,632 5-minutes activity

measurements in total). The activity of each chamois was monitored during 18 to 129 days in

total, corresponding to 11 to 49 days (*median* = 36, *SD* = 10) during a single summer period

for a given chamois (Supporting Information 2).

We assessed daily characteristics (see below) from 01:39 UTC of day *d* to 01:39 UTC

of day *d+1*. This allowed us to explore potential compensation processes during the night

following the day of interest. We selected the starting hour in accordance with the median

value of the beginning of the crepuscular period in the morning during our study period. We

used a micro-climate model that integrates ERA5 hourly weather data produced by the

Copernicus Climate Change service at ECMWF (spatial resolution 30 km x 30 km) and a

digital elevation model (30 m) for downscaling atmospheric data at the population scale

(micro_era5 function in NicheMapR package; Kearney & Porter, 2017; Maclean, Mosedale,

& Bennie, 2019; Kearney, Gillingham, Bramer, Duffy, & Maclean, 2020). This allowed us to

overcome the issue of incomplete weather data collected in the valley by local weather station

and to get standardized temperatures within the population range over the study period. Mean

daily temperatures were predicted from this model at 1 m high on flat north-facing terrain for

a spatial location and elevation corresponding to the centroid of all chamois GPS locations.

We explored the effect of temperature on female chamois activity through a categorical

variable with two levels defined according to the quantiles of the mean daily temperature

distribution. Days with mean temperature ≤ 25% quantile (*i.e.,* 10.8°C) were considered as

cold days and days with mean temperature > 75% quantile (*i.e.,* 15.5°C) were considered as

hot days. Cold days experienced by chamois ranged between 3.4°C and 10.8°C (*median =*

8.9°C, *SD* = 1.6°C), while hot days ranged between 15.5°C and 20.9°C (*median* = 17.1°C, *SD*

= 1.2°C, see also Supporting Information 1).

231   Temporal disturbance associated with hiking practice was defined as a categorical

variable with two levels. Weekends and public holidays, *i.e.,* vacation days such as Ascension

Day, were considered as **highly disturbed days**. Days between Monday and Friday, *i.e.,*

working days including weekdays during school holidays, were considered as **weakly**

disturbed days. To check for the reliability of these two categories in the field, we compared

the daily attendance of hikers (number of hikers recorded on the trail) between 00:00 UTC

and 23:59 UTC and the hour of first arrival for both temporal disturbance levels using an eco-

counter (PYRO sensor Eco-Compteur, www.eco-compteur.com; 90 cm above ground)

located on Armenaz trail (45°37'18.60" N, 6°13'2.53" E), on a subset of the study period

(2016 and 2017 summers). Our study period encompassed 189 theoretically highly disturbed

days and 440 weakly disturbed days. During the same period of the year in 2016 and 2017,

hikers were almost two times more on the trails during highly (*median* = 73 hikers, *SD* = 53

hikers) than during weakly (*median*= 40 hikers, *SD* = 34 hikers) disturbed days (*Mann-*

Whitney test statistic = 2,558, *p-value* = 0.006). Hikers also tended to arrive one hour earlier

during highly (*median* = 07:00 UTC, *SD* = 2 hours) compared to weakly (*median* = 08:00

UTC, *SD* = 3 hours) disturbed days although this difference was not statistically significant

(*Mann-Whitney test statistic* = 1,559, *p-value* = 0.188; Supporting Information 1).

248   Potential spatial exposure to hikers' disturbance was evaluated within each female

chamois home range using frequentation rates of the trails extracted from Strava Global

Heatmap (Strava, 2022; www.strava.com, download date: 2022/03/09, data from the two

previous years, see also Courbin *et al.,* 2022 for a similar approach on this study site). Strava

heatmap provides a value ranging from 0 for no attendance to 255 for the highest attendance

for each pixel (pixel size = 25 m x 25 m) and is a good proxy of relative human frequentation

(Supporting Information 3; Thorsen *et al.,* 2022; Venter, Gundersen, Scott & Barton, 2023).

The lowest and highest values recorded for a trail pixel in our study site were 51 and 255

respectively. We thus defined frequentation classes according to the quantiles of the [51; 255]

distribution in order to cover trail pixels exclusively: 1 = [51; 119] (between 0% and 33%

quantile, low frequentation); 2 =]119; 187] (between 33% and 66% quantiles, intermediate

frequentation); 3 =]187; 255] (between 66% and 100% quantiles, high frequentation). We

counted the number of trail pixels of each class in a given home range, and multiplied this

number by the corresponding class value (*i.e.,* 1, 2 or 3) to give a different weight to each

trail pixel according to its frequentation level. We then summed these values for a given

home range and normalised the total by dividing it by the home range surface to obtain our

spatial disturbance index. We used the median spatial disturbance index (*i.e.,* 0.00577) as a

threshold to discriminate between **highly exposed chamois** and **weakly exposed chamois**.

As our study period ran over almost three months, day length varied widely and is

known to influence chamois activity (Brivio *et al.*, 2016; Fattorini *et al.,* 2019). Similarly to

Semenzato *et al.* 2021, we accounted for this effect by including day length in our models.

We evaluated day length as the time elapsed between dawn and dusk using the *crepuscule*

function from the *maptools* package (Bivand & Lewin-Koh, 2022; reference angle of the sun

below the horizon = 6°, *i.e.,* civil dawn/dusk).

Statistical analyses: predicting activity patterns and calculating activity metrics

We first modelled the daily activity of chamois as a series of values of either 0 (inactive) or 1

(active) for each 5-minutes boot using a generalised additive mixed model (GAMM) with a

binomial distribution of errors (*mgcv* package, Wood, 2011; Wood, Goude, & Shaw 2015;

Wood, 2017). We included three fixed effects in the model: 1) *time of the day*, *i.e.,* exact time

every 5 minutes, and 2) *day length*, as well as their interaction (using the *ti* function dedicated

to interactions in generalised additive models), as continuous variables with a cyclic cubic

regression splines as smoothing basis for terms including *day length* exclusively; 3) a

synthesis 8-levels categorical variable representing all possible combinations between

temperature classes, levels of temporal disturbance and of spatial disturbance, in interaction

with the *time of the day* (using the *by* argument of the *s* function in the implementation of the

GAMM). The levels were ordered, with the reference level corresponding to a chamois living

in a weakly exposed area during a cold and weakly disturbed day. The interaction between

day length and *time of the day* allowed us to account for the fact that the activity pattern is

determined by the timing of sunrise and sunset (Ashby, 1972; Cederlund, 1989), which varies

widely according to the date. We defined the unique annual id-year ID of each chamois as a

random effect (using the parameter "re" for the *bs* argument in the *s* function in the

implementation of the GAMM) to account for inter-individual variations on circadian activity

pattern (Supporting Information 4). We set the maximum number of knots for the smoothing

functions to 24. We also fitted the null model, which was similar to the model described

above but excluded the *temperature-disturbance* variable, for comparison purposes. All

details about model fitting are provided in Supporting Information 5.

We used the predictions of the model to investigate the effect of temperature and hikers'

disturbance on several activity metrics derived from this model (described below). We

controlled for the effect of *day length* by fixing it to the median of the complete dataset (*i.e.,*

963 minutes) when predicting chamois activity for each level of the *temperature-disturbance*

combination. We derived the *total*, *diurnal*, *nocturnal* and both *morning* and *evening*

crepuscular activity as the area under the activity curve (*AUC* function of the *DescTools*

package; Signorell, 2023) for each level of the *temperature-disturbance* variable. We defined

daytime as the period elapsed between two hours after dawn and two hours before dusk, and

night-time as the period elapsed between two hours after dusk and two hours before dawn

(see Bonnot *et al.,* 2020 for a similar approach). Crepuscular periods corresponded to the

remaining periods between daytime and night-time. We assessed the *timing of the morning*

and *evening activity peaks* as the time when the highest activity probability occurs in the

morning and the evening, respectively. We finally evaluated *morning* and *evening peak*

duration as the period elapsed between the first time when the activity probability reached

0.5 and the last time before it went underneath 0.5 for both activity peaks (for a similar

approach see Bourgoin *et al.*, 2011). The percentage of variation between two situations was

calculated as $100*(value_2 - value_1)/value_1$. We performed all analyses using the R software

(R Core Development Team 2022, version R 4.3.0). Data is available from the Zenodo

Repository: https://zenodo.org/record/8123239.

Results

Model adjustment

The full generalised additive mixed model (*AIC* = 1,578,865, *explained deviance* = 11.4%,

adjusted R2 = 0.149) was better than the null model including *time of the day* and *day length*

effects and excluding the *temperature-disturbance* variable (*AIC* = 1,591,999, *explained*

deviance = 10.7%, *adjusted R2* = 0.140, Supporting Information 5) to predict activity pattern

(Fig. 1). All smooth terms (fixed and random effects) were significant (*p-values* < 0.001).

Figure 1: predicted circadian activity patterns (activity ranges between 0 for inactivity, to 1 for active) of 62 female northern chamois in the National Game and Hunting Reserve of Les Bauges massif, in summer (15th June to 7th September, 2004-2020), according to temperature (blue: cold day; red: hot day) and human disturbance: a) chamois living in a weakly exposed area during a weakly disturbed day; b) chamois living in a weakly exposed area during a highly disturbed day; c) chamois living in a highly exposed area during a weakly disturbed day; d) chamois living in a highly exposed area during a highly disturbed day (see also pictograms directly in the figure). Solid lines represent predicted values from the model, shaded areas around the curves represent 95% confidence intervals. Background shades represent the periods used to calculate the area under the activity curve for morning and evening crepuscules (grey), daytime (yellow) and night-time (blue; see also pictograms directly in the figure). Time of the day represents hours in Coordinated Universal Time (UTC).

Temperature

Whatever the level of spatial exposure and temporal disturbance, chamois increased their

total activity during hot days compared to cold days of 5% on average (Fig. 2). This change

corresponds to an average increase of 22% in the nocturnal activity and a decrease of 10% in

the diurnal activity, from cold to hot days (Fig. 2). In addition, crepuscular activity was

increased from cold to hot days by more than 31% in the morning and 18% in the evening on

average (Fig. 2). Chamois also advanced their morning peak by 17 minutes from cold to hot

days, and they only slightly delayed their evening peak by 5 minutes (Fig. 3). Morning and

evening peaks lasted longer during hot compared to cold days of almost 1 hour and of 20

minutes respectively (Fig. 4). Chamois maintained a certain amount of activity in the middle

of the day during cold days (*i.e.,* above 40%), whereas the activity probability at the same

hours was lower and more variable during hot days (Fig. 1).

Figure 2: Area Under Curve (AUC, unitless) normalised by the duration of each period for

each situation of interest, namely a combination of different levels of temperature (blue: cold

day; red: hot day), spatial (dot: low exposure; triangle: high exposure) and temporal (low:

week days; high: weekends; see also pictograms directly in the figure) disturbance: a) total

(24 hours), b) diurnal (12.08 hours), c) nocturnal (3.92 hours), d) morning crepuscular (4

hours) and e) evening crepuscular (4 hours) activity (see also pictograms directly in the

figure). AUC are calculated from predicted circadian activity patterns of northern chamois of

the National Game and Hunting Reserve of Les Bauges massif, in summer (June to

September, 2004-2020). We defined "diurnal" as the period elapsed between two hours after

dawn and two hours before dusk, "nocturnal" as the period elapsed between two hours after

dusk and two hours before dawn, and "crepuscule" as the remaining periods in the morning

and in the evening (see text for details).

Figure 3: Timing of the Activity Peak (APT, time of the day when the activity peaks happen,

i.e., the time when the highest probability of activity occurs) for each situation of interest,

namely a combination of different levels of temperature (blue: cold day; red: hot day), spatial

(dot: low exposure; triangle: high exposure) and temporal (low: week days; high: weekends;

see also pictograms directly in the figure) disturbance: a) morning peak and b) evening peak.

APT are calculated from predicted circadian activity patterns of northern chamois of National

Game and Hunting Reserve of Les Bauges massif, in summer (June to September, 2004-

2020). Note the change of interval on the y axis between figures, although the scale stays the

same.

Figure 4: Peak Duration (PD, in minutes) for each situation of interest, namely a combination

of different levels of temperature (blue: cold day; red: hot day), spatial (dot: low exposure;

triangle: high exposure) and temporal (low: week days; high: weekends; see also pictograms

directly in the figure) disturbance: a) morning and b) evening peak. PD are calculated from

predicted circadian activity patterns of northern chamois of National Game and Hunting

Reserve of Les Bauges massif, in summer (June to September, 2004-2020).

Hikers' disturbance

All the summer home ranges used by the studied chamois were crossed by at least one hiking

trail. Our spatial disturbance index ranged between 0.00317 and 0.02912 (*median* = 0.00577,

SD = 0.00433), with only seven chamois above 0.01000 (Supporting Information 1). Overall,

the effect of temperature was much more pronounced than the effect of disturbance and very

little difference between the general activity patterns was discernible between the different

levels of hikers' disturbance for a given temperature (Figs. 1 and 5). This translated by very

similar amounts of total, diurnal, nocturnal and crepuscular activity whatever the level of

disturbance (Fig. 2). Similarly, the changes in terms of timing of the activity peaks according

to hikers' disturbance was not biologically significant (< 7 minutes, Fig. 3). However, the

duration of the activity peaks during hot days seemed to be affected by the level of hikers'

disturbance (Fig. 4). Indeed, the morning activity peak decreased by 46 minutes from

chamois living in a weakly disturbed area during weakly disturbed days to chamois living in

a highly disturbed area during highly disturbed days. The evening activity peak increased by

almost 20 minutes for the same groups.

Figure 5: predicted circadian activity patterns (activity ranges between 0 for inactivity, to 1

for active) of female northern chamois in the National Game and Hunting Reserve of Les

Bauges massif, in summer (15th June to 7th September, 2004-2020), according to temperature

and human disturbance: a) cold day and b) hot day. In both panels, brown represents chamois

living in a weakly exposed area during a weakly disturbed day; green represents chamois

living in a weakly exposed area during a highly disturbed day; orange represents chamois

living in a highly exposed area during a weakly disturbed day; purple represents chamois

living in a highly exposed area during a highly disturbed day (see also pictograms directly in

the figure). Solid lines represent predicted values from the model, shaded areas around the

curves represent 95% confidence intervals. Background shades represent the periods used to

calculate the area under the curve for morning and evening crepuscules (grey), daytime

(yellow) and night-time (blue; see also pictograms directly in the figure). Time of the day

represents hours in Coordinated Universal Time (UTC).

Discussion

Our objective was to determine and compare the relative contribution of summer

temperatures and hikers' disturbance on the activity pattern of females of a wild ungulate, the

northern chamois. During hot days, chamois adjusted their circadian activity by shifting their

activity peak earlier in the morning and by increasing its duration. During hot days, they also

increased their nocturnal and crepuscular activity while being less active during the hottest

hours of daytime (in agreement with **H1** stating that chamois should adjust their circadian

activity pattern by avoiding being active during the hottest hours during the hottest days, with

a compensation of activity loss during daytime). Surprisingly, their total daily activity was

higher during hot than cold days. On the other hand, chamois only seldomly adjust their

activity pattern according to hikers' disturbance, disproving **H2** stating that chamois should

adjust their circadian activity pattern by avoiding being active during the most disturbed days.

Nevertheless, during hot days, we observed a decrease in the duration of the morning activity

peak and a slight increase in the duration of the evening activity peak from the less exposed

chamois to the most exposed chamois, as well as a tenuous increase in the nocturnal and

crepuscular activity at the expense of the diurnal activity. Hikers' disturbance had very little

effect on chamois activity peaks. These results provide only a partial support for **H3** stating

that the chamois the most exposed to hikers' disturbance should have a more marked

response.

435   Though chamois seem to be less sensitive to high temperatures than other alpine

ungulate species (see Darmon *et al.,* 2014 for a comparison between chamois and mouflon;

see also Semenzato *et al.* 2021 for Alpine ibex), their faecal glucocorticoid metabolites

concentration increases during drought conditions in summer (Anderwald, Campell Andri, &

Palme 2021), as well as the time they spend in thermal refuges during hot days (Loison *et al.,*

2022). This may explain that chamois responded much more strongly to temperature than to

hikers' disturbance. Moreover, in our study site, hikers' attendance was significantly higher

when temperature was high (*median* = 21, *SD* = 56 hikers during cold days; *median* = 54, *SD*

= 37 hikers during hot days; *Mann-Whitney test statistic* = 333, *p-value* = 0.007). Hence, the

behavioural changes to buffer from high temperatures may also protect chamois from hikers'

disturbance, as both factors are correlated. Animals have to face several sources of

disturbance (*e.g.,* human disturbance, predation, high temperatures) and they eventually

cannot indefinitely adjust their behavioural response as disturbance factors accumulate

(Benoist, Garel, Cugnasse, & Blanchard, 2013). Such a limitation seems to appear in our

population when chamois highly exposed to disturbance experience high hikers'

frequentation during hot days. Instead of maintaining a longer morning activity peak to

compensate for the loss of diurnal activity due to the heat, they reduce it likely because of

hikers already present on the trails (mainly between 08:00 UTC and 15:00 UTC, Supporting

Information 1).

Recent studies provided evidence of a non-negligeable nocturnal activity peak in

chamois (Carnevali, Lovari, Monaco, & Mori, 2016). Here, we also highlighted that

nocturnal activity significantly increased while diurnal activity decreased during hot days,

confirming the influence of high temperatures and human disturbance on chamois activity.

Similarly to numerous other species, chamois decreased their diurnal activity when exposed

to high temperatures and/or human disturbance and attempted to compensate for the activity

loss when it is absent (*e.g.,* later in the day or at night, Gaynor, Hojnowski, Carter, &

Brashares, 2018). However, an increase in nocturnal activity could expose individuals to

other threats, such as nocturnal predators (Bonnot *et al.,* 2020). This is particularly true in our

study site with the progressive return of wolves recorded since the recent years. Large

herbivores may face the difficult challenge of avoiding being exposed to high temperature

while limiting their exposure to human disturbance, but without shifting completely their

activity at night because of predation risk. The shift to more crepuscular activity observed in

this study could be an answer to this necessary trade-off.

Interestingly and in contrast with Brivio *et al.*, (2016) and Grignolio *et al.*, (2018) in

another chamois population, total activity was higher during hot than cold days in our

chamois population. Such a finding is not necessarily surprising as other studies have found

similar results (in the same population; Loison *et al.*, 2022; in Alpine ibex maintaining a

constant activity both during cold and hot days, Semenzato *et al.*, 2021). In addition, this does

not necessarily translate into an increase in feeding behaviour, but might be because chamois

display diel migrations between refuges and feeding spots (Courbin *et al.,* 2022), particularly

if both habitats are distant or heterogeneous. With the increasing temperatures in summer,

chamois could be forced to commute more frequently or move further to find shelter from the

heat while maintaining the same level of feeding activity, eventually leading to the overall

increase observed. Unfortunately, our data provides no clue about the nature of the

behaviours displayed (*i.e.,* foraging, moving or social behaviours) and we cannot ascertain if

this rise of activity effectively corresponds to spatial displacement.

We observed only minor differences between the most and least disturbed chamois,

either because chamois respond similarly to human presence regardless of the encounter risk,

because hikers are present almost every day, or because they shift their activity in habitats

further away from the main source of disturbance, *i.e.,* the trails. Conversely, Samia,

Nakagawa, Nomura, Rangel, & Blumstein (2015) showed that weakly exposed populations

can be less tolerant to human disturbance because they are less used to it. In our population,

85% of the individuals move temporarily away from trails every day when hikers are present

(Courbin *et al.,* 2022), demonstrating that even the less exposed chamois adjust their behaviour to human disturbance. This low tolerance affects the whole population and not just the most exposed individuals and might explain the weak difference observed between groups.

Mountain animals face both rising temperatures from ongoing climate warming, and modifications in their exposure to actual and perceived risks from the development of recreational activities and the return of predators in these landscapes. The interaction between those factors could eventually jeopardise their demography if particular attention is not paid in conserving habitats providing both refuge against summer heat and human disturbance (Frid & Dill, 2002; Chirichella, Stephens, Mason, & Apollonio, 2021). Moreover, both constraints can lead either to similar responses (*e.g.,* rest in refuge areas), or to opposite reactions (*e.g.,* reduce activity to avoid thermal stress or escape to limit predation risk), which complicates our understanding of their behavioural response. Here we highlight a strong effect of temperature but almost no effect of hikers' disturbance on the activity pattern of northern female chamois. Identifying the effect of human disturbance on wildlife remains a difficult task, particularly at the population scale (Tablado & Jenni, 2017), and an integrated approach is needed to consider all forms of response, such as variations in activity level and timing, behavioural budgets and spatial displacements. Our study contributes to filling the knowledge gap on the cumulative effects of weather and human disturbance on chamois circadian activity in this context of ongoing socio-environmental changes (Corlatti *et al.,* 2022).

References

Anderwald, P., Campell Andri, S., & Palme, R. (2021). Reflections of ecological differences?

Stress responses of sympatric Alpine chamois and red deer to weather, forage quality,

and human disturbance. Ecology and Evolution, 11(22), 15740–15753.

https://doi.org/10.1002/ece3.8235

Arlettaz, R., Patthey, P., Baltic, M., Leu, T., Schaub, M., Palme, R., & Jenni-Eiermann, S.

(2007). Spreading free-riding snow sports represent a novel serious threat for wildlife.

Proceedings of the Royal Society B: Biological Sciences, 274(1614), 1219–1224.

Ashby, K. (1972). Patterns of daily activity in mammals Mammal Review, 1, 171–185.

Beniston, M. (2003). Climatic change in mountain regions: a review of possible impacts.

Climate Variability and Change in High Elevation Regions: Past, Present & Future,

5–31.

Benoist, S., Garel, M., Cugnasse, J.-M., & Blanchard, P. (2013). Human disturbances, habitat

characteristics and social environment generate sex-specific responses in vigilance of

Mediterranean mouflon. PLoS One, 8(12), e82960.

Bivand, R., & Lewin-Koh, N. (2022). maptools: Tools for Handling Spatial Objects.

https://cran.r-project.org/package=maptools

Bonnot, N. C., Couriot, O., Berger, A., Cagnacci, F., Ciuti, S., De Groeve, J. E., Gehr, B.,

Heurich, M., Kjellander, P., Kröschel, M., & others. (2020). Fear of the dark?

Contrasting impacts of humans versus lynx on diel activity of roe deer across Europe.

Journal of Animal Ecology, 89(1), 132–145. https://doi.org/10.1111/1365-2656.13161

Bonnot, N., Morellet, N., Verheyden, H., Cargnelutti, B., Lourtet, B., Klein, F., & Hewison,

A. J. (2013). Habitat use under predation risk: hunting, roads and human dwellings

influence the spatial behaviour of roe deer. European Journal of Wildlife Research,

59(2), 185–193.

Bourgoin, G, Garel, M., Van Moorter, B., Dubray, D., Maillard, D., Marty, E., & Gaillard, J.-

M. (2008). Determinants of seasonal variation in activity patterns of mouflon.

Canadian Journal of Zoology, 86(12), 1410–1418.

Bourgoin, G., Garel, M., Blanchard, P., Dubray, D., Maillard, D., & Gaillard, J.-M. (2011).

Daily responses of mouflon (*Ovis gmelini musimon* x *Ovis sp.*) activity to summer

climatic conditions. Canadian Journal of Zoology, 89(9), 765–773.

https://doi.org/10.1139/Z11-046

Brivio, F., Bertolucci, C., Tettamanti, F., Filli, F., Apollonio, M., & Grignolio, S. (2016). The

weather dictates the rhythms: Alpine chamois activity is well adapted to ecological

conditions. Behavioral Ecology and Sociobiology, 70(8), 1291–1304.

Calenge, C. (2006). The package adehabitat for the R software: tool for the analysis of space

and habitat use by animals. Ecological Modelling, 197, 1035.

Carnevali, L., Lovari, S., Monaco, A., & Mori, E. (2016). Nocturnal activity of a diurnal

species, the northern chamois, in a predator-free Alpine area. Behavioural Processes,

126, 101–107.

Cederlund, G. (1989). Activity patterns in moose and roe deer in a north boreal forest.

Ecography, 12, 39–45.

Chirichella, R., Stephens, P. A., Mason, T. H. E., & Apollonio, M. (2021). Contrasting

effects of climate change on Alpine chamois. The Journal of Wildlife Management,

85(1), 109–120.

Cloud Parc des Bauges. (2014). Observatoire touristique du Parc Naturel Régional du Massif

des Bauges - Présentation 25 novembre 2014. Retrieved from

https://cloud.parcdesbauges.com/index.php/login. Accessed July 3, 2023

Clutton-Brock, T., Albon, S. & Guinness, F. (1989). Fitness costs of gestation and lactation in

wild mammals. Nature, 337, 260–262.

Corlatti, L., Iacolina, L., Safner, T., Apollonio, M., Buzan, E., Ferretti, F., Hammer, S. E.,

Herrero, J., Rossi, L., Serrano, E., & others. (2022). Past, present and future of

chamois science. Wildlife Biology, e01025. https://doi.org/10.1002/wlb3.01025

Courbin, N., Garel, M., Marchand, P., Duparc, A., Debeffe, L., Börger, L., & Loison, A.

(2022). Interacting lethal and nonlethal human activities shape complex risk tolerance

behaviors in a mountain herbivore. Ecological Applications, e2640.

https://doi.org/10.1002/eap.2640

Darmon, G., Bourgoin, G., Marchand, P., Garel, M., Dubray, D., Jullien, J.-M., & Loison, A.

(2014). Do ecologically close species shift their daily activities when in sympatry? A

test on chamois in the presence of mouflon. Biological Journal of the Linnean

Society, 111(3), 621–626. https://doi.org/10.1111/bij.12228

Duparc, A., Redjadj, C., Viard-Crétat, F., Lavorel, S., Austrheim, G., & Loison, A. (2013).

Co-variation between plant above-ground biomass and phenology in sub-alpine

grasslands. Applied Vegetation Science, 16(2), 305–316.

Fattorini, N., Brunetti, C., Baruzzi, C., Chiatante, G., Lovari, S., & Ferretti, F. (2019).

Temporal variation in foraging activity and grouping patterns in a mountain-dwelling

herbivore: Environmental and endogenous drivers. Behavioural Processes, 167,

103909.

French, S. S., González-Suárez, M., Young, J. K., Durham, S., & Gerber, L. R. (2011).

Human disturbance influences reproductive success and growth rate in California sea

lions (*Zalophus californianus*). PloS One, 6(3), e17686.

https://doi.org/10.1371/journal.pone.0017686

Frid, A., & Dill, L. (2002). Human-caused disturbance stimuli as a form of predation risk.

Conservation Ecology, 6(1). https://doi.org/10.5751/es-00404-060111

Gaillard, J.-M., Festa-Bianchet, M., Yoccoz, N. G., Loison, A., & Toigo, C. (2000).

584   Temporal variation in fitness components and population dynamics of large

585   herbivores. Annual Review of Ecology and Systematics, 31(1), 367–393.

586   https://doi.org/10.1146/annurev.ecolsys.31.1.367

587   Garel, M., Loison, A., Gaillard, J. M., Cugnasse, J. M., & Maillard, D. (2004). The effect of a

588   severe drought on mouflon lamb survival. Proceedings of the Royal Society of

589   London. Serie B: Biological Sciences, 27(suppl_6), S471–S473.

590   Gaynor, K. M., Hojnowski, C. E., Carter, N. H., & Brashares, J. S. (2018). The influence of

591   human disturbance on wildlife nocturnality. Science, 360(6394), 1232–1235.

592   https://doi.org/10.1126/science.aar7121

593   Gehr, B., Hofer, E., Pewsner, M., Ryser, A., Vimercati, E., Vogt, K. & Keller, L. F. (2018).

594   Hunting--mediated predator facilitation and superadditive mortality in a European

595   ungulate. Ecology and Evolution, 8, 109–119.

596   Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., & Stoffel, M. (2014). 21st

597   century climate change in the European Alps - A review. Science of the Total

598   Environment, 493, 1138–1151.

599   Greives, T. J., Kingma, S. A., Kranstauber, B., Mortega, K., Wikelski, M., van Oers, K.,

600   Mateman, A. C., Ferguson, G. A., Beltrami, G., & Hau, M. (2015). Costs of sleeping

601   in: circadian rhythms influence cuckoldry risk in a songbird. Functional Ecology,

602   29(10), 1300–1307.

603   Grignolio, S., Brivio, F., Apollonio, M., Frigato, E., Tettamanti, F., Filli, F. & Bertolucci, C.

604   (2018). Is nocturnal activity compensatory in chamois? A study of activity in a

605   cathemeral ungulate. Mammalian Biology, 93, 173-181.

606   Hamel, S., & Côté, S. D. (2008). Trade-offs in activity budget in an alpine ungulate:

607   contrasting lactating and nonlactating females. Animal Behaviour, 75(1), 217–227.

Kearney, M. R., Gillingham, P. K., Bramer, I., Duffy, J. P., & Maclean, I. M. D. (2020). A

method for computing hourly, historical, terrain-corrected microclimate anywhere on

Earth. Methods in Ecology and Evolution, 11(1), 38–43.

Kearney, M. R., & Porter, W. P. (2017). NicheMapR - an R package for biophysical

modelling: the microclimate model. Ecography, 40(5), 664–674.

Larson, C. L., Reed, S. E., Merenlender, A. M., & Crooks, K. R. (2016). Effects of recreation

on animals revealed as widespread through a global systematic review. PloS One,

11(12), e0167259.

Lesmerises, F., Déry, F., Johnson, C. J., & St-Laurent, M.-H. (2018). Spatiotemporal

response of mountain caribou to the intensity of backcountry skiing. Biological

Conservation, 217, 149–156.

Lima, S. L., & Bednekoff, P. A. (1999). Temporal variation in danger drives antipredator

behavior: the predation risk allocation hypothesis. The American Naturalist, 153(6),

649–659.

Lindstedt, S. L., & Boyce, M. S. (1985). Seasonality, fasting endurance, and body size in

mammals. The American Naturalist, 125(6), 873–878. https://doi.org/10.1086/284385

Loison, A., Jullien, J.-M., & Menaut, P. (1999). Subpopulation structure and dispersal in two

populations of chamois. Journal of Mammalogy, 80(2), 620–632.

Loison, A., Malagnino, A., Borger, L., Marchand, P., Courbin, N., Bonnot, N. & Morellet, N.

(2022). It's getting hot! Chamois activity patterns and habitat selection during

summer - Evidence of a functional response to temperature. 8[th] World Conference on

Mountain Ungulates.

Long, R. A., Martin, T. J., & Barnes, B. M. (2005). Body temperature and activity patterns in

free-living arctic ground squirrels. Journal of Mammalogy, 86(2), 314–322.

Lopez, J. F. (2001). Inventaire systématique de la flore et de la végétation du Massif des

Bauges.

Loupfrance. (2021). Situation du loup en France. Retrieved from

https://www.loupfrance.fr/suivi-du-loup/situation-du-loup-en-france/. Accessed July

3, 2023

Maclean, I. M. D., Mosedale, J. R., & Bennie, J. J. (2019). Microclima: An r package for

modelling meso-and microclimate. Methods in Ecology and Evolution, 10(2), 280–

290.

Marchand, P., Garel, M., Bourgoin, G., Dubray, D., Maillard, D., & Loison, A. (2014).

Impacts of tourism and hunting on a large herbivores spatio-temporal behavior in and

around a French protected area. Biological Conservation, 177, 1–11.

Marchand, P., Garel, M., Bourgoin, G., Dubray, D., Maillard, D., & Loison, A. (2015). Sex-

specific adjustments in habitat selection contribute to buffer mouflon against summer

conditions. Behavioral Ecology, 26(2), 472–482.

Pepin, N., Bradley, R. S., Diaz, H. F., Baraër, M., Caceres, E. B., Forsythe, N., Fowler, H.,

Greenwood, G., Hashmi, M. Z., Liu, X. D., & others. (2015). Elevation-dependent

warming in mountain regions of the world. Nature Climate Change, 5(5), 424–430.

Phillips, G. E., & Alldredge, A. W. (2000). Reproductive success of elk following

disturbance by humans during calving season. The Journal of Wildlife Management,

521–530.

Richard, Q., Toïgo, C., Appolinaire, J., Loison, A., & Garel, M. (2017). From gestation to

weaning: Combining robust design and multi-event models unveils cost of lactation in

a large herbivore. Journal of Animal Ecology, 86, 1497–1509.

Rogala, J. K., Hebblewhite, M., Whittington, J., White, C. A., Coleshill, J., & Musiani, M.

(2011). Human activity differentially redistributes large mammals in the Canadian

Rockies National Parks. Ecology and Society, 16(3).

Ruckstuhl, K. & Ingold, P. (1998). Baby-sitting in chamois: a form of cooperation in

females? Mammalia, 62, 125-128.

Samia, D. S. M., Nakagawa, S., Nomura, F., Rangel, T. F., & Blumstein, D. T. (2015).

Increased tolerance to humans among disturbed wildlife. Nature Communications,

6(1), 1–8.

Schmeller, D. S., Urbach, D., Bates, K., Catalan, J., Cogălniceanu, D., Fisher, M. C., Friesen,

J., Füreder, L., Gaube, V., Haver, M. & others (2022). Scientists' warning of threats to

mountains. Science of the Total Environment, 853, 158611.

Semenzato, P., Cagnacci, F., Ossi, F., Eccel, E., Morellet, N., Hewison, A. J. M., Sturaro, E.,

& Ramanzin, M. (2021). Behavioural heat-stress compensation in a cold-adapted

ungulate: Forage-mediated responses to warming Alpine summers. Ecology Letters,

24(8), 1556–1568.

Signorell, A. (2023). DescTools: Tools for Descriptive Statistics.

Strava. 2022. Strava Global Heatmap. https://www.strava.com/heatmap.

Tablado, Z., & Jenni, L. (2017). Determinants of uncertainty in wildlife responses to human

disturbance. Biological Reviews, 92(1), 216–233.

Thorsen, N. H., Bischof, R., Mattisson, J., Hofmeester, T. R., Linnell, J. D. C., & Odden, J.

(2022). Smartphone app reveals that lynx avoid human recreationists on local scale,

but not home range scale. Scientific Reports, 12(1), 1–13.

van Beest, F. M., & Milner, J. M. (2013). Behavioural responses to thermal conditions affect

seasonal mass change in a heat-sensitive northern ungulate. PloS One, 8(6), e65972.

Venter, Z., Gundersen, V., Scott, S. L., & Barton, D. N. (2023). Bias and precision of

crowdsourced recreational activity data from Strava. Landscape and Urban Planning,

232, 104686.

Watanuki, Y., & Nakayama, Y. (1993). Age difference in activity pattern of Japanese

monkeys: effects of temperature, snow, and diet. Primates, 34(4), 419–430.

Wheaton, B. (2010). Introducing the consumption and representation of lifestyle sports. Sport

in Society, 13(7–8), 1057–1081.

Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood

estimation of semiparametric generalized linear models. Journal of the Royal

Statistical Society (B), 73(1), 3–36.

Wood, S. N. (2017). Generalized Additive Models: An Introduction with R (2nd ed.).

Chapman and Hall/CRC.

Wood, S. N, Goude, Y., & Shaw, S. (2015). Generalized additive models for large data sets.

Journal of the Royal Statistical Society: Series C (Applied Statistics), 64(1), 139–155.

Worton, B. J. (1989). Kernel methods for estimating the utilization distribution in home-

range studies. Ecology, 70(1), 164–168.

Xu, W., Gong, Y., & Wang, H. (2021). Alert time reflects the negative impacts of human

disturbance on an endangered bird species in Changbai Mountain, China. Global

Ecology and Conservation, 28, e01709.

Acknowledgments

This study was funded by Agence Nationale de la Recherche (French National Research

Agency) (ANR) Grant Mov-It No.16-CE02-0010 coordinated by Anne Loison, Centre

National de la Recherche Scientifique (Centre National de la Recherche Scientifique)

(CNRS). The authors thank all participants for helping with the capture and marking of

animals, for the recovery of GPS collars, and for collecting GPS data on human activities:

volunteers and professionals from the Office Français de la Biodiversité (formerly Office

National de la Chasse et de la Faune Sauvage), the Office National des Forêts, the

Groupement d'Intérêt Cynégétique des Bauges, the Parc Naturel Régional du massif des

Bauges, the Observatoire Grande Faune et Habitats, and the laboratory Environnements

Dynamiques Territoires Montagnes (EDYTEM). This work was performed using the

computing facilities of the IFB Cloud.

Conflict of Interest

15　The authors declared they have no conflict of interest.

Data availability statement

18　Data available from the Zenodo Repository: https://zenodo.org/record/8123239.

Authors' contributions

LT, MG, PM, AL, GB conceived the ideas and designed methodology; MG, PM and AL organised data collection; LT analysed the data; LT led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.