MICA: A multi-omics method to predict gene regulatory networks in early human embryos

Gregorio Alanis-Lobato[1,*], Thomas E. Bartlett[2,*], Qiulin Huang[1,3*], Claire Simon[1], Afshan
McCarthy[1], Kay Elder[4], Phil Snell[4], Leila Christie[4] & Kathy K. Niakan[1,3,5,6]

1 Human Embryo and Stem Cell Laboratory, The Francis Crick Institute, London, UK.
2 Department of Statistical Science, University College, London WC1E 7HB, UK.
3 The Centre for Trophoblast Research, Department of Physiology, Development and
Neuroscience, University of Cambridge, Cambridge CB2 3EG, UK.
4 Bourn Hall Clinic, Bourn, Cambridge, CB23 2TN.
5 Wellcome – Medical Research Council Cambridge Stem Cell Institute, University of
Cambridge, Jeffrey Cheah Biomedical Centre, Puddicombe Way, Cambridge CB2 0AW, UK
6 Epigenetics Programme, Babraham Institute, Cambridge CB22 3AT, UK
*Equal contribution
Corresponding authors: kkn21@cam.ac.uk; g.alanis.lobato@gmail.com;
thomas.bartlett.10@ucl.ac.uk

Abstract

Recent advances in single-cell -omics have been transformative to characterise cell types in
challenging to study biological contexts. It is technically difficult to infer transcription factor-
gene regulatory interactions from these datasets, especially in contexts with limited single-cell
sample number such as the early human embryo. Here we systematically assessed the
application of four different linear or non-linear gene regulatory network (GRN) prediction
strategies to single-cell simulated and human embryo transcriptome datasets. We also
compared how gene expression normalisation methods impact on regulatory network
predictions, finding that TPM (transcripts per million reads) outperformed alternative methods.
We identified more reproducible GRN inferences using a non-linear method based on mutual
information (MI) applied to single-cell transcriptome datasets refined with chromatin
accessibility (CA) (called MICA), compared to alternative network prediction methods tested.
MICA captures complex non-monotonic dependencies and feedback loops. Using MICA, we
generated the first GRN inferences in early human development. MICA predicted co-
localisation of the AP-1 transcription factor subunit proto-oncogene JUND and the TFAP2C
transcription factor AP-2γ in human preimplantation embryos. These interactions were
supported by previous findings in other developmental and stem cell contexts. Overall, our
comparative analysis of gene regulatory network prediction methods defines a pipeline that
can be applied to single-cell multi-omics datasets in especially challenging contexts to infer
interactions between transcription factor expression and target gene regulation.

Introduction

Following the fusion of the oocyte and sperm, the zygote undergoes a series of cell divisions until it forms a blastocyst prior to implantation into the uterus. A human blastocyst is formed of a fluid-filled cavity and approximately 200 cells that comprise three distinct cell types: the trophectoderm (TE), which gives rise to fetal components of the placenta; the primitive endoderm (PE), which forms the yolk sac; and the pluripotent epiblast (EPI), which gives rise to the embryo proper (Blakeley et al., 2015). The specification of these three lineages represents the earliest cell-fate decisions in humans. Understanding the molecular mechanisms that regulate these decisions is important for applications including stem cell biology, regenerative medicine, and reproductive technologies (Niakan et al., 2012).

We do not yet understand how cell fate specification is regulated in the human embryo. Transcription factor (TF) and target gene regulatory interactions associated with cell fate specification in this context would be informative and has not been elucidated. Defining the gene regulatory networks (GRNs) associated with a given cell type at a distinct time in development facilitates characterisation of cell type identity and prediction of the transcriptional regulators and cis-regulatory DNA sequences that may underlie cell fate specification (Davidson and Erwin, 2006; Materna and Davidson, 2007; Peter and Davidson, 2011). Challenges to determining GRNs in early embryos include the small number of cells that contribute to the distinct cell lineages of the human embryo at these early stages of development. Moreover, while single cell -omics technologies facilitate the characterisation of genome-wide gene expression and chromatin accessibility changes in human blastocysts (Blakeley et al., 2015; Li et al., 2018; Liu et al., 2019; Petropoulos et al., 2016; Yan et al., 2013), it is unclear if resolution at this level would allow for accurate predictions of GRNs contributing to cell type specific identity.

The most common computational approach to infer TF-gene regulatory interactions is to model the expression of each target gene as either a linear or non-linear combination of the expression of a set of potential regulators (e.g., the TFs in the dataset of interest) (Dobra et al., 2004; Marbach et al., 2012). The 'target gene' method uses advanced regression methods, such as sparse penalised regression or random forests, to model the expression level of a particular 'target' gene (the response), conditional on the expression levels of a set of other genes (predictors, such as TFs) (Haury et al., 2012; Huynh-Thu et al., 2010). By combining these local network model fits genome-wide, with each gene taking a turn as the response or 'target gene' (Dobra et al., 2004), the GRN is constructed. The linear regression model has previously been used in combination with chromatin accessibility to infer GRNs in human brain organoids (Fleck et al., 2022). Bayesian networks also model the expression of each gene conditional on a set of regulating genes (i.e., parents in the network) by computing how likely it is that each TF in the dataset was responsible for the expression of a certain gene (Kamimoto et al., 2020; Marbach et al., 2012; Pe'er, 2005; Tipping, 2001). The Bayesian/Bagging Ridge model in combination with chromatin accessibility has been used effectively to predict the consequence of gene perturbations and to understand cell fate transitions in the context of development and cellular reprogramming (Argelaguet et al., 2022; Kamimoto et al., 2023). Of note, when fitting Bayesian models it is usually necessary to assume global compatibility (of the local network models), which in turn typically requires that the graph contains no loops (acyclicity). This an assumption which is challenging for biological systems, where feedback loops are a persistent feature (Brandman and Meyer, 2008). In another important group of methods, TF-gene (or more generally gene-gene) interactions are ranked based on a pairwise similarity measure of their expression across cell samples, such as a variant of correlation or mutual information (Daub et al., 2004; Krishnaswamy et al., 2014). Again, alternatives are available that consider conditional statistics (e.g., partial correlation or partial information decomposition) (Chan et al., 2017). Finally, a range of 'black-box' machine learning methods are available that model the expression of each gene in hard to visualise ways using for

94 example sophisticated non-linear combinations of TF expression values (Shu et al., 2021;
95 Wang et al., 2020).
96
Although GRN inference methods have been comprehensively compared and developed for
whole-tissue or bulk transcriptome analysis (Cahan et al., 2014; Chai et al., 2014; Marbach et
al., 2012; Morris et al., 2014; Thompson et al., 2015), the interest in cell type-specific
regulatory networks has prompted the application of these methods to single cell RNA-seq
(scRNA-seq) datasets (Aibar et al., 2017; Chen and Mar, 2018; Fiers et al., 2018; Iacono et
al., 2019; Kang et al., 2021; Mochida et al., 2018; Nguyen et al., 2021; Pratapa et al., 2020;
Stone et al., 2021). These previous studies showed that the approaches with the best overall
performance belong to the regression and mutual information categories and highlighted the
challenges of performing GRN inference on single cell datasets. This is attributed mainly to
the heterogeneous and stochastic nature of gene expression in individual cells (Nguyen et al.,
2021). Of note is that even state-of-the-art methods that have been evaluated on well-defined
benchmark datasets predict a considerable number of false positive interactions (Kang et al.,
2021; Marbach et al., 2012; Nguyen et al., 2021; Pratapa et al., 2020; Stone et al., 2021). Most
of these false predictions originate from indirect associations, for example a path $a \rightarrow b \rightarrow c$
can result in the prediction of $a \rightarrow c$ even if there is no direct link between those nodes (i.e., a
'transitive edge' in the network). Several methods to eliminate such effects have been
proposed (Barzel and Barabási, 2013; Chan et al., 2017; Feizi et al., 2013; Margolin et al.,
2006; Wang et al., 2018) but they can be computationally demanding and sensitive to the
choice of hyper-parameters (Feizi et al., 2013).
116
Given the stage of development of human preimplantation embryos and their precious nature,
together with the restrictions on such research in some countries, the -omics datasets from
human blastocysts are very small compared to those from other biological contexts. This
makes it challenging to mine these datasets using GRN inference methods, which require a
sufficiently large number of cells to produce reproducible results (Kang et al., 2021; Pratapa
et al., 2020). This tight restriction on sample sizes places corresponding restrictions on
statistical power, and means that the optimal statistical network inference methodology may
be specific to this context. Here, we assessed whether the integration of other types of -omics
datasets with transcriptomic-based predictions could help to reduce indirect TF-gene
relationships and thereby produce more reliable gene regulatory networks in this setting,
especially with restricted sample sizes. We utilised a low number of cell samples in high quality
-omics data from early human embryos at the blastocyst stage (6-7 days post-fertilisation)
(Blakeley et al., 2015; Petropoulos et al., 2016; Yan et al., 2013) to evaluate the plausibility of
our predictions: this has the advantage of being a biological context with only three well-
defined cell types to evaluate the plausibility of our predictions.
132
The aims of this work were (i) to evaluate whether it is possible to infer reliable cell type-
specific regulatory networks for each one of the cell types in the early human embryo in spite
of the size of the available -omics data, (ii) to determine if the integration of chromatin
accessibility data with transcriptome analysis would better inform predictions of GRNs, and
(iii) to predict and validate previously unidentified gene regulations in the human blastocyst.
Overall, we demonstrate that the available single cell transcriptomic data was most robustly
analysed by a non-linear mutual information-based inference method which had been refined
with chromatin accessibility data (MICA). The resulting analysis predicted the first GRN in
human preimplantation development and showed that the interactions were consistent with
the transcripts and proteins that are known to be enriched in specific lineages. MICA predicted
a novel putative regulatory interaction between the TFAP2C transcription factor AP-2γ and the
AP-1 transcription factor subunit proto-oncogene JUND in human preimplantation embryos.
Overall, we propose that MICA will be an informative method to make gene regulatory network
predictions in other challenging-to-study biological contexts with limitations in the number of
cells that can be analysed.
148

Results

Assessing the conditions for optimal GRN inference on synthetic data

We generated synthetic gene-expression data with known ground-truth GRN structure for a variety of sample sizes (see Methods) to determine the characteristics of the transcriptomic data on which GRN inference methods perform best and, importantly, to compare these conditions with our human blastocyst scRNA-seq datasets (see Fig. 1A). We then applied four different GRN prediction strategies to the simulated transcriptomes (see Methods for more details). 1) We applied GENIE3 (Aibar et al., 2017; Huynh-Thu et al., 2010), a random forests regression approach which takes into account expression levels of the regulated or target gene and is used for regulatory network inference in the SCENIC GRN prediction pipeline (Aibar et al., 2017; Kang et al., 2021; Marbach et al., 2012; Nguyen et al., 2021; Pratapa et al., 2020; Stone et al., 2021). 2) We compared this to the target gene approach using the L0L2 sparse penalised regression model (Hazimeh and Mazumder, 2018; Tibshirani, 1996) which minimizes the total size of the linear model coefficients so that a minimal set of regulatory TFs receive the largest linear model coefficients. 3) We also compared with the correlation coefficient as a reference: we used Spearman's rank correlation coefficient (Fieller et al., 1957) as we expect non-linear associations in the data. 4) Lastly, we applied a non-linear alternative measure of pairwise association based on mutual information (MI) that we speculated may be more appropriate for non-linear response functions in single-cell data (Faith et al., 2007; Krishnaswamy et al., 2014) (Fig. S1A). For the MI method, we used the empirical distribution of the MI values for each gene (Faith et al., 2007). We also varied the number of transcriptional regulators of each target gene in the synthetic data, to study the impact of this parameter on their performance. Further details of the analysis methods and step-by-step code are also available at: https://github.com/galanisl/early_hs_embryo_GRNs

Two main conclusions emerged from these simulations. First, using synthetic datasets, the performance of all methods increases from the sample size of $n = 10$ to $n = 1000$, as assessed statistically by the area under the receiver operating characteristic curve (AUROC) (Fig. 1B). However, we noted that the prediction accuracy of the GRNs only marginally improves as the number of cell samples surpasses $n = 100$ (Fig. 1B). Therefore, while sample size is important, increasing this beyond a threshold sample size does not further improve inference of the predicted transcriptional regulations. We further note that low sequencing depth (such as in 10x Genomics single-cell studies) may increase this optimal value of n. Second, limiting the number of potential transcriptional regulators of a target gene positively impacts the ability of a chosen inference method to recover the ground-truth GRN (Figs. 1B,C). From our simulations, 50 or fewer TFs yield the highest AUROC values for $n < 100$ (Fig. 1B). This would be a reasonable prediction for a given gene in a specific cell type based on our analysis of low input chromatin accessibility (CA) data from the human blastocyst (Liu et al., 2019), where the median number of TF motifs per gene is 35 for the inner cell mass (ICM) and 40 for the TE (Fig. S1B).

Based on these results and given the size of our human blastocyst datasets (Fig. 1C EPI: 26 cells, PE: 33 cells, TE: 45 cells with 1,366 TFs among 25,098 genes (Blakeley et al., 2015; Petropoulos et al., 2016; Yan et al., 2013)), we reasoned that TF-gene interactions predicted from experimentally collected gene expression data could be refined with complementary, context-specific epigenomic datasets. Specifically, we used low-input CA analysis of human blastocyst TE cells and the inner cell mass (ICM), comprised of EPI and PE cells, to refine the GRNs of the respective cell types with putative cis-regulatory interactions (Liu et al., 2019). Peak calling and annotation were performed with nf-core/atacseq to identify regions of open chromatin (Ewels et al., 2020). TF motif enrichment analysis in these open regions of open chromatin were identified using rgt-hint with TF binding models from HOCOMOCO and JASPAR (Li et al., 2019). Finally, the open chromatin regions enriched for TF motifs along with the predicted downstream target genes were determined based on the nearest transcriptional

204 start sites (most distances ranging from -5kbp to 10kbp). This narrowed down the number of
205 genes (to 12,780 for EPI and PE, 12,981 for TE) and TFs (514) in the datasets, thus bringing
206 our sample sizes and potential TFs per gene to the ranges suggested by our simulations.
207
208
209 **Statistical evaluation of inferred human blastocyst GRNs**
210
We next integrated the single-cell transcriptome data (Fig. S1C) from human blastocysts
generated using the SMART-seq2 library preparation protocol (Blakeley et al., 2015;
Petropoulos et al., 2016; Picelli et al., 2014; Yan et al., 2013) where fewer cells are sequenced
but better transcript coverage and sequencing depth is obtained (~15,000 genes and ~7
million reads per cell) compared to more conventional scRNA-seq methods, such as 10x
Genomics (~3,000 genes and ~50,000 reads per cell) (Wang et al., 2021). Due to the lack of
genome-scale experimentally derived GRNs in the human embryo context, we evaluated the
robustness of the predictions made by four inference approaches (L0L2 regression, GENIE3,
Spearman correlation and MI) using three different strategies (see Methods). Since we did not
find reports about the impact of gene expression normalisation choice on GRN predictions,
we also assessed this parameter and considered the application of GRN predictors to
log(TPM+1), log(FPKM+1), log-count and batch-corrected expression data.
223
We calculated a reproducibility score R for each putative regulatory interaction following the
application of the network inference methods to human blastocyst data (see Methods) and
investigated the distribution of the reproducibility values for the top-100,000 predicted edges
(Figs. 2A and S2). The reproducibility estimator R estimates the posterior probability of seeing
a network edge given the data; it quantifies the robustness or stability of the inference of this
network edge. The accuracy quantified by the R reproducibility statistic relates to the stability
of the model predictions to perturbation of the data. Fig. 2A reports the difference between the
median of this distribution and the median of the distribution produced by a predictor that
generates a random ranking of all possible TF-gene interactions (ΔR). We found that the most
reproducible regulatory interactions were inferred by MI followed by a filtering process in which
only TF-gene associations supported by CA data were considered in the final network (MICA)
(Fig. 2A). As predicted by our simulations, the size of each cell type-specific dataset had a
clear impact on GRN inference with methods such as GENIE3 producing more reproducible
interactions in the TE. Interestingly, most inference methods produced better results with gene
expression values following log(TPM+1) or log(FPKM+1) normalisation (Fig. 2A).
239
We also assessed robustness at the level of network structures, features, or subnetworks. To
do so, we randomly split each dataset (EPI, PE and TE) into two groups with the same number
of cells, then applied the inference methods to each one (with or without CA refinement), and
finally benchmarked them either focussing on the top regulatory interactions (early
recognition) or on all regulatory interactions (see Methods). We repeated this 10 times for 10
random splits of the data. For the first comparison, we identified the top 1% (by network score)
of all possible regulatory interactions from each group, and then quantified the overlap of these
top interactions between the groups using the area under the precision-recall curve (AUPRC)
(Figs. 2B and S3). By comparing the differences between the median of ten resulting AUPRCs
for each method to the random predictions described above, we found that MICA
outperformed the other network prediction approaches and gave the highest difference in
AUPRC when applied to log(TPM+1)- and log(FPKM+1)-normalised expression values (Fig.
2B). For the second comparison, we divided the data into two portions and then calculated a
normalised L2 loss between the network scores over the whole network, rather than just the
top 1% of interactions (Figs. 2C and S4). We found that the inverse value of this metric also
confirms the robustness of MICA (Figs. 2C and S4).
256
Overall, our statistical evaluation strategies showed that the most robust GRN inference
method to analyse the limited number of human blastocyst cells was MICA. Importantly, if

transcriptome-based predictions are not refined with CA data, most methods perform just
slightly better than random (Figs. 2 and S2-S4), underscoring the importance of integrating
multi-omics analysis in inference models. It was also important to assess the impact of gene
expression normalisation on GRN prediction because we observed apparent effects in our
benchmarks depending on the normalisation method used, with log(TPM+1) or log(FPKM+1)
being the most suitable gene expression units in this context.
265
266
267 **Association of inferred GRNs to human blastocyst cell lineages**
268
We next evaluated the inferred GRNs to determine if they could recapitulate interactions of
molecular markers of the three cell types that comprise the human blastocyst. First, we
computed the overlap between the GRN edges predicted by each inference method for each
blastocyst cell type to identify interactions associated with the EPI, PE and TE, as well as the
interactions common to the three cell types (see Methods). We then used the out-degree of
NANOG, GATA4 and CDX2, TFs which respectively mark the EPI, PE and TE (Figs. 3A,B),
as a proxy for their activity in each one of the four networks. Our prediction is that these TFs
should be actively regulating genes in the cell type-specific networks and participate in only a
few interactions in the common GRN, based on their known expression pattern in the
blastocyst and function in other mammalian contexts such as the mouse (Arceci et al., 1993;
Chambers et al., 2003; Dietrich and Hiiragi, 2007; Mitsui et al., 2003; Niakan and Eggan, 2012;
Roode et al., 2012; Strumpf et al., 2005). The common network did not show marker activity
in the predicted GRNs (Figs. 3B and S5-S8), which is expected given that the expression of
these markers is known to be mutually exclusive at this stage. The expected pattern for
NANOG (active in the EPI and inactive in the PE and TE, Fig. 3A), was observed in the GRNs
with TF-gene interactions supported by CA data and inferred with GENIE3, Spearman
correlation and MI when applied to batch-corrected, FPKM and TPM data with some instances
of the non-refined L0L2, GENIE3 and Spearman GRNs also matching the expected pattern
(Fig. 3B). In the PE, the expected activity pattern for GATA4 (active in the PE and inactive in
EPI and TE, Fig. 3A) and lack of detectable networks for NANOG and CDX2, were predicted
by applying GENIE3+CA, Spearman correlation+CA and MICA to the FPKM and TPM
datasets (Fig. 3B). Finally, the TE-expected pattern for CDX2 (active in the TE and inactive in
the EPI and PE) and lack of detectable network for NANOG and GATA4, Fig. 3A) was only
observed when using Spearman correlation or MICA on log-counts and batch-corrected data,
MICA or MICA+CA on FPKM or TPM data (Fig. 3B). L0L2, with or without CA refinement,
consistently performed poorly at predicting GRNs compared to the other methods used. Log-
count normalisation failed to recover cell type specific GRNs for these TFs.
296
We next manually curated a list of gene sets representing the most relevant pathways and
biological processes in the EPI (e.g., regulation of pluripotency) and the TE (e.g., embryonic
placenta development) to perform a gene set enrichment analysis with the genes involved in
the 500 regulatory interactions with the best prediction scores in the EPI and TE GRNs of the
25 most active TFs (Tables S1, S2, Methods). We restricted this analysis to the EPI and TE,
because similar lists of gene-sets were not available for PE, where it is currently unclear which
pathways are most relevant to this cell type. Next, we computed a validation score V as the
number of relevant gene sets that were enriched at a significance level of 10% ($p<0.1$) in the
EPI and TE GRNs (Fig. S9). We note that a low significance level was set because the
resulting inferences are aggregated, mitigating the effect of false positive hits for individual
gene groups. To facilitate the comparison between V scores across GRNs from different
inference and normalisation methods, we normalised this score to the maximum score
attained at the cell type level (Fig. 3C). We found that the CA-refined data agree better with
the cell type-specific gene sets for both the top-predicted EPI and TE interactions, i.e. the
interactions with the highest prediction scores (Fig. 3B). The GRNs inferred by Spearman+CA
produced the highest V scores in the EPI and TE, and these were not impacted by the
normalisation method employed. The second-best inference method was MICA, with

consistent *V* scores across normalisation methods. We note that both MICA and Spearman+CA are non-linear methods which do not involve regression, indicating that advanced regression methods may not be the most effective choice for biological discovery with these restricted sample-sizes. Taken together, both biological evaluation metrics that we considered confirmed the importance of refining single-cell transcriptome-based GRN inferences with CA data and underscored the robustness of MICA in predicting GRNs from the analysed human blastocyst datasets. Based on these results, we decided to focus on the GRNs predicted by this method for the analyses presented in the following sections.

Predicted TF networks for NANOG, GATA4 and CDX2

Using MICA, we constructed GRNs for TFs expressed in the EPI, PE and TE. All MICA GRN predictions can be found on FigShare: doi.org/10.6084/m9.figshare.21968813. For visualization, the predicted GRN for each of the TFs is separated into target and regulator TF networks. Target networks contain a maximum of 25 top potential target TFs of the hub (or central) TF, whereas regulator networks include a maximum of 25 top TFs that potentially regulate the hub. The average expression of the network members across samples of the cell type of interest is represented by the size of the node. MI scores are represented by the thickness of the edges in the network and edge colour highlights. To further refine the MICA-predictions we used the Spearman's rank correlation coefficient between the expression levels of the source and target nodes across samples of the cell type of interest to define correlated or anti-correlated expression. Correlated or anti-correlated node pairs correspond to positive or negative Spearman's rank coefficient with p-value smaller than 0.1, whereas node pairs having p-value equals or larger than 0.1 were defined as uncorrelated.

Among the top NANOG targets, TFs RREB1, NCAO3, ZNF343, ZFP42, and NME2 are predicted to be positively regulated by NANOG (Fig. 4A). ZFP42 is a pluripotency marker encoding the REX1 protein and has been shown to be a direct target of NANOG in mouse pluripotent stem cells (Shi et al., 2006). ZNF343 is a less well-characterized TF, but multiple NANOG ChIP-seq datasets in both naïve and primed human embryonic stem cells (ESCs) showed high binding score (MACS2 score > 1000) in the proximal region of the transcription start site (TSS) of ZNF343 (Fig. S10A; (Barakat et al., 2018; Chovanec et al., 2021; Lyu et al., 2018)), which suggests direct regulation of ZNF343 by NANOG. Interestingly, NME2 was previously predicted to be a regulator instead of a target of NANOG in mouse pluripotent stem cells using the TENET GRN inference method (Kim et al., 2021). This inconsistency may be due to the lack of chromatin accessibility data for robust directionality inference in the TENET method, or a species difference. Indeed, some mouse naïve pluripotency regulators, such as ESRRB, which is also a direct target of NANOG in mouse pluripotent stem cells (Festuccia et al., 2012), were not expressed in the human EPI (Blakeley et al., 2013). Putative regulators of NANOG predicted from our MICA network analysis contain multiple KLF factors, including KLF3, KLF5, KLF9 and KLF16 (Fig. 4B). Spearman correlation analysis suggest that KLF9 and KLF16 potentially down-regulate NANOG expression, based on their anti-correlated expression, whereas KLF3 and KLF5 are identified as potential regulators of NANOG by MICA, but not significantly correlated by Spearman correlation.

In the PE, GATA4 was predicted to positively regulate the expression of SP8, TET1, and SKIL while repressing the expression of ELF3, TFDP2 POGK, ZNF770, and NR2F2 (Fig. 4C). It is also predicted to be a target of HNF1B and SALL4 and repressed by ETV4 and E2F6 (Fig. 4D). These interactions have not been experimentally validated or inferred. However, NR2F2 was previously identified as a maturation marker of polar TE (Meistermann et al., 2021). The repression of NR2F2 by GATA4 predicted in the PE suggests a role for GATA4 in maintaining the PE cell identity by inhibiting the polar TE program. Furthermore, it has been shown that SALL4 is required for mouse PE-derived extra-embryonic endoderm cell derivation and knockout of SALL4 in these cells cause down-regulation of GATA4 (Lim et al., 2008). In

369 addition, multiple endoderm genes such as GATA4, GATA6, and SOX17 were shown to be
370 SALL4-bound genes by ChIP-seq (Lim et al., 2008). These findings are consistent with the
371 inferred GATA4 regulatory network.
372
Interestingly, TBX3 is a target of CDX2 (Fig. 4E), and the role of TBX3 has been implicated in
trophoblast cell differentiation (Lv et al., 2019). CDX2 is predicted to be regulated by both
KLF5 and KLF6, which are also molecular markers of the trophectoderm lineage (Fig. 4F).
KLF5 is necessary for trophectoderm formation in the mouse pre-implantation embryo and is
required for CDX2 expression in mouse ESCs (Ema et al., 2008; Lin et al., 2010). Interestingly,
further analysis of the CDX2 interactions showed that TBX3 and FOXH1 are also potential
positive regulators of KLF5 and KLF6. Positive feedback loops are predicted between CDX2,
TBX3 and KLF5, as well as CDX2, FOXH1 and KLF6 (Fig. S10B). Other key developmental
regulators including HAND1, GATA3 and GATA6, are also predicted to regulate CDX2
expression (Fig. 4F). This is consistent with the timing of GATA3 protein expression preceding
that of CDX2 (Gerri et al., 2020). Further investigation will be needed to understand the
differences between the CDX2-high and CDX2-low TE cells and how the positive feedback
loops formed within the CDX2 network enhance and stabilize CDX2 expression in the CDX2-
high population.
387
388
389 **Maintenance of TFAP2C expression by JUND in all lineages of human blastocyst**
390
We next sought to determine if the MICA network modelling predicts novel interactions or
associations between TFs in early human embryo development. We focused on TFAP2C as
an example for the network comparison. TFAP2C is a molecular marker that is initially
expressed in all the cells at the morula stage in mouse embryos and later specifically restricted
to the TE at the blastocyst stage and it is not expressed in other lineages (Cao et al., 2015;
Gerri et al., 2020). However, in human embryos, TFAP2C is expressed in the TE, EPI and PE
at the blastocyst stage (Blakeley et al., 2015) and has been shown to maintain pluripotency in
naïve human embryonic stem cells by regulating OCT4 expression (Chen et al., 2018; Pastor
et al., 2018). By performing comparisons of TFAP2C networks in human EPI, TE, and PE
cells, we found a conserved putative interaction consisting of TFAP2C, JUND, SOX4, and
GCM1 (Fig. 5A). In addition, all four factors showed significant positive correlations in their
expression (Figs. 5B, S11). MICA predicts that JUND and SOX4 regulate TFAP2C, while
TFAP2C targets GCM1 in all three lineages. In the EPI and PE, SOX4 and GCM1 formed a
feedback loop. Interestingly, in the TE, the correlation between GCM1 and SOX4 is absent,
and SOX4 is predicted to target JUND, which in turn may regulate TFAP2C. Our network
predicts that interactions between these four TFs maintains TFAP2C expression in all three
lineages.
408
To determine if TFAP2C and JUND protein expression is positively correlated in embryos, as
predicted by MICA, we performed immunofluorescence analysis of human preimplantation
blastocysts approximately 6.5 days after fertilisation. We observed that TFAP2C and JUND
were co-expressed in cells of the human blastocysts (Fig. 5C, S13A; n=4). We next performed
3D nuclear segmentation and calculated the Pearson correlation coefficient between TFAP2C
and JUND based on the DAPI-normalized protein intensity (Fig. 5D). TFAP2C intensity
showed a linear increase with the increase of JUND intensity in all analysed human embryos,
and their intensities are highly correlated (0.68-0.85 Pearson correlation; $p < 0.001$; Fig. 5D).
Overall, this demonstrates the informativeness of the MICA network analysis to predict
correlations and possible regulatory relationships between transcription factors that can be
experimentally tested.
420
In addition to the MICA predictions, we also performed Spearman correlation analysis of the
significant edges identified by the MICA analysis. Surprisingly, around half of the interactions
are in non-linear fashion, highlighting the informativeness of MICA to capture complex non-

monotonic dependencies. From the TFAP2C network predicted in the EPI lineage, we found
that TFAP2C potentially regulates FOXO3 and ZFP42 in a non-linear manner (Fig. S12). The
expression of FOXO3 and ZFP42 seems to fit better with the expression of TFAP2C on an
exponential curve rather than a linear line between TFAP2C and JUND (Fig. S13B). This
suggests that when analysing scRNAseq or low input multi-omics analyses similar non-linear
correlations may exist and this may have biological significance. It would be interesting to
know whether and which types of non-linear interaction predominate and the biological
significance of these non-linear regulations.
432
433
434 **Discussion**
435
The relationship between genome-wide transcriptomic and epigenomic changes and cell fate
specification in human embryogenesis is unclear. Studies of human pre-implantation
development rely on the donation of surplus embryos derived from assisted reproduction
technologies, and the use of such embryos for research is tightly regulated and subject to
significant limitations, such as a lack of ability to conduct such research in some jurisdictions
(Niakan et al., 2012). In addition, the collection of single cells from such precious embryos is
technically challenging and requires specialist expertise and micropipettes to disaggregate
microscopic embryos. Therefore, currently available -omics datasets from human blastocysts
comprise only a few tens of samples per cell type and are therefore very limited. This contrasts
with single-cell analyses in other cellular and developmental contexts that are based on tens
of thousands of samples (or more) per cell type (Zheng et al., 2017).
447
Sample size is one of the most important considerations when selecting or designing statistical
methodology, for example to infer networks of regulations of TFs and their target genes.
Hence, GRN inference in -omics data from human pre-implantation embryos presents unique
statistical challenges. In particular, methodology that can leverage information about gene
regulations from small sample-sizes is required for this context. On the other hand, the lack of
heterogeneity in early human embryos compared to adult tissue makes this a good context in
which to assess GRN inference methodologies, because there is less unmeasured variability
arising from environmental factors. To assess whether GRN inference method can be
informative in this challenging-to-study context, we have systematically compared several
popular methodologies. Furthermore, we have tested how incorporation of complementary
cis-regulatory epigenomic data from ATAC-seq improves GRN inferences. Consistent with
other contexts (Argelaguet et al., 2022; Kamimoto et al., 2020), we found that incorporating
chromatin accessibility/TF motif analysis together with transcriptional inferences improves the
accuracy of GRN inference, by first narrowing down the choice of TF targets from which to
infer the mRNA transcript co-expression network, a principle that is likely to be especially
important with small sample sizes. Notably, we analyse RNA-seq data generated using
SMART-seq2 and the sequencing method and depth of sequencing will likely impact on the
choice of GRN inference method.
466
Here we showed that incorporating complementary epigenomic data with transcriptomic data
improves the reproducibility of inferred GRNs. Furthermore, it has enabled us to make
predictions about GRNs operational in early human embryos that are consistent with an
understanding of the function and association of the regulators in other developmental and
stem cell contexts. We suggest that for network inference using advanced regression
methods, it may be preferable to pre-filter using epigenomic data to first narrow down the TF
targets in scRNA-seq datasets, because this gives the network inference algorithm an easier
time of selecting the regulators from transcriptome data, which is consistent with the
experience of others using alternative GRN prediction methods (Fleck et al., 2022; Kamimoto
et al., 2020; Kartha et al., 2022) . Alternative methods incorporate epigenomic data after
inference of the mRNA transcript co-expression network (González-Blas et al., 2022) and this
has been applied successfully to infer GRNs in the fly brain (Janssens et al., 2022). It will be

interesting to determine how the type and sequential order of incorporating multi-omics
datasets impacts on GRN predictions.
481
We also note some limitations on the interpretation of the GRNs predicted. For example, one
ATAC-seq peak could cover multiple transcription factor binding sites in a region. In such
cases, we include all TFs with motifs mapping to this region as potential regulators of the gene
with the closest TSS (transcriptional start site) to the region, for subsequent transcript co-
expression network modelling. However, it will be interesting to determine if this can be
refined, by applying the recently developed chromVAR-Multiome method to human
blastocysts to generate an *in silico* ChIP-seq library for this context (Argelaguet et al., 2022).
Moreover, the ATAC-seq data applied in this study analysed the ICM data in bulk, without
distinguishing between EPI and PE cells. Therefore, cell-type specific chromatin accessibility
could not be considered, and specific interactions may have been missed due to the
heterogeneity of the data or if the ICM CA data failed to reflect developmentally cis-regulations
of more developmentally progressed EPI and PE cells. In addition, predictions based on the
nearest TSS will miss long-range enhancers that are known to be important for gene regulation
(Schoenfelder and Fraser, 2019). In the future, integration of single-cell or nuclei matched
transcriptome and ATAC-seq chromatin accessibility data, like recent studies in the mouse
(Argelaguet et al., 2022), would be preferable to apply in the human blastocyst context. We
note that the GRN interference methods predicted edges that overlap between the four
inference methods. Identifying overlapping inferences by comparing more than one GRN
inference method may be a strategy to identify network edges with more confidence due to
the agreement between several inference methodologies. However, this strategy may also
miss some edges which can only be detected by one method and not another.
503
We also note that so far, we have separately modelled GRN structure to specific pre-
implantation embryonic cell types constrained to a single developmental time-point. In the
future, we seek to model dynamic GRN structure in transcriptionally distinct human blastocyst
lineages. Moreover, we seek to integrate CUT&RUN or CUT&Tag TF-DNA binding analysis
(Kaya-Okur et al., 2019; Meers et al., 2019; Skene and Henikoff, 2017) for key putative
developmental regulators, such as GATA3, to further narrow down experimentally validated
occupancy from the ATAC-seq predictions we used in this study, similarly to a recent
application in mouse blastocysts (Hainer et al., 2019; Hayashi and Inoue, 2023), though we
note that in this context TF occupancy studies will be restricted to a few TFs with good quality
antibodies. It will also be important to determine which cis regulatory regions are required for
target gene regulation through systematic perturbation studies.
515
516 In summary, the MICA network analysis pipeline we developed is a tool that can be applied to
517 challenging-to-study developmental contexts with limited sample size, such as the human
518 blastocyst, to make predictions about TF interactions that can be experimentally tested in the
519 future. As more datasets become available, we anticipate that the networks predicted will be
520 further refined.
521
522
523 **Methods**
524
525 **Ethics statement**
526
527 All human embryo experiments followed all relevant institutional and national guidelines and
528 regulations.
529
530 This study was approved by the UK Human Fertilisation and Embryology Authority (HFEA):
531 research licence number R0162, and the Health Research Authority's Research Ethics
532 Committee (Cambridge Central reference number 19/EE/0297).
533

The process of licence approval entailed independent peer review along with consideration by
the HFEA Licence and Executive Committees. Our research is compliant with the HFEA code
of practice and has undergone inspections by the HFEA since the licence was granted.
Research donors were recruited from patients at Bourn Hall Clinic, Homerton University
Hospital, The Bridge Centre and IVF Hammersmith.

Informed consent was obtained from all couples that donated surplus embryos following IVF
treatment. Before giving consent, people donating embryos were provided with all of the
necessary information about the research project, an opportunity to receive counselling and
the conditions that apply within the licence and the HFEA code of practice. Donors were
informed that, in the experiments, embryo development would be stopped before 14 days
post-fertilization, and that subsequent biochemical and genetic studies would be performed.
Informed consent was also obtained from donors for all the results of these studies to be
published in scientific journals. No financial inducements were offered for donation. Consent
was not obtained to perform genetic tests on patients and no such tests were performed. The
patient information sheets and consent document provided to patients are publicly available
(https://www.crick.ac.uk/research/a-z-researchers/researchers-k-o/kathy-niakan/hfea-
licence/). Donated embryos surplus to the IVF treatment of the patient were cryopreserved
and were transferred to the Francis Crick Institute where they were thawed and used in the
research project.

ATAC-seq data processing and analysis

Chromatin accessibility profiles from the inner cell mass (ICM) and TE were obtained from the
data produced by Liu *et al.* with the LiCAT-seq protocol (Liu et al., 2019). Alignment to the
reference genome (GRCh38), peak calling and annotation were performed with nf-
core/atacseq v1.1.0 (Ewels et al., 2020). Then, we carried out footprinting followed by TF motif
enrichment analysis in the regions of open chromatin using rgt-hint v0.13.0 with TF binding
models from HOCOMOCO and JASPAR (Li et al., 2019). Finally, we associated the TFs that
exhibited over-represented motifs in these regions with their closest transcription starting sites
(most distances ranging from -5kbp to 10kbp).

scRNA-seq data processing and analysis

We integrated scRNA-seq data from three different studies (Blakeley et al., 2015; Petropoulos
et al., 2016; Yan et al., 2013) focusing on the three cell types present at the late blastocyst
stage (Fig. 1, Fig. S3A). Cell type annotations were taken from the work by Stirparo *et al*.
Alignment to the reference genome (GRCh38) and calculation of gene counts and TPM-
normalised counts were performed on each dataset separately with nf-core/rnaseq v1.4.2
(Ewels et al., 2020). The resulting gene expression matrices were integrated and normalised
(log-counts and batch-corrected counts) using Bioconductor tools (Amezquita et al., 2019).
The final set of cells was manually curated based on the UMAP representation of the batch-
corrected data (see Fig. S3 and Table S1). The list of 25,098 genes in the expression matrices
was reduced to the unique set of TFs and transcription starting sites (TSS) derived from the
motif enrichment analysis applied to the ATAC-seq data. We used the ICM TFs and TSS for
the EPI and PE matrices (12,780 genes) and the TE TFs and TSS for the TE matrices (12,981
genes).

Network inference methods

For network inference, we compared the best performing strategy in the DREAM5 challenge,
GENIE3 (Huynh-Thu et al., 2010), with a non-linear alternative based on MI (Faith et al., 2007),
the Spearman's rank correlation coefficient, and the L0L2 sparse regression method that we
applied using recent advances in sparse multivariate statistical modelling (Bartlett et al., 2019;
Hazimeh and Mazumder, 2018), as follows.

589
For GENIE3, transcription factors are ranked according to the degree of variability in their
expression level and how the expression of the putative regulator correlates with a target gene.
This ranking is then used to construct the co-expression network for all genes and transcription
factors, by thresholding the algorithm's variance reduction score at the tenth percentile of its
empirical distribution. The GRN is then inferred as the intersection of the edges in this co-
expression network with the edges in the network of all possible gene regulations derived from
the chromatin accessibility / DNA binding motif data. This network of all possible gene
regulations is defined as all network edges from regulating TF to regulated gene, where an
edge represents a DNA binding motif for the TF in regulatory DNA within open chromatin in
the regulated gene. For GENIE3, all default settings were used.

600
601 With the Spearman correlation coefficient, a weighted co-expression network is inferred as
602 the absolute value of the correlation coefficient. The gene regulatory network is then inferred
603 as the intersection of the edges in this co-expression network with the edges in the network of
604 all potential gene regulations derived from the chromatin accessibility / DNA binding motif
605 data.

606
For L0L2 regression (described in more detail below), the model automatically chooses a
ranked subset of TFs from those predicted as regulators of the target gene in the chromatin
accessibility / DNA binding motif data inferences. In this way, each target gene takes its turn
for a model to be fitted around that target gene. After the model has been fitted to every target
gene, the global gene regulatory network can be constructed by combining the local networks
fitted around each target gene. For L0L2 regression, sparsity hyperparameters were chosen
using the L0Learn package's internal cross-validation.

614
In more detail, for L0L2 regression we start with a linear model of the expression level y of the
regulated target gene (Dobra et al., 2004), in terms of the expression levels $x_1, x_2, \ldots , x_p$, of
p transcription factors. We want to use the size of the fitted model coefficients b_1, b_2, etc. to
measure the strength of regulation of the target gene by transcription factors (TFs) 1, 2, etc.
We use sparse regression to find b_1, b_2, etc., as this specifically minimises the number of non-
zero model coefficients, by requiring that the coefficients b_q are set to zero as much as
possible. This leads to a more parsimonious model, in which a relatively small number $p' \ll p$
of transcription factors is inferred to be regulating the target gene, as a result of non-zero b_q.
Sparse regression minimizes:

624
625 $$[\, y - (a + b_1x_1 + b_2x_2 + \ldots + b_px_p)\,]^2 + \psi, \tag{1}$$

626
where ψ 'penalises' models with values of b_q $(q=1,\ldots,p)$ further from zero. The most popular
choices for ψ include $\psi = \lambda \Sigma\, b_j^2$, which is called 'ridge regression' (L2 regression), and $\psi = \lambda$
$\Sigma\, |b_j|$, which is called 'the lasso' (L1 regression). It can be shown that Eq.1 is equivalent to
applying the constraint $\Sigma\, b_j^2 < t^2$ for $\psi = \lambda \Sigma\, b_j^2$ (L1 regression) or $\Sigma\, |b_j| < t$ for $\psi = \lambda \Sigma\, |b_j|$ (L0
regression). Then, 'best-subset regression' (L0 regression) specifies the number of
transcription factors that can have non-zero b_j. It does this by using the constraint $\Sigma b_j^0 \leq k$,
which is equivalent to $\psi = \lambda \Sigma\, b_j^0$ in Eq.1. Combinations of these constraints are also often
more effective, such as L1L2 regression (also called 'the elastic net' (Zou and Hastie, 2005),
which was very successful in genomics), with penalty term $\psi = \gamma \Sigma\, |b_j| + \lambda \Sigma\, b_j^2$. Recently,
L0L2 regression has been proposed as an improvement (Hazimeh and Mazumer, 2018), with
penalty term $\psi = \gamma \Sigma\, b_j^0 + \lambda \Sigma\, b_j^2$. We use L0L2 regression in this study, for reasons as follows.
Sparse regression using the L0 penalty is an ideal model for inferring a minimal set of
regulating transcription factors, because it specifically selects the best set of k transcription
factors for the model (i.e., the 'best subset', of the available transcription factors). Combining
with the L2 penalty in L0L2 makes the model better specified for the data, by minimizing the
total size of the linear model coefficients so that the most important TFs receive the largest
linear model coefficients. We use sparse L0L2 regression to infer the best subset of regulators

of each target gene, from the full list of TF-gene associations supported by the chromatin
accessibility data.
646
647 For MI-based inference, the co-expression network is estimated first, as follows. We use an
648 empirical estimate of the distribution of the MI values for each gene (Daub et al., 2004; Faith
649 et al., 2007). Writing the estimated MI between the expression levels of genes x and y as
650 M_{xy}, we calculate the approximately $N(0,1)$ distributed variable z_{xy} according to the equation:
651
652 $$Z_{xy}^2 = F^{-1}[\Phi_{y|x}(M_{xy})]^2 + F^{-1}[\Phi_{x|y}(M_{xy})]^2 ,$$
653
where F is the $N(0,1)$ cumulative distribution function (c.d.f.), and $\Phi_{x|y}$ and $\Phi_{y|x}$ are the empirical
c.d.f.s of M_{xy} conditioned on x and y, respectively. From these $N(0,1)$ distributed variables z_{xy},
we calculate p-values, and then threshold at a false discovery rate of 5%. Again, the gene
regulatory network is inferred as the intersection of the edges in this co-expression network
with the edges in the network of all possible gene regulations derived from the chromatin
accessibility / DNA binding motif data.
660
661 We also considered a "random predictor" that outputs a random ordering of all the possible
662 TF-gene interactions.
663
The putative regulatory links predicted by each of these methods using the scRNA-seq data
were evaluated *as is* but also subjected to a filtering process in which only TF-gene
associations supported by the chromatin accessibility data were considered in the final
network (for GENIE3, MI and Spearman correlation methods). In the case of L0L2 sparse
regression, only TF-gene associations supported by the chromatin accessibility data were
considered as potential regulators of the target gene in the regression model. To identify these
refined predictions, we added the *+CA* suffix to the name of the GRN inference methods. In
both cases, to generate the final network, we selected the top 100,000 edges by ranking edges
according to their network score. We define network score as the absolute correlation
coefficient or linear regression coefficient (in the case of Spearman's correlation and sparse
regression respectively), or as -log10p for mutual information, or the GENIE3 score.
675
676 **Simulation study**
677
For our simulation study, we used the `daggitty` package in `R` to generate synthetic gene
expression data based on pre-defined GRNs with pre-determined network structure. These
GRNs were generated with network edge density of ρ=0.07, i.e., 7% of all possible edges (or
gene regulations), are present in the network (this value was estimated from the available
ATAC-seq+TF motif data). The synthetic datasets were generated with a range of sample size
n and number of potential regulators of each node (gene) p, using linear models, as follows.
For each combination of n and p, we generated 100 GRNs; then corresponding observed
gene-expression data-sets were generated for each of the GRNs by specifying that the
expression level of a downstream gene should depend on a linear combination of the
expression levels of its upstream regulators:
688
689 $$x_d = b_1{}^*x_{u1} + b_2{}^*x_{u2} + b_3{}^*x_{u3} + \ldots b_g{}^*x_{ug} + e,$$
690
where x_d is the expression level of the regulated (downstream) gene, and $x_{u1 \ldots} x_{up}$ are the
expression levels of the g regulating (upstream) TFs, b_1, $b_2 \ldots b_g$ are linear model coefficients,
and e is random noise. The different network inference methods (GENIE3, L0L2 sparse
regression, MI, Spearman correlation) were then applied to these generated data, and the
results were compared with the known pre-determined network structure. This comparison
was assessed by the AUC statistic, the 'area under the ROC (receiver-operator characteristic)
curve'.
698

Statistical evaluation metrics

We calculated a reproducibility score R as a bootstrap estimate of the posterior probability of observing an edge E given the dataset D, $R=P(E|D)$ (Pe'er, 2005), for the top 100,000 predicted edges. We also carried out 2-fold cross-validation in two ways as follows. We randomly split each dataset (EPI, PE and TE) in two groups with the same number of cells ten times (repeated 2-fold cross-validation), applied the inference methods (with or without chromatin accessibility refinement), took the top-1% of all possible regulatory interactions from one group and quantified the extent to which the top interactions inferred from the other group coincide with that reference using the area under the precision-recall curve (AUPRC). In addition, we calculated a normalised L2 loss comparing all the network inferences from the two network fits on the 2 folds of the data (at the level of the network scores). The normalised L2 loss is then defined as the L2 loss comparing the network scores from each of the 2 data folds, divided by the product of the square roots of the L2 norms of the network scores of each of the 2 data folds.

Biological evaluation metrics

For each combination of inference method and normalisation approach, we computed the intersection between the GRN edges predicted to identify interactions that are specific to the EPI, PE and TE, as well as the interactions common to the three cell types. In set notation this corresponds to EPI-specific = (EPI \ PE) \ TE, PE-specific = (PE \ EPI) \ TE, TE-specific = (TE \ EPI) \ PE and Common = EPI ∩ PE ∩ TE. Then, we used the out-degree of NANOG, GATA4 and CDX2 (marker TFs of the EPI, PE and TE, respectively) as a proxy for their activity in each one of the four networks. We considered that a marker was active if its normalised out-degree (i.e., the proportion of all genes it regulates in that GRN) was at least the median normalised out-degree across all networks.

We calculated a validation score V as the number of relevant gene sets that were enriched at a significance level of 10% ($p<0.1$) in a gene-set enrichment analysis performed with the genes regulated by the 25 most connected TFs and that were part of the 500 interactions with the highest prediction scores produced by the GRN inference methods for the EPI and TE. The list of gene sets for the EPI and TE were manually curated and represent the most relevant pathways and biological processes in each one of these cell types (see Tables S1 and S2). The starting list of gene sets comprised all the pathways and Gene Ontology Biological Process (GO BP) terms collated by the Bader Lab at the University of Toronto on 01 October 2020 (http://download.baderlab.org/EM_Genesets/October_01_2020/Human/symbol/Human_GO BP_AllPathways_no_GO_iea_October_01_2020_symbol.gmt). To facilitate the comparison between V scores from different GRN inference methods within a cell type, we normalised V to the maximum score attained by a network predictor for that cell type (i.e., EPI or TE).

Human Embryo Culture

Human embryos were cultured as previously described (Gerri et al., 2020). Vitrified human blastocyst stage embryos were thawed using Kitazato Thawing Media (VT602, order number 91121) following the manufacturer's instructions.

Human embryos were cultured in drops of pre-equilibrated Global medium (LifeGlobal; LGGG-20) supplemented with 10% human serum albumin (LifeGlobal; GHSA-125) and overlaid with mineral oil (Origio; ART-4008-5P). Preimplantation embryos were incubated at 37 °C and 5% CO_2 in an EmbryoScope+ time-lapse incubator (Vitrolife) and cultured for up to 24hrs for human blastocyst.

Immunofluorescence staining

754
Embryos were fixed with freshly prepared 4% paraformaldehyde in PBS. Fixation was
performed for 20 min at room temperature for embryos. Embryos were then washed 3 times
in 1× PBS with 0.1% Tween-20 to remove residual paraformaldehyde. Permeabilization was
performed with 1× PBS with 0.5% Tween-20 and followed by blocking in blocking solution
(10% donkey serum in 1× PBS with 0.1% Tween-20) for 1 h at room temperature on a rotating
shaker. Then, antibody incubation was performed with primary antibodies diluted in blocking
solution overnight at 4 °C on rotating shaker. The following day, embryos and cell cultures
were washed in 1× PBS with 0.1% Tween-20 for 3 times, and then incubated with secondary
antibodies diluted in blocking solution for 1 h at room temperature on a rotating shaker in the
dark. Next, embryos and cell cultures were washed in 1× PBS with 0.1% Tween-20 for 3 times.
Finally, embryos were placed in 1× PBS with 0.1% Tween-20 with Vectashield and DAPI
mounting medium (Vector Lab; H-1200) (1:30 dilution). Embryos were placed on µ-Slide 8-
well dishes (Ibidi; 80826) for confocal imaging. The antibodies and concentrations used are
reported in Table S3.
769
770 **Confocal imaging**
771
772 Confocal immunofluorescence images were taken with Leica SP8 confocal microscope. 2-µm-
773 thick optical sections were collected for embryo Z-stack imaging.
774
775 **Nuclei segmentation and quantification**
776
Nuclei segmentation and quantification were performed on Z-stack confocal images taken at
2-µm-thick optical sections. Stardist (Weigert et al., 2020) was used for nuclei segmentation
followed by CellProfiler (Stirling et al., 2021) for nuclei tracking and fluorescence intensity
quantification based on a customized pipeline modified from Lea et al., 2021. First, LIF format
confocal images were exported into multi-channel Z-stack TIF format for each Z-stack image
using ImageJ (Schneider et al., 2012). Then, Stardist was used to identify the nuclei based on
the DAPI channel (358 nm). Finally, Stardist outputs were split into single-channel single-stack
images and loaded into the CellProfiler v4.2.5 to track nuclei across image slices and quantify
the fluorescence intensity. Customized pipeline and scripts can be found here:
https://github.com/galanisl/early_hs_embryo_GRNs
787
788 **Data and code availability**
789

794
795
796 **Acknowledgements**
797

We thank Zimeng Chen for contributing to the initial analysis of the synthetic datasets. We
thank members of the Niakan lab for helpful discussions and feedback on the manuscript.
Work in the laboratory of KKN was supported by the Wellcome (221856/Z/20/Z) and the
Wellcome Human Developmental Biology Initiative (215116/Z/18/Z). Work in the laboratory of
KKN was also supported by the Francis Crick Institute, which receives its core funding from
Cancer Research UK (CC2074), the UK Medical Research Council (CC2074) and the
Wellcome Trust (CC2074). The work carried out by TEB was partly supported by MRC grant
(MR/P014070/1). For the purpose of Open Access, the authors have applied a CC BY public
copyright licence to any Author Accepted Manuscript version arising from this submission.

807
808

Competing Interests

No competing interests declared.

Author contributions

GAL, TEB and KKN conceived the project. GAL and TEB developed the analytical network interference and statistical methods and analysed data. QH developed the network model representation pipeline, analysed data and performed immunostaining analysis and quantification. AM performed human embryo thaw and culture. CS provided advice on image quantification and analysed data. KKN analysed data and funded the project. KE, PS and LC coordinate the donation of embryos to the research project. GAL, TEB, QH, CS and KKN wrote the manuscript.

References

Aibar, S., González-Blas, C. B., Moerman, T., Huynh-Thu, V. A., Imrichova, H., Hulselmans, G., Rambow, F., Marine, J.-C., Geurts, P., Aerts, J., et al. (2017). SCENIC: single-cell regulatory network inference and clustering. *Nature Methods* 14, 1083–1086.

Amezquita, R. A., Lun, A. T. L., Becht, E., Carey, V. J., Carpp, L. N., Geistlinger, L., Marini, F., Rue-Albrecht, K., Risso, D., Soneson, C., et al. (2019). Orchestrating single-cell analysis with Bioconductor. *Nature Methods* 17, 137–145.

Arceci, R. J., King, A. A., Simon, M. C., Orkin, S. H. and Wilson, D. B. (1993). Mouse GATA-4: a retinoic acid-inducible GATA-binding transcription factor expressed in endodermally derived tissues and heart. *Mol Cell Biol* 13, 2235–2246.

Argelaguet, R., Lohoff, T., Li, J. G., Nakhuda, A., Drage, D., Krueger, F., Velten, L., Clark, S. J. and Reik, W. (2022). Decoding gene regulation in the mouse embryo using single-cell multi-omics. *bioRxiv* 2022.06.15.496239.

Barakat, T. S., Halbritter, F., Zhang, M., Rendeiro, A. F., Perenthaler, E., Bock, C. and Chambers, I. (2018). Functional Dissection of the Enhancer Repertoire in Human Embryonic Stem Cells. *Cell Stem Cell* 23, 276-288.e8.

Bartlett, T. E., Kosmidis, I. and Silva, R. (2019). Two-way sparsity for time-varying networks, with applications in genomics. *arXiv* arXiv:1802.08114.

Barzel, B. and Barabási, A.-L. (2013). Network link prediction by global silencing of indirect correlations. *Nature Biotechnology* 31, 720–725.

Blakeley, P., Fogarty, N. M. E., del Valle, I., Wamaitha, S. E., Hu, T. X., Elder, K., Snell, P., Christie, L., Robson, P. and Niakan, K. K. (2015). Defining the three cell lineages of the human blastocyst by single-cell RNA-seq. *Development* 142, 3151–3164.

Brandman, O. and Meyer, T. (2008). Feedback Loops Shape Cellular Signals in Space and Time. *Science* 322, 390–395.

Boroviak, T., Loos, R., Lombard, P., Okahara, J., Behr, R., Sasaki, E., Nichols, J., Smith, A., Bertone, P. (2015). Lineage-specific profiling delineates the emergence and progression of naïve pluripotency in mammalian embryogenesis. *Developmental Cell* 35, 366–82.

Butte, A. J. and Kohane, I. S. (1999). Mutual information relevance networks: functional genomic clustering using pairwise entropy measurements. In *Biocomputing 2000*, pp. 418–429. Honolulu, Hawaii, USA: WORLD SCIENTIFIC.

Cahan, P., Li, H., Morris, S. A., Lummertz da Rocha, E., Daley, G. Q. and Collins, J. J. (2014). CellNet: Network biology applied to stem cell engineering. *Cell* 158, 903–915.

Cao, Z., Carey, T. S., Ganguly, A., Wilson, C. A., Paul, S. and Knott, J. G. (2015). Transcription factor AP-2γ induces early Cdx2 expression and represses HIPPO signaling to specify the trophectoderm lineage. *Development* 142, 1606–1615.

Chai, L. E., Loh, S. K., Low, S. T., Mohamad, M. S., Deris, S. and Zakaria, Z. (2014). A review on the
computational approaches for gene regulatory network construction. *Comput Biol Med* 48,
55–65.

Chambers, I., Colby, D., Robertson, M., Nichols, J., Lee, S., Tweedie, S. and Smith, A. (2003).
Functional expression cloning of Nanog, a pluripotency sustaining factor in embryonic stem
cells. *Cell* 113, 643–655.

Chan, T. E., Stumpf, M. P. H. and Babtie, A. C. (2017). Gene Regulatory Network Inference from
Single-Cell Data Using Multivariate Information Measures. *Cell Systems* 5, 251-267.e3.

Chen, S. and Mar, J. C. (2018). Evaluating methods of inferring gene regulatory networks highlights
their lack of performance for single cell gene expression data. *BMC Bioinformatics* 19, 232.

Chen, D., Liu, W., Zimmerman, J., Pastor, W. A., Kim, R., Hosohama, L., Ho, J., Aslanyan, M., Gell, J. J.,
Jacobsen, S. E., et al. (2018). The TFAP2C-Regulated OCT4 Naive Enhancer Is Involved in
Human Germline Formation. *Cell Rep* 25, 3591-3602.e5.

Chovanec, P., Collier, A. J., Krueger, C., Várnai, C., Semprich, C. I., Schoenfelder, S., Corcoran, A. E.
and Rugg-Gunn, P. J. (2021). Widespread reorganisation of pluripotent factor binding and
gene regulatory interactions between human pluripotent states. *Nat Commun* 12, 2098.

Daub, C. O., Steuer, R., Selbig, J. and Kloska, S. (2004). Estimating mutual information using B-spline
functions - an improved similarity measure for analysing gene expression data. *BMC
Bioinformatics* 5, 118.

Davidson, E. H. and Erwin, D. H. (2006). Gene regulatory networks and the evolution of animal body
plans. *Science* 311, 796–800.

Dietrich, J.-E. and Hiiragi, T. (2007). Stochastic patterning in the mouse pre-implantation embryo.
Development 134, 4219–4231.

Dobra, A., Hans, C., Jones, B., Nevins, J. R., Yao, G. and West, M. (2004). Sparse graphical models for
exploring gene expression data. *Journal of Multivariate Analysis* 90, 196–212.

Ema, M., Mori, D., Niwa, H., Hasegawa, Y., Yamanaka, Y., Hitoshi, S., Mimura, J., Kawabe, Y., Hosoya,
T., Morita, M., et al. (2008). Krüppel-like factor 5 Is Essential for Blastocyst Development and
the Normal Self-Renewal of Mouse ESCs. *Cell Stem Cell* 3, 555–567.

Ewels, P. A., Peltzer, A., Fillinger, S., Patel, H., Alneberg, J., Wilm, A., Garcia, M. U., Di Tommaso, P.
and Nahnsen, S. (2020). The nf-core framework for community-curated bioinformatics
pipelines. *Nature Biotechnology* 38, 276–278.

Faith, J. J., Hayete, B., Thaden, J. T., Mogno, I., Wierzbowski, J., Cottarel, G., Kasif, S., Collins, J. J. and
Gardner, T. S. (2007). Large-Scale Mapping and Validation of Escherichia coli Transcriptional
Regulation from a Compendium of Expression Profiles. *PLoS Biology* 5, e8.

Feizi, S., Marbach, D., Médard, M. and Kellis, M. (2013). Network deconvolution as a general method
to distinguish direct dependencies in networks. *Nature Biotechnology* 31, 726–733.

Festuccia, N., Osorno, R., Halbritter, F., Karwacki-Neisius, V., Navarro, P., Colby, D., Wong, F., Yates,
A., Tomlinson, S. R. and Chambers, I. (2012). Esrrb Is a Direct Nanog Target Gene that Can
Substitute for Nanog Function in Pluripotent Cells. *Cell Stem Cell* 11, 477–490.

Fieller, E. C., Hartley, H. O. and Pearson, E. S. (1957). Tests for rank correlation coefficients. I.
Biometrika 44, 470–481.

Fiers, M. W. E. J., Minnoye, L., Aibar, S., Bravo González-Blas, C., Kalender Atak, Z. and Aerts, S.
(2018). Mapping gene regulatory networks from single-cell omics data. *Brief Funct Genomics*
17, 246–254.

Fleck, J. S., Jansen, S. M. J., Wollny, D., Zenk, F., Seimiya, M., Jain, A., Okamoto, R., Santel, M., He, Z.,
Camp, J. G., et al. (2022). Inferring and perturbing cell fate regulomes in human brain
organoids. *Nature*.

Gerri, C., McCarthy, A., Alanis-Lobato, G., Demtschenko, A., Bruneau, A., Loubersac, S., Fogarty, N.
M. E., Hampshire, D., Elder, K., Snell, P., et al. (2020). Initiation of a conserved
trophectoderm program in human, cow and mouse embryos. *Nature* 587, 443–447.

González-Blas, C. B., De Winter, S., Hulselmans, G., Hecker, N., Matetovici, I., Christiaens, V.,
Poovathingal, S., Wouters, J., Aibar, S. and Aerts, S. (2022). SCENIC+: single-cell multiomic
inference of enhancers and gene regulatory networks. *bioRxiv* 2022.08.19.504505.

Hainer, S. J., Bošković, A., McCannell, K. N., Rando, O. J. and Fazzio, T. G. (2019). Profiling of
Pluripotency Factors in Single Cells and Early Embryos. *Cell* 177, 1319-1329.e11.

Haury, A.-C., Mordelet, F., Vera-Licona, P. and Vert, J.-P. (2012). TIGRESS: Trustful Inference of Gene
REgulation using Stability Selection. *BMC Systems Biology* 6, 145.

Hayashi, R. and Inoue, A. (2023). Low-Input CUT&RUN for Mouse Oocytes and Preimplantation
Embryos. *Methods Mol Biol* 2577, 83–92.

Hazimeh, H. and Mazumder, R. (2018). Fast Best Subset Selection: Coordinate Descent and Local
Combinatorial Optimization Algorithms. *arXiv* arXiv:1803.01454,.

Huynh-Thu, V. A., Irrthum, A., Wehenkel, L. and Geurts, P. (2010). Inferring Regulatory Networks
from Expression Data Using Tree-Based Methods. *PLoS ONE* 5, e12776.

Iacono, G., Massoni-Badosa, R. and Heyn, H. (2019). Single-cell transcriptomics unveils gene
regulatory network plasticity. *Genome Biology* 20, 110.

Janssens, J., Aibar, S., Taskiran, I. I., Ismail, J. N., Gomez, A. E., Aughey, G., Spanier, K. I., De Rop, F. V.,
González-Blas, C. B., Dionne, M., et al. (2022). Decoding gene regulation in the fly brain.
Nature 601, 630–636.

Kamimoto, K., Stringa, B., Hoffmann, C. M., Jindal K., Solnica-Krezel, L., and Morris, S. A. (2023).
Dissecting cell identity via network inference and in silico gene perturbation. *Nature* 614,
742-751.

Kang, Y., Thieffry, D. and Cantini, L. (2021). Evaluating the Reproducibility of Single-Cell Gene
Regulatory Network Inference Algorithms. *Front. Genet.* 12, 617282.

Kartha, V. K., Duarte, F. M., Hu, Y., Ma, S., Chew, J. G., Lareau, C. A., Earl, A., Burkett, Z. D., Kohlway,
A. S., Lebofsky, R., et al. (2022). Functional inference of gene regulation using single-cell
multi-omics. *Cell Genom* 2, 100166.

Kaya-Okur, H. S., Wu, S. J., Codomo, C. A., Pledger, E. S., Bryson, T. D., Henikoff, J. G., Ahmad, K. and
Henikoff, S. (2019). CUT&Tag for efficient epigenomic profiling of small samples and single
cells. *Nat Commun* 10, 1930.

Kim, J., T. Jakobsen, S., Natarajan, K. N. and Won, K.-J. (2021). TENET: gene network reconstruction
using transfer entropy reveals key regulatory factors from single cell transcriptomic data.
Nucleic Acids Research 49, e1–e1.

Krishnaswamy, S., Spitzer, M. H., Mingueneau, M., Bendall, S. C., Litvin, O., Stone, E., Pe'er, D. and
Nolan, G. P. (2014). Conditional density-based analysis of T cell signaling in single-cell data.
Science 346, 1250689.

Li, L., Guo, F., Gao, Y., Ren, Y., Yuan, P., Yan, L., Li, R., Lian, Y., Li, J., Hu, B., et al. (2018). Single-cell
multi-omics sequencing of human early embryos. *Nature Cell Biology* 20,.

Li, Z., Schulz, M. H., Look, T., Begemann, M., Zenke, M. and Costa, I. G. (2019). Identification of
transcription factor binding sites using ATAC-seq. *Genome Biology* 20,.

Lim, C. Y., Tam, W.-L., Zhang, J., Ang, H. S., Jia, H., Lipovich, L., Ng, H.-H., Wei, C.-L., Sung, W. K.,
Robson, P., et al. (2008). Sall4 Regulates Distinct Transcription Circuitries in Different
Blastocyst-Derived Stem Cell Lineages. *Cell Stem Cell* 3, 543–554.

Lin, S.-C. J., Wani, M. A., Whitsett, J. A. and Wells, J. M. (2010). Klf5 regulates lineage formation in
the pre-implantation mouse embryo. *Development* 137, 3953–3963.

Liu, L., Leng, L., Liu, C., Lu, C., Yuan, Y., Wu, L., Gong, F., Zhang, S., Wei, X., Wang, M., et al. (2019). An
integrated chromatin accessibility and transcriptome landscape of human pre-implantation
embryos. *Nature Communications* 10, 364.

Lv, B., An, Q., Zeng, Q., Zhang, X., Lu, P., Wang, Y., Zhu, X., Ji, Y., Fan, G. and Xue, Z. (2019). Single-cell
RNA sequencing reveals regulatory mechanism for trophoblast cell-fate divergence in human
peri-implantation conceptuses. *PLoS Biol* 17, e3000187.

Lyu, X., Rowley, M. J. and Corces, V. G. (2018). Architectural Proteins and Pluripotency Factors
Cooperate to Orchestrate the Transcriptional Response of hESCs to Temperature Stress.
Molecular Cell 71, 940-955.e7.
Marbach, D., Costello, J. C., Küffner, R., Vega, N. M., Prill, R. J., Camacho, D. M., Allison, K. R., The
DREAM5 Consortium, Kellis, M., Collins, J. J., et al. (2012). Wisdom of crowds for robust gene
network inference. *Nature Methods* 9, 796–804.
Margolin, A. A., Nemenman, I., Basso, K., Wiggins, C., Stolovitzky, G., Favera, R. D. and Califano, A.
(2006). ARACNE: An Algorithm for the Reconstruction of Gene Regulatory Networks in a
Mammalian Cellular Context. *BMC Bioinformatics* 7, S7.
Materna, S. C. and Davidson, E. H. (2007). Logic of gene regulatory networks. *Curr Opin Biotechnol.* 4,
351–354.
Meers, M. P., Bryson, T. D., Henikoff, J. G. and Henikoff, S. (2019). Improved CUT&RUN chromatin
profiling tools. *eLife* 8,.
Meistermann, D., Bruneau, A., Loubersac, S., Reignier, A., Firmin, J., François-Campion, V., Kilens, S.,
Lelièvre, Y., Lammers, J., Feyeux, M., et al. (2021). Integrated pseudotime analysis of human
pre-implantation embryo single-cell transcriptomes reveals the dynamics of lineage
specification. *Cell Stem Cell* 28, 1625-1640.e6.
Mitsui, K., Tokuzawa, Y., Itoh, H., Segawa, K., Murakami, M., Takahashi, K., Maruyama, M., Maeda,
M. and Yamanaka, S. (2003). The homeoprotein Nanog is required for maintenance of
pluripotency in mouse epiblast and ES cells. *Cell* 113, 631–642.
Mochida, K., Koda, S., Inoue, K. and Nishii, R. (2018). Statistical and Machine Learning Approaches to
Predict Gene Regulatory Networks From Transcriptome Datasets. *Front Plant Sci* 9, 1770.
Morris, S. A., Cahan, P., Li, H., Zhao, A. M., San Roman, A. K., Shivdasani, R. A., Collins, J. J. and Daley,
G. Q. (2014). Dissecting engineered cell types and enhancing cell fate conversion via CellNet.
Cell 158, 889–902.
Nguyen, H., Tran, D., Tran, B., Pehlivan, B. and Nguyen, T. (2021). A comprehensive survey of
regulatory network inference methods using single cell RNA sequencing data. *Briefings in
Bioinformatics* 22, bbaa190.
Niakan, K.K. and Eggan K. (2012). Analysis of human embryos from zygote to blastocyst reveals
distinct gene expression patterns relative to the mouse. *Developmental Biology* 375, 54-64.
990
Niakan, K. K., Han, J., Pedersen, R. A., Simon, C. and Pera, R. A. R. (2012). Human pre-implantation
embryo development. *Development* 139, 829–841.
Pastor, W. A., Liu, W., Chen, D., Ho, J., Kim, R., Hunt, T. J., Lukianchikov, A., Liu, X., Polo, J. M.,
Jacobsen, S. E., et al. (2018). TFAP2C regulates transcription in human naive pluripotency by
opening enhancers. *Nat Cell Biol* 20, 553–564.
Pe'er, D. (2005). Bayesian Network Analysis of Signaling Networks: A Primer. *Science Signaling* 2005,
pl4–pl4.
Peter, I. S. and Davidson, E. H. (2011). Evolution of Gene Regulatory Networks Controlling Body Plan
Development. *Cell* 144, 970–985.
Petropoulos, S., Edsgärd, D., Reinius, B., Deng, Q., Panula, S. P., Codeluppi, S., Plaza Reyes, A.,
Linnarsson, S., Sandberg, R. and Lanner, F. (2016). Single-Cell RNA-Seq Reveals Lineage and X
Chromosome Dynamics in Human Preimplantation Embryos. *Cell* 165, 1012–1026.
Picelli, S., Faridani, O. R., Björklund, A. K., Winberg, G., Sagasser, S. and Sandberg, R. (2014). Full-
length RNA-seq from single cells using Smart-seq2. *Nat Protoc* 9, 171–181.
Pratapa, A., Jalihal, A. P., Law, J. N., Bharadwaj, A. and Murali, T. M. (2020). Benchmarking
algorithms for gene regulatory network inference from single-cell transcriptomic data.
Nature Methods 17, 147–154.
Roode, M., Blair K., Snell, P., Elder, K., Marchant, S., Smith, A., Nichols, J. (2012). Human hypoblast
formation is not dependent on FGF signalling. *Developmental Biology* 361, 358-363.
1010

Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* 9, 671–675.
Schoenfelder, S. and Fraser, P. (2019). Long-range enhancer-promoter contacts in gene expression
control. *Nat Rev Genet* 20, 437–455.
Shi, W., Wang, H., Pan, G., Geng, Y., Guo, Y. and Pei, D. (2006). Regulation of the Pluripotency
Marker Rex-1 by Nanog and Sox2. *Journal of Biological Chemistry* 281, 23319–23325.
Shu, H., Zhou, J., Lian, Q., Li, H., Zhao, D., Zeng, J. and Ma, J. (2021). Modeling gene regulatory
networks using neural network architectures. *Nature Computational Science* 1, 491–501.
Skene, P. J. and Henikoff, S. (2017). An efficient targeted nuclease strategy for high-resolution
mapping of DNA binding sites. *eLife* 6, e21856.
Stirling, D. R., Swain-Bowden, M. J., Lucas, A. M., Carpenter, A. E., Cimini, B. A. and Goodman, A.
(2021). CellProfiler 4: improvements in speed, utility and usability. *BMC Bioinformatics* 22,
433.
Stone, M., Li, J., McCalla, S. G., Siahpirani, A. F., Periyasamy, V., Shin, J. and Roy, S. (2021). Identifying
strengths and weaknesses of methods for computational network inference from single cell
RNA-seq data. *bioRxiv* 2021.06.01.446671,.
Strumpf, D., Mao, C.-A., Yamanaka, Y., Ralston, A., Chawengsaksophak, K., Beck, F. and Rossant, J.
(2005). Cdx2 is required for correct cell fate specification and differentiation of
trophectoderm in the mouse blastocyst. *Development* 132, 2093–2102.
Thompson, D., Regev, A. and Roy, S. (2015). Comparative analysis of gene regulatory networks: from
network reconstruction to evolution. *Annu Rev Cell Dev Biol* 31, 399–428.
Tibshirani, R. (1996). Regression Shrinkage and Selection Via the Lasso. *Journal of the Royal
Statistical Society: Series B (Methodological)* 58, 267–268.
Tipping, M. E. (2001). Sparse Bayesian Learning and the Relevance Vector Machine. *Journal of
Machine Learning Research* 1, 211–244.
Wang, B., Pourshafeie, A., Zitnik, M., Zhu, J., Bustamante, C. D., Batzoglou, S. and Leskovec, J. (2018).
Network enhancement as a general method to denoise weighted biological networks.
Nature Communications 9,.
Wang, J., Ma, A., Ma, Q., Xu, D. and Joshi, T. (2020). Inductive inference of gene regulatory network
using supervised and semi-supervised graph neural networks. *Computational and Structural
Biotechnology Journal* 18, 3335–3343.
Wang, X., He, Y., Zhang, Q., Ren, X. and Zhang, Z. (2021). Direct Comparative Analyses of 10X
Genomics Chromium and Smart-seq2. *Genomics Proteomics Bioinformatics* 19, 253–266.
Weigert, M., Schmidt, U., Haase, R., Sugawara, K. and Myers, G. (2020). Star-convex Polyhedra for 3D
Object Detection and Segmentation in Microscopy. In *2020 IEEE Winter Conference on
Applications of Computer Vision (WACV)*, pp. 3655–3662. Snowmass Village, CO, USA: IEEE.
Yan, L., Yang, M., Guo, H., Yang, L., Wu, J., Li, R., Liu, P., Lian, Y., Zheng, X., Yan, J., et al. (2013).
Single-cell RNA-Seq profiling of human preimplantation embryos and embryonic stem cells.
Nature Structural & Molecular Biology 20, 1131–1139.
Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., Ziraldo, S. B., Wheeler, T.
D., McDermott, G. P., Zhu, J., et al. (2017). Massively parallel digital transcriptional profiling
of single cells. *Nat Commun* 8, 14049.
Zou, H. and Hastie, T. (2005). Regularization and variable selection via the elastic net. *J Royal
Statistical Soc B* 67, 301–320.

1060 **Figure legends**
1061

Figure 1. An overview of the regulatory network inference, evaluation, and validation framework and inference on simulated scRNA-seq data. (A) Footprinting analysis was applied to ATAC-seq data from human embryos at the late blastocyst stage to identify potential transcription factor (TF) binding sites and TF-gene regulations. The TFs and genes in this list of chromatin accessibility (CA)-predicted regulations were used to refine the size of scRNA-seq data from each of the cell types in embryos where cells were collected at the same developmental stage (EPI: Epiblast, 26 cells; PE: Primitive Endoderm, 33 cells; TE: Trophectoderm, 45 cells). Four gene regulatory network (GRN) inference approaches and a random approach were applied to the scRNA-seq data. The scRNA-seq data were normalised using four different methods for comparison (log(TPM+1), log(FPKM+1), batch-corrected and log-counts).The reproducibility and biological context of the predicted GRNs were evaluated using several statistical tests. GRNs refined by keeping CA-predicted regulations only were also evaluated. Squares represent genes that regulate other genes, and circles represent genes that are regulated by other genes. **(B)** Comparison of different GRN inference methods (L0L2, GENIE3, Spearman correlation and MI) to recover the ground-truth GRN structure from simulated gene expression data as measured with the average area under the receiver operating characteristic curve (AUROC) from 100 simulations. The average AUROC is shown as a function of different sample sizes (n=10 to 1000) and the number of potential regulators of each gene in the simulated datasets is also varied from 10 to 100. The range of sizes of the three human blastocyst datasets that we analysed is highlighted in grey as a reference. Error bars correspond to standard errors of the mean. **(C)** Box-and-whisker plots show comparison of AUROC values for all simulations for each method with number of samples n=100 and number of regulators p=20, 50 and 100.

Figure 2. Statistical comparison of four different methods to infer early human embryo GRNs. Robustness of the GRNs predicted by four inference methods with or without chromatin accessibility (CA) refinement was evaluated by three different metrics. **(A)** ∆R measures the difference between the median edge-level reproducibility for a GRN inference method and a random predictor. **(B)** ∆AUPRC quantifies the extent to which the interactions inferred from half of a dataset coincide with the top-1% interactions inferred from the second half. This 2-fold cross-validation experiment was repeated ten times and compared to a random predictor. **(C)** The inverse of the normalised L2 loss between the network scores from the two folds was computed. L0L2 did not converge when applied to the TE dataset. All data used was generated using the SMART-seq2 sequencing technique.

Figure 3. Evaluation of the biological relevance of inferred GRNs in the context of early human development. (A) Two-dimensional UMAP representation of the scRNA-seq data from each of the three cell types in the early human at the blastocyst stage. The expression of marker genes for each cell type is also represented in UMAP space: NANOG is used as an Epiblast marker, CDX2 as a Trophectoderm marker and GATA4 as a Primitive endoderm marker. **(B)** Marker activity in the cell-type specific GRNs (EPI: Epiblast, PE: Primitive endoderm; TE: Trophectoderm) and the network common to the three cell types in the human blastocyst. The expected pattern, i.e. the marker that was expected to be active in each GRN, is shown for reference. **(C)** Normalised V-statistic across datasets and inference methods. V corresponds to the number of gene sets that were enriched at a significance level of 10% in a gene set enrichment analysis performed with the genes involved in the top-500 regulations by the 25 most-connected TFs in the EPI or TE GRNs. All sequencing was performed using the SMART-seq2 technique.

Figure 4. TF GRNs predicted by MICA to be regulated by, or to regulate, NANOG, GATA4 or CDX2. Target networks contain a maximum of 25 potential target TFs of the hub (or central) TF, whereas regulator networks include a maximum of 25 TFs that potentially regulate the hub. The average expression of the transcript is represented by the size of the TF node. MI scores are represented by the thickness of the line in the predicted network. To further refine the MICA-predictions we used the Spearman's rank correlation coefficient between the

expression levels of the source and target nodes to define correlated or anti-correlated
expression. Correlated or anti-correlated node pairs correspond to positive or negative
Spearman's rank coefficient with p-value smaller than 0.1. Node pairs with a p-value equal to
or larger than 0.1 were defined as uncorrelated, though they are predicted by MICA. **(A)**
Network of TFs predicted to be targeted by NANOG. **(B)** Network of TFs that are predicted to
regulate NANOG. **(C)** Network of TFs predicted to be targeted by GATA4. **(D)** Network of TFs
that that are predicted to regulate GATA4. **(E)** Network TFs predicted to be targeted by CDX2.
(F) Network of TFs that are predicted to regulate CDX2.
1125
1126
Figure 5. MICA predicted conserved TFAP2C interactions in human early embryo EPI,
TE, and PE cells. (A) Network composed of TFAP2C, JUND, SOX4, and GCM1 in EPI
(left), TE (middle), and PE (right). **(B)** Correlation plots between TFAP2C, JUND, SOX4, and
GCM1 in TE cells. Bottom left: log transformed RNA expression of genes in single cells.
Diagonal: Distribution of log transformed RNA expression for TFAP2C, JUND, SOX4 and
GCM1. Top right: Spearman correlation between TF pairs. (***: $p < 0.001$). **(C)**
Immunofluorescence staining of TFAP2C and JUND in E6.5 human blastocysts. **(D)**
Correlation plot of TFAP2C and JUND protein expression intensity in nuclei of human E6.5
blastocysts. Numbers on the plot are Pearson correlation value. (***: $p < 0.001$).
1136
1137

Figure 1

1140 **Figure 2**
1141

1142

Figure 3

Figure 4

Figure 5