Brainstem *Dbh*⁺ Neurons

Control Chronic Allergen-Induced Airway Hyperreactivity

Authors: Yujuan Su[1], Jinhao Xu[1,2], Ziai Zhu[1], Ji Chin[1], Le Xu[1], Haoze Yu[1], Victoria Nudell[3], Barsha Dash[4], Esteban A. Moya[5], Li Ye[3], Axel Nimmerjahn[6], Xin Sun[1,2, *]

Affiliations:

[1]Department of Pediatrics, School of Medicine, University of California San Diego, La Jolla, CA 92093, USA

[2]Department of Biological Sciences, University of California San Diego, La Jolla, CA 92093, USA

[3]Department of Neuroscience, Scripps Research Institute, La Jolla, CA 92037, USA

[4]La Jolla Institute for Immunology, La Jolla, CA 92037, USA

[5]Division of Pulmonary, Critical Care & Sleep Medicine, Department of Medicine, University of California, San Diego, CA 92093, USA

[6]Waitt Advanced Biophotonics Center, The Salk Institute for Biological Studies, La Jolla, CA 92037, USA

*Correspondence author: xinsun@health.ucsd.edu

Abstract: Chronic exposure of the lung to irritants such as allergen is a primary cause of asthma characterized by exaggerated airway constriction, also called hyperreactivity, which can be life-threatening. Aside from immune cells, vagal sensory neurons are important for airway hyperreactivity [1-4]. However, the identity and signature of the downstream nodes of this adaptive circuit remains poorly understood. Here we show that a single population of *Dbh*⁺ neurons in the nucleus of the solitary tract (nTS) of the brainstem, and downstream neurons in the nucleus ambiguous (NA), are both necessary and sufficient for chronic allergen-induced airway hyperreactivity. We found that repeated exposures of mice to inhaled allergen activates nTS neurons in a mast cell-, interleukin 4 (IL-4)- and vagal nerve-dependent manner. Single-nucleus RNA-seq of the nTS at baseline and following allergen challenges reveals that a *Dbh*⁺ population is preferentially activated. Ablation or chemogenetic inactivation of *Dbh*⁺ nTS neurons blunted, while chemogenetic activation promoted hyperreactivity. Viral tracing indicates that *Dbh*⁺ nTS neurons, capable of producing norepinephrine, project to the NA, and NA neurons are necessary and sufficient to relay allergen signals to postganglionic neurons that then directly drive airway constriction. Focusing on transmitters, delivery of norepinephrine antagonists to the NA blunted allergen-induced hyperreactivity. Together, these findings provide molecular, anatomical and functional definitions of key nodes of a canonical allergen response circuit. The knowledge opens the possibility of targeted neural modulation as an approach to control refractory allergen-induced airway constriction.

Interoception, a concept first introduced over 100 years ago [5], defines a fundamental biological process whereby the nervous system senses and responds to the inner state of the body including that of internal organs [6]. Recent work has begun to unravel the neural mechanisms for interoception in many tissues, including the larynx, heart, intestine, and bladder [7-11]. However, knowledge remains limited on complete multi-synaptic circuits that sense signals from organs, integrate them in the nervous system, and respond to them through modulation of organ function.

The lung, given its large surface area, rich nerve innervation and direct exposure to aerosol environment, is a prominent source of interoceptive signals [12-15]. In lung, airway constriction, triggered by exposure to irritants including allergen, is a normal protective physiologic response. It is driven by contraction of smooth muscles that wrap around the airways [16]. Following chronic exposure, allergen would trigger exacerbated airway constriction, also termed airway hyperreactivity. The resulting restricted airflow is a primary asthma morbidity and a major cause of asthma-associated deaths.

It has been shown that in repeated allergen-challenged mice but not in naïve mice, vagotomy abolished airway hyperreactivity as assayed in the presence of methacholine, a synthetic acetylcholine analogue, suggesting dependence of the hyperreactivity on vagal reflex [2]. Similarly, ablation or repression of $Trpv1^+$ or $Nav1.8^+$ vagal afferent neurons led to reduced airway hyperreactivity in allergen-challenged mice, but not methacholine-induced acute constriction in naïve mice [3,4]. In the context of repeated allergen-induced airway hyperreactivity, the identity of the nodes downstream of vagal afferent neurons, their molecular signatures, and the nature of neurotransmitters and signals remain poorly defined.

A key knowledge gap is the brainstem neurons important for transmitting the chronic allergen signal. Mapping data from us and others show that lung-innervating vagal sensory neurons project exclusively to the nucleus of the solitary tract (nTS) region of the brainstem [8,13,15,17]. Aside from lung, nTS integrates sensory inputs from many organs including the heart, gut and intestine [8,9,14,18,19]. As a key signal relay center, neurons in the nTS project to higher order brain nuclei, as well as effector areas in the brainstem, including the dorsal motor nucleus of the vagus nerve (DMV) and nucleus ambiguous (NA) [20-22]. Systematic analyses of the neuronal diversity of the nTS and their downstream nodes are critical for interrogating functional heterogeneity and specificity.

In this study, we dissect the mechanism of how normal lungs react to repeated allergen exposures, a disease-relevant adaptive response that is distinct from acute airway constriction. In wild-type mice, we show that repeated, but not single, allergen challenges led to localized activation of nTS neurons. This activation is reduced in either mast cell mutants, after treatment with interleukin 4 (IL-4) neutralizing antibody, or upon vagotomy. Genetic ablation of activated nTS neurons blunted allergen-induced airway hyperreactivity, but not goblet cell metaplasia or type 2 immune cell recruitment. Using single-nucleus RNA sequencing (snRNA-seq), we identified 13 distinct neuron types in the nTS. Using this dataset, we identified a single Dbh^+ sub-population that is enriched for allergen-activated neurons. Ablating or disrupting the function of Dbh^+ nTS neurons blunted airway hyperreactivity, while activating them in place of allergen exposure promoted hyperreactivity in sensitized airways. On the descending efferent side, Dbh^+ nTS neurons project to parasympathetic neurons in the NA region of the brainstem. Chemogenetic inactivation of $Chat^+$ neurons in the NA, but not the DMV, attenuated airway hyperreactivity, while activation of NA neurons in place of allergen exposure promoted hyperreactivity in sensitized airways. Blocking norepinephrine receptors in the NA blunted allergen-induced airway hyperreactivity. Instead of projecting directly to lung, NA neurons project to postganglionic neurons in the trachea and extrapulmonary bronchi, which in turn target airway smooth muscles. Taken together, these *in vivo* data delineate a complete interoception

88 neural circuit in the normal lung that is conditioned by repeated allergen exposures, and is
89 composed of nodes in the brainstem that are both necessary and sufficient for allergen-induced
90 airway hyperreactivity, mimicking exacerbated airway constriction in asthma.
91
nTS neurons were activated upon repeated, but not single, inhaled allergen challenges
To investigate neurons involved in allergen response, we used a mouse model of asthma with
chronic intranasal instillation of house dust mites (HDM), a known trigger for asthmatic
responses in human and rodents [23]. We employed a well-established regimen, with multiple
doses of allergen administrations to lung (Fig. 1a) [24,25]. This triggered the expected asthmatic
responses including exacerbated airway smooth muscle constriction, also termed hyperreactivity,
goblet cell metaplasia and type 2 immune cell infiltration (Fig. 1q, r, Extended Data Fig. 1a, 6d-
f). Extending beyond previous findings indicating that allergen acts through vagal neurons [3,4,26],
we addressed if allergen administered to lung could trigger brainstem neuron activation.
Immunostaining for immediate early factor FOS, a neuronal activation marker, two hours after
each of the challenges in serial sections of the whole brainstem revealed an increase of FOS[+]
cells in HDM-treated mice compared to saline-treated control mice (Fig. 1b, c, Extended Data
Fig. 1b-v). Increased FOS[+] cells were enriched in the dorsomedial part of the nTS region
between Bregma -7.20 mm and Bregma -8.08 mm (Extended Data Fig. 1b-u) in comparison to
other regions such as the adjoining area postrema (AP), DMV and hypoglossal nucleus (12N)
(Extended Data Fig. 2a, b). This enrichment in the nTS is consistent with the findings that lung
innervating sensory nerves project specifically to the same dorsomedial part of the nTS at these
Bregma regions [8,13,15,17]. The increase is found to be statistically significant only after the 4th
HDM challenge, but not after the previous three doses (Extended Data Fig. 3a-i), further
corroborated by *Fos* transcript signal (Extended Data Fig. 4a, b). These findings suggest that
there is conditioning of nTS neurons by repeated, but not single-dose administration of allergen.
It is reported that mast cells are required for the development of allergen-induced airway
constriction [27-29]. Using the same mast cell deficient *c-Kit^w-sh/w-sh* mice, we found a statistically
significant decrease of FOS[+] neurons in the nTS following HDM challenge compared to
heterozygous *c-Kit^w-sh/+* controls (Fig. 1d, e). These data suggest that signals relayed through
mast cells are an important contributor to allergen-induced nTS activation.
Mast cells are among immune cells that produce type 2 immune signals, including interleukin
4 (IL-4), which are critical for allergen-induced airway hyperreactivity [30,31]. It is reported that
neutralization of IL-4 using antibodies can abrogate allergen-induced airway hyperreactivity [30].
Using the same IL-4 neutralizing antibody and regimen, we found that administration of anti-IL-
4, but not isotype control antibody significantly decreased the number of FOS[+] neurons in the
nTS (Fig. 1f-h).
To determine if vagal nerves are essential for allergen activation of nTS, we performed
unilateral vagotomy before allergen administration. This led to a statistically significant decrease
of FOS[+] neurons on the ipsilateral operated side compared to the contralateral side following
HDM challenge (Fig. 1i, j), suggesting that vagal nerves are required for transmitting allergen
signals to the nTS.
To further quantify cumulative allergen-induced activation and also test a tool to manipulate
these neurons, we crossed *Fos^2A-iCreER* (*TRAP2*) mice to *Rosa-lxl-tdTomato* (*Ai14*) mice to label
activated neurons with tdTomato. Following 4-OH tamoxifen injection after each of the HDM
challenge or saline administration (2nd, 3rd and 4th doses, Fig. 1k), we cleared the whole brain and
brainstem using CLARITY [32]. We found a statistically significant increase of activated
tdTomato[+] neurons in the nTS of HDM-treated mice compared to saline-treated controls (Fig. 1l,

m). This was further confirmed through serial sectioning and quantification (Extended Data Fig.
5a-c). These findings paved the way for manipulating these allergen-activated nTS neurons.
137
Ablation of allergen-activated nTS neurons blunted airway hyperreactivity
To determine if activation of nTS neurons plays a role in allergen-induced responses, we
crossed *Fos[2A-iCreER]; Rosa-lxl-tdTomato* (*TRAP2; Ai14*) mice to *Rosa-lxl-DTR* mice to express
diphtheria toxin receptor (DTR) in allergen-activated neurons (*TRAP2; Ai14; DTR,* Fig. 1n).
Following bilateral injection of diphtheria toxin (DTX) into nTS, we confirmed a decrease of
allergen-activated neurons in the nTS (Fig. 1o, p).
To determine if targeted nTS neuron ablation affects airway hyperreactivity, we utilized a
well-established assay that is tailored to test this exacerbated adaptive physiological response as
a result of repeated allergen challenges [1,33]. Following the last dose of HDM or saline
administration, increasing doses of methacholine, a synthetic analogue of acetylcholine, was
intratracheally administered into the airway to mimic what occurs in asthma patients, namely
how acute triggers elicit chronically heightened airway exacerbation, the so-called "asthma
attack". Aside from directly acting on airway smooth muscles to trigger contraction,
methacholine also signals to vagal afferent fibers to activate them which enhance smooth muscle
constriction [2-4,34]. In rodents, lung mechanics including airway hyperreactivity are commonly
assayed using the flexiVent system.
Using this assay, in wild-type mice, as expected, we found that multiple doses of HDM
sensitization and challenge led to further increase of flexiVent measurement of respiratory
system resistance (Rrs) and elastance (Ers) compared to the saline control group, demonstrating
allergen-induced airway hyperreactivity (Fig. 1q, r, N=5 for Saline, N=6 for HDM). As this
hyperreactivity is abrogated by vagotomy [2], the differential airway constriction between the
HDM and saline groups is primarily due to methacholine-activated vagal reflex rather than its
direct effect on airway smooth muscles. Thus, by comparing the extent of airway constriction
triggered by the same concentration of methacholine, we used the differences between
experimental groups and corresponding controls as a measure of the impact of the chronically
adaptive vagal circuit.
In the DTR-DTX experimental group, DTX ablation of allergen-activated nTS neurons led to
blunted airway hyperreactivity compared to vehicle-injected group (Fig. 1s, t, N=4 for vehicle
injection control, N=7 for DTX injection), suggesting that allergen-activated nTS neurons are
part of the circuit conditioned by HDM to control airway hyperreactivity. In contrast, there was
no change in HDM-induced goblet cell metaplasia, type 2 immune cell recruitment or expression
of key type 2 cytokine genes *Il4*, *Il5* and *Il13* (Extended Data Fig. 6a-i). These data together
suggest that while allergen-activated nTS neurons do not substantially modulate goblet
metaplasia or type 2 immune recruitment, they are critical for amplifying allergen-induced
airway hyperreactivity.
173
Single nucleus transcriptomic signatures of the nTS allow dissection of diversity and
selectivity
The nTS is known to receive diverse signals from many organs, including lung, heart, gut,
intestine, etc [8,9,14,18,19]. To identify which specific subset of the nTS neurons were activated by
allergen response in lung, we started with defining the overall molecular diversity of the nTS
through single-nucleus RNA sequencing (snRNA-seq) using flash frozen tissues. This method
was chosen over single-cell RNA sequencing to avoid tissue digestion, cell dissociation and
sorting which reduces neuronal viability and can also artificially induce cellular stress signatures
[35]. Using anatomical landmarks to minimize incorporation of the adjacent AP and DMV regions,

we dissected the whole nTS region from adult wild-type mice (Fig. 2a). Transcriptome data from
12,966 nTS nuclei were obtained using 10x Genomics technology. Unsupervised clustering
analysis revealed 8148 neurons and 4818 glial cells. Focusing on neurons, we further purified
nTS neurons in silico by excluding AP and DMV neurons as identified by their profiles from two
recently published snRNA-seq datasets of these regions [36,37]. The remaining nTS neurons
segregated into 13 clusters as shown in uniform manifold approximation and projection (UMAP)
plots (Fig. 2b), a comparable number to a recent published nTS single cell-RNAseq dataset [38].
All 13 neuronal clusters express pan-neuronal markers *Rbfox3* and *Snap25* (Fig. 2c, d), with little
expression of glial marker genes (*GFAP*, *Olig1*, *Aqp4* or *Tmem119,* Extended Data Fig. 7a-d).
Five neuronal clusters (Clusters 1, 2, 5, 9,12) are *Scl17a6*$^+$ glutamatergic excitatory neurons,
while the remaining eight clusters (Clusters 3, 4, 6, 7, 8, 10, 11,13) are *Slc32a1*$^+$ GABAergic
inhibitory neurons (Fig. 2e, f). Individual neuronal populations were clearly distinguished by a
set of marker genes (Fig. 2g). We further validated the expressions of marker genes using UMAP
plots and either our own RNAscope *in situ* hybridization assay (for excitatory clusters, Fig. 2h-q)
or data from the Allen Brain Atlas (for inhibitory clusters, Extended Data Fig. 7e-l). A
comprehensive analysis of genes encoding neurotransmitter/neuropeptide receptors,
neurotransmitter synthesis/secretion machineries, neuropeptides and hormones revealed the
molecular capabilities of each neuronal population in receiving and transmitting signals
(Extended Data Fig. 7m-o).
202
Dbh$^+$ neurons in the nTS were preferentially activated upon allergen challenge
To determine the molecular signature of allergen-activated nTS neurons, we compared nTS
snRNA-seq signatures of wild-type mice 1.5 hours after the 4th saline or 4th HDM challenge.
Compared to naïve nTS, the same 13 neuron subtypes were detected in saline- and HDM-
challenged nTS (Fig. 3a, Extended Data Fig. 8a). Among the five clusters (Clusters 1, 6, 7, 10
and 12) that show Log2 average expression levels of *Fos* above 0, Cluster 1 is the only excitatory
cluster that showed an increase of the percentage of *Fos*$^+$ neurons (Fig. 3b, Extended Data Fig.
8b). Hence, we focused on Cluster 1 to determine its role in allergen-induced responses.
To validate that Cluster 1 neurons were activated upon HDM challenge, we performed
double *in situ* hybridization with *Fos* and Cluster 1 marker gene *Dbh* (Fig. 3c). By *Dbh*
RNAscope *in situ* hybridization on serial sections, we found that within nTS, *Dbh* expression is
consistently detected between Bregma -6.96 mm and Bregma -8.08 mm, encompassing the
regions where allergen-induced Fos$^+$ cells were enriched (Extended Data Fig. 1b-u, Extended
Data Fig. 9a). On individual sections, we observed a substantial overlap between *Dbh* and *Fos*,
where 59.11% *Dbh*$^+$ neurons were *Fos*$^+$, and 38.18% *Fos*$^+$ neurons were *Dbh*$^+$ (Fig. 3d, N=12
sections). To better assay the spectrum of allergen-activated neurons, we also screened a number
of additional known nTS neuronal markers [38-40], including *Cnr1*, *Th* and *Cck*. We found that after
HDM challenge, there were some *Fos* overlap with *Cnr1* and *Th*, but not *Cck* (Extended Data
Fig. 10a-c). Among *Fos*$^+$ neurons, the *Dbh*$^+$ neurons appear more enriched compared to *Cnr1*$^+$ or
Th$^+$ neurons.
To further confirm that *Dbh*$^+$ neurons were activated, and to validate a driver line to
manipulate these neurons, we crossed transgenic *Dbh-cre* mice to *Rosa-lxl-tdTomato* (*Ai14*)
mice. In *Dbh-cre; Ai14* mice, double *in situ* hybridization of *Dbh* and *tdTomato* showed >95%
overlap of signals, confirming specificity of cre expression (Extended Data Fig. 11a, b, N=6
sections). After HDM administrations in *Dbh-cre; Ai14* mice, we observed a substantial overlap
between tdTomato$^+$ and FOS$^+$ neurons in the nTS (41.99% tdTomato$^+$ neurons were Fos$^+$, and
57.59% Fos$^+$ neurons were tdTomato$^+$), further confirming that *Dbh*$^+$ neurons were activated
upon allergen challenges to lung (Fig. 3e, f, N=8 sections).

Ablation or inactivation of *Dbh+* neurons in the nTS blunted airway hyperreactivity

To determine if *Dbh+* nTS neurons are necessary for mediating allergen response, we employed three approaches. First, we performed chemical ablation by injecting anti-DBH antibody conjugated to the toxin Saporin (SAP). DBH-SAP was previously validated to specifically ablate DBH+ neurons with no effects on neighboring neurons [41]. We injected anti-DBH-SAP stereotactically into both sides of the nTS of wild-type mice before allergen challenge (Fig. 4a). Compared to scrambled peptide-conjugated SAP (blank-SAP) control, anti-DBH-SAP group showed a clear reduction of DBH+ neurons after injection, confirming ablation efficiency (Fig. 4b). There is little change in the size of other excitatory nTS neuronal subsets as labeled by their specific markers, including *Gli3* (Cluster 2), *Sncg* (Cluster 5), *Slc17a7* (Cluster 9) or *Mafa* (Cluster 12), confirming ablation specificity (Extended Data Fig. 12a-f). Following HDM treatment, the anti-DBH-SAP injected group showed reduced FOS+ cells compared to blank-SAP control (Extended Data Fig. 13a-c). As a baseline control, in flexiVent assays, DBH+ neuron ablation had no effect on methacholine responses in naïve mice not exposed to HDM (Extended Data Fig. 13d-e, N=3 for blank-SAP+Saline, N=4 for anti-DBH-SAP+Saline). In contrast, following HDM challenge, the anti-DBH-SAP treated group showed statistically significant blunted airway hyperreactivity compared to the blank-SAP control group (Fig. 4c, d, N=4 for blank-SAP+HDM, N=9 for anti-DBH-SAP+HDM).

Second, we performed genetic ablation by crossing *Dbh-cre* mice to *Rosa-lxl-DTR* mice (*Dbh-cre; DTR*) and then injected DTX bilaterally into nTS (Fig. 4e). This led to efficient loss of *Dbh+* neurons in the nTS (Fig. 4f). Following this targeted genetic ablation, we carried out HDM challenge and found that allergen-induced airway hyperreactivity was blunted in the DTX group compared to vehicle-injected control group (Fig. 4g, h, N=10 for vehicle+HDM, N=10 for DTX+HDM).

Third, we performed chemogenetic inactivation using Designer Receptors Exclusively Activated by Designer Drugs (DREADD) system. We injected AAV-flex-hM4D(Gi)-mCherry bilaterally into nTS of *Dbh-cre* mice, eliciting efficient and specific expression in the nTS, but not nearby regions such as AP (Fig. 4i, Extended Data Fig. 16a, b). Following robust expression, we started the HDM regimen and administered Clozapine-N-Oxide (CNO) to activate hM4D(Gi) before the 2nd to the 4th HDM, sparing the first HDM to allow for sensitization of the immune system which is required for subsequent activation of the nervous system [3]. We found that the CNO group showed blunted airway hyperreactivity response compared to vehicle group (Fig. 4j, k, N=5 for HDM+vehicle, N=9 for HDM+CNO). Taking together, data from chemical ablation, genetic ablation and chemogenetic inactivation demonstrate that *Dbh+* nTS neurons constitute an essential node of the circuit critical for allergen-induced airway hyperreactivity.

Activation of *Dbh+* neurons in the nTS induced airway hyperreactivity

To address if *Dbh+* neurons in the nTS are sufficient for mediating allergen response, we used DREADD targeted activation. We injected AAV-DIO-hM3D(Gq)-mCherry bilaterally into nTS of *Dbh-cre* mice, eliciting efficient and specific expression in the nTS (Fig. 4l, Extended Data Fig. 17a). Following this robust expression, we started the HDM regimen and administrated CNO in place of the 4th HDM challenge while all groups received the 1st to the 3rd doses of HDM. The flexiVent data showed that while not to the full extent as HDM, CNO activation of *Dbh+* neurons in place of the 4th dose was sufficient to induce increased airway hyperreactivity compared to saline control (Fig. 4m, n, N=5 for 4th HDM, N=11 for 4th CNO, N=8 for 4th Vehicle). In contrast, in naïve mice not exposed to HDM, CNO activation (the 4th dose) of *Dbh+* neurons had no effect (Extended Data Fig. 17b-d, N=3 for 4th Saline+vehicle, N=5 for 4th

Saline+CNO). Taking both loss- and gain-of-function data together, these findings suggest that the activity of *Dbh*[+] neurons in the nTS is both necessary and sufficient for mediating allergen-induced airway hyperreactivity.

In none of the three loss-of-function experiments, nor the chemogenetic gain-of-function experiment did we observe any changes in HDM-induced goblet cell metaplasia, increased immune cell infiltration or expression of *Il4*, *Il5* and *Il13* (Extended Data Fig. 14a-f, 15a-f, 16c-i, 18a-h). These findings together confirm findings from *TRAP2; Ai14; DTR* mice (Extended Data Fig. 6a-i). We also assayed for possible effects on other aspects of lung function. Following either HDM challenge or CNO-activation, compared to saline controls, we found no significant difference in minute ventilation, respiratory frequency, tidal volume, or metabolic rate, as measured by plethysmography (Extended Data Fig. 19a-g). This is consistent with previous report that HDM challenge in mice did not change respiratory parameters [42]. These results indicate that CNO activation of *Dbh*[+] nTS neurons induced airway hyperreactivity in sensitized airways without affecting respiration.

Parasympathetic neurons in the NA are downstream of *Dbh*[+] nTS and upstream of airway-innervating postganglionic neurons

To map the downstream targets of *Dbh*[+] nTS neurons and their role in mediating allergen-induced airway hyperreactivity, we injected AAV-flex-tdTomato to the nTS of *Dbh*-cre mice (Fig. 5a). No fiber was detected in either the trachea or the lung, indicating that *Dbh*[+] nTS neurons do not innervate the airway directly. We then sectioned and screened the whole brainstem and found that tdTomato[+] fibers project to the NA which is composed of *Chat*[+] neurons (Fig. 5b). Our data suggest that *Dbh*[+] nTS neurons project to the NA.

To address if NA neurons project directly to lung, we first stereotaxically injected AAV-flex-tdTomato to the NA of *Chat-cre* mice (Fig. 5c). Both light-sheet imaging of intact trachea/bronchi/lung (Fig. 5d, e) and confocal imaging of thick sections (Fig. 5f) revealed that NA-originated tdTomato[+] fibers project to postganglionic parasympathetic ganglia residing in both the trachea and the bronchi, rather than directly to smooth muscle cells in lung (Extended Data Fig. 20a-e). In comparison, stereotaxic injection of AAV-flex-Tom into the DMV and the nearby 12N of *Chat-cre* mice labeled fibers that passed by the space between the trachea and the esophagus without innervating the trachea or the bronchi (Extended Data Fig. 20f, g).

To confirm NA to postganglionic neuron connection from the retrograde direction, we introduced retrograde tracing dye CTB488 dorsal to the trachea, where postganglionic parasympathetic ganglia are enriched. In whole brainstem sections, we found CTB488 labeling of *Chat*[+] neurons in the NA region, with little cell body labeling in other regions including the DMV and the 12N (Fig. 5g).

To directly address the role of *Chat*[+] neurons in the NA in mediating allergen-induced airway hyperreactivity, we utilized the DREADD system. To inactivate *Chat*[+] neurons in the NA, we injected AAV-flex-hM4D(Gi)-mCherry bilaterally into NA of *Chat-cre* mice (Fig. 5h, i). We found that the CNO group showed blunted airway resistance and elastance responses compared to the vehicle-injected control group (Fig. 5j, k, N=7 for HDM+vehicle, N=7 for HDM+CNO). Conversely, to specifically activate *Chat*[+] neurons in the NA, we injected AAV-DIO-hM3D(Gq)-mCherry bilaterally into NA of *Chat-cre* mice (Fig. 5l, m). While not as potent as HDM, activation of *Chat*[+] NA neurons by CNO is sufficient to induce increased airway hyperreactivity compared to saline (Fig. 5n, o, N=6 for 4th HDM, N=13 for 4th CNO, N=6 for 4th Vehicle). In naïve mice not exposed to HDM, we found that CNO activation of *Chat*[+] NA neurons had no added effect on airway constriction under increasing doses of methacholine (Extended Data Fig. 21a-c, N=3 for 4th Saline+vehicle, N=6 for 4th Saline+CNO).

Previous studies have shown that DMV, a key node of autonomic control [43], is also a target of nTS neurons [43,44]. To test if DMV may also act downstream of *Dbh*[+] nTS neurons in the allergen circuit, we performed DREADD-based inactivation or activation of *Chat*[+] neurons in the DMV (Extended Data Fig. 22a, d). In contrast to results from the NA, these DMV perturbations showed little effect in modulating HDM-induced airway hyperreactivity (Extended Data Fig. 22b, c, e, f).

Our data together indicate that *Chat*[+] neurons in the NA are necessary and sufficient in mediating allergen-induced airway hyperreactivity.

Single nucleus transcriptomic signatures of the NA allow dissection of allergen signaling relay

To determine the molecular nature of the signaling relay leading to NA activation, we utilized snRNA-seq to define NA neuron signatures. As all NA neurons express *Chat*, in the brainstem of *Chat-cre; CAG-Sun1/sfGFP* mice, we collected around the small GFP[+] region that corresponds to NA, and avoided DMV and 12N regions which are also GFP[+]. Among all the neurons identified in the snRNAseq dataset, we focused on the 190 *Chat*[+] neurons, which segregated into 3 clusters as shown in the UMAP plot (Fig. 6a). Individual neuronal populations were distinguished by marker genes (Fig. 6b), which were verified by UMAP plots and caudal brainstem *in situ* hybridization data from the Allen Brain Atlas (Fig. 6c-h). Our neuron number and neuron type clusters are comparable to the two recent published NA single cell datasets [45,46].

The addition of snRNA-seq signature of this downstream node of the allergen circuit allowed us to define an in silico molecular pathway that starts from the lung and back to the lung (Extended Data Fig. 23a-c). Given our finding that allergen-induced *Fos* activation is reduced in mast cell mutants and following anti-IL-4 antibody neutralization (Fig. 1d-h), we started with key immune signals from type 2 immune cells. Our previous findings also indicate that pulmonary neuroendocrine cells (PNECs) are essential for amplifying allergen-induced responses [25]. Thus, we added key PNEC products. In the next node the vagal ganglia, we focused on *Trpv1*[+] population because its ablation led to reduced airway hyperreactivity in allergen-challenged mice [3]. By analyzing published snRNA-seq datasets of the vagal ganglia [7,8,47,48], we found multiple corresponding receptors for immune cell and PNEC ligands in *Trpv1*[+] vagal populations (Extended Data Fig. 23a).

We then paired ligands from *Trpv1*[+] vagal neurons and receptors in *Dbh*[+] neurons in the nTS from our snRNA-seq dataset, and ligands from *Dbh*[+] nTS neurons and receptors in the next downstream node using our snRNA-seq datasets (Extended Data Fig. 23b, c). Given that both NA neurons and their downstream airway smooth muscle-innervating postganglionic neurons are *Chat*[+], the ligands from both of these neuron groups are likely acetylcholine [49-51]. Taken together, this analysis revealed candidate signals for this neuroimmune circuit that relays from lung immune cells, neurons in the vagal ganglia, nTS and NA of the brainstem, trachea/bronchi, and airway smooth muscle cells.

Blocking norepinephrine receptors in the NA blunted allergen-induced airway hyperreactivity

To test the function of signals predicted from our ligand-receptor analysis, we focused on the nTS-NA relay. *Dbh* encodes a key enzyme that converts dopamine to norepinephrine, leading to the hypothesis that within the allergen circuit, norepinephrine may be the signal between *Dbh*[+] nTS and NA. From analysis of the NA snRNA-seq dataset and RNAscope assays, among all norepinephrine receptor genes, only *Adra1a* and *Adra1b* are expressed in *Chat*[+] neurons in the NA (Fig. 6i-l). Furthermore, we found that retrograde tracing by tracheal injection of CTB488

labeled $Adra1a^+$ and $Adra1b^+$ neurons in the NA (Fig. 6m, n). These data support that norepinephrine receptor-expressing NA neurons are in the airway circuit, and may receive norepinephrine signals from Dbh^+ nTS neurons in response to allergen challenges.

To test this hypothesis, we addressed whether blocking norepinephrine signal reception in the NA can blunt airway hyperreactivity. Either Prazosin or Terazosin, two norepinephrine receptor antagonists specific for ADRA1, were individually infused into NA bilaterally through implanted cannula (Fig. 6o, Extended Data Fig. 24a). Compared to vehicle control, targeted administration of Prazosin or Terazosin significantly blunted airway hyperreactivity response (Fig. 6p, q, N=6 for HDM+vehicle, N=7 for HDM+Prazosin, N=6 for HDM+Terazosin). These results indicate that the activity of norepinephrine receptor in the NA is necessary for mediating allergen-induced airway hyperreactivity, demonstrating the molecular nature of this key relay in the allergen response circuit.

Discussion

In this study, our findings delineate a full neural circuit where chronic allergen challenges to lung are transmitted through immune cells such as IL-4 producing immune cells and ascending vagal afferents to a Dbh^+ subset of the nTS central integrator, which acts through descending efferent neurons ($Adra1a^+$/$Adra1b^+$) in the NA before projecting back to postganglionic neurons innervating the airway to control airway hyperreactivity (Fig. 6r). This is a circuit identified in wild-type mice following repeated conditioning by allergen, suggesting the presence of an intrinsic molecular and cellular response machinery. Recurring irritation is a common cause of chronic diseases and repeated allergen exposure is central to asthma pathogenesis. Our findings reveal a disease-relevant circuit with players from the immune system, the nervous system and functional cells of the organ.

We found that only repeated, but not single, allergen challenges led to a statistically significant increase in activated nTS neurons. This exemplifies a cardinal feature of chronic conditioning, a behavior that is distinct from mechanisms that drive acute airway constriction [2,52,53]. These results provide an example of experience-driven neural plasticity in peripheral interoceptive circuits, which are analogous to features classically showcased by central nervous system neurons in learning and memory.

While methacholine can induce acute airway constriction in both groups, repeated allergen challenge led to exacerbated response above the level in naïve animals. Our data demonstrate that this elevated airway resistance was blunted by ablation or chemogenetic inactivation of Dbh^+ nTS neurons, demonstrating for the first time the critical role of the nTS neurons in this response. Despite differences, airway hyperreactivity and acute airway constriction may share overlapping nodes. Activation of nervous system in naïve animals, by either electrical stimulation of vagal C-fiber, chemical induction of $MrgprC11^+$ vagal neurons, or optogenetic stimulation of thoracic cholinergic nerves, induced acute airway constriction in the absence of exogenous methacholine [26,52,53]. These findings suggest that repeated allergen exposures may hijack these existing nodes, condition them through molecular changes or anatomical remodeling, to elevate airway constriction above the threshold level of acute response, mimicking asthma. Our results further define the identity of the brainstem neurons that connect the afferent and efferent nodes to control airway constriction.

While the combined connectivity is necessary and sufficient for allergen-induced hyperactivity, individual nodes of this allergen circuit may also function in other physiological or pathological settings. For example, Dbh-expressing nTS neurons were implicated in appetite control by projecting to the lateral parabrachial nucleus or in the arcuate nucleus of the hypothalamus [54-56]. Dissecting the anatomical connectivity and the pairing of

neurotransmitter/neuropeptide to their receptors are key to unraveling the different roles of the same neuron type in distinct circuits. Towards this goal, our snRNA-seq datasets of the nTS and NA provide the basis for dissecting molecular connectivity and specificity. In the context of the allergen circuit, these two datasets, together with existing lung and vagal neuron single cell datasets, and the knowledge that acetylcholine is the likely neurotransmitter from NA to airway smooth muscle cells, complete the connectome of the allergen circuit (Extended Data Fig. 23a-c) [7,8,47,48]. Two recent studies delineated diverse functions of NA neurons in targeting the esophagus, the heart and/or the lung [45,46]. It is likely that the cardiopulmonary subset of the NA neurons is responsible for mediating allergen response.

The molecular signatures of the allergen hyperreactivity circuit enables a novel dimension for modulating airway tone in asthma. Neuromodulation could bypass the systemic side effects of current asthma treatments that relay on administration of steroids and biologics. This adaptive circuit showcases the importance of neuroimmune control of organ function. Beyond the lung, mapping how organs act through their peripheral and central nerve connections to receive, process and respond to chronic inputs will open a new dimension in the modulation of disease-relevant interoceptive activities.

Methods
Mice
All experimental procedures were performed in the American Association for Accreditation of Laboratory Animal Care (AAALAC)-certified laboratory animal facility at the University of California, San Diego (UCSD). All animal husbandry and experiments were conducted under approved Institutional Animal Care and Use Committee (IACUC) guidelines. Adult (8-16 weeks) mice from strains: C57BL/6J (wild-type, JAX 000664), c-$Kit^{w\text{-}sh/w\text{-}sh}$ (JAX 030764), $Fos^{2A\text{-}iCreER}$ (*TRAP2*, JAX 030323), *Rosa-lxl-tdTomato* (*Ai14*, JAX 007914), *Rosa-lxl-DTR* (JAX 016603), *Chat-cre* (JAX 031661) and *CAG-Sun1/sfGFP* (JAX 030952) lines were purchased from the Jackson lab. *Dbh-cre* (MMRRC 036778) line was purchased from the Mutant Mouse Resource and Research Center (MMRRC). All the *cre* lines we used in this study were kept in B6 background. All *cre* driver lines used are viable and fertile, and no abnormal phenotypes were detected. We crossed *cre* males to B6 wild-type females, and both male and female mice were used in the experiment. Mice were at least six-week-old when performing HDM challenge or doing stereotaxic injection or surgery.

House Dust Mite (HDM) challenge
Adult mice were anesthetized using isoflurane, 50 μg/20 μl of HDM extract (Dermatophagoides pteronyssinus, GREER Labs) was introduced intranasally weekly continuous for 4 weeks on Days 0, 7, 14, and 21. For controls, 20 μl saline was administered intranasally on the same schedule instead of HDM. Mice were sacrificed 1.5 hours after the 4th challenge for *Fos in situ* hybridization or nTS single-nucleus RNA sequencing, 2 hours after the 4th challenge for FOS antibody immunostaining, 24 hours after the 4th challenge for flexiVent assay, periodic acid–schiff (PAS) staining and quantitative PCR (qPCR), 3 days after the 4th challenge for flow cytometry.

Histology preparation and PAS staining
Mice were euthanized by CO_2 inhalation. Lungs were inflated with 4 % paraformaldehyde (PFA) at 35 cm H_2O airway pressure, fixed overnight and then prepared for paraffin sections at 6 μm. Goblet cells were stained using a PAS staining kit (Sigma).

Tissue collection and immunofluorescence staining

Mice were euthanized by CO_2 inhalation. Mice were then transcardinally perfused with Phosphate-buffered saline (PBS) followed by perfusion fixation with 4% PFA. The brainstems were harvested and post-fixed overnight at 4°C in 4 % PFA. After fixing, tissues were washed three times in PBS followed by overnight sucrose dehydration. The brainstem or the lung were then cryo-embedded in OCT compound following standard procedure. Brainstem blocks were sectioned at 20 μm or 40 μm in rostral to caudal sequence, lung blocks were sectioned at 300 μm for parasympathetic neurons/airway staining. All the sections were then processed for immunostaining following a standard protocol. Primary antibodies used include rabbit anti-c-FOS (SYSY, 226 003), rabbit anti-DBH (Sigma, AB1585), rabbit anti-Dsred (Takara, 632496), rabbit anti-VAChT (SYSY, 139 103), mouse anti-alpha Smooth Muscle Actin-FITC (Sigma, F3777). Secondary antibodies used include goat anti-rabbit FITC, goat anti-rabbit Cy3, goat anti-rabbit Cy5 (all from Jackson Immuno Research Labs). Slides were mounted using Vectashield (Vector Labs) and imaged using the Olympus VS200 Slide Scanner or the Leica SP8 confocal microscope. For quantification of FOS^+ (Fig. 1c, e, h, j, Extended Data Fig. 3c, f, i, Extended Data Fig. 13c) or $tdTomato^+$ (Fig. 1p and Extended Data Fig. 5c) cells in the nTS, or Fos^+ cells after RNAscope *in situ* hybridization (Extended Data Fig. 12f), we did serial section of the whole nTS and counted from three cryosections (three sections were chosen coordinate between Bregma -7.20 mm and -8.08 mm with the same stereotaxic, where HDM-induced FOS^+ cells are elevated) with the maximal numbers of FOS^+ cells and count the average number as one data point for one animal. We used Qupath for overlap quantification in the nTS (Fig. 3d, f).

RNAscope *in situ* hybridization

To prepare nTS sections for RNAscope *in situ* hybridization, fresh brainstems were cryo-embedded in OCT compound. Sections at 12 μm were prepared using a cryostat, collected on Superfrost Plus slides, and left to dry in the cryostat for at least 30 minutes before staining. All staining procedures were performed using RNAscope Fluorescent Multiplex Kit (Advanced Cell Diagnostics, 320850) following the manufacturer's instruction. Briefly, sections were post-fixed in 4% PFA for 1 hour at room temperature, washed in PBS, dehydrated in a series of ethanol washes, and then dried. A hydrophobic barrier was drawn around the section with an ImmEdge pen (Vector Lab, H-4000). The sections were treated with Protease IV in a HybEZ Humidity Control Tray for 30 minutes at room temperature, incubated with target probes in a HybEZ Oven for 2 hours at 40°C, and then treated with Hybridize Amp buffer. Slides were mounted using Fluoromount-G with DAPI (Southern Biotech). The following probes from Advanced Cell Diagnostics were used: Mm-*Fos* (316921), Mm-*Dbh* (407851), Mm-*Gil3* (445511), Mm-*Sncg* (482741), Mm-*Slc17a7* (431201), Mm-*Mafa* (556931), Mm-*Fos* (316921), Mm-*Cnr1*(420721), Mm-*Th* (317621), Mm-*Cck* (402271), Mm-*tdTomato* (317041), Mm-*Chat* (408731), Mm-*Adra1a* (408611), Mm-*Adra1b* (413561), Mm-*Adra2a* (425341), Mm-*Adra2b* (425321), Mm-*Adrb1* (449761), Mm-*Adrb2* (449771) and Mm-*Adrb3* (495521).

Vagotomy

Mice were anesthetized with a mixture of ketamine (100 mg/kg) and xylazine (10 mg/kg) via intraperitoneal injection. Fully anesthetized mice were secured in the sternal position to the homoeothermic blanket using lab tape. Spread 75 % EtOH on the throat of the mouse to wet the fur. Lift the skin to make a vertical cut (1 cm) on the throat, isolate one side of vagal nerve and tease away from carotid artery using small, curved forceps. Pass fine, curved forceps underneath

the vagal nerve and slowly open forceps to separate the nerve fiber from the surrounding tissue. Unilateral vagotomy was conducted by lifting up the vagal nerve and cutting with straight scissors. Control sham operations are simulated by lifting the vagal nerve and releasing it intact. After vagotomy, the wounds were closed carefully with sterilized dissecting scissors and suture strings. The area will be cleaned with 75% EtOH and disinfected with Povidone-Iodine. The time between fully anesthesia and completion of the surgery should take less than 10 minutes. Buprenorphine SR (0.1 mg/kg, subcutaneous injection) was used for pain relief following surgery. One week after surgery, mice were used for allergen challenge.

4-OH tamoxifen injection and *TRAP2* labeling

4-hydroxytamoxifen (4-OHT, Sigma H6278) was dissolved at 20 mg/mL in ethanol by shaking at 37°C for 15 minutes and was then aliquoted and stored at -20°C for up to several weeks. Before use, 4-OHT was redissolved in ethanol by shaking at 37°C for 15 minutes, a 1:4 mixture of castor oil: sunflower seed oil (Sigma, 259853 and S5007) was added to give a final concentration of 10 mg/mL 4-OHT, and the ethanol was evaporated by vacuum under centrifugation. All injections were delivered intraperitoneally to adult *TRAP2* mice. Mice were kept in their home cage for additional 1 week to allow the expression of tdTomato before treatment.

CLARITY-based brain clearing

A hydrogel based on 1 % acrylamide (1 % acrylamide, 0.125 % Bis, 4 % PFA, 0.025 % (w/v) VA-044 initiator, in 1X PBS) was used for all CLARITY preparations. Mice were transcardially perfused with 4 % PFA. After perfusion, brains were post-fixed in 4% PFA overnight at 4°C and then transferred to 1 % hydrogel for 48 hours to allow monomer diffusion. The samples were degassed and polymerized (4-5 hours at 37°C) in a 50 ml tube. The brains were removed from hydrogel and washed with 200 mM NaOH-Boric buffer (pH=8.5) containing 8 % sodium dodecyl sulfate (SDS) for 6-12 hours to remove residual PFA and monomers. Brains could then be transferred to a flow-assisted clearing device using a temperature-control circulator or a simpler combination of 50 ml tube and heated stirring plate. 100 mM Tris-Boric Buffer (pH=8.5) containing 8 % SDS was used to accelerate the clearing (at 40°C). A whole mouse brain can be cleared in 12-16 days. After clearing, the brain was washed in PBST (PBS+0.2% Triton X) for at least 24 hours at 37°C to remove residual SDS. Brains were incubated in a refractive index matching solution (RI=1.45) for 8 hours (up to 1 day) at 37°C and then 6-8 hours at room temperature. After the incubation, brainstem hemispheres were imaged using confocal microscope.

Brainstem stereotaxic injection

Fully anesthetized mice (ketamine at 100 mg/kg and xylazine at 10 mg/kg via intraperitoneal injection) were placed in a stereotaxic frame with the head angled at 45° and body temperature maintained with a heating pad. A midline incision was made though the animal's skin, posterior neck muscles, and dura mater to expose the medulla between the occipital bone and C1 vertebra. Based on the stereotaxic coordinates of mouse brain [57] and using Obex as a reference point, bilateral injections were made into the nTS (0.1 mm rostral to Obex, 0.2 mm lateral to midline, 0.25 mm under the medullary surface), the DMV (0.05 mm caudal to Obex, 0.1 mm lateral to midline, 0.1 mm under the medullary surface), or the NA (0.65 mm caudal to Obex, 1.25 mm lateral to midline, 0.45 mm under the medullary surface) using a calibrated glass micropipette attached to a Nanoject II injector (Drummond) microprocessor pump (Pneumatic PicoPump; WPI). Over 5 minutes, DTX (2 ng/200 nl), anti-DBH-SAP (42 ng/200 nL, advanced targeting

system, IT-03), blank-SAP (advanced targeting system, IT-21), or viruses (50 nl for nTS or
DMV, 20 nl for NA, Boston Children's Hospital Vector Core) were gradually injected at the
targeted site of *TRAP2* mice, wild-type mice, *Dbh-cre* or *Chat-cre* mice, respectively. Viruses
include AAV2/9-flex-tdTomato (1.3×10^{13} genome copies (gc)/ml), AAV2/8-flex-hM4D(Gi)-
mCherry (2.07×10^{13} gc/ml), AAV2/8-DIO-hM3D(Gq)-mCherry (8×10^{12} gc/ml). After injection,
the glass micropipette was left in place for 10 minutes to limit nonspecific spread of the toxin or
virus. The micropipette was pulled out very slowly. Buprenorphine SR (0.1 mg/kg, subcutaneous
injection) was used for pain relief following surgery. The mice were allowed to recover from
anesthesia on a heating blanket before returning to the home cage.
574
Airway hyperreactivity assayed by flexiVent
Anesthetized (100 mg/kg ketamine and 10 mg/kg xylazine, intraperitoneal injection) mice
were paralyzed with acepromazine (10 mg/kg, intraperitoneal injection) and then tracheotomized
with a 20G sterile catheter, attached to a flexiVent pulmonary mechanics apparatus (SCIREQ).
Mice were ventilated at a tidal volume of 9 ml/kg with a frequency of 150 bpm. Positive end-
expiratory pressure was set at 300 mm H_2O. Lung resistance (Rrs) and elastance (Ers) of the
respiratory system were determined in response to aerosolized methacholine challenges (0, 6, 12
and 24 mg/ml). All mice used were adults of similar age and weight. The weight of each animal
was entered into flexiVent at the start of each round of assay. As part of the flexiVent program,
pre-scans were carried out that led to calculation of lung size. Methacholine was dissolved in
sterile saline. The nebulizer was on for 10 seconds to deliver each dose of methacholine. The
mean maximal elastance and resistance of twelve measurements by doses was calculated.
Samples were compared using two-way ANOVA. Statistical analysis at each methacholine
concentration was performed separately. Error bars represent means ± SEM. Statistical
significance called at $p < 0.05$.
590
Tissue dissociation and flow cytometry
Three days after last challenge, mice were anesthetized (100 mg/kg ketamine and 10 mg/kg
xylazine, intraperitoneal injection) and then injected with AF700-counjugated CD45 through
intravenous injection to distinguish circulating immune cells and immune cells resident within
the lung. Mice were euthanized 5 minutes later for lung harvest. Whole lungs were mechanically
dissociated in GentleMACS C tubes (Miltenyi Biotec) containing 5 ml of PRMI 1640 (Thermo
Scientific) with 10 % FBS, 1 mM HEPES (Life Technology), 1 mM $MgCl_2$ (Life Technology), 1
mM $CaCl_2$ (Sigma-Aldrich), 0.525 mg/ml collagenase/dispase (Roche) and 0.25 mg DNase I
(Roche) by running mouse lung 1-2 program on GentleMACS (Miltenyi Biotec). Lung pieces
were then digested by shaking (~150 rpm) for 30 minutes at 37°C. After incubation, Lung pieces
were mechanically dissociated further using mouse lung 2-1 program on GentleMACS, followed
by straining through a 70 mm filter. Red blood cells were removed by adding 1 mL RBC lysis
buffer (Biolegend) to each tube and incubate at room temperature for 1 minute.
The single-cell suspensions from above were then pelleted (1500 rpm, 4°C, 5 minutes),
counted with hemocytometer and diluted to around 1×10^6 cells per ml. They were stained with
Fc blocking antibody (5 mg/ml, BD) at 4°C for 30 minutes. The cells were washed with DPBS
and then incubated with surface marker antibody cocktail for 30 minutes at 4°C. For flow
cytometry of lung myeloid, the following antibodies were used (all antibodies were from
BioLegend): 1:100 AF488-conjugated anti-F4/80; 1:500 BV510-conjugated anti-CD45; 1:500
BV605-counjugated anti-Siglec-F; 1:100 V450-counjugateed anti-TCR-beta; 1:200 V450-
counjugated anti-B220; 1:1000 PerCP-Cy5.5-conjugated anti-Ly6C; 1:2000 PE-conjugated anti-
MHC-II; 1:1000 APC-conjugated anti-CD11c; 1:1000 PE-Cy7 conjugated anti-Ly6G; 1:2000

PE-CF594-counjugated CD11b. For flow cytometry of lung lymphoid, the following antibodies were used (all antibodies were from BioLegend): 1:200 FITC-conjugated anti-CD45; 1:100 AP-Cy7-counjugateed anti-IL7Ra; V450-counjugated Lineage mix (1:200 anti-CD19; 1:500 anti-CD11c; 1:500 anti F4/80; 1:100 anti-NK1.1; 1:100 anti-TER119; 1:100 anti-TCR gamma delta); 1:100 BV510-conjugated anti-ST2; 1:200 PE-Cy7-counjugated anti-TCR-beta; 1:100 BV604-counjugated anti-CD4; 1:2000 PerCP-Cy5.5-conjugated anti-CD90.2; 1:100 APC-conjugated anti-cKit; 1:2000 PE-CF594-counjugated anti-CD11b; 1:200 PE-conjugated anti-MAR.1. The cells were then stained using live/dead dye (1:1000, Ghost Dye Red 780, TONBO biosciences for myeloid, 1:500, Ghost Dye Violet 450, TONBO biosciences for lymphoid) for 30 minutes at 4°C. The cells were fixed using BD Stabilizing Fixative for 30 minutes at 4°C before transferring to FACs tubes. Flow cytometry was analyzed on BD FACS Canto RUO - ORANGE with three lasers (405 nm, 488 nm and 640 nm) using the Flow Cytometry Core at the VA San Diego Health Care System and the San Diego Veterans Medical Research Foundation. All data were further analyzed and plotted with FlowJo software (Tree Star). Eosinophils, innate lymphoid cells (ILC2s) and Th2 cells were gated on live, resident CD45$^+$ singlets.

Isolation of nuclei from frozen nTS

We dissected out the whole nTS region from adult C57BL6/J male mice brainstem. Mice were euthanized using CO_2 inhalation. Brains were acutely harvested from two groups of wild-type males (WT), or from two adult saline- or HDM-challenged males at 1.5 hours after the 4th challenge, respectively (two technical replicates in every group). The nTS was visualized by microscopy and harvested based on anatomical landmarks. Dissected nTS were put in liquid N_2 immediately and store at -80°C or proceed directly to nuclei isolation. On the day of nuclei dissociation, plunge the dounce and pestle in ice. Transfer douncing buffer (DB, 0.25 M sucrose, 25 mM KCl, 5 mM $MgCl_2$ and 10 mM Tris-HCl, pH=7.5) to the dounce in ice and let them chill. Transfer dissected nTS into dounce which includes 1 mL DB with 1 mM DTT (D9779, Sigma), 1x cOmplete EDTA-free protease inhibitor (05056489001, Roche, DB-DP) with 0.1% Triton-X. Use loose pestle 5-10 times gently followed by 15-20 stokes with tight pestle, without introducing bubbles. After filter with 30 μm CellTrics, transfer the contents to pre-chilled low bind Eppendorf tube. Spin at 1,000 g for 10 minutes at 4°C. Remove supernatant and gently resuspend in 1 mL of DB-DP. Spin at 1,000 g for 10 minutes at 4°C. Remove supernatant and gently resuspend in 1 mL permeabilization buffer (BSA, 5% IGEPAL-CA630 (I8896, Sigma), 0.2% DTT, 1 mM cOmplete EDTA-free protease inhibitor, 1xPBS). Resuspend in 50-500 μL of 1.11x tagmentation buffer (66 mM Tris-acetate (pH=7.8, BP-152, Thermo Fisher Scientific), 132 mM K-acetate (P5708, Sigma), 20 mM Mg-acetate (M2545, Sigma) and 32 mM DMF (DX1730, EMD Millipore). Count with hemocytometer.

Single-nucleus RNA sequencing and data analysis

Single-nucleus RNA sequencing experiments were provided by the Center for Epigenomics, University of California, San Diego. nTS nuclei were processed into cDNA libraries using Chromium Single Cell 3' v3 kit (10XGenomics). Sequencing was carried out on NovaSeq (Illumina) platform at the Institute for Genomic Medicine, University of California, San Diego (UCSD). CellRanger package (version 3.0.2) was used to align the raw reads onto the mouse reference genome (GRCm38) and generate the feature-barcode matrix. CellBender was used next to model and remove systematic biases and background noise, and to produce improved estimates of gene expression. Next, R package Seurat (version 4.0) was used to perform data quality control, normalization, principal components analysis and uniform manifold approximation and projection (UMAP method). For raw data from WT, Saline or HDM

conditions, we firstly used SCTransform to regress our batch effects and did doublets removal separately. We then integrated all the data from three conditions (WT, Saline, HDM) using Harmony. Cells with 3,500 to 5,500 unique feature counts were considered high quality cells and used for further analysis. The integrated dataset was normalized, scaled, and then analyzed for variable genes using Seurat's NormalizeData, ScaleData, and FindVariableFeatures functions, respectively. A total of 2,000 top variable features were identified by function FindVariableFeatures and selected for subsequent principal components analysis. Top 20 significant components were chosen to conduct cell clustering by using the algorithm of uniform manifold approximation and projection (UMAP) with default settings. The expression level and feature of selected genes were profiled and visualize by R package ggplot2 (Version 3.3.2). We then read snRNA-seq data from WT condition and profiled dot plot and UMAP plot of marker gene expression in all the nTS clusters (Fig. 2a-q, Extended Data Fig. 7a-o, Extended Data Fig. 8a). We compared snRNA-seq data from Saline and HDM conditions and profiled dot plot of *Fos* in all the nTS clusters (Fig. 3a, b, Extended Data Fig. 8b).

Quantitative PCR (qPCR)

Total RNA was extracted from lungs using Trizol (Invitrogen) and RNeasy Mini RNA extraction kit (Qiagen). RT-PCR was then performed to obtain corresponding cDNA using iScript Select cDNA Synthesis Kit (Bio-Rad). qPCR was performed by CFX ConnectTM system (Bio-Rad) using SYBR Green (Bio-Rad). At least three technical and three biological replicates were performed for each gene if not otherwise notated. Samples were compared using unpaired student's t-test. Error bars represent means ± SEM. Statistical significance called at $p<0.05$.

Primers

Gene name	Sequence
Actb (β-actin)	5′-CGGCCAGGTCATCACTATTGGCAAC-3′ 5′-GCCACAGGATTCCATACCCAAGAAG-3′
Muc5ac	5′-TGACTCAATCTGCGTGCCTT-3′ 5′-AGGCCTTCTTTTGGCAGGTT-3′
Il4	5'-GGTCTCAACCCCCAGCTAGT-3' 5'-GCCGATGATCTCTCTCAAGTGAT-3'
Il5	5′-CCTCTTCGTTGCATCAGGGT-3′ 5′-GATCCTCCTGCGTCCATCTG-3′
Il13	5′-AAAGCAACTGTTTCGCCACG-3′ 5′-CCTCTCCCCAGCAAAGTCTG-3′

nTS-targeted SAP injections

Mice received bilateral injections of either unconjugated blank-SAP or SAP conjugated to an antibody directed to DBH (DBH-SAP; 42 ng/200 nL, pH=7.4; MAB394; Millipore, Temecula, CA, USA), using a Hamilton syringe directed toward the nTS (0.1 mm rostral to Obex, 0.2 mm lateral to midline, 0.25 mm under the medullary surface). Blank-SAP cannot enter cells and is non-toxic to neurons, serving as the appropriate control for DBH-SAP administration. Over 5 minutes, 50 nL of blank-SAP or DBH-SAP was gradually injected to the nTS. DBH-SAP was previously validated to specifically ablate DBH[+] neurons with no effects on neighboring neurons [41]. Upon specific binding to DBH at the synaptic cleft of DBH[+] cells, the complex including

696   conjugated anti-DBH-SAP antibody is endocytosed and transported back to the cell body [58]. In
697   the cell body, the SAP toxin inactivates ribosomes, leading to cell death [59]. As previously
698   published studies indicate that 2 weeks is necessary to eliminate *Dbh*[+] neurons using DBH-SAP
699   [58], mice were injected 1 week before the 1st HDM sensitization and 2 weeks before the 2nd
700   challenge.
701
Plethysmography
Ventilatory parameters during normoxia was measured in unrestrained male mice using a
whole-body barometric plethysmograph (500 mL) modified for continuous flow [60-62]. Flow was
maintained constant through the chamber while a pressure transducer (45 mMP with 2 cm H_2O
diaphragm, Validyne, Northridge CA) recorded the changes attributable to the warming and
expansion of inhaled gases. On the experimental day, the mice were weighed and sealed into an
individual plethysmograph chamber along with a temperature and humidity probe (Thermalert
TH5, Physitemp, Clifton NJ). A constant gas flow (335 ml/min) was delivered with a rotameter
(603, Matheson, Montgomeryville PA) and measured with a flow meter (Sables System
International, Inc. Las Vegas, NV) upstream of the chamber. Gases exited the chamber through a
valve and into a vacuum pump (Model 25, Precision Scientific Co, Chicago IL) to isolate
pressure changes from breathing in the chamber during constant flow with high input and output
impedances. This also allows us to maintain chamber pressure near-atmospheric pressure and
reference pressure measurements (Validyne) in the chamber to atmosphere.
All ventilatory parameters were recorded on an analog-digital acquisition system (PowerLab
8SP, AD Instruments) and analyzed with the LabChart 8-Pro Software, sampling at a rate of 1
kHz. Mice were allowed to acclimatize to the chamber and constant air flow for 45 minutes (air
with 21% of O_2) and then exposed to a 5-minute challenge of hypoxic gas (10% O_2) to test
responsiveness. After that we exposed the mice to normoxic gas for 15 minutes. We analyzed a
minimum of 30 seconds as part of the region of interest between 10 and 15 minutes of the las
exposure to normoxic gas. The frequency (breaths/min) and tidal volume (ml) were calculated
from cyclic peaks in the plethysmograph pressure pulses, and tidal volume was measured using
0.02 ml calibration pulses and the equations were developed by Drorbaugh and Fenn [63]. We
measured minute ventilation ($\dot{V}_I$), respiratory frequency (Rf) and tidal volume (Vt). All of them
were measured under 21% of oxygen. Minute ventilation ($\dot{V}_I$) is the product of frequency and
tidal volume, which was normalized to body mass ventilation (ml/min·kg). We also calculated
oxygen consumption ($\dot{V}_{O_2}$) and carbon dioxide production ($\dot{V}_{CO_2}$) during the ventilation
measurement by recording the inspired and expired oxygen and carbon dioxide fractions were
measured with an O_2/CO_2 analyzer (FOXBOX, Sables System International, Inc. Las Vegas,
NV) sampling from the chamber, and then calculating inspired and expired O_2 and CO_2 fractions,
chamber flow rate and correcting for water vapor and respiratory quotient (eqn. 10.6 and 10.7 in
J.R.B. Lighton, Measuring Metabolic Rates, Oxford Univ. Press, Oxford, 2008). Incurrent gases
are dry from tanks. Water vapors were scrubbed from excurrent gas when it was analyzed with
the O_2/CO_2 analyzer. Thus, we calculated the ratio of $\dot{V}_I / \dot{V}_{O_2}$ and $\dot{V}_I / \dot{V}_{CO_2}$ to have a more
precise estimation of the mice ventilation avoiding confounding factors produced by changes in
metabolic rate.
738
CUBIC tissue clearing and light sheet fluorescence microscopy

CUBIC (Clear, Unobstructed Brain/Body Imaging Cocktails and Computational analysis
protocol) buffers were prepared according to Susaki et al [64]. After sufficient tissue clearing in R1
buffer, tissues were embedded in 2 % low melting agarose and then incubated in R2 solution at
room temperature overnight prior to imaging. Cleared samples were imaged using a Zeiss Z.1
light sheet fluorescence microscope (LSFM). Lung samples were imaged using a 2.5x objective
(LSFM 2.5x NA 0.1) and a 1.45 2.5X CLARITY specific chamber (Translucence Biosystems).
746
Isolation of nuclei from frozen NA
We dissected out the NA region (GFP[+]) from adult *Chat-cre; CAG-Sun1/sfGFP* mice
brainstem under fluorescence microscope. Mice were euthanized using CO2 inhalation. Brains
were acutely harvested. The NA was visualized by nucleus localized GFP fluorescence signals
and harvested based on anatomical landmarks. We collected GFP+ neurons around NA region
and avoided DMV and 12N regions which also express Chat. NA from 7 mice were dissected
and pulled together and put in liquid N_2 immediately and store at -80°C or proceed directly to
nuclei isolation. We used the same dounce dissociation method that we isolated nTS nuclei (see
method-Isolation of nuclei from frozen nTS).
756
757   **NA single-nucleus RNA sequencing and data analysis**
758   NA nuclei were processed, sequenced and analyzed using the same kit, platform and
759   software that were used for nTS (see method-nTS single-nucleus RNA sequencing and data
760   analysis). For raw data, we firstly confirmed the mitochondrial counts was < 5%, and cells above
761   200 unique feature counts were considered high quality cells and used for further analysis.
762
Under trachea injection
Mice were anesthetized with a mixture of ketamine (100 mg/kg) and xylazine (10 mg/kg) via
intraperitoneal injection. After fully anesthetized, mice were placed ventral side up in a
stereotaxic frame with body temperature maintained with a heating pad. Ophthalmic ointment
was applied to maintain eye lubrication. The extra thoracic trachea was exposed via a ventral
incision in the animal's neck. The trachea was lifted up by forceps carefully without bleed the
trachea. Using a 10 μl Hamilton glass micropipette fitted with a 32G needle, 10 μl CTB488
(concentration at 1 mg/ml, Thermo Scientific, C22841) was injected into the dorsal side of the
trachea from bilateral sides, 3 to 5 cartilage rings caudal to the larynx. The injection was done
within 15 minutes per mouse. Mouse skin was sutured and recovered on thermal pad before
returned to housing. Buprenorphine SR (0.1 mg/kg, subcutaneous injection) was used
for pain relief following surgery. For brainstem collection, mice were sacrificed 1 week after
CTB488 injection.
776
Cannula implant and microinjection into bilateral NA
Mice were anesthetized with a mixture of ketamine (100 mg/kg) and xylazine (10 mg/kg) via
intraperitoneal injection. After fully anesthetized, mice were mounted in a stereotaxic frame. A
midline incision was made to reveal Bregma, the skull was cleaned with hydrogen peroxide, and
small holes were drilled through the skull at the designated stereotaxic coordinates of NA[57].
Bilateral guide cannula affixed with two 26G stainless-steel tubing (P1 Technologies Inc.,
Roanoke, VA, USA) was stereotaxically implanted 0.5 mm above the NA region (0.7 mm caudal
to Bregma, 1.25 mm lateral to midline, 0.4 mm under the dura surface). Guide cannulas were
secured to the skull using super glue and dental cement. Matching dummy cannula were inserted
into the guide cannula and secured with a dust cap to ensure guide cannula patency. Post-

operative analgesia was provided with Buprenorphine SR (0.1 mg/kg, subcutaneous injection).
Mice recovered in their home cages for one week before challenge and drug delivery.
Starting from the baseline measurement of flexiVent assay, Prazosin (0.4 mg/ml in ultra-pure
water, P7791, Sigma), or Terazosin (0.4mg/ml in ultra-pure water, T4680, Sigma) or ultra-pure
water control was continuously microinjected (total volume 1 μl, continuously injecting for 17.5
minutes) into bilateral NA through the guided cannula using two-channel syringe pump (R462,
RWD, USA). For all the mice for NA microinjection, we kept the brainstem and confirmed
targeted cannula implantation in NA region through cryosection.
795
Statistical analysis
All statistics were calculated using Microsoft Excel and performed using GraphPad Prism
(GraphPad Software Inc., CA, USA). Unpaired student's t-test, one-way or two-way ANOVA
Tukey's multiple comparisons test were used as indicated in figure legends to examine
significant differences across mice and experimental conditions. Values of $p<0.05$ were deem
significant and lack of significant differences ($p>0.05$) are indicated by N.S. (not significant).
802
803 **Data and materials availability:** The accession number for the raw and fully processed nTS
804 single-nucleus RNA sequencing data reported in this paper is GEO: GSE200003. The accession
805 number for the raw and fully processed NA single-nucleus RNA sequencing data reported in this
806 paper is GEO: GSE211538. Both are publicly available upon publication.

References

1.	Undem BJ, Taylor-Clark T. Mechanisms underlying the neuronal-based symptoms of allergy. *J Allergy Clin Immunol*. Jun 2014;133(6):1521-34. doi:10.1016/j.jaci.2013.11.027

2.	McAlexander MA, Gavett SH, Kollarik M, Undem BJ. Vagotomy reverses established allergen-induced airway hyperreactivity to methacholine in the mouse. *Respir Physiol Neurobiol*. Jul 2015;212-214:20-4. doi:10.1016/j.resp.2015.03.007

3.	Tränkner D, Hahne N, Sugino K, Hoon MA, Zuker C. Population of sensory neurons essential for asthmatic hyperreactivity of inflamed airways. *Proc Natl Acad Sci U S A*. Aug 2014;111(31):11515-20. doi:10.1073/pnas.1411032111

4.	Talbot S, Abdulnour RE, Burkett PR, et al. Silencing Nociceptor Neurons Reduces Allergic Airway Inflammation. *Neuron*. Jul 2015;87(2):341-54. doi:10.1016/j.neuron.2015.06.007

5.	Sherrington CS. *The integrative action of the nervous system*. C. Scribner's sons; 1906:xvi, 411 p.

6.	Chen WG, Schloesser D, Arensdorf AM, et al. The Emerging Science of Interoception: Sensing, Integrating, Interpreting, and Regulating Signals within the Self. *Trends Neurosci*. 01 2021;44(1):3-16. doi:10.1016/j.tins.2020.10.007

7.	Prescott SL, Umans BD, Williams EK, Brust RD, Liberles SD. An Airway Protection Program Revealed by Sweeping Genetic Control of Vagal Afferents. *Cell*. 04 2020;181(3):574-589.e14. doi:10.1016/j.cell.2020.03.004

8.	Zhao Q, Yu CD, Wang R, et al. A multidimensional coding architecture of the vagal interoceptive system. *Nature*. Mar 2022;603(7903):878-884. doi:10.1038/s41586-022-04515-5

9.	Kaelberer MM, Buchanan KL, Klein ME, et al. A gut-brain neural circuit for nutrient sensory transduction. *Science*. 09 21 2018;361(6408)doi:10.1126/science.aat5236

10.	Muller PA, Schneeberger M, Matheis F, et al. Microbiota modulate sympathetic neurons via a gut-brain circuit. *Nature*. 07 2020;583(7816):441-446. doi:10.1038/s41586-020-2474-7

11.	Keller JA, Chen J, Simpson S, et al. Voluntary urination control by brainstem neurons that relax the urethral sphincter. *Nat Neurosci*. 09 2018;21(9):1229-1238. doi:10.1038/s41593-018-0204-3

12.	Scott GD, Fryer AD, Jacoby DB. Quantifying nerve architecture in murine and human airways using three-dimensional computational mapping. *Am J Respir Cell Mol Biol*. Jan 2013;48(1):10-6. doi:10.1165/rcmb.2012-0290MA

13.	McGovern AE, Davis-Poynter N, Yang SK, Simmons DG, Farrell MJ, Mazzone SB. Evidence for multiple sensory circuits in the brain arising from the respiratory system: an anterograde viral tract tracing study in rodents. *Brain Struct Funct*. Nov 2015;220(6):3683-99. doi:10.1007/s00429-014-0883-9

14.	Mazzone SB, Undem BJ. Vagal Afferent Innervation of the Airways in Health and Disease. *Physiol Rev*. 07 2016;96(3):975-1024. doi:10.1152/physrev.00039.2015

15.	Su Y, Barr J, Jaquish A, Xu J, Verheyden JM, Sun X. Identification of Lung Innervating Sensory Neurons and Their Target Specificity. *Am J Physiol Lung Cell Mol Physiol*. Nov 10 2021;doi:10.1152/ajplung.00376.2021

16.	Taylor-Clark T, Undem BJ. Transduction mechanisms in airway sensory nerves. *J Appl Physiol (1985)*. Sep 2006;101(3):950-9. doi:10.1152/japplphysiol.00222.2006

17.	Kim SH, Hadley SH, Maddison M, et al. Mapping of Sensory Nerve Subsets within the Vagal Ganglia and the Brainstem Using Reporter Mice for Pirt, TRPV1, 5-HT3, and Tac1 Expression. *eNeuro*. 2020 Mar/Apr 2020;7(2)doi:10.1523/ENEURO.0494-19.2020

18.   Cutsforth-Gregory JK, Benarroch EE. Nucleus of the solitary tract, medullary reflexes,
and clinical implications. *Neurology*. 03 2017;88(12):1187-1196.
doi:10.1212/WNL.0000000000003751
19.   Han W, Tellez LA, Perkins MH, et al. A Neural Circuit for Gut-Induced Reward. *Cell*. 10
2018;175(3):887-888. doi:10.1016/j.cell.2018.10.018
20.   Wang J, Irnaten M, Neff RA, et al. Synaptic and neurotransmitter activation of cardiac
vagal neurons in the nucleus ambiguus. *Ann N Y Acad Sci*. Jun 2001;940:237-46.
doi:10.1111/j.1749-6632.2001.tb03680.x
21.   Travagli RA, Hermann GE, Browning KN, Rogers RC. Brainstem circuits regulating
gastric function. *Annu Rev Physiol*. 2006;68:279-305.
doi:10.1146/annurev.physiol.68.040504.094635
22.   Affleck VS, Coote JH, Pyner S. The projection and synaptic organisation of NTS afferent
connections with presympathetic neurons, GABA and nNOS neurons in the paraventricular
nucleus of the hypothalamus. *Neuroscience*. Sep 06 2012;219:48-61.
doi:10.1016/j.neuroscience.2012.05.070
23.   Calderón MA, Linneberg A, Kleine-Tebbe J, et al. Respiratory allergy caused by house
dust mites: What do we really know? *J Allergy Clin Immunol*. Jul 2015;136(1):38-48.
doi:10.1016/j.jaci.2014.10.012
24.   Miller M, Vuong C, Garcia MF, et al. Does reduced zona pellucida binding protein 2
(ZPBP2) expression on chromosome 17q21 protect against asthma? *J Allergy Clin Immunol*. 08
2018;142(2):706-709.e4. doi:10.1016/j.jaci.2018.04.011
25.   Sui P, Wiesner DL, Xu J, et al. Pulmonary neuroendocrine cells amplify allergic asthma
responses. *Science*. 06 2018;360(6393)doi:10.1126/science.aan8546
26.   Han L, Limjunyawong N, Ru F, et al. Mrgprs on vagal sensory neurons contribute to
bronchoconstriction and airway hyper-responsiveness. *Nat Neurosci*. 03 2018;21(3):324-328.
doi:10.1038/s41593-018-0074-8
27.   Bradding P, Walls AF, Holgate ST. The role of the mast cell in the pathophysiology of
asthma. *J Allergy Clin Immunol*. Jun 2006;117(6):1277-84. doi:10.1016/j.jaci.2006.02.039
28.   Weigand LA, Myers AC, Meeker S, Undem BJ. Mast cell-cholinergic nerve interaction
in mouse airways. *J Physiol*. Jul 2009;587(Pt 13):3355-62. doi:10.1113/jphysiol.2009.173054
29.   Hitomi K, Tahara-Hanaoka S, Miki H, et al. Allergin-1 on mast cells suppresses house
dust mite-induced airway hyperresponsiveness in mice. *Int Immunol*. 08 30 2018;30(9):429-434.
doi:10.1093/intimm/dxy025
30.   Corry DB, Folkesson HG, Warnock ML, et al. Interleukin 4, but not interleukin 5 or
eosinophils, is required in a murine model of acute airway hyperreactivity. *J Exp Med*. Jan 01
1996;183(1):109-17. doi:10.1084/jem.183.1.109
31.   Tomkinson A, Duez C, Cieslewicz G, et al. A murine IL-4 receptor antagonist that
inhibits IL-4- and IL-13-induced responses prevents antigen-induced airway eosinophilia and
airway hyperresponsiveness. *J Immunol*. May 01 2001;166(9):5792-800.
doi:10.4049/jimmunol.166.9.5792
32.   Tomer R, Ye L, Hsueh B, Deisseroth K. Advanced CLARITY for rapid and high-
resolution imaging of intact tissues. *Nat Protoc*. Jul 2014;9(7):1682-97.
doi:10.1038/nprot.2014.123
33.   Undem BJ, Kajekar R, Hunter DD, Myers AC. Neural integration and allergic disease. *J
Allergy Clin Immunol*. Nov 2000;106(5 Suppl):S213-20. doi:10.1067/mai.2000.110153
34.   Potenzieri C, Meeker S, Undem BJ. Activation of mouse bronchopulmonary C-fibres by
serotonin and allergen-ovalbumin challenge. *J Physiol*. Nov 01 2012;590(21):5449-59.
doi:10.1113/jphysiol.2012.237115

35.   Wu YE, Pan L, Zuo Y, Li X, Hong W. Detecting Activated Cell Populations Using
Single-Cell RNA-Seq. *Neuron*. Oct 11 2017;96(2):313-329.e6.
doi:10.1016/j.neuron.2017.09.026
36.   Zhang C, Kaye JA, Cai Z, Wang Y, Prescott SL, Liberles SD. Area Postrema Cell Types
that Mediate Nausea-Associated Behaviors. *Neuron*. 02 03 2021;109(3):461-472.e5.
doi:10.1016/j.neuron.2020.11.010
37.   Tao J, Campbell JN, Tsai LT, Wu C, Liberles SD, Lowell BB. Highly selective brain-to-
gut communication via genetically defined vagus neurons. *Neuron*. 07 07 2021;109(13):2106-
2115.e4. doi:10.1016/j.neuron.2021.05.004
38.   Ludwig MQ, Cheng W, Gordian D, et al. A genetic map of the mouse dorsal vagal
complex and its role in obesity. *Nat Metab*. 04 2021;3(4):530-545. doi:10.1038/s42255-021-
00363-1
39.   Maley BE. Immunohistochemical localization of neuropeptides and neurotransmitters in
the nucleus solitarius. *Chem Senses*. Jun 1996;21(3):367-76. doi:10.1093/chemse/21.3.367
40.   Baptista V, Zheng ZL, Coleman FH, Rogers RC, Travagli RA. Characterization of
neurons of the nucleus tractus solitarius pars centralis. *Brain Res*. Aug 09 2005;1052(2):139-46.
doi:10.1016/j.brainres.2005.05.073
41.   Ritter S, Bugarith K, Dinh TT. Immunotoxic destruction of distinct catecholamine
subgroups produces selective impairment of glucoregulatory responses and neuronal activation. *J
Comp Neurol*. Apr 02 2001;432(2):197-216. doi:10.1002/cne.1097
42.   Mailhot-Larouche S, Deschênes L, Lortie K, et al. Assessment of Respiratory Function in
Conscious Mice by Double-chamber Plethysmography. *J Vis Exp*. 07 10
2018;(137)doi:10.3791/57778
43.   Travagli RA, Anselmi L. Vagal neurocircuitry and its influence on gastric motility. *Nat
Rev Gastroenterol Hepatol*. 07 2016;13(7):389-401. doi:10.1038/nrgastro.2016.76
44.   Babic T, Browning KN, Travagli RA. Differential organization of excitatory and
inhibitory synapses within the rat dorsal vagal complex. *Am J Physiol Gastrointest Liver Physiol*.
Jan 2011;300(1):G21-32. doi:10.1152/ajpgi.00363.2010
45.   Veerakumar A, Yung AR, Liu Y, Krasnow MA. Molecularly defined circuits for
cardiovascular and cardiopulmonary control. *Nature*. Jun 01 2022;doi:10.1038/s41586-022-
04760-8
46.   Coverdell TC, Abraham-Fan RJ, Wu C, Abbott SBG, Campbell JN. Genetic encoding of
an esophageal motor circuit. *Cell Rep*. Jun 14 2022;39(11):110962.
doi:10.1016/j.celrep.2022.110962
47.   Kupari J, Häring M, Agirre E, Castelo-Branco G, Ernfors P. An Atlas of Vagal Sensory
Neurons and Their Molecular Specialization. *Cell Rep*. 05 2019;27(8):2508-2523.e4.
doi:10.1016/j.celrep.2019.04.096
48.   Mazzone SB, Tian L, Moe AAK, Trewella MW, Ritchie ME, McGovern AE.
Transcriptional Profiling of Individual Airway Projecting Vagal Sensory Neurons. *Mol
Neurobiol*. Feb 2020;57(2):949-963. doi:10.1007/s12035-019-01782-8
49.   Nadel JA, Barnes PJ. Autonomic regulation of the airways. *Annu Rev Med*. 1984;35:451-
67. doi:10.1146/annurev.me.35.020184.002315
50.   Fryer AD, Jacoby DB. Muscarinic receptors and control of airway smooth muscle. *Am J
Respir Crit Care Med*. Nov 1998;158(5 Pt 3):S154-60.
doi:10.1164/ajrccm.158.supplement_2.13tac120
51.   Sweeney M, McDaniel SS, Platoshyn O, et al. Role of capacitative Ca2+ entry in
bronchial contraction and remodeling. *J Appl Physiol (1985)*. Apr 2002;92(4):1594-602.
doi:10.1152/japplphysiol.00722.2001

52.   Canning BJ, Fischer A. Neural regulation of airway smooth muscle tone. *Respir Physiol*. Mar 2001;125(1-2):113-27. doi:10.1016/s0034-5687(00)00208-5

53.   Pincus AB, Adhikary S, Lebold KM, Fryer AD, Jacoby DB. Optogenetic Control of Airway Cholinergic Neurons. *Am J Respir Cell Mol Biol*. 04 2020;62(4):423-429. doi:10.1165/rcmb.2019-0378MA

54.   Roman CW, Derkach VA, Palmiter RD. Genetically and functionally defined NTS to PBN brain circuits mediating anorexia. *Nat Commun*. 06 15 2016;7:11905. doi:10.1038/ncomms11905

55.   Chen J, Cheng M, Wang L, et al. A Vagal-NTS Neural Pathway that Stimulates Feeding. *Curr Biol*. 10 19 2020;30(20):3986-3998.e5. doi:10.1016/j.cub.2020.07.084

56.   Aklan I, Sayar Atasoy N, Yavuz Y, et al. NTS Catecholamine Neurons Mediate Hypoglycemic Hunger via Medial Hypothalamic Feeding Pathways. *Cell Metab*. 02 04 2020;31(2):313-326.e5. doi:10.1016/j.cmet.2019.11.016

57.   Franklin KBJ, Paxinos G. *Paxinos and Franklin's The mouse brain in stereotaxic coordinates*. Fourth edition. ed. Academic Press, an imprint of Elsevier; 2013:1 volume (unpaged).

58.   Wrenn CC, Picklo MJ, Lappi DA, Robertson D, Wiley RG. Central noradrenergic lesioning using anti-DBH-saporin: anatomical findings. *Brain Res*. Nov 18 1996;740(1-2):175-84. doi:10.1016/s0006-8993(96)00855-4

59.   Ippoliti R, Lendaro E, Bellelli A, Brunori M. A ribosomal protein is specifically recognized by saporin, a plant toxin which inhibits protein synthesis. *FEBS Lett*. Feb 24 1992;298(2-3):145-8. doi:10.1016/0014-5793(92)80042-f

60.   Moya EA, Go A, Kim CB, Fu Z, Simonson TS, Powell FL. Neuronal HIF-1α in the nucleus tractus solitarius contributes to ventilatory acclimatization to hypoxia. *J Physiol*. 05 2020;598(10):2021-2034. doi:10.1113/JP279331

61.   Reid SG, Powell FL. Effects of chronic hypoxia on MK-801-induced changes in the acute hypoxic ventilatory response. *J Appl Physiol (1985)*. Dec 2005;99(6):2108-14. doi:10.1152/japplphysiol.01205.2004

62.   Jacky JP. A plethysmograph for long-term measurements of ventilation in unrestrained animals. *J Appl Physiol Respir Environ Exerc Physiol*. Oct 1978;45(4):644-7. doi:10.1152/jappl.1978.45.4.644

63.   DRORBAUGH JE, FENN WO. A barometric method for measuring ventilation in newborn infants. *Pediatrics*. Jul 1955;16(1):81-7.

64.   Susaki EA, Tainaka K, Perrin D, Yukinaga H, Kuno A, Ueda HR. Advanced CUBIC protocols for whole-brain and whole-body clearing and imaging. *Nat Protoc*. Nov 2015;10(11):1709-27. doi:10.1038/nprot.2015.085

Acknowledgments: We thank members of the Sun lab for inputs and discussions. We thank Drs. Takaki Komiyama, David Kleinfeld and Byungkook Lim at the University of California San Diego (UCSD), Dr. Samuel Pfaff at the Salk Institute for inputs and discussions. We thank the Center for Epigenomics at UCSD for performing the single nucleus RNA-seq experiments. Work at the Center for Epigenomics was supported in part by the UCSD School of Medicine. This publication includes data generated at the UCSD Institute for Genomic Medicine utilizing an Illumina NovaSeq 6000 that was purchased with funding from a National Institutes of Health SIG grant (#S10 OD026929). Flow cytometry was performed with the support of the Flow Cytometry Research Core at the San Diego Center for AIDS Research (P30 AI036214), the VA San Diego Health Care System, and the San Diego Veterans Medical Research Foundation. We thank Jennifer Santini and Marcella Erb at the UCSD School of Medicine Microscopy Core (National Institute of Neurological Disorders and Stroke Grant P30 NS047101) for assistance with imaging. Fundings were provided by the National Institutes of Health 1R01AT011676 (X.S.), National Institutes of Health OT2OD023857 (X.S.), Tobacco-Related Disease Research Program (TRDRP) T29IR0475 (X.S.) and American Heart Association 19POST34450103 (Y.S.).

Author contributions:

Y.S. and X.S. conceived and designed experiments; Y.S. performed experiments; Y.S., A.N. and X.S. interpreted data; Y.S. prepared figures and drafted manuscript. J.X. helped with *TRAP2* and flexiVent experiments, and performed PAS staining; Z.Z., H.Y. and Y.S. analyzed single-nucleus RNA-seq data and generated figures; V.N. and L.Y. performed CLARITY of *TRAP2* mice and did confocal imaging; B.D. designed antibody cocktails for flow cytometry; E.M. performed plethysmography and analyzed the data; Y.S. and X.S. edited and revised the manuscript with input from the other authors; Y.S., J.X., Z.Z., H.Y., V.N., B.D., E.M., L.Y., A.N. and X.S. approved final version of manuscript.

Competing interests: Authors declare that they have no competing interests.

Materials & Correspondence: Correspondence and requests for materials should be addressed to Xin Sun.

All figures and extended data figures have been deposited in figshare, and the private sharing links are provided here to access.

Fig. 1: https://figshare.com/s/c93fff81efdb06965a55
Fig. 2: https://figshare.com/s/45c42b6036ecf8f24a7d
Fig. 3: https://figshare.com/s/366a0e6b67eb9abef37f
Fig. 4: https://figshare.com/s/37ccb9018f4653d0ecfd
Fig. 5: https://figshare.com/s/95a526de44919eeaff88
Fig. 6: https://figshare.com/s/1bd52dd2ff58a5853cca
Extended Data Fig. 1: https://figshare.com/s/67ca20595b020a9b61cf
Extended Data Fig. 2: https://figshare.com/s/f8a2cf4e4029a0bf806d
Extended Data Fig. 3: https://figshare.com/s/2ce8910e15a77da11a90
Extended Data Fig. 4: https://figshare.com/s/07566d37b1a37b109190
Extended Data Fig. 5: https://figshare.com/s/f0d231f3fd9c71da8167
Extended Data Fig. 6: https://figshare.com/s/be7a508b61b645125db9

Extended Data Fig. 7: https://figshare.com/s/480e03f3d08e98464bd4
Extended Data Fig. 8: https://figshare.com/s/4f1489d96416fc9071d6
Extended Data Fig. 9: https://figshare.com/s/1a1bc40336f6c2f61996
Extended Data Fig. 10: https://figshare.com/s/4fe23454fc61b4e7388a
Extended Data Fig. 11: https://figshare.com/s/612f575fec98429ae61e
Extended Data Fig. 12: https://figshare.com/s/05e0065abc0baac90be1
Extended Data Fig. 13: https://figshare.com/s/2105e42d4cf4c82b2a2d
Extended Data Fig. 14: https://figshare.com/s/dbefeabb2dcfe5e58731
Extended Data Fig. 15: https://figshare.com/s/4e0c530edb25cd77617e
Extended Data Fig. 16: https://figshare.com/s/79b7907b8190d8ce2e7e
Extended Data Fig. 17: https://figshare.com/s/1c1bfd80b34556126476
Extended Data Fig. 18: https://figshare.com/s/70f4f3730099261788be
Extended Data Fig. 19: https://figshare.com/s/1903a369125fcfb9e701
Extended Data Fig. 20: https://figshare.com/s/d50fe7c0874ca27fa01d
Extended Data Fig. 21: https://figshare.com/s/086052bb8b27fe4dd2bb
Extended Data Fig. 22: https://figshare.com/s/a19716980cd22d3cf2cf
Extended Data Fig. 23: https://figshare.com/s/ad9bc446b926845a0942
Extended Data Fig. 24: https://figshare.com/s/789a21c020440c3fdbfb

Figures and figure legends

Su et al. Fig. 1

Figures and figure legends

Su et al. Fig. 1

a HDM or Saline — wild-type. 1st Day0, 2nd Day7, 3rd Day14, 4th Day21, analysis 2hrs after

b FOS Saline / FOS HDM (-7.92mm)

c FOS+ cells/40um — Saline, HDM ***

d FOS c-Kit^w-sh/+ Saline / HDM; FOS c-Kit^w-sh/w-sh Saline / HDM (-7.92mm)

e FOS+ cells/40um — c-kit^w-sh/+, c-kit^w-sh/w-sh; Saline, HDM **** **** *

f HDM, anti-IL-4/Ctrl — wild-type. anti-IL-4/Ctrl i.p. before each HDM. 1st Day0, 2nd Day7, 3rd Day14, 4th Day21, flexiVent 24hrs after

g FOS Ctrl+HDM / FOS anti-IL-4+HDM (-7.92mm)

h FOS+ cells/40um — Ctrl, anti-IL-4 / HDM **

i Contralateral Ipsilateral FOS Sham+HDM / Contralateral Ipsilateral FOS Vagotomy+HDM (-7.92mm)

j FOS+ cells/40um — Sham, Vagotomy / HDM; Ipsilateral, Contralateral ***

k HDM or Saline — TRAP2;Ai14. 4-OHT i.p. right after each challenge. 1st Day0, 2nd Day7, 3rd Day14, 4th Day21, analysis Day28

l tdTom Saline / tdTom HDM

m cells/mm^3 — Saline, HDM; region of interest ***

n HDM, 4-OHT — TRAP2;Ai14;DTR. 4-OHT i.p. right after each HDM. 1st Day0, 2nd Day7, 3rd Day14, 4th Day21, DTX/vehicle Day28, flexiVent 24hrs after 5th HDM Day35

o tdTom vehicle+HDM / tdTom DTX+HDM (-7.92mm)

p tdTom+ cells/40um — vehicle, DTX / HDM ****

q Rrs (cm H2O.s/ml) Differences over baseline vs Mch (mg/ml) — Saline N=5, HDM N=6 ** ***

r Ers (cm H2O.s/ml) Differences over baseline vs MCh (mg/ml) — Saline N=5, HDM N=6 * ***

s Rrs (cm H2O.s/ml) Differences over baseline vs MCh (mg/ml) — vehicle+HDM N=4, DTX+HDM N=7 ** ****

t Ers (cm H2O.s/ml) Differences over baseline vs MCh (mg/ml) — vehicle+HDM N=4, DTX+HDM N=7 * ****

Fig. 1. nTS neurons were activated upon repeated allergen challenges in lung. (**a**) Experiment scheme for HDM treatment in wild-type mice. (**b**, **c**) Representative FOS antibody staining (b) and quantification (c) showing increased FOS$^+$ cells after 4th HDM. Dashed circles outline nTS at the indicated Bregma level, as in subsequent related staining panels. Each data point represents the result from one animal, as in all subsequent graphs. (**d**, **e**) Representative FOS antibody staining (d) and quantification (e) showing decreased FOS$^+$ cells in the nTS of *c-Kit$^{w-sh/w-sh}$*, compared to *c-Kit$^{w-sh/+}$* mice after HDM. (**f**) Experiment scheme for treatment with IL-4 neutralizing antibody in wild-type mice. (**g**, **h**) Representative FOS antibody staining (g) and quantification (h) showing decreased FOS$^+$ cells in the nTS with anti-IL-4 neutralizing antibody treatment compared to isotype control antibody treatment. (**i**, **j**) Representative FOS antibody staining (i) and quantification (j) showing reduced signals on ipsilateral side with vagotomy, compared to contralateral side. (**k**) Experiment scheme for labeling allergen-activated neurons in *TRAP2; Ai14* mice. (**l**, **m**) Coronal view of CLARITY cleared hemisphere brainstem (l, dashed circles) and quantification (m) showing increased tdTomato$^+$ cells after HDM. (**n**) Experiment scheme for ablating allergen-activated neurons in *TRAP2; Ai14; DTR* mice. (**o**, **p**) Representative nTS sections (o) and quantification (p) showing reduced tdTomato signals following DTX injection. (**q**, **r**) FlexiVent-measured maximal resistance (Rrs, q) and elastance (Ers, r) of the wild-type airway following increasing doses of MCh, demonstrating hyperreactivity after HDM challenge compared to saline control. (**s**, **t**) Maximal resistance (Rrs, s) and elastance (Ers, t) showing blunted airway hyperreactivity in DTX+HDM group of *TRAP2; Ai14; DTR* mice compared to vehicle controls. Statistical analysis was performed separately for each methacholine (MCh) concentration. Scale bars, 200 μm in b, d, g, i, o. Unpaired student's t test was used for C, H, M and P, two-way ANOVA was used for e, j and q-t. Error bars represent means ± SEM. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Fig. 2. Single nucleus transcriptomic signatures of the nTS. (**a**) Diagram illustrating the relative locations of the nucleus of solitary tract (nTS), area postrema (AP), dorsal motor nucleus of the vagus (DMV) and nucleus ambiguous (NA). (**b**) Uniform manifold approximation and projection (UMAP) plots of snRNA-seq data showing 13 nTS neuron types. (**c-f**) UMAP plots showing expression of pan-neuronal markers *Rbfox3* (c) and *Snap25* (d) in all clusters, excitatory neuronal marker *Scl17a6* enriched in Clusters 1, 2, 5, 9 and 12 (e), inhibitory neuronal marker *Slc32a1* enriched in Clusters 3, 4, 6,7, 8, 10, 11 and 13 (f). (**g**) Dot plot showing top marker genes for each cluster. Gene names in red were used for validation by UMAP or RNAscope *in situ* hybridization as shown in Fig. 2h-q and Extended Data Fig. 7e-l. (**h-q**) UMAP plots (h, j, l, n, p) and RNAscope (i, k, m, o, q) showing expression of marker genes in the nTS excitatory clusters. Bregma levels shown are tailored to the layers with optimal signal for each marker. For i, k, m, o, q, outlines indicate nTS regions at the indicated Bregma level. Scale bars, 200 μm.

Su et al. Fig. 3

Fig. 3. *Dbh*⁺ neurons in the nTS were preferentially activated upon allergen challenge in lung. (**a**) UMAP plots showing the corresponding 13 neuron clusters in saline- (left) and HDM-challenged (right) nTS. (**b**) Dot plot of Fos expression in excitatory nTS clusters. Numbers on the upper row denote Log2 average expression levels, and numbers on the bottom row denote percentages of expressing cells. (**c**, **d**) Representative double *in situ* (c) and quantification (d) showing overlap between *Dbh* and *Fos* expression. (**e**, **f**) Representative FOS staining (e) and quantification (f) of *Dbh-cre; Rosa-lxl-tdTomato* (*Dbh-cre; Ai14*) nTS showing partial overlap between FOS and tdTomato. For c, e, outlines indicate nTS at the indicated Bregma, boxed areas magnified on the right, arrowheads indicate overlapped expression. Scale bars, 200 μm.

Su et al. Fig. 4

Fig. 4. *Dbh⁺* neurons in the nTS mediates airway hyperreactivity. (**a**) Experiment scheme for chemical ablation of *Dbh⁺* nTS neurons. (**b**) Representative DBH antibody staining showing reduced DBH⁺ cells after ablation. Dashed circles outline dorsomedial part of nTS region at the indicated Bregma levels. (**c**, **d**) FlexiVent data showing blunted airway hyperreactivity after DBH-SAP. (**e**) Experiment scheme for genetic ablation of *Dbh⁺* nTS neurons. (**f**) Representative *Dbh in situ* hybridization showing reduced expression after DTX. (**g**, **h**) FlexiVent data showing blunted airway hyperreactivity after DTX. (**i**) Experiment scheme for chemogenetic inhibition of *Dbh⁺* neurons. (**j**, **k**) FlexiVent data showing blunted airway hyperreactivity after CNO injection. (**l**) Experiment scheme for chemogenetic activation of *Dbh⁺* neurons. (**m**, **n**) FlexiVent data showing increased airway hyperreactivity after CNO, in place of the 4th HDM. All groups received the 1st-3rd HDM challenge. Scale bars, 200 μm in b, f. Two-way ANOVA was used for c, d, g, h, j, k, m, n. Statistical analysis at each MCh concentration was performed separately. Error bars represent means ± SEM. *p<0.05, **p<0.01, ****p<0.0001.

Su et al. Fig. 5

1192

Fig. 5. Parasympathetic neurons in the NA downstream of the nTS are necessary and sufficient for allergen-induced airway hyperreactivity. (**a**) Representative brainstem section at indicated Bregma level showing specificity of the injection of AAV-flex-tdTomato into bilateral nTS (outlined) of *Dbh-cre* mouse. (**b**) In the same mouse shown in A, tdTomato$^+$ nerves project to NA region (VAChT$^+$). Boxed areas are magnified below. (**c**) Injection of AAV-flex-tdTomato into NA of *Chat-cre* mouse. (**d**, **e**) In the same mouse shown in c, light sheet images showing tdTomato$^+$ fibers innervate the trachea (d, arrowheads) and extrapulmonary bronchi (e, arrowheads). (**f**) Confocal stack (300 µm) of extrapulmonary bronchi showing NA-originated tdTomato$^+$ fibers innervate postganglionic parasympathetic ganglia (VAChT$^+$, arrowheads). Boxed areas are magnified on the right. (**g**) In *Chat-cre; Ai14* mice, injection of CTB488 in dorsal trachea led to overlapping CTB488$^+$ and tdTomato$^+$ signals in the NA (arrowheads). (**h**) Scheme for chemogenetic inhibition of NA *Chat$^+$* neurons. (**i**) AAV-flex-hM4D-mCherry signals in the NA of *Chat-cre* mouse. (**j**, **k**) FlexiVent measurements indicate blunted airway hyperreactivity after CNO injection. (**l**) Scheme for chemogenetic activation of NA *Chat$^+$* neurons. (**m**) AAV-DIO-hM3D-mCherry signals in the NA of *Chat-cre* mouse. (**n**, **o**) FlexiVent measurements indicate increased airway hyperreactivity after CNO, in place of the 4th HDM. Scale bars, 100 µm in c, i, m, 200 µm in f (50 µm in magnified views), 500 µm in a, b (50 µm in magnified views) and g (100 µm in magnified views). Two-way ANOVA analysis for j, k, n and o were performed separately at each MCh concentration. Error bars represent means ± SEM. *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

1213
1214

Fig. 6. Blocking norepinephrine receptors in the NA blunted allergen-induced airway hyperreactivity. (**a**) UMAP plot of NA snRNA-seq data showing 3 NA neuron types. (**b**) Dot plot showing top marker genes for each cluster. Gene names in red were used for validation by UMAP plots or *in situ* hybridization as shown in c-h. (**c-h**) UMAP plots (left) and *in situ* hybridization images from Allen Brain Atlas (right) showing expression of marker genes. (**i, j**) Double *in situ* hybridization showing *Chat*[+] neurons in the NA express norepinephrine receptors *Adra1a* and *Adra1b* (arrowheads). (**k, l**) UMAP plots showing expression of *Adra1a* (k) and *Adra1b* (l) in all NA clusters. (**m, n**) Injection of CTB488 to dorsal trachea in wild-type mice labeled neurons in the NA that express *Chat* and *Adra1a* (m) and *Adra1b* (n) by RNAscope *in situ* hybridization (arrowheads). (**o**) Experiment scheme for norepinephrine receptor antagonist treatment. (**p, q**) FlexiVent data showing decreased hyperreactivity after Prazosin or Terazosin microinjection into bilateral NA, compared to vehicle control injection. (**r**) A diagram illustrating the complete allergen neuroimmune circuit. Scale bars, 100 µm in i, j, m, n, 200 µm in d, f, h. Two-way ANOVA analysis for p and q were performed separately at each MCh concentration. Error bars represent means ± SEM. *p<0.05, ****p<0.0001.

Extended Data-figures and figure legends

Su et al. Extended Data Fig.1

Extended Data Fig. 1. nTS neurons were activated after allergen challenge in lung.
(**a**) PAS staining shows that HDM regimen effectively induced mucus-secreting airway goblet cells as expected. (**b-u**) FOS antibody staining two hours after the 4th challenges in serial sections of the whole brainstem. Increased FOS$^+$ cells were enriched in the dorsomedial part of the nTS region from Bregma -7.20 mm to -8.08 mm in HDM-treated mice (yellow outlined) compared to the saline-treated control mice. (**t**) Quantification of total number of FOS$^+$ cells after HDM or Saline control in the whole nTS regions (Bregma -6.24 mm to -8.24 mm). Each serial section was at 20 µm thick, 80 µm interval in-between sections, so the number represent ¼ (500 µm) of the whole nTS regions. Each data point represents result from one animal. Scale bars, 200 µm in b-u. Unpaired student's t test was used for t. Error bars represent means ± SEM. ***$p<0.001$.

Extended Data Fig. 2. Allergen-activated neurons were not located in the adjoining AP, DMV or 12N regions in the brainstem. (**a**) *Fos in situ* hybridization from brainstem section of naïve mice after HDM challenge in lung showing most *Fos* RNA signals do not overlap with genes (*Gfral* or *Glp1r*) that are enriched in the AP. (**b**) FOS antibody staining from brainstem section of *Chat-cre; Ai14* mice showing most FOS[+] cells locate in the outlined nTS region and are not tdTomato[+]. There were strong tdTomato signals in adjoining DMV and 12N regions. Outlines in a and b indicate nTS regions at different Bregma. Scale bars, 200 μm.

Su et al. Extended Data Fig.3

Extended Data Fig. 3. nTS neurons were activated only after repeated allergen challenge in lung. (**a**, **d**, **g**) Experiment scheme for the 1st, 2nd, or 3rd HDM challenge in wild-type mice. (**b**, **c**, **e**, **f**, **h**, **i**) FOS antibody staining (b, e, h) and quantification (c, f, i) showing the lack of statistically significant increase in FOS$^+$ cells after the 1st, 2nd, or 3rd HDM. Outlines in b, e and h indicate nTS region at Bregma -7.48mm. Scale bars, 200 μm. Unpaired student's *t* test was used for c, f, i. Error bars represent means ± SEM. N.S., not significant, P ≥ 0.05.

Su et al. Extended Data Fig.4

1333
1334
1335 **Extended Data Fig. 4.** *Fos in situ* **hybridization shows increased signals after allergen**
1336 **challenge in lung.** (**a**) Experiment scheme for HDM challenge and tissue collection in wild-type
1337 mice for *Fos in situ* hybridization. (**b**) *Fos* RNA signal in the nTS is increased in the HDM
1338 group, compared to saline control. Outlines in B indicate nTS region at Bregma -7.92mm. Scale
1339 bars, 200 μm.

1350
1351
Extended Data Fig. 5. Serial sectioning and quantification of *TRAP2; Ai14* mice nTS
confirmed an increase of activated tdTomato[+] neurons in the nTS of HDM-treated mice
compared to saline-treated controls. (**a**) Experiment scheme for labeling HDM-activated
neurons in *TRAP2; Ai14* mice. (**b**, **c**) Representative image (b) and quantification (c) from
sections of *TRAP2; Ai14* mice showing increased tdTomato signals in the nTS after HDM at
Bregma -7.92 mm. Outlines in b indicate nTS region. Scale bars, 200 μm. Unpaired student's *t*
test was used for c. Error bars represent means ± SEM, ***p<0.001.

Extended Data Fig. 6. Genetic ablation of HDM-activated neurons in the nTS did not affect goblet cell metaplasia or immune cell infiltration. (a) Experimental scheme for DTX ablation of HDM-activated nTS neurons in *TRAP2; Ai14; DTR* mice. **(b**, **c)** PAS staining (b) and qPCR of *Muc5ac* (c) showing no change in HDM-induced goblet cell metaplasia response between nTS vehicle-injected and DTX-injected *TRAP2; Ai14; DTR* mice. **(d-i)** Flow cytometry analyses of eosinophils (Eos, d), innate lymphoid cells (ILC2s, e) and T-helper type 2 cells (Th2, f), and qPCR of *Il4* (g), *Il5* (h) and *Il13* (i) from whole lungs of vehicle-injected or DTX-injected mice showing no statistically significant differences between groups. Scale bars, 100 µm in b. Unpaired student's *t* test was used for c-i. Error bars represent means ± SEM. N.S., not significant.

1371

Extended Data Fig. 7. snRNA-seq provided a transcriptomic atlas of nTS. (a-d) UMAP
plots showing little to no expression of glia marker genes-*GFAP* (a), *Olig1* (b), *Aqp4* (c) or
Tmem119 (d) in the neuron clusters. (**e-l**) UMAP plots (left) and RNA *in situ* hybridization
images from the Allen Brain Atlas (right) showing expression of marker genes in the nTS
inhibitory clusters as labeled. Genes plotted here were indicated in red in Fig. 2g. Scale bars, 100
μm for e-l. (**m-o**) Dot plot showing nTS neuron expression of factors involved in neuronal signal
reception and transmission. Gene names in red are ones with notable varying expression across
the 13 nTS clusters.

1393
**Extended Data Fig. 8. *Dbh*[+] neurons in the nTS were activated upon allergen challenge in
lung.** (**a**) UMAP plot pseudo-colored to show representation of neurons in all 13 nTS clusters
from wild-type, saline-treated or HDM-treated mice. (**b**) Dot plot of *Fos* expression in individual
inhibitory nTS clusters after either saline or HDM challenge to the lung. Numbers on upper row
denote Log_2 average expression levels, and numbers on bottom row denote percentages of
expressing cells. Clusters 6, 7 and 10 populations (boxed) show decreased proportions after
HDM challenge compared to saline control.

1401
1402
1403 **Extended Data Fig. 9. Expression pattern of *Dbh* in the whole nTS. (a)** *Dbh* RNAscope *in*
1404 *situ* hybridization on serial sections of naïve brainstem that covers most of the whole nTS
1405 regions from Bregma -6.96 mm to Bregma -8.08 mm. Scale bar, 200 µm.

1406
1407
1408 **Extended Data Fig. 10. Double *in situ* hybridization between *Fos* and *Cnr1*, or *Th*, or *Cck***
1409 **after HDM challenge. (a-c)** Double RNAscope *in situ* hybridization showing some overlap
1410 between *Fos* and *Cnr1* (a) or *Th* (b), and little to no overlap between *Fos* and *Cck* (c) in the nTS
1411 at 1.5 hours after HDM challenge to lung. Outlines in a-c indicate nTS region at Bregma -7.92
1412 mm. Scale bars, 200 μm.

1413
1414
1415 **Extended Data Fig. 11. *Dbh*[+] neurons are genuinely captured by *Dbh-cre; tdTom* labeling.**
1416 (**a**, **b**) Double *in situ* hybridization representative images (a) and quantification (b) showing high
1417 overlap between *Dbh* and *tdTomato* in the nTS of *Dbh-cre; tdTomato* mice. Outlines in a
1418 indicate nTS region at Bregma -7.84 mm. Arrowheads in a indicate cells with overlapped
1419 signals. Scale bars, 200 μm.
1420

1421
1422
1423
Extended Data Fig. 12. No change in the expression of other excitatory nTS neuronal
subsets after anti-DBH-SAP injection into nTS. (**a-f**) Representative *in situ* hybridization of
nTS excitatory cluster marker genes *Dbh*, *Gli3*, *Sncg*, *Slc17a7*, *Mafa* (a-e) and quantification (f,
each datapoint represents the average number from one mouse) showing positive cells in nTS
region following DBH-SAP bilateral injection into wild type nTS, compared to blank-SAP
treatment. Outlines in a-e indicate nTS regions at different Bregma. Scale bars, 200 μm in a-e.
Unpaired student's *t* test was used for f. Error bars represent means ± SEM. N.S., not significant,
P ≥ 0.05; ***P < 0.001.

Su et al. Extended Data Fig.13

1432
1433
Extended Data Fig. 13. Chemical depletion of *Dbh*+ neurons in the nTS reduced FOS+ cells
after HDM challenge in lung while had no effect on airway hyperreactivity on naïve mice.
(**a**) Experiment scheme for detection of FOS+ cells after DBH-SAP treatment in wild-type mice.
(**b**, **c**) Representative staining (b) and quantification (c) showing reduced FOS+ cells after
ablation of *Dbh*+ neurons in the nTS following DBH-SAP treatment scheme. (**d**, **e**) FlexiVent
data showing that DBH-SAP bilateral injection into nTS of naïve mice had no effect on airway
constriction in response to increasing doses of methacholine. Outlines in b indicate nTS region at
Bregma -7.92 mm. Scale bars, 200 μm in b. Unpaired student's *t* test was used for c, two-way
ANOVA was used for d, e, statistical analysis was performed separately at each MCh
concentration. Error bars represent means ± SEM, *P < 0.05; N.S., not significant, P ≥ 0.05.

1444
1445
Extended Data Fig. 14. Chemical depletion of *Dbh*[+] neurons did not affect allergen-induced
goblet cell metaplasia or cytokine genes expressions. (**a**) Experiment scheme for flexiVent
measurement after DBH-SAP treatment in wild-type mice. (**b**) PAS staining showing no change
in HDM-induced goblet cell metaplasia response between blank-SAP-injected and DBH-SAP-
injected wild-type mice. (**c-f**) Whole lung qPCR of goblet cell marker *Muc5ac* (c) and key type 2
immunity cytokine genes *Il4* (d), *Il5* (e) and *Il13* (f) showing no statistic difference between
blank-SAP-injected and anti-DBH-SAP-injected wild-type mice. Scale bars, 100 μm in b.
Unpaired student's *t* test was used for c-f. Error bars represent means ± SEM. N.S., not
significant, P ≥ 0.05.

Su et al. fig. Extended Data Fig.15

1455
1456
Extended Data Fig. 15. Genetic depletion of *Dbh*+ neurons did not affect allergen-induced
goblet cell metaplasia or cytokine gene expressions. (**a**) Experiment scheme for DTX injection
on *Dbh-cre; DTR* mice. (**b**) PAS staining showed no change in HDM-induced goblet cell
metaplasia response between vehicle- and DTX-injected *Dbh-cre; DTR* mice. (**c-f**) qPCR of
goblet cell marker *Muc5ac* (c), key type 2 immunity cytokines *Il4* (d), *Il5* (e) and *Il13* (f) in
vehicle- or DTX-injected *Dbh-cre; DTR* mice. Scale bars, 100 µm in b. Unpaired student's *t* test
was used for c-f. Error bars represent means ± SEM. N.S., not significant.
1464

Su et al. Extended Data Fig.16

1465
1466
Extended Data Fig. 16. Genetic inactivation of *Dbh*⁺ neurons in the nTS did not affect
allergen-induced goblet cell metaplasia or immune cell infiltration. (a) Experiment scheme
for *Dbh*⁺ neuron inhibition following AAV-flex-hM4D-mCherry injection in *Dbh-cre* mice. (b)
Representative mCherry signals (with Dsred antibody staining, at Bregma -7.48 mm) following
bilateral AAV-hM4D-mCherry injection in the nTS (but not AP). (c, d) PAS staining (c) and
qPCR of *Muc5ac* (d) showed no change in HDM-induced goblet cell metaplasia response
between vehicle- and CNO-injected lungs after AAV-flex-hM4D-mCherry injection into
bilateral nTS. (e-i) Flow cytometry analyses of immune cells (e-g), and qPCR of *Il5* (h) and *Il13*

(i) from whole lungs showing no change in HDM-induced immune cell infiltration between vehicle- and CNO-injected lungs after AAV-flex-hM4D-mCherry injection into bilateral nTS. Outlines in b indicate dorsomedial part of nTS region at Bregma -7.48 mm. Scale bars, 200 μm in b, 100 μm in c. Unpaired student's *t* test was used for d-i. Error bars represent means ± SEM. N.S., not significant.

Extended Data Fig. 17. Genetic activation of *Dbh*+ neurons in the nTS had no effect on airway hyperreactivity on naïve mice. (**a**) Representative mCherry signals (with Dsred antibody staining) following bilateral AAV-DIO-hM3D-mCherry injection in the nTS (but not AP). (**b**) Experiment scheme for CNO activation of nTS *Dbh*+ neurons on naïve mice. Both groups received the 1st-3rd saline challenge. (**c**, **d**) FlexiVent data showing that CNO activation (4th dose) of *Dbh*+ nTS neurons had no effect on airway constriction in response to increasing doses of methacholine. Outlines in a indicate nTS region at Bregma -7.48 mm. Scale bars, 200 µm in a. Two-way ANOVA was used for c, d, statistical analysis was performed separately at each MCh concentration. Error bars represent means ± SEM. N.S., not significant, $P \geq 0.05$.

1507
1508
Extended Data Fig. 18. Genetic activation of *Dbh*+ neurons in the nTS did not affect
allergen-induced goblet cell metaplasia or immune cell infiltration. (**a**) Experiment scheme
for CNO activation of nTS *Dbh*+ neurons on sensitized mice. Both groups received the 1st-3rd
HDM challenge. (**b, c**) PAS staining (b) and qPCR of *Muc5ac* (c) from whole lungs showing no
change in HDM-induced goblet cell metaplasia response following 4th *Dbh*+ neuronal activation
in the nTS compared to 4th saline controls. (**d-h**) Flow cytometry analyses of immune cells (d-f),
and qPCR of *Il5* (g) and *Il13* (h) from whole lungs showing no change in HDM-induced immune
cell infiltration following 4th *Dbh*+ neuronal activation in the nTS compared to 4th saline controls.
Scale bars, 100 μm in b. One-way ANOVA was used for c-h. Error bars represent means ± SEM.
N.S., not significant, P ≥ 0.05; *P < 0.05; **P < 0.01.

Su et al. Extended Data Fig.19

1519

1520

Extended Data Fig. 19. Genetic activation of *Dbh*$^+$ neurons in the nTS did not change respiratory parameters in response to normal levels of O_2 and CO_2, as measured by plethysmograph. (a-g) Measurements of ventilation showing no change in respiratory parameters including minute ventilation ($\dot{V}_I$, a) , respiratory frequency (R_f, b) and tidal volume (V_t, c), as well as measurements for metabolic rate that includes the $\dot{V}_{O_2}$ (d) and $\dot{V}_{CO_2}$ (e), and the "real" ventilation considering the metabolic rate (Vi/VO$_2$, f and Vi/VCO$_2$, g) across groups. $\dot{V}I$, inspired ventilation, product of frequency and tidal volume which was normalized to body mass ventilation (ml/min·kg). $\dot{V}_{O_2}$, oxygen consumption, $\dot{V}_{CO_2}$, carbon dioxide. The ratio of Vi/VO$_2$ and Vi/VCO$_2$ provide a precise assessment of ventilation by avoiding confounding factors produced by changes in metabolic rate. One-way ANOVA was used for a-g. Error bars represent means ± SEM. N.S., not significant, $P \geq 0.05$.

Su et al. Extended Data Fig.20

1532
1533
Extended Data Fig. 20. Parasympathetic neurons in the NA, but not the DMV or 12N,
project to the airway-innervating postganglionic neurons. (**a-d**) tdTomato[+] neurons/ganglia
(arrowheads) reside in the trachea, main bronchi (a) and main airway entrance (hilum) of lung
lobes (b-d) of *Chat-cre; tdTomato* mice. (**e**) tdTomato[+] nerves from NA innervate postganglionic
parasympathetic ganglia (labeled with VAChT antibody, arrowhead) in the extrapulmonary
airway, rather than directly on smooth muscle cells (labeled with a-SMA antibody). (**f**)
Stereotaxic injection of AAV-flex-tdTomato into bilateral DMV (also labeled neurons in ventral
adjoining 12N) of *Chat-cre* mice. Boxed area in #1 was magnified below. Ovals in #1 outline
DMV region. (**g**) tdTomato[+] nerves from DMV were found passing through the space between
trachea and esophagus (arrowheads). Scale bars, 100 μm in e, f, 400 μm in g.

Su et al. Extended Data Fig.21

Extended Data Fig. 21. Genetic activation of *Chat*[+] neurons in the NA had no effect on
airway hyperreactivity on naïve mice. (**a**) Experiment scheme for CNO activation of NA
Chat[+] neurons on naïve mice. Both groups received the 1st-3rd saline challenge. (**b**, **c**) FlexiVent
data showing that CNO activation (4th dose) of *Chat*[+] NA neurons had no effect on airway
constriction in response to increasing doses of methacholine. Two-way ANOVA was used for b,
c, statistical analysis was performed separately at each MCh concentration. Error bars represent
means ± SEM. N.S., not significant, $P \geq 0.05$.

Su et al. Extended Data Fig.22

1558

1559

Extended Data Fig. 22. Parasympathetic neurons in the DMV or 12N did not mediate

allergen-induced airway hyperreactivity. (**a**) Experiment scheme for chemogenetic inhibition

of *Chat+* neurons in the DMV. (**b**, **c**) FlexiVent data showing no change in airway

hyperreactivity after CNO injection. (**d**) Experiment scheme for chemogenetic activation of

Chat+ neurons in the DMV. (**e**, **f**) FlexiVent data showing no change in airway hyperreactivity

after CNO injection, in place of the 4th HDM. Two-way ANOVA was used for b, c, e and f,

analysis was performed separately at each MCh concentration. Error bars represent means $\pm$

SEM. N.S., not significant, $P \geq 0.05$.

1568

Su et al. Extended Data Fig.23

a

Ligands in lung → Receptors in vagal ganglia

b

Ligands in vagal ganglia → Receptors in nTS

c

Ligands in nTS → Receptors in NA

1569
1570

Extended Data Fig. 23. A predicted neuro-immune interaction network between the lung and the brainstem. (**a-c**) A Predicted neuron-immune interaction network between the lung, *Trpv1*$^+$ neurons in the vagal ganglia (a), *Dbh*$^+$ neurons in the nTS (b) and *Chat*$^+$ neurons in the NA (c) of the brainstem (connected by lines, arrows at top show the signaling direction). Red dashed lines outline genes participate in immune responses and blue dashed lines outline genes participate in neural signaling pathways.

Su et al. Extended Data Fig.24

Extended Data Fig. 24. Cannula plantation in bilateral NA. (a) Representative cryosection (99 μm) of the NA showing bilateral cannula implantation to target NA. Arrowheads indicate the tract of cannula implantation. *Chat*[+] neurons in the NA (circled) were labeled with VAChT antibody. Scale bar, 0.5 mm.