Human FAM111A inhibits vaccinia virus replication by degrading viral uncoating factor I3 and is antagonized by poxvirus host range factor SPI-1

Junda Zhu[1*], Xintao Gao[2*], Zihui Zhang[1], Yining Deng[1], Shijie Xie[1], Shuning Ren[1], Yarui Li[1], Hua Li[1], Kang Niu[1], Shufang Fu[1], Yinü Li[2], Bernard Moss[3], Wenxue Wu[1#], Chen Peng[1#]

[1]National Key Laboratory of Veterinary Public Health, College of Veterinary Medicine, China Agricultural University, Beijing, China 100193

[2]Biotechnology Research Institute, Chinese Academy of Agricultural Sciences, Beijing, China 100081

[3]Laboratory of Viral Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, United States of America 20892

[*]Contributed equally

[#]Corresponding authors: E-mail: wuwenxue@cau.edu.cn (W.W.); pengchenea@cau.edu.cn (C.P.)

Abstract

Poxviruses are large double-stranded DNA viruses that infect a wide range of animals including humans. Smallpox virus, the most notorious poxvirus, was eradicated globally in

the 1970s. Since then, other members of the poxvirus family, such as monkeypox virus (MPXV) are still posing a great threat to public health. Vaccinia virus (VACV) is a prototypic poxvirus and was used as the vaccine strain for smallpox eradication. VACV encodes a serine protease inhibitor 1 (SPI-1) conserved in all orthopoxviruses, which has been recognized as a host range factor for modified vaccinia virus Ankara (MVA), an approved smallpox vaccine and a promising vaccine vector. FAM111A, a nuclear protein that regulates host DNA replication, was previously identified as a putative target of VACV-ΔSPI-1, which was unable to replicate in human cells unless FAM111A was repressed. Nevertheless, the detailed antiviral mechanisms of FAM111A were unresolved. Here, we show that FAM111A is a potent antiviral factor for VACV-ΔSPI-1 and MVA. Deletion of FAM111A was able to rescue the replication of MVA and VACV-ΔSPI-1. Overexpression of FAM111A significantly reduced viral DNA replication and virus titers but did not affect viral early gene expression. The antiviral effect of FAM111A required its functional serine protease domain and DNA binding domain but not the PCNA-interacting motif. We further discovered that FAM111A translocated into the cytoplasm upon VACV infection and this process was activated through the cGAS-STING signaling pathway. Infection-triggered FAM111A degraded the nuclear pore complex via its serine protease activity, translocated to the cytoplasm, and interacted with and promoted the degradation of virus uncoating factor I3 in a DNA-dependent manner. Interestingly, the serine protease activity of FAM111A was only needed for nuclear export but not I3 degradation. Further analysis suggested I3 was degraded through autophagy. Moreover, VACV SPI-1 was found primarily in the nucleus of infected cells and antagonized FAM111A by prohibiting its

nuclear export. Mutations in the SPI-1 reactive-site loop (RSL) abolished its anti-FAM111A activity. SPI-1 from MPXV and lumpy skin disease virus could also inhibit human FAM111A. Our findings reveal the detailed mechanism by which FAM111A functions to restrict a cytoplasmic DNA virus and provide explanations for the immune evasive function of VACV SPI-1.

Introduction

During their co-evolution, viruses and their hosts have evolved numerous strategies to antagonize each other. Upon viral infection, cells initiate various programs, such as the induction of type I interferons and the expression of an armory of antiviral factors that can restrict viral invasion, gene expression, DNA replication, and virion assembly. More recent findings point to the fact that some host defense factors are multifunctional and can display potent antiviral activity in a way that is distinct from their regular cellular functions[1-4]. One such example is FAM111A (family with sequence similarity 111 member A), a protein found exclusively in mammals harboring a fully functional trypsin-like C-terminal protease domain. FAM111A is expressed in the nucleus and contains an N-terminal proliferating cell nuclear antigen (PCNA) interacting peptide (PIP) box and plays a role in PCNA loading during DNA replication[5]. In addition, FAM111A has been shown to prevent replication forks from stalling via its protease domain. FAM111A containing disease-associated mutations display aberrant nuclear morphology due to the disruption of nuclear pore complex (NPC) as a result of its hyperactive protease activity[6]. Recently, more studies have pinpointed the function of FAM111A in restricting virus infection. Fine et al. identified FAM111A as a host restriction factor for SV40 and showed

FAM111A was specifically targeted by the C-terminal region of Simian Virus 40 (SV40) large T (LT) antigen[7]. Furthermore, in a separate report, they demonstrated that FAM111A restricted the formation of replication center in the nucleus for SV40 host range mutant[6]. Moreover, Panda et al. discovered through a whole genome RNAi screening with VACV-ΔSPI-1 that FAM111A may be a cellular target of viral serine protease inhibitor SPI-1 (designated C12 in VACV-cop)[8]. Nevertheless, detailed mechanisms by which a nuclear protein FAM111A restricts viral infections, especially infections of poxviruses that replicate exclusively in the cytoplasm, remain largely elusive.

Poxviruses are large cytoplasmic DNA viruses, and as a family, can infect a wide variety of different animal species. The most notorious members of the poxvirus family include variola virus, the causative agent of smallpox, and monkeypox virus (MPXV), which is responsible for the recent global outbreaks of monkeypox infections since 2021[9-12]. Vaccinia virus was used as the vaccine strain for the eradication of smallpox and is also the prototypic poxvirus that is best characterized[13,14]. The genome of vaccinia virus encodes more than 180 genes, nearly half of which are involved in host-virus interactions and a third of them responsible for evading host antiviral defenses[15]. Vaccinia virus encodes three serine protease inhibitors (SPI-1, SPI-2, and SPI-3). Deletion of SPI-1 from vaccinia virus caused a host range defect and insertion of it into the modified vaccinia virus Ankara (MVA)[16,17], the highly attenuated vaccine strain unable to replicate in most mammalian cells, was able to rescue MVA's replicative capability in mammalian cells such as BS-C-1 and MRC-5[17]. As a previous study identified FAM111A as one of the major cellular targets of SPI-1[8], more efforts are needed to characterize the role of SPI-1 in

antagonizing FAM111A's antiviral effect on vaccinia virus.

Here, we demonstrate that FAM111A, a component of the replication fork primarily localized in the nucleus under normal conditions, translocated into the cytoplasm in VACV-infected cells upon the activation of the cGAS-STING signaling pathway by VACV infection. The export of FAM111A necessitated the disruption of NPC protein NUP62, which is dependent on its peptidase activity. Cytoplasmic FAM111A moved to the viral factories, bound and mediated the degradation of a viral DNA binding protein and uncoating factor I3 through autophagy, leading to the arrest of viral DNA replication, post-replicative protein expression, and eventually virus replication. We also show that poxvirus SPI-1 from MPXV and lumpy skin disease virus also blocked FAM111A-mediated I3 degradation. Upon infection, SPI-1 entered the nucleus of infected cells and prevented the nuclear export of FAM111A by inhibiting the peptidase activity of the latter, and a functional SPI-1 reactive-site loop (RSL) was indispensable for this inhibitory effect[18]. Our findings uncover the detailed molecular mechanism by which FAM111A suppresses vaccinia virus replication and demonstrate the mode of action of SPI-1 in antagonizing the antiviral function of FAM111A.

Results

FAM111A is a host restriction factor for VACV lacking SPI-1

Panda et al. previously reported FAM111A as a restriction factor for rabbit poxvirus and VACV lacking SPI-1. To verify the antiviral effect of FAM111A, we cloned human FAM111A into a mammalian expression vector and expressed it ectopically in human A549 cells before infecting the cells with MVA and MVA engineered to express SPI-1 or/and C16

(MVA+SPI-1, MVA+SPI-1/C16), VACV-WR or VACV-ΔSPI-1. We examined the expression of viral genes and viral replication by Western blotting analysis (0.3 PFU/cell) and virus quantification with plaque assay (0.03 PFU/cell), respectively. Transfected FAM111A was detectable with an anti-FAM111A antibody and overexpression of FAM111A dramatically decreased viral protein expression detected using an anti-VACV polyclonal antiserum (Fig. 1A). Notably, the decrease of viral protein expression was more phenomenal in VACV-ΔSPI-1 infected cells than in cells infected with VACV-WR or MVA+SPI-1/C16, suggesting SPI-1 may counteract with the function of FAM111A. In conformity with the results of Western blotting, over-expressing FAM111A significantly attenuated the replication of VACV in A549 cells and the effects were more pronounced for viruses that lack SPI-1 (VACV-ΔSPI-1 and MVA, Fig. 1B), The antiviral effect of FAM111A improved as we increased the amount of FAM111A transfected in MVA and VACV-ΔSPI-1-infected cells (Fig. 1C). To test the role of FAM111A on viral DNA replication[19], we examined the abundance of viral DNA using a set of primers for E11 in cells transfected with FAM111A and infected with VACV, MVA and their recombinants indicated in the figure at 3 PFU/cell in A549 cells (Fig. 1D), and the blockage of viral DNA replication was observed in the presence of FAM111A. In agreement with the above results, viral DNA loads were decreased significantly in both MVA and VACV-ΔSPI-1-infected cells in a FAM111A dose-dependent manner (Fig. 1E), further confirming the blockage of FAM111A on viral DNA replication.

Next, we aimed to determine if depletion of endogenous FAM111A could relieve the replication of VACV lacking SPI-1. We first transiently knocked down the expression of

FAM111A by transfecting A549 cells with siFAM111A or the negative control siNC. The expression of viral proteins determined by anti-VACV antiserum (Fig. 1F), virus titers (Fig. 1B), and viral DNA loads (Fig. 1D) were much higher in cells transfected with siFAM111A than in those transfected with only siNC as a negative control, and the comparison was more dramatic in MVA and VACV-ΔSPI-1-infected cells than in cells infected with SPI-1-expressing viruses. To further confirm our findings, A549$^{FAM111A-KO}$ cell lines were generated using CRISPR-Cas9 technology with sgRNA targeting the 5' region of FAM111A. Western blotting analysis verified the complete depletion of the protein in our FAM111A-KO clones (Fig. S1) and the deletion was confirmed by Sanger sequencing using genomic DNA collected from the FAM111A-KO clones. We then infected A549^{WT} and A549$^{FAM111A-KO}$ cells with the above-described viruses and examined the expression of viral proteins, virus titers, and viral DNA loads using the methods described. Depletion of FAM111A increased viral protein expression (Fig. 1G), virus titers (Fig. 1B), and viral DNA loads (Fig. 1D) dramatically. Replication of VACV was however not affected by FAM111A depletion. Notably, even though VACV-ΔSPI-1 replicated better in FAM111A-KO cells than in WT cells, the titer was still lower than what VACV could achieve in either A549^{WT} or A549$^{FAM111A-KO}$ cells, indicating additional restricting factors may exist for VACV-ΔSPI-1. To test if FAM111A may be a species-specific host restriction factor, we transfected monkey BS-C-1 cells and avian DF-1 cells with human FAM111A and found human FAM111A worked to restrict virus replication in neither of those tested cells, this results suggested the antiviral function of FAM111A may require other participating factors that could be species-specific and are absent in non-human cells (Fig. S2). To inspect if

viral infections lead to induction or repression of FAM111A, we determined FAM111A expression by Western blotting analysis post MVA or VACV infections and found no changes in FAM111A expression were observed overtime (Fig. S3). In summary, our results illustrated that FAM111A was a host restriction factor for VACV viruses that do not possess a functional SPI-1.

Inhibition of VACV-ΔSPI-1 and MVA by FAM111A requires its functional protease domain

The C-terminal domain of FAM111A contains a trypsin-like motif that constitutes a catalytic triad consisting of His385, Aps439, and Ser541[20]. This catalytic triad is highly conserved among different mammals (Fig. S4). Mutation of S541A jeopardized the enzymatic function of FAM111A and mutation of R569H is a hyperactive mutation found in patients with Kenny–Caffey syndrome (KCS) and osteocraniostenosis (OCS) which also exhibited an increased self-cleavage activity[20]. To test if FAM111A's antiviral effect is dependent on its protease activity, we generated vectors containing a loss-of-function mutant FAM111A[S541A,] a hyperactive mutant FAM111A[R569H] or a double mutant FAM111A[S541A+R569H] that was previously shown to behave like a loss-of-function mutant (illustrated in Fig. 2A). When transfected into human cells, FAM111A[S541A] and FAM111A[S541A+R569H] mutants caused accumulation of the protein comparing to FAM111A[WT](Fig. 2B), a result of the decreased self-cleavage activity due to loss of peptidase activity[5]. The hyperactive mutation FAM111A[R569H] expressed at a lower level compared to FAM111A[WT], possibly due to excessive self-cleavage rather than proteasome-mediated degradation as the addition MG132 (20 µM), a proteasome inhibitor,

showed no effect on FAM111A protein levels (Fig. 2C). These results confirmed that the

mutations generated indeed altered its protease activity. We next infected the cells

transfected with either FAM111A^{WT} or mutated FAM111As with MVA or VACV-ΔSPI-1 and

inspected viral protein expression, virus growth, and DNA replication in these cells. The

results showed that the reduction of VACV protein caused by overexpressing FAM111A

was eliminated when the enzymatic activity was disrupted through S541 mutation, even

though the mutants without a functional protease domain were expressed at much higher

levels (Fig. 2D). It is worth noting that the hyperactive mutation FAM111A^{R569H} did not

further decrease viral protein expression comparing to FAM111A^{WT}, which may be the

result of low protein accumulation due to stronger self-degradation of FAM111A^{R569H}. In

addition to the results of protein expression, expression of FAM111As with loss-of-function

mutations was no longer able to suppress virus replication (Fig. 2E) and viral DNA

replication (Fig. 2F) in A549 cells, while the gain-of-function mutation did not show a

significantly altered antiviral activity compared to the FAM111A^{WT}. As the PCNA

interaction motif is required for its physiological function for host DNA replication, we

generated a mutant that contained mutations in the PIP box (Fig. 2A) but found the loss of

PCNA binding did not compromise the antiviral function of FAM111A (Fig 2G). These

results exhibited that the enzymatic activity of the protease domain of FAM111A was

indispensable for its antiviral function.

VACV infection leads to the nuclear export of FAM111A through

disruption of the nuclear pore complex (NPC)

Unlike most DNA viruses, poxvirus replication is restricted to the cytoplasm of infected

cells. We thus deduce that FAM111A, a protein that existed in the nucleus under normal conditions, would have to be exported to the cytosol to exert its antiviral effect. We sought to determine if viral infection triggers any changes in the subcellular localization of FAM111A. A549 cells plated on coverslips were mock-infected or infected with MVA and the subcellular localization of FAM111A was determined by immunofluorescence using anti-FAM111A antibodies and Hoechst was used to stain nucleus and virus factories. In mock-infected cells, transfected FAM111A were exclusively found in the nucleus and the hyperactive mutation of FAM111A caused aberrant morphology of the nucleus, consistent with a previous report[6] (Fig. 3A). When infected with MVA, however, FAM111A^{WT} was found to be localized primarily in the cytosol (Fig. 3B). In contrast, the two mutants with inactivated protease domain failed to translocate and were accumulated in the nucleus. FAM111A^{R569H}, on the other hand, was also found mostly in the cytosol but with a more concentrated distribution close to the viral factories (Fig. 3B). The subcellular localization of FAM111A and its mutants upon viral infection was also quantitated and the results were shown in Fig. 3C. We further isolated nuclear and cytosolic proteins and examined the abundance of FAM111A in each part and were able to confirm the nuclear export of FAM111A upon MVA infection in a time-dependent manner (Fig. 3D-F). The above experiments were repeated in Hela cells and the results were consistent (Fig. S5).

Previous reports showed overexpression of FAM111A could destroy NPC by degrading nuclear pore complex proteins[6]. We reasoned the nuclear export of FAM111A may be achieved through the destruction of the nuclear pore complex by its protease activity. To test this hypothesis, A549 cells plated on coverslips were transfected with FAM111A or the

loss-of-function and hyperactive mutants, and Mab414 was used to stain NPC for confocal microscopic analysis. As expected, transfection of FAM111A^{WT} and FAM111A^{R569H} disrupted normal NPC structure and FAM111A^{S541A} and FAM111A$^{S541A+R569H}$ showed no such effect (Fig. S6A). In addition, NUP62, a core component of the NPC was found to be degraded by ectopically expressed FAM111A (Fig. S6B). To test if endogenous FAM111A could also degrade NPC upon viral infection, A549 cells were mock-infected or infected with MVA, MVA+SPI-1 at 3 PFU/cell for 6 hours and Mab414 antibody was used to examine NPC integrity by confocal microscopy. MVA infection led to disruption of the NPC in areas where FAM111A was concentrated (highlighted in the white box) and the nuclear membrane was undisturbed in mock-infected cells or cells infected with MVA+SPI-1 (Fig. 4A). The results from the confocal microscopic analysis were also quantitated as shown in Fig. 4B. These data revealed VACV infection triggered FAM111A-mediated NPC disruption and the subsequent nuclear export of FAM111A in the absence of SPI-1.

The cGAS-STING signaling pathway mediated the nuclear export of FAM111A upon VACV infection

As the lifecycle of poxvirus is complete in the cytosol of infected cells, we wondered how could FAM111A sense the invasion of VACV and initiate its migration to the cytosol. Cyclic GMP-AMP synthase (cGAS) is a crucial pattern recognition receptor activated by cytosolic DNA[21,22]. To determine if cGAS affect the nuclear export of FAM111A, we designed siRNA specific for cGAS and confirmed that sicGAS could lessen the level of cGAS efficiently (Fig. 5A). A549 cells transfected with siNC or sicGAS for 24 hours were

left untreated or transfected with an empty vector, extracted genomic DNA from MVA, poly (dA:dT), a repetitive synthetic dsDNA, or poly (I:C), a molecule used to simulate dsRNA. Eight hours later, total protein was collected and components of nuclear and cytosolic proteins were isolated using the method described in figure 3. As shown in Fig 5A, sicGAS was able to repress the expression of cGAS and phosphorylation of STING but had no effect on the expression of FAM111A, total STING, PCNA, or histone H3. Transfection of vector DNA, viral genomic DNA, or poly (dA:dT) all led to increased nuclear export of FAM111A while the transfection of poly (I:C) failed to do so (Fig. 5C). Transfection of poly (dA:dT) showed strong induction of FAM111A expression and the reason was unknown. Importantly, the cytosolic accumulation of FAM111A caused by transfection of plasmid vector or viral genomic DNA was largely reduced in cells transfected with sicGAS, indicating the nuclear export of FAM111A was induced by the presence of dsDNA in the cytosol through the cGAS-STING signaling pathway. We concluded from these results that the cytosolic accumulation of virus DNA triggered the export of FAM111A from the nucleus to the cytosol in VACV-infected cells.

FAM111A targets virus DNA binding protein I3 for degradation

The above-mentioned experiments demonstrated that VACV infection induced the export of FAM111A to the cytosol through the cGAS signaling pathway and the export of FAM111A was achieved through NPC destruction via its protease activity. It was hence important to investigate how FAM111A exerted its antiviral activity. We first hypothesized that FAM111A could directly bind to and catalyze the degradation of certain viral proteins via its active protease domain. We designed an experiment to pull down the viral

interactome of FAM111A in virus-infected cells and characterize its components with mass spectrometric analysis (Fig. 6A). In this assay, VACV I3 showed the most significant difference in protein abundance between FAM111A-transfected and vector-transfected cells and was further investigated as a putative target of FAM111A (Fig. 6B). The interaction between FAM111A and I3 was first confirmed by co-immunoprecipitation (co-IP) analysis by using either FAM111A or I3 as the bait. To determine if the interaction was DNA-dependent as FAM111A contains a double-stranded DNA binding domain (DBD), cell lysates were incubated with or without 250U/ml of benzonase, an endonuclease known to degrade all forms of DNA without base preference. As expected, total lysates and proteins captured with anti-FAM111A antibody revealed an interaction between FAM111A and I3 (Fig. 6C) and this interaction could be further verified by capturing I3-interacting proteins with anti-I3 antibodies (Fig. 6D). However, the association of I3 and FAM111A disappeared when benzonase was added. As FAM111A contains a DBD and no discernible RNA binding motifs, we concluded that the interaction between FAM111A and I3 was DNA-dependent. Additional experiments were carried out to validate the co-localization of I3 and FAM111A within cells (Fig. 6E). A549 cells on coverslips were co-transfected with vectors expressing viral I3 and FAM111A and both were stained for confocal microscopy. As expected, virus I3 was associated with Hoechst-stained virus factories in the cytoplasm of infected cells and the majority of FAM111A colocalized with I3 (Fig. 6E). Therefore, the association of FAM111A and I3 was shown by immunoaffinity purification and further suggested by their intracellular colocalization in the cytoplasm.

Next, experiments were performed to investigate if FAM111A resulted in virus I3

degradation. A549 cells transfected with increasing amounts of FAM111A were mock-infected or infected with MVA or MVA+SPI-1 at 3 PFU/cell and virus protein expression was examined by Western blotting analysis. Expression of viral proteins was reduced as the amounts of FAM111A transfected increased in MVA and MVA+SPI-1 infected cells (Fig. 6F). Unexpectedly, FAM111A was also able to lessen viral protein expression of MVA+SPI-1. One possible explanation is that the amount of SPI-1 present was not sufficient to fully restrict FAM111A when it was transfected at a high concentration as SPI-1 was expressed from the virus using a virus promoter. Nevertheless, it was apparent that I3 in MVA+SPI-1-infected cells was more resistant to FAM111A than that in MVA-infected cells (Fig. 6F). Importantly, virus I3 decreased as the amount of FAM111A increased while virus E3, another early protein that binds dsRNA, was insensitive to the presence of FAM111A (Fig. 6F), furture confirming the antiviral effect of FAM111A occurred post-replicatively. To further verify the specificity of I3 degradation by FAM111A, we co-transfected A549 cells with FAM111A and flag-tagged I3 or H5, another DNA binding protein, and found like E3, H5 was unaffected by FAM111A (Fig. 6G). These results suggested FAM111A was specifically associated with virus uncoating factor I3 and mediated its degradation upon virus infection.

FAM111A promotes I3 degradation through autophagy but not its peptidase activity

To investigate if FAM111A directly degrades virus I3 through its C-terminal peptidase activity, we simultaneously co-transfected A549 cells with VACV I3 and FAM111A or its hyper- and hypo-active mutants described above. Surprisingly, I3 was degraded when not

only FAM111A^{WT} but also when FAM111A mutants with inactivated (FAM111A^{S541A} and FAM111A$^{S541A+R569H}$) protease activity were transfected (Fig. 7A). As we previously discovered that the protease activity of FAM111A was crucial for its nuclear export, these combined results suggested that the FAM111A mutants with inactivated peptidase domain may bypass nuclear export and associate with I3 directly when they were co-transfected. The co-transfection of FAM111A and I3 may lead to a pre-mature association of FAM111A and I3 in the cytoplasm and result in I3 degradation. To determine whether the subcellular localization of FAM111A could affect its antiviral function, we decided to alter our strategy by transfecting cells with FAM111A or its derivative mutants for 24 hours before transfecting I3. By doing this, FAM111A expressed could be localized to the correct subcellular position before interacting with I3 (Fig. 7B). In addition, we generated more FAM111A mutants by either introducing a nucleus export signal (NES) or disrupting its nucleus localization signal (NLS-mut) (Fig. 7C). Next, A549 cells were transfected with FAM111A^{WT}, FAM111A$^{DBD\text{-}mutant}$, FAM111A^{S541A}, FAM111A$^{S541A+NES}$ or FAM111A$^{NLS\text{-}mut}$ for 24 hours, and then transfected with a vector expressing virus I3 and the levels of FAM111A and I3 were examined by Western blotting analysis. As expected, mutants with disrupted double-stranded DNA binding domain or inactivated serine protease domain abolished their ability to degrade I3 (Fig. 7D) and were found primarily in the nucleus (Fig. S7). Remarkably and interestingly, the mutant with an inactivated protease domain combined with an NES motif was found mostly in the cytosol and colocalized with I3 (Fig. S7), and it was still able to cause I3 degradation (Fig. 7D). Similarly, the mutant with a mutated NLS was found primarily in the cytoplasm (Fig. S7) and was also able to mediate

I3 degradation. These results demonstrated that the serine protease domain of FAM111A was only required for nuclear export but not its antiviral function and FAM111A designed to be expressed in the cytoplasm could associate with and mediate the degradation of I3 even when its peptidase domain was inactivated.

Next, we decided to explore the nature of I3 degradation by using chemical inhibitors targeting various cellular pathways known to modulate protein degradation. A549 cells were sequentially transfected with FAM111A and I3 in the presence or absence of small molecule chemicals inhibiting autophagy [10 µM chloroquine (CQ), 5 mM 3-methyladenine (3-MA), 0.5 µM bafilomycin A1 (BAF)], proteasomal degradation (20 µM MG132) or pan-caspase activity (20 µM Z-VAD-FMK). DMSO was used to dissolve the above drugs and was added as a negative control. While the addition of CQ and MG132 both alleviated FAM111A-induced degradation of I3, the addition of 3-MA, BAF or Z-VAD-FMK showed no discernible effects (Fig. 7E). Proteasome-mediated I3 degradation was ruled out as I3 co-purified with FAM111A exhibited no sign of ubiquitination (Fig. S8). The result of MG132 was likely due to an unknown off-target effect. We next performed a dose-dependent experiment and further confirmed that the addition of CQ (10/50/100 µM) caused the accumulation of LC3B-II and led to the rescue of FAM111A-mediated I3 degradation (Fig. 7F). Moreover, CQ slightly increased the expression I3 when cells were transfected with only I3 but not FAM111A (Fig. 7G), this was possibly due to I3 degradation stimulated by endogenous FAM111A, as this phenomenon was not observed in A549$^{FAM111A-KO}$ cells (Fig. 7G). To rule out the possibility that the rescue of I3 degradation was due to the off-target effect of CQ, we manipulated

the level of ATG16L1, a key regulator of autophagic flow, by generating an A549 cells line in which the endogenous level of ATG16L1 was stably knocked down by shRNA and named it A549$^{ATG16L1-KD}$ (Fig. 7H). Similar to the effect observed with CQ, depression of ATG16L1 was able to rescue FAM111A-mediated I3 degradation, suggesting the essential role of autophagy in such a process. It is worth noting that in the above experiments, the accumulation of LC3B-II decreased slightly when FAM111A was transfected, indicating that FAM111A may only utilize the autophagy pathway for I3 degradation but did not initiate the autophagic flow. The above results showed that FAM111A interacted with virus I3 and promoted I3 degradation by utilizing the autophagy pathway, but not its peptidase activity.

Poxvirus SPI-1 blocks the nuclear export of FAM111A through its serpin activity

SPI-1 (C12 in VACV-cop) has previously been shown to function as a host range factor for MVA[17]. To explore its role in FAM111A inhibition, we first observed its subcellular localization by infecting FAM111A-transfected A549 cells with MVA or MVA+SPI-1 at 3 PFU/cell and probed SPI-1 using a myc antibody at 12 hpi as the SPI-1 inserted into MVA contains a myc tag at its N-terminus[17]. The confocal microscopic analysis showed that SPI-1 was mainly localized in the nucleus during MVA+SPI-1 infection (Fig. S9). In addition, nuclear export of FAM111A activated by MVA was no longer observed when cells were infected with MVA+SPI-1 (Fig. 8A), suggesting SPI-1 prevented FAM111A from exporting to the cytoplasm. To investigate if SPI-1 was associated with FAM111A, we performed immunoaffinity purification using either SPI-1 or FAM111A as the bait and

observed a strong interaction of FAM111A and SPI-1 (Fig. 8B-C). As FAM111A restricted

viral replication by degrading virus I3, we next inspected if SPI-1 could interfere with

FAM111A-mediated I3 degradation. A549 cells were co-transfected vectors encoding

FAM111A, VACV I3, and VACV-SPI-1. Consistent with previous results, transfection of

FAM111A led to a prominent I3 degradation when SPI-1 was absent, however, this effect

was alleviated when plasmid expressing VACV-SPI-1 was co-transfected (Fig. 8D).

Interestingly, SPI-1 from MPXV, a close relative of VACV, and lumpy skin disease virus, a

member of the capripoxviruses, displaying 96.4% and 37.8% sequence identity with

VACV SPI-1, respectively, could also revert FAM111A-mediated I3 degradation. A

previous report identified cathepsin G as one of the cellular targets of rabbitpox virus

SPI-1 and showed mutations in the reactive-site loop (RSL) of SPI-1 abolished its

interaction with cathepsin G[18]. We thus introduced the two mutations T305R, F318A

separately or in combination into VACV SPI-1 and examined their effects on FAM111A

inhibition (Fig. 8E). We used the same transfection strategy described in Fig. 7B by

transfecting FAM111A 24 hours before introducing I3 and SPI-1 to allow FAM111A to be

localized to the nucleus. As shown in Fig. 8E, we again observed that wt-SPI-1 was able

to rescue FAM111A-mediated I3 degradation, however, the SPI-1 mutants with mutations

in the RSL region failed to inhibit FAM111A-promoted I3 degradation even though

expressed at comparable levels with the wt-SPI-1. These results suggested that SPI-1

may interact with FAM111A via the RSL region and demonstrated that SPI-1 antagonized

FAM111A's antiviral function by inhibiting the serine protease activity of FAM111A. We

summarized our findings in a diagram presented in Fig. 9.

Discussion

FAM111A has been postulated to be a serine protease regulated in a cell-cycle-dependent fashion and conserved in mammals[5]. It displays autocleavage and was identified as a component of replication fork by nascent chromatin proteomics. Nuclear FAM111A protein co-localized with PCNA at replication sites and functions to load PCNA via its PIP motif[23]. Recent studies suggested FAM111A could restrict the replication of SV40 mutant by interfering with the formation of replication centers in the nucleus[7]. A human genome-wide RNAi screening using VACV-ΔSPI-1 identified FAM111A as a restrictive factor that influences VACV that lacks SPI-1 replication in human cells. However, the detailed mechanism by which FAM111A restricts poxvirus, a virus that normally replicates in the cytoplasm, was poorly understood. To investigate the role of FAM111A in prohibiting VACV infection, we first examined the effect of overexpressing or depressing the expression of FAM111A on virus replication using VACV that lacks a functional SPI-1 (VACV-ΔSPI-1) or MVA, which lost its SPI-1 during repeated passages in avian cells. We were able to verify the results reported previously that deletion of FAM111A was able to enhance VACV replication in human A549 cells. In addition, we discovered that its antivral effect occurred post-replicatively as a viral early gene E3 was expressed in the presence of FAM111A. Moreover, transient transfection of FAM111A in human cells exhibited a strong inhibitory effect on viral DNA replication, viral protein expression, and virus replication，further confirming the antiviral function of FAM111A for SPI-1-lacking VACV. The trypsin-like domain of FAM111A contains a catalytic triad including His385, Aps439, and Ser541[24,25] and a DNA-binding domain that plays essential

roles for its localization at the replication forks[5,23]. FAM111A was also shown to bind to the large T antigen of SV40 and inhibit the formation of the replication center of SV40 mutants, demonstrating its broad antiviral activity for DNA viruses[7]. However, the replication of VACV differs from that of SV40 as VACV completes its entire replication cycle in the cytoplasm of infected cells. As a result, nuclear export is the first step for FAM111A to confer its antiviral function during a VACV infection. Mutations of the catalytic triad of FAM111A abolished its antiviral activity for VACV-ΔSPI-1 and the reason was that the nuclear export of FAM111A requires a functional trypsin-like domain. This was further validated by the observation that FAM111A mutant with inactivated peptidase domain was unable to degrade NUP62 and cause NPC disruption (Fig. S6). In addition, transfection of FAM111A with a hyperactive peptidase domain disrupted normal nuclear morphology, which was previously reported and further demonstrated that the exportation of FAM111A is dependent on its catalytic peptidase domain[6]. Following MVA infection, the subcellular localization of FAM111A changed, showing transportation from the nucleus to the cytoplasm and it co-localized with the viral factories stained by Hoechst or I3 (Fig. 3 and 6). In addition, this change in FAM111A localization was dependent on its DNA-binding domain but not its PCNA-interaction motif. The catalytic triad at the C-terminus of FAM111A is highly conserved among different species, suggesting that its protease activity is critical for its cellular function.

In uninfected cells, FAM111A was exclusively found in the nucleus, therefore, we were interested in exploring how infection of a DNA virus that replicates in the cytosol could trigger the nuclear exportation of FAM111A. Panda et al. postulated that replication

factor C (RFC) 1-5 could load PCNA and act as a DNA sensor recruiting FAM111A to endogenous replication sites or viral genomes that utilize/attract PCNA[8]. However, we found that FAM111A mutant with a dysfunctional PCNA-interectin motif (FAM111A[PCNA-mut]) could still restrict VACV replication, suggesting the recruitment of FAM111A to the viral factor is not PCNA mediated. Rather, mutations in the DBD abolished the nuclear export of FAM111A and resulted in its nuclear accumulation, suggesting the involvement of DNA binding in FAM111A recruitment (Fig.S7). Interestingly, we found that the expression level of FAM111A increased significantly after transfecting the cells with DNA isolated from MVA or poly(dA:dT) but not poly (I:C). In addition, the presence of dsDNA alone, but not dsRNA triggered the nuclear export of FAM111A without virus infection. Both the increase of FAM111A protein abundance and nuclear export were diminished in cells where cGAS was repressed (Fig. 5). These results all suggested the existence of a regulatory axis that associated cGAS with FAM111A that needs to be further explored. More needs to be done to clarify components in the cGAS-FAM111A signaling pathway and to explore the dynamics of FAM111A exportation. Previous studies showed that infection of an SV40 host range mutant enhanced the protease activity of FAM111A, leading to impairment of nuclear barrier function, cell morphology, and normal distribution of the nuclear pore [6]. These results were in line with our observation but the upstream activating signal for FAM111A may be different as SV40 replicates in the nucleus.

Using an immunoprecipitation assay coupled with mass-spectrometry, we were able to identify the viral binding partner of FAM111A during MVA infection. We discovered that cytosolic FAM111A interacted with viral I3 in a DNA-dependent manner. VACV I3 is a 34

kDa phosphoprotein essential for poxvirus genome replication[26,27]. Like FAM111A, I3 can also bind DNA with its single-stranded DNA binding domain (SSB)[28]. The interaction between FAM111A and I3 was terminated with benzonase treatment, and the FAM111A mutant that lacks a functional DNA binding domain no longer displayed any antiviral activity (Fig. 6 and 7). These observations highlight the importance of DNA binding in its interaction with I3 and the subsequent antiviral function of FAM111A. Remarkably, our results showed both the FAM111A^{WT} and the mutants that lack a functional peptidase domain could lead to efficient virus I3 degradation, and this degradation was protein specific as virus H5, another VACV early protein, was not affected by FAM111A (Fig. 6). We were surprised by these results as we initially predicted that the catalytic activity of FAM111A was responsible for I3 degradation. By manipulation of the subcellular localization of FAM111A using NES or interruption of NLS signals within FAM111A, we found that the catalytic activity of FAM111A was only required for nuclear exportation but not I3 degradation as the FAM111A^{S541A} containing an NES (FAM111A$^{S541A+NES}$) was still able to induce I3 degradation (Fig. 7). To examine the nature of I3 degradation, we used pharmaceutical inhibitors that target various cellular protein-degradation pathways. We found the degradation of I3 by FAM111A did not rely on its protease activity, nor did it use the apoptotic pathway. Instead, the I3 degradation was alleviated when autophagy was inhibited by chloroquine. Furthermore, we also observed that the depression of ATG16L1 by shRNA was able to relax the degradation of I3 by FAM111A. In our assay, 3-MA and bafilomycin A1, two different chemical inhibitors of autophagy failed to negate FAM111A-mediated I3 degradation. The exact reason was unknown but we speculated

that it may be because bafilomycin A1, 3-MA, and chloroquine inhibit different steps of the autophagic flow and VACV may possess strategies to compensate for the blockade of autophagosome nucleation and lysosomal acidification, which were inhibited by 3-MA and bafilomycin A1, respectively[29].

VACV SPI-1 encoded by the C12 gene in RPXV was previously thought to affect the virus host range by inhibiting the serine protease activity of cathepsin G[18]. SPI-1 of VACV was mainly distributed in the nucleus after virus infection and interacted with FAM111A in a co-IP assay. The presence of SPI-1 was able to prevent FAM111A's nuclear export by inhibiting the protease activity of FAM111A, thereby antagonizing its antiviral function. FAM111A-mediated I3 degradation was also inhibited by SPI-1. In RPXV, SPI-1 was predicted to be associated with viral proteins essential for DNA replication[30]. Our results were in agreement with this hypothesis as we observed interactions and co-localization of SPI-1 and FAM111A within the nucleus of infected cells, which mainly inhibited viral DNA replication by degrading virus uncoating factor I3. Furthermore, we found that SPI-1 proteins from different poxviruses [VACV, monkeypox virus (MPXV), and lumpy skin disease virus (LSDV)] could all inhibit FAM111A-mediated I3 degradation, indicating a conserved role of SPI-1 in antagonizing human FAM111A. Importantly, mutations of the two residues in the RSL of SPI-1 (T305R, F318A) abrogated its function in mitigating FAM111A-promoted I3 degradation without significantly affecting SPI-1 expression. As these two residues were previously reported to be essential for cathepsin G binding, we speculated that SPI-1 interacted with FAM111A via a similar region in its RSL[18] (Fig. 8).

As far as we know, our work is the first report of a detailed mechanism by which FAM111A prohibited cytoplasmic DNA virus replication. Our results indicated that nuclear components normally involved in mammalian cell DNA replication could translocate from the nucleus to the cytoplasm and restrict poxvirus replication by degrading key viral uncoating factors using cellular autophagic machinery. In addition, we presented an explanation for the immune evasion function of VACV SPI-1.

Materials and methods

Cell Culture and Generation of Stable Cell Lines. Human A549 cells (ATCC CCL-185) were purchased from ATCC and maintained in Dulbecco's Modified Eagle Medium (DMEM)/F-12 (Gibco) supplemented with 10% fetal bovine serum (FBS, Gibco) and 1% penicillin-streptomycin solution (Solarbio). HeLa cells (ATCC CCL-2), 293T cells (ATCC CRL-3216), and DF-1 cells (ATCC CRL-12203) were obtained from ATCC and propagated in DMEM supplemented as above. BS-C-1 cells (ATCC CCL-26) were purchased from ATCC and grown in minimum essential medium with Eagle's balanced salts (EMEM) supplement as above.

The FAM111A-KO cell line was constructed using the CRISPR-Cas9 technology described previously[31]. Briefly, two different sgRNAs were designed to target human FAM111A (5'-AAAAATCCAGAAGACCAGACCA-3', 5'-TGGGAAAATGTAATGACCACCT-3') and were inserted into pSpCas9(BB)-2A-GFP, a gift from Feng Zhang (Addgene plasmid #48138). A549 cells were transfected with the above plasmids at 60% confluency and sorted by flow cytometry based on the expression of GFP at 48 h post-transfection. GFP-positive cells were serially diluted for clonal

selection and successful inactivation of FAM111A was confirmed by Sanger sequencing and Western blotting analysis with a FAM111A antibody (Abcam).

The A549-ATG16L1-KD cell line stably expressing ATG16L1-shRNA was constructed with retroviral transduction. ATG16L1 shRNA (5'-GTCATCGACCTCCGGACAAAT-3') was inserted into pLKO.1 puro (Addgene plasmid #8453) to generate pLKO.1-ATG16L1-shRNA. Retroviruses were packaged in 293T cells by co-transfecting the cells with 1ug pLKO.1-ATG16L1-shRNA, 1µg pMD2.G (Addgene plasmid #12259) and 1µg psPAX2 (Addgene plasmid #12260) in each well of a 6-well plate. Virus particles were collected 48 h post-transfection and filtered (0.22 µm) to remove cell debris. A549 cells were transduced with retroviruses at various dilutions in the presence of 5 µg/mL of polybrene for 48 h. Cells were passaged three times in a complete medium supplemented with 1 µg/mL of puromycin (Sigma-Aldrich). The A549-ATG16L1-KD and the A549-shControl cells were both maintained in the presence of 1 µg/mL puromycin and 500 µg/mL geneticin. The expression of ATG16L1 was determined by Western blotting using an anti-ATG16L1 antibody (Cell Signaling Technology).

Antibodies. Rabbit antibody to VACV-WR strain and mouse antibody to VACV-I3 or VACV-E3 were described previously[32,33]. Mouse anti-β-actin antibody (#AA128), mouse anti-histone H3 antibody (#AF0009), mouse anti-Flag antibody (#AF519), mouse anti-HA antibody (#AH158), HRP-labeled goat anti-mouse IgG (#A0216) and HRP-labeled goat anti-rabbit IgG (#A0208) were purchased from Beyotime Biotechnology; rabbit anti-FAM111A antibody (#ab184572), mouse anti-PCNA antibody (#ab29) and mouse anti-nuclear pore complex proteins antibody (#ab24609) were purchased from Abcam;

rabbit anti-cGAS antibody (#79978), rabbit anti-STING antibody (#13647), rabbit anti-p-STING antibody (#19781), rabbit anti-LC3B antibody (#3868) and rabbit anti-ATG16L1 antibody (#8089) were purchased from Cell Signaling Technology; Goat anti-mouse IgG Alexa Fluor 488 (#A32723) and Goat anti-rabbit IgG Alexa Fluor 555 (#A32732) were purchased from Thermo Fisher Scientific.

Generation of recombinant vaccinia viruses. Recombinant viruses were constructed by homologous recombination using fluorescent reporter genes (mCherry or eGFP) for plaque selection. MVA+SPI-1 was previously described[31]. Briefly, a DNA cassette containing the C12L ORF regulated by the mH5 promoter followed by the GFP ORF regulated by the P11 promoter was generated and flanked by 5' and 3' sequences around the deletion III locus of MVA. CEF cells were infected with MVA at 3 PFU/cell and then transfected with the DNA cassette described above. Viruses were harvested 24 hpi and recombinant viruses were selected by selecting mCherry or eGFP positive clones and plaque-purified 3× times in CEF cells. Viral DNA was isolated from the clones with the DNeasy Blood & Tissue kit from Qiagen and the insertion of genes was verified by PCR amplification followed by Sanger sequencing. To generate MVA+SPI-1/C16, the C16L DNA segment along with the upstream 339 bp presumed to contain the natural promoter was copied by PCR from the genome of MVA-51.2 virus[17] and inserted into a cassette that contained the mCherry ORF regulated by the P11 promoter with flanking sequences comprised of ORFs 069 and 070. Homologous recombination was performed as above. All recombinant MVAs were propagated in CEF. VACV-ΔSPI-1 recombinant viruses were generated as follows. First, 500 bp up- and downstream of C12 ORF were cloned to the 5'

and 3' end, respectively, of eGFP ORF regulated by the VACV p11 promoter. The resultant 1,720-bp PCR product was then transfected into BS-C-1 cells infected with VACV-WR at 3 PFU/cell for 1 hour. A recombinant virus lacking SPI-1 was isolated by selecting GFP-positive virus clones and clonally purified by repeated plaque isolation. The loss of the C12 gene was confirmed by PCR.

Western Blotting analysis. Human A549 cells were washed once with ice-cold PBS and lysed in wells with 1× cell lysis buffer and 1× PMSF (Beyotime Biotechnology) for western blotting analysis on ice. Cell lysates were sonicated for 1 min to reduce viscosity. Proteins were resolved on 4–20% FastPAGE precast gels (Tsingke Biotechnology) and transferred to nitrocellulose membranes in a Trans-Blot Turbo machine (Bio-Rad). Membranes were blocked with 5% nonfat milk dissolved in Tris-buffered saline (TBS) for 1 h at RT and then incubated with primary antibodies diluted 1× TBS with 0.1% Tween 20 (TBST) containing 5% nonfat milk overnight at 4 °C. The membranes were washed three times with 1× TBST and incubated with a secondary antibody conjugated with horseradish peroxidase diluted 1:5,000 in 1× TBST containing 5% nonfat milk for 1 hour at RT. ECL signals were detected with SuperSignal West Dura substrates (Thermo Fisher Scientific) and visualized by Image J.

Immunoprecipitation. Human A549 cells were co-transfected with vectors expressing myc-tagged FAM111A and flag-tagged I3 or SPI-1 and total protein was harvested 24 hours post-transfection. Cells were first washed once with ice-cold PBS and lysed in wells with 150 µL 1× cell lysis buffer and 1× PMSF (Beyotime Biotechnology) in each well of a 6-well plate on ice. Protein lysates were centrifuged for 15 min at 15,000 × g at 4 □C and

then incubated with control magnetic beads, flag-trap magnetic beads, or myc-trap magnetic beads at 4 □C for 3 hours with rotation. The beads were then washed 6× times with 1× lysis buffer and the bound proteins were eluted by boiling with 30 µL LDS sample buffer containing 0.05 M DTT for 15 min. The proteins were then resolved on 4–20% FastPAGE precast gels (Tsingke Biotechnology) and analyzed by Western blotting with antibodies described previously.

Determination of virus yield. Cell monolayers in 24-well plates were infected in triplicates with viruses at a multiplicity of 0.03 PFU/cell in DMEM/F12 containing 2.5% FBS and 1% pen/strep for 2 hours. Cells were then washed twice with 1× phosphate-buffered saline (PBS) and incubated for 48 h. Viruses were harvested by scraping and followed by three freeze–thaw cycles. Duplicate DF-1 (for MVAs) or BS-C-1 (for VACV-WR and VACV-ΔSPI-1) monolayers were infected with serially diluted viruses for 2 h, and then the inoculum was aspirated and replaced with medium containing 2.5% FBS supplemented with 0.5% methylcellulose followed by 48h incubation. For quantification of MVA, DF-1 cells were fixed with methanol–acetone (1:1, vol/vol) and washed with distilled water, and incubated with anti-VACV-WR antibody diluted 1:1,000 in 3% FBS-PBS for 1 h. Cells were then washed twice with distilled water and incubated with a 1:3,000 dilution of protein A-HRP (Thermo Fisher Scientific). After 1 hour, protein A-HRP was aspirated and replaced with ethanol-saturated dianisidine, diluted 1:50 in PBS containing 0.3% H_2O_2. The dianisidine solution was removed after 10 min of incubation at room temperature, and cells were washed with distilled water. Plaques were counted manually. For quantification of VACV-WR and VACV-ΔSPI-1, cells were fixed with 1%

crystal violet diluted in 20% EtOH, plaques were counted and average titers of triplicates were shown.

Quantification of viral genomic DNA by quantitative PCR. Monolayers of A549 cells in 24-well plates were infected with indicated viruses at 3 PFU/cell for 1 hour in triplicates. Cells were washed twice with 1× PBS and total DNA was extracted at 6 hpi with DNeasy Blood/Tissue DNA mini kit (Qiagen) and quantitated by a nanodrop spectrophotometer (Thermo Fisher Scientific). Equal amounts (1µg) of DNA from each sample were then serially diluted and subjected to quantitative PCR with gene-specific primers for virus E11 and host GAPDH as the housekeeping control. The results of virus genomic DNA were normalized to that of GAPDH and relative DNA abundance was calculated.

Confocal microscopy. Human A549 cells and HeLa cells were cultured on glass coverslips and infected with viruses indicated in figures at 3 PFU/cell. Cells at various hpi were fixed with 4% paraformaldehyde (PFA) for 15 min, permeabilized with 0.5% Triton X-100 for 15 min, and blocked with 3% bovine serum albumin (BSA) in 1× PBS for 30 min at room temperature. Primary antibodies were diluted in 3% BSA-PBS and incubated with cells at 4 □C overnight. Samples were then washed with 1× PBS, and incubated with secondary antibodies conjugated with Alexa Fluor 488 or Alexa Fluor 555. Nuclei and virus factories were stained with Hoechst and cells were washed multiple times before mounting on slides using ProLong Diamond Antifade reagent (Thermo Fisher Scientific). Images were taken on a Leica SP8 confocal microscope and images were processed with the Leica software (Leica Biosystems).

Availability Statement. Materials and additional details regarding methods and protocols

may be obtained by contacting the corresponding author via email.

Acknowledgments

This work was funded by the National Key Research and Development Program (2021YFD1800700), the National Natural Science Foundation of China (32172822), and the Beijing STRS program (Z211100002121021) to Chen Peng.

Contributions

Conceived and designed the study: Bernard Moss, Chen Peng, Junda Zhu

Investigation: Junda Zhu, Xintao Gao, Zihui Zhang, Yining Deng, Shijie Xie, Shuning Ren, Yarui Li, Hua Li, Kang Niu, Shufang Fu, Yinü Li.

Methodology: Junda Zhu.

Resources: Chen Peng.

Supervision: Wenxue Wu, Chen Peng.

Writing – original draft: Chen Peng.

Writing – review & editing: Junda Zhu, Chen Peng, Bernard Moss

Conflict interests

The authors declare no conflict of interest.

Reference

Schelle, L., Corte-Real, J. V., Esteves, P. J., Abrantes, J. & Baldauf, H. M. Functional cross-species conservation of guanylate-binding proteins in innate immunity. *Med Microbiol Immunol*, 1-12, doi:10.1007/s00430-022-00736-7 (2022).

Evans, J. P. & Liu, S. L. Multifaceted Roles of TIM-Family Proteins in Virus-Host Interactions. *Trends Microbiol* **28**, 224-235, doi:10.1016/j.tim.2019.10.004 (2020).

Wang, B., Timilsena, Y. P., Blanch, E. & Adhikari, B. Lactoferrin: Structure, function, denaturation and digestion. *Crit Rev Food Sci Nutr* **59**, 580-596, doi:10.1080/10408398.2017.1381583 (2019).

Seo, J. Y., Yaneva, R. & Cresswell, P. Viperin: a multifunctional, interferon-inducible protein that regulates virus replication. *Cell Host Microbe* **10**, 534-539, doi:10.1016/j.chom.2011.11.004 (2011).

Kojima, Y. *et al.* FAM111A protects replication forks from protein obstacles via its trypsin-like domain. *Nat Commun* **11**, 1318, doi:10.1038/s41467-020-15170-7 (2020).

Nie, M. *et al.* FAM111A induces nuclear dysfunction in disease and viral restriction. *EMBO Rep* **22**, e50803, doi:10.15252/embr.202050803 (2021).

Tarnita, R. M., Wilkie, A. R. & DeCaprio, J. A. Contribution of DNA Replication to the FAM111A-Mediated Simian Virus 40 Host Range Phenotype. *J Virol* **93**, doi:10.1128/JVI.01330-18 (2019).

Panda, D., Fernandez, D. J., Lal, M., Buehler, E. & Moss, B. Triad of human cellular proteins, IRF2, FAM111A, and RFC3, restrict replication of orthopoxvirus SPI-1 host-range mutants. *Proc Natl Acad Sci U S A* **114**, 3720-3725, doi:10.1073/pnas.1700678114 (2017).

Vauhkonen, H. *et al.* Intrahost Monkeypox Virus Genome Variation in Patient with Early Infection, Finland, 2022. *Emerg Infect Dis* **29**, doi:10.3201/eid2903.221388 (2023).

Burdon, R. M. *et al.* Sustained Mpox Proctitis with Primary Syphilis and HIV Seroconversion, Australia. *Emerg Infect Dis* **29**, doi:10.3201/eid2903.221845 (2023).

Dung, N. T. *et al.* Monkeypox Virus Infection in 2 Female Travelers Returning to Vietnam from Dubai, United Arab Emirates, 2022. *Emerg Infect Dis* **29**, doi:10.3201/eid2904.221835 (2023).

McCollum, A. M. *et al.* Epidemiology of Human Mpox - Worldwide, 2018-2021. *MMWR Morb Mortal Wkly Rep* **72**, 68-72, doi:10.15585/mmwr.mm7203a4 (2023).

Moss, B. Genetically engineered poxviruses for recombinant gene expression, vaccination, and safety. *Proc Natl Acad Sci U S A* **93**, 11341-11348, doi:10.1073/pnas.93.21.11341 (1996).

Paoletti, E. Applications of pox virus vectors to vaccination: an update. *Proc Natl Acad Sci U S A* **93**, 11349-11353, doi:10.1073/pnas.93.21.11349 (1996).

Haller, S. L., Peng, C., McFadden, G. & Rothenburg, S. Poxviruses and the evolution of host range and virulence. *Infect Genet Evol* **21**, 15-40, doi:10.1016/j.meegid.2013.10.014 (2014).

Volz, A. & Sutter, G. Modified Vaccinia Virus Ankara: History, Value in Basic Research, and Current Perspectives for Vaccine Development. *Adv Virus Res* **97**, 187-243, doi:10.1016/bs.aivir.2016.07.001 (2017).

Liu, R. *et al.* SPI-1 is a missing host-range factor required for replication of the attenuated modified vaccinia Ankara (MVA) vaccine vector in human cells. *PLoS Pathog* **15**, e1007710, doi:10.1371/journal.ppat.1007710 (2019).

Moon, K. B., Turner, P. C. & Moyer, R. W. SPI-1-dependent host range of rabbitpox

virus and complex formation with cathepsin G is associated with serpin motifs. *J Virol*

73, 8999-9010, doi:10.1128/JVI.73.11.8999-9010.1999 (1999).

19   Americo, J. L., Earl, P. L. & Moss, B. Droplet digital PCR for rapid enumeration of viral

genomes and particles from cells and animals infected with orthopoxviruses. *Virology*

511, 19-22, doi:10.1016/j.virol.2017.08.005 (2017).

20   Unger, S. *et al.* FAM111A mutations result in hypoparathyroidism and impaired

skeletal development. *Am J Hum Genet* **92**, 990-995, doi:10.1016/j.ajhg.2013.04.020

(2013).

21   Xiao, T. S. & Fitzgerald, K. A. The cGAS-STING pathway for DNA sensing. *Mol Cell*

51, 135-139, doi:10.1016/j.molcel.2013.07.004 (2013).

22   Chen, C. & Xu, P. Cellular functions of cGAS-STING signaling. *Trends Cell Biol*,

doi:10.1016/j.tcb.2022.11.001 (2022).

23   Alabert, C. *et al.* Nascent chromatin capture proteomics determines chromatin

dynamics during DNA replication and identifies unknown fork components. *Nat Cell*

Biol **16**, 281-293, doi:10.1038/ncb2918 (2014).

24   Ruggiano, A. & Ramadan, K. DNA-protein crosslink proteases in genome stability.

Commun Biol **4**, 11, doi:10.1038/s42003-020-01539-3 (2021).

25   Welter, A. L. & Machida, Y. J. Functions and evolution of FAM111 serine proteases.

Front Mol Biosci **9**, 1081166, doi:10.3389/fmolb.2022.1081166 (2022).

26   Greseth, M. D., Czarnecki, M. W., Bluma, M. S. & Traktman, P. Isolation and

Characterization of vDelta3 Confirm that Vaccinia Virus SSB Plays an Essential Role

in Viral Replication. *J Virol* **92**, doi:10.1128/JVI.01719-17 (2018).

Senkevich, T. G., Katsafanas, G. C., Weisberg, A., Olano, L. R. & Moss, B. Identification of Vaccinia Virus Replisome and Transcriptome Proteins by Isolation of Proteins on Nascent DNA Coupled with Mass Spectrometry. *J Virol* **91**, doi:10.1128/JVI.01015-17 (2017).

Tseng, M., Palaniyar, N., Zhang, W. & Evans, D. H. DNA binding and aggregation properties of the vaccinia virus I3L gene product. *J Biol Chem* **274**, 21637-21644, doi:10.1074/jbc.274.31.21637 (1999).

Mauthe, M. *et al.* Chloroquine inhibits autophagic flux by decreasing autophagosome-lysosome fusion. *Autophagy* **14**, 1435-1455, doi:10.1080/15548627.2018.1474314 (2018).

Luttge, B. G. & Moyer, R. W. Suppressors of a host range mutation in the rabbitpox virus serpin SPI-1 map to proteins essential for viral DNA replication. *J Virol* **79**, 9168-9179, doi:10.1128/JVI.79.14.9168-9179.2005 (2005).

Peng, C. *et al.* Zinc-finger antiviral protein (ZAP) is a restriction factor for replication of modified vaccinia virus Ankara (MVA) in human cells. *PLoS Pathog* **16**, e1008845, doi:10.1371/journal.ppat.1008845 (2020).

Davies, D. H. *et al.* Antibody profiling by proteome microarray reveals the immunogenicity of the attenuated smallpox vaccine modified vaccinia virus ankara is comparable to that of Dryvax. *J Virol* **82**, 652-663, doi:10.1128/JVI.01706-07 (2008).

Lin, Y. C. *et al.* Vaccinia virus DNA ligase recruits cellular topoisomerase II to sites of viral replication and assembly. *J Virol* **82**, 5922-5932, doi:10.1128/JVI.02723-07 (2008).

Figure 1. FAM111A prohibits the replication of vaccinia viruses lacking SPI-1 in A549 cells.

(A) Human A549 cells were transfected with FAM111A or empty vector for 24 hours and infected in triplicates with MVA, MVA+SPI-1, MVA+SPI-1/C16, VACV or VACV-ΔSPI-1 at 0.3 PFU/cell. Cell lysates were analyzed by SDS-PAGE and Western blotting with anti-FAM111A, anti-VACV, or anti-β-actin antibodies at 12 hpi. **(B and D)** Human A549 cells, A549 cells transiently transfected with FAM111A expression vector for 24 h, A549 cells transfected with siNC or siFAM111A for 48 hours, or A549$^{FAM111A-KO}$ cells were infected in triplicates with viruses described above at 0.03 PFU/cell. Viruses were harvested at 24 hpi and virus titers were measured by plaque assay on DF-1 (MVA, MVA+SPI-1, and MVA+SPI-1/C16) or BS-C-1 cells (VACV, VACV-ΔSPI-1). The bars represent the mean values of virus titers **(B)**; Indicated cells were infected in triplicates with viruses described above at 3 PFU/cell. Total cellular DNA was harvested at 6 hpi and viral genomic DNA levels were determined by RT-qPCR using primers for E11 **(D)**. **(C and E)** Human A549 cells were transfected with a vector expressing FAM111A at concentrations of 0.1, 0.5, and 1 µg/mL for 24 hours and then infected in triplicates with MVA or VACV-ΔSPI-1 at 0.03 PFU/cell. Viruses were harvested at 24 hpi and virus titers were determined by plaque assay using DF-1 cells (MVA) or BS-C-1 cells (VACV-ΔSPI-1). Bars represent the mean values of virus titers **(C)**; Indicated cells were infected in triplicates with MVA or VACV-ΔSPI-1 at 3 PFU/cell. Total DNA was harvested at 6 hpi and viral genomic DNA levels were determined by RT-qPCR using primers for E11 **(E)**. **(F)** Human A549 cells were transfected with indicated siRNA for 48 hours and then infected in triplicates with viruses described above at 0.3 PFU/cell. Cell lysates were analyzed by SDS-PAGE and Western blotting with anti-FAM111A, anti-VACV, or anti-β-actin antibodies at 12 hpi. **(G)** Human A549^{WT} or A549$^{FAM111A-KO}$ cells were infected in triplicates with viruses described above at 0.3 PFU/cell and cell lysates were collected and analyzed by SDS-PAGE followed by Western Blotting analysis with anti-FAM111A, anti-VACV or anti-β-actin antibodies at 12 hpi.

Data in **(B-E)** represent the mean values ± SD (standard deviation) of 3 independent experiments. Data in **(A, F, and G)** are representative of 3 independent experiments. Statistics: ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$ by two-sided student's t-test.

Figure 2. The antiviral effect of FAM111A is dependent on its serine protease activity. **(A)** Schematic of the protein structure of FAM111A with previously defined domains, the trypsin catalytic triplet mutants, and PCNA interaction domain mutant. Trypsin-like: the protease domain. PIP box: PCNA interaction domain. DBD: DNA binding domain. Catalytic residues in the protease domain are underlined. U1 and U2 are potential UBLs in FAM111A. **(B)** Human A549 cells transfected with the empty vector or vectors expressing FAM111A^{WT}, FAM111A^{S541A}, FAM111A^{R569H,} or FAM111A$^{S541A+R569H}$ at the final concentration of 1 µg/mL. Cell lysates were collected and analyzed by SDS-PAGE followed by Western blotting analysis using anti-FAM111A or anti-β-actin antibodies. **(C)** Human A549 cells transfected with the empty vector or vectors expressing FAM111A^{WT}, FAM111A^{S541A}, FAM111A^{R569H,} or FAM111A$^{S541A+R569H}$ at the final concentration of 1 µg/mL in the presence or absence of MG132 at 20 µM added 6h post-transfection. Cell lysates were collected and analyzed by SDS-PAGE followed by Western blotting analysis using anti-FAM111A or anti-β-actin antibodies. **(D-F)** Human A549 cells were transfected with an empty vector or vectors expressing FAM111A^{WT}, FAM111A^{S541A}, FAM111A^{R569H,} or FAM111A$^{S541A+R569H}$ at 1µg/ml for 24 hours and then infected in triplicates with MVA or VACV-ΔSPI-1 at 0.3 PFU/cell. Cell lysates were analyzed by SDS-PAGE and Western blotting with anti-FAM111A, anti-VACV, or anti-β-actin antibodies at 12 hpi **(D)**; Indicated cells were infected in triplicates with MVA at 0.03 PFU/cell. Viruses were harvested at 24 hpi and virus titers were determined by plaque assay using DF-1 cells (MVA) or BS-C-1 cells (VACV-ΔSPI-1). Bars represent the mean values of virus titers **(E)**; Indicated cells were infected in triplicates with MVA or VACV-ΔSPI-1 at 3 PFU/cell. Total DNA was harvested at 6 hpi and viral genomic DNA levels were determined by RT-qPCR using primers for E11 **(F)**. **(G)** Human A549 cells were transfected with an empty vector or vectors expressing FAM111A^{WT} or FAM111A$^{PCNA-mut}$ and then infected in triplicates with MVA at 0.3 PFU/cell. Protein lysates were harvested and analyzed with SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-VACV, and anti-β-actin antibodies. Data in **(E and F)** represent the mean values ± SD (standard deviation) of 3 independent experiments. Data in **(B-G)** are representative of 3 independent experiments. Statistics: ns, not significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$ by two-sided student's t-test.

Figure 3. MVA infection prompts protease-dependent relocalization of FAM111A.

(A-C) Human A549 cells plated on coverslips were transfected with vectors expressing FAM111A^{WT}, FAM111A^{S541A}, FAM111A^{R569H,} or FAM111A$^{S541A+R569H}$ and then mock infected or infected in triplicates with MVA at 3 PFU/cell. At 12 hpi, cells were then fixed, permeabilized, blocked, and stained with primary antibodies to FAM111A and VACV followed by fluorescent conjugated secondary antibodies. Hoechst was used to stain DNA **(A and B)**. The confocal analyses were performed in triplicates and the localization of FAM111A was quantified in 100 randomly selected cells and bars represent mean values +/- SD **(C)**. **(D-F)** Human A549 cells were infected in triplicates with MVA at 3 PFU/cell and protein was harvested at time points indicated in the figure. Protein components in the nucleus and cytosol were separated by nuclear and cytoplasmic protein extraction kit and analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, ati-VACV antibodies. Anti-β-actin and anti-histone H3 were used for cytosolic and nuclear loading controls, respectively.

Data in **(C)** represent the mean values ± SD (standard deviation) of 3 independent experiments. Data in **(A, B, D, E, and F)** are representative of 3 independent experiments.

A

	Mab414	FAM111A	Hochest	Merge

Mock

(Zoom) Mock

MVA

(Zoom) MVA

MVA+SPI-1

(Zoom) MVA+SPI-1

— 30 μM or 5 μM (Zoom)

B

Figure 4. FAM111A aggregates near the nuclear membrane and degrades nuclear pore complex proteins after MVA infection in A549 cells.

(A-B) Human A549 cells were mock infected or infected in triplicates with MVA or MVA+SPI-1 at 3 PFU/cell. At 6 hpi, cells were then fixed, permeabilized, blocked, and stained with primary antibodies to Mab414 and FAM111A followed by fluorescent conjugated secondary antibodies. Hoechst was used to stain DNA **(A)**. The confocal analyses were performed in triplicates and the localization of FAM111A was quantified in 100 randomly selected cells and bars represent mean values +/- SD **(B)**;

Data in **(B)** represent the mean values ± SD (standard deviation) of 3 independent experiments. Data in **(A and B)** are representative of 3 independent experiments.

A WCL

FAM111A ——70

cGAS ——70

p-STING ——40

STING ——35

PCNA ——35 ——25

β-actin ——40

Histone H3 ——15

B Nuclear

FAM111A ——70

cGAS ——70

p-STING ——40

STING ——35

PCNA ——35 ——25

β-actin ——40

Histone H3 ——15

C Cytosolic

FAM111A ——70

cGAS ——70

p-STING ——40

STING ——35

PCNA ——35 ——25

β-actin ——40

Histone H3 ——15

Figure 5. The antiviral effect and relocalization of FAM111A are enabled by the activation of the cGAS-STING pathway.

(A-C) Human A549 cells were transfected in triplicates with siNC or sicGAS for 48 hours and then transfected with empty vector, DNA collected from MVA virus particles, poly (dA:dT), or poly (I:C) for 8 hours. Nuclear and cytosolic proteins were separated using nuclear and cytoplasmic protein extraction kits and analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-cGAS, anti-p-STING, anti-STING, anti-PCNA, anti-β-actin, and anti-histone H3 antibodies. Anti-β-actin and anti-histone H3 were used for cytosolic and nuclear loading controls, respectively.

Data in **(A-C)** are representative results of 3 independent experiments.

A

Mass spectrometric screening of viral proteins interacting with FAM111A

A549 cells were transfected with empty vector or flag-tagged FAM111A for 24 h

↓

Confluent A549 cells + 3 PFU/cell of MVA for 24 h

↓

Cells were lysed with IP lysis buffer

12,000 rpm, 10 min → Pellet

↓

Supernatant

↓

For pre-clear, the lysate was incubated with mouse IgG magnetic beads by shaking for 2 h in 4°

Separation on a magnetic stand, 10 s → Pellet

↓

Supernatant

↓

The lysate was incubated with anti-flag magnetic beads by shaking for 18 h in 4°C

Separation on a magnetic stand, 10 s → Pellet

↓

Supernatant

↓

Wash the beads 5 times with TBS

Separation on a magnetic stand, 10 s → Supernatant

↓

Pellet

↓

Mass spectrometry

B emPAI

	I3	E3	D7	H5	A37	F4	A44	F13	A3	J6
Vec+MVA	3.61	5.88	1.43	0.45	1.00	2.09	1.33	1.98	3.00	0.12
FAM111A+MVA	7.03	2.85	0.81	0.75	0.74	0.57	0.11	0.10	0.06	0.03

Figure 6. FAM111A mediates the degradation of viral I3 in a DNA-dependent manner

(A) Mass spectrometric screening of viral proteins interacting with FAM111A. **(B)** Mass spectrometry was used to identify the MVA viral proteins that interacted with FAM111A, and the relative content of different viral proteins was expressed by the exponentially modified protein abundance indice (emPAI). **(C and D)** Human A549 cells were transfected with vectors expressing myc-tagged FAM111A or flag-tagged I3 for 24 hours. Cell lysates were first incubated with or without Benzonase (250 U/mL) at 37°C for 20 min and then incubated with control magnetic beads or myc-/flag- conjugated beads at 4°C for 18 hours. Beads were extensively washed and proteins were eluted with SDS-loading buffer and analyzed by SDS-PAGE and Western blotting analysis. The arrow indicates the correct band size for I3, and the light chain or heavy chain antibody bands are at 25 kDa or 55 kDa, respectively. **(E)** A549 cells plated on coverslips were co-transfected with VACV-I3 and FAM111A for 24 hours and then infected in triplicates with MVA at 3 PFU/cell. At 12 hpi, cells were then fixed, permeabilized, blocked, and stained with primary antibodies to I3 and FAM111A followed by fluorescent conjugated secondary antibodies. Hoechst was used to stain DNA. Scale bars were shown at the bottom-right corners. **(F)** Human A549 cells were transfected with a vector expressing FAM111A at concentrations of 0.1, 0.5, and 1 μg/mL for 24 hours and then infected in triplicates with MVA or MVA+SPI-1 at 0.3 PFU/cell. Cell lysates were analyzed by SDS-PAGE and Western blotting with anti-FAM111A, anti-VACV, anti-I3, anti-E3, or anti-β-actin antibodies at 24 hpi. **(G)** Human A549 cells were co-transfected with flag-tagged I3 or flag-tagged H5 and vectors expressing FAM111A for 24 hours. Cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-flag, and anti-β-actin antibodies.

Data in **(C-G)** are representative of 3 independent experiments.

Figure 7. Cytosolic FAM111A activates DNA-dependent I3 degradation through autophagy.

(A) Human A549 cells were co-transfected with VACV-I3 and vectors expressing FAM111A^{WT} and lab-generated mutants indicated in figure 2(A) for 24 hours. Cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-I3, and anti-β-actin antibodies. **(B)** Human A549 cells were transfected with vectors expressing FAM111A for 24 hours. After waiting for FAM111A to enter the nucleus, indicated cells were transfected with VACV-I3. After 24 hours, cells were analyzed by SDS-PAGE and Western blotting or confocal microscopy. **(C)** FAM111A DNA binding domain mutant, FAM111A^{S541A} mutant with nuclear export signal, and FAM111A nuclear localization signal mutant. **(D)** Human A549 cells were transfected with vectors expressing FAM111A^{WT} and lab-generated mutants indicated in the figure for 24 hours. Then indicated cells were transfected with VACV-I3. After 24 hours, cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-I3, anti-LC3B, and anti-β-actin antibodies. **(E)** Human A549 cells were transfected with an empty vector or vector expressing FAM111A for 24 hours. Then indicated cells were transfected with VACV-I3. Different drugs were added at the same time. After 24 hours, cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-I3, and anti-β-actin antibodies. **(F)** Human A549 cells were transfected with an empty vector or vector expressing FAM111A for 24 hours. Then indicated cells were transfected with VACV-I3. At the same time, different concentrations of CQ were added. After 24 hours, cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-I3, anti-LC3B, and anti-β-actin antibodies. **(G)** Human A549^{WT} or A549$^{FAM111A-KO}$ cells were transfected with an empty vector or vector expressing FAM111A for 24 hours. Then indicated cells were transfected with VACV-I3. At the same time, DMSO or CQ was added. After 24 hours, cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-I3, anti-LC3B, and anti-β-actin antibodies. **(H)** Human A549^{WT} or A549$^{ATG16L1-KD}$ cells were transfected with an empty vector or vector expressing FAM111A for 24 hours. Then indicated cells were transfected with VACV-I3. At the same time, DMSO or CQ was added. After 24 hours, cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-I3, anti-ATG16L1, anti-LC3B, and anti-β-actin antibodies.

Data in **(A and D-H)** are representative of 3 independent experiments.

A

FAM111A | Anti-Vaccinia | Hochest | Merge

FAM111A MVA

FAM111A MVA+SPI-1

— 5 μM

B

FAM111A − +
SPI-1 + +

IP: FAM111A

SPI-1
— 55
— 40 ←
— 35
— 25

FAM111A
— 70

Input

FAM111A — 70
SPI-1 — 40
β-actin — 40

C

FAM111A − +
SPI-1 + +

IP: SPI-1

FAM111A — 70

SPI-1
— 55
— 40 ←
— 35
— 25

Input

FAM111A — 70
SPI-1 — 40
β-actin — 40

D

SPI-1 − VACV − VACV MPXV LSDV
I3 − + + + + +
FAM111A − − + + + +

FAM111A — 70
SPI-1 — 40
I3 — 35
β-actin — 40

E

SPI-1 − − WT T305R F318A T305R+F318A
FAM111A − + + + + +

FAM111A — 70
SPI-1 — 40
I3 — 35
β-actin — 40

Figure 8. Vaccinia virus SPI-1 antagonizes the nuclear export and antiviral activity of FAM111A.

(A) Human A549 cells were transfected with an empty vector or vector expressing FAM111A for 24 hours. Then indicated cells were infected with MVA or MVA+SPI-1 at 3 PFU/cell. After 12 hours, the cells were then fixed, permeabilized, blocked, and stained with primary antibodies to FAM111A and VACV followed by fluorescent conjugated secondary antibodies. Hoechst was used to stain DNA. Scale bar at bottom. **(B and C)** Human A549 cells were transfected with vectors expressing myc-tagged FAM111A or flag-tagged SPI-1 for 24 hours. Cell lysates were incubated with control magnetic beads or myc-/flag-conjugated beads at 4°C for 18 hours. Beads were extensively washed and proteins were eluted with SDS-loading buffer and analyzed by SDS-PAGE and Western blotting analysis. The arrow indicates the correct band size for SPI-1, and the light chain or heavy chain antibody bands are at 25 kDa or 55 kDa, respectively. **(D)** Human A549 cells were co-transfected with flag-tagged SPI-1 of different viruses and a vector expressing FAM111A for 24 hours. Then indicated cells were transfected with VACV-I3. After 24 hours, cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-flag, anti-I3, and anti-β-actin antibodies. **(E)** Human A549 cells were co-transfected with flag-tagged SPI-1 of different mutants and a vector expressing FAM111A for 24 hours. Then indicated cells were infected with MVA at 0.3 PFU/cell. After 24 hours, cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, anti-flag, anti-I3, and anti-β-actin antibodies;

Data in **(A-E)** are representative of 3 independent experiments.

Figure 9. cGAS causes FAM111A protease-dependent nuclear export and inhibits MVA replication through autophagic degradation of I3.

(A) After SPI-1-deficient MVA or VACV invaded cells, the viral genomic DNA was recognized by the cGAS-STING signaling pathway, and its downstream signal activated the upregulation of FAM111A expression, allowing FAM111A in the nuclear to enter the cytosolic by degrading the NPC. **(B)** After exporting from the nucleus, FAM111A is recruited by DNA to the virus factory located in the cytoplasm and binds to viral protein I3, which is degraded by the autophagy-lysosome pathway together. **(C)** SPI-1 of different poxviruses can inhibit the serine protease activity of FAM111A and prevent its nuclear export to degrade I3.

Figure S1. Monoclonal Identification of A549 FAM111A-KO Cell Line.

A549^FAM111A-KO cell line was constructed using the CRISPR-Cas9 method. After picking different monoclonal cells and culturing the cells into sheets, cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A and anti-β-actin antibodies.

Data are representative of 3 independent experiments.

Figure S2. The antiviral function of human-FAM111A in different species of cells.

(A) Monkey BS-C-1 cells were transfected with a vector expressing FAM111A at concentrations of 0.1, 0.2, 0.5, 1, and 2 µg/mL for 24 hours and then infected in triplicates with MVA 47.1 at 0.3 PFU/cell. Cell lysates were analyzed by SDS-PAGE and Western blotting with anti-FAM111A, anti-VACV, anti-I3, or anti-β-actin antibodies at 24 hpi. **(B)** Chicken DF-1 cells were transfected with a vector expressing FAM111A at concentrations of 1 µg/mL for 24 hours and then infected in triplicates with MVA or MVA+SPI-1 at 0.3 PFU/cell. Cell lysates were analyzed by SDS-PAGE and Western blotting with anti-FAM111A, anti-VACV, anti-I3, or anti-β-actin antibodies at 4 hpi. **(C)** Chicken DF-1 cells were transfected with a vector expressing FAM111A at concentrations of 1 µg/mL for 24 hours and then infected in triplicates with MVA or MVA+SPI-1 at 0.03 PFU/cell. Viruses were harvested at 24 hpi and virus titers were determined by plaque assay using DF-1 cells. Bars represent the mean values of virus titers;

Data in **(A-C)** are representative of 3 independent experiments.

Figure S3. Expression levels of endogenous FAM111A after MVA and VACV-WR infection.

Human A549 cells were infected with MVA or VACV-WR for 0, 6, 12, and 18 hours at 3 PFU/cell. Cell lysates were analyzed by SDS-PAGE and Western blotting with anti-FAM111A, anti-VACV, or anti-β-actin antibodies;

Data are representative of 3 independent experiments.

Figure S4. The serine protease catalytic triad at the C-terminus of FAM111A is highly conserved across species.

The FAM111A amino acid sequences of 68 mammals were downloaded from NCBI (https://www.ncbi.nlm.nih.gov/), and Snapgene was used for multiple sequence alignment. The FAM111A C-terminal trypsin-like protease domain contains a typical H385-D439-S541 catalytic triad.

Figure S5. Virus infection triggers the protease-dependent relocalization of FAM111A in HeLa.

(A-C) Human HeLa cells plated on coverslips were transfected with vectors expressing FAM111A^{WT}, FAM111A^{S541A}, FAM111A^{R569H,} or FAM111A$^{S541A+R569H}$ and then mock infected or infected in triplicates with MVA at 3 PFU/cell. At 12 hpi, cells were then fixed, permeabilized, blocked, and stained with primary antibodies to FAM111A and VACV followed by fluorescent conjugated secondary antibodies. Hoechst was used to stain DNA **(A and B)**. The confocal analyses were performed in triplicates and the localization of FAM111A was quantified in 100 randomly selected cells and bars represent mean values +/- SD **(C)**;

Data in **(C)** represent the mean values ± SD (standard deviation) of 3 independent experiments. Data in **(A and B)** are representative of 3 independent experiments.

Figure S6. Protease-dependent disruption of the NPC upon overexpression of FAM111A. **(A)** A549 cells were transfected with vectors expressing FAM111A[WT] and lab-generated mutants indicated in figure 2(A) for 24 hours. The cells were then fixed, permeabilized, blocked, and stained with primary antibodies to Mab414 and FAM111A followed by fluorescent conjugated secondary antibodies. Hoechst was used to stain DNA. Scale bar at bottom. **(B)** A549 cells were transfected with vectors expressing FAM111A[WT] and indicated mutants for 24 hours. Cell lysates were analyzed by SDS-PAGE and Western blotting analysis using anti-FAM111A, Mab414 (anti-NUP62), and anti-β-actin antibodies; Data in **(A and B)** are representative of 3 independent experiments.

Figure S7. Colocalization of different FAM111A mutants with I3 in A549 cells

Human A549 cells were transfected with vectors expressing FAM111A^{WT} and indicated mutants for 24 hours. Then indicated cells were transfected with VACV-I3. After 24 hours, cells were then fixed, permeabilized, blocked, and stained with primary antibodies to I3 and FAM111A followed by fluorescent conjugated secondary antibodies. Hoechst was used to stain DNA. Scale bars are shown at bottom.

Data are representative of 3 independent experiments.

Figure S8. FAM111A does not degrade I3 through the ubiquitin-proteasome pathway. A549 cells were co-transfected with myc-tagged FAM111A, flag-tagged I3, and HA-tagged Ub for 24 hours. Cell lysates were incubated with control magnetic beads or flag-conjugated beads at 4°C for 18 hours. Beads were extensively washed and proteins were eluted with SDS-loading buffer and analyzed by SDS-PAGE and Western blotting analysis. The light chain or heavy chain antibody bands are at 25 kDa or 55 kDa, respectively;

Data are representative of 3 independent experiments.

Figure S9. Subcellular localization of SPI-1 in A549 cells.

Human A549 cells were infected with MVA+SPI-1 at 3 PFU/cell. After 24 hours, the cells were then fixed, permeabilized, blocked, and stained with primary antibodies to SPI-1 followed by fluorescent conjugated secondary antibodies. Hoechst was used to stain DNA. Scale bar at bottom.

Data are representative of 3 independent experiments.