Targeting sterol regulatory element binding proteins (SREBPs) activation lowers intraocular pressure
by halting its mechanosensing function in the trabecular meshwork

Ting Wang[1,2], Avinash Soundararajan[1], Jeffery Rabinowitz[3], Anant Jaiswal[4], Timothy Osborne[4],
Padmanabhan Pattabiraman[1,2]

[1]Glick Eye Institute, Department of Ophthalmology, Indiana University School of Medicine, 1160 West
Michigan Street, Indianapolis, Indiana, 46202, United States of America

[2]Stark Neuroscience Research Institute, 320 W. 15th Street, Indiana University School of Medicine,
Indiana, 46202, United States of America

[3]Departments of Ophthalmology, Case Western Reserve University, Cleveland, Ohio , United States of
America

[4]Institute for Fundamental Biomedical Research, Departments of Medicine and Biological Chemistry,
Johns Hopkins University School of Medicine, St. Petersburg, Florida, 33701, United States of America

Significance

Elevated intraocular pressure (IOP) is a major risk factor for glaucoma, which is an optic neuropathy and a leading cause of irreversible blindness. Targeting trabecular meshwork (TM) outflow pathway tissue is a significant modality to lower the IOP. The contractile properties of TM tissue play an important role in the regulation of IOP. Though lipids are known to contribute to TM contractility and IOP regulation, the contributions of the lipogenic pathway are not clear. In this study, we identify the mechanosensing role of the lipogenic transcription factors sterol regulatory element binding proteins (SREBPs) in TM. Further, we show that pharmacological or molecular inactivation of SREBPs can lower IOP by modifying actin contractility and ECM remodeling. Thus signifying a novel relationship between SREBP activation status and achieving IOP homeostasis.

Abstract

Trabecular meshwork (TM) cells are highly contractile and mechanosensitive to aid in maintaining intraocular pressure (IOP) homeostasis. Lipids are attributed to modulating the TM contractility with poor mechanistic understanding. In this study using human TM cells, we identify the mechanosensing role of the transcription factors sterol regulatory element binding proteins (SREBPs) involved in lipogenesis. By constitutively activating SREBPs and pharmacologically inactivating SREBPs, we have mechanistically deciphered the attributes of SREBPs in regulating the contractile properties of TM. The pharmacological inhibition of SREBPs by fatostatin and molecular inactivation of SREBPs *ex vivo* and *in vivo* respectively results in significant IOP lowering. As a proof of concept, fatostatin significantly decreased the SREBPs responsive genes and enzymes involved in lipogenic pathways as well as the levels of the phospholipid, cholesterol, and triglyceride. Further, we show that fatostatin mitigated actin polymerization machinery and stabilization, and decreased ECM synthesis and secretion. We thus postulate that lowering lipogenesis in the TM outflow pathway can hold key to lowering IOP by modifying the TM biomechanics.

1. Introduction:

Chronic and sustained elevation in intraocular pressure (IOP) is a significant risk factor for
primary open angle glaucoma (POAG) (1-3). A rise in IOP above normal causes mechanical stress and
reduced blood flow to the optic nerve(4). This increases the risk of glaucoma with significant
consequences on vision and quality of life (5-7). Glaucoma is a major public health concern and the
second leading cause of irreversible blindness worldwide afflicting mainly the aging population (8, 9).
The estimated number of glaucoma patients in the world is around 80 million with over 2 million in the
United States alone (10, 11). Interestingly lowering IOP is neuroprotective and the most effective way to
delay the onset of POAG and to halt the progression towards vision loss (12-14). A 20% decrease in IOP
significantly alleviates the risk of developing glaucoma (15). The IOP is maintained by the balance
between the generation of AH via the ciliary body and drainage mainly via the conventional outflow
pathway, which includes trabecular meshwork (TM), juxtacanalicular TM tissue (JCT), and Schlemm's
canal (SC) (16). The TM contains an extracellular matrix (ECM) organized into a network of beams lined
by highly contractile and mechanosensitive TM cells sensing pressure changes (17-19). Increased actin
contractility in TM and ECM hyperdeposition in TM outflow pathway augment AH outflow resistance
and IOP elevation (20-22). In POAG, the increased IOP is directly correlated with the increased outflow
resistance, and the most characteristic structural change in TM is increased tissue stiffness including
ECM accumulation, increased cell contractility, and cell-ECM connections (18, 23-25).

Cellular lipids play important roles in tissue biomechanics by modulating the plasma membrane
remodeling, signal transduction cascades, and actin cytoskeleton-cell adhesions linked to the ECM (26).
Moreover various lipids found in TM and AH are known to impact IOP homeostasis (27-29). This is
attributed via modulating the signaling events including altering the TM contractility and tissue stiffness.
Hyperlipidemia is associated with an increased risk of elevated IOP and glaucoma with poor mechanistic
understanding (30-32). Interestingly, cholesterol-lowering statins lower IOP and reduce the incidence of
glaucoma (33-36). Mechanistic studies in TM point at statin-mediated disruption of membrane
association of Rho GTPase due to inhibition of the isoprenylation of Rho (36, 37). We recently found that
cyclic mechanical stretch, which mimics mechanical stress on TM, significantly altered lipid contents and
increased the expression of cholestogenic enzymes like squalene synthase, 3-hydroxy-3-methylglutaryl
coenzyme A (HMG-CoA) synthase (HMGCS) and the transcriptional regulator of lipid biogenesis sterol
regulatory element binding protein 2 (SREBP2) in human TM (HTM) cells (38). The SREBPs are master
regulator of cellular lipogenesis and occur as three isoforms – SREBP1a, SREBP1c, and SREBP2 (39-41).
SREBP1a is involved in overall lipid biosynthesis, SREBP1c regulates fatty acid synthesis, whereas SREBP2
modulates sterol biosynthesis specifically (39-42). SREBPs are basic-helix-loop-helix leucine zipper
(bHLH-LZ) transcription factors. Each SREBP precursor has three domains: (1) an NH2-terminal domain
including transactivation domain, which is rich in serine and proline and the bHLH-LZ region for DNA
binding and dimerization; (2) two hydrophobic transmembrane spanning segments; (3) a COOH-terminal
domain (43, 44). Within the cell, following SREBPs mRNA translation, the proform SREBPs (Pro-SREBPs)
are bound to SREBP cleavage activating protein (SCAP) forming a SREBP-SCAP complex located on the ER
membrane. Stimuli including low lipids levels promote SCAP to guide the whole complex to leave the ER
membrane to the Golgi membrane. In the Golgi, the SREBP-SCAP complex is cleaved by site 1 protease
(S1P) and site 2 protease (S2P), releasing SCAP from the complex and releasing the active or nuclear
SREBPs (N-SREBPs) from Golgi to translocate into the nucleus. Inside the nucleus, N-SREBPs bind to the
sterol response element (SRE) on the promoter region of the responsive genes and promote their
transcription. These target genes are involved in the fatty acid, triglycerides, and cholesterol
biosynthetic pathways by modulating the expression of rate limiting enzymes like acetyl-CoA
carboxylase (ACC), fatty acid synthase, and 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA)

reductase (HMGCR) (42, 44-47). Interestingly, it was earlier reported that there is an isoprenylation of
RhoA-dependent and cholesterol-independent actin cytoskeleton contraction, which prevents SREBP1
dependent lipogenesis in human immortalized cancer lines and *Drosophila* (48). Such regulatory
mechanistic evaluation of lipogenic pathways in AH outflow tract and the role of SREBPs in TM biology
lacks understanding. By regulating lipid biogenesis in TM, we believe that SREBPs activities play a
significant role in defining actin polymerization and ECM remodeling (49-52). Investigating the
mechanosensing function via SREBPs activation paradigm in TM can elucidate how TM cells actively
sense and transduce mechanical signals via SREBPs. Thus, we report the biological relevance of SREBPs
activation in TM for active pressure sensing to convert mechanical stimuli into TM contractility and
mechanistically decipher how the inactivation of SREBPs in the TM outflow pathway lowers IOP.

2. Material and Methods:

2.1 Materials:

The reagents and antibodies used in this study are documented in Table 1.

TABLE 1:

Reagents and antibodies	Catalog number	Company
Vinculin	V9131	R&D system
β-actin	A1798	R&D system
Collagen 1α (COL1A)	ab34710	R&D system
LIMK1	sc515585	Santa Cruz Biotechnology
SREBP1	sc13551	Santa Cruz Biotechnology
SREBP2	sc13552	Santa Cruz Biotechnology
Paxillin	sc365379	Santa Cruz Biotechnology
GAPDH	sc32233	Santa Cruz Biotechnology
FAS	sc-48357	Santa Cruz Biotechnology
Fascin 1	sc46675	Santa Cruz Biotechnology
SREBP1	NB100-2215	NOVUS Biologicals
SREBP2	NBP1-54446	NOVUS Biologicals
SREBP2	LS-B1609-BCD1-1	Life Span Bioscience
SCAP	12266-1-AP	Proteintech
SCAP	MABS2288	Sigma-Aldrich
Collagen 1α (COL1A)	ABT257	Sigma-Aldrich
HMGCR	PA5-37367	Invitrogen/ThermoFisher
Lamin B1	PA5-19468	Invitrogen/ThermoFisher
Total-paxillin (T-paxillin)	2542S	Cell Signaling Technology
Phospho-paxillin (P-paxillin)	69363S	Cell Signaling Technology
Total-cofilin (T-cofilin)	5175S	Cell Signaling Technology
Phospho-cofilin (P-cofilin)	3311S	Cell Signaling Technology
ACC	3662S	Cell Signaling Technology
Fibronectin		Gift from Dr. Harold Erickson, Duke University
Donkey-anti-goat IgG	705-035-003	Jackson ImmunoResearch
Donkey-anti-mouse IgG	715-035-150	Jackson ImmunoResearch
Donkey-anti-rabbit IgG	715-035-144	Jackson ImmunoResearch

Goat-anti-Mouse IgG Alexa Fluro Plus 488	A32723	ThermoFisher
Goat-anti-Mouse IgG Alexa Fluor Plus 594	A32742	ThermoFisher
Goat-anti-Rabbit IgG Alexa Fluor Plus 488	A32731	ThermoFisher
Goat-anti-Rabbit IgG Alexa Fluor Plus 594		ThermoFisher
Phalloidin (Alexa Fluor Plus 488 PHA)	A12379	Invitrogen/ThermoFisher
Phalloidin (Alexa Fluor Plus 647 PHA)	A22287	Invitrogen/ThermoFisher
DAPI	UH2820551	Sigma-Aldrich
Fluoromount-G	E139999	Invitrogen/ThermoFisher
Fatostatin	341329	Sigma-Aldrich

2.2 Primary TM cell culture:

Primary human TM (HTM) cells were cultured from TM tissue isolated from the leftover donor corneal
rings after they had been used for corneal transplantation at the Indiana University Clinical Service,
Indianapolis, and characterized as described previously (20, 53). HIPPA compliance guidelines were
adhered for the use of human tissues. The usage of donor tissues was exempt from the DHHS regulation
and the IRB protocol (1911117637), approved by the Indiana University School of Medicine IRB review
board.

Primary porcine TM (PTM) cells were cultured from fresh porcine TM globes obtained from Indiana
Packers, Delphi, In, USA, as described previously (20).

All experiments were conducted using confluent HTM or PTM cultures as mentioned, using cells in
between passage four to six, and were performed after overnight serum starvation unless mentioned,
otherwise. All experiments in this manuscript were performed using biological replicates.

2.3 Construction of replication-deficient recombinant adenovirus expressing N-SREBPs

Generation of replication-defective recombinant adenovirus expressing N-SREBPs under CMV promoter
was performed using ViraPower Adenoviral Expression System (#K4930-00, Invitrogen) as described
earlier (54). N-SREBP1a (1470 bps) , N-SREBP1c (1398 bps), and N-SREBP2 (1440 bps) were amplified
using high-fidelity PCR (Advantage-HF 2 PCR kit, #639123; Clontech) from human pcDNA3.1-2xFLAG-
SREBP1a (#26801, Addgene), pcDNA3.1-2xFLAG-SREBP1c (#26802, Addgene) and pcDNA3.1-2xFLAG-
SREBP-2 (#26807, Addgene). Amplified N-SREBP1a (Ad5-N-SREBP1a), N-SREBP1c (Ad5-N-SREBP1c), and
N-SREBP2 (Ad5-N-SREBP2) were purified and titres determined.

2.4 Adenovirus-mediated gene transduction in HTM cells

Primary HTM cells were grown on gelatin-coated glass coverslips or in plastic Petri dishes and were
infected with Ad5-N-SREBP1a, Ad5-N-SREBP1c, and Ad5-N-SREBP2 or a control empty virus (AdMT) at 50
multiplicities of infection for 24 h followed by 48 h serum starvation. Cells were then washed with 1X

PBS and were fixed for immunofluorescence (IF) analysis and collected using either TRIZOL for RNA
extraction or 1X RIPA for protein extraction.

2.5 Optimization fatostatin for *in vitro* dosing

Fatostatin directly binds to SCAP and inhibits the SREBP-escorting function of SCAP, thereby restricting
the translocation of SREBPs from ER to the Golgi and subsequently causing SREBPs inactivation and
reducing lipogenesis and fat accumulation (55-57). Overnight serum starvation (~16 h) was carried out
on PTM and HTM cultures before treatments. To identify the optimal concentration of fatostatin for
lowering SREBPs activation, PTM cells were subjected to 0.2, 2, and 20 µM fatostatin reconstituted in
serum-free DMEM for 24 h respectively. Post-treatment, PTM cells were either used for cell viability
assay or protein collected for total and nuclear fraction. HTM cells were subjected to the optimal
concentration of fatostatin and post-treatment, cells were collected in TRIZOL for RNA isolation or in 1X
radioimmunoprecipitation assay (RIPA) buffer for protein isolation along with the cell culture media and
stored in -80 °C until used.

2.6 Cell viability assay

The cell viability assay was performed using fluorescein diacetate (FDA) and propidium iodide (PI)
staining. PTM cells were grown in 6-well plates until they attained 90% confluency. The cells were
treated with a final concentration of 20 µM fatostatin for 24 h followed by overnight serum starvation.
FDA-PI double staining was performed based on a previously published protocol with minor
modifications (58).

2.7 Cyclic mechanical Stretch:

HTM cells were subjected to cyclic mechanical stretch as published earlier (38). 2 h post stretch, cells
were fixed and processed for IF analysis.

2.8 Porcine *ex vivo* elevated IOP model

Elevated IOP was modeled using porcine *ex vivo* perfusion system as described earlier (20, 54). Post
perfusion the TM tissue was carefully isolated and frozen at -80°C until further processing.

**2.9 *Ex vivo* porcine anterior segment perfusion culture to assess the effect of SREBPs inactivation on
IOP**

The effect of pharmacological SREBPs inactivation on IOP regulation was studied using *ex vivo* porcine
anterior segment perfusion culture as described earlier (20). After achieving a stable baseline IOP, the
IOP was continuously recorded in the control eyes that received a sham DMSO treatment, and the
experimental eyes received 20 µM fatostatin for up to 30h. The changes in IOP (∆IOP) were calculated as
the change in pressure relative to the baseline of drug/sham perfusion. At the end of the perfusion
studies, TM tissues were either sectioned for histology studies and fixed with 4% paraformaldehyde, or
the TM was collected for protein analysis.

2.10 SCAP-Floxed (*SCAP^{f/f}*) mice

All experiments were performed in compliance with the Association for Research in Vision and
Ophthalmology (ARVO) Statement for the Use of Animals in Ophthalmic and Vision Research. Mice
carrying the floxed Scap allele (B6;129-*Scap^{tm1Mbjg}/J*) (59) were obtained from Dr. Tim Osborne, bred,

and housed under a standard 12 h light and dark cycle with food and water provided *ad libitum* in the
laboratory animal resource center (LARC). Exon 1 of the SCAP gene is flanked with LoxP sites, which are
targets for bacteriophage P1 Cre recombinase originally made by Dr. Horton, UTSW Med Center, Dallas
(59, 60). They are viable, fertile, normal in size, and do not display any physical or behavioral
abnormalities. The genotype was identified by PCR analysis of tail biopsies. All animal procedures were
approved by the Institutional Animal Care and Use Committee at the Indiana University School of
Medicine and conducted following the Declaration of Helsinki.

2.11 Molecular inactivation of SREBPs *in vivo* in *SCAP^{f/f}* mice

Molecular inactivation of SREBPs was achieved by knocking down SCAP in *SCAP^{f/f}* mice via intravitreal
injection of adenovirus expressing Cre [Ad5.CMV.iCre-eGFP] (1.6×10^{11} pfu/ml). The Ad5.CMV.eGFP
(2.6×10^{11} pfu/ml) was used as an adenovirus control and saline injection was used for injection control.
These adenoviruses were purchased from the Vector Development Lab, Baylor College of Medicine. Four
groups were included in the study: untouched, saline injection, Ad5.CMV.eGFP injection, and
Ad5.CMV.Cre-eGFP injection. A 33-gauge needle with a glass microsyringe (10 µL volume; Hamilton
Company) was used for the intravitreal injection. Intravitreal injections were performed only in the left
eye under a surgical microscope (Trinocular Stereo Zoom Microscope, AmScope). The needle was
inserted through the equatorial sclera into the vitreous chamber at an angle of $\sim 45°$, carefully avoiding
contact with the posterior lens capsule or the retina. The right eyes in all groups were left untouched.
Viral vectors or saline in a volume of 2 µL were slowly injected into the vitreous over the course of 30
seconds. The needle was then left in place for an additional 45 s to avoid leakage before being
withdrawn. Before and during intravitreal injection, mice were anesthetized with isoflurane (2.5%)
(#NDC 46066-755-04, Patterson Veterinary, Loveland, CO, USA) and 100% oxygen using SomnoSuite
(Kent Scientific Corporation, Torrington, CT, USA). Initially, a total of 15 animals/group were used, but
post-injection due to infection, cataract formation, and death the final numbers were reduced to 9-10
mice/group for the data acquisition with roughly equal number of adult (10-14 months) and old (18-24
months) mice.

2.12 Measurement of IOP

The mice were anesthetized with 2.5% isoflurane and 100% oxygen using SomnoSuite. A previously
validated commercial rebound tonometer (TonoLab, Colonial Medical Supply) was used to take three
sets of six measurements of IOP in each eye. The IOP measurement and recording were carried out in a
blinded manner by the experimenter to minimize outcome bias. Right and left eye measurement sets
were alternated with the initial eye selected randomly. All measurements were taken between 3 and 5
min after isoflurane inhalation based on prior studies. As indicated by the manufacturer, the tonometer
was fixed horizontally for all measurements, and the tip of the probe was 2–3 mm from the eye. The
probe contacted the eye perpendicularly over the central cornea. The average of a set of measurements
was accepted only if the device indicated that there was "no significant variability" (as per the protocol
manual; Colonial Medical Supply). Daytime measurements were taken between 11:00 and 15:00 h.
Baseline IOP was measured before saline and viral vector injection. After injection, IOP was measured
every 10 days until 50 days post-injection.

2.13 Gene expression analysis

Total RNA was extracted from HTM cells using the Trizol method following the manufacturer's protocol.
The RNA amounts measurement, cDNA conversion, and qPCR reaction were conducted as published
before (54). Sequence-specific forward and reverse oligonucleotide primers for the indicated genes are
provided in **Table 2**.

q PCR reaction was performed and calculated by the delta-delta Ct method as described earlier (54, 61).
The normalization was performed using 18S ribosomal RNA (18S rRNA) for HTM.

TABLE 2:

SREBP1 Transcript 1 (F1V1)	Forward	CTGACCGACATCGAAGGTGA
	Reverse	AAGTGCAATCCATGGCTCCG
SREBP1 Transcript 2 (F1V2)	Forward	GCTGACCGACATCGAAGACA
	Reverse	CAGGAGGTGGAGACAAGCTG
SREBP1 Transcript 3 (F1V3)	Forward	AACGGCTTCAAAAATCCGCC
	Reverse	CCAGCATAGGGTGGGTCAAA
SREBP2 Transcript 1 (F2V1)	Forward	GGTTGTCGGGTGTCATGGG
	Reverse	TTGCAGCATCTCGTCGATGT
SCAP Transcript 1	Forward	CTGGAGAAGCTGGACTGAGC
	Reverse	CCCAAAGTGCCTGACAGATG
Total-SREBP1	Forward	CAACACAGCAACCAGAAACTC
	Reverse	CTCCACCTCAGTCTTCACG
Total-SREBP2	Forward	ATTCAGCACCACTCCGCAG
	Reverse	GCCCCATCATTACAGGCATTG
LDLR	Forward	TTCACTCCATCTCAAGCATCG
	Reverse	ACTGAAAATGGCTTCGTTGATG
HMGCS1	Forward	TTCCCTTGCATCTGTTCTAGC
	Reverse	TTTTATCAAGAGCAGACCCCG
HMGCR	Forward	ACAGATACTTGGGAATGCAGAG
	Reverse	CTGTCGGCGAATAGATACACC
ACC	Forward	ACAGTGGAGCAAGAATCGG
	Reverse	AATGGACAGAGTTGAGAGCAC
FASN	Forward	AAGCTCTTTCACTTCGGAGG
	Reverse	GGGCATTAACACTTTTGGACG
PPARG	Forward	ATCAAAGTGGAGCCTGCATC
	Reverse	CGACATTCAATTGCCATGAG
ACS	Forward	GGGAAGAGTTGGAAGGTGAAG
	Reverse	GCAGCCATCTCCTGTAACATAG
GPAT	Forward	GTGGAGTGTAGCAAGAGGTG
	Reverse	CAGGGAAAGTAGAGCAGACAC
FN	Forward	GGTGACACTTATGAGCGTCCTAAA
	Reverse	AACATGTAACCACCAGTCTCATGTG
COL1α	Forward	GAGAGCATGACCGATGGATT
	Reverse	CCTTCTTGAGGTTGCCAGTC
COL4α	Forward	TGTGGATCGGCTACTCTTTTG
	Reverse	TAGTAATTGCAGGTCCCACG
	Forward	TCAAGTCCTTCACCAAGCG

COL6α	Reverse	ATCTCCACCTCGTCACTGTA
TNC	Forward	CACTACACAGCCAAGATCCAG
	Reverse	TCGTGTCTCCATTCAGCATTG
TGFβ2	Forward	TTGACGTCTCAGCAATGGAG
	Reverse	TTCGCCTTCTGCTCTTGTTT
SSPN	Forward	AGGTTGACGAACGGACATG
	Reverse	TGTGAGCTGCGAATAGTGG
18S	Forward	CTACCACATCCAAGGAAGCA
	Reverse	TTTTTCGTCACTACCTCCCCG

2.14 Protein Analysis by Immunoblotting

For nuclear fraction, cells were collected from a 10 cm plate using fractionation buffer (20 mM HEPES,
10 mM KCl, 2 mM MgCl2, 1 mM EDTA, 1 mM EGTA, 1 mM DTT). After using a 1 mL syringe to pass cell
suspension through a 27 gauge needle several times, the sample was centrifuged and a pellet that
contains nuclei was collected. The pellet was washed, dispersed, and centrifuged again, then the nuclear
fraction was collected by TBS with 0.1% SDS.

The whole cell lysates containing total protein were prepared using 1X RIPA buffer composed of 50 mM
Tris-HCL (PH 7.2), 150 mM NaCl, 1% NP-40, 0.1% SDS, 1 mM EDTA, and 1 mM PMSF with protease and
phosphatase inhibitors (#A32961, ThermoFisher Scientific) and then sonicated. Nuclear extraction was
collected using 20mM HEPES (pH 7.4), 10mM KCl, 2mM MgCl2, 1mM EDTA, and 1mM EGTA. Media was
collected and concentrated using Nanosep® Centrifugal Devices with Omega™ Membrane 10K
(#OD010C34, Pall), and add 1X RIPA to collect the media protein.

After determining the protein concentration and the immunoblotting process was performed as
published earlier (38, 54). Blots were stripped using mild stripping buffer if required to reprobe for the
loading control and multiple proteins within the same molecular weight range. The data were
normalized to GAPDH, β-actin, LaminB1, or Ponceau S. Semi-quantitative analyses and fold changes
were calculated from the band intensities measured using ImageJ software.

2.15 Protein distribution analysis by immunohistochemistry (IHC) and IF

Tissue sections from formalin-fixed, paraffin-embedded human donor whole globes eyes were prepared
at Case Western Reserve University, Cleveland, Ohio. Paraffin-embedded porcine TM tissue slides and
mice eyeball slides were prepared at the Histology Core, Indiana University, and immunolabeling was
performed. IF staining was done on HTM cells grown on 2% gelatin-coated glass coverslip. The
methodologies used for IHC and IF has been published from the lab earlier (20, 54).

All the slides were observed under a Zeiss LSM 700 confocal microscope, and z-stack images were
obtained and processed using Zeiss ZEN image analysis software.

2.16 Quantitative image analysis

ImageJ (version 1.53a) software was used to analyze the SREBPs nuclear fluorescence intensity in IF
HTM cell images and SREBPs and ECM proteins in IHC tissue images. Specifically, the
immunofluorescence image obtained from the HTM cells and tissue section was converted into an 8-bit
image, then the threshold default setup under Adjust was used to convert the image from grayscale into

a binary image. Next, the region of interest (ROI) tool was chosen for analysis. In this analysis, the

nuclear area in the HTM cells or different ROIs (equal area) in the TM outflow pathway was chosen in an

image, and the intensity was measured and compared between the control and fatostatin.

2.17 Quantitative filamentous actin/globular actin (F-actin/G-actin) ratio measurement

The F-actin/G-actin ratio was measured using G-Actin/F-Actin In Vivo Assay Biochem Kit (#BK037,

Cytoskeleton, Inc.), following the manufacturer's instructions.

2.18 Lipid content measurement

HTM cells were plated on 100 mm plates. Post-treatment, cells were counted and equal numbers were

collected to be frozen at -80°C. Then the phospholipid, cholesterol, and triglyceride from HTM cells were

extracted and measured with a phospholipid quantification assay kit (#CS0001, Sigma-Aldrich),

cholesterol quantitation kit (#MAK043, Sigma-Aldrich) and triglyceride quantification kit (#MAK266,

Sigma-Aldrich) respectively, following manufacturer's instructions.

2.19 Statistical analysis

All data presented were based on greater than three independent observations. The graphs represent

the mean ± standard error. All statistical analyses were performed using Prism 9.3.1 (GraphPad). The

difference between the two groups was assessed using paired Student's t-test when parametric

statistical testing was applicable. When multiple groups were compared, a two-way analysis of variance

followed by the Tukey post hoc test was applied. The p-value < 0.05 was considered a statistically

significant difference between the test and control samples.

3. Results:

3.1 Mechanosensing via SREBPs activation in HTM cells aids in increasing the actin-based contractility

In our recent work (38) on HTM cells subjected to mechanical stress, we showed a significant increase in

the N-SREBP2 that is involved in cholesterol biogenesis. To better understand the mechanosensing

functionality of SREBPs, here we provide empirical evidence by utilizing multiple systems to qualitatively

and quantitatively assess the potential changes in SREBPs localization under mechanical stress. Firstly,

we performed IF analysis post cyclic mechanical stretch to look at the changes in the localization of

SREBPs. After 2 h of stretch (ST), HTM cells showed strong nuclear localization of SREBP1 (top panel

second-row third column, green staining) and 2 (bottom panel second-row third column, green staining)

(Figure 1A). However, there was no significant difference in actin distribution in HTM cells after 2 h ST

(top and bottom panel second-row fourth column, red staining) **(Figure 1A)**. Further substantial

evidence for increased nuclear localization of SREBPs under mechanical stress was obtained from the

acute pressure stress model (20, 54, 62). In porcine anterior segments subjected to two times (2x)

elevated pressure (30 mmHg) compared to normal pressure (15 mmHg), we found a significant increase

in the N-SREBP1 (n = 4, p = 0.04) and N-SREBP2 (n = 4, p = 0.03) **(Figure 1B)**. Thus, confirming that

mechanical stress promoted both SREBP1 and SREBP2 activation by inducing its nuclear translocation.

This prompted us to examine the role of SREBPs in the AH drainage outflow pathway and the regulation

of IOP. The expression of SREBP isoforms and SCAP transcripts in HTM cells was identified using reverse-

transcription polymerase chain reaction (RT-PCR) in primary HTM cell cultures **(Supplementary Figure**

1A). Additionally, the SREBP isoforms and SCAP protein were found in HTM and PTM cells. We observed

pro-SREBP1 and pro-SREBP2 at ~ 125 kDa, N-SREBP1 and N-SREBP2 at ~ 68 kDa, and SCAP at ~ 140 kDa
(**Supplementary Figure 1B and 1C**). IF analysis and confocal imaging of SREBP1, 2, and SCAP distribution
in HTM cells and the conventional human AH outflow pathway showed their distinct localization. In HTM
cells, both SREBP1 and 2 showed cytoplasmic and nuclear distribution (**Supplementary Figure 1D**).
Relatively, SREBP1 was located in the more cytoplasmic and perinuclear region around the ER than in
the nucleus. On the other hand, SREBP2 showed a more pronounced baseline nuclear localization than
cytoplasmic and SCAP showed cytoplasmic distribution (**Supplementary Figure 1D**). In the anterior
chamber angle of a normal human eye, the SREBP1, SREBP2, and SCAP immunopositive cells were seen
in the TM-JCT region and the endothelial lining of the SC (**Supplementary Figure 1E**). Interestingly,
SREBP1 and 2, as well as SCAP showed specific distribution in the TM-JCT region in the anterior chamber
angle. Negative control in the presence of secondary antibody alone did not show any staining
(**Supplementary Figure 1F**).

Since we found increased SREBPs activation under mechanical stress, to provide more functional
outcome of SREBPs nuclear localization, we assessed the effects of constitutively activating SREBPs in
HTM cells. For this, we expressed Ad5-N-SREBPs in primary HTM cells *in vitro*. In serum-starved HTM
cells treated with Ad5-N-SREBPs, Ad5-N-SREBP1a and Ad5-N-SREBP1c significantly increased (n = 4, p =
0.01 and n = 4, p = 0.046, respectively) N-SREBP1 transcripts, Ad5-N-SREBP2 significantly increased (n =
4, p = 0.01) N-SREBP2 mRNA expression (**Figure 2A**) in comparison to AdMT control. Similarly, N-SREBP1
protein expression significantly increased under Ad5-N-SREBP1a (n = 4, p = 0.0001) and Ad5-N-SREBP1c
(n = 4, p = 0.03) treatments and N-SREBP2 (n = 4, p = 0.0005) significantly increased under Ad5-N-
SREBP2 treatment (**Figure 2B**). Such increases were also confirmed via IF imaging studies. Increased N-
SREBP1 distribution (red staining) inside the nucleus (**Figure 2C and 2D**) due to Ad5-N-SREBP1a (second-
row third column) and Ad5-N-SREBP1c (third-row third column) compared with AdMT (first-row third
column) and increased N-SREBP2 distribution (red staining) inside the nucleus (**Figure 2E and** 2**F**) due to
Ad5-N-SREBP2 (second-row third column) compared with AdMT (first-row third column). A striking
feature observed in these cells treated with Ad5-N-SREBPs was altered cell shape with increased
lamellipodia and filopodial extensions observed with remarkable changes in the actin cytoskeleton
(**Supplementary Figure 2A**, denoted by black arrows). To check the effects of constitutive activation of
SREBPs on actin polymerization and focal adhesion (FA) formation in HTM, we used phalloidin to stain
the filamentous actin (F-actin) in HTM cells along with FA proteins – vinculin and paxillin. In comparison
to AdMT (**Figure 2C, 2D, 2E, and 2F**, first-row fifth column), Ad5-N-SREBP1a (**Figure 2C and 2D**, second-
row fifth column), Ad5-N-SREBP1c (**Figure 2C and 2D**, third-row fifth column), and Ad5-N-SREBP2 (**Figure
2E and 2F**, second-row fifth column) increased F-actin distribution inside HTM cells (purple staining in
the merged/individually in grayscale) indicating increased actin polymerization. Interestingly, we also
observed that under constitutive activation of SREBPs, there was increased lamellipodia and filopodia
formation with a strong lamellipodial actin network (**Figure 2C,** 2**D, 2E, and 2F**, denoted by yellow
arrows). Increased distribution of vinculin and paxillin in green staining in the merged/individually in
grayscale (denoted by white arrows) at the edge of F-actin was observed in Ad5-N-SREBP1a (**Figure 2C
and 2D**, second-row fourth column), Ad5-N-SREBP1c (**Figure 2C and 2D**, third-row fourth column), and
Ad5-N-SREBP2 (**Figure 2E and 2F**, second-row fourth column) indicating increased FA formation,
contractility inside the cells, and mechanosensing by HTM cells. Thus, augmentation of nuclear SREBPs
in TM cells helps sense mechanical stress by modulating actin polymerization and FA formation.

3.2 Pharmacological inhibition of SREBPs activation lowers IOP by altering the ECM architecture of the TM AH outflow pathway in *ex vivo* porcine perfusion cultures

After deriving a linear relationship between mechanical stress, SREBPs activation, and increased actin polymerization, we hypothesized that loss of SREBPs activity can have a significant effect on IOP. The pharmacological inhibition of SREBPs activity was accomplished using fatostatin (56). To test the inhibitory effect of fatostatin on SREBPs activity in TM, the dose-dependency (0.2, 2, 20 µM) of fatostatin on PTM cells was performed. We found that 20 µM fatostatin for 24 h treatment significantly decreased both N-SREBP1 (n = 4, p = 0.02) and N-SREBP2 (n = 4, p = 0.02) from the total protein extract (**Supplementary Figure 3A and 3B**). The 20 µM fatostatin did not show cytotoxic effects when tested using FDA and PI staining (**Supplementary Figure 3C**). The 20 µM fatostatin treatment for 24 h in PTM cells showed a significant decrease in N-SREBP1 (n = 4, p = 0.01) and N-SREBP2 (n = 4, p = 0.02) protein levels in the nuclear fraction (**Figure 3A**) and same treatment in HTM cells also significantly decreased N-SREBP1 (n = 4, p = 0.03) and N-SREBP2 (n = 4, p = 0.0001) protein expression (**Figure 3B**). The SCAP protein expression did not change (n = 4, p = 0.29) (**Figure 3B**). The IF localization of SREBP1 and SREBP2 distribution after fatostatin treatment demonstrated a marked decrease inside of the nucleus compared to the controls (**Figure 3C and D**). Histograms on the right of **Figure 3C and D** represent the ImageJ-based quantification of nuclear SREBPs fluorescence intensities showing a significant decrease in N-SREBP1 (n = 5, p = 0.02) and N-SREBP2 (n = 5, p = 0.001) fluorescent staining. Thus, providing the proof-of-concept for fatostatin can inhibit SREBPs activity in TM cells.

We then tested the effect of SREBPs inactivation via fatostatin on IOP *ex vivo* using the porcine organ culture system. Perfusion of 20 µM fatostatin significantly decreased IOP by 12 h (n = 5, p = 0.04), and sustained until 22 h (n = 5, p < 0.05) (**Figure 3E**). To determine the effects of fatostatin on the TM outflow pathway structures, we analyzed the histological changes. The H and E staining did not show any marked difference in the AH outflow pathway (**Figure 3F**). Confocal imaging of the TM outflow pathway perfused with fatostatin resulted in an appreciable decrease in the immunostaining of SREBP1 and SREBP2 (**Figure 3G**). ImageJ-based quantification of fluorescence intensity in the immunofluorescence image showed a significant decrease in SREBP1 staining (n = 5, p = 0.04) and SREBP2 staining (n = 5, p = 0.04) (**Figure 3G, right panel**). We provided evidence for changes in the contractile properties of TM cells upon SREBPs activation. On the corollary, we investigated the ECM protein changes upon SREBPs inactivation as the ECM remodeling is affected due to changes in the cellular contractile machinery (63). We found that both COL1A and FN in the TM-JCT region showed a marked decrease compared to the control (**Figure 3H**), which is represented in the histogram for COL1A staining (n = 5, p = 0.0001), and FN staining (n = 5, p = 0.0001) in the TM outflow pathway (**Figure 3H, right panel**). To get more insights into the quantification of protein levels, we performed immunoblotting with the protein extracted from the TM tissue obtained from DMSO and fatostatin-perfused eyes. In agreement with IHC results, we found that compared to DMSO-perfused porcine TM tissue, fatostatin-perfused porcine TM tissue shows significantly decreased expression of N-SREBP1 (n = 4, p = 0.005), N-SREBP2 (n = 4, p = 0.03), COL1A (n = 4, p = 0.01), and FN (n = 4, p = 0.03) (**Figure 3I**). Thereby, we demonstrate that the direct inhibition of SREBPs activity significantly lowers the IOP by decreasing ECM in the TM outflow pathway.

3.3 Molecular inactivation of SREBPs by knocking down SCAP in *SCAP^{f/f}* mice lowers IOP *in vivo*

We checked the effects of SREBPs inactivation on IOP regulation in *SCAP^{f/f}* mice *in vivo*. Conditional knockdown of SCAP was achieved using intravitreal injection of Ad5.CMV.iCre-eGFP in *SCAP^{f/f}* mice inducing the loss of SCAP in the TM tissue due to Cre/lox recombination. The cohorts were divided into

four groups - three injection groups: Ad5.CMV.iCre-eGFP, Ad5.CMV.eGFP, and saline and one uninjected
or the untouched group. Among the injected groups, Ad5.CMV.eGFP injection served as sham virus
control and injection of saline served as the sham injection control. The IOP was measured over time in
anesthetized *SCAP^{f/f}* mice before injection, and every 10 days after injection until 50 days post-injection.
Data presented here show the mean IOP ± SEM in mmHg. The Ad5.CMV.iCre-eGFP injection group
displayed a significant IOP lowering (12.97 ± 0.71, n = 10) compared to Ad5.CMV.eGFP injection group
(21.74 ± 1.47, n = 9, p = 0.0001), saline injection group (20.5 ± 1.00, n = 10, p = 0.0001), and the
untouched group (19.67 ± 0.83, n = 9, p = 0.0001) (**Figure 4A**) 10 days after injection. The IOP decrease
in Ad5.CMV.iCre-eGFP injection group compared to the control groups sustained up to 30 days post-
injection (n = 10, p < 0.05). At post-injection days 40 and 50, IOP in Ad5.CMV.iCre-eGFP injection group
was significantly decreased (n = 10, p < 0.05) compared to saline injection and Ad5.CMV.eGFP injection
groups. The IOP was also lower than the untouched group, but not statistically significant (**Figure 4A**).
We also compared the IOP changes within each group to their baseline normal IOP (before
injection/time point 0 days), and the result shows that there was no significant change in IOP levels in
untouched, saline injection, and Ad5.CMV.eGFP injection groups. However, within Ad5.CMV.iCre-eGFP
injection group, the IOP was significantly decreased 10 days post-injection and sustained till 50 days
post-injection (n = 10, p < 0.05) (**Figure 4B**). Upon analyzing percentage changes in IOP (delta IOP %) in
each group at each time point, Ad5.CMV.iCre-eGFP injection group showed a decrease of 40.38%
compared to the baseline normal (before injection/time point 0 days) with an average decrease of
27.85% (**Figure 4C**). The delta IOP % in the control groups were less than 13% changes of baseline
normal IOP. (**Figure 4C**). We evaluated IOP changes after SREBPs knock down in adult and old mice to
check if aging in combination with SREBPs knockdown influenced IOP. We found IOP was consistently
and significantly decreased in Ad5.CMV.iCre-eGFP group compared to Ad5.CMV.eGFP group (n = 5, p <
0.05), and it was also lower than the untouched group and saline injection group in the old and adult
cohorts (**Supplementary Figure 4A and 4D, respectively**). In both old and adult cohorts, there was no
significant change in IOP levels in all sham control groups, but IOP was significantly decreased 10 days
post-injection in Ad5.CMV.iCre-eGFP injection group and sustained till 50 days post-injection (n = 5, p <
0.05) (**Supplementary Figure 4B and 4E, respectively**). The IOP in Ad5.CMV.iCre-eGFP injection groups
showed a decrease of as much as 42.72 % and 43.14 % of baseline normal IOP in old and adult mice,
respectively (**Supplementary Figure 4C and 4F, respectively**). All sham control groups showed less than
17 % changes of baseline normal IOP in both old and adult mice (**Supplementary Figure 4C and 4F**). At
the end of the longitudinal IOP measurement study, the mice eyes were enucleated and processed for
histological and immunohistochemical analysis. The H and E staining showed normal open iridocorneal
angles and no marked differences in TM morphology in all the groups (**Figure 4D**). Immunostaining in
Ad5.CMV.iCre-eGFP injection group compared to all control groups for - SCAP showed a marked
decrease in TM (**Figure 4E, green arrow in the inset**), SREBP1 and SREBP2 showed decreased
distribution in the nuclear regions in TM (**Figure 4F and 4G, green arrow in the inset**). This data proves
that Ad5.CMV.iCre-eGFP injection selectively knockdown SCAP in TM and further inhibited SREBPs
activation in TM consistent with decreased IOP.

3.4 Cholesterol metabolism is greatly affected under SREBP inactivation

Since SREBPs are master regulators of lipid biosynthesis, as a proof-of-concept study, we looked at
changes in lipid metabolism in HTM cells under fatostatin treatment. HTM cells treated with 20 µM
fatostatin for 24 h significantly decreased SREBPs responsive mRNA expression involved in - 1) genes

related to cholesterol biosynthesis: low-density lipoprotein receptor (LDLR) (n = 3, p = 0.05),
hydroxymethylglutaryl-CoA synthase 1 (HMGCS1) (n = 3, p = 0.10), 3-Hydroxy-3-Methylglutaryl-CoA
Reductase (HMGCR) (n = 3, p = 0.049), SREBP2 (n = 3, p = 0.0008); 2) genes involved in fatty acid
biosynthesis: acetyl-CoA carboxylase (ACC) (n = 3, p = 0.0002), fatty acid synthase (FASN) (n = 3, p =
0.0055), SREBP1 (n = 3, p = 0.0001), peroxisome proliferator-activated receptor gamma (PPARG) (n = 3, p
= 0.01), acetyl-CoA synthetase (n = 3, p = 0.0002); and 3) gene regulating triglyceride and phospholipid
biosynthesis: glycerol-3-phosphate acyltransferase (GPAT) (n = 3, p = 0.0017) (**Figure 5A**). We also used
immunoblotting to check protein expression levels of FAS, HMGCR, and ACC under control and
fatostatin treatment. The immunoblotting results show that fatostatin significantly decreased FAS (n = 4,
p = 0.049), HMGCR (n = 4, p = 0.02), and ACC (n = 4, p = 0.049) expression (**Figure 5B**). To further confirm
the lipid changes under fatostatin treatment, we treated HTM cells with 20 µM fatostatin for 24 h, and
after treatment, we counted the number of cells under each treatment and measured normalized
phospholipid/cell, normalized cholesterol/cell, and normalized triglyceride/cell. The results show that
phospholipid (n = 4, p = 0.0001) (**Figure 5C**), cholesterol (n = 4, p = 0.04) (**Figure 5D**), and triglyceride (n =
4, p = 0.04) (**Figure 5E**) were significantly reduced under fatostatin treatment. Interestingly the
magnitude of the phospholipid decrease was the highest (~95%). All these results indicate that SREBPs
inactivation using fatostatin can lower the total lipid biosynthesis in HTM cells by decreasing the
expression of SREBPs responsive genes.
432
3.5 Induction of actin depolymerization machinery and decreasing ECM-based interactions in TM
defines the cellular basis of SREBPs inhibition on IOP lowering
Next, we checked if SREBPs inactivation by fatostatin modulated actin and FA proteins. We probed for
filamentous actin (F-actin) using phalloidin labeling in HTM cells along with FA proteins – vinculin and
paxillin. As seen in **Figures 6A and 6B**, compared to DMSO control, 24 h of 20 µM fatostatin treatment
on HTM cells demonstrated a marked decrease in the F-actin stress fibers (green staining in
merged/grayscale) and sparse distribution of vinculin and paxillin at the edges of F-actin fibers with
greater cytoplasmic accumulation (red staining in the merged/grayscale, indicated by white arrows). To
further confirm this result, we extracted F-actin and globular actin (G-actin) from control and fatostatin-
treated HTM cells separately and checked the changes in F-actin/G-actin ratio in each condition. **Figure**
6C shows that fatostatin significantly decreased F-actin/G-actin ratio (n = 4, p = 0.04), implying the
decreased actin polymerization under fatostatin treatment. Since we also observed decreased FA
formation in IF. We next checked the expression of proteins involved in FA formation, F-actin
polymerization, and their stabilization. Fascin is a major F-actin bundling protein and regulates the
maintenance and stability of parallel bundles of F-actin in various cell types including TM (64),(20).
Immunoblotting showed that fatostatin significantly decreased fascin-1 expression (n = 4, p = 0.03)
(**Figure 6D**). We did not see any change in vinculin expression (n = 4, p = 0.61) (**Figure 6D**). Thus
suggesting that fatostatin changes vinculin distribution at the edge of F-actin fibers, but not its
expression. We found that fatostatin significantly decreased LIM kinase 1 (LIMK1) expression (n = 4, p =
0.04) (**Figure 6D**), an actin polymerization and stabilization protein(65). Looking at cofilin, we found no
significant change in the levels of total-cofilin (T-cofilin) (n = 4, p = 0.97) and phosphorylated cofilin at
serine 3 (P-cofilin) (n = 4, p = 0.05), but the ratio of P-cofilin/T-cofilin was significantly decreased in
fatostatin treatment (n = 4, p = 0.01) (**Figure 6E**). Therefore, providing evidence for weakening the
LIMK1-cofilin-driven actin polymerization. Moreover, we also found that fatostatin treatment increased
total-paxillin (T-paxillin) levels (n = 4, p = 0.07), and significantly decreased its tyrosine 118
phosphorylated form (P-paxillin) (n = 4, p = 0.03). The ratio of P-paxillin/T-paxillin was significantly
decreased in fatostatin treatment (n = 4, p = 0.01) (**Figure 6F**), indicating decreased cell adhesiveness
and cellular contractility.

Alterations in actin and focal adhesions can have a direct impact on ECM adhesion (66). Consequently,
we evaluated the effect of SREBPs activity on ECM protein expression and distribution in TM, we treated
HTM cells with 20 μM fatostatin or vehicle DMSO for 24 h. Interestingly, we found that fatostatin
significantly decreased all ECM mRNA expression, including FN (n = 4, p = 0.0001), COL1α (n = 4, p =
0.0001), COL6α (n = 4, p = 0.0001), COL4α (n = 4, p = 0.0001), tenascin C (TNC) (n = 4, p = 0.0001),
transforming growth factor β2 (TGFβ2) (n = 4, p = 0.0001), and sarcospan (SSPN) (n = 4, p = 0.0001)
(**Figure 7A**). Consequently, upon fatostatin treatment, the whole cell lysate (CL) protein expression
analysis showed a significant reduction of FN (n = 4, p = 0.01) and COL1A (n = 4, p = 0.001) (**Figure 7B**). In
addition, secretory ECM levels in conditioned media (CM) demonstrated a significant decrease in FN (n =
4, p = 0.02) and COL1A (n = 4, p = 0.001) expression in the fatostatin treatment compared to the control
(**Figure 7C**). These results were confirmed using IF. Under fatostatin treatment, FN (**Figure 7D**) and
COL1A (**Figure 7E**) immunostaining were comprehensively decreased. Altogether these data indicate
that SREBPs inactivation significantly impacts actin polymerization and FA formation. Additionally lowers
the intracellular and secreted ECM levels, as well as their distribution. Thus explaining the contribution
of the SREBPs activity on the maintenance of IOP by balancing the actin cytoskeletal and ECM
remodeling in the TM outflow pathway.

4. Discussion:

This is the first study to provide the initial evidence and elucidate the effects of SREBPs on lipid
metabolism, cell contractility, ECM remodeling in TM, and IOP. In this study, we report that SREBPs and
SCAP are highly expressed in the TM-JCT region compared to other components in the ocular anterior
segment. We found that mechanical stress on TM including acute elevation of IOP significantly increased
SREBPs activation. Augmenting SREBP activation induced contractility indicating a perpetual
feedforward signaling from mechanical stress sensing by SREBPs resulting in induction of contractile
force in the TM. Conversely inhibiting SREBPs activation using fatostatin *ex vivo* or in the *SCAP^{f/f}* mice *in*
vivo significantly lowered the IOP. We hypothesize that such IOP decrease due to SREBPs inactivation
lowers the TM contractility and ECM-based stiffness potentially by downregulating lipogenesis. SREBPs
inactivation significantly reduced the expression of their responsive lipid genes and enzymes related to
lipogenic pathways in HTM cells. Here, we showed that SREBPs inactivation mitigated actin
polymerization and actin bundling by lowering the levels of fascin-1, LIMK1, and phosphorylation of
paxillin and cofilin. This study strengthens the correlation between lipid changes and tissue stiffness and
reinforces the significance and role of actin-cell adhesive interactions in the regulation of IOP
homeostasis. Thus, suggesting that the SREBP-SCAP complex could play an important role in the
lipogenesis in the TM tissue.

Here, we inactivated the SREBP-SCAP pathway using two different approaches to decipher the role of
SREBPs in the regulation of IOP. *Ex vivo* perfusion studies using pharmacological inhibition of SREBPs by
fatostatin significantly lowered IOP. A significant IOP lowering was achieved only by 12h, which we
believe is due to the low concentration (20 μM) used for perfusion. We also predict that the effect of
inhibiting a transcription factor will require a longer time to produce the effect since it will require
regulation at the gene expression level first followed by changes in protein/enzyme levels and finally
effecting physiological changes. On the other hand, *in vivo* studies in *SCAP^{f/f}* mice injected with AdCre
confirmed that the molecular inactivation of SREBPs by knocking down SCAP significantly lowered IOP by
10 days post-injection. In our recent study, we observed that TM cells under mechanical stress showed
increased SREBP2 protein levels (38). The SREBP2 is primarily known to contribute to cholesterol

production. Based on the levels of cholesteryl ester levels under mechanical stretch, we hypothesized
that cholesterol could be used for rapid membrane remodeling and remaining stored as cholesteryl
esters (38). As an extension, here we provide the proof that acute elevation of ocular pressure using *ex*
vivo porcine anterior segment perfusion cultures induced SREBPs activation in TM. Thereby confirming
that SREBPs activation is correlated with mechanical stress on TM. The TM adapts to increased
mechanical stress by modifying the actin cytoskeleton and the ECM (38, 62). Upon inducing SREBPs
activation, we found that there were increased actin stress fibers indicating a strong connection
between SREBPs activation and actin polymerization and an impact on IOP homeostasis. Therefore, we
speculated that inhibiting SREBPs activation in TM will modulate IOP. Using both *ex vivo* and *in vivo*
studies, we provide strong evidence that attenuating the SREBP-SCAP pathway results in significant IOP
lowering. *Ex vivo* evidence indicated a significant loss in the ECM components like collagen and
fibronectin in the outflow pathway which concurred in the *in vitro* experiments in HTM cells. *In vitro*
analysis provided further proof that the inactivation of SREBPs decreased actin fibers and retraction of
FAs to the cytosol with very little found at the edges of the cells. The FAs aid in linking the intracellular
cytoskeleton to the extracellular ECM (67). A compromise in vinculin and paxillin localization at the edge
of F-actin fibers dictates the dissolution of FAs and decreased cell-ECM connections.

It has been characterized that the FA vinculin but not paxillin directly interacts with actin (68, 69).
Additional mechanistic evidence in our study confirms that SREBPs inactivation regulates vinculin
distribution but not expression confirming the requirement of SREBPs activity in modulating the
recruitment of vinculin to the FA. Also, a decrease in paxillin phosphorylation suggests a decreased
inside-out signaling towards integrin binding to ECM and recruitment of FA (70, 71). Inside the cells, we
identified that loss of SREBPs activation lowered the actin-bundling protein fascin-1 (20, 64, 72). The
decreased fascin-1 expression under fatostatin treatment can result in a slower rate of actin
polymerization with lesser filopodia formation (73, 74). Within this context, the decreased LIMK1
resulting in the loss of phosphorylation of cofilin-1 at serine 3 can induce F-actin destabilization and this
can prevent contraction and attenuate hypertension (65, 75). Thus, we propose that the loss of cell
adhesive interaction upon SREBPs inactivation is an important mechanism in lowering the IOP.
Interestingly, we found that there was a strong negative regulation of most of the ECM gene expression
upon fatostatin treatment in HTM cells. Though it is not yet clear why SREBPs does not directly regulate
the transcription of ECM genes, we believe this to be a result of negative feedback outside-in signaling
towards transcribing ECM genes due to the disengagement of ECM.

The major function of SREBPs is to regulate lipogenesis. In line, upon inactivation of the SREBP-SCAP
pathway, we found a significant and remarkable decrease in total phospholipids in TM. Additionally,
both cholesterol and triglycerides were significantly decreased. The current understanding of the
correlation between the regulation of lipid levels in the TM outflow pathway with IOP changes is
unclear. Lipids levels are known to be varied in aqueous humor from ocular hypertensives (76) (28, 77)
and can participate in the pathogenesis of elevated IOP – the bioactive lysophospholipids (lipid growth
factors) (29, 78-81), and cholesterol (30-32). Yet the causality has not been completely attributed to
increased lipids on IOP elevation. Based on recent epidemiologic studies, there is an association
between cholesterol and glaucoma, although the findings have not been conclusive if hyperlipidemia is a
cause or an effect of the disease. Some studies found that hyperlipidemia, especially high cholesterol
levels, is significantly associated with an increased risk of glaucoma and increased IOP (31, 32, 82) (83)
and others show the opposite (84) (85). Interestingly, even though it is not clear how cholesterol is

related to the glaucoma risk, there is mounting evidence showing that HMG-CoA reductase inhibitor
statin can decrease POAG risk by mitigating disease progression (35, 86, 87). The proposed mechanism
for statins to decrease IOP is by inhibiting the isoprenylation of Rho GTPase thus resulting in decreased
actomyosin contractile activity and ECM synthesis/assembly (36, 88-90). All these studies imply that
there is an association between lipids, ocular hypertension, and POAG. However, the contributions of
lipid metabolism in TM for the regulation of TM biomechanics and IOP are blurred (91). We have
recently identified that TM cholesterol plays a significant role in the maintenance of the polymerized
state of actin, focal adhesion recruitment, and TM membrane tension (*Unpublished study*). Thus,
suggesting a strong role of lipids including cholesterol in regulating TM actin-adhesion complex-ECM
modulated via the SREBP-SCAP pathway to efficiently maintain the IOP homeostasis. These findings
indicate that SREBPs and their downstream lipid biosynthesis pathways have an essential role in TM
functions and contribute significantly to IOP regulation. Moreover, multiple rate-limiting enzymes in
lipid biogenic pathways like FAS (49, 92), acetyl-CoA carboxylase(ACC) (93, 94), 3-hydroxy-3-
methylglutaryl coenzyme A (HMG-CoA) reductase (HMGCR), and SREBPs are implicated in increasing
ECM and fibrosis in various tissue types (95-97). ACC catalyzes the carboxylation of acetyl-CoA into
malonyl-CoA. This reaction is the first and the rate-limiting step in the biosynthesis of fatty acids.
Significantly, Acetyl-CoA metabolism is implicated in POAG (98). Further investigation on the role of
these rate limiting enzymes in TM biomechanics is essential to decode their function in IOP regulation.
We predict that induction of SREBPs activation can induce IOP elevation by modifying the contractile
properties of the TM, which warrants further studies. Though out of scope for this manuscript, we
predict that SREBPs can modulate ECM remodeling via lipid-independent pathways. SREBP1 has been
shown to bind to the promoter region of TGFβ (99), a major pro-fibrotic factor. Also, SREBP1 can act as a
cell surface retention factor for the TGFβ receptor, TβRI, by preventing its secretion in exosomes or
directly changing COL VI transcription (100) (101). Thus, pointing to the idea that SREBPs activation can
potentially increase ECM production and deposition and IOP elevation in both lipid-dependent and -
independent pathways.

5. Conclusion

574 In summary, this study presents a novel way of lowering IOP by inactivating SREBPs through decreasing
575 lipogenesis, actin-based tension, and attenuating ECM accumulation in TM. Thus, demonstrating the
576 important role of SREBPs in the regulation of TM contractility and stiffness, and IOP regulation. Further
577 modalities of SREBPs regulating the actin cytoskeleton and ECM in TM will aid in identifying better
578 targets for IOP lowering and glaucoma therapy.

6. Conflict of Interest

580 The authors declare no conflict of interest. The funders had no role in the design of the study; in the
581 collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to
582 publish the results.

7. Funding

584 This project was supported by the National Institutes of Health/ National Eye Institute (R01EY029320)
585 (PPP), Glick Research Endowment Funds (PPP), Cohen AMD Research Pilot Grant (PPP), RPB
586 Departmental Pilot Grant (PPP), Sigma Xi Grant-in-aid (TW) and grant from Research to Prevent
587 Blindness to IU.

10. Acknowledgments

We would like to acknowledge Dr. Nuria Morral, Dr. Benjamin Perrin, Dr. Gary Landreth, and Dr. Tim Corson from IUSM for their useful discussions on the project.

11. References

1.	R. N. Weinreb, P. T. Khaw, Primary open-angle glaucoma. *Lancet* **363**, 1711-1720 (2004).
2.	W. M. Grant, Clinical tonography. *Trans Am Acad Ophthalmol Otolaryngol* **55**, 774-781 (1951).
3.	W. M. Grant, Experimental aqueous perfusion in enucleated human eyes. *Arch Ophthalmol* **69**, 783-801 (1963).
4.	N. A. Sharif, Therapeutic Drugs and Devices for Tackling Ocular Hypertension and Glaucoma, and Need for Neuroprotection and Cytoprotective Therapies. *Frontiers in pharmacology* **12**, 729249 (2021).
5.	Y. Shiose, Intraocular pressure: new perspectives. *Surv Ophthalmol* **34**, 413-435 (1990).
6.	L. Bonomi *et al.*, Prevalence of glaucoma and intraocular pressure distribution in a defined population. The Egna-Neumarkt Study. *Ophthalmology* **105**, 209-215 (1998).
7.	I. Dielemans *et al.*, Primary open-angle glaucoma, intraocular pressure, and systemic blood pressure in the general elderly population. The Rotterdam Study. *Ophthalmology* **102**, 54-60 (1995).
8.	H. A. Quigley, Open-angle glaucoma. *The New England journal of medicine* **328**, 1097-1106 (1993).
9.	J. B. Saaddine, K. M. Narayan, F. Vinicor, Vision loss: a public health problem? *Ophthalmology* **110**, 253-254 (2003).
10.	Y. C. Tham *et al.*, Global prevalence of glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-analysis. *Ophthalmology* **121**, 2081-2090 (2014).
11.	H. A. Quigley, A. T. Broman, The number of people with glaucoma worldwide in 2010 and 2020. *British journal of ophthalmology* **90**, 262-267 (2006).
12.	M. A. Kass *et al.*, The Ocular Hypertension Treatment Study: a randomized trial determines that topical ocular hypotensive medication delays or prevents the onset of primary open-angle glaucoma. *Arch Ophthalmol* **120**, 701-713; discussion 829-730 (2002).
13.	A. F. Clark, T. Yorio, Ophthalmic drug discovery. *Nat Rev Drug Discov* **2**, 448-459 (2003).
14.	A. J. Lee, I. Goldberg, Emerging drugs for ocular hypertension. *Expert Opin Emerg Drugs* **16**, 137-161 (2011).
15.	L. Crawley, S. M. Zamir, M. F. Cordeiro, L. Guo, Clinical options for the reduction of elevated intraocular pressure. *Ophthalmol Eye Dis* **4**, 43-64 (2012).
16.	D. K. Sunderland, A. Sapra, "Physiology, Aqueous Humor Circulation" in StatPearls. (Treasure Island (FL), 2022).
17.	W. D. Stamer, The cell and molecular biology of glaucoma: mechanisms in the conventional outflow pathway. *Invest Ophthalmol Vis Sci* **53**, 2470-2472 (2012).
18.	D. W. Abu-Hassan, T. S. Acott, M. J. Kelley, The Trabecular Meshwork: A Basic Review of Form and Function. *J Ocul Biol* **2** (2014).
19.	T. Carreon, E. van der Merwe, R. L. Fellman, M. Johnstone, S. K. Bhattacharya, Aqueous outflow - A continuum from trabecular meshwork to episcleral veins. *Prog Retin Eye Res* **57**, 108-133 (2017).
20.	A. Soundararajan, S. A. Ghag, S. S. Vuda, T. Wang, P. P. Pattabiraman, Cathepsin K Regulates Intraocular Pressure by Modulating Extracellular Matrix Remodeling and Actin-Bundling in the Trabecular Meshwork Outflow Pathway. *Cells* **10**, 2864 (2021).

21. P. P. Pattabiraman, C. B. Toris, The exit strategy: Pharmacological modulation of extracellular
matrix production and deposition for better aqueous humor drainage. *Eur J Pharmacol* **787**, 32-
42 (2016).
22. P. P. Pattabiraman *et al.*, RhoA GTPase-induced ocular hypertension in a rodent model is
associated with increased fibrogenic activity in the trabecular meshwork. *The American journal*
of pathology **185**, 496-512 (2015).
23. K. Wang, A. T. Read, T. Sulchek, C. R. Ethier, Trabecular meshwork stiffness in glaucoma. *Exp Eye*
Res **158**, 3-12 (2017).
24. J. A. Vranka, M. J. Kelley, T. S. Acott, K. E. Keller, Extracellular matrix in the trabecular meshwork:
intraocular pressure regulation and dysregulation in glaucoma. *Exp Eye Res* **133**, 112-125 (2015).
25. E. R. Tamm, B. M. Braunger, R. Fuchshofer, Intraocular Pressure and the Mechanisms Involved in
Resistance of the Aqueous Humor Flow in the Trabecular Meshwork Outflow Pathways. *Prog*
Mol Biol Transl Sci **134**, 301-314 (2015).
26. J. D. Humphrey, E. R. Dufresne, M. A. Schwartz, Mechanotransduction and extracellular matrix
homeostasis. *Nature reviews. Molecular cell biology* **15**, 802-812 (2014).
27. N. V. Azbukina *et al.*, Targeted Lipidomic Analysis of Aqueous Humor Reveals Signaling Lipid-
Mediated Pathways in Primary Open-Angle Glaucoma. *Biology (Basel)* **10**, 658 (2021).
28. G. Edwards *et al.*, Endogenous ocular lipids as potential modulators of intraocular pressure. *J*
Cell Mol Med **24**, 3856-3900 (2020).
29. P. V. Rao, Bioactive lysophospholipids: role in regulation of aqueous humor outflow and
intraocular pressure in the context of pathobiology and therapy of glaucoma. *J Ocul Pharmacol*
Ther **30**, 181-190 (2014).
30. K. M. Madjedi *et al.*, The Association between Serum Lipids and Intraocular Pressure in 2 Large
United Kingdom Cohorts. *Ophthalmology* **129**, 986-996 (2022).
31. L. Posch-Pertl *et al.*, Cholesterol and glaucoma: a systematic review and meta-analysis. *Acta*
Ophthalmol **100**, 148-158 (2022).
32. S. Wang, X. Bao, Hyperlipidemia, Blood Lipid Level, and the Risk of Glaucoma: A Meta-Analysis.
Invest Ophthalmol Vis Sci **60**, 1028-1043 (2019).
33. J. H. Kang *et al.*, Association of Statin Use and High Serum Cholesterol Levels With Risk of
Primary Open-Angle Glaucoma. *JAMA ophthalmology* **137**, 756-765 (2019).
34. M. W. Marcus *et al.*, Cholesterol-lowering drugs and incident open-angle glaucoma: a
population-based cohort study. *PloS one* **7**, e29724 (2012).
35. J. D. Stein *et al.*, The relationship between statin use and open-angle glaucoma. *Ophthalmology*
119, 2074-2081 (2012).
36. J. Song, P. F. Deng, S. S. Stinnett, D. L. Epstein, P. V. Rao, Effects of cholesterol-lowering statins
on the aqueous humor outflow pathway. *Invest Ophthalmol Vis Sci* **46**, 2424-2432 (2005).
37. C. L. Von Zee, M. P. Richards, P. Bu, J. I. Perlman, E. B. Stubbs, Jr., Increased RhoA and RhoB
protein accumulation in cultured human trabecular meshwork cells by lovastatin. *Invest*
Ophthalmol Vis Sci **50**, 2816-2823 (2009).
38. A. Soundararajan *et al.*, Multiomics analysis reveals the mechanical stress-dependent changes in
trabecular meshwork cytoskeletal-extracellular matrix interactions. *Front Cell Dev Biol* **10**,
874828 (2022).
39. H. Shimano, R. Sato, SREBP-regulated lipid metabolism: convergent physiology - divergent
pathophysiology. *Nat Rev Endocrinol* **13**, 710-730 (2017).
40. H. Shimano, Sterol regulatory element-binding proteins (SREBPs): transcriptional regulators of
lipid synthetic genes. *Prog Lipid Res* **40**, 439-452 (2001).
41. D. Eberle, B. Hegarty, P. Bossard, P. Ferre, F. Foufelle, SREBP transcription factors: master
regulators of lipid homeostasis. *Biochimie* **86**, 839-848 (2004).

42. J. D. Horton, J. L. Goldstein, M. S. Brown, SREBPs: activators of the complete program of
cholesterol and fatty acid synthesis in the liver. *J Clin Invest* **109**, 1125-1131 (2002).
43. J. L. Goldstein, R. A. DeBose-Boyd, M. S. Brown, Protein sensors for membrane sterols. *Cell* **124**,
35-46 (2006).
44. D. Dorotea, D. Koya, H. Ha, Recent Insights Into SREBP as a Direct Mediator of Kidney Fibrosis via
Lipid-Independent Pathways. *Front Pharmacol* **11**, 265 (2020).
45. M. Matsuda *et al.*, SREBP cleavage-activating protein (SCAP) is required for increased lipid
synthesis in liver induced by cholesterol deprivation and insulin elevation. *Genes Dev* **15**, 1206-
1216 (2001).
46. J. Sakai, A. Nohturfft, J. L. Goldstein, M. S. Brown, Cleavage of sterol regulatory element-binding
proteins (SREBPs) at site-1 requires interaction with SREBP cleavage-activating protein. Evidence
from in vivo competition studies. *J Biol Chem* **273**, 5785-5793 (1998).
47. S. Daemen, M. Kutmon, C. T. Evelo, A pathway approach to investigate the function and
regulation of SREBPs. *Genes Nutr* **8**, 289-300 (2013).
48. R. Bertolio *et al.*, Sterol regulatory element binding protein 1 couples mechanical cues and lipid
metabolism. *Nat Commun* **10**, 1326 (2019).
49. X. Zhao, J. Y. Y. Kwan, K. Yip, P. P. Liu, F. F. Liu, Targeting metabolic dysregulation for fibrosis
therapy. *Nat Rev Drug Discov* **19**, 57-75 (2020).
50. G. Proctor *et al.*, Regulation of renal fatty acid and cholesterol metabolism, inflammation, and
fibrosis in Akita and OVE26 mice with type 1 diabetes. *Diabetes* **55**, 2502-2509 (2006).
51. J. H. Lee *et al.*, SREBP-1a-stimulated lipid synthesis is required for macrophage phagocytosis
downstream of TLR4-directed mTORC1. *Proc Natl Acad Sci U S A* **115**, E12228-e12234 (2018).
52. G. Sorrentino *et al.*, Metabolic control of YAP and TAZ by the mevalonate pathway. *Nat Cell Biol*
16, 357-366 (2014).
53. K. E. Keller *et al.*, Consensus recommendations for trabecular meshwork cell isolation,
characterization and culture. *Exp Eye Res* **171**, 164-173 (2018).
54. A. Soundararajan, T. Wang, S. A. Ghag, M. H. Kang, P. P. Pattabiraman, Novel insight into the
role of clusterin on intraocular pressure regulation by modifying actin polymerization and
extracellular matrix remodeling in the trabecular meshwork. *J Cell Physiol* **237**, 3012-3029
(2022).
55. X. Cheng, J. Li, D. Guo, SCAP/SREBPs are Central Players in Lipid Metabolism and Novel
Metabolic Targets in Cancer Therapy. *Curr Top Med Chem* **18**, 484-493 (2018).
56. S. Kamisuki *et al.*, A small molecule that blocks fat synthesis by inhibiting the activation of
SREBP. *Chem Biol* **16**, 882-892 (2009).
57. S. H. Lee, J. H. Lee, S. S. Im, The cellular function of SCAP in metabolic signaling. *Exp Mol Med* **52**,
724-729 (2020).
58. L. Jiajia *et al.*, Assessment of Neuronal Viability Using Fluorescein Diacetate-Propidium Iodide
Double Staining in Cerebellar Granule Neuron Culture. *J Vis Exp* 10.3791/55442 (2017).
59. M. Matsuda *et al.*, SREBP cleavage-activating protein (SCAP) is required for increased lipid
synthesis in liver induced by cholesterol deprivation and insulin elevation. *Genes & development*
15, 1206-1216 (2001).
60. Y. Kidani *et al.*, Sterol regulatory element–binding proteins are essential for the metabolic
programming of effector T cells and adaptive immunity. *Nature Immunology* **14**, 489-499 (2013).
61. K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408 (2001).
62. P. P. Pattabiraman, T. Inoue, P. V. Rao, Elevated intraocular pressure induces Rho GTPase
mediated contractile signaling in the trabecular meshwork. *Exp Eye Res* **136**, 29-33 (2015).

63. P. P. Pattabiraman, P. V. Rao, Mechanistic basis of Rho GTPase-induced extracellular matrix
synthesis in trabecular meshwork cells. *Am J Physiol Cell Physiol* **298**, C749-763 (2010).

64. J. Zanet *et al.*, Fascin promotes filopodia formation independent of its role in actin bundling.
Journal of Cell Biology **197**, 477-486 (2012).

65. O. Bernard, Lim kinases, regulators of actin dynamics. *Int J Biochem Cell Biol* **39**, 1071-1076
(2007).

66. C. Wu, Focal adhesion: a focal point in current cell biology and molecular medicine. *Cell Adh*
Migr **1**, 13-18 (2007).

67. B. Geiger, A. Bershadsky, R. Pankov, K. M. Yamada, Transmembrane crosstalk between the
extracellular matrix--cytoskeleton crosstalk. *Nat Rev Mol Cell Biol* **2**, 793-805 (2001).

68. J. D. Humphries *et al.*, Vinculin controls focal adhesion formation by direct interactions with talin
and actin. *The Journal of cell biology* **179**, 1043-1057 (2007).

69. M. Reinhard, M. Rüdiger, B. M. Jockusch, U. Walter, VASP interaction with vinculin: a recurring
theme of interactions with proline-rich motifs. *FEBS Lett* **399**, 103-107 (1996).

70. S. L. Bellis, J. T. Miller, C. E. Turner, Characterization of tyrosine phosphorylation of paxillin in
vitro by focal adhesion kinase. *The Journal of biological chemistry* **270**, 17437-17441 (1995).

71. A. M. López-Colomé, I. Lee-Rivera, R. Benavides-Hidalgo, E. López, Paxillin: a crossroad in
pathological cell migration. *J Hematol Oncol* **10**, 50 (2017).

72. L. M. Machesky, A. Li, Fascin: Invasive filopodia promoting metastasis. *Commun Integr Biol* **3**,
263-270 (2010).

73. R. Jaiswal *et al.*, The Formin Daam1 and Fascin Directly Collaborate to Promote Filopodia
Formation. *Current Biology* **23**, 1373-1379 (2013).

74. D. Vignjevic *et al.*, Role of fascin in filopodial protrusion. *Journal of Cell Biology* **174**, 863-875
(2006).

75. M. Morales-Quinones *et al.*, LIMK (LIM Kinase) Inhibition Prevents Vasoconstriction- and
Hypertension-Induced Arterial Stiffening and Remodeling. *Hypertension* **76**, 393-403 (2020).

76. G. Edwards, K. Aribindi, Y. Guerra, S. K. Bhattacharya, Sphingolipids and ceramides of mouse
aqueous humor: Comparative profiles from normotensive and hypertensive DBA/2J mice.
Biochimie **105**, 99-109 (2014).

77. H. Wang, G. Edwards, C. Garzon, C. Piqueras, S. K. Bhattacharya, Aqueous humor phospholipids
of DBA/2J and DBA/2J-Gpnmb(+)/SjJ mice. *Biochimie* **113**, 59-68 (2015).

78. P. S. Mettu *et al.*, Role of lysophospholipid growth factors in the modulation of aqueous humor
outflow facility. *Invest Ophthalmol Vis Sci* **45**, 2263-2271 (2004).

79. L. T. Y. Ho *et al.*, Role of the autotaxin-lysophosphatidic acid axis in glaucoma, aqueous humor
drainage and fibrogenic activity. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of*
Disease **1866**, 165560 (2020).

80. M. Honjo *et al.*, Autotaxin–Lysophosphatidic Acid Pathway in Intraocular Pressure Regulation
and Glaucoma Subtypes. *Investigative Ophthalmology & Visual Science* **59**, 693-701 (2018).

81. P. Iyer *et al.*, Autotaxin-lysophosphatidic acid axis is a novel molecular target for lowering
intraocular pressure. *PloS one* **7**, e42627 (2012).

82. H. C. Lin, C. W. Chien, C. C. Hu, J. D. Ho, Comparison of comorbid conditions between open-angle
glaucoma patients and a control cohort: a case-control study. *Ophthalmology* **117**, 2088-2095
(2010).

83. S. Pavljasević, M. Asćerić, Primary open-angle glaucoma and serum lipids. *Bosn J Basic Med Sci* **9**,
85-88 (2009).

84. P. A. Newman-Casey, N. Talwar, B. Nan, D. C. Musch, J. D. Stein, The relationship between
components of metabolic syndrome and open-angle glaucoma. *Ophthalmology* **118**, 1318-1326
(2011).

85.   W. C. Stewart, C. Sine, S. Sutherland, J. A. Stewart, Total cholesterol and high-density lipoprotein
levels as risk factors for increased intraocular pressure. *Am J Ophthalmol* **122**, 575-577 (1996).
86.   P. McCann, R. E. Hogg, R. Fallis, A. Azuara-Blanco, The Effect of Statins on Intraocular Pressure
and on the Incidence and Progression of Glaucoma: A Systematic Review and Meta-Analysis.
Investigative Ophthalmology & Visual Science **57**, 2729-2748 (2016).
87.   N. Talwar, D. C. Musch, J. D. Stein, Association of Daily Dosage and Type of Statin Agent With
Risk of Open-Angle Glaucoma. *JAMA ophthalmology* **135**, 263-267 (2017).
88.   X. Y. Song *et al.*, Atorvastatin reduces IOP in ocular hypertension in vivo and suppresses ECM in
trabecular meshwork perhaps via FGD4. *Int J Mol Med* **49** (2022).
89.   L. Cong, S. Fu, J. Zhang, J. Zhao, Y. Zhang, Effects of atorvastatin on porcine aqueous humour
outflow and trabecular meshwork cells. *Exp Ther Med* **15**, 210-216 (2018).
90.   C. L. Von Zee, E. B. Stubbs, Jr., Geranylgeranylation facilitates proteasomal degradation of rho G-
proteins in human trabecular meshwork cells. *Invest Ophthalmol Vis Sci* **52**, 1676-1683 (2011).
91.   M. Lakk *et al.*, Membrane cholesterol regulates TRPV4 function, cytoskeletal expression, and the
cellular response to tension. *Journal of lipid research* **62**, 100145-100145 (2021).
92.   M. Y. Jung *et al.*, Fatty acid synthase is required for profibrotic TGF-beta signaling. *FASEB J* **32**,
3803-3815 (2018).
93.   M. Lee *et al.*, Phosphorylation of Acetyl-CoA Carboxylase by AMPK Reduces Renal Fibrosis and Is
Essential for the Anti-Fibrotic Effect of Metformin. *J Am Soc Nephrol* **29**, 2326-2336 (2018).
94.   J. Bates *et al.*, Acetyl-CoA carboxylase inhibition disrupts metabolic reprogramming during
hepatic stellate cell activation. *J Hepatol* **73**, 896-905 (2020).
95.   S.-N. Lin *et al.*, Development of antifibrotic therapy for stricturing Crohn's disease: lessons from
randomized trials in other fibrotic diseases. *Physiological Reviews* **102**, 605-652 (2022).
96.   M. Martin *et al.*, Role of endothelial cells in pulmonary fibrosis via SREBP2 activation. *JCI Insight*
6 (2021).
97.   M. Matsumoto *et al.*, Acetyl-CoA carboxylase 1 and 2 inhibition ameliorates steatosis and
hepatic fibrosis in a MC4R knockout murine model of nonalcoholic steatohepatitis. *PloS one* **15**,
e0228212-e0228212 (2020).
98.   J. N. C. Bailey *et al.*, Hypothesis-independent pathway analysis implicates GABA and Acetyl-CoA
metabolism in primary open-angle glaucoma and normal-pressure glaucoma. *Hum Genet* **133**,
1319-1330 (2014).
99.   L. Uttarwar, B. Gao, A. J. Ingram, J. C. Krepinsky, SREBP-1 activation by glucose mediates TGF-β
upregulation in mesangial cells. *Am J Physiol Renal Physiol* **302**, F329-341 (2012).
100. A. Ferrari, S. Maretto, D. Girotto, D. Volpin, G. M. Bressan, SREBP contributes to induction of
collagen VI transcription by serum starvation. *Biochem Biophys Res Commun* **313**, 600-605
(2004).
101. R. Van Krieken *et al.*, Sterol Regulatory Element Binding Protein (SREBP)-1 is a novel regulator of
the Transforming Growth Factor (TGF)-β receptor I (TβRI) through exosomal secretion. *Cell*
Signal **29**, 158-167 (2017).

Figure Legends:

Figure 1: Mechanical stress on trabecular meshwork induces SREBPs activation

(**A**) Localization of SREBPs in HTM cells subjected to cyclic mechanical stretch was checked using
immunofluorescence (IF). After 2 h of mechanical stress, HTM cells showed strong nuclear localization of
both SREBP1 (top panel second-row third column) and SREBP2 (bottom panel second-row third column).
Phalliodin was used to stain the distribution of filamentous actin (F-actin) fiber in the cells. After 2 h of

mechanical stress, there was no observable difference in actin distribution inside the HTM cells (top and
bottom panel second-row fourth column). The nucleus was stained with DAPI in blue. Images were
captured in z-stack in a confocal microscope, and stacks were orthogonally projected. Scale bar 20
micron. (**B**) Protein expression of both nuclear form SREBP1 (N-SREBP1) and SREBP2 (N-SREBP2) were
significantly increased in TM derived from enucleated porcine anterior segments perfused under the
elevated pressure of 30 mmHg for 5 h compared with 15 mmHg. The results were based on semi-
quantitative immunoblotting with subsequent densitometric analysis. β-actin was used as a loading
control. Values represent the mean ± SEM, where n = 4-5 (biological replicates). *$p < 0.05$ was
considered statistically significant. CTL: un-stretched control HTM cells, ST: stretched HTM cells.

Figure 2: Constitutive induction of SREBPs activation modulates actin and focal adhesion dynamics

(**A**) Compared to AdMT, Ad5-N-SREBP1a and Ad5-N-SREBP1c treatment on HTM cells significantly
increased N-SREBP1 mRNA expression, and Ad5-N-SREBP2 significantly increased N-SREBP2 mRNA
expression in HTM cells. 18S were used as internal controls for qPCR analysis. (**B**) Compared to AdMT,
Ad5-N-SREBP1 and Ad5-N-SREBP2 significantly increased N-SREBP1 protein expression, and Ad5-N-
SREBP2 significantly increased N-SREBP2 protein expression. The results were based on semi-
quantitative immunoblotting with subsequent densitometric analysis. β-actin was used as a loading
control. (**C**), (**D**), (**E**), and (**F**) Immunofluorescence (IF) shows the distribution of SREBP1, SREBP2,
filamentous actin (F-actin) fibers, vinculin, and paxillin in HTM cells under AdMT and Ad5-N-SREBPs
treatments. (**C**) and (**D**) Ad5-N-SREBP1a (second-row third column), and Ad5-N-SREBP1c (third-row third
column) induced strong staining of SREBP1 in the nucleus in HTM cells compared to AdMT (first-row
third column). Similarly, (**E**) and (**F**) Ad5-N-SREBP2 (second-row third column) induced strong staining of
SREBPs in the nucleus in HTM cells compared to AdMT (first-row third column). Compared to AdMT
(first-row fifth column), (**C**) and (**D**) Ad5-N-SREBP1a (second-row fifth column) and Ad5-N-SREBP1c
(third-row fifth column), and (**E**) and (**F**) Ad5-N-SREBP2 (second-row fifth column) caused the increased
distribution of F-actin fibers stained by phalloidin (purple/grayscale) in HTM cells, and induced increased
lamellipodia and filopodia formation (indicated by yellow arrows). (**C**) and (**D**) Ad5-N-SREBP1a (second-
row fourth column), Ad5-N-SREBP1c (third-row fourth column), and (**E**) and (**F**) Ad5-N-SREBP2 (second-
row fourth column) also induced more distribution of vinculin (green/grayscale) and paxillin
(green/grayscale) at the edges of F-actin fibers (indicated by white arrows) in HTM cells compared to the
AdMT (first-row fourth column). The nucleus was stained with DAPI in blue. Images were captured in z-
stack in a confocal microscope, and stacks were orthogonally projected. Scale bar 20 micron. Values
represent the mean ± SEM, where n = 4 (biological replicates). *$p < 0.05$ was considered statistically
significant.

Figure 3: Pharmacological inhibition of SREBP-SCAP pathway lowers IOP

(**A**) Nuclear fraction from PTM cells after 20 µM fatostatin for 24 h treatment. Immunoblotting result
shows that both N-SREBP1 (left panel) and N-SREBP2 (right panel) were significantly decreased. Lamin
B1 was used as loading control. (**B**) Primary HTM cells were treated with 20 µM fatostatin for 24 h, and
immunoblotting results show that both N-SREBP1 (left panel) and N-SREBP2 (middle panel) were
significantly decreased under fatostatin treatment, but SCAP protein expression level was no significant
difference in two groups (right panel). GAPDH was used as loading control. (**C**) and (**D**)
Immunofluorescence (IF) show the changes of SREBP1 and SREBP2 distribution in HTM cells after 20 µM
fatostatin treatment (left panel). In the fatostatin treated HTM cells, less SREBP1 (green staining) and

SREBP2 (green staining) were distributed inside the nucleus (second row), compared to DMSO treated
control HTM cells (first row). The nucleus was stained with DAPI in blue. Images were captured in z-stack
in a confocal microscope, and stacks were orthogonally projected. Scale bar 20 microns. Quantification
of immunofluorescence images using ImageJ-based fluorescence intensity measurements show a
significant decrease in both nuclear SREBP1 and nuclear SREBP2's mean fluorescence intensity in
fatostatin treated HTM cells (right panel). Freshly enucleated porcine eyes were perfused with 20 μM
fatostatin or DMSO control after baseline was established with perfusion media containing D-glucose at
37 °C. (**E**) Graphical representation of relative fold change in IOP showed a significant decrease post 12 h
of fatostatin perfusion (red line) and remained significant until 22 h compared to vehicle DMSO perfused
control (blue line). (**F**) Histological examination of the outflow pathway tissues using Hematoxylin and
Eosin (H and E) staining showed no gross morphological alterations between fatostatin perfusion
compared to the DMSO control. Scale bar 100 micron. (**G**) and (**H**) Immunolocalization of SREBP1,
SREBP2, ECM proteins, including COL1A and FN (green staining) in the outflow pathway tissue in DMSO
control or fatostatin. In the fatostatin perfused eyes, SREBP1, SREBP2, COL1A, and FN (left panel second
column) showed a decreased distribution in the TM-JCT region. TM indicates trabecular meshwork. AP
indicates aqueous plexus. Images were captured in z-stack in a confocal microscope, and stacks were
orthogonally projected. Scale bar 50 micron. Quantification of immunofluorescence images using
ImageJ-based fluorescence intensity measurements showed a significant decrease in the fluorescence
intensity of SREBP1, SREBP2, COL1A and FN in the TM outflow pathway (right panel). (**I**) Quantitative
analysis of protein expression from 20 μM fatostatin perfused TM tissues showed a significant decrease
in N-SREBP1, N-SREBP2, COL1A and FN compared to vehicle DMSO perfused control. Lamin B1 was used
as loading control. Values represent the mean ± SEM, where n = 4-5 (biological replicates). *$p < 0.05$ was
considered statistically significant.

Figure 4: Molecular inactivation of SREBPs by SCAP knockdown lowers IOP

(**A**) Start from 10 days after saline and viral injection, IOP was significantly decreased in Ad5.CMV.iCre-
eGFP injection group compared to untouched group, saline injection group, and Ad5.CMV.eGFP
injection group. This significant reduction in IOP in Ad5.CMV.iCre-eGFP injection group was sustained
until 30 days post-injection. On 40 days and 50 days post-injection, compared to saline injection group
and Ad5.CMV.eGFP injection group, IOP in Ad5.CMV.iCre-eGFP injection group was significantly
decreased, it was also lower than untouched group, but not statistically significant. (**B**) IOP changes in
each group was analyzed and compared to their baseline normal IOP (before injection/time point 0
days), the graphical representation shows that IOP was significantly decreased in Ad5.CMV.iCre-eGFP
injection group starting from 10 days after injection and sustained until 50 days post-injection. There
were no significant changes in IOP level in other control groups. (**C**) IOP percentage change (Delta IOP)
compared to their baseline normal IOP in each group was calculated, graphical representation shows
that IOP in Ad5.CMV.iCre-eGFP injection group was decreased as much as 40.38 % of baseline normal
(before injection/time point 0 days) with an average decrease of 27.85 %. The IOP in other control
groups showed less than 13 % changes of baseline normal IOP. (**D**) Histological examination of the TM
using Hematoxylin and Eosin (H and E) staining showed no gross morphological alterations between four
groups. Scale bar 100 micron. (**E**), (**F**) and (**G**) Immunolocalization of SCAP, SREBP1 and SREBP2 proteins
in TM in four groups. SCAP showed decreased distribution in TM in Ad5.CMV.iCre-eGFP injection group
compared to all other shame control groups. SREBP1 and SREBP2 showed decreased distribution in the
nuclear regions in TM in Ad5.CMV.iCre-eGFP injection group compared to other shame control groups.

TM indicates trabecular meshwork and marked by white color arrow. Images were captured in z-stack in
a confocal microscope, and stacks were orthogonally projected. Scale bar 50 micron. Top left corner in
each image shows magnification of the framed area showing the nuclear regions in TM and marked by
green color arrow. Scale bar 10 micron. Values represent the mean ± SEM, where n = 9-10 (biological
replicates). *$p < 0.05$ was considered statistically significant.

Figure 5: Inactivation of SREBPs inhibits lipogenesis in HTM cells

(**A**) Fatostatin treatment significantly decreased SREBPs responsive genes mRNA expression involved in
1) cholesterol biosynthesis: low-density lipoprotein receptor (LDLR), hydroxymethylglutaryl-CoA
synthase 1 (HMGCS1), 3-hydroxy-3-methylglutaryl-CoA Reductase (HMGCR), SREBP2; 2) fatty acid
biosynthesis: acetyl-CoA carboxylase (ACC), fatty acid synthase (FASN), SREBP1, peroxisome proliferator-
activated receptor gamma (PPARG), acetyl-CoA synthetase ; 3) triglyceride and phospholipid
biosynthesis: glycerol-3-phosphate acyltransferase (GPAT). 18S were used as internal controls for qPCR
analysis. (**B**) Immunoblotting show that fatostatin significantly reduced FAS, HMGCR, and ACC protein
expression compared to the control. GAPDH was used as loading control. (**C**), (**D**) and (**E**) show that
fatostatin significantly reduced total lipid biosynthesis in HTM cells, including normalized
phospholipid/cell, normalized cholesterol/cell, and normalized triglyceride/cell. Values represent the
mean ± SEM, where n = 4 (biological replicates). *$p < 0.05$ was considered statistically significant.

Figure 6: SREBPs inactivation alters actin polymerization machinery and redistributes focal adhesions

(**A**) and (**B**) Immunofluorescence (IF) show the distribution of filamentous actin (F-actin) fiber, vinculin
and paxillin distribution in HTM cells under DMSO and fatostatin treatment. Fatostatin caused reduced
distribution of F-actin fiber stained by phalloidin (green/grayscale) in HTM cells (first and third column)
compared to the control. Fatostatin treatment also caused less distribution of vinculin (red/grayscale)
and paxillin (red/grayscale) at the edges of F-actin fiber, but more cytoplasmic distribution (indicated by
white arrows) in HTM cells (first and forth column) compared to the control. The nucleus was stained
with DAPI in blue. Images were captured in z-stack in a confocal microscope, and stacks were
orthogonally projected. Scale bar 20 micron. (**C**) Globular actin (G-actin) and F-actin were extracted from
HTM cells treated with DMSO or fatostatin. Immunoblotting shows that fatostatin significantly
decreased F-actin/G-actin ratio in HTM cells compared to the control. (**D**) Immunoblotting shows that
fatostatin treatment significantly reduced Fascin-1 and LIMK1 protein expression, but no difference in
vinculin expression compared to the control. GAPDH was used as loading control. (**E**) Immunoblotting
shows that there was no significant difference in phosphor-cofilin (P-cofilin) and total-cofilin (T-cofilin)
protein expression level in DMSO and fatostatin treated HTM cells. Ratio of P-cofilin/T-cofilin was
significantly decreased in fatostatin treated HTM cells compared to the control. GAPDH was used as
loading control. (**F**) Immunoblotting shows that fatostatin significantly decreased phosphor-paxillin (P-
paxillin) protein expression level, increased total-paxillin (T-paxillin) protein expression level, and
significantly reduced the ratio of P-paxillin/T-paxillin compared to the control. GAPDH was used as
loading control. Values represent the mean ± SEM, where n = 4 (biological replicates). *$p < 0.05$ was
considered statistically significant, and ns denotes non-significant.

Figure 7: SREBPs inactivation lowers ECM production, secretion, and alters remodeling

(**A**) Regulation of ECM mRNA expression under fatostatin treatment. Fatostatin treatment significantly
decreased mRNA expression in fibronectin (FN), collagen 1α (COL1α), collagen 6α (COL6α), collagen 4α

(COL4α), tenascin C (TNC), transforming growth factor β2 (TGFβ2), and sarcospan (SSPN) in HTM cells
compared to DMSO treated control HTM cells. (**B**) Immunoblotting for whole cell lysate (CL) shows that
fatostatin significantly decreased FN, COL1A protein expression in fatostatin treated HTM cells
compared to DMSO treated control HTM cells. GAPDH was used as loading control. (**C**) Immunoblotting
for conditioned media (CM) shows that fatostatin significantly decreased FN and COL1A protein
expression in fatostatin treated HTM CM compared to DMSO treated control HTM CM. PonceauS was
used as loading control. (**D**) and (**E**) Immunofluorescence (IF) show the FN and COL1A expression and
distribution in HTM cells after DMSO or fatostatin treatment. Fatostatin reduced FN (green staining) and
COL1A (green staining) distribution compared to the control. The nucleus was stained with DAPI in blue.
Images were captured in z-stack in a confocal microscope, and stacks were orthogonally projected. Scale
bar 20 micron. Values represent the mean ± SEM, where n = 4 (biological replicates). *p < 0.05 was
considered statistically significant.

Supplementary Figure 1: Expression and distribution of SREBP isoforms and SCAP in TM

(**A**) Expression profile of SREBP isoforms and SCAP in primary human trabecular meshwork (HTM) cells
by reverse‑transcription polymerase chain reaction (RT‑PCR) showing that SREBP1 transcript 2, SREBP2
transcript 1, and SCAP were expressed in HTM. (**B**) Protein expression of proform SREBP1 (Pro-SREBP1)
and SREBP2 (Pro-SREBP2), and nuclear form SREBP1 (N-SREBP1) and SREBP2 (N-SREBP2), and SCAP in
primary HTM cells. GAPDH was used as the loading control. (**C**) Protein expression of Pro-SREBP1 and
Pro-SREBP2, N-SREBP1 and N-SREBP2, and SCAP in primary porcine trabecular meshwork (PTM) cells.
GAPDH was used as the loading control. (**D**) Immunofluorescence (IF) shows the cytosolic distribution of
SREBP1 (green staining), SREBP2 (green staining) and SCAP (green staining) in primary HTM cells. The
nucleus was stained with DAPI in blue. Images were captured in z-stack in a confocal microscope, and
stacks were orthogonally projected. Scale bar 20 microns. (**E**) Tissue distribution of SREBP1 (green
staining) and SREBP2 (green staining) and SCAP (green staining) in the aqueous humor (AH) outflow
pathway of a normal human eye specimen by IF. The nucleus was stained with DAPI in blue. Images
were captured in z-stack in a confocal microscope, and stacks were orthogonally projected. Scale bar 50
microns. (**F**) The negative control for SREBPs and SCAP distribution in AH outflow pathway (in the
presence of secondary antibody alone) did not show any significant staining (green panel).

Supplementary Figure 2: HTM cell changes after Ad5-N-SREBPs treatment

HTM cells were treated with either AdMT or Ad5-N-SREBPs for 24 h followed by 48 h serum starvation.
(**A**) Bright-field cell culture images were captured by Nikon TS100 Inverted Phase Contrast Microscope.
After treatment, Ad5-N-SREBPs treated HTM cells displayed increased lamellipodia and filopodial
extension formation (indicated by black arrows), compared to AdMT treatment.

Supplementary Figure 3: Dose-dependent effects of fatostatin on SREBPs activation in PTM cells

(**A**) and (**B**) Dose-dependent treatment of fatostatin on PTM cells to test the optimal concentration of
fatostatin on TM cells. Immunoblotting shows 20 μM fatostatin for 24 h treatment significantly
decreased both N-SREBP1 and N-SREBP2 expression. GAPDH was used as a loading control. (**C**)
Examination of the effects of 20 μM fatostatin on primary PTM cell viability. Cell viability assay in serum-
starved PTM cells treated with 20 μM fatostatin for 24 h was performed using fluorometric live/dead
staining of FDA-PI staining. The first row represents the control DMSO treatment, and the second row
represents 20 μM fatostatin treatment of serum-starved PTM cells. The first, second, and the third

columns are bright field image of PTM cells 24 h after treatment, FDA, and PI staining respectively. The
green channel shows viable cells that take up FDA and emit green fluorescence, and the red channel
shows dead cells, which take up PI and give out red florescence. The graphical representation denotes
the mean percentage ratio of FDA/PI, including the viable cells (green bar) and the non-viable cells (red
bar) in both control and 20 μM fatostatin treated cells. No significant cell death was observed after 20
μM fatostatin for 24 h treatment. Values represent the mean ± SEM, where n = 4 (biological replicates).
*p < 0.05 was considered statistically significant.

Supplementary Figure 4: Molecular inactivation of SREBPs by knocking down SCAP in old and adult
SCAP^{f/f}* mice lowers IOP *in vivo

(**A**) In old *SCAP^{f/f}* mice, start from 10 days after saline and viral injection, IOP was consistently and
significantly decreased in Ad5.CMV.iCre-eGFP injection group compared to Ad5.CMV.eGFP injection
group, and sustained until 50 days post-injection. Compared to untouched and saline injection group,
IOP was also lower in Ad5.CMV.iCre-eGFP injection group, but not consistently statistally significant. (**B**)
In old *SCAP^{f/f}* mice, IOP changes in each group was analyzed and compared to their baseline normal IOP
(before injection/time point 0 days), the graphical representation shows that IOP was significantly
decreased in Ad5.CMV.iCre-eGFP injection group starting from 10 days after injection and sustained
until 50 days post-injection. There was no significant changes in IOP level in other control groups. (**C**) In
old *SCAP^{f/f}* mice, IOP percentage change (Delta IOP) compared to their baseline normal IOP in each
group was calculated, graphical representation shows that IOP in Ad5.CMV.iCre-eGFP injection group
was decreased as much as 42.72 % of baseline normal (before injection/time point 0 days) with an
average decrease of 27.61 %. The IOP in other control groups showed less than 17 % changes of baseline
normal IOP. (**D**) In adult *SCAP^{f/f}* mice, start from 10 days after saline and viral injection, IOP was
consistently and significantly decreased in Ad5.CMV.iCre-eGFP injection group compared to
Ad5.CMV.eGFP injection group, and sustained until 50 days post-injection. Compared to untouched,
saline injection group, IOP was also lower in Ad5.CMV.iCre-eGFP injection group, but not consistently
statitally significant. (**E**) In adult *SCAP^{f/f}* mice, IOP was significantly decreased in Ad5.CMV.iCre-eGFP
injection group starting from 10 days after injection and sustained until 50 days post-injection. There
were no significant changes in IOP level in other control groups. (**F**) In adult *SCAP^{f/f}* mice, IOP in
Ad5.CMV.iCre-eGFP injection group was decreased as much as 43.14 % of baseline normal (before
injection/time point 0 days) with an average decrease of 28.10 %. The IOP in other control groups
showed less than 17 % changes of baseline normal IOP. Values represent the mean ± SEM, where n = 4-5
(biological replicates). *p < 0.05 was considered statistically significant.

1020

1021

1022 **FIGURE 1**

1023

1024

1025

 FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

 SUPPLEMENTARY MATERIAL:

 Supplementary Figure 1:

1043   **Supplementary Figure 2:**

1044

 Supplementary Figure 3:

 Supplementary Figure 4:

1048

1049

1050

1051