At the core of salinity: convergent and divergent transcriptome response pathways to neutral

and alkaline salinity in natural populations of *Arabidopsis thaliana*

Maria Almira-Casellas[1]; Sílvia Busoms[1*]; Laura Pérez-Martín[1]; Glòria Escolà[1], Álvaro López-Valiñas[2]; Antoni

Garcia-Molina[3]; Mercè Llugany[1*], Charlotte Poschenrieder[1]

[1]Plant Physiology Unit, Biosciences Faculty, Universitat Autònoma de Barcelona (UAB), 08193 Bellaterra, Spain

[2] Unitat Mixta d'Investigació IRTA-UAB en Sanitat Animal, Centre de Recerca en Sanitat Animal (CReSA), Campus UAB,

08193 Bellaterra, Spain

[3]Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Barcelona, Spain

*Corresponding author: Mercè Llugany (0000-0002-9940-0383; merce.llugany@uab.cat) and Sílvia Busoms (0000-0002-

9917-7118; silvia.busoms@uab.cat)

Submitting author:

ORCID and e-mail of all authors:

Maria José Almira Casellas; 0000-0002-4035-8766; mariajose.almira@uab.cat

Sílvia Busoms; 0000-0002-9917-7118; silvia.busoms@uab.cat

Laura Pérez-Martín; 0000-0001-6066-3939; laura.perezmartin@unige.ch

Glòria Escolà; 0000-0003-4231-4334; gloria.escola@uab.cat

Álvaro López-Valiñas; 0000-0002-7492-5108; alvaro.lopezv@irta.cat

Antoni Garcia; 0000-0002-7790-9736; antoni.garcia@cragenomica.es

Mercè Llugany: 0000-0002-9940-0383; merce.llugany@uab.cat

Charlotte Poschenrieder: ORCID 0000-0002-3818-0874; charlotte.poschentrieder@uab.es

Abstract

More than 70% of land's cultivated area is affected by alkaline salinity stress. As 98% of plants are

glycophytes – unable to successfully reproduce under salinity – our previous research focused on

comparative studies of *Arabidopsis thaliana* demes with differential performance under neutral and

alkaline salinity *(neuSAL* and *alkSAL*) due to local adaptation processes. Here, an integrated analysis

on leaf tissue was performed, including physiological indicators, nutritional status, endogenous

phytohormonal concentration and transcriptome profiling, to further understand differences in

molecular mechanisms underlying *neuSAL* and *alkSAL* responses. The results support that *alkSAL* is

more detrimental to plant performance than *neuSAL* and indicate higher sensitivity to *alkSAL* in demes

locally adapted to coastal siliceous soils. A decreased internal Fe use efficiency in coastal demes under

alkSAL is proposed to be the driver of their enhanced sensitivity, and sequence variation at β-CA1 and

α-CA1 locus is hypothesized to contribute to the imbalance of Fe homeostasis. Dissection on the down-

regulated transcripts shared by *neuSAL* and *alkSAL* confirmed enhanced inhibition of central features

on primary and secondary metabolism in coastal individuals under *alkSAL*. The cell wall and vacuolar

β-galactosidase *BGAL4* was revealed as a candidate for conferring tolerance to *neuSAL* by favoring

stress-regulated cell wall rearrangement, but not to *alkSAL*, probably due to pH-restricted enzymatic

activity. In addition, differential modulation of endogenous phytohormonal cues was reported among
salinity types and demes, by which higher alteration of the auxinic, ethylene and jasmonic acid
signaling pathways was exerted by alkSAL but sustained ABA biosynthesis was detected only in
coastal plants under neuSAL. Weighted correlation network analysis (WGCNA) confirmed the
involvement of the identified candidate genes in co-expression modules significantly correlating with
favorable responses to *neuSAL* and *alkSAL*. Overall, the present study provides useful insights into key
targets for breeding improvement in alkaline saline soils.
48

49 **Keywords**
50 *Arabidopsis* demes, abiotic stress, alkaline salinity, transcriptome, WGCNA
51

52 **Introduction**
53

Saline stress is a major agricultural constraint and its co-occurrence with alkalinity is an arising
environmental concern, specially under arid and semiarid climates. Based on the soil map of the world
by FAO/UNESCO, more than one million hectares of the Earth's land are affected by alkaline salinity
stress. This is the case of the Mediterranean region, where excess of soluble Na^+, Ca^{2+}, Mg^{2+}, K^+, CO_3^{2-}
and HCO^{3-} ions get to the superficial soil layers due to high evapotranspiration rates (Singh 2021).
Besides the induction of osmotic, ion and ROS stress by neutral salinity, alkaline salinity exacerbates
nutrient deficiency (Liu et al., 2021) and is reported to exert greater inhibition on plant germination,
growth, photosynthesis, and root system activity (Guo et al., 2010; Marconi et al. 2013; Guo et al.,
2017). Although there is increasing evidence that alkaline salinity is more detrimental for overall plant
performance than neutral salinity, the mechanisms driving such synergistic effects are not yet clearly
established. In this regard, more attention should be paid to the environmental factors that generate
local-scale diversity patterns in the model plant *Arabidopsis thaliana*, since understanding natural
genetic and phenotypic variation is essential for ecological and evolutionary studies (Bomblies et al.,
2010).
Genetic differentiation occurs at all geographic scales in *A. thaliana*, and the study of local populations
provides a valuable genetic tool for detecting adaptive traits (Bakker et al., 2006). To this end,
differences in tolerance to salinity and moderate levels of carbonate as individual stress factors have
been described among natural populations of *A. thaliana* in the NE region of Catalonia, Spain (Busoms
et al., 2015; Terés et al., 2019). Although T6[CS] and Ro2[CS] were shown to perform better under salinity
on siliceous soils than V1[PC] and LM2[PC], it was proven that tolerance to neutral salinity does not confer
tolerance to alkaline salinity in the studied individuals, and that populations from soils with
intermediate salinity and carbonate levels performed better under alkaline salinity stress due to
enhanced phenotypic plasticity (Pérez-Martín et al., 2022). However, the transcriptomic basis of such
differential response were not explored. Plants are exposed to the co-occurrence of biotic and abiotic

stresses in their natural habitats but the effects of combined stresses at the genetic level are much less

extensively reviewed than plant responses to single stress factors. Still, there are several studies

targeting the existing interconnection of signaling pathways that regulate the coordination of plant

responses to different stresses (Sharma et al., 2018; Hosseini et al., 2021; Georgii et al., 2017; Suzuki

et al., 2016; Forieri et al., 2017; Rasmussen et al., 2013), whose results highlight that combinatorial

stress triggers responses not evoked by single stressors. Despite the scarcity of studies on the effects

of co-occurrence of salinization and alkalization at the transcriptomic level in *Arabidopsis*,

transcriptome responses to alkaline salinity have been assessed in several crops. In cotton (*Gossypium*

hirsutum L.), shared responses in transcripts involved in ion homeostasis were observed under NaCl

and Na_2CO_3 conditions, while differential expression of genes involved in starch and sucrose

metabolism was reported to be exclusive of Na_2CO_3 stress due to high pH (Zhang et al.,

2018). Transcriptomics on alkalinity-tolerant and sensitive wheat varieties treated for nine days with

100 mM mixed salts (9:1 $NaHCO_3$:Na_2CO_3 at pH 8.90) revealed enhanced ability of nutrient uptake,

intracellular pH regulation and ROS scavenging activation as the main traits conferring tolerance to

alkaline salinity (Sun et al., 2020). Similarly, recent transcriptome analysis in *Brassica napus* revealed

higher expression of genes from carbohydrate metabolism, photosynthetic processes, ROS regulating,

and response to salt stress attributing tolerance to alkaline salinity (Meng et al., 2017). Enhanced

antioxidant capacity was also reported to be a major player in the tolerance response of alfalfa to saline-

alkaline stress (An et al., 2016). In rice, alkaline tolerant and sensitive cultivars were compared at the

transcriptomic level under 0.5% Na_2CO_3 solution and pH 11, and differentially expressed genes from

plant hormone signal transduction pathway were primarily related to the tolerance response (Li et al.,

2018).

Linking changes in gene expression profiles to functional consequences is the ultimate goal of

transcriptomic analysis on stress tolerance-related traits (Yan et al., 2020). The combination of

physiological studies with transcriptomic analyses provides a deeper understanding of biological

processes (Yang et al., 2022) and, based on the available studies, some of which are reported above,

nutrient homeostasis and changes in endogenous phytohormone levels are considered among the most

important mechanisms conferring tolerance in the context of neutral and alkaline salinity. While

maintaining nutrient homeostasis starts from ion sensing, uptake, transport, and activation of response

mechanisms through regulation of gene networks (Farooq et al., 2018), the alteration in the content of

phytohormone implies a specific crosstalk between endogenous plant growth regulators. This leads to

the triggering of signal transduction pathways in response to external conditions and can affect

germination (Atici et al., 2005), morphology of root and aerial structures (Mao et al., 2018, Wen et al.,

2018, Song et al., 2019), and reproduction, specially flowering (Guo et al., 2019, Shu et al., 2018, Bao

et al., 2019). These considerations reinforce the idea that differences in mineral nutrition and

phytohormone profile orchestrate adaptive processes in populations grown on specific local

environments by modulating their growth and reproductive responses.

The present study aims to identify the levels and functions of plant stress regulatory networks
underlying two contrasted types of soil salinity: neutral and alkaline. To this end, we seeked for shared
and exclusive *Arabidopsis* responses to both stress conditions among genotypes locally adapted to
either siliceous (neutral) or calcareous (alkaline) saline substrates. . Four *A. thaliana* local populations
(demes) were subjected to neutral (*neuSAL*) and alkaline salinity (*alkSAL*) under the same light and
temperature conditions. The physiological and transcriptomic responses of the 4 demes under each
type of salinity were analyzed, and convergent and singular metabolic pathways involved in each type
of salinity are described. Finally, co-expression network analysis correlating transcriptomics with
physiological data were performed to identify hub pathways in shared and exclusive plant responses
to neutral and alkaline salinity.
125
126 **Results & Discussion**
127
128 *1.   Physiological responses of A. thaliana demes under neutral and alkaline salinity*
129
To see if the adaptation to specific soils confer tolerance to neutral and alkaline salinity, two *A. thaliana*
demes from coastal siliceous soils (cs), NaCl-rich and pH between 5.7-6.2 (T6^{CS} and Ro2^{CS}); and 2
demes from pre-litoral calcareous soils (pa), considered non-saline, and with moderate carbonate
content (10-15%) (V1^{PC}, LM2^{PC}) were selected. Plants were grown on potting mix soil irrigated with
neuSAL (100 mM NaCl, pH 5.9) or *alkSAL* (85 mM NaCl; 15 mM NaHCO$_3$, pH 8.3) for 2 weeks. The
physiological, nutritional, and phytohormonal responses of the studied demes were assessed and
multivariate analyses of all measured variables were performed to establish the association between
traits and plant stress tolerance. Figures S1 and S2 detail the geographic location and source of the
studied demes and the study workflow, respectively.
139
140 *Demes from coastal siliceous soils display reduced growth and photosynthesis under alkaline salinity*
141
Overall, *neuSAL* and *alkSAL* caused 50% decrease in relative rosette diameter (RD) [(RD$_{Treatment}$)/
(RD$_{Control}$)] regardless of deme and treatment, although LM2^{PC} showed significantly better growth
maintenance than the other demes under *alkSAL* conditions (p-value $\leq$ 0.05, Tukey's HSD) (Figure
1A; Dataset S1). The chlorophyll (Chl) fluorescence parameter, Fv/Fm, reflects the maximum
quantum efficiency of PSII photochemistry in the dark-adapted state and has been widely used for the
detection of early stress responses in plants. Under control and *neuSAL* conditions, no significant
differences in Fv/Fm among demes were observed and mean Fv/Fm values were comprised in the
reading considered as optimal (between 0.79 and 0.84) for stress free plants (Maxwell and Johnson
2004). Under *alkSAL*, T6^{CS} displayed a dramatic decrease in F$_v$/F$_m$ values (mean F$_v$/F$_m$ = 0.520) when
compared to the rest, while V1^{PC} (mean F$_v$/F$_m$ = 0.770) and LM2^{PC} (mean F$_v$/F$_m$ = 0.820) showed better

F_v/F_m maintenance (p-value ≤ 0.05, Tukey's HSD). Contrarily, under *neuSAL*, T6[CS] and Ro2[CS] (0.860

and 0.810, respectively) displayed significantly higher F_v/F_m values than V1[PC] (0.740) and LM2[PC]

(0.730) (Figure 1B; Dataset S2). Reduction of total photosynthetic activity is a common feature of

plant stress, and the results confirm that the ability to maintain the photosynthetic capacity is a good

indicator of tolerance to either *neuSAL* or *alkSAL* in *A. thaliana*. These observations point to a better

performance of demes from pre litoral calcareous soils (V1[PC], LM2[PC]) under *alkSAL* in terms of growth

maintenance and photosynthesis, and agree with the severe reduction in rosette diameter previously

reported on coastal demes under *alkSAL* (Pérez-Martín et al., 2022).

K⁺ and Fe uptake efficiency are indicators of plant tolerance to salinity

Mean-standardized values (– 1< value >1) of leaf and soil elemental contents were used for the

analysis. Overall, significantly differentiated leaf nutrient profiles (p-value ≤ 0.05, Tukey's HSD)

were observed for all demes in most elements under control, *neuSAL* and *alkSAL* treatments (Figure

1C; Datasets S3). Whereas nutrient profiles were clearly genotype-specific under control conditions,

neuSAL highlighted a significantly higher K^+ levels in T6[CS] and Ro2[CS] when compared to V1[PC] and

LM2[PC], on one hand, and opposite trends in leaf Na^+ accumulation in coastal demes, on the other: T6[CS]

accumulated significantly less leaf Na^+ when compared to all other demes, but Ro2[CS] increased leaf

Na^+ due to reported allelic variation at the HKT1;1 locus, which leads to higher shoot Na^+

accumulation in individuals bearing the weak *HKT1;1[HLS]* allele (Rus et al., 2006; Busoms et al., 2018).

Still, highest leaf Na^+ accumulation under neuSAL was reported in V1[PC] (p-value = 1,40E-04; Tukey's

HSD). Under *alkSAL*, V1[PC] outrated all other demes in leaf nutritional status, while LM2[PC] displayed

a general significant reduction in leaf nutrient concentration.

V1[PC] and LM2[PC] have been previously classified as good performers under *alkSAL* based on growth,

reproductive and physiological traits when compared to coastal T6[CS] and Ro2[CS], and the most tolerance

phenotypes under neutral and alkaline salinity were observed for T6 and V1, respectively (Pérez-

Martín et al., 2022). Therefore, nutrient ratios with reported biological importance under salinity

conditions were compared in T6[CS] – as tolerant to *neuSAL* but sensitive to *alkSAL* – and V1[PC] – as

sensitive to *neuSAL* but good performer under *alkSAL*, to further explain the implications of the

differential nutrient accumulation under each type of salinity (Figure 1D; Dataset S4). Na:K ratio in

T6[CS] (1.51) was significantly lower (p-value ≤ 0.05, Tukey's HSD) than V1[PC] (2.88) under *neuSAL*,

but higher (2.56 and 1.86 for T6[CS] and V1[PC], respectively) under *alkSAL*. It has been previously

concluded that salt tolerance is driven more by a plants ability to retain K^+ under salt stress, than its

ability to simply exclude Na^+ ions (Wang et al., 2007; Shabala & Cuin, 2008). The Na:Ca ratio

followed the same trend as Na:K, being significantly lower in T6[CS] (1.49) than in V1[PC] (2.88) under

neuSAL. Cellular Na^+ and Ca cell transport pathways are linked via SOS2 (a ser/thr protein kinase

activated by SOS3, a calcium sensor protein), as SOS2 regulates SOS1 and NHX1 (plasma membrane

and tonoplast Na^+/H^+ antiporters) but also CAX1 (a vacuolar Ca/H^+ antiporter) (Cheng et al., 2004).

Decreasing Na:Ca ratio is shown to enhance K^+ uptake and reduce Na^+ uptake, ultimately decreasing

Na:K (Garg et al., 2015). This seems to be the case for V1^{PC} under *alkSAL* but not under *neuSAL*,

which might be causing its sensitivity to neutral but not to alkaline salinity. Moreover, Fe uptake

efficiency is critical in a context of alkalinity, as high pH decreases Fe availability and leads to plant

chlorosis (Hsieh et al., 2016). Under *neuSAL*, the Fe:Mn ratio was increased in T6^{CS} (0.77) due to a

higher leaf Fe accumulation when compared to V1^{PC} (0.01 vs -0.20). Under *alkSAL*, the maintenance

of Fe:Mn despite the high increase in leaf Mn concentration displayed by V1^{PC} (0.30) indicates higher

Fe uptake or translocation efficiency (Figure 1D). This suggests that T6^{CS} – adapted to moderate levels

of neutral salinity - could be hampered by the alkalinity component of *alkSAL*. Overall, the observed

response of each deme to *alkSAL* seems driven by other factors than the Na^+ content in their native

soils, as it is the case for *neuSAL* (Busoms et al., 2018), and provides evidence that tolerance to salinity

in siliceous soils does not confer tolerance to alkaline salinity (Pérez-Martín et al., 2022).

Differential elicitation of phytohormonal signaling pathways depends on salinity type and deme

Considering that regulation of phytohormone levels is one of the most important mechanisms

conferring tolerance to a wide spectrum of abiotic stresses (Zhang et al., 2019), endogenous

phytohormones were quantified in leaf tissue of the studied demes. Average concentrations of the

analyzed phytohormones can be found in Dataset S5. A significant elicitation (p-value $\leq$ 0.01,

Student's *t*-Test) of IAA, ACC, and JA was observed under *alkSAL* when compared to control and

neuSAL treatment (Figure 2 A-C), and a decrease in leaf SA and an increase in leaf ABA concentrations

were caused by both *neuSAL* and *alkSAL* conditions (Figure 2 D-E), which supports previous

observations on ABA-SA and JA-SA antagonistic signaling (Kazan & Manners, 2012; Moeder et al.,

2010). In addition, the studied demes exhibited an origin-dependent accumulation profile for ABA

under *neuSAL*, by which coastal individuals (T6^{CS}, Ro2^{CS}) displayed higher ABA accumulation when

compared to non-coastal V1^{PC} and LM2^{PC} (Figure 2 E). Salinity-induced ABA biosynthesis causing

stomatal closure, reducing stomatal conductance and water consumption, and promoting overall plant

tolerance to salinity is well reported (Lovelli et al., 2012). Here, alkaline salinity elicits higher

alteration of hormonal networks than neutral salinity, whereas ABA accumulation seems to contribute

to the enhanced response of demes locally adapted to *neuSAL* conditions by increasing water use

efficiency (Cardoso et al., 2020) and restricting K^+ efflux (Chen et al., 2022). In fact, ABA

concentration in coastal demes under *neuSAL* correlated with increased plant relative water content

(RWC) (Figure 2 F), suggesting that ABA enhances leaf salt resistance by improving the plant water

status through reduced stomatal conductance.

Multivariate analyses reveal new players defining the alkalinity component of salinity

Multivariate analyses on all measured traits were performed for all samples by means of Principal

Component (PCA) and Multiple Pearson Correlation (MPC) analyses. The PCA showed that

eigenvalues of the two first principal components represented up to 48% (PC1 25%; PC2 23%) of the

total variance (*p*-value = 1.13E-09 for PC1 and 0.0002 for PC2 (Dataset S6). A general look at the

PC1–PC2 axes plane plot (Fig 3A) shows that alkalinity contribute to the construction of component

1 (samples submitted to neutral or alkaline salinity differentiate along the x-axis), whereas salinity

contributes to PC2 construction (control samples differentiate from *neuSAL* and *alkSAL* samples along

the y-axis). The loading matrix revealed that well correlated mineral nutrient traits (leaf Zn and leaf

Mg), as well as endogenous JA, were the variables contributing to PC1 with a loading factor above

0,70 (Dataset S7). In fact, *alkSAL* samples contained significantly higher Zn than *neuSAL* samples (*p*-

value ≤ 0.05, Tukey's HSD) (Figure 3B). In turn, MPC results showed that Zn displayed the highest

correlation with Mg and JA under *alkSAL* ($R^2 = 0.8$; $R^2 = 0.6$), but correlation with Mg was lost ($R^2 =$

0.15), and correlation with JA was inverted ($R^2 = -0.48$) under *neuSAL* (Figure 3C).

Leaf Na^+, leaf K^+ and leaf B contributed to the construction of component 2, being leaf B dramatically

decreased in the presence of salinity regardless of neutral or alkaline (Figure 3D). MPC showed how

Na-B correlated negatively under *neuSAL* ($R^2 = - 0.42$) but positively under alkSAL ($R^2 = 0.35$),

whereas no correlation was found in control ($R^2 = 0.03$) (Figure 3E). While co-toxicity of salinity and

B is reported (Pandey et al., 2019), few studies on neutral or alkaline salinity causing B deficiency are

found. Boron alleviates salinity and saline-sodicity stress by improving leaf water and soluble

carbohydrate contents as well as maintaining osmotic potential (Yousefi et al., 2020; Mehmood et al.,

2009). Remarkably, V1 displayed the highest B and K^+ maintenance under *alkSAL* and the highest B

and K^+ deficiency under *neuSAL* (Figure 3E-F).

In summary, the reported results suggest that growth and PSII efficiency maintenance are indicators

of differential tolerance to neutral and to alkaline salinity in the studied demes. Whereas neutral salinity

tolerance in coastal individuals is driven by a decreased Na^+ shoot translocation in $T6^{CS}$ and by the

ability to retain K^+ in $Ro2^{CS}$, the maintenance of leaf K^+ homeostasis by increasing leaf Ca^{2+}

accumulation is a requirement for a better response to alkaline salinity. Moreover, enhanced tolerance

to *alkSAL* in $V1^{PC}$ is partially explained by higher Fe, B and K^+ internal use efficiency. Differential

modulation of auxin, ABA and CKs among genotypes is likely involved in tolerance to either *neuSAL*

or *alkSAL* in the studied demes; and differences in leaf Zn concentration under *neuSAL* and *alkSAL*

may affect endogenous phytohormone content through the stimulation of tryptophan biosynthesis,

which is a precursor of indoleacetic acid and a potential conjugate of IAA and JA.

2. Transcriptome profiling of A. thaliana under neutral and alkaline salinity

Leaf transcriptome profiling of the 4 demes cultivated under control, *neuSAL* or *alkSAL* conditions was performed to characterize the mechanisms underlying the differential response to neutral and alkaline salinity at the gene expression level. Multi-Dimensional Scaling (MDS) was used to visualize global transcriptional similarity among samples (Figure 4A). The transcriptome of each sample was shaped by deme and colored by treatment. Accordingly, individuals showed higher transcriptional similarity between samples corresponding to the same treatment, indicating that the studied transcriptomic response is treatment specific.

Differentially expressed genes (DEGs) per deme and treatment were declared based on absolute $\log_2$ fold-change (FC) > 2 under each treatment compared to control conditions (*neuSAL*/Control, *alkSAL*/Control) (*q*-value < 0.05, Wald test). When considering all demes, a total of 6164 DEGs (2559 up-regulated and 3605 down-regulated) were identified under neuSal and 10487 DEGs (4995 up-regulated and 5492 down-regulated) under *alkSAL*. A 65% of total DEGs from *neuSAL* were found in T6[CS] and V1[PC], while 29% and 34% in Ro2[CS] and LM2[PC]. Under *alkSAL*, T6[CS], Ro2[CS], V1[PC] and LM2[PC] displayed 1.9, 4.1, 2.2 and 1.53-times more DEGs than under *neuSAL*. Under *alkSAL*, T6[CS], still displayed the most altered transcriptome, followed by Ro2[CS] (73% and 68% of total *alkSAL* DEGs, respectively), while V1[PC] and LM2[PC] transcriptomic response was more attenuated (58% and 44% of total DEGs) (Figure 4B-C).

Exclusive and shared molecular components of neutral and alkaline salinity

To gain more insight into commonalities and particularities between treatments, the subset of DEGs shared by at least 3 out of 4 demes in *neuSAL* and *alkSAL* were selected and merged, thus obtaining individual and common DEGs under both treatments (Figure 4D). On the one hand, 345 of DEGs were found in response to *neuSAL* but not to *alkSAL* and were referred as *onlyNEUSAL* DEGs set (Table S1). From these, 141 and 186 (41% and 54% of transcripts, respectively) were commonly upregulated or downregulated among demes, and 18 (5%) showed ambiguous expression trends depending on the deme (Figure 5A). On the other hand, 3585 DEGs changed under *alkSAL* but not under *neuSAL* and were referred to as *onlyALKSAL* DEGs set (Table S2). From these, 1621 and 1935 (45% and 54% of transcripts, respectively) were upregulated or downregulated, and only a minimal fraction of 29 (0,8%) showed opposite expression trends in at least 1 deme (Figure 5B). Finally, the 1219 of the selected DEGs were shared under both *neuSAL* and *alkSAL* and were referred as *coreSAL* DEGs set (Table S3). From these, 406 (65%) were upregulated, 786 (33%) downregulated and 27 (2%) with opposite expression trends in at least one deme (Figure 7A). It is well known that plants facing complex stress conditions trigger cascade of signals unique to specific stress components as well as shared responses (Zandalinas et al., 2020; Skalak et al., 2021). Studies assessing *Arabidopsis* response to combinatorial stresses described antagonistic expression trends for only 5% to 10% of transcripts, which is in line

with the expression trends found for each DEGs subset described above. Moreover, 86% of DEGs
from the *coreSAL* were found in the 4 studied demes (Figure S3), which reveals that contribution to
the *coreSAL* DEGs set is high for all studied demes and, thus, similar strategies in the studied
individuals lead to differential responses to *neuSAL* and *alkSAL*.

The overall results obtained from DEG analysis and classification provide evidence that the stress
imposed on plants by alkaline salinity is distinct from that imposed by salinity in a neutral soil (Guo
et al., 2010) and that alkaline salinity involves more complex processes than neutral salinity et the
transcriptomic level (Fang et al., 2021).

KEGG Erichment analysis of onlyNEUSAL and onlyALKSAL DEGs

Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis was performed in
onlyNEUSAL and *onlyALKSAL* DEGs to identify which categories were significantly enriched in
metabolic pathways (q-value < 0.05), and thus target the presence of alkalinity-dependent salinity
effects in the studied demes. *onlyNEUSAL* and *onlyALKSAL* enriched pathways are listed in Table S4.

Upregulated transcripts in onlyNEUSAL and onlyALKSAL reveal higher demand for
antioxidant protection under alkaline salinity. In *onlyNEUSAL*, only 2 pathways containing 6 and 2
upregulated transcripts were significantly enriched: "ribosome biogenesis" and "anthocyanin
biosynthesis". This could indicate that expression of such transcripts is triggered by the NaCl
component, likely Cl^- ion toxicity, which is present in higher extent under *neuSAL* than under *alkSAL*
is reported to be the most significant toxic component of the saline solution in some species (Munns
& Tester, 2008) and, more specifically, strongly decreases ribosome stability (Brady et al., 1984). The
role of anthocyanin accumulation in the positive regulation of tolerance mechanisms to abiotic factors
is well reviewed (Perea-Resa et al., 2017; Naing & Kim, 2021; Kubra et al., 2021; Zhang et al., 2019),
among others, has been reported. Here, activation of anthocyanin biosynthesis genes (AT4G14090 and
AT5G54060) could take place for scavenging ROS excess due to plant oxidative damage (Naing &
Kim, 2021). In *onlyALKSAL*, the most enriched pathways with up-regulated transcripts were
phenylpropanoid biosynthesis (25 DEGs) and glutathione metabolism (18 DEGs) (Figure S4).
Phenylpropanoid metabolism is elicited by abiotic stresses such as drought and salinity and, moreover,
is reported to contribute to salinity-alkalinity tolerance (Gan et al., 2021). The pH-dependent increase
in expression of DEGs from phenylpropanoid and glutathione signaling pathways is in line with that
reported in recent studies (Waters et al., 2018, Gan et al., 2021; Han et al., 2018).

The above results point to a pH-dependent response in the studied demes and a higher requirement of
the plant antioxidant machinery in the presence of alkaline salinity, which would lead to an enhanced
activation of transcripts from the phenylpropanoid and glutathione metabolism.

Salinity-triggered phytohormone responses are alkalinity-dependent. Endogenous phytohormone

quantification of the studied demes under *neuSAL* and *alkSAL* treatments revealed a sustained

biosynthesis of auxins, ACC and JA under prolonged plant exposure to alkaline salinity. In turn, higher

ABA after 2 weeks of *neuSAL* irrigation was observed only on genotypes locally adapted to neutral

salinity (Figure 2). To identify activation or repression patterns in signaling pathways accounting for

the differential phytohormone profiles, the expression levels of phytohormone-responsive genes under

each treatment were analyzed (*q*-value < 0.05; Student *t*-Test) (Dataset 13). A general trend towards

gene repression was present (Figure 6), as previously reported for other abiotic stress long term

transcriptional responses (Leyva-Pérez et al., 2015).

Higher inhibition of auxin (SAURs, Aux/IAAs) and JA (LOX, MYCs) responsive transcripts was

observed after 2 weeks of *alkSAL* exposure when compared to neutral salinity (Figure 6 B-C). In

contrast, *alkSAL* treatment caused an induction of the transcription factor WRKY and ABI (*ABA-*

insensitive) families, of repressors of the JA pathway (JAZs), and of protein phosphatases activating

abscisic acid-inducible gene expression (PP2C5 & PP2CG1), whereas their average expression levels

showed to be attenuated after prolonged exposure to *neuSAL* (Figure 6 D-G). Opposite expression

trends were displayed among treatments in ethylene and ABA responsive transcripts (ERFs and

PYL/PYRs), in which significant activation under *alkSAL* but inhibition under *neuSAL* were observed

(Figure 6 H-I). Finally, DREBs (*Dehydration responsive element binding*) showed sustained activation

under both *neuSAL* and *alkSAL,* with significantly higher transcript expression under *alkSAL* (Figure

J).

The downregulation of SAURs during the late phase of the plant salinity response agrees with a protein

degradation targeted after the observed auxin accumulation (Figure 2). Meanwhile, the reduced

expression of the early auxin-responsive SAURs and the auxin co-receptors Aux/IAAs under *alkSAL*

suggest further repression of auxin signaling mediated by the presence of JA and ABA rather than the

removal of the inducer (IAA) (Markakis et al., 2013; van).

Under alkaline salinity, an antagonistic interaction between ABA and JA signaling pathways

modulating stress-responsive gene expression (Anderson et al., 2004; Kazan & Manners, 2013; Sah et

al., 2016) is revealed, as depicted by the significant upregulation of ABA receptors (PYLs), ABA

responsive genes (ABIs) and other ABA-dependent MAPK phosphatases (PP2Cs), and the enhanced

inhibition of JAs responsive transcripts (LOXs and MYCs) in *onlyALKSAL.* Moreover, the expression

of JAZs, targeting MYCs repression, was only induced by *alkSAL.* This supports a negative regulatory

feedback loop occurring after long exposure to alkaline salinity to turn off the transcriptional

reprogramming caused by JA accumulation under *alkSAL* and to minimize JA inhibition on plant

growth, in an attempt of acclimation (Riemann et al., 2015). In turn, the stimulation of ABA signaling

and accumulation, and the NaCl tolerance observed in coastal demes under *neuSAL* are likely mediated
by WRKYs (Hussain et al., 2021), which remained active after 2 weeks of *neuSAL* treatment. Among
them, enhancers of the osmotic, ROS and hormone signaling pathways under salinity stress have been
recently reported (Price et al., 2022).

Most DREBs regulate the expression of multiple stress-inducible genes and play pivotal roles in the
plant abiotic stress response. All identified DREBs were strongly upregulated after 2-week exposure
to both *neuSAL* and *alkSAL.* Among them, DREB1A and DREB2C, reported as two of the DREBs
containing the most types of abiotic stress-related motifs in the promoter region (Xie et al., 2019), were
found. Upregulated DREBs in the plant long-term *coreSAL* gene set provide further evidence that
DREBs form a central network to control the plant salinity response also at the late stage (Sazegari et
al., 2015). Contrarily, sustained upregulation of twelve DEGs from another subfamily of AP2/ERF
transcription factors, ERF-IX, seems to be required solely by alkaline salinity stress and associates
with the plant significant ACC accumulation under *alkSAL* (Wang etal., 2022).

Overall, differences in the transcriptional regulation of phytohormones-related genes under prolonged
plant exposure to either neutral or alkaline salinity were observed. The above-mentioned expression
patterns are consistent with a higher alteration in the plant transcriptome caused by alkaline salinity
(Figure 4B) leading to a more extensive rearrangement in the phytohormonal signaling network. In
this regard, transcript inhibition for protein turnover seems to be a regulation mechanism for
phytohormone signaling pathways such as auxin and JA under *alkSAL,* and ABA under *neuSAL*
(McClellan & Chang, 2008). An implication of differential auxin and ABA regulation mechanisms
for proton extrusion, through H^+-ATPase activation by inhibition of PP2C proteins should be further
addressed, as it is an essential Fe-deficiency plant response (Sun et al., 2020; Spartz et al., 2014).

KEGG Erichment analysis of coreSAL DEGs

KEGG pathway analysis was performed in *coreSAL* DEGs to identify significantly enriched metabolic
pathways (*q*-value < 0.05) in the general salinity stress response and the presence of genotype-
dependent salinity effects in the studied demes. *coreSAL* enriched pathways are listed in Table S6.

coreSAL transcript activation reflects shared salinity stress responses. The significantly
enriched pathways in the *coreSAL* set were "Ribosome biogenesis in eukaryotes", "Protein processing
in endoplasmic reticulum" and "Cutin, suberin and wax biosynthesis" (Table S6). Ribosome
biogenesis and protein synthesis in the ER are the two pathways comprising cellular processes with
highest metabolic cost (Piazzi et al., 2019). Here, we report a consistent upregulation of genes involved
in pre rRNA and RNA processing and maturation (FIB2, PWP2, AT4G04940, AT1G67120,

AT3G06530), nucleolytic cleavage (AT5G22100), and cell differentiation (AT2G18900) in all 4
demes under neutral and alkaline salinity, followed by the activation of 11 Heat Shock Proteins –
which encode for molecular chaperones that interact with ribosomes ensuring their stability (Piazzi et
al., 2019) (Figure S5A). This indicates nucleolar stress (Yang et al., 2018) triggered by both neuSAL
and alkSAL that leads to the activation of the protein synthetic machinery in the studied demes (Figure
S5B). Moreover, enhanced activation of 2 genes involved in suberin deposition (FAR1, FAR5) –
previously reported to be transcriptionally induced by salinity (Domergue et al., 2010) - were observed
in all demes under *neuSAL* and *alkSAL* (Figure SC), which suggests that increase of suberin deposition
may enhance tolerance to neutral and alkaline salinity (Zhang et al., 2021).

coreSAL transcript inhibition reveals differential sensitivities to alkSAL in the studied demes.
The significantly enriched pathways containing down-regulated transcripts detected in the *coreSAL* set
were 'Photosynthesis' and 'Photosynthesis antenna proteins', 'Glucosinolate biosynthesis'
'Glyoxylate and dicarboxylate metabolism', 'Carbon fixation', 'Carbon metabolism', '2-
Oxocarboxylic acid metabolism', 'Nitrogen metabolism', and 'Glycine, serine and threonine
metabolism' (Figure 5C; Table S6). $T6^{CS}$ suffered the highest inhibition of all DEGs from
"Photosynthesis" and "photosynthesis antenna proteins" (an average of 1.5 and x1.8 times more
downregulation than $V1^{PC}$ and $LM2^{PC}$, respectively), followed by $Ro2^{CS}$ in 2 transcripts: PSAH1-2 (an
average of x1.5 times more downregulation in $T6^{CS}$, $Ro2^{CS}$ when compared to $V1^{PC}$ and $LM2^{PC}$ under
alkSAL) (Figure 7A). PSAH1 and PSAH2 encode subunits H of photosystem I reaction center and
their up-regulation under neutral salinity has been reported in halophytes (Hao et al., 2020). The
downregulation of photosynthesis-related transcripts in the *coreSAL* response agrees with previous
transcriptome studies under neutral salinity reporting a correlation between down regulation of antenna
proteins and higher salt sensitivity (Guo et al., 2017; Jing et al., 2019). Similarly, DEGs from
"glucosinolate biosynthesis" and a DEG subset from "carbon metabolism" displayed an average of
x1.7 times more downregulation in $T6^{CS}$ and $Ro2^{CS}$ when compared to $V1^{PC}$ and $LM2^{PC}$ under *alkSAL*
(Figure 7B).
Glucosinolates are secondary metabolites mainly found in Brassicaceae. Downregulation of
genes involved in glucosinolates biosynthesis has been reported under several types of salts
(Aghajanzadeh et al., 2017). A previous study (Pérez-Martín et al., 2021) characterized the response
to bicarbonate exposure of $T6^{CS}$ and $A1_i$ – as a carbonate sensitive and carbonate tolerant deme,
respectively – and reported the downregulation of several GSTU genes involved in glucosinolate
metabolism only in $T6^{CS}$. Two Ribulose Biphosphate Carboxylase Subunits (RBCS1-2B), an
Hydroxypyruvate Reductase (HPR) and a Fumarase (FUM2) comprised the DEG subset from "carbon
metabolism" sharing expression pattern with the glucosinolate pathway in coastal demes (Figure 7C).
The last three KEGG pathways significantly enriched in the downregulated transcript set from *coreSAL*
were "glyoxylate and decarboxylate metabolism" (19 DEGs), "glycine, serine and threonine
metabolism" (10 DEGs) and "nitrogen metabolism" (8 DEGs) (Figure S6; Table S4). *PGLP1*
(AT5G36700, from the glyoxylate decarboxylate metabolism) suffered x1.4 times more

downregulation in coastal demes under *alkSAL*. PGLP1 is one of the core metabolite repair enzymes
of plant photosynthetic carbon assimilation, as it degrades 2-phosphoglycolate (2PG) for its
reconversion to Calvin cycle metabolites (Flügel et al., 2017). After 2PG degradation to glycolate,
glyoxylate is generated, ultimately being catalyzed to glycine (Dellero et al., 2016). Thus, *coreSAL*
inhibition of glyoxylate and decarboxylate transcripts in the studied demes is likely affecting glycine,
serine, and threonine metabolism. In this regard, 10 DEGs enriched in the "glycine, serine and
threonine metabolism" pathways were downregulated in the 4 demes under neutral and alkaline
salinity (Table S6), and *AK3* (AT3G02020) was x1.6 times more downregulated in T6^{CS} and Ro2^{CS}
when compared to V1^{PC} and LM2^{PC} under *alkSAL* (Figure S6). AK3 codes for an aspartate kinase that
catalyzes the first step in aspartate-derived amino acids (Clark et al., 2015). *AK3* upregulation was also
reported in A1 – carbonate tolerant deme – under bicarbonate exposure when compared to control, but
not in T6^{CS} – carbonate sensitive deme (Pérez-Martín et al., 2021).

Finally, inhibition of transcripts from the α-CA and β-CA (Carbonic Anhydrases) (α-CA1, α-CA2, β-
CA1, β-CA2 and β-CA4), and the glutamine synthetases 1;4 (GLN1;4) and 2 (GS) family proteins
comprised the "Nitrogen metabolism" down-regulated set (Table S6). From these, *βCA-1*
(At3G01500) and *α-CA1* (At3G52720) displayed x1.7 times more inhibition in T6^{CS} when compared
to V1^{PC} and LM2^{PC} under *alkSAL* (Figure S6). βCA-1 is reported to regulate CO_2 controlled stomatal
movements in guard cells together with *βCA4* (AT1G70410). Previous electrophysiological assays
reported *βCA4* reduction in T6^{CS} under bicarbonate conditions (Pérez-Martín, 2021), and *βca1ca4*
mutant under $NaHCO_3$ conditions show reduced photosynthetic efficiency (F_v/F_m) and increased total
ion leakage (as reporter for cell death) under $NaHCO_3$ treatment (Dąbrowska-Bronk et al., 2016). α-
CA1 belongs to the α-CA family and is involved in the transformation of HCO^{3-} to CO_2 in chloroplast
stroma to supply it at the active site of Rubisco (Burén 2010).

Considering the high demand of Fe cofactors by photosynthetic enzymes (Kroh & Pilon, 2020), the
above results raise the hypothesis that the defective Fe use efficiency in coastal demes under *alkSAL*
(especially T6^{CS}) leads to a stronger photosynthetic inhibition. This would result in the observed
unbalance in carbon fixation and metabolism in T6^{CS} and Ro2^{CS}, which ultimately would compromise
their total energetic investment into secondary metabolism. In such a scenario, the stronger
downregulation of PGLP1 (entry enzyme into one of the major pathways of primary plant metabolism)
and the chloroplast carbonic anhydrases β-CA1 and α-CA1 in coastal demes under *alkSAL* is consistent
with their reduced photosynthesis efficiency (F_v/F_m values) and stronger downregulation of transcripts
from photosynthesis (PSAH1-2), carbon fixation and metabolism (RBCS1-2B, HPR and FUM2).
FUM2 (Fumarate hydratase 2 - AT5G50950, encoding a cytosolic chloroplast fumarase) would link
primary metabolism to secondary metabolism (Figure S7), leading to the downregulation of all
transcripts involved in glucosinolate metabolism and some transcripts from the glycine, serine and
threonine metabolism reported to enhance alkalinity tolerance, like AK3. Likewise, the results confirm

that a significant inhibition of glucosinolate signaling in coastal demes is caused in an alkalinity-
specific manner, likely by decreased availability of ferrous ions (Fahey et al., 2001; Wittstock et al.,
2002).

A reduced coreSAL DEG subset orchestrate contrasted plant response to neuSAL and alkSAL. All
shared DEGs displaying opposite expression trends between neutral and alkaline salinity in more
than 2 demes were analyzed to seek for contrasted transcriptomic responses among the studied demes
potentially driving their differential performance under each salinity type (Figure 7D). *BGAL4*
(AT5G56870), *DIN10* (AT5G20250) and *SEN1* (AT4G35770) exhibited contrasted expression
trends among V1^{PC} and LM2^{PC} (down-regulated) and T6^{CS} and Ro2^{CS} (up-regulated) under *neuSAL*
(Figure 8A). This suggests that activation of *BGAL4*, *DIN10* and/or *SEN1* is advantageous under
neuSAL but not under *alkSAL*. Therefore, the genomic sequences of *BGAL4*, *DIN10* and *SEN1* in the
4 study demes were extracted from the Catalan demes germplasm Whole Genome Sequencing
(WGS) data (Busoms et al., 2018) to identify sequence variation likely causal of the contrasted
expression trends in the studied demes under *neuSAL* and *alkSAL* (Figure S8; Table S7). A SNP
located only 4 bp upstream the 5'UTR at *BGAL4* promoter was consistent for all replicates of each
deme and, moreover, differentiated coastal from non-coastal demes (Figure 8B).

BGAL4 encodes a glycosyl hydrolase regulated by SNF1-related Kinase 1 (SnRK1), a kinase
involved in key plant stress responses (Peixoto et al., 2021). Degradation of structural
polysaccharides in plant cell wall by cell wall-remodeling enzymes - like BGAL4 – forms part of the
stress-regulated cell wall rearrangement (Moneo-Sánchez et al., 2016). A study comparing *A.*
thaliana Wild Type vs salt-tolerant mutant roots exposed to NaCl stress reported higher expression
of glycosyl hydrolases in the salt-tolerant mutant root, which was hypothesized to increase its
potential of salt resistance (Guo et al., 2014). In contrast, β-galactosidases are reported to
dramatically decrease enzymatic activity at high pH and to have pH optima ranging from 3.5 to 5.0
(Ross et al., 1994; Smith & Gross, 2000; Kotake et al., 2005, Hussien et al., 2021). Thus, enhanced
upregulation of *BGAL4* in T6^{CS} and Ro2^{CS} could point to an enhanced activation of plant energy
management functions under *neuSAL*, whereas a lack of ability to modulate BGAL4 activity for the
prioritization of compounds more suitable under alkaline conditions could compromise their energy
budget (Tsai & Schmidt, 2020). Moreover, the decline in photosynthesis and carbon fixation
efficiency observed in coastal demes under *alkSAL* may cause further stimulation of the hydrolase
enzymes, as these are triggered by sugar starvation (Lee et al., 2007). Transcription factor binding is
the strongest contributor to variation in mRNA levels (Pai et al., 2015). This might be the case for
BGAL4, where the SNP change in the promoter region of V1^{PC} and LM2^{PC} could contain cis-
regulatory elements responsible for its consistently lower expression.

2. WGCNA Network

To combine transcriptomic and physiological datasets and detect possible correlations between genes
determining traits of interest, all DEGs detected in each deme under *neuSAL* and *alkSAL* were retained
for WGCNA unsigned co-expression network analysis. This involved a total of 11432 DEGs (11K
WGCNA hereafter). The soft threshold power of 7 ($\beta = 7$) was selected according to the preconditions
of approximate scale-free topology (Figure S9). WGCNA 11K analysis identified sixteen distinct co-
expression modules which were assigned to different colors shown in the dendrogram (Figure S10).
The obtained module-trait correlation matrix displayed two major contrasted patterns defined by
correlation trends in 3 main traits: leaf Na^+, leaf B and relative rosette biomass (Figure 9). This
confirmed that correlation patterns observed in the performed multivariate analyses of physiological
traits (PCA and MPC) were maintained when integrating co-expressed transcriptome data into trait
correlations (WGCNA). Besides, most co-expression modules that negatively correlated with leaf Na^+
but positively correlated with leaf B and sample biomass were also displaying positive correlation with
leaf K^+ and F_v/F_m values, confirming their biological relevance in the context of salinity stress.
544
Considering all the above, modules showing significant negative correlation with Na^+ and positive
correlation with B, K^+ or FW were picked for further analysis of their implication in tolerance
mechanisms to salinity stress and GO and KEGG enrichment analysis of each selected co-expression
module was performed for each significant selected module. After that, DEGs comprised in the
significantly enriched pathways of each module were merged with the representative set of DEGs from
coreSAL, *onlyNEUSAL* and *onlyALKSAL*, and results confirmed that DEGs selection criteria carried
for transcriptome analysis – picking only DEGs detected in at least 3 out of the 4 demes comprised in
the shared and the exclusive response to neutral and alkaline salinity - accurately reflected the behavior
of the entire dataset (Table S8). Moreover, all the genes highlighted from previous analysis have also
been identified in the 3 modules explored (Figure 10).
555
Overall, the combination of multivariate analysis on physiological traits, transcriptome analysis and
WGCNA results prove that the selection criteria performed to assess effects of salinity in the studied
demes are an accurate reflection of the whole dataset behavior. This combination provides knowledge
on the hub biological processes and metabolic pathways that reflect the differential responses of
neutral-salinity-adapted demes when compared to demes with tolerance to moderate alkaline-salinity.
Moreover, it gives evidence supporting the involvement of key nutritional (K^+ homeostasis, B
sufficiency, Na^+ uptake reduction and Zn limitation) and phytohormonal cues driving tolerance to
salinity.
564
565 **Conclusions**
566
By means of an integrative physiological, ionomic, metabolomic and transcriptomic approach,
convergent and divergent biological pathways conferring shared and exclusive responses to neutral
(*neuSAL*) and alkaline salinity (*alkSAL*) have been revealed in natural populations of *Arabidopsis*

thaliana. From the analysis of up-regulated transcripts at the core of salinity, we conclude that
activation of the protein synthetic machinery and changes in cell wall composition by FARs -mediated
suberin deposition processes are common responses to salinity. Whereas under *neuSAL* antioxidant
mechanisms are boosted mainly through the anthocyanin biosynthesis pathway, the phenylpropanoid
and glutathione pathways are raised under *alkSAL*. Through screening of DEGs with contrasted
expression profiles among demes, we propose that a smart regulation of BGAL4, a cell wall and
tonoplast hydrolase, is key to tolerate different salt stresses, and a SNP detected in the 5'UTR of V1^{PC}
and LM2^{PC} is proposed as a candidate for the differential regulation of gene expression levels. In
contrast, dissection on the down-regulated transcripts at the *coreSAL* set confirms that tolerance to
neutral salinity does not confer advantageous features under alkaline salinity. Key genes involved in
the photorespiratory cycle – PGLP1, βCA1, RBCS1-2 -, organic acid accumulation – FUM2 -, and
other secondary metabolism processes for abiotic stress resistance – AK3, glucosinolate biosynthesis
genes – could be targets for alkaline salinity resistance breeding. WGCNA analysis supports that the
presented candidates are comprised in co-expression modules significantly correlating with favorable
responses to *alkSAL*. In this concern, increasing internal Fe use efficiency for maintenance of
photosynthesis rate, and rising Ca, B and K accumulation for lowering Na-K, Na-Ca, Na-B ratios
might be essential when facing alkaline salinity stress. Moreover, the enhanced activation of ABA-
dependent signaling is highlighted as a key trait for plant performance under *alkSAL*, likely by
PYL/PYRs binding to PP2Cs and consequent activation of H$^+$ATP-ase activity, allowing apoplastic
acidification. In turn, sustained ABA accumulation and signaling through WRKYs regulation are
shown exclusively in coastal demes T6^{CS} and Ro2^{CS} under *neuSAL* and are proposed to be adaptive
strategies for these *A.thaliana* demes.
592
593 **Material & Methods**
594
595   *Plant material*
596 Seeds of 2 coastal-siliceous (T6^{CS} and Ro2^{CS}) and 2 intermediate-calcareous (V1^{PC}, LM2^{PC}) *A. thaliana*
597 demes were collected from the natural habitat (Figure S1) and propagated in the laboratory. The
598 flowchart in Figure S2 summarizes the experimental procedures of the study.
599
Irrigation experiment
Seeds were surface sterilized by soaking them in 30% (v/v) commercial Clorox bleach and 1 drop of
Tween-20 for 15 min and rinsed 5 times in 18 MΩmilli-Q sterile water. Seeds were stratified at 4ºC
for 1 week in the dark and sowed on wet soil. Pots were covered with PVC film until the seedlings
germinated. Pots with germinated seedlings were placed in a growth chamber (Conviron CMP5090)
with 8 h light / 16 h dark photoperiod, an irradiance of 80 mmol·m^{-2}·s^{-1} and a constant temperature of
22 ºC. Plants were watered with 0,5-strength Hoagland solution every 3 days. After 2 weeks, all plants

were irrigated every 3 days with 0,5-strength Hoagland solution containing 0,5-strength Hoagland

solution alone (control), with 100 mM NaCl (neutral salinity: **neuSAL**) or with 85 mM NaCl and 15

mM $NaHCO_3$ at pH 8.3 (alkaline salinity, **alkSAL**) for two weeks. The Rosette Diameter (RD) and the

PSII efficiency of each sample were measured after two weeks of treatment. After harvest, samples

for ionomic analysis were dried and samples for RNA extraction and hormone determination were

immediately frozen in liquid nitrogen and stored at -80 ºC.

Leaf nutrition

Plant tissues were sampled by removing 2–3 leaves or cutting roots (1–5 mg dry weight). They were

then washed with 18 MΩ water before placing them in Pyrex digestion tubes. The sampled plant

material was dried for 2 days at 60 ºC and weighed before open-air digestion in Pyrex tubes using 0.7

mL concentrated HNO_3 at 110 ºC for 5 h in a hot-block digestion system (SC154-54-Well Hot Block,

Environmental Express, Charleston, SC, United States). Concentrations of the selected elements (Ca,

K, Mg, Na, P, S, Mo, Cu, Fe, Mn, Zn) were determined by ICP-MS (Perkin ElmerInc., ELAN 6000,

MA, United States) or ICP-OES (Thermo Jarrell-Ash, model 61E Polyscan, England) (n=6 per deme

and treatment).

PSII efficiency

The maximal photochemical efficiency of PSII was calculated from the ratio of variable (Fv) to

maximum (Fm) fluorescence [$F_v/F_m = (Fm - F_o)/F_m$] using a MINI-PAM-II/B portable chlorophyll

fluorometer (Walz) connected to an Arabidopsis Leaf Clip holder (model 2060-B). The minimum

fluorescence yield (F_o) was measured under measuring light (650 nm) with very low intensity (0.8

mmol m^{-2} s^{-1}). To estimate the maximum fluorescence yield (F_m), a saturating pulse of white light

(3000 mmol m^{-2} s^{-1} for 1 s) was applied. Leaves were dark-adapted for 30 min before measurements

and the analysis was performed in a dark room with stable ambient conditions (n=6 per deme and

treatment).

Endogenous phytohormones

Phytohormones extraction was done according to the protocol of Pan et al. (2008) with modifications

from Delatorre et al. (2017). The phytohormones analyzed were abscisic acid (ABA), indoleacetic acid

(IAA), 1-aminocyclopropane-1-carboxylic acid (ACC), jasmonic acid (JA), and salicylic acid (SA).

The following deuterated standards were employed: ABA-^{2}H6, IAA-d5, ACC-d4, JA-d5, and SA-d6.

All of them were supplied by Olchemim Ltd. (Czech Republic) except SA-d6 obtained from Sigma

Aldrich (St. Louis, MO, USA). Briefly, 0.05 g fresh weight of leaf samples were homogenized in liquid

N_2 and extracted with 250 µL of extraction buffer (2- propanol/Mili Q water/Hydrochloric acid;

2:1:0.002; v/v/v) in 1.5 mL Eppendorf tubes (Eppendorf Safe Lock England) and agitated by repeated

inversion (60 rpm) for 20 min, at 4 ºC, in the dark. The resulting homogenate was transferred to a

Teflon tube (Thermo Scientific, England) with 450 µL dichloromethane and reextracted by repeated
and inverted agitation for 30 min, at 4 ℃ in the dark. Three phases were obtained, an aqueous, a
material debris and an organic phase. The organic phase (bottom) was recovered, and the intermediate
debris phase was also re-extracted. The combined extracts were evaporated under N_2 flow to remove
propanol and resuspended 150 µL of 100% methanol. Resuspended extracts were filtered through a
0.22 µm cellulose acetate Spin-X centrifuge tube filter (Costar, Corning Inc., Salt Lake City,
USA). Samples were injected into a QTrap4000 LC–ESI($-$)–MS/MS system following the method
described by Segarra et al. (2006) and modified by Llugany et al. (2013) using HPLC Agilent 1100
(Waldrom, Germany) on a Discovery C18 2.1 × 150 mm ID, 5 μm column (Supelco, Bellefonte, USA)
at 50 ℃, a constant flow rate of 0.6 mL min−1 and 10 μL injected volume. MS/MS experiments were
performed on an API 3000 triple quadrupole mass spectrometer (PE Sciex, Concord, Ontario, Canada).
All the analyses were performed using the Turbo Ionspray source in negative ion mode for SA, JA,
ABA, IAA, and PAA and in positive mode for ACC. Quantification was performed by injection of
extracted and spiked samples in multiple reaction monitoring (MRM) mode (n=6 per deme and
treatment).
659
660   *Transcriptome profiling*
661

Total RNA Exaction and RNA-Seq Library Preparation

662
Total RNA was extracted using Maxwell plant RNA kit (Promega Corporation, Madison, WI, USA)
following the manufacturer's instruction. The RNA concentration was measured using Qubit® 2.0
Fluorometer (Invitrogen). 4 μg total RNA was used to prepare RNA-seq libraries using the TruSeq
Stranded mRNA Library Prep Kit (Illumina, San Diego, US). Strand specific paired-end mRNA
sequencing was performed on DNBSEQ platform at the Next-Generation Sequencing Core of the BGI
Genomics Service (Hong Kong, China) (n=3 per deme and treatment).
669

RNA-Seq analysis

670
Bowtie2 (Langmead & Salzberg, 2012) was used to map the clean reads to the reference gene sequence
(transcriptome), and then used RSEM (Li & Dewey, 2011) to calculate the transcript abundance of
each sample. Quality check of raw sequence data was performed using Fastqc. Reads containing
adapters and poly-N and low-quality reads from raw data were removed with Qualimap.
Simultaneously, Q20 (percentage of bases with the quality score greater than 20; sequencing error rate
less than 1%), Q30 (percentage of bases with the quality score greater than 30; sequencing error rate
less than 0.1%), and GC (or guanine-cytosine) content of the clean data were calculated. Paired-end
reads were mapped to the TAIR10 reference genome using the STAR aligner and the .sam output of
STAR was then converted to its compressed format .bam and sorted by gene identifier with samtools.

The overlap of reads with annotation features found in the reference .gtf was calculated using HT-seq
(Putri et al., 2022).
The output computed for each sample (raw read counts) were imported into BGI data visualization
system (https://www.bgi.com/global/service/dr-tom) and parsed using DESeq2 (v 1.8.1). Raw counts
were normalized using DESeq2's function "rlog," which normalises sequences according to library
size to make them comparable among replicates and transforms the original count data to the log scale.
The DEseq2 method is based on the principle of negative binomial distribution (Love et al., 2014).
This method was used to perform differential gene expression analysis (DEG) with an absolute |FC| >
2. The p-values were corrected for multiple error testing according to Benjamini-Hochberg (q-value)
and q-value threshold of 0.05 was set. According to the results of differential gene detection, the R
package *pheatmap* was used to perform hierarchical clustering analysis and extract expression trends
on the union set differential genes.
692
Gene functional annotation
Gene Onthology (GO) functional significant enrichment analysis - used to analyze molecular
functions, cellular components and biological processes - was performed by mapping all candidate
genes to each entry in the Gene Ontology database (http://www.geneontology.org/). A hypergeometric
test was applied to find the GO function significantly enriched in candidate genes compared to all
background genes of the species. p-value was calculated using the basis function phyper
(https://stat.ethz.ch/R-manual/R-devel/library/stats/html/Hypergeometric.html) of R and then
corrected for multiple error testing according to Benjamini-Hochberg
(https://bioconductor.org/packages/release/bioc/html/qvalue.html). Finally, a q-value threshold of
0.05 was set, and the GO term that satisfied this condition was defined as the GO term that was
significantly enriched in candidate genes. Kyoto Encyclopedia of Genes and Genomes (KEGG)
pathway-based enrichment analysis – which links gene sets with a network of interacting molecules in
the cell, such a pathway or complex - was performed following the same methodology and significance
threshold than for GO functional enrichment analysis. Pathways significantly enriched in differentially
expressed genes were identified and visualized through BGI data visualization system
(https://www.bgi.com/global/service/dr-tom)
709
Protein-Protein Interaction (PPI) Network Analysis
Protein–protein Interaction (PPI) networks of differential expressed genes were performed on
STRING version 11.0. Active interaction sources, including text mining, experiments, databases, and
co-expression as well as species limited to "*Arabidopsis thaliana*" and an interaction score > 0.4 were
applied to construct the PPI networks.
WGCNA network analysis

Weighted gene co-expression network analysis (WGCNA) was conducted with the "WGCNA"
package (Langfelder and Horvath, 2008) on BGI's Dr.Tom system (http://biosys.bgi.com). Gene
expression file and trait file were combined, and the soft thresholding power (β value) was filtered
based on the calculation of scale-free topological fit index and mean connectivity. The topological
overlap matrix (TOM) was constructed based on the topological overlap between pairwise genes, and
hierarchical clustering analysis was performed. Cluster dendrogram plot and clustering tree of co-
expression modules of DEGs were created. The co-expression relationships among different modules
were analyzed and modules with high similarity were merged at a similarity threshold of 0.25 and a
minimum module size of 20 genes.
725

Statistical Analysis
All statistical analyses were performed using JMP SAS software (SAS Institute, Cary, NC, United
States). Significant differences for multiple comparisons were determined by one-way or two-way
ANOVA as indicated in figure legends, treatment or treatment relative to control (neuSAL/Control;
alkSAL/Control) and deme (T6^{CS}, Ro2^{CS}, V1^{PC}, LM2^{PC}) or deme origin [coastal-siliceous$_{(T6+Ro2)}$;
intermediate-calcareous$_{(V1+LM2)}$] as independent factors. Comparisons among demes were followed by
post hoc Tukey HSD test (adjusted p-value < 0.05). Multivariate Correlation Analysis (MCA) was
performed to determine the level of association among the measured physiological traits for each deme
and treatment. Pearson's correlation coefficients were calculated for all pairs of measured parameters
under neuSAL and alkSAL at a level of 5% significance. Principal Component Analysis, eigenvalues
and relative proportion of the variance explained by each trait included in MCA were calculated.
737

Acknowledgements. Special thanks to Rosa Padilla for processing ICP and HPLC samples. This
research was supported by the Spanish Ministry of Science and Innovation (MICINN) (PID 2019
104000 RB I00).

Authors contribution. MJ-A performed the lab experiments. MJ-A, LP-M, GE-O and AL-V
performed data analysis. MJ-A, SB, LP-M, AG, ML and CP conceived the study. All authors
contributed to the writing of the manuscript.
745

746 **Declaration of interests.** The authors declare that they have no conflict of interest.
747

748 **References**
749

750 **Achard, P., Cheng, H., De Grauwe, L., Decat, J., Schoutteten, H., Moritz, T., ... & Harberd, N. P.** (2006).
751   Integration of plant responses to environmentally activated phytohormonal signals. *Science*, *311*(5757), 91-94.
752   https://doi.org/10.1126/science.1118642

Aghajanzadeh, T. A., Reich, M., Kopriva, S., & De Kok, L. J. (2018). Impact of chloride (NaCl, KCl) and sulphate (Na2SO4, K2SO4) salinity on glucosinolate metabolism in Brassica rapa. *Journal of Agronomy and Crop Science*, *204*(2), 137-146. https://doi.org/10.1111/jac.12243

Ahuja, I., de Vos, R. C., Bones, A. M., & Hall, R. D. (2010). Plant molecular stress responses face climate change. *Trends in plant science*, *15*(12), 664-674. https://doi.org/10.1016/j.tplants.2010.08.002

Ahuja, I., de Vos, R. C., Bones, A. M., & Hall, R. D. (2010). Plant molecular stress responses face climate change. *Trends in plant science*, *15*(12), 664-674. https://doi.org/10.1104/pp.108.129791

An, Y. M., Song, L. L., Liu, Y. R., Shu, Y. J., & Guo, C. H. (2016). De novo transcriptional analysis of alfalfa in response to saline-alkaline stress. Frontiers in Plant Science, 7, 931. https://doi.org/10.3389/fpls.2016.00931

Anderson, J. P., Badruzsaufari, E., Schenk, P. M., Manners, J. M., Desmond, O. J., Ehlert, C., ... & Kazan, K. (2004). Antagonistic interaction between abscisic acid and jasmonate-ethylene signaling pathways modulates defense gene expression and disease resistance in Arabidopsis. The Plant Cell, 16(12), 3460-3479. https://doi.org/10.1105/tpc.104.025833

Atici, Ö., Ağar, G., & Battal, P. E. Y. A. M. İ. (2005). Changes in phytohormone contents in chickpea seeds germinating under lead or zinc stress. *Biologia plantarum*, *49*(2), 215-222. https://doi.org/10.1007/s10535-005-5222-9

Bakker, E. G., Stahl, E. A., Toomajian, C., Nordborg, M., Kreitman, M., & Bergelson, J. (2006). Distribution of genetic variation within and among local populations of Arabidopsis thaliana over its species range. *Molecular ecology, 15(5),* 1405-1418. https://doi.org/10.1111/j.1365-294X.2006.02884.x

Bao, S., Hua, C., Huang, G., Cheng, P., Gong, X., Shen, L., & Yu, H. (2019). Molecular basis of natural variation in photoperiodic flowering responses. *Developmental cell*, *50*(1), 90-101. https://doi.org/10.1016/j.devcel.2019.05.018

Bomblies, K., Yant, L., Laitinen, R. A., Kim, S. T., Hollister, J. D., Warthmann, N., ... & Weigel, D. (2010). Local-scale patterns of genetic variability, outcrossing, and spatial structure in natural stands of Arabidopsis thaliana. *PLoS genetics, 6(3),* e1000890. https://doi.org/10.1371/journal.pgen.1000890

Burén, S. (2010). *Targeting and function of CAH1: Characterization of a novel protein pathway to the plant cell chloroplast* (Doctoral dissertation, Institutionen för fysiologisk botanik, Umeå universitet). DiVA, id: diva2:284338

Busoms, S., Teres, J., Huang, X. Y., Bomblies, K., Danku, J., Douglas, A., ... & Salt, D. E. (2015). Salinity is an agent of divergent selection driving local adaptation of Arabidopsis to coastal habitats. *Plant Physiology, 168(3),* 915-929.https://doi.org/10.1104/pp.15.00427

Busoms, S., Paajanen, P., Marburger, S., Bray, S., Huang, X. Y., Poschenrieder, C., ... & Salt, D. E. (2018). Fluctuating selection on migrant adaptive sodium transporter alleles in coastal Arabidopsis thaliana. *Proceedings of the National Academy of Sciences, 115(52)*, E12443-E12452. https://doi.org/10.1073/pnas.1816964115

Cackett, L., Cannistraci, C. V., Meier, S., Ferrandi, P., Pěnčík, A., Gehring, C., ... & Donaldson, L. (2022). Salt-specific gene expression reveals elevated auxin levels in Arabidopsis thaliana plants grown under saline conditions. *Frontiers in plant science, 15.* https://doi.org/10.3389/fpls.2022.804716

Cardoso, A. A., Gori, A., Da-Silva, C. J., & Brunetti, C. (2020). Abscisic acid biosynthesis and signaling in plants: Key targets to improve water use efficiency and drought tolerance. *Applied Sciences*, *10*(18), 6322. https://doi.org/10.3390/app10186322

Chen, Q., Bai, L., Wang, W., Shi, H., Ramón Botella, J., Zhan, Q., ... & Song, C. P. (2021). COP1 promotes ABA-induced stomatal closure by modulating the abundance of ABI/HAB and AHG3 phosphatases. *New Phytologist*, *229*(4), 2035-2049. https://doi.org/10.1111/nph.17001

Chen, L., Xiang, S., Chen, Y., Li, D., & Yu, D. (2017). Arabidopsis WRKY45 interacts with the DELLA protein RGL1 to positively regulate age-triggered leaf senescence. *Molecular Plant*, *10*(9), 1174-1189. https://doi.org/10.1016/j.molp.2017.07.008

Chen, G., Zheng, D., Feng, N., Zhou, H., Mu, D., Zhao, L., ... & Huang, A. (2022). Physiological mechanisms of ABA-induced salinity tolerance in leaves and roots of rice. *Scientific Reports, 12(1)*, 8228. https://doi.org/10.1038/s41598-022-11408-0

Cheng, N. H., Liu, J. Z., Nelson, R. S., & Hirschi, K. D. (2004). Characterization of CXIP4, a novel Arabidopsis protein that activates the H+/Ca2+ antiporter, CAX1. *FEBS letters*, *559*(1-3), 99-106. https://doi.org/10.1016/S0014-5793(04)00036-5

Che-Othman, M. H., Millar, A. H., & Taylor, N. L. (2017). Connecting salt stress signalling pathways with salinity-induced changes in mitochondrial metabolic processes in C 3 plants. *Plant, cell & environment, 40(12)*, 2875-2905. https://doi.org/10.1104/pp.106.085555

Chew, Y. H., & Halliday, K. J. (2011). A stress-free walk from Arabidopsis to crops. *Current Opinion in Biotechnology*, *22*(2), 281-286. https://doi.org/10.1016/j.copbio.2010.11.011

Clark, T. J., & Lu, Y. (2015). Analysis of loss-of-function mutants in aspartate kinase and homoserine dehydrogenase genes points to complexity in the regulation of aspartate-derived amino acid contents. *Plant Physiology*, *168*(4), 1512-1526. https://doi.org/10.1104/pp.15.00364

Colebrook, E. H., Thomas, S. G., Phillips, A. L., & Hedden, P. (2014). The role of gibberellin signalling in plant responses to abiotic stress. *Journal of experimental biology*, *217*(1), 67-75. https://doi.org/10.1242/jeb.089938

Coolen, S., Proietti, S., Hickman, R., Davila Olivas, N. H., Huang, P. P., Van Verk, M. C., ... & Van Wees, S. C. (2016). Transcriptome dynamics of Arabidopsis during sequential biotic and abiotic stresses. *The Plant Journal*, *86*(3), 249-267. https://doi.org/10.1111/tpj.13167

Dąbrowska-Bronk, J., Komar, D. N., Rusaczonek, A., Kozłowska-Makulska, A., Szechyńska-Hebda, M., & Karpiński, S. (2016). β-carbonic anhydrases and carbonic ions uptake positively influence Arabidopsis photosynthesis, oxidative stress tolerance and growth in light dependent manner. *Journal of plant physiology*, *203*, 44-54. https://doi.org/10.1016/j.jplph.2016.05.013

Davila Olivas, N. H., Coolen, S., Huang, P., Severing, E., van Verk, M. C., Hickman, R., ... & Dicke, M. (2016). Effect of prior drought and pathogen stress on *Arabidopsis* transcriptome changes to caterpillar herbivory. *New Phytologist*, *210*(4), 1344-1356. https://doi.org/10.1111/nph.13847

de Kraker, J. W., Luck, K., Textor, S., Tokuhisa, J. G., & Gershenzon, J. (2007). Two Arabidopsis genes (IPMS1 and IPMS2) encode isopropylmalate synthase, the branchpoint step in the biosynthesis of leucine. *Plant physiology, 143(2)*, 970–986. https://doi.org/10.1104/pp.106.085555

Dellero, Y., Jossier, M., Schmitz, J., Maurino, V. G., & Hodges, M. (2016). Photorespiratory glycolate–glyoxylate metabolism. *Journal of Experimental Botany*, *67*(10), 3041-3052. https://doi.org/10.1093/jxb/erw090

Domergue, F., Vishwanath, S. J., Joubès, J., Ono, J., Lee, J. A., Bourdon, M., ... & Rowland, O. (2010). Three Arabidopsis fatty acyl-coenzyme A reductases, FAR1, FAR4, and FAR5, generate primary fatty alcohols associated with suberin deposition. Plant physiology, 153(4), 1539-1554. https://doi.org/10.1104/pp.110.158238

Fahey, J. W., Zalcmann, A. T., & Talalay, P. (2001). The chemical diversity and distribution of glucosinolates and isothiocyanates among plants. *Phytochemistry*, *56*(1), 5-51. https://doi.org/10.1016/S0031-9422(00)00316-2

Fang, S., Hou, X., & Liang, X. (2021). Response mechanisms of plants under saline-alkali stress. Frontiers in Plant Science, 12, 667458. https://doi.org/10.3389/fpls.2021.667458

Farooq, S., Ahmad, S., Hussain, S., & Hussain, M. (2018). Nutrient homeostasis and salt stress tolerance. *In Plant Nutrients and Abiotic Stress Tolerance* (pp. 391-413). Springer, Singapore. https://doi.org/10.1007/978-981-10-9044-8_17

Flügel, F., Timm, S., Arrivault, S., Florian, A., Stitt, M., Fernie, A. R., & Bauwe, H. (2017). The photorespiratory metabolite 2-phosphoglycolate regulates photosynthesis and starch accumulation in Arabidopsis. *The Plant Cell*, *29*(10), 2537-2551. https://doi.org/10.1105/tpc.17.00256

Forieri, I., Sticht, C., Reichelt, M., Gretz, N., Hawkesford, M. J., Malagoli, M., ... & Hell, R. (2017). System analysis of metabolism and the transcriptome in Arabidopsis thaliana roots reveals differential co-regulation upon iron, sulfur and potassium deficiency. *Plant, cell & environment*, *40*(1), 95-107. https://doi.org/10.1111/pce.12842

Gan, X., Stegle, O., Behr, J., Steffen, J. G., Drewe, P., Hildebrand, K. L., ... & Mott, R. (2011). Multiple reference genomes and transcriptomes for *Arabidopsis thaliana*. *Nature*, *477*(7365), 419-423. https://doi.org/10.1038/nature10414

Gan, T., Lin, Z., Bao, L., Hui, T., Cui, X., Huang, Y., ... & Qian, Y. (2021). Comparative proteomic analysis of tolerant and sensitive varieties reveals that phenylpropanoid biosynthesis contributes to salt tolerance in mulberry. *International journal of molecular sciences*, *22*(17), 9402. https://doi.org/10.3390/ijms22179402

Garg, N., & Pandey, R. (2015). Effectiveness of native and exotic arbuscular mycorrhizal fungi on nutrient uptake and ion homeostasis in salt-stressed Cajanus cajan L.(Millsp.) genotypes. *Mycorrhiza*, *25*(3), 165-180. https://doi.org/10.1007/s00572-014-0600-9

Georgii, E., Jin, M., Zhao, J., Kanawati, B., Schmitt-Kopplin, P., Albert, A., ... & Schäffner, A. R. (2017). Relationships between drought, heat and air humidity responses revealed by transcriptome-metabolome co-analysis. *BMC plant biology*, *17*(1), 1-23. https://doi.org/10.1186/s12870-017-1062-y

Guo, R., Shi, L., Ding, X., Hu, Y., Tian, S., Yan, D., ... & Yang, Y. (2010). Effects of saline and alkaline stress on germination, seedling growth, and ion balance in wheat. Agronomy journal, 102(4), 1252-1260. https://doi.org/10.2134/agronj2010.0022

Guo, M., Li, H., Li, L., Cheng, X., Gao, W., Xu, Y., ... & Liu, X. (2014). Comparative proteomic analysis of Arabidopsis thaliana roots between wild type and its salt-tolerant mutant. *Journal of Plant Interactions*, *9*(1), 330-337. https://doi.org/10.1080/17429145.2013.833653

Guo, R., Shi, L., Yan, C., Zhong, X., Gu, F., Liu, Q., ... & Li, H. (2017). Ionomic and metabolic responses to neutral salt or alkaline salt stresses in maize (Zea mays L.) seedlings. *BMC Plant Biology*, *17*(1), 1-13. https://doi.org/10.1186/s12870-017-0994-6

Guo, L., Wang, J., Li, M., Liu, L., Xu, J., Cheng, J., ... & Luedeling, E. (2019). Distribution margins as natural laboratories to infer species' flowering responses to climate warming and implications for frost risk. *Agricultural and Forest Meteorology*, *268*, 299-307. https://doi.org/10.1016/j.agrformet.2019.01.038

Gupta, A., Hisano, H., Hojo, Y., Matsuura, T., Ikeda, Y., Mori, I. C., & Senthil-Kumar, M. (2017). Global profiling of phytohormone dynamics during combined drought and pathogen stress in *Arabidopsis thaliana* reveals ABA and JA as major regulators. *Scientific reports*, *7*(1), 1-13. https://doi.org/10.1038/s41598-017-03907-2

Gupta, A., Sarkar, A. K., & Senthil-Kumar, M. (2016). Global transcriptional analysis reveals unique and shared responses in *Arabidopsis thaliana* exposed to combined drought and pathogen stress. *Frontiers in plant science*, *7*, 686. https://doi.org/10.3389/fpls.2016.00686

Han, H., Wang, Q., Wei, L., Liang, Y., Dai, J., Xia, G., & Liu, S. (2018). Small RNA and degradome sequencing used to elucidate the basis of tolerance to salinity and alkalinity in wheat. *BMC plant biology*, *18*(1), 1-17. https://doi.org/10.1186/s12870-018-1415-1

Hao, X., Li, J., Gao, S., Tuerxun, Z., Chang, X., Hu, W., ... & Huang, Q. (2020). SsPsaH, a H subunit of the photosystem I reaction center of Suaeda salsa, confers the capacity of osmotic adjustment in tobacco. *Genes & Genomics*, *42*(12), 1455-1465. https://doi.org/10.1007/s13258-020-00970-4

Heidari, P., Entazari, M., Ebrahimi, A., Ahmadizadeh, M., Vannozzi, A., Palumbo, F., & Barcaccia, G. (2021). Exogenous EBR ameliorates endogenous hormone contents in tomato species under low-temperature stress. *Horticulturae*, *7*(4), 84. https://doi.org/10.3390/horticulturae7040084

Hirayama, T., & Shinozaki, K. (2010). Research on plant abiotic stress responses in the post-genome era: past, present and future. *The Plant Journal*, *61*(6), 1041-1052. https://doi.org/10.1111/j.1365-313X.2010.04124.x

Hosseini, S. Z., Ismaili, A., Nazarian-Firouzabadi, F., Fallahi, H., Nejad, A. R., & Sohrabi, S. S. (2021). Dissecting the molecular responses of lentil to individual and combined drought and heat stresses by comparative transcriptomic analysis. *Genomics*, *113*(2), 693-705.https://doi.org/10.1016/j.ygeno.2020.12.038

Hsieh, E. J., Cheng, M. C., & Lin, T. P. (2013). Functional characterization of an abiotic stress-inducible transcription factor AtERF53 in Arabidopsis thaliana. Plant molecular biology, 82(3), 223-237. https://doi.org/10.1007/s11103-013-0054-z

Hsieh, E. J., & Waters, B. M. (2016). Alkaline stress and iron deficiency regulate iron uptake and riboflavin synthesis gene expression differently in root and leaf tissue: implications for iron deficiency chlorosis. *Journal of Experimental Botany, 67(19),* 5671-5685. https://doi.org/10.1093/jxb/erw328

Hussain, A., Shah, M., Hamayun, M., Iqbal, A., Murad, W., Irshad, M., ... & Kim, H. Y. (2021). *Pseudocitrobacter anthropi* reduces heavy metal uptake and improves phytohormones and antioxidant system in *Glycine max L. World Journal of Microbiology and Biotechnology*, *37*(11), 1-19. https://doi.org/10.1007/s11274-021-03156-6

Hussain, Q., Asim, M., Zhang, R., Khan, R., Farooq, S., & Wu, J. (2021). Transcription factors interact with ABA through gene expression and signaling pathways to mitigate drought and salinity stress. *Biomolecules, 11(8)*, 1159. https://doi.org/10.3390/biom11081159

Hussien, S. A., & Doosh, K. (2021, May). Extraction, Purification and Characterization of β-Galactosidase from Tomato (Lycopersicom esculentum). In *IOP Conference Series: Earth and Environmental Science* (Vol. 761, No. 1, p. 012123). IOP Publishing. https://doi.org/10.1088/1755-1315/761/1/012123

Jin, H., Plaha, P., Park, J. Y., Hong, C. P., Lee, I. S., Yang, Z. H., ... & Lim, Y. P. (2006). Comparative EST profiles of leaf and root of Leymus chinensis, a xerophilous grass adapted to high pH sodic soil. Plant Science, 170(6), 1081-1086. https://doi.org/10.1016/j.plantsci.2006.01.002

Jing, Y., Shi, L., Li, X., Zheng, H., Gao, J., Wang, M., ... & Zhang, W. (2019). OXS2 is required for salt tolerance mainly through associating with salt Inducible genes, CA1 and Araport11, in Arabidopsis. *Scientific reports*, *9*(1), 1-11. https://doi.org/10.1038/s41598-019-56456-1

Julkowska, M. M., & Testerink, C. (2015). Tuning plant signaling and growth to survive salt. Trends in plant science, 20(9), 586-594. https://doi.org/10.1016/j.tplants.2015.06.008

Kazan, K., & Manners, J. M. (2012). JAZ repressors and the orchestration of phytohormone crosstalk. *Trends in plant science*, *17*(1), 22-31. https://doi.org/10.1016/j.tplants.2011.10.006

Kazan, K., & Manners, J. M. (2013). MYC2: the master in action. Molecular plant, 6(3), 686-703.

Khare, R., Dhar, Y. V., Sandhu, G., Singh, S., Kumar, S., Khan, A., ... & Trivedi, P. K. (2022). Genome-wide
expression and variation in nucleotide sequences lead to differential response of Arabidopsis thaliana ecotypes
towards arsenic stress under sulfur limiting condition. *Environmental and Experimental Botany*, *195*, 104764.
https://doi.org/10.1016/j.envexpbot.2021.104764

Kotake, T., Dina, S., Konishi, T., Kaneko, S., Igarashi, K., Samejima, M., ... & Tsumuraya, Y. (2005). Molecular
cloning of a β-galactosidase from radish that specifically hydrolyzes β-(1→ 3)-and β-(1→ 6)-galactosyl residues
of arabinogalactan protein. *Plant Physiology*, *138*(3), 1563-1576. https://doi.org/10.1104/pp.105.062562

Kroh, G. E., & Pilon, M. (2020). Regulation of iron homeostasis and use in chloroplasts. *International Journal of*
Molecular Sciences, *21*(9), 3395. https://doi.org/10.3390/ijms21093395

Kubra, G., Khan, M., Munir, F., Gul, A., Shah, T., Hussain, A., ... & Amir, R. (2021). Expression characterization
of flavonoid biosynthetic pathway genes and transcription factors in peanut under water deficit
conditions. *Frontiers in Plant Science*, *12*, 680368. https://doi.org/10.3389/fpls.2021.680368

Kumar, D., Datta, R., Hazra, S., Sultana, A., Mukhopadhyay, R., & Chattopadhyay, S. (2015). Transcriptomic
profiling of *Arabidopsis thaliana* mutant pad 2.1 in response to combined cold and osmotic stress. *PLoS One*,
10(3), e0122690. https://doi.org/10.1371/journal.pone.0122690

Kumar, D., Hazra, S., Datta, R., & Chattopadhyay, S. (2016). Transcriptome analysis of *Arabidopsis* mutants
suggests a crosstalk between ABA, ethylene and GSH against combined cold and osmotic stress. *Scientific reports*,
6(1), 1-13. https://doi.org/10.1038/srep36867

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature methods, 9(4)*, 357-359.
https://doi.org/10.1038/nmeth.1923

Lee, E. J., Matsumura, Y., Soga, K., Hoson, T., & Koizumi, N. (2007). Glycosyl hydrolases of cell wall are induced
by sugar starvation in Arabidopsis. *Plant and cell physiology*, *48*(3), 405-413.**Li, B., & Dewey, C. N.** (2011).
RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC*
bioinformatics, *12*(1), 1-16. https://doi.org/10.1093/pcp/pcm009

Lee, H. Y., Chen, Y. C., Kieber, J. J., & Yoon, G. M. (2017). Regulation of the turnover of ACC synthases by
phytohormones and heterodimerization in *Arabidopsis*. *The Plant Journal*, *91*(3), 491-504.
https://doi.org/10.1111/tpj.13585

Leyva-Perez, M. D. L. O., Valverde-Corredor, A., Valderrama, R., Jiménez-Ruiz, J., Muñoz-Merida, A.,
Trelles, O., ... & Luque, F. (2015). Early and delayed long-term transcriptional changes and short-term transient
responses during cold acclimation in olive leaves. *DNA research, 22(1)*, 1-11.
https://doi.org/10.1093/dnares/dsu033

Li, N., Liu, H., Sun, J., Zheng, H., Wang, J., Yang, L., ... & Zou, D. (2018). Transcriptome analysis of two
contrasting rice cultivars during alkaline stress. *Scientific reports, 8(1),* 1-16. https://doi.org/10.1038/s41598-018-
27940-x

Liu, Q., Zhang, Y. C., Wang, C. Y., Luo, Y. C., Huang, Q. J., Chen, S. Y., ... & Chen, Y. Q. (2009). Expression
analysis of phytohormone-regulated microRNAs in rice, implying their regulation roles in plant hormone signaling.
FEBS letters, *583*(4), 723-728. https://doi.org/10.1016/j.febslet.2009.01.020

Liu, P., Huang, R., Hu, X., Jia, Y., Li, J., Luo, J., ... & Chen, Z. (2019). Physiological responses and proteomic
changes reveal insights into Stylosanthes response to manganese toxicity. *BMC plant biology*, *19*(1), 1-21.
https://doi.org/10.1186/s12870-019-1822-y

Liu, L., & Wang, B. (2021). Protection of halophytes and their uses for cultivation of saline-alkali soil in China.
Biology, 10(5), 353. https://doi.org/10.3390/biology10050353

Llanes, A., Andrade, A., Masciarelli, O., Alemano, S., & Luna, V. (2016). Drought and salinity alter endogenous
hormonal profiles at the seed germination phase. *Seed Science Research*, *26*(1), 1-13.
https://doi.org/10.1017/S0960258515000331

Llugany, M., Martin, S. R., Barceló, J., & Poschenrieder, C. (2013). Endogenous jasmonic and salicylic acids
levels in the Cd-hyperaccumulator *Noccaea (Thlaspi) praecox* exposed to fungal infection and/or mechanical
stress. Plant cell reports, 32(8), 1243-1249. https://doi.org/10.1007/s00299-013-1427-0

Love, M.I., Huber, W. & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. *Genome Biol 15*, 550. https://doi.org/10.1186/s13059-014-0550-8

Lovelli, S., Scopa, A., Perniola, M., Di Tommaso, T., & Sofo, A. (2012). Abscisic acid root and leaf concentration
in relation to biomass partitioning in salinized tomato plants. Journal of plant physiology, 169(3), 226-233.
https://doi.org/10.1016/j.jplph.2011.09.009

Ma, H. & Wang, P. (2021). Regulation of plant responses to salt stress. International *Journal of Molecular Sciences,*
22(9), 4609. https://doi.org/10.3389/fpls.2019.00498

Mao, B., Yin, H., Wang, Y., Zhao, T. H., Tian, R. R., Wang, W., & Ye, J. S. (2017). Combined effects of O3 and
UV radiation on secondary metabolites and endogenous hormones of soybean leaves. *PloS one*, *12*(*8*), e0183147.
https://doi.org/10.1371/journal.pone.0183147

Mao, F., Nan, G., Cao, M., Gao, Y., Guo, L., Meng, X., & Yang, G. (2018). The metal distribution and the change
of physiological and biochemical process in soybean and mung bean plants under heavy metal stress. *International*
journal of phytoremediation, *20*(11), 1113-1120. https://doi.org/10.1080/15226514.2017.1365346

Marconi, G., Pace, R., Traini, A., Raggi, L., Lutts, S., Chiusano, M., ... & Albertini, E. (2013). Use of MSAP
markers to analyse the effects of salt stress on DNA methylation in rapeseed (Brassica napus var. oleifera). *PloS*
one, 8(9), e75597. https://doi.org/10.1371/journal.pone.0075597

Markakis, M. N., Boron, A. K., Van Loock, B., Saini, K., Cirera, S., Verbelen, J. P., & Vissenberg, K. (2013).
Characterization of a small auxin-up RNA (SAUR)-like gene involved in Arabidopsis thaliana development. *PloS*
one, 8(11), e82596. https://doi.org/10.1371/journal.pone.0082596

Martos, S., Gallego, B., Cabot, C., Llugany, M., Barceló, J., & Poschenrieder, C. (2016). Zinc triggers signaling
mechanisms and defense responses promoting resistance to *Alternaria brassicicola* in *Arabidopsis thaliana*. *Plant*
Science, 249, 13-24. https://doi.org/10.1016/j.plantsci.2016.05.001

Maxwell K., Johnson G.N. (2000). Chlorophyll fluorescence - a practical guide. *J. Exp. Bot. 51(345)*: 659-668.
https://doi.org/10.1093/jexbot/51.345.659

McClellan, C. A., & Chang, C. (2008). The role of protein turnover in ethylene biosynthesis and response. *Plant*
Science, 175(1-2), 24-31. https://doi.org/10.1016/j.plantsci.2008.01.004

Mehmood, E., R. Kausar, M. Akram and S.M. Shahzad. (2009). Is boron required to improve rice growth and yield
in saline environment. *Pak. J. Bot., 41(*3):1339-1350.

Meng, C., Quan, TY., Li, ZY. et al. (2017). Transcriptome profiling reveals the genetic basis of alkalinity tolerance
in wheat. *BMC Genomics* 18-24.   https://doi.org/10.1186/s12864-016-3421-8

Moeder, W., Ung, H., Mosher, S., & Yoshioka, K. (2010). SA-ABA antagonism in defense responses. *Plant*
signaling & behavior, *5*(10), 1231-1233. https://doi.org/10.4161/psb.5.10.12836

Moneo-Sánchez, M., Izquierdo, L., Martín, I., Labrador, E., & Dopico, B. (2016). Subcellular location of
Arabidopsis thaliana subfamily a1 β-galactosidases and developmental regulation of transcript levels of their
coding genes. *Plant Physiology and Biochemistry*, *109*, 137-145. https://doi.org/10.1016/j.plaphy.2016.09.016

Munns, R. (1993). Physiological processes limiting plant growth in saline soils: some dogmas and hypotheses. *Plant,*
Cell & Environment, 16(1), 15-24. https://doi.org/10.1111/j.1365-3040.1993.tb00840.x

Munns, R., & Tester, M. (2008). Mechanisms of salinity tolerance. Annual review of plant biology, 59, 651.
https://doi.org/10.1146/annurev.arplant.59.032607.092911

Naing, A. H., & Kim, C. K. (2021). Abiotic stress-induced anthocyanins in plants: Their role in tolerance to abiotic
stresses. Physiologia Plantarum, 172(3), 1711-1723. https://doi.org/10.1111/ppl.13373

Nam, Y. J., Herman, D., Blomme, J., Chae, E., Kojima, M., Coppens, F., ... & Gonzalez, N. (2017). Natural
variation of molecular and morphological gibberellin responses. *Plant Physiology*, *173*(1), 703-714.
https://doi.org/10.1104/pp.16.01626

Pai, A. A., Pritchard, J. K., & Gilad, Y. (2015). The genetic and mechanistic basis for variation in gene
regulation. *PLoS genetics*, *11*(1), e1004857. https://doi.org/10.1016/j.plaphy.2016.09.016

Pandey, A., Khan, M. K., Hakki, E. E., Gezgin, S., & Hamurcu, M. (2019). Combined boron toxicity and salinity
stress - An insight into its interaction in plants. *Plants, 8(10)*, 364. https://doi.org/10.3390/plants8100364

Peixoto, B., Moraes, T. A., Mengin, V., Margalha, L., Vicente, R., Feil, R., ... & Baena-González, E. (2021).
Impact of the SnRK1 protein kinase on sucrose homeostasis and the transcriptome during the diel cycle. *Plant*
physiology, *187*(3), 1357-1373. https://doi.org/10.1093/plphys/kiab350

Perea-Resa, C., Rodríguez-Milla, M. A., Iniesto, E., Rubio, V., & Salinas, J. (2017). Prefoldins negatively regulate
cold acclimation in Arabidopsis thaliana by promoting nuclear proteasome-mediated HY5 degradation. Molecular
plant, 10(6), 791-804. https://doi.org/10.1016/j.molp.2017.03.012

Pérez-Martín, L., Busoms, S., Tolrà, R., & Poschenrieder, C. (2021). Transcriptomics reveals fast changes in
salicylate and jasmonate signaling pathways in shoots of carbonate-tolerant Arabidopsis thaliana under bicarbonate
exposure. International journal of molecular sciences, 22(3), 1226. https://doi.org/10.3390/ijms22031226

Pérez-Martín, L., Busoms, S., Almira, M. J., Azagury, N., Terés, J., Tolrà, R., ... & Barcelo, J. (2022). Evolution
of salt tolerance in Arabidopsis thaliana on siliceous soils does not confer tolerance to saline calcareous soils. Plant
and Soil, 1-21. https://doi.org/10.1007/s11104-022-05439-9

Piazzi, M., Bavelloni, A., Gallo, A., Faenza, I., & Blalock, W. L. (2019). Signal transduction in ribosome biogenesis:
a recipe to avoid disaster. International journal of molecular sciences, 20(11), 2718.
https://doi.org/10.3390/ijms20112718

Prerostova, S., Dobrev, P. I., Gaudinova, A., Hosek, P., Soudek, P., Knirsch, V., & Vankova, R. (2017). Hormonal
dynamics during salt stress responses of salt-sensitive *Arabidopsis thaliana* and salt-tolerant *Thellungiella*
salsuginea. *Plant Science*, *264*, 188-198. https://doi.org/10.1016/j.plantsci.2017.07.020

Price, L., Han, Y., Angessa, T., & Li, C. (2022). Molecular Pathways of WRKY Genes in Regulating Plant Salinity
Tolerance. *International Journal of Molecular Sciences*, *23*(18), 10947. https://doi.org/10.3390/ijms231810947

Putri, G. H., Anders, S., Pyl, P. T., Pimanda, J. E., & Zanini, F. (2022). Analysing high-throughput sequencing
data in Python with HTSeq 2.0. *Bioinformatics, 38(10)*, 2943-2945. https://doi.org/10.1093/bioinformatics/btac166

Rasmussen, S., Barah, P., Suarez-Rodriguez, M. C., Bressendorff, S., Friis, P., Costantino, P., ... & Mundy, J.
(2013). Transcriptome responses to combinations of stresses in Arabidopsis. *Plant physiology*, *161*(4), 1783-1794.
https://doi.org/10.1104/pp.112.210773

Riemann, M., Dhakarey, R., Hazman, M., Miro, B., Kohli, A., & Nick, P. (2015). Exploring jasmonates in the
hormonal network of drought and salinity responses. *Frontiers in plant science*, *6*, 1077.
https://doi.org/10.3389/fpls.2015.01077

Ross, G. S., Wegrzyn, T., MacRae, E. A., & Redgwell, R. J. (1994). Apple [beta]-Galactosidase (Activity against
Cell Wall Polysaccharides and Characterization of a Related cDNA Clone). *Plant Physiology*, *106*(2), 521-528.
https://doi.org/10.1104/pp.106.2.521

Rus, A., Baxter, I., Muthukumar, B., Gustin, J., Lahner, B., Yakubova, E., & Salt, D. E. (2006). Natural variants
of At HKT1 enhance Na+ accumulation in two wild populations of Arabidopsis. *PLoS genetics*, *2*(12), e210.
https://doi.org/10.1371/journal.pgen.0020210

Sah, S. K., Reddy, K. R., & Li, J. (2016). Abscisic acid and abiotic stress tolerance in crop plants. *Frontiers in plant*
science, *7*, 571. https://doi.org/10.3389/fpls.2016.00571

Saini, S., Kaur, N., & Pati, P. K. (2021). Phytohormones: Key players in the modulation of heavy metal stress
tolerance in plants. *Ecotoxicology and Environmental Safety*, *223*,
112578.https://doi.org/10.1016/j.ecoenv.2021.112578

Sazegari, S., Niazi, A., & Ahmadi, F. S. (2015). A study on the regulatory network with promoter analysis for
Arabidopsis DREB-genes. Bioinformation, 11(2), 101. https://doi.org/10.6026/97320630011101

Segarra, G., Jáuregui, O., Casanova, E., & Trillas, I. (2006). Simultaneous quantitative LC–ESI-MS/MS analyses
of salicylic acid and jasmonic acid in crude extracts of Cucumis sativus under biotic stress. *Phytochemistry, 67(4),*
395-401. https://doi.org/10.1016/j.phytochem.2005.11.017

Shabala, S., & Cuin, T. A. (2008). Potassium transport and plant salt tolerance. *Physiologia plantarum, 133(4),* 651-
669. https://doi.org/10.1111/j.1399-3054.2007.01008.x

Sharma, R., Singh, G., Bhattacharya, S., & Singh, A. (2018). Comparative transcriptome meta-analysis of
Arabidopsis thaliana under drought and cold stress. *PLoS one*, *13*(9),
e0203266.https://doi.org/10.1371/journal.pone.0203266

Shu, J., Liu, Y., Zhang, L., Li, Z., Fang, Z., Yang, L., ... & Lv, H. (2018). QTL-seq for rapid identification of
candidate genes for flowering time in broccoli× cabbage. *Theoretical and applied genetics*, *131*(4), 917-928.
https://doi.org/10.1007/s00122-017-3047-5

Sibole, J. V., Cabot, C., Michalke, W., Poschenrieder, C., & Barceló, J. (2005). Relationship between expression
of the PM H+-ATPase, growth and ion partitioning in the leaves of salt-treated Medicago species. *Planta, 221(4),*
557-566. https://doi.org/10.1007/s00425-004-1456-6

Singh, A. (2021). Soil salinization management for sustainable development: A review. *Journal of environmental*
management, 277, 111383. https://doi.org/10.1016/j.jenvman.2020.111383

Skalak, J., Nicolas, K. L., Vankova, R., & Hejatko, J. (2021). Signal integration in plant abiotic stress responses
via multistep phosphorelay signaling. *Frontiers in Plant Science, 12, 644823.*
https://doi.org/10.3389/fpls.2021.644

Smith, D. L., & Gross, K. C. (2000). A family of at least seven β-galactosidase genes is expressed during tomato
fruit development. *Plant Physiology*, *123*(3), 1173-1184. https://doi.org/10.1104/pp.123.3.1173

Song, Y., Lv, J., Ma, Z., & Dong, W. (2019). The mechanism of alfalfa (Medicago sativa L.) response to abiotic
stress. *Plant Growth Regulation*, *89*(3), 239-249. https://doi.org/10.1007/s10725-019-00530-1

Spartz, A. K., Ren, H., Park, M. Y., Grandt, K. N., Lee, S. H., Murphy, A. S., ... & Gray, W. M. (2014). SAUR
inhibition of PP2C-D phosphatases activates plasma membrane H+-ATPases to promote cell expansion in
Arabidopsis. *The Plant Cell, 26(5),* 2129-2142. https://doi.org/10.1105/tpc.114.126037

Sun, Y., Hao, P., Lv, X., Tian, J., Wang, Y., Zhang, X., ... & Wu, T. (2020). A long non-coding apple RNA,
MSTRG. 85814.11, acts as a transcriptional enhancer of SAUR32 and contributes to the Fe-deficiency
response. *The Plant Journal*, *103*(1), 53-67. https://doi.org/10.1111/tpj.14706

Suzuki, N., Bassil, E., Hamilton, J. S., Inupakutika, M. A., Zandalinas, S. I., Tripathy, D., ... & Mittler, R.
(2016). ABA is required for plant acclimation to a combination of salt and heat stress. *PloS one*, *11*(1),
e0147625. https://doi.org/10.1371/journal.pone.0147625

Tang, X., Chang, L., Wu, S., Li, P., Liu, G., & Wang, N. N. (2008). Auto-regulation of the promoter activities of
Arabidopsis 1-aminocyclopropane-1-carboxylate synthase genes AtACS4, AtACS5, and AtACS7 in response to
different plant hormones. *Plant Science*, *175*(1-2), 161-167.https://doi.org/10.1016/j.plantsci.2008.01.002

Terés, J., Busoms, S., Perez Martín, L., Luís-Villarroya, A., Flis, P., Álvarez-Fernández, A., ... &
Poschenrieder, C. (2019). Soil carbonate drives local adaptation in Arabidopsis thaliana. *Plant, cell &*
environment, 42(8), 2384-2398. https://doi.org/10.1111/pce.13567

Torun, H., Novák, O., Mikulík, J., Strnad, M., & Ayaz, F. A. (2022). The Effects of exogenous salicylic acid on
endogenous phytohormone status in Hordeum vulgare L. under salt stress. *Plants*, *11*(5),
618.https://doi.org/10.3390/plants11050618

Tsai, H. H., & Schmidt, W. (2020). pH-dependent transcriptional profile changes in iron-deficient Arabidopsis
roots. *BMC genomics*, *21*(1), 1-11. https://doi.org/10.1186/s12864-020-07116-6

Van Veen, H., Vashisht, D., Akman, M., Girke, T., Mustroph, A., Reinen, E., ... & Sasidharan, R. (2016).
Transcriptomes of eight *Arabidopsis thaliana* accessions reveal core conserved, genotype-and organ-specific
responses to flooding stress. *Plant Physiology*, *172*(2), 668-689. https://doi.org/10.1104/pp.16.00472

Wang, X. S., & Han, J. G. (2007). Effects of NaCl and silicon on ion distribution in the roots, shoots and leaves of
two alfalfa cultivars with different salt tolerance. *Soil Science & Plant Nutrition, 53(3),* 278-285.
https://doi.org/10.1111/j.1747-0765.2007.00135.x

Wang, N., Fan, X., Lin, Y., Li, Z., Wang, Y., Zhou, Y., ... & Ma, J. (2022). Alkaline stress induces different
physiological, hormonal and gene expression responses in diploid and autotetraploid rice. *International Journal*
of Molecular Sciences, 23(10), 5561. https://doi.org/10.3390/ijms23105561

Waters, B. M., Amundsen, K., & Graef, G. (2018). Gene expression profiling of iron deficiency chlorosis sensitive
and tolerant soybean indicates key roles for phenylpropanoids under alkalinity stress. Frontiers in plant science,
9, 10. https://doi.org/10.3389/fpls.2018.00010

Wen, R., Lv, H. N., Jiang, Y., & Tu, P. F. (2018). Anti-inflammatory flavone and chalcone derivatives from the
roots of Pongamia pinnata (L.) Pierre. *Phytochemistry*, *149*, 56-63.
https://doi.org/10.1016/j.phytochem.2018.02.005

Wittstock, U., & Halkier, B. A. (2002). Glucosinolate research in the Arabidopsis era. *Trends in plant science*, *7*(6),
263-270. https://doi.org/10.1016/S1360-1385(02)02273-2

Xie, Z., Nolan, T. M., Jiang, H., & Yin, Y. (2019). AP2/ERF transcription factor regulatory networks in hormone
and abiotic stress responses in Arabidopsis. *Frontiers in plant science*, *10*, 228.
https://doi.org/10.3389/fpls.2019.00228

Yan, F., Zhu, Y., Zhao, Y., Wang, Y., Li, J., Wang, Q., & Liu, Y. (2020). De novo transcriptome sequencing and
analysis of salt-, alkali-, and drought-responsive genes in Sophora alopecuroides. BMC genomics, 21(1), 1-15.
https://doi.org/10.1186/s12864-020-06823-4

Yang, H., Wu, Y., Duan, Y., Zhang, C., Huang, Z., Wu, W., ... & Li, W. (2022). Metabolomics combined with
physiological and transcriptomic analyses reveal regulatory features associated with blueberry growth in different
soilless substrates. *Scientia Horticulturae*, *302*, 111145. https://doi.org/10.1016/j.scienta.2022.111145

Yang, K., Yang, J., & Yi, J. (2018). Nucleolar Stress: hallmarks, sensing mechanism and diseases. *Cell Stress*, *2*(6),
125. https://doi.org/10.15698/cst2018.06.139

Yousefi, H., Dalir, N., Rahnemaie, R., & Babaei, A. (2020). The alleviation of salinity-induced stress by using
boron in soilless grown rose. *Journal of plant nutrition*, *43*(4), 526-537.
https://doi.org/10.1080/01904167.2019.1685103

Yu, Y., Li, Y., Yan, Z., & Duan, X. (2021). The role of cytokinins in plant under salt stress. Journal of Plant
Growth Regulation, 1-13. https://doi.org/10.1007/s00344-021-10441-z

Zandalinas, S. I., Sengupta, S., Fritschi, F. B., Azad, R. K., Nechushtai, R., & Mittler, R. (2021). The impact of
multifactorial stress combination on plant growth and survival. New Phytologist, 230(3), 1034-1048.
https://doi.org/10.1111/nph.17232

Zhang, L., Chen, L., & Yu, D. (2018). Transcription factor WRKY75 interacts with DELLA proteins to affect
flowering. Plant Physiology, 176(1), 790-803. https://doi.org/10.1104/pp.17.00657

Zhang, B., Chen, X., Lu, X., Shu, N., Wang, X., Yang, X., ... & Ye, W. (2018). Transcriptome analysis of
Gossypium hirsutum L. reveals different mechanisms among NaCl, NaOH and Na2CO3 stress tolerance.
Scientific reports, 8(1), 1-14. https://doi.org/10.1038/s41598-018-31668-z
Zhang, W., Wang, J., Huang, Z., Mi, L., Xu, K., Wu, J., ... & Jiang, D. (2019). Effects of low temperature at
booting stage on sucrose metabolism and endogenous hormone contents in winter wheat spikelet. *Frontiers in*
Plant Science, *10*, 498.
Zhang, Q., Zhai, J., Shao, L., Lin, W., & Peng, C. (2019). Accumulation of anthocyanins: an adaptation strategy
of Mikania micrantha to low temperature in winter. *Frontiers in Plant Science, 10*, 1049.
https://doi.org/10.3389/fpls.2019.01049
Zhang, S., Quartararo, A., Betz, O. K., Madahhosseini, S., Heringer, A. S., Le, T., ... & Drakakaki, G. (2021).
Root vacuolar sequestration and suberization are prominent responses of Pistacia spp. rootstocks during salinity
stress. Plant Direct, 5(5), e00315. https://doi.org/10.1002/pld3.315
Zhao, S., Zhang, Q., Liu, M., Zhou, H., Ma, C., & Wang, P. (2021). Regulation of plant responses to salt stress.
International Journal of Molecular Sciences, 22(9), 4609. https://doi.org/10.3390/ijms22094609
Zheng, L., Ma, H., Jiao, Q., Ma, C., & Wang, P. (2020). Phytohormones: important participators in plant salt
tolerance. *International Journal of Agriculture and Biology*, *24*(2), 319-332.
https://doi.org/10.17957/ijab.15.1441
Zou, Y. N., Wang, P., Liu, C. Y., Ni, Q. D., Zhang, D. J., & Wu, Q. S. (2017). Mycorrhizal trifoliate orange has
greater root adaptation of morphology and phytohormones in response to drought stress. *Scientific reports*, *7*(1),
1-10. https://doi.org/10.1038/srep41134

1155
1156
1157 **Figure legends**

1158

Figure 1. Leaf growth, photosynthesis efficiency and mineral nutrition in *A. thaliana* demes under control
(C), neutral (*neuSAL*) and alkaline (*alkSAL*) salinity. (A) Mean $\pm$ SE of growth (relative rosette diameter) of
each deme and treatment. Each dot represents a biological replicate (n=6). Mean values with different letters
indicate significant differences (Tukey's HSD, adj. p-value < 0.05). **(B)** Mean $\pm$ SE of Maximum PSII
photochemical efficiency (F_v/F_m) of each deme and treatment. Each dot represents the mean value of 2 technical
replicates (2 measurements on each biological replicate) (n=6). Mean values with different letters indicate
significant differences (Tukey's HSD, adj. p-value < 0.05). **(C)** Radial plots showing normalized differences of
10 elements in plant leaf samples from each deme and treatment. Axes display Z-scores calculated per element,
and elements exhibiting significant differences in at least 1 deme (Tukey's HSD, adj. p-value < 0.05) are marked
with an asterisk on each radial plot. **(D)** Mean $\pm$ SE of analyzed nutrient ratios in the two contrasted demes T6[CS]
and V1[PC]. Each dot represents a biological replicate (n=6). Nutrient ratios exhibiting significant differences for
each deme between salinity treatments (Student *t*-Test, adj. p-value < 0.05) are marked with an asterisk (*: adj.
p-value < 0.05; **: adj. p-value < 0.01). Plants were grown in potting mix soil and irrigated with C (0.5-
Hoagland, pH 5.9), *neuSAL* (100 mM NaCl, pH 5.9) or *alkSAL* (85 mM NaCl, 15 mM NaHCO$_3$, pH 8.3) for 2
weeks.

1174

Figure 2. Endogenous phytohormone content in *A.thaliana* demes under control (C), neutral (*neuSAL*)
and alkaline (*alkSAL*) salinity. Figure 2. Endogenous phytohormone content in A.thaliana demes under neutral
(neuSAL) and alkaline (alkSAL) salinity. Mean $\pm$ SE of relative (A) Indole acetic acid (IAA), (B) 1-
Aminocyclopropane-1-carboxylic acid (ACC), (C) Jasmonic Acid (JA), (D) Salicylic Acid (SA) and (E) Abscisic
Acid (ABA) concentration in all demes under control and salinity treatments. Each dot represents a biological
replicate (n=6), and treatments are color-coded (see legend). Zoomed boxplots show relative phytohormone
concentration for each deme. Mean values with different letters indicate significant differences (Tukey's HSD,
adj. p-value < 0.05). Plants were grown in potting mix soil and irrigated with C (0.5-Hoagland, pH 5.9), neuSAL
(100 mM NaCl, pH 5.9) or alkSAL (85 mM NaCl, 15 mM NaHCO$_3$, pH 8.3) for 2 weeks.

1184

1185 **Figure 3. Multivariate analysis on physiological traits in *A. thaliana* demes under control (C), neutral**
1186 **(*neuSAL*) and alkaline (*alkSAL*) salinity. (A)**: Principal Component Analysis (PCA). Left: individual factor
1187 map with treatment-colored samples distribution on the 2 most informative axes (PC1 and PC2) of the PCA.

Right: variable factor map presenting the amount of variance from each trait on the total variance in the PCA. Longer arrows represent larger amount from total variance. **(B)** Mean ± SE of leaf Zn (μg g^{-1} DW) in all demes under control and salinity treatments. **(C)** Scatter plot matrix section from multivariate correlation analysis of all physiological traits. Scatter plots show correlation between leaf Zn [(μg g^{-1} DW), y-axis] and leaf JA and Mg [(ng g^{-1} FW), left; (μg g^{-1} DW), right; x-axis] in all samples. **(D)** Mean ± SE of leaf B (μg g^{-1} DW) in all demes under control and salinity treatments. **(E)** Scatter plot matrix section from multivariate correlation analysis of all physiological traits. Scatter plots show correlation between leaf Na [(μg g^{-1} DW), y-axis] and leaf B (left) and K (right) (μg g^{-1} DW); x-axis] in all samples. r = correlation coefficient. Each dot represents a biological replicate (n=3 per deme) and treatments are color-coded (see legend). Letters indicate significant differences (Tukey's HSD, adj. p-value < 0.05). Plants were grown in potting mix soil and irrigated with C (0.5-Hoagland, pH 5.9), *neuSAL* (100 mM NaCl, pH 5.9) or *alkSAL* (85 mM NaCl, 15 mM NaHCO$_3$, pH 8.3) for 2 weeks.

Figure 4. Differential Gene Expression (DEG) analysis in *A. thaliana* demes T6^{CS} (yellow), Ro2^{CS} (black), V1^{PC} (blue), and LM2^{PC} (grey). (A) Multidimensional scaling (MDS) plot of RNA expression profiles in the studied samples. Sample relations are plotted using a multidimensional scaling plot (MDS) generated with edgeR showing the variability between replicates and treatments in log2-fold-change distance. The axes represent gene expression levels between the different experimental factors. Each dot represented a sample, color coded by treatment (Black: Control, Blue: *neuSAL*, Yellow: *AlkSAL*) and shape coded by deme (Color-filled square: T6^{CS}, empty square: Ro2^{CS}, color-filled octagon: V1^{PC}, empty octagon: LM2^{PC}). **(B)** Histogram showing the number of differentially expressed genes for each deme (n=3) and treatment (*neuSAL*/Control; *alkSAL*/Control). **(C)** Venn diagram showing the mutual overlaps of DEGs among demes under each treatment. DEGs were filtered at log fold change (LFC > 2, LFC < -2), and *q*-value (FDR-adjusted *p*-value) < 0.05.

Figure 5. Characterization of the onlySAL and onlyALKSAL sets. DEGs were filtered at log fold change (LFC > 2, LFC < -2), and q-value (FDR-adjusted p-value) < 0.05. Chart pie showing expression trend of total **(A)** onlySAL, **(B)** onlyALKSAL and **(C)** coreSAL DEGs. KEGG pathway enrichment analyses of downregulated (bubble plot, left) and upregulated DEGs (bubble plot, right) are shown for each corresponding DEG set. In bubble plots, pathways belonging to different classifications are listed on the left. Only the top 4 are presented, which are sorted by the q-value. Rich ratio (x-axis) is the ratio of the DEG number to the total gene number in a certain pathway. Bubble size represents number of DEGs included in each pathway (see legend) and enclosed bubbles indicate significant pathway enrichment. In heatmap: genes were clustered according to their expression pattern (Euclidean distance) and column clustering groups samples based on gene expression similarity. Color scale indicates the expression levels (yellow, low expression; blue, high expression).

Figure 6. Global overview of phytohormone signaling dynamics. (A) Protein-protein interaction (PPI) network of the analyzed DEGs involved in phytohormone signaling based on the STRING online database (https://string-db.org/). Nodes are color-coded according to each DEGs subset, arrows inside nodules indicate expression trend and colored surrounding areas distinguish between different hormone classes (see legends). (B) Expression levels [mean ± SE log2(FC)] of DEGs from (B) SAURs & IAAs (n=56), (C) LOXs/MYCs (n=20), (D) WRKYs (n=44), (E) ABIs (n=16), (F) JAZs (n=8), (G) PP2Cs (n=8), (H) IX clade ERFs (n=12), (I) PYLs/PYRs (n=23) and (J) DREBs (n=24) in all demes among treatments. Asterisks indicate significant differences in mean expression values among treatments (Student t-test, adj. p-value < 0.05; *: adj. p-value < 0.05; **: adj. p-value < 0.01). Neutral salinity: *neuSAL*; alkaline salinity: *alkSAL*.

Figure 7. Characterization of the coreSAL DEGs set. DEGs were filtered at log fold change (LFC > 2, LFC < -2), and *q*-value (FDR-adjusted *p*-value) < 0.05. Expression levels [mean ± SE log2(FC)] of **(A)** "photosynthesis", **(B)** "glucosinolate metabolism" and **(C)** "carbon metabolism" DEGs in each deme (n=3) and treatment. Letters indicate significant differences in mean expression values among demes (Tukey's HSD, adj. *p*-value < 0.05). **(D)** Heat map showing the expression profile of coreSAL DEGs (1219 genes) for each deme between neutral and alkaline salinity when compared to control (neuSAL/C; alkSAL/C). Genes were clustered according to their expression pattern (Euclidean distance). In frame, heatmap showing the expression profiles of coreSAL DEGs with contrasted expression trends among salinity types and/or demes. The color scale indicates the expression levels (grey, low expression; pink, high expression). Genes were clustered according to their

1241 expression pattern (Euclidean distance), while column clustering groups samples based on gene expression
1242 similarity.
1243
Figure 8. Characterization of *BGAL4* expression and sequence variation in *A. thaliana* demes. (A)
Expression levels [mean ± SE log2(FC)] of *BGAL4, DIN10* and *SEN1* in each deme (n=3) and treatment. Letters
indicate significant differences in mean expression values among demes (Tukey's HSD, adj. *p*-value < 0.05).
T6^{CS} (yellow), Ro2^{CS} (black), V1^{PC} (blue), and LM2^{PC} (grey). **(B)** SNP polymorphisms surrounding the *BGAL4*
locus. Triangles: candidate SNPs position. Amino acid change is indicated below each SNP. Gene orientation is
indicated with an arrow on the right. Exons are indicated with boxes and introns with lines connecting them.
Chromosome positions are indicated at the top left (picture from https://www.arabidopsis.org/index.jsp).
Reference and alternative allele nucleotide changes, variant type and deme bearing the SNP can be found in
Figure S8.
1253
1254 **Figure 9. WGCNA analysis.** Relationships of co-expression modules and analyzed physiological traits. Each
1255 row in the table corresponds to a co-expression module, and each column to a trait. Numbers in the matrix report
1256 the correlations of the corresponding module genes and each trait, with the p values below the correlations in
1257 parentheses. The table is color coded by correlation according to the color legend.
1258
1259 **Figure 10. Overview of WGCNA co-expression modules significantly correlating to alkaline salinity**
1260 **tolerance traits.** Protein-protein interaction (PPI) network of all DEGs comprised in darkgrey, plum1 and blue
1261 WGCNA co-expression modules, based on the STRING online database (https://string-db.org/). Proposed
1262 candidates identified in each module and appearing on each PPI Network are indicated, and main signaling
1263 pathways enriched per module encircled.
1264
1265 **Figure S1. Plant material and soil of origin. (A)** Distribution of saline soils in Europe. Red area is categorized
1266 as saline (>50% surface). Frame shows studied demes in Catalonia's map **(B)** Table lists deme UTM coordinates
1267 and source.
1268
1269 **Figure S2. Workflow.** Schematic overview of the compared sample growth conditions and performed analyses.
1270
1271 **Figure S3. Contribution of each deme to the coreSAL DEGs set.** Venn diagrams show overlapping DEGs in
1272 *coreSAL* (grey) and total DEGs per deme (yellow: total DEGs from *neuSAL* treatment; blue: total DEGs from
1273 *alkSAL* treatment). Table summarizes how many coreSAL DEGs are altered in each deme ("Total"), the salinity
1274 type causing the alteration ("*neuSAL*", "*alkSAL*", "*Shared*") and number of upregulated, downregulated and
1275 mixed-trends DEGs.
1276
1277 **Figure S4. Dissection on significantly enriched pathways from *onlyALKSAL* upregulated DEGs.**
1278 Expression levels [mean ± SE log2(FC)] of DEGs from **(A)** "Phenylpropanoid biosynthesis" (n=100) and **(B)**
1279 "Glutathione metabolism" (n=72) pathways in all demes among salinity treatments. Asterisks indicate significant
1280 differences in mean expression values (Student *t*-test, adj. *p*-value < 0.05; *: adj. *p*-value < 0.05; **: adj. *p*-value
1281 < 0.01) among treatments: ***neuSAL*** (neutral salinity in blue), ***alkSAL*** (alkaline salinity in yellow).
1282
Figure S5. Dissection on significantly enriched pathways from *coreSAL* upregulated DEGs under neutral
salinity (*neuSAL* in blue) and alkaline salinity (*alkSAL* in yellow) treatments. (A) Expression levels [mean ±
SE log2(FC)] of DEGs from "Ribosome biogenesis" pathway in each deme and treatment (n=7). Letters indicate
significant differences in mean expression values among demes under *neuSAL* (capital) and *alkSAL* (lower case)
(Tukey's HSD, adj. *p*-value < 0.05). Asterisks indicate significant differences in mean expression values among
treatments for each deme (Student *t*-test, adj. *p*-value < 0.05 **(B)** Expression levels [mean ± SE log2(FC)] of
DEGs from "Protein synthesis in ER" pathway in each location group (T6^{CS};Ro2^{CS}/V1^{PC};LM2^{PC}) and treatment
(n=4). Letters indicate significant differences in mean expression values among location groups under *neuSAL*
(capital) and *alkSAL* (lower case) (Student *t*-test, adj. *p*-value < 0.05). Asterisks indicate significant differences
in mean expression values among treatments for each location group (Student *t*-test, adj. *p*-value < 0.05; *: adj.
p-value < 0.05; **: adj. *p*-value < 0.01). **(C)** Expression levels [mean ± SE log2(FC)] of DEGs subset (FAR1

and FAR5) from "Cutin, suberin & wax biosynthesis" pathway in each location group (T6^{CS};Ro2^{CS}/V1^{PC};LM2^{PC}) and treatment (n=4). Letters indicate significant differences in mean expression values among location groups under *neuSAL* (capital) and *alkSAL* (lower case) (Student *t*-test, adj. *p*-value < 0.05).

Figure S6. Dissection on significantly enriched pathways from *coreSAL* downregulated DEGs under neutral salinity (*neuSAL* in blue) and alkaline salinity (*alkSAL* in yellow) treatments. Expression levels [mean ± SE log2(FC)] of DEGs from **(A)** "Photosynthesis" (n=2), **(B)** "Photosynthesis Antenna proteins" (n=6), **(C)** "Glucosinolate biosynthesis" (n=9), **(D)** "Carbon metabolism" (n=3), **(E)** "Carbon fixation" (n=2), **(F)** "Glioxylate and dicarboxylate metabolism", **(G)** "Glycine, serine and threonine metabolism" and **(H)** "Nitrogen metabolism" pathways in each deme and treatment. Letters indicate significant differences in mean expression values among demes under *neuSAL* (capital) and *alkSAL* (lower case) (Tukey's HSD, adj. *p*-value < 0.05).

Figure S7. Key DEGs in the inhibition response of the studied demes. Protein-protein interaction (PPI) network of the down-regulated DEGs from enriched KEGG pathways in the *coreSAL* response. Performed at the STRING online database (https://string-db.org/).

Figure S8. Gene models for BGAL4 and DIN10. Location of the SNPs identified at the coding sequence.

Figure S9. WGCNA analysis. Soft thresholding Network topology for different soft-thresholding powers. Numbers in the plots indicate the corresponding soft thresholding powers.

Figure S10. WGCNA analysis. Gene modules identified by Weighted gene co-expression network analysis. Gene dendrogram obtained by clustering the dissimilarity based on consensus topological overlap with the corresponding module colors indicated by the color row. Each colored row represents a color-coded module which contains a group of highly connected genes.

Figure S11. WGCNA analysis. Eigengene dendrogram and adjacency heatmap. Dendrogram of consensus module eigengenes obtained by WGCNA on the consensus correlation. The red line is the merging threshold, and groups of eigengenes below the threshold represent modules whose expressions profiles should be merged due to their similarity. Heatmap plotting adjacencies of modules. Red represents high adjacency (positive correlation) and blue represents low adjacency (negative correlation).

Figure S12. WGCNA analysis. Topological Overlap Matrix (TOM) showing the interaction of co-expression genes and the cluster dendrogram of 1,000 randomly selected genes. The intensity of the yellow inside the heatmap represents the degree of overlap (orange: high; yellow: low).

Figure 1. Leaf growth, photosynthesis efficiency and mineral nutrition in *A. thaliana* demes under control (C), neutral (neuSAL) and alkaline (alkSAL) salinity. (A) Mean ± SE of growth (relative rosette diameter) of each deme and treatment. Each dot represents a biological replicate (n=6). Mean values with different letters indicate significant differences (Tukey's HSD, adj. p-value < 0.05). (B) Mean ± SE of Maximum PSII photochemical efficiency (Fv/Fm) of each deme and treatment. Each dot represents the mean value of 2 technical replicates (2 measurements on each biological replicate) (n=6). Mean values with different letters indicate significant differences (Tukey's HSD, adj. p-value < 0.05). (C) Radial plots showing normalized differences of 10 elements in plant leaf samples from each deme and treatment. Axes display Z-scores calculated per element, and elements exhibiting significant differences in at least 1 deme (Tukey's HSD, adj. p-value < 0.05) are marked with an asterisk on each radial plot. (D) Mean ± SE of analyzed nutrient ratios in the two contrasted demes T6CS and V1PC. Each dot represents a biological replicate (n=6). Nutrient ratios exhibiting significant differences for each deme between salinity treatments (Student t-Test, adj. p-value < 0.05) are marked with an asterisk (*: adj. p-value < 0.05; **: adj. p-value < 0.01). Plants were grown in potting mix soil and irrigated with C (0.5- Hoagland, pH 5.9), neuSAL (100 mM NaCl, pH 5.9) or alkSAL (85 mM NaCl, 15 mM NaHCO3, pH 8.3) for 2 weeks.

Figure 2. Endogenous phytohormone content in *A.thaliana* demes under control (C), neutral (neuSAL) and alkaline (alkSAL) salinity. Figure 2. Endogenous phytohormone content in A.thaliana demes under neutral (neuSAL) and alkaline (alkSAL) salinity. Mean ± SE of relative (A) Indole acetic acid (IAA), (B) 1-Aminocyclopropane-1-carboxylic acid (ACC), (C) Jasmonic Acid (JA), (D) Salicylic Acid (SA) and (E) Abscisic Acid (ABA) concentration in all demes under control and salinity treatments. Each dot represents a biological replicate (n=6), and treatments are color-coded (see legend). Zoomed boxplots show relative phytohormone concentration for each deme. Mean values with different letters indicate significant differences (Tukey's HSD, adj. p-value < 0.05). Plants were grown in potting mix soil and irrigated with C (0.5-Hoagland, pH 5.9), neuSAL (100 mM NaCl, pH 5.9) or alkSAL (85 mM NaCl, 15 mM NaHCO3, pH 8.3) for 2 weeks.

Figure 3. Multivariate analysis on physiological traits in *A. thaliana* demes under control (C), neutral (neuSAL) and alkaline (alkSAL) salinity. (A): Principal Component Analysis (PCA). Left: individual factor map with treatment-colored samples distribution on the 2 most informative axes (PC1 and PC2) of the PCA. Right: variable factor map presenting the amount of variance from each trait on the total variance in the PCA. Longer arrows represent larger amount from total variance. (B) Mean ± SE of leaf Zn (μg g-1 DW) in all demes under control and salinity treatments. (C) Scatter plot matrix section from multivariate correlation analysis of all physiological traits. Scatter plots show correlation between leaf Zn [(μg g-1 DW), y-axis] and leaf JA and Mg [(ng g-1 FW), left; (μg g-1 DW), right; x-axis] in all samples. (D) Mean ± SE of leaf B (μg g-1 DW) in all demes under control and salinity treatments. (E) Scatter plot matrix section from multivariate correlation analysis of all physiological traits. Scatter plots show correlation between leaf Na [(μg g-1 DW), y-axis] and leaf B (left) and K (right) (μg g-1 DW); x-axis] in all samples. r = correlation coefficient. Each dot represents a biological replicate (n=3 per deme) and treatments are color-coded (see legend). Letters indicate significant differences (Tukey's HSD, adj. p-value < 0.05). Plants were grown in potting mix soil and irrigated with C (0.5-Hoagland, pH 5.9), neuSAL (100 mM NaCl, pH 5.9) or alkSAL (85 mM NaCl, 15 mM NaHCO3, pH 8.3) for 2 weeks.

A

B

Figure 5. Characterization of the onlySAL and onlyALKSAL sets. DEGs were filtered at log fold change (LFC > 2, LFC < -2), and q-value (FDR-adjusted p-value) < 0.05. Chart pie showing expression trend of total (A) onlySAL, (B) onlyALKSAL and (C) coreSAL DEGs. KEGG pathway enrichment analyses of downregulated (bubble plot, left) and upregulated DEGs (bubble plot, right) are shown for each corresponding DEG set. In bubble plots, pathways belonging to different classifications are listed on the left. Only the top 4 are presented, which are sorted by the q-value. Rich ratio (x-axis) is the ratio of the DEG number to the total gene number in a certain pathway. Bubble size represents number of DEGs included in each pathway (see legend) and enclosed bubbles indicate significant pathway enrichment. In heatmap: genes were clustered according to their expression pattern (Euclidean distance) and column clustering groups samples based on gene expression similarity. Color scale indicates the expression levels (yellow, low expression; blue, high expression).

Figure 4. Differential Gene Expression (DEG) analysis in *A. thaliana* demes. (A) Multidimensional scaling (MDS) plot of RNA expression profiles in the studied samples. Sample relations are plotted with edgeR showing the variability between replicates and treatments in log2-fold-change distance. The axes represent gene expression levels between the different experimental factors.

Each dot represents a sample, color coded by treatment (Black: Control, Blue: neuSAL, Yellow: AlkSAL) and shape coded by deme (Color-filled square: T6CS, empty square: Ro2CS, color-filled octagon: V1PC, empty octagon: LM2PC). (B) Histogram showing the number of differentially expressed genes for each deme (n=3) and treatment (neuSAL/Control; alkSAL/Control). (C) Venn diagram showing mutual overlaps of DEGs among demes under each treatment. DEGs were filtered at log fold change (LFC > 2, LFC < -2), and q-value (FDR-adjusted p-value) < 0.05.

Figure 6. Global overview of phytohormone signaling dynamics. (A) Protein-protein interaction (PPI) network of the analyzed DEGs involved in phytohormone signaling based on the STRING online database (https://string-db.org/). Nodes are color-coded according to each DEGs subset, arrows inside nodules indicate expression trend and colored surrounding areas distinguish between different hormone classes (see legends). (B) Expression levels [mean ± SE log2(FC)] of DEGs from (B) SAURs & IAAs (n=56), (C) LOXs/MYCs (n=20), (D) WRKYs (n=44), (E) ABIs (n=16), (F) JAZs (n=8), (G) PP2Cs (n=8), (H) IX clade ERFs (n=12), (I) PYLs/PYRs (n=23) and (J) DREBs (n=24) in all demes among treatments. Asterisks indicate significant differences in mean expression values among treatments (Student t-test, adj. p-value < 0.05; *: adj. p-value < 0.05; **: adj. p-value < 0.01). Neutral salinity: neuSAL; alkaline salinity: alkSAL.

Figure 7. Characterization of the coreSAL DEGs set. DEGs were filtered at log fold change (LFC > 2, LFC < -2), and q-value (FDR-adjusted p-value) < 0.05. Expression levels [mean ± SE log2(FC)] of (A) "photosynthesis", (B) "glucosinolate metabolism" and (C) "carbon metabolism" DEGs in each deme (n=3) and treatment. Letters indicate significant differences in mean expression values among demes (Tukey's HSD, adj. p-value < 0.05). (D) Heat map showing the expression profile of coreSAL DEGs (1219 genes) for each deme between neutral and alkaline salinity when compared to control (neuSAL/C; alkSAL/C). Genes were clustered according to their expression pattern (Euclidean distance). In frame, heatmap showing the expression profiles of coreSAL DEGs with contrasted expression trends among salinity types and/or demes. The color scale indicates the expression levels (grey, low expression; pink, high expression). Genes were clustered according to their expression pattern (Euclidean distance), while column clustering groups samples based on gene expression similarity.

Figure 8. Characterization of BGAL4 expression and sequence variation in A. thaliana demes. (A) Expression levels [mean ± SE log2(FC)] of BGAL4, DIN10 and SEN1 in each deme (n=3) and treatment. Letters indicate significant differences in mean expression values among demes (Tukey's HSD, adj. p-value < 0.05). $T6_{CS}$ (yellow), $Ro2_{CS}$ (black), $V1_{PC}$ (blue), and $LM2_{PC}$ (grey). (B) SNP polymorphisms surrounding the BGAL4 locus. Triangles: candidate SNPs position. Amino acid change is indicated below each SNP. Gene orientation is indicated with an arrow on the right. Exons are indicated with boxes and introns with lines connecting them. Chromosome positions are indicated at the top left (picture from https://www.arabidopsis.org/index.jsp). Reference and alternative allele nucleotide changes, variant type and deme bearing the SNP can be found in Figure S8.

Co-expression module-trait correlation

Figure 9. WGCNA analysis. Relationships of co-expression modules and analyzed physiological traits. Each row in the table corresponds to a co-expression module, and each column to a trait. Numbers in the matrix report the correlations of the corresponding module genes and each trait, with the p values below the correlations in parentheses. The table is color coded by correlation according to the color legend.

Darkgrey Module

JA signaling

IAA signaling

Glucosinolate signaling

PEP-Carboxylase
(Photosynthesis)

AK3

FUM2

α-CA1

Plum1 Module

PGLP1

HPR

Photorespiration

Blue Module

Photosynthesis
&
Antenna Proteins

DNA replication
&
Cell cycle

RBCS1B

PSAH-1

LHCs

PP2C
(AT2G40860)

MAP kinases

Serine & threonine kinases

Figure 10. Overview of WGCNA co-expression modules significantly correlating to alkaline salinity tolerance traits. Protein-protein interaction (PPI) network of all DEGs comprised in darkgrey, plum1 and blue WGCNA co-expression modules, based on the STRING online database (https://string-db.org/). Proposed candidates identified in each module and appearing on each PPI Network are indicated, and main signaling pathways enriched per module encircled.

(A)

Saline category
> 50% saline

Pre-litoral range

● T6CS ●Ro2CS ● V1PC ● LM2PC

(B)

Deme	UTM (X)	UTM (Y)	Source
T6CS	496641	4621417	K. Bomblies
Ro2CS	515998	4680193	Anthos
V1PC	512474	4686157	Anthos
LM2PC	489123	4641204	K. Bomblies

Figure S1. Plant material and soil of origin. (A) Distribution of saline soils in Europe. Red area is categorized as saline (>50% surface). Frame shows studied demes in Catalonia's map (B) Table lists deme UTM coordinates and source.

Figure S2. Workflow. Schematic overview of the compared sample growth conditions and performed analyses.

Deme	Total	neuSAL	Trend	alkSAL	Trend	Shared	Trend
T6	1214	10	2 ⬇ 8 ⬆	22	6 ⬆ 16 ⬇	1182	446 ⬆ 772 ⬇ 4 sal ⬇⬆ alksal
Ro2	1170	30	19 ⬇ 11 ⬆	193	99 ⬆ 94 ⬇	947	269 ⬆ 672 ⬇ 6 sal ⬇⬆ alksal
LM2	1133	31	11 ⬇ 20 ⬆	193	56 ⬆ 137 ⬇	909	342 ⬆ 566 ⬇ 1 sal ⬇⬆ alksal
V1	1188	30	9 ⬇ 21 ⬆	65	19 ⬆ 49 ⬇	1090	382 ⬆ 702 ⬇ 1 sal ⬇⬆ alksal ; 6 sal ⬆⬆ alksal

Figure S3. Contribution of each deme to the coreSAL DEGs set. Venn diagrams show overlapping DEGs in coreSAL (grey) and total DEGs per deme (yellow: total DEGs from neuSAL treatment; blue: total DEGs from alkSAL treatment). Table summarizes how many coreSAL DEGs are altered in each deme ("Total"), the salinity type causing the alteration ("neuSAL", "alkSAL", "Shared") and number of upregulated, downregulated and mixed-trends DEGs.

Figure S4. Dissection on significantly enriched pathways from onlyALKSAL upregulated DEGs. Expression levels [mean ± SE log2(FC)] of DEGs from (A) "Phenylpropanoid biosynthesis" (n=100) and (B) "Glutathione metabolism" (n=72) pathways in all demes among salinity treatments. Asterisks indicate significant differences in mean expression values (Student t-test, adj. p-value < 0.05; *: adj. p-value < 0.05; **: adj. p-value < 0.01) among treatments: neuSAL (neutral salinity in blue), alkSAL (alkaline salinity in yellow).

Figure S5. Dissection on significantly enriched pathways from coreSAL upregulated DEGs under neutral salinity (neuSAL in blue) and alkaline salinity (alkSAL in yellow) treatments. (A) Expression levels [mean ± SE log2(FC)] of DEGs from "Ribosome biogenesis" pathway in each deme and treatment (n=7). Letters indicate significant differences in mean expression values among demes under neuSAL (capital) and alkSAL (lower case) (Tukey's HSD, adj. p-value < 0.05). Asterisks indicate significant differences in mean expression values among treatments for each deme (Student t-test, adj. p-value < 0.05 (B) Expression levels [mean ± SE log2(FC)] of DEGs from "Protein synthesis in ER" pathway in each location group (T6CS;Ro2CS/V1PC;LM2PC) and treatment (n=4). Letters indicate significant differences in mean expression values among location groups under neuSAL (capital) and alkSAL (lower case) (Student t-test, adj. p-value < 0.05). Asterisks indicate significant differences in mean expression values among treatments for each location group (Student t-test, adj. p-value < 0.05; *: adj. p-value < 0.05; **: adj. p-value < 0.01). (C) Expression levels [mean ± SE log2(FC)] of DEGs subset (FAR1 and FAR5) from "Cutin, suberin & wax biosynthesis" pathway in each location group (T6CS;Ro2CS/V1PC;LM2PC) and treatment (n=4). Letters indicate significant differences in mean expression values among location groups under neuSAL (capital) and alkSAL (lower case) (Student t-test, adj. p-value < 0.05).

Figure S6. Dissection on significantly enriched pathways from coreSAL downregulated DEGs under neutral salinity (neuSAL in blue) and alkaline salinity (alkSAL in yellow) treatments. Expression levels [mean ± SE log2(FC)] of DEGs from (A) "Photosynthesis" (n=2), (B) "Photosynthesis Antenna proteins" (n=6), (C) "Glucosinolate biosynthesis" (n=9), (D) "Carbon metabolism" (n=3), (E) "Carbon fixation" (n=2), (F) "Glioxylate and dicarboxylate metabolism", (G) "Glycine, serine and threonine metabolism" and (H) "Nitrogen metabolism" pathways in each deme and treatment. Letters indicate significant differences in mean expression values among demes under neuSAL (capital) and alkSAL (lower case) (Tukey's HSD, adj. p-value < 0.05).

Figure S7. Key DEGs in the inhibition response of the studied demes. Protein-protein interaction (PPI) network of the down-regulated DEGs from enriched KEGG pathways in the coreSAL response. Performed at the STRING online database (https://string-db.org/).

Figure S8. Gene models for BGAL4 and DIN10. Location of the SNPs identified at the coding sequence.

Figure S9. WGCNA analysis. Soft thresholding Network topology for different soft-thresholding powers. Numbers in the plots indicate the corresponding soft thresholding powers.

Gene dendrogram and module colors

Module colors

Figure S10. WGCNA analysis. Gene modules identified by Weighted gene co-expression network analysis. Gene dendrogram obtained by clustering the dissimilarity based on consensus topological overlap with the corresponding module colors indicated by the color row. Each colored row represents a color-coded module which contains a group of highly connected genes.

Figure S11. WGCNA analysis. Eigengene dendrogram and adjacency heatmap. Dendrogram of consensus module eigengenes obtained by WGCNA on the consensus correlation. The red line is the merging threshold, and groups of eigengenes below the threshold represent modules whose expressions profiles should be merged due to their similarity. Heatmap plotting adjacencies of modules. Red represents high adjacency (positive correlation) and blue represents low adjacency (negative correlation).

Figure S12. WGCNA analysis. Topological Overlap Matrix (TOM) showing the interaction of co-expression genes and the cluster dendrogram of 1,000 randomly selected genes. The intensity of the yellow inside the heatmap represents the degree of overlap (orange: high; yellow: low).