# Implicit auditory memory in older listeners: from encoding to 6-month retention

Roberta Bianco[1,2], Edward T. R. Hall[3], Marcus. T. Pearce[3,4], Maria Chait[1]

[1] Ear Institute, University College London, WC1X 8EE London, United Kingdom

[2] Neuroscience of Perception and Action Laboratory, Italian Institute of Technology, 00161 Rome, Italy

[3] School of Electronic Engineering and Computer Science, Queen Mary University of London, E1 4NS London, United Kingdom

[4] Department of Clinical Medicine, Aarhus University, 8000 Aarhus C, Denmark

**Declaration of interests.** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Corresponding author:**

Roberta Bianco

roberta.bianco@iit.it

Neuroscience of Perception & Action Laboratory, Italian Institute of Technology (IIT)

Viale Regina Elena 291, 00161 Rome, Italy

# Abstract

Any listening task, from sound recognition to sound-based communication, rests on auditory memory which is known to decline in healthy ageing. However, how this decline maps onto multiple components and stages of auditory memory remains poorly characterised. In an online unsupervised longitudinal study, we tested ageing effects on implicit auditory memory for rapid tone patterns. The test required participants (younger, aged 20-30, and older adults aged 60-70) to quickly respond to rapid regularly repeating patterns emerging from random sequences. Patterns were novel in most trials (REGn), but unbeknownst to the participants, a few distinct patterns reoccurred identically throughout the sessions (REGr). After correcting for processing speed, the response times (RT) to REGn should reflect the information held in echoic and short-term memory before detecting the pattern; long-term memory formation and retention should be reflected by the RT advantage (RTA) to REGr vs REGn which is expected to grow with exposure. Older participants were slower than younger adults in detecting REGn and exhibited a smaller RTA to REGr. Computational simulations using a model of auditory sequence memory indicated that these effects reflect age-related limitations both in early and long-term memory stages. In contrast to ageing-related accelerated forgetting of verbal material, here older adults maintained stable memory traces for REGr patterns up to 6 months after the first exposure. The results demonstrate that ageing is associated with reduced short-term memory and long-term memory formation for tone patterns, but not with forgetting, even over surprisingly long timescales.

*Keywords:* sequential pattern, sound perception, forgetting, statistical learning, Prediction by Partial Matching (PPM)

# 1. Introduction

Memory loss is one of the most significant changes to cognitive processing experienced in healthy ageing (Cansino, 2009; Füllgrabe, 2020; Nyberg et al., 2012; Raz and Lindenberger, 2011; Salthouse, 2011). The most pronounced memory deficits are related to direct recall of episodic memory (Koen and Yonelinas, 2016, 2014), but evidence is increasingly revealing impairment in older listeners also in tasks that draw on automatic sensory memory processes (Cheng et al., 2013; Humes et al., 2013; Janacsek et al., 2012; Rieckmann and Bäckman, 2009; Rimmele et al., 2012; Schneider and Pichora-Fuller, 2000; Sluming et al., 2002; Wayne and Johnsrude, 2015).

Sensory memory is at the core of auditory processing (Atkinson and Shiffrin, 1968; Cowan, 1984; Nees, 2016; Winkler and Cowan, 2005). The nature of the unfolding signal is such that any listening task, from sound recognition to sound-based communication, depends on the ability to store successive events in memory to derive a coherent representation (Baldeweg, 2006; Conway, 2020; Heilbron and Chait, 2018; Massaro and Cohen, 1975; Rimmele et al., 2015; Santolin and Saffran, 2018). Age-related deficits in implicit auditory memory are increasingly being documented (Bianco and Chait, 2023; Fogerty et al., 2016; Jääskeläinen et al., 1999; Pekkonen et al., 1996; Rimmele et al., 2012), but remain poorly characterised due to limited computational tractability and paucity of longitudinal research designs. Recently emerging links between auditory processing and dementia are also making it urgent to quantify and understand these impairments (Griffiths et al., 2020; Johnson et al., 2021). In particular, the brain networks that are thought to underlie auditory memory – involving auditory, frontal and hippocampal areas (Barascud et al., 2016; Bonetti et al., 2022; Brown et al., 2004; Burunat et al., 2014; Kumar et al., 2014; Schapiro et al., 2014; Schmithorst, 2005; Watanabe et al., 2008) – are the same networks that exhibit age-related alterations (Mander et al., 2013) and the earliest decline in Alzheimer's disease (Benhamou and Warren, 2020; Griffiths et al., 2020; Johnson et al., 2021). This makes auditory memory decline a

promising proximity marker of dementia that is worth investigating (Fleischman, 2007; Swords

et al., 2018).

Traditionally, auditory memory is conceived as a sequence of stores (Atkinson and

Shiffrin, 1968): an echoic buffer stores detailed unprocessed information for several hundred

milliseconds to allow successive sounds to be linked to the representation of sequences; then

information passes to short-term memory for a few seconds and strengthens in long-term

memory upon repeated presentations of the to-be-remembered sound (Winkler and Cowan,

2005). Impairments can potentially arise at any of these different stages.

Older, compared to younger, adults exhibit diminished amplitude and longer latencies

of the mismatch negativity (MMN) – an automatic brain response evoked by a rare deviant

sound in a sequence of standard sounds  (Cooper et al., 2006; Jääskeläinen et al., 1999;

Kiang et al., 2009; Pekkonen et al., 2001, 1996; Rimmele et al., 2012). The MMN is

hypothesised to reflect the process of comparing incoming inputs with sensory-memory traces

(Atienza et al., 2002; Baldeweg, 2006; Haenschel et al., 2005; Näätänen et al., 2007; Squires

et al., 1976). Diminished MMN in older listeners suggests reduced echoic buffer and/or short-

term memory (Näätänen et al., 2012). Age-related performance decline is also reported in

probabilistic sequence learning tests using artificial auditory material and is hypothesised to

reflect short- and long-term implicit memory deficits in older adults (Lukács and Kemény,

2015). In such tests, target sequences of arbitrary syllables or tones (presentation rate ~2Hz)

are structured according to certain probabilistic or deterministic statistics and repetitively

presented to listeners unaware of the structure underlying the sequence (Christiansen, 2019;

Petkov and ten Cate, 2020). Relative to novel sequences, a memory benefit for the target

sequences is reflected by better immediate sequence reproduction or higher familiarity ratings

after the exposure phase. This benefit declines with age and drastically at the age of around

65 (Herff et al., 2020; Lukács and Kemény, 2015; Schevenels et al., 2021).

Age-related decline has also been described in verbal memory tests asking

participants to memorise a list of words, nonsense words, or stories to a minimum required

level of accuracy. Memory is then typically probed with free recall at delayed sessions. Older

compared with younger adults exhibit a reduced primacy effect suggesting age-related

impairment in short-term memory (Mander et al., 2013; Murphy et al., 2000), as well as

accelerated long-term forgetting (Elliott et al., 2014; Mary et al., 2013; McGibbon et al., 2022;

Wearn et al., 2020) implicating impairments of long-term memory consolidation (Hoefeijzers

et al., 2013). Notably, accelerated long-term forgetting has gained particular traction in the

clinical field because of its potential as a predictor of Alzheimer's pathology (Weston et al.,

2018).

However, general confounding issues may emerge when tasks involve the

presentation of stimuli at relatively slow rates (typically ~ 2-4 Hz) and require engagement of

the participant with the to-be-remembered information (e.g., the requirement to actively recall,

judge familiarity etc). Factors such as experience, attention- or executive processing ability

might conceal a core informational aspect of age-related memory decline. Indeed, age

differences in immediate or long-term recall might reflect the effects of attentional load (Palmer

et al., 2018), availability of feedback (Herff et al., 2020), or vocabulary knowledge (Schneider

et al., 2002), as well as rehearsal strategies or interference (Manes et al., 2008; Mary et al.,

2013; Wearn et al., 2020; Weston et al., 2018). Additionally, complex material, such as words,

limits the cross-linguistic and translational diagnostic potential of verbal tests and, critically, it

is difficult to informationally quantify and model. Consequently, whilst it is generally accepted

that ageing is associated with auditory sensory memory impairment, the properties of this

mnemonic decline are poorly understood. An additional persistent methodological hurdle lies

in the ability to monitor the trajectory of memory processes over time – from the initial encoding

of memories to their enduring retention in the long term. To achieve this, it is necessary to use

standardized assessment techniques throughout these distinct stages.

We tested younger (aged between 20 and 30) and older (aged between 60 and 70)

participants with an online paradigm that allows us to quantify short-term memory, the

dynamics of long-term memory formation and long-term retention of tone patterns in an

unsupervised manner. The "Auditory pattern Memory test" (ApMEM) (Bianco et al., 2020a)

employs arbitrary rapid pure-tone sequences spanning the acoustic time scale of speech (20

Hz tone presentation rate, 1 Hz pattern rate) (Rosen, 1992). In 50% of the sequences, a

pattern (REG; a repeating sequence of 20 tones) emerges partway, and listeners are required

to detect it as quickly as possible (Fig. 1A). The rate at which successive tones are presented

precludes deliberate tracking of the sequence structure, instead the REG patterns pop-out

perceptually (Warren and Ackroff, 1976). Implicit auditory memory plays an obligatory role in

pattern detection (Winkler et al., 2009). The brain's ability to detect a REG pattern is

hypothesised to arise from an automatic process that scans the unfolding sequence and

compares incoming information with the just-heard sounds held in short-term memory and

stored representations of the longer-term context (Barascud et al., 2016). Therefore, after

adjusting for individual variability in detecting a simple sound change (see methods), the

response time (RT) associated with REG detection becomes a quantitative measure of the

information automatically held in memory until sufficient evidence has accumulated to trigger

the observer's behavioural response (Bianco et al., 2020a). In the task, most REG patterns

are novel on each trial (REGn). The associated RT can thus be taken to reflect the combined

contribution of implicit **_echoic and short-term memory_** to pattern detection. Crucially,

unbeknownst to the participants, a few different patterns reoccur sparsely (REGr; every 2

minutes). Memory for REGr strengthens through repetitive exposure, as reflected by the

gradual emergence of a RT advantage (RTA) in REGr pattern detection compared to REGn.

This relative measure is used to quantify **_long-term memory formation_**. Previous results from

young adults have shown that this effect is implicit, in that it is not driven by explicit familiarity

(Bianco et al., 2020a). To measure **_long-term memory retention_** of REGr patterns, RT to

REGr is also measured 8 days and 6 months after the first exposure.

Utilizing such arbitrary stimuli presents several notable advantages when investigating

auditory memory. Beyond overcoming potential linguistic barriers, these stimuli are too rapid

to be actively monitored (Warren and Ackroff, 1976), and unlikely to be encountered in real-

world settings. Consequently, they effectively mitigate mnemonic biases stemming from prior

linguistic familiarity, deliberate rehearsal tactics, or the interference posed by routinely

encountered auditory inputs. Importantly, these stimuli are also well suited for establishing a

systematic link between implicit listener performance (measured through RTs to patterns) and the information-theoretic attributes of the stimuli. This facilitates the computational simulation and quantification of how mnemonic subcomponents constrain listener behaviour.

The Prediction by Partial Matching (PPM) model, has successfully explained listeners' performance with a variety of discrete musical and artificial auditory sequences (Barascud et al., 2016; Bianco et al., 2020b; Di Liberto et al., 2020; Harrison et al., 2020; Kern et al., 2022; Omigie et al., 2019; Pearce et al., 2010; Quiroga-Martinez et al., 2020). Here, we use its memory-constrained variant (PPM-decay, see (Harrison et al., 2020), previously used to simulate performance in the ApMEM task in young listeners' (Bianco et al., 2020a). The model encodes sequences by weighting sub-sequences (n-grams) of multiple orders based on recency. Each auditory event is stored as a single count with a certain weight which decays over time (following the parameters set by a customizable decay kernel; Fig. 1C) and increases if the event is re-encountered. The non-linear decay profile simulates the contribution of three subcomponents to memory formation corresponding to echoic buffer, short- and long-term memory decay. Based on stored observations, the model estimates the information content (IC) for each event, corresponding to the degree to which the event is expected based on prior context. When a random sequence transitions into a novel REG pattern (REGn), the IC drops due to the match between incoming information and prior observations held in short-term memory. The brain is hypothesized to be sensitive to this change in IC (Barascud et al., 2016) as an indicator for changes in underlying stimulus statistics. For a previously encountered REG pattern (REGr), there will be a stronger match with information stored in long-term memory, resulting in earlier change in IC and consequently a faster response (Fig 1C). Within this framework, the RT to a pattern may encapsulate a combination of both speed in accessing stored information and associated confidence indirectly reflecting the weights this information has in memory.

Fitting ApMEM RT data from young and older adults with parameters associated with the decay kernel and memory weights, allowed us to identify the potential sources of memory impairment in the ageing cohort.

***Figure 1. Stimuli, experimental design, and a model of auditory memory. A) Example***

***spectrograms of the ApMEM stimuli***. *RAN sequences contained a random arrangement of*

*tone pips. RANREG sequences contained a transition from a random (RAN) to regularly*

*repeating cycles of 20 tone-pips (REGn). The repetition of the REG pattern becomes*

*detectable after the first cycle ('effective transition'). RAN and REGn sequences were*

*generated anew on each trial. Three different regular patterns (REGr) were each presented*

*identically trice within a block. Reoccurrences were spaced ~2 min apart. STEP stimuli,*

*containing a step change in frequency, (and their 'no change' control, CONT) were also*

*included in the stimulus set as a form of attentional checks, and to control for age-related*

*differences in speed of processing reflected in RT-based measures. To distil the computation*

*time required to detect the patterns RTs to RANREG were corrected by the median STEP RTs. **B) Experiment design and task order.** This was a 3-session study, conducted on day 1 (d1) to test memory formation over 3 blocks of ApMEM task, and on day 8 (d8) and month 6 (m6) to test memory retention with 1 block of ApMEM. Additional cognitive tests (random order) were included on d1: Sustained attention to response (SART), Choice reaction time task (CRT), and visual sequence memory task (Corsi blocks). The ApMEM familiarity surprise task was administered at the end of d8. **C) Auditory sequence memory model.** A schematic representation of the parameters implemented in the model to simulate auditory memory. **Memory decay.** Sequence statistics are memorised through partitioning the unfolding sequence into events and sub-sequences of increasing order (n-grams) that are thereon stored in memory. The salience of these observations ('weight') decays over time through 3 phases indicated by the shaded areas: (1) buffer, (2) short-term memory (STM, with a fast exponential decay from the weight of the buffer to the starting weight of the next phase over a given duration), and (3) the long-term memory (LTM, with a slow exponential-decay defined by a starting weight and a half-life). **Information tracking:** The model uses these stored statistics to quantify the information content (IC, where high IC corresponds to low predictability and low IC corresponds to high predictability) of incoming tones. A newly encountered pattern (REGn, in black) is detected when incoming information matches the information held in the buffer and the STM resulting in a drop of IC. Once an n-gram is encountered, its representation enters a slow LTM-decay phase. Pattern reoccurrence (REGr, illustrated with different shades of purple to indicate the same pattern after 3, 6 and 9 reoccurrences), leads to weight increase in memory as an index of increasing memory strength. This causes the fall in IC associated with pattern recognition to occur progressively earlier. Previous observations (Barascud et al., 2016; Bianco et al., 2020a) support the hypothesis that listeners track and leverage the drop in IC as an indication of the emergence of regularity.*

## 2. Methods

**2.1 Power analysis.**

We initially ran an online pilot experiment (N = 20, age between 20-30 years old). The RTA effects size across 3 blocks was $h_p^2$ = .22. We expected the difference between groups to be potentially small ($h_p^2$ = .02). A prospective power calculation (beta = 0.8; alpha = .05) for an ANOVA within-between interaction yielded a required total sample size of N = 41 per group. We set our online target sample size to N = 90 per group to account for dropouts (expected ~30%) due to headphone check exclusion and the unsupervised and longitudinal nature of the experiment. Experimental procedures were approved by the research ethics committee of University College London and informed consent was obtained from each participant.

**2.2 Participants.**

Two participant groups were recruited via the Prolific platform (https://www.prolific.co/). A group of younger participants (age range 20-30 years old; N=93) and a group of older participants (age range 60-70 years old; N=98). Inclusion criteria included being a native speaker of British English, generally good health, no known hearing problems, or cognitive impairment (all based on self-report). Participants using low-quality audio equipment, or those suffering from binaural hearing loss, were screened out using the test introduced by Milne et al. (2020). 29 participants in the older group, and 24 of the younger participants failed the screen and their data were therefore not analysed. Additional exclusion criteria were: (a) poor performance in the main apMEM test (mean d' < 1.5 across blocks in day 1 or day 8; N = 1 in the older group, N = 4 in the younger group excluded.) (b) poor performance on the attentional checks (mean RT to STEP changes larger than 2 STD away from the group means; N=1 in the younger group excluded). A final N = 132 was analysed: N = 68 (28 female) in the older group, and N = 64 (33 female) in the younger group (Fig. 2A). Six months later we ran an additional (surprise) session. From the original pool, N = 104 participants (N = 63 older, N =

41 younger group) signed up, and, after exclusion as mentioned above, data from 93 subjects

were analysed.

**2.3 General procedure.**

This study was implemented in the Gorilla Experiment Builder platform

(www.gorilla.sc) (Anwyl-Irvine et al., 2020) and delivered across three sessions: day 1 (d1),

day 8 (d8) and month 6 (m6) (Fig. 1B). Participants were initially recruited only for d1 and d8.

They were later invited to participate in the m6 session. Participants were recruited via the

Prolific platform and remunerated based on an hourly wage of £ 8. To enhance the precision

of online RT measurements and minimize the potential interference arising from variances in

devices and browsers (as outlined by Anwyl-Irvine et al., 2020; Bridges et al., 2020), we

implemented a controlled approach. Only participants equipped with a computer (excluding

tablets or phones) and using the 'Chrome' browser were eligible to take part in the study.

Offline checks of the OS used by participants revealed no systematic bias between older and

younger participants (see Table S1 in supplementary material). Potential biases in laptop vs

desktop computers or OS versions were not assessed. Participants who performed below

70% accuracy in the practice of the main ApMEM task (see below) were prevented from

continuing and received partial compensation for the time spent on the experiment.

On **day 1 (60 minutes)**, participants first completed a headphone check (Milne et al

2020; strict test version). The test is based on a binaural pitch signal that is only audible over

headphones (i.e., where L and R audio channels are delivered separately to each ear).

Passing the test requires reasonable quality audio equipment (headphones with separate R

and L channels) and preserved binaural hearing (Sanchez Lopez et al., 2018). People who

failed this very first stage were excluded from the analysis. Next, participants performed the

Auditory pattern Memory task (ApMEM; 3 blocks). The main task was preceded by a short

practice with a simplified version of the stimuli (see below). People who did not reach 70%

accuracy in this practice stage were stopped from continuing the experiment and received

partial compensation. ApMEM was followed by a series of cognitive tests - Sustained Attention

to Response Task (SART), the Corsi blocks task, and the Choice Reaction Time (CRT) task -

presented in random order across participants (Fig. 6A). More details about each task are

provided below. At the end of the session, participants completed a short questionnaire about

their listening environment and equipment. This included 4 choice questions about how old (1

year/2-4 years/5-7 years/8 or more), and how expensive their computer was (less than £700

/ between £700 and £1000 / between £1000 and £2000 / more than £2000). We observed no

between-group differences associated with these two ratings (W = 2100, p-value = 0.711; W

= 1855, p-value = 0.100). Participants were also asked about their physical activity habits

(numbers of hours per week) and years of musical training (ranked as 0, 0.5, 1, 2, 3-5, 6-9, 10

or more).

On **day 8 (15 minutes)**, participants completed the headphone check followed by a

single ApMEM block. The session ended with a surprise familiarity test for REGr (see details

below). **6 Months** later participants who completed d1 and d8 were re-invited for another

surprise session **(15 minutes)**. This included the headphone check and 1 block of ApMEM.

**2.4 Tasks.**

**2.4.1 Headphone check:** This test was used to exclude from the analysis participants with

poor sound environments. We used the strict version of the test freely available online

(https://gorilla.sc/ openmaterials/100917) and thoroughly documented in (Bianco et al., 2021;

Milne et al., 2020).

**2.4.2 ApMEM task.** The ApMEM task was used to measure multiple stages of auditory

memory (Bianco et al., 2020a). Stimuli (Fig. 1A) were sequences of contiguous 50-ms tone-

pips of different frequencies generated at a sampling rate of 22.05 kHz and gated on and off

with 5-ms raised cosine ramps. Twenty frequencies (logarithmically spaced values between

222 and 2,000 Hz; 12% steps; loudness normalised based on iso226) were arranged in

sequences with a total duration varying between 5.5 and 6 s. The specific order in which these

frequencies were successively distributed defined different conditions that were otherwise

identical in their spectral and timing profiles. We created 5 different stimulus sets which

participants were randomly assigned to. Each set contained five conditions as follows. RAN

('random') sequences consisted of tone-pips arranged in random order. This was implemented

by sampling uniformly from the pool with the constraint that adjacent tones were not of the

same frequency. Each frequency was equiprobable across the sequence duration. The

RANREG (random-to-regular) sequences contained a transition between a random (RAN),

and a regularly repeating pattern: Sequences with initially randomly ordered tones changed

into regularly repeating cycles of 20 tones (an overall cycle duration of 1 s; new on each trial).

The change occurred between 2.5 and 3 s (random jitter) after sequence onset such that each

RANREG sequence contained 3 REG cycles. RAN and RANREGn (RANREG novel)

conditions were generated anew for each trial and occurred equiprobably. Additionally, and

unbeknownst to participants, 3 different REG patterns reoccurred identically several times

within the d1, d8 and m6 sessions (RANREGr condition, reoccurring). The RAN portion of

RANREGr trials was always novel. Each of the 3 regular patterns (REGr) reoccurred 3 times

per block (every ~ 2 minutes, i.e., 9 presentations in d1, 3 in d8, and 3 in m6). Reoccurrences

were distributed within each block such that they occurred at the beginning (first third), middle

and end of each block. Two control conditions were also included: sequences of tones of a

fixed frequency (CONT), and sequences with a step change in frequency partway through the

trial (STEP). The STEP trials served as a lower-bound measure of individuals' RT to simple

acoustic changes. They were also used as attention checks – no, or very slow (see below)

responses to STEP trials indicated insufficient task engagement.

Each session of the main task was preceded by a volume adjustment stage.

Participants heard a few sounds from the main task and were instructed to adjust the volume

to a comfortable listening level. In the main task, participants were instructed to monitor for

transitions (50% of trials) from random to regular patterns (RANREG) and frequency changes

in STEP stimuli and press a keyboard button as soon as possible upon pattern detection. On

day 1, to acquaint participants with the task, two practice runs were administered. The first

practice contained 24 sequences consisting of simplified versions of the stimuli (10 RAN, 10

RANREGn, 2 STEP, 2 CONT), in that sequences were presented at a slower tempo (10 Hz)

and contained regularities of 10 tones. The second practice consisted of 21 sequences (9

RAN, 9 RANREGn, 2 STEP, 1 CONT) presented at a faster tempo (20 Hz) and containing

regularities of 20 tones, as in the main task. The main task consisted of 3 blocks on d1, 1 on

d8 and 1 on m6 sessions. Each block lasted about 6 minutes and contained 43 stimuli (18

RAN, 9 RANREGn, 9 RANREGr, 5 STEP, 2 CONT), with an inter-trial-interval of 1 s.

Participants were instructed to respond as quickly and accurately as possible both to the

transition from random to regular pattern and to the step frequency change. Feedback on

accuracy and speed was provided at the end of each trial as in our previous work (Bianco et

al., 2020a): a red cross for incorrect responses, and a tick after correct responses. The colour

of the tick was green if responses were 'fast' (< 2200 ms from transition to REG or <500 ms

from the step frequency change), and orange otherwise. The inter-block intervals were set to

have a maximum duration of 3 minutes to keep the overall duration of the exposure equal

across participants. Altogether, on day 1 instructions and practice took approximately 20 min

and the main task lasted 18 minutes. On day 8 and month 6, the ApMEM task took 8 min, 2

of which consisted of 20 trials of practice.

d' (computed per session across RANREGn and RANREGr conditions) served as a

general measure of sensitivity to regularity. Responses were marked as hits when they

occurred after the pattern begins to repeat (i.e., after the first cycle; effective transition).

Responses that occurred earlier or during random trials were marked as false alarms.

Participants whose d' was smaller than 1.5 in at least one session were excluded from the

analysis as this indicated poor pattern sensitivity and made RT uninterpretable. The core

analysis focused on the RTs to the onset of regular patterns and followed the pipeline adopted

in (Barascud et al., 2016; Bianco et al., 2020a). RT was defined as the time difference between

the onset of the regular pattern or the frequency step change and the participant's button

press. For each participant and block, RTs beyond ±2 SD from the mean were discarded

(OLD: 4.3% in REG, 4.1 in REGr; YOUNG: 4.1% in REG, 3.8% in REGr). Individuals identified

as outliers in the RTs to the STEP condition were excluded from the analysis as this indicated

low task engagement. The median STEP RT computed per session was used as a measure of the latency of the response to a simple acoustic change. This measure of individual baseline was stable across sessions (day 1 vs day 8 OLD: V = 991, p = .267; YOUNG: V = 1012, p = .854), and was subtracted from the RTs to RANREGn and RANREGr to yield a lower-bound estimate of the computation time required for pattern detection. Lastly, for each subject, we computed indexes of RT advantage (RTA) of REGr over REGn to quantify memory at different time points. To do so, we first corrected the RTs to REGr trials by the median RTs to REGn in each block. Then, to calculate the RTA by block, we computed the median RTA across the 3 intra-block presentations and the 3 different REGr patterns.

**2.4.3 ApMEM familiarity task.** Explicit memory for REGr was examined with a surprise task at the end of day 8. The 3 REGr patterns presented in the ApMEM (only one instance per REGr) were intermixed with 18 REGn patterns, as in (Bianco et al., 2020a). Participants were instructed to indicate which patterns sounded 'familiar'. The task took approximately 2 minutes to complete. The classification was evaluated using the Matthews Correlation Coefficient (MCC) score which ranges between 1 (perfect classification) to -1 (total misclassification) (Boughorbel et al., 2017; Powers, 2007). Before starting the task, participants were played a few sounds similar to those played in the upcoming task and asked to adjust the volume to a comfortable listening level.

**2.4.4 Choice reaction time task (CRT).** The CRT task is an established measure of individual variability in processing speed and is known to be linked with age-related decline in higher-level cognitive functions (Salthouse, 1996). Subjects were required to respond as soon as possible with the index or middle finger to a cue appearing with equal probability on the left or right box displayed on the screen. The task comprised 20 trials and took approximately 1 minute to complete. The task has two outcome measures: The central tendency (the median RT), and intraindividual variability (the raw standard deviation of the RTs), known to show a marked increase with age (Der and Deary, 2006; Hultsch et al., 2002).

**2.4.5 Corsi blocks (visual-sequence memory) task.** Nine identical black squares were

presented on the screen. On each trial, following a fixation duration (500 ms) a few blocks

flashed (briefly changed colour from black to yellow; flash duration 500ms; inter-flash-interval

250 ms) in a sequence. Participants had to reproduce the order of the sequence by mouse-

clicking on the correct blocks. The initial sequence length was 2 blocks. Correct responses

resulted in a length increase and incorrect responses in a length decrease. Overall participants

completed 20 trials. The task took approximately 5 minutes to complete. As an outcome

measure, we computed the mean sequence length. This score is considered to reflect the

ability to remember the temporal order of spatial sequences and it is known to deteriorate with

ageing (Beigneux et al., 2007; Bianco and Chait, 2022; Fournet et al., 2012).

**2.4.6 Sustained Attention to Response test (SART).** The ApMEM task is attentionally

demanding and memory formation may be affected by the listener's capacity to sustain

focused attention. The SART task was used to measure individual vigilance and propensity to

inattention (Manly et al., 2000). Participants were asked to respond by pressing a button to

serially and frequently presented 'go' visual stimuli (digits from 0 to 9, except 3) but maintain

a readiness to withhold a response to rare and unpredictable no-go trials (the digit 3). The

task took approximately 8 minutes to complete. The key outcome measure was the % 'no-go'

fail – quantifying listeners' ability to successfully stay "on task".

**2.6 Statistical analyses.** Performance was statistically tested with linear analyses of variance

(ANOVA) implemented in the R environment using the 'ezANOVA' function (Michael

Lawrence, 2016). P-values were Greenhouse-Geisser adjusted when sphericity assumptions

were violated. Post hoc t-tests were used to test for differences in performance between

conditions across blocks and groups. A Bonferroni correction was applied by multiplying p

values by the number of comparisons. Resulting values below the significance level of .05 are

indicated as n.s. – non-significant. Non-parametric tests were used where the normality of the

outcome distribution and homogeneity of variances were violated based on the Shapiro-Wilk

test. To isolate the contributions of different tasks to ApMEM performance, we used

hierarchical linear regressions.

**2.6 PPM-decay modelling.** Observed data from the ApMEM task were computationally

modelled using a memory-constrained Prediction by Partial Matching (PPM) model. PPM is a

variable-order Markov modelling technique that estimates the likelihood of the occurrence of

symbolic sequential events, given the number of occurrences of n-grams of varying size within

a training sequence, smoothing between models of different orders (Bunton, 1996; Cleary and

Witten, 1984).

Conventional models using PPM possess a perfect memory for all events in their

training data, regardless of proximity to the modelled event. To model the effects of human

memory on learning, Harrison et al. (2020) implemented a PPM model with the ability to down-

weight occurrences in the model over time, based on a customisable decay kernel. As used

here, the kernel contained three phases: (1) a high-fidelity echoic memory buffer, defined by

a weight and a duration; (2) a short-term memory (STM) phase that decays exponentially from

the weight of the buffer to the starting weight of the next phase over a given duration; and (3)

an exponentially decaying long-term memory (LTM) phase, defined by a starting weight and

a half-life; (examples of decay kernels and their phases can be seen in Fig. 3C). Additionally,

varying levels of noise were added to event probabilities, fulfilling the role of a general

processing speed parameter, and replicating similar imperfections in human memory. This

noise was parameterised by the SD of a normal distribution, from which sampled absolute

values were added to the weights on memory retrieval.

All stimuli in blocks 1 to 3 of day 1 were modelled, as presented for each stimulus set,

maintaining the tone, stimulus, and block timings of the task. Models were trained dynamically,

estimating a probability for each tone, given the sequence preceding it and all preceding

stimuli, which was converted into information content (negative log-base-2 probability). Models

were limited to a maximum n-gram length of 5 symbols (an order bound of 4). As in (Bianco

et al., 2020a), changes in information content were identified for REGn and REGr stimuli using

the nonparametric change-point detection algorithm of (Ross et al., 2011), a sequential

application of the Mann-Whitney test, while controlling for a Type I error rate of 1 in 10,000.

Model parameters were optimised to find the decay configuration that best reproduced

the observed data of the younger group using Rowan's Subplex algorithm, as implemented in

the NLopt package (Johnson, 2020; Rowan, 1990). Initial parameter values were adapted

from the manually fitted parameters by Bianco et al. (2020a). To account for the increased

variability of change points due to modelling prediction noise, for every optimisation iteration,

the modelling was repeated 100 times, refreshing model memory between each. Repeated

change points were then averaged for individual stimuli. Optimisation sought to minimise the

root-mean-square error (RMSE) between observed RTs and modelled change points, when

averaged for each of the REGn and REGr conditions within blocks 1 to 3, for each different

stimulus set. To facilitate a comparison of goodness of fit between models using different

numbers of parameters to fit the same data, Akaike Information Criterion (AIC) values were

calculated under the assumption of normally distributed errors. A correction was applied to

penalize potential overfitting when using small sample sizes (AICc).

To characterise differences between the older and younger groups, first, observed

data for the older group were modelled by optimising a single parameter while holding all

others to the values obtained for the younger group. The fit of these models is shown in Fig.

3B. As no single change of only an individual parameter adequately reproduced the observed

data, pairs of parameters were then fitted. For each pair, the parameter giving the best fit when

optimised individually was initialised at the value obtained in that optimisation. The other

parameter in the pair was initialised at, and all remaining parameters were held at, the values

obtained for the younger group. The parameters that produced the best-fitting of these models,

based on RMSE, were selected as those characterising the older group.

**3. Results**

With an online version of ApMEM, we characterised implicit memory for rapid tone

patterns in young and older participants over multiple timescales, from early mnemonic stages

required to detect novel patterns to long-term memory formation and retention at 1-week (d8)

and 6 months (m6). Using a computational model of auditory sequence processing, we then

distilled and quantified how different memory parameters contribute to between-group

differences in performance on day 1 (d1).

To account for age-related effects on general aspects of the ApMEM task (ApMEM

task is RT-based, measures memory for sequences and requires focused attention) we also

included tests of processing speed (Salthouse, 1996), spatial-visual sequence memory (Corsi,

1972; Kessels et al., 2000), and attention (Manly et al., 2000) (Fig. 1B). We explored whether

these measures might explain response differences in ApMEM performance between the age

groups.

**441    3.1 No group difference in accuracy of pattern detection**

Overall, the ability to detect the emergence of a pattern from a random sequence, as quantified

with d' (collapsed across REGn and REGr), was consistently high across blocks (Fig. 2B),

though a drop between d8 and m6 was seen in the younger group (YOUNG: W = 796, p =

0.003, CI [-0.642 -4.953e-05; OLD: W = 1628, p = 0.208, CI [-0.321 3.74e-05]). d' was similar

between groups on d1 (W = 2524.5, p = .113, CI [-.006 .231], mean OLD: 3.3 ± .40, YOUNG:

3.17 ± .454), on d8 (W = 2036, p =.513, CI [-0.321 7.035e-06], mean OLD: 3.34 ± .48, YOUNG:

3.36 ± .558), and on m6 (W = 1265, p = 0.082, CI [-1.999e-05  .496], mean OLD: 3.22 ± .55,

YOUNG: 2.97 ± .67). This confirms high sensitivity to the presence of regularities and allows

us to confidently interpret the between-group differences in RTs as a measure of memory

strength.

The hit rate for novel (REGn) and reoccurring patterns (REGr) was computed for each

session (d1, d8, m6). Hits were higher for REGr than REGn on d1 (mean over blocks 1,2 and

3) in both groups (REGr: 96.6±4.36, REGn: 94.1±7.7 in OLD; REGr: 97±4.57, REGn: 93.6±7.9

in YOUNG; one-sample Wilcoxon test of REGr – REGn hit per cent, OLD: V = 596, p = .006;

YOUNG: V = 807, p < .001), indicating a memory effect for REGr. The hit advantage of REGr

over REGn did not differ between the groups on d1 (W = 2439.5, p = .218, CI [-2.86e-05 3.70]),

on d8 (W = 2341.5, p= .39, CI [-1.38e-05 3.97e-05]), and m6 (W = 1118.5, p = .531, CI [-

3.47e-05 3.44e-05]). Finally, the hit advantage did not change over time: similar effects were

found in the last block of d1 (b3_d1), d8 and m6 in both groups (all p < .78).

Overall, both groups were highly accurate in detecting the patterns and showed a

memory advantage for REGr. However, this advantage was similar between groups and did

not change over time, likely because it quickly reached ceiling effects. The ApMEM task was

designed to yield high levels of accuracy consistently over time. This enabled us to focus on

analysing the RT to capture processes related to dynamic sequence processing. Below we

demonstrate that between-group differences associated with these effects are sensitively

captured when focusing on RT.

**3.2 Age-related decline in early mnemonic stages**

RTs to simple frequency changes (STEP) collapsed across d1, d8 and m6 showed

substantial inter-individual variability and were also generally slower in the OLD than YOUNG

group (Fig. 2C; W = 2766.5, p = .007, CI [15.9 92.9], mean OLD: 482 ± 115, YOUNG: 434 ±

108 ms). For each subject, the RT to the pattern emergence (RANREGn and RANREGr) was

corrected by the RT to the STEP (median per session). This was done to control for individual

variability in RT to a simple stimulus change, and thus isolate the computation time required

to detect an emerging pattern.

We first analysed responses to REGn, as a measure of early mnemonic stages.  We

computed the median RT to REGn, collapsed across all sessions (Fig. 2D). The OLD group

took longer than the YOUNG to detect the REGn patterns (RTs to REG: W = 2654.5, p = .03,

CI [5.49 106.25]; mean OLD: 1807 ± .152; YOUNG: 1747 ± 137 ms), suggesting age-related

decline of early mnemonic components (echoic/short-term memory) supporting pattern

detection.

**3.3 Age-related decline in long-term memory formation**

Fig. 2E shows the median RT to REGn vs REGr for each block and session. We first

focused on day 1 (memory formation stage). A mixed ANOVA with factors condition

(RANREGn / RANREGr), block (block 1-3 of d1) and between-subjects factor group

(OLD/YOUNG) yielded a main effect of condition [$F(1, 130) = 82.52$, $p < .001$, $\eta_p^2 = .39$], a

main effect of block [$F(2, 260) = 6.87$, $p = .002$, $\eta_p^2 = .05$] and an interaction of condition by

block [$F(2, 260) = 7.57$, $p = .001$, $\eta_p^2 = .06$]. This confirms the general pattern previously

observed for this task (Bianco et al., 2020a): whilst RT to REGn patterns remains stable across

blocks (no significant difference between blocks), RT to REGr becomes progressively faster

with repeated exposure (block 1 vs 2 $p = .004$; block 1 vs 3 $p < .001$).

A main effect of group [$F(1, 130) = 8.71$, $p = .004$, $\eta_p^2 = .06$] and interaction of condition

with group [$F(1, 130) = 4.74$, $p = .031$, $\eta_p^2 = .04$] confirmed that, across the 3 blocks of d1, the

older group were slower overall than the younger group in detecting the REGr patterns (RT to

REGr: OLD, mean 1725 ± 182 ms, vs YOUNG, 1622 ± 164, [$t(130) = 3.40$, $p < .001$]). An

effect of ageing on REGn RT did not quite reach significance when focusing on the first day

only (RT to REGn: OLD, mean 1809 ± 173 ms, vs YOUNG, mean 1761 ± 138 ms, [$t(130) =$

1.75, $p = .08$]), possibly due to some noise in the YOUNG data in block 2. No three-way

interaction was found [$F(2, 260) = 0.76$, $p = .467$, $\eta_p^2 = .01$].

*Figure 2. Auditory memory formation in young and older adults (ApMEM task). A)*
*Participant age distribution. B) d' (sensitivity to the emergence of regularity) for the OLD and*
*YOUNG groups. Shaded areas indicate the different stages of memory formation (d1 in grey)*
*and retention (d8, m6 in yellow). C) Median RT to STEP trials (frequency step changes)*
*averaged across the d1, d8 and m6 sessions. (D) Across all sessions (d1, d8 and m6), older*
*listeners exhibited slower REGn RT than young controls. E) RTs to REGr and REGn across*
*the 3 blocks of d1, and the 1 block of d8 and m6. Error bars represent the standard error of*
*the mean.*

**3.4 Computational modelling indicates that reduced performance among older listeners**
**can be explained by reduced echoic buffer weight and faster LTM decay**

Using a memory-constrained variant of Prediction by Partial Matching (Harrison et al.,
2020), a computational model was optimised to fit the observed data on day 1 over blocks 1
to 3 of the ApMEM task for both OLD and YOUNG groups. We use this model to provide a
formal simulation of early memory encoding and long-term memory formation characterising
differences between the groups.

Fig. 3A shows the simulated RTs and the parameters optimised to fit the data of the

YOUNG group (RMSE = 56.85). The decay kernel that these parameters generate is

illustrated in Fig. 3C (right plot). Qualitatively, these parameters show close correspondence

to those obtained when simulating responses of young participants on the same task in

(Bianco et al., 2020a).

The optimisation of the data of the OLD group first examined whether a single

parameter change from the values obtained for YOUNG could explain the differences in RT.

The individually optimised parameters and their fit are given in Fig. 3B. It should be noted that,

in several cases, the change of a single parameter affects the characteristics of multiple

memory phases. No optimisation of only a single parameter managed to adequately fit the

observed data for the OLD group, with all models possessing both high RMSEs reflecting an

inability to recreate the trajectories of REGn and REGr responses, as displayed in Fig. 3B (left

plot). In particular, parameters affecting the buffer, STM, or overall prediction noise, were

unable to reproduce the decrease in learning rate exhibited in block 3 of the observed data for

the REGr condition. While LTM weight was better able to account for this effect, it could not

sufficiently increase simulated RTs for the REGn condition at the same time. Manipulating

LTM half-life on its own was unable to produce a fit for either condition.

Next, pairs of parameters were optimised in turn to fit the data for the OLD group (a

full list of parameter values and fit of models is given in supplementary material). Four

parameter sets had better goodness of fit than the best-fitting single parameter optimisations

when accounting for the increased number of parameters (i.e., with an AICc lower than that

of LTM weight at 249.54). Each could plausibly fit the observed data, and each contained one

parameter controlling properties of the buffer or short-term memory, and one controlling an

aspect of long-term memory. The best-fitting combination (RMSE = 57.22; AICc = 247.26)

was the optimisation of buffer weight and LTM half-life (0.93 and 232.86 s, respectively). The

decay kernel described by these parameters diverges from that of the YOUNG group by

having some down-weighting of memories formed within the buffer and more rapid long-term

decay, as shown in Fig. 3C (right). These differences, and those of the other low-RMSE

models fitting the OLD data (buffer duration and LTM half-life; buffer duration and LTM weight;

STM duration and LTM half-life), indicate that the older group possesses weaker memory

formation in both the immediate and long-term mnemonic phases and that both deficits are

required to explain the differences observed between the two groups.

***Figure 3. Modelling contribution of buffer, STM and LTM to auditory memory formation***

***in young and older adults. A)*** *Simulated and observed RTs to REGn and REGr conditions*

*for the YOUNG group (left), and the parameter values of the optimised model with RMSE =*

*56.85 (right).* ***B)*** *Simulated RTs and parameter values for models fitting OLD RTs by optimising*

*individual parameter changes from those of YOUNG.* ***C)*** *Simulated and observed RTs to*

*REGn and REGr conditions for the OLD group using parameter values obtained for YOUNG,*

*modifying buffer weight and LTM half-life yielding RMSE = 57.22 (left), and the memory decay*

*kernels of best fitting models for both groups on a time scale of 30 seconds (right). The small*

 *insert illustrates the LTM phase on a timescale of 1 hour. See Supplementary Material for*

*simulated RTs using tone intervals, instead of absolute tone values.*

**3.5 Auditory memory retained for up to 6 months in both older and younger listeners**

Fig. 4A displays the RT advantage between REGr and REGn across all experimental sessions. The data revealed that the difference in RTA observed at the end of d1 (and modelled above) persisted when probed at d8 and at m6.

To explicitly test for long-term memory retention, for each subject we compared the RTA at d8 and m6 to that observed in b1 (block 1 of d1, b1_d1). If listeners retained a lasting memory of REGr we expected RTA in d8 and m6 to be different from that in b1_d1. An ANOVA with factors block (b1_d1/ d8) and group yielded main effects of group [$F(1, 130) = 8.69$, $p = .004$, $\eta_p^2 = .06$], block [$F(1, 130) = 26.30$, $p < .001$, $\eta_p^2 = .17$], and no interaction [$F(1, 130) = 3.24$, $p = .074$, $\eta_p^2 = .02$], indicating a greater RTA in d8 than in b1_d1 in both groups [$t(131) = 5.03$, $p < .001$; mean RTA OLD: b1_d1 36.4 ± 189, d8 115 ± .190 ms; YOUNG: b1_d1 71.6 ± 215, d8 234 ± 215 ms], and overall greater in the YOUNG than in the OLD group [$t(130) = 2.94$, $p = .004$].

An ANOVA with factors block (b1_d1 / m6) and group yielded main effects of group [$F(1, 91) = 10.82$, $p = .001$, $\eta_p^2 = .11$], block [$(1, 91) = 16.59$, $p < .001$, $\eta_p^2 = .15$], and no interaction [$F(1, 91) = 1.52$, $p = .221$, $\eta_p^2 = .02$], indicating a greater RTA in m6 than in b1_d1 in both groups [$t(92) = 3.90$, $p < .001$; mean RTA OLD: b1_d1 30.2 ± 198, m6 119 ± 189 ms; YOUNG: b1_d1 89.5 ± 197, d8 255 ± 239 ms], and an overall greater RTA in the YOUNG group [$t(91) = 3.28$, $p = .001$]. This analysis shows that weaker LTM memory is formed in the OLD than in the YOUNG group, but both groups maintain non-decaying memories of the REGr patterns up to 6 months following initial memory formation.

Lastly, we compared the RTA in d1 block 3 (b3_d1), to those in d8 and m6 (Fig. 4B). A repeated measures ANOVA revealed a main effect of group only [$F(1, 91) = 13.86$, $p < .001$, $\eta_p^2 = .12$], consistent with the overall larger RTA among the young listeners. There was no main effect of block, nor an interaction (block: [$F(2, 182) = .97$, $p = .381$, $\eta_p^2 = .01$];

interaction: [$F(2, 182) = .93$, $p = .397$, $\eta_p^2 = .01$]), confirming a plateauing of the RTA after d1

in both groups – consistent with an enduring memory trace.

We also found that RTA in b3_d1 positively correlated with RTA in d8 (spearman's rho

= 0.198, $p = 0.022$) and that RTA in d8 correlated with RTA in m6 (spearman's rho = 0.230, p

= 0.026). This indicates good reliability of individual effects even in online settings.

Finally, we found no correlations between the age of the participants and the RTA in

b3_d1, d8, or m6, in either the OLD (all $p < .06$), nor the YOUNG group (all $p < .19$). This may

be due to the narrow age ranges used in the current study. However, the observed negative

trend in the correlation between age and RTA in m6 in the older group (rho = -0.255, $p = .06$)

suggests that future studies with larger sample sizes and a broader age range may potentially

reveal critical age periods in implicit auditory memory decay.

To summarise, an effect of group (young vs. older) emerged across different time

scales. In addition to weaker early mnemonic stages (Fig. 2D), the older group exhibited

weaker long-term memory, as reflected by smaller RTA in b3_d1, d8 and m6. However, there

was no evidence of a decline in memory e.g., between b3_d1 and d8 or d8 and m6.

**Figure 4. Auditory memory formation and retention in the young and older groups**

**(ApMEM task). (A) Long-term** *memory dynamics across d1, d8 and m6 in young and older*

*listeners, quantified as RT advantage (RTA) of REGr over REGn patterns.* **(B)** *Violin plots*

*showing the individual data for the indexes extracted from the ApMEM task to quantify memory*

*formation (the RTA computed over the last block of the first exposure on d1) and memory*

*retention (the RTA computed on d8 and m6). Statistically significant differences between*

*group means (Mann-Whitney U) are indicated.*

**3.6 No link between explicit and implicit auditory memory formation.** Explicit memory for

the REGr patterns was assessed at the end of d8 with a surprise familiarity task (Fig. 5A).

Each REGr was presented once only amongst a large set of foils (REGn) and participants

judged if the pattern was familiar. MCC (Matthews correlation coefficient) was used to

measure the quality of subjects' binary classification. Both groups exhibited above-chance

performance (OLD: V = 1734, p < .001; YOUNG: V = 1765.5, p < .001), but the performance

was poorer in older listeners (W = 1589, p = .040, CI [-.137 -5.16e-05], mean MCC OLD: .173

±.217, YOUNG: .253 ± .183) (Fig. 5A). As in our previous findings (Bianco et al., 2020a),

explicit memory scores did not correlate with the RTA observed in the last block of d1 (b3_d1;

spearman's Rho = -.025; p = .777) nor with that in d8 (spearman's Rho = .056; p = .170).

These analyses confirm the implicit nature of the RTA measures.

**3.7 Age-related decline in visual-sequence memory and processing speed, but no link**

**with ApMEM RTA.** At the group level, older participants showed slower median RTs in the

CRT (W = 4000, p < .001, CI [68.80 100.09], mean OLD: 421 ± 97.2; YOUNG: 319 ± 40.5),

and greater variability (SD of trials: W = 3227, p < .001, CI [11.63 27.06]) (Fig. 5B-C), reflecting

a well-known effect of age-related processing speed impairment (Hultsch et al., 2002). The

median RTs in CRT correlated with the RTs in our control STEP condition (Rho = .312, p <

.001) confirming that STEP RTs are a good measure for correcting differences in baseline

speed.

The older group exhibited worse performance in the Corsi blocks task (OLD vs

YOUNG: t(130) = -3.56, p < .001, mean OLD: 4.54 ± .59; YOUNG: 4.80 ± .537), confirming

the expected age-related decline in visual sequence memory (Fig. 5D) (Beigneux et al., 2007;

Bianco and Chait, 2023; Fournet et al., 2012).

In line with previous findings on the SART (de Kerangal et al., 2021), there was no

age-related decline in the sustained attention accuracy (% 'no-go' fail: W = 1967, p = .343, CI

[-11.10    3.10], mean OLD: 33.5 ± 20.3; YOUNG: 38.1 ± 24.5) (Fig. 5E). As expected, RTs

were slower in the OLD vs the YOUNG group in the 'go-trials' (W = 11073, p < .001, CI: [19.62

52.54], mean OLD: 378 ± 66.5; YOUNG: 348 ± 85.7). The speed-accuracy trade-off was

similar between groups: accuracy in 'no go' trials was predicted by RT ($\chi^2$(1) = 25.29, p <

.001), but not by group or the interaction between RTs and group (p > 0.1).

We conducted linear regression analyses to understand to what extent group-specific

variance in ApMEM is predicted by performance on these cognitive tasks. We also included

weekly hours of physical activity and years of musical training as possible predictors of

ApMEM performance. Physical activity has been listed among the factors reducing the risk of

cognitive and memory decline (Nyberg and Pudas, 2019; Sofi et al., 2011). Evidence from a

meta-analysis has linked musical practice in healthy ageing with cognitive benefits both in

domain-specific functions (auditory perception) and more general ones (Román-Caballero et

al., 2018). For each outcome measure of ApMEM (Fig. 4B) and group, we performed a linear

regression analysis with the predictors: CRT standard deviation, SART RTs, Corsi mean

sequence length, and the above-mentioned demographic scores. None of the models was

significant in the older group (all p-values > .11). A similar analysis in the younger cohort also

yielded non-significant models (all p-values > 0.14). Overall, this pattern of results indicates

that the variability in ApMEM is not driven by general processing speed, visuospatial

sequential memory, or sustained attention, and might thus reflect age-related deficits specific

to auditory memory.

*Figure 5. Distribution of performance, across the various tasks, in the young and older*

*groups. Statistically significant differences between group means (Mann-Whitney U) are*

*indicated. Worse performance in the OLD than in the YOUNG group was observed in (A)*

*ApMEM explicit familiarity scores computed as Matthews correlation coefficient (MCC) on the*

*reoccurring patterns and obtained at the end of day 8. The coefficient ranges between 1 (perfect classification) to -1 (total misclassification). **(B)** CRT both in terms of median and **(C)** variance of RTs, and **(D)** Corsi-blocks. Accuracy on the SART task **(E)** did not differ between groups. The two groups did not differ in terms of **(F)** musical training (W = 2076, p = 0.580).*

## 4. Discussion

Despite its role in supporting fundamental aspects of auditory perception, how implicit sensory memory is affected by ageing remains poorly understood. Existing work has predominantly focused on verbal material and tasks that require cued reporting which are susceptible to factors such as rehearsal strategies or interference during long retention periods. For example, it is not known whether the "accelerated forgetting" effect recently characterised in older listeners (Elliott et al., 2014; Wearn et al., 2020) is specific to explicit memory or also extends to more basic auditory mnemonic representations.

Here, we introduce a paradigm that distils and quantifies age-related deficits in core memory mechanisms underlying auditory scene analysis. We compared younger (aged between 20 and 30 years old) and older adults (aged between 60 and 70 years old) with an RT-based implicit memory test. Participants are required to detect emerging regular patterns from random rapid sequences of tones. Patterns are novel in most trials (REGn), but unbeknownst to the participants, a few distinct patterns reoccur identically throughout the experimental sessions (REGr). The progressively growing RT advantage for REGr vs REGn demonstrates that mnemonic traces for the specific reoccurring patterns become more salient in memory through repeated presentation. The test is repeated 8 days and 6 months after initial exposure. Notably, the stimuli are arbitrary, and too fast to allow conscious tracking of the sequence events; thus, minimizing active tracking processes and mnemonic interference with real-world sounds.

We found that compared to young adults, older participants were slower in detecting novel patterns and exhibited a smaller RT advantage in detecting reoccurring patterns,

indicating deficits in echoic/short-term memory and long-term memory formation. A

computational model of auditory sequence memory fit to the data on the first day of exposure

also suggested age-related limitations in both early and long-term mnemonic components.

However, in contrast to mounting demonstrations of accelerated forgetting of verbal material

with ageing, here older adults maintained stable memory traces for the reoccurring patterns –

an unaltered RT advantage – up to 6 months after the first exposure. An absence of a

correlation between the RT advantage and the active recall abilities measured with familiarity

ratings confirmed the implicit nature of the RT effects.

**4.1 Processing speed does not explain the reduced auditory memory effect in the older**

**cohort**

The between-group difference in the derived RT-based measures of memory cannot

be explained by general reduced processing speed in the aged cohort (Salthouse, 1996; Zajac

and Nettelbeck, 2018). This is supported by three arguments: First, RTs were corrected by

RT to a simple stimulus change (STEP condition) interspersed in the main ApMEM task. This

ensured that performance in pattern detection was controlled for inter-individual biological

(e.g., subject's general state of vigilance, or the time taken to perceive an auditory change, to

generate a response) or equipment-based differences (e.g., keyboard latency) introducing

non-memory specific variability. Second, the worse performance of older than younger adults

in the control choice-RT task (CRT) showed the expected age effect on processing speed,

and it correlated with the ApMEM STEP control condition, confirming the latter as a valid

measure for correcting general differences in baseline speed. Finally, the regression analyses,

including control tasks as predictors of ApMEM performance, showed that CRT did not

contribute to variability in any of ApMEM-related memory measures. This, together with the

absence of correlations between the ApMEM memory measures and tasks associated with

sustained attention (SART) and visual-sequence memory suggest that between-group

differences in ApMEM reflects age-related deficits specific to auditory memory.

Online RT measurements are subject to limitations due to the variability in the

participants' equipment (Anwyl-Irvine et al., 2020; Bridges et al., 2020) and engagement

(Bianco et al., 2021). Although we implemented strict equipment requirements (see methods),

residual components of equipment-driven RT delay might still contribute to the RT variance

when measured online. Importantly, this variance is likely to be equally distributed across the

groups (see methods and table S2).

**4.2 Deficits in both early and long-term mnemonic stages contribute to age-related**

**performance decline**

Our measure of memory was RT which can be related to the computations inherent to

dynamic sequence processing, and hence to the model output. Whilst accuracy measures

were at ceiling by task design, older compared with younger participants exhibited overall

slower RTs in response to novel patterns and, over time, formed a smaller RTA for the

reoccurring patterns. Slower RTs in the older group could reflect a combination of slower

access to stored information and less confidence due to noisier memory encoding. Whilst

these factors could be reflected in both RT to REGr and REGn, the difference between the

two conditions (the RTA) is likely to specifically reflect long-term memory representations.

Modelling demonstrated that optimisation of parameters associated with both early and later

mnemonic stages was necessary to accurately model older listeners' performance.

We modelled the effect of ageing on memory by optimising the model memory decay

kernel pertaining to: (1) a high-fidelity echoic memory buffer; (2) an STM phase; and (3) an

exponentially decaying LTM phase.  Each of these is associated with parameters describing

their duration, relative weight, and rate of decay. Amongst the four best-fitting models, each

contained one parameter controlling properties of the buffer or short-term memory, and one

controlling an aspect of long-term memory. The best fitting model suggested that weaker

echoic buffer weight (YOUNG: 1; OLD: .93), as well as more rapid long-term decay (YOUNG:

502.96 s; OLD: 232.86 s), contributed to the age-related performance decline on day 1.

The first 'pre-perceptual' stage of temporarily holding auditory information allows

listeners to bind incoming events with the just heard ones to perceive a coherent

representation of sequential sounds (e.g., a sequence of single tones as a motive). The

computational modelling results support the interpretation that the overall slower detection of

REGn patterns in older adults is a consequence of limited buffer / short-term memory

components. This is in line with previous hypotheses (Herrmann et al., 2022) and consistent

with neurophysiological evidence showing a somewhat weaker ability to represent information

in echoic memory in older than young adults (Cooper et al., 2006; Pekkonen et al., 1996).

Parameters affecting the early stages of auditory memory alone could reproduce the

age effect on responses to novel patterns but were unable to reproduce the diminished RTA

for reoccurring patterns. A combination of reduced memory buffer and more rapid long-term

decay (LTM half-life) best accounted for both aspects of performance differences between

groups.

The decay kernel optimised for older adults' data provides an initial model of how

limitations at multiple stages of memory may explain different cognitive performance between

populations. We focused on PPM-decay as the primary model for benchmarking auditory

sequence tracking and learning. This model has successfully explained behavioural and

neural responses to stimuli like those used here (Barascud et al., 2016; Bianco et al., 2020a;

Harrison et al., 2020). Notably, relative to neural network (Huang et al., 2018) and Gestalt

models (Narmour, 1990), PPM excels in simulating brain responses to more complex signals,

such as music (Kern et al., 2022). Moreover, unlike alternative models (e.g., Skerritt-Davis

and Elhilali, 2021), to the best of our knowledge, PPM-decay is the only model that has explicit

decay parameters. In particular, the long-term decay seems to be necessary to simulate

listeners' performance on different auditory tasks (Bianco et al., 2020a; Harrison et al., 2020).

The PPM model also provides some inherent reproduction of interference effects which are

thought to co-exist with decay (Hardt et al., 2013; Lewandowsky et al., 2009).

The parameters for this model were optimised to fit blocks 1 to 3 of the ApMEM task

performed on d1. As the constant long-term decay of the model predicts that memory should

eventually reduce to zero, d8 and m6 are beyond the scope of this modelling in its current

form. Modelling such time spans, while still being able to recreate effects within the first three

blocks, would require a non-trivial addition that could account for memory consolidation over
the intervening time periods.

Could the observed age-related memory effects arise from poorer hearing sensitivity
in older listeners? We consider this unlikely for several reasons: firstly, the sensitivity to the
presence of regularities was high among older adults and did not differ between older and
young listeners. Secondly, during the instructions stage, participants were encouraged to
adjust the sound volume to as high a level as needed, and the tone frequency range of 220-
2000 Hz was intentionally chosen to be less affected by age-related hearing loss. This further
guaranteed adequate audibility for all sounds. Finally, only participants who passed the
headphone test (see methods) were included.

**4.3 No evidence of long-term forgetting with ageing: memory traces to arbitrary tone**
**patterns are retained for up to 6 months from initial exposure.**

The older compared with the young group formed weaker memory during the first day
of exposure as quantified by a smaller RTA. However, just as observed in the younger group,
the RTA in older listeners persisted for 8 days and 6 months after the initial exposure. This
very long-lasting effect observed in both groups is noteworthy considering that the RTA was
not driven by the explicit familiarity judgments obtained 6 months earlier.

There is growing interest in tests taxing memory circuit functionality at delayed recall
because memory problems at this stage could indicate incipient dementia (Ryan and
Frankland, 2022; Weston et al., 2018). That auditory patterns were not forgotten at delays of
8 days nor 6 months in the older cohort is in contrast to the body of work on accelerated long-
term forgetting (ALF) for verbal material in ageing (Davis et al., 2003; Elliott et al., 2014; Manes
et al., 2008; Mary et al., 2013; Wearn et al., 2020). ALF of verbal material has been reported
also in pre-symptomatic autosomal dominant Alzheimer's disease (Weston et al., 2018),
patients with temporal lobe epilepsy (Blake et al., 2000), and it may represent a failure of
memory consolidation processes (Hoefeijzers et al., 2013) due to altered integrity of
hippocampal-neocortical (temporal) connections (Alvarez and Squire, 1994). One explanation

for the discrepancy between verbal tasks and ApMEM may reside in the very low probability,

compared to verbal material, that subjects encountered ApMEM-like sequences outside of the

experimental sessions. This might have minimised phenomena such as forgetting due to

interference with real-world stimuli (Davis and Zhong, 2017), which perhaps affects older more than

younger adults (Wais and Gazzaley, 2014).

An alternative explanation can be found in the dichotomy between the active and

implicit nature of the memorization processes inherent in verbal memory tasks versus the

ApMEM task. While the former requires participants' active engagement in memorization and

subsequent recall, the latter relies on implicit memorization achieved through repetitive

exposure. Notably, the memory effects observed in the ApMEM task, as quantified by RTA,

did not correlate with the active recall of the REGr sequences in the familiarity task. This

observation is in line with demonstrations in the visuomotor domain that implicit learning

through repetition leads to memory retention for a remarkably long time (Kobor et al., 2017;

Nemeth and Janacsek, 2011; Tóth-Fábera et al., 2023). These findings collectively suggest

that implicit memory is rooted in robust biological substrates (Bailey et al., 2004; Ohno et al.,

2011) less vulnerable to the availability of processing resources, attention, or interference,

and so more preserved by ageing.

One important open question is why such long-term auditory implicit memory is overall

resilient to time decay even later in life.  The remarkable examples of preserved auditory

memory for music in severe cases of dementia (Baird and Samson, 2009; Benhamou et al.,

2021; Jacobsen et al., 2015) suggest that implicit auditory memory has a privileged status in

the brain. In young listeners, implicit auditory memory based on repeated exposure has been

demonstrated for many sound types, ranging from white noise (Agus et al., 2010; Dauer et al.,

2022; Ringer et al., 2022), click trains (Kang et al., 2017), discrete sequences of tones (Bianco

et al., 2020a; Bonetti et al., 2022; Herrmann et al., 2021; Leek and Watson, 1988), tone clouds

(Agus and Pressnitzer, 2021), and naturalistic textures (Ringer et al., 2022; Woods and

Mcdermott, 2018). All these studies capitalise on Hebb-type learning tasks (Hebb, 1961;

Reber, 1989), whereby regardless of subject awareness, recognition of reoccurring patterns

improves compared to novel ones simply due to reinforcement through repetition. Repetition

is perhaps the simplest cue-inducing learning because it indicates the presence of patterns

potentially relevant to behaviour (McDermott et al., 2011). Patterns have often a

communicative function and are indeed implicitly learned through repetition in humans (Aslin,

2017; Saffran et al., 1996; Smalle et al., 2018) and non-human animals (Cazala et al., 2019;

Hauser et al., 2001; Lu and Vicario, 2014; Soyman and Vicario, 2019; Wilson et al., 2013).

The long-term memory of tone-sequences observed here even in older adults might thus

reflect this primordial predisposition of the brain to remember patterns even when they

sparsely reoccur.

## 5. Conclusions

Presenting a multi-stage auditory memory test that is scalable and easy to administer,

we propose a paradigm shift in memory research beyond traditional verbal memory tests. We

demonstrated that ageing is associated with poorer auditory echoic/short-term memory and

long-term memory formation than young listeners, but not with forgetting. We speculate that

ageing might affect frontal-auditory and hippocampal circuits underlying memory formation,

but once formed auditory memories of rapid tone patterns remain accessible for months after

the initial exposure even in older listeners. This result might be explained by the absence of

interference with memory traces of arbitrary stimuli, unlikely to be encountered in daily life

(Hardt et al., 2013; Ryan and Frankland, 2022), and suggests preserved long-term implicit

auditory memory in ageing. Future studies combining human neuroimaging, animal models

and synaptic simulations should shed light on the underlying circuits and neuronal

mechanisms (Gershman, 2022; Ohno et al., 2011; Poeppel and Idsardi, 2022). Expanding the

work to include a wider age range, from children to the elderly, could offer valuable insights

into the currently understudied development and decline of auditory memory.

## Funding

857    This work was supported by the NIHR UCLH BRC Deafness and Hearing Problems Theme,

858    a BBSRC grant (BB/P003745/1) to M. C., an ARUK ECR bridging award and Marie

859    Skłodowska-Curie Individual Fellowship (MSCA-PF 101064334) to R. B.

## Data availability

861    The datasets for this study are shared via

862    https://osf.io/4d5se/?view_only=aa13d54c3ec248489bde41b9b562f470 and will be made

863    publicly available upon publication.

## Authors contribution

RB: Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation;

Methodology; Validation; Visualization; Roles/Writing - original draft; Writing - review & editing.

ETRH: Formal analysis; Methodology; Software; Validation; Visualization; Roles/Writing -

original draft; Writing - review & editing.

MTP: Methodology; Resources; Software; Supervision; Validation; Visualization;

Roles/Writing - original draft; Writing - review & editing.

MC: Conceptualization; Funding acquisition; Methodology; Project administration; Resources;

Supervision; Validation; Visualization; Roles/Writing - original draft; Writing - review & editing.

## Supplementary material

**Table S1. Counts of participants in the OLD and YOUNG groups per OS**

|         | Windows | Mac | Linux | Chromium |
|---------|---------|-----|-------|----------|
| OLD     | 54      | 7   | 2     | 5        |
| YOUNG   | 55      | 10  | 0     | 0        |

**Table S2. Optimising parameter pairs to OLD ApMEM blocks 1 to 3**

| Parameter 1 | Value 1 | Parameter 2 | Value 2 | RMSE | AICc |
|-------------|---------|-------------|---------|------|------|
| Buffer weight | 0.86 | Buffer duration | 0.97 | 64.76 | 254.69 |
| Buffer weight | 0.90 | STM duration | 14.90 | 63.20 | 253.22 |
| Buffer weight | 1.00 | LTM weight | 0.006 | 61.34 | 251.43 |
| **Buffer weight** | **0.93** | **LTM half-life** | **232.86** | **57.22** | **247.26** |
| Buffer weight | 1.00 | Noise | 1.43 | 62.08 | 252.15 |
| Buffer duration | 0.80 | STM duration | 7.40 | 64.70 | 254.63 |
| **Buffer duration** | **0.69** | **LTM weight** | **0.009** | **58.52** | **248.60** |
| **Buffer duration** | **0.54** | **LTM half-life** | **185.94** | **57.53** | **247.59** |
| Buffer duration | 1.01 | Noise | 1.47 | 62.70 | 252.75 |
| STM duration | 13.07 | LTM weight | 0.005 | 61.67 | 251.75 |
| **STM duration** | **8.10** | **LTM half-life** | **217.05** | **58.62** | **248.71** |
| STM duration | 32.42 | Noise | 1.49 | 61.28 | 251.37 |
| LTM weight | 0.005 | LTM half-life | 544.37 | 61.63 | 251.72 |
| LTM weight | 0.005 | Noise | 1.31 | 61.72 | 251.80 |
| LTM half-life | 237.87 | Noise | 1.40 | 59.51 | 249.62 |

Values of parameters and their RMSE and AICc fit to RTs of the OLD group, when optimising pairs of parameters in turn to those of YOUNG. Multiple pairs of parameters produced plausible possibilities for differences between groups, with the best fitting parameters being buffer weight and LTM half-life. The best-fitting combinations (buffer weight and LTM half-life; buffer duration and LTM half-life; buffer duration and LTM weight; STM duration and LTM half-life) each contained one parameter controlling properties of buffer or short-term memory, and one controlling an aspect of long-term memory.

**Comparison of tone and pitch interval modelling**

The simulation of participants' responses reported in this paper modelled sequences using the absolute pitches of their tones. However, these sequences may also be represented by the size and direction of intervals between tones. While there are many situations where the intervalic content of an auditory sequence may be the more salient feature (Saffan et al., 2004) (*e.g.,* in more complex stimuli, such as music, where relative pitch may be an advantage in pattern recognition), the nature of the RANREG stimuli used in the present experiment results in similar outputs when using either representation of stimuli sequences.

In the RANREG stimuli used in this experiment, repetition of patterns in interval occur

concurrently with repetitions in absolute tone pitch. When a REG pattern is established, it repeats

both its tone and interval information exactly; RAN sections contain both random pitch and interval

content. These stimuli do not contain the common scenarios in which a distinction between absolute

pitch and interval provides different information, such as the transposition of a pattern's pitches but

maintaining interval sizes. As a result, simulated RTs using absolute tone values and intervals are highly

correlated and produce a similar fit to the task data. YOUNG: $r(268) = 0.86$, $p < .001$; OLD: $r(268) =$

$0.86$, $p < .001$.

913
914 **Figure S1. Comparison of simulated RTs using representations of absolute tone values and tone**
915 **intervals.** Modelling for YOUNG and OLD groups used the parameters obtained through optimisation
916 to observed RTs when using the tone representation (see Figure 3). Parameters of the OLD group
917 differ from those obtained for the YOUNG in *buffer weight* and *LTM half-life*.
918
919

921 **Figure S2. Observed and simulated RTs using representations of absolute tone values and tone**
922 **intervals.** Modelling for YOUNG and OLD groups used the parameters obtained through optimisation
923 to observed RTs when using the tone representation.
924

# References

Agus, T.R., Pressnitzer, D., 2021. Repetition detection and rapid auditory learning for
stochastic tone clouds. J. Acoust. Soc. Am. 150, 1735–1749. doi:10.1121/10.0005935
Agus, T.R., Thorpe, S.J., Pressnitzer, D., 2010. Rapid Formation of Robust Auditory
Memories: Insights from Noise. Neuron 66, 610–618. doi:10.1016/j.neuron.2010.04.014
Alvarez, P., Squire, L.R., 1994. Memory consolidation and the medial temporal lobe: {A}
simple network model. Proc. Natl. Acad. Sci. U. S. A. 91, 7041–7045.
doi:10.1073/pnas.91.15.7041
Anwyl-Irvine, A.L., Dalmaijer, E., Hodges, N., Evershed, J., 2020. Online Timing Accuracy
and Precision: A comparison of platforms, browsers, and participant's devices.
PsyArXiv doi:10.31234/osf.io/jfeca. doi:10.31234/osf.io/jfeca
Aslin, R.N., 2017. Statistical learning: a powerful mechanism that operates by mere
exposure. Wiley Interdiscip. Rev. Cogn. Sci. 8, 1–7. doi:10.1002/wcs.1373
Atienza, M., Cantero, J.L., Dominguez-Marin, E., 2002. Mismatch negativity ({MMN}): {An}
objective measure of sensory memory and long-lasting memories during sleep. Int. J.
Psychophysiol. 46, 215–225. doi:10.1016/S0167-8760(02)00113-7
Atkinson, R.C., Shiffrin, R.M., 1968. Human Memory: A Proposed System and its Control
Processes. Psychol. Learn. Motiv. - Adv. Res. Theory 2, 89–195. doi:10.1016/S0079-
7421(08)60422-3
Baird, A., Samson, S., 2009. Memory for music in Alzheimer's disease: Unforgettable?
Neuropsychol. Rev. 19, 85–101. doi:10.1007/s11065-009-9085-2
Baldeweg, T., 2006. Repetition effects to sounds: {Evidence} for predictive coding in the
auditory system [1]. Trends Cogn. Sci. 10, 93–94. doi:10.1016/j.tics.2006.01.010
Barascud, N., Pearce, M.T., Griffiths, T., Friston, K., Chait, M., 2016. Brain responses in
humans reveal ideal-observer-like sensitivity to complex acoustic patterns. Proc. Natl.
Acad. Sci. 113, E616-25. doi:10.1073/pnas.1508523113
Beigneux, K., Plaie, T., Isingrini, M., 2007. Aging effect on visual and spatial components of
working memory. Int. J. Aging Hum. Dev. 65, 301–314. doi:10.2190/AG.65.4.b
Benhamou, E., Warren, J.D., 2020. Disorders of music processing in dementia. Music Aging
Brain 107–149. doi:10.1016/b978-0-12-817422-7.00004-3
Benhamou, E., Zhao, S., Sivasathiaseelan, H., Johnson, J.C.S., Requena-Komuro, M.-C.,
Bond, R.L., van Leeuwen, J.E.P., Russell, L.L., Greaves, C. V, Nelson, A., Nicholas,
J.M., Hardy, C.J.D., Rohrer, J.D., Warren, J.D., 2021. Decoding expectation and
surprise in dementia: the paradigm of music. Brain Commun. 3.
doi:10.1093/braincomms/fcab173
Bianco, Chait, M., 2022. No link between Speech-in-noise perception and Auditory short-
term memory – evidence from a large cohort of older and younger listeners [WWW
Document]. PsyArXiv Prepr. doi:10.31234/osf.io/mjg4q
Bianco, Mills, G., de Kerangal, M., Rosen, S., Chait, M., 2021. Reward Enhances Online
Participants' Engagement With a Demanding Auditory Task. Trends Hear. 25.
doi:10.1177/23312165211025941
Bianco, R., Harrison, P.M.C.C., Hu, M., Bolger, C., Picken, S., Pearce, M.T., Chait, M.,
2020a. Long-term implicit memory for sequential auditory patterns in humans. Elife
9:e56073. doi:10.1101/2020.02.14.949404
Bianco, R., Ptasczynski, L.E., Omigie, D., 2020b. Pupil responses to pitch deviants reflect
predictability of melodic sequences. Brain Cogn. 138, 103621.
doi:10.1016/j.bandc.2019.103621

Blake, R. V., Wroe, S.J., Breen, E.K., McCarthy, R.A., 2000. Accelerated forgetting in
patients with epilepsy. Evidence for an impairment in memory consolidation. Brain 123,
472–483. doi:10.1093/brain/123.3.472
Bonetti, L., Brattico, E., Bruzzone, S.E.P., Donati, G., Deco, G., Pantazis, D., Vuust, P.,
Kringelbach, M.L., 2022. Brain recognition of previously learned versus novel temporal
sequences : a differential simultaneous. Cereb. Cortex 33, 1–14.
Boughorbel, S., Jarray, F., El-Anbari, M., 2017. Optimal classifier for imbalanced data using
Matthews Correlation Coefficient metric. PLoS One 12, 1–17.
doi:10.1371/journal.pone.0177678
Bridges, D., Pitiot, A., MacAskill, M.R., Peirce, J.W., 2020. The timing mega-study:
Comparing a range of experiment generators, both lab-based and online. PeerJ 8.
doi:10.7717/peerj.9414
Brown, S., Martinez, M.J., Parsons, L.M., 2004. Passive music listening spontaneously
engages limbic and paralimbic systems. Neuroreport 15, 2033–2037.
doi:10.1097/00001756-200409150-00008
Bunton, S., 1996. On-line stochastic processes in data compression. University of
Washington, Seattle, WA.
Burunat, I., Alluri, V., Toiviainen, P., Numminen, J., Brattico, E., 2014. Dynamics of brain
activity underlying working memory for music in a naturalistic condition. Cortex 57, 254–
269. doi:10.1016/j.cortex.2014.04.012
Cansino, S., 2009. Episodic memory decay along the adult lifespan: {A} review of behavioral
and neurophysiological evidence. Int. J. Psychophysiol. 71, 64–69.
doi:10.1016/j.ijpsycho.2008.07.005
Cazala, A., Giret, N., Edeline, J.-M., Del Negro, C., 2019. Neuronal Encoding in a High-Level
Auditory Area: From Sequential Order of Elements to Grammatical Structure. J.
Neurosci. 39, 6150–6161. doi:10.1523/jneurosci.2767-18.2019
Cheng, C.H., Hsu, W.Y., Lin, Y.Y., 2013. Effects of physiological aging on mismatch
negativity: A meta-analysis. Int. J. Psychophysiol. 90, 165–171.
doi:10.1016/j.ijpsycho.2013.06.026
Christiansen, M.H., 2019. Implicit Statistical Learning: A Tale of Two Literatures. Top. Cogn.
Sci. 11, 468–481. doi:10.1111/tops.12332
Cleary, J., Witten, I., 1984. Data compression using adaptive coding and partial string
matching. EEE Trans. Commun. 32, 396–402. doi:https://doi.org/10.1109/TCOM.1984.
1096090 18.
Conway, C.M., 2020. How does the brain learn environmental structure? Ten core principles
for understanding the neurocognitive mechanisms of statistical learning. Neurosci.
Biobehav. Rev. 112, 279–299. doi:10.1016/j.neubiorev.2020.01.032
Cooper, R.J., Todd, J., McGill, K., Michie, P.T., 2006. Auditory sensory memory and the
aging brain: {A} mismatch negativity study. Neurobiol. Aging 27, 752–762.
doi:10.1016/j.neurobiolaging.2005.03.012
Corsi, P.M., 1972. Human memory and the medial temporal region of the brain. Diss. Abstr.
Int. Sect. B. Sci. Eng. 34, 891.
Cowan, N., 1984. On short and long auditory stores. Psychol. Bull. 96, 341–370.
doi:10.1037/0033-2909.96.2.341
Dauer, T., Henry, M.J., Herrmann, B., 2022. Auditory perceptual learning depends on
temporal regularity and certainty. J. Exp. Psychol. Hum. Percept. Perform. 48, 755–
770. doi:10.1037/xhp0001016
Davis, H.P., Small, S.A., Stern, Y., Mayeux, R., Feldstein, S.N., Keller, F.R., 2003.
Acquisition, recall, and forgetting of verbal information in long-term memory by young,
middle-aged, and elderly individuals. Cortex 39, 1063–1091. doi:10.1016/S0010-
9452(08)70878-5
Davis, R.L., Zhong, Y., 2017. The Biology of Forgetting—A Perspective. Neuron 95, 490–
503. doi:10.1016/j.neuron.2017.05.039
de Kerangal, M., Vickers, D., Chait, M., 2021. The effect of healthy aging on change
detection and sensitivity to predictable structure in crowded acoustic scenes. Hear.

Res. 399, 108074. doi:10.1016/j.heares.2020.108074
Der, G., Deary, I.J., 2006. Age and Sex Differences in Reaction Time in Adulthood: Results
From the United Kingdom Health and Lifestyle Survey. Psychol. Aging 21, 62–73.
doi:10.1037/0882-7974.21.1.62.supp
Di Liberto, G.M., Pelofi, C., Bianco, R., Patel, P., Mehta, A.D., Herrero, J.L., de Cheveigné,
A., Shamma, S., Mesgarani, N., 2020. Cortical encoding of melodic expectations in
human temporal cortex. Elife 9:e51784.
Elliott, G., Isaac, C.L., Muhlert, N., 2014. Measuring forgetting: A critical review of
accelerated long-term forgetting studies. Cortex 54, 16–32.
doi:10.1016/j.cortex.2014.02.001
Fleischman, D.A., 2007. Repetition priming in aging and Alzheimer's disease: An integrative
review and future directions. Cortex 43, 889–897. doi:10.1016/S0010-9452(08)70688-9
Fogerty, D., Humes, L.E., Busey, T.A., 2016. Age-related declines in early sensory memory:
Identification of rapid auditory and visual stimulus sequences. Front. Aging Neurosci. 8,
1–16. doi:10.3389/fnagi.2016.00090
Fournet, N., Roulin, J.-L., Vallet, F., Beaudoin, M., Agrigoroaei, S., Paignon, A., Dantzer, C.,
Desrichard, O., 2012. Evaluating short-term and working memory in older adults:
{French} normative data. Aging Ment. Health 16, 922–930.
doi:10.1080/13607863.2012.674487
Füllgrabe, C., 2020. On the Possible Overestimation of Cognitive Decline: The Impact of
Age-Related Hearing Loss on Cognitive-Test Performance. Front. Neurosci. 14.
doi:10.3389/fnins.2020.00454
Gershman, S.J., 2022. The molecular memory code and synaptic plasticity: a synthesis 1–
51.
Griffiths, T.D., Lad, M., Kumar, S., Holmes, E., McMurray, B., Maguire, E.A., Billig, A.J.,
Sedley, W., 2020. How Can Hearing Loss Cause Dementia? Neuron 1–12.
doi:10.1016/j.neuron.2020.08.003
Haenschel, C., Vernon, D.J., Dwivedi, P., Gruzelier, J.H., Baldeweg, T., Haeaschel, C.,
Vernon, D.J., Dwivedi, P., Gruzelier, J.H., Baldeweg, T., 2005. Event-related brain
potential correlates of human auditory sensory memory-trace formation. J. Neurosci.
25, 10494–10501. doi:10.1523/JNEUROSCI.1227-05.2005
Hardt, O., Nader, K., Nadel, L., 2013. Decay happens: The role of active forgetting in
memory. Trends Cogn. Sci. 17, 111–120. doi:10.1016/j.tics.2013.01.001
Harrison, P.M.C.C., Bianco, R., Chait, M., Pearce, M.T., 2020. PPM-Decay: A computational
model of auditory prediction with memory decay, PLoS Computational Biology.
doi:10.1371/journal.pcbi.1008304
Hauser, M.D., Newport, E.L., Aslin, R.N., 2001. Segmentation of the speech stream in a
non-human primate: Statistical learning in cotton-top tamarins. Cognition 78, B53–B64.
doi:10.1016/S0010-0277(00)00132-3
Hebb, D.O., 1961. Brain mechanisms and learning., in: Delafresnaye, J.F. (Ed.), Distinctive
Features of Learning in the Higher Animal. New York: Oxford University Press., pp. 37–
46.
Heilbron, M., Chait, M., 2018. Great Expectations: Is there Evidence for Predictive Coding in
Auditory Cortex? Neuroscience 389, 54–73. doi:10.1016/j.neuroscience.2017.07.061
Herff, S.A., Zhen, S., Yu, R., Agres, K.R., 2020. Age-dependent statistical learning
trajectories reveal differences in information weighting. Psychol. Aging 35, 1090–1104.
doi:10.1037/pag0000567
Herrmann, B., Araz, K., Johnsrude, I.S., 2021. Sustained neural activity correlates with rapid
perceptual learning of auditory patterns. Neuroimage 238, 118238.
doi:10.1016/j.neuroimage.2021.118238
Herrmann, B., Maess, B., Johnsrude, I.S., 2022. A neural signature of regularity in sound is
reduced in older adults. Neurobiol. Aging 109, 1–10.
doi:10.1016/j.neurobiolaging.2021.09.011
Hoefeijzers, S., Dewar, M., Della Sala, S., Zeman, A., Butler, C., 2013. Accelerated long-
term forgetting in transient epileptic amnesia: {An} acquisition or consolidation deficit?

Neuropsychologia 51, 1549–1555. doi:10.1016/j.neuropsychologia.2013.04.017

Huang, C.-Z.A., Vaswani, A., Uszkoreit, J., Shazeer, N., Simon, I., Hawthorne, C., Dai, A.M., Hoffman, M.D., Dinculescu, M., Eck, D., 2018. Music Transformer.

Hultsch, D.F., MacDonald, S.W.S.S., Dixon, R.A., 2002. Variability in reaction time performance of younger and older adults. Journals Gerontol. - Ser. B Psychol. Sci. Soc. Sci. 57, 101–115. doi:10.1093/geronb/57.2.P101

Humes, L.E., Busey, T.A., Craig, J., Kewley-Port, D., 2013. Are age-related changes in cognitive function driven by age-related changes in sensory processing? Attention, Perception, Psychophys. 75, 508–524. doi:10.3758/s13414-012-0406-9

Jääskeläinen, I.P., Varonen, R., Näätänen, R., Pekkonen, E., 1999. Decay of cortical preattentive sound discrimination in middle-age. Neuroreport 10, 123–126.

Jacobsen, J.H., Stelzer, J., Fritz, T.H., Chételat, G., La Joie, R., Turner, R., 2015. Why musical memory can be preserved in advanced Alzheimer's disease. Brain 138, 2438–2450. doi:10.1093/brain/awv135

Janacsek, K., Fiser, J., Nemeth, D., 2012. The best time to acquire new skills: Age-related differences in implicit sequence learning across the human lifespan. Dev. Sci. 15, 496–505. doi:10.1111/j.1467-7687.2012.01150.x

Johnson, J.C.S.S., Marshall, C.R., Weil, R.S., Bamiou, D.E., Hardy, C.J.D.D., Warren, J.D., 2021. Hearing and dementia: from ears to brain. Brain 144, 391–401. doi:10.1093/brain/awaa429

Johnson, S.G., 2020. The NLopt nonlinear-optimization package. [WWW Document]. http://github.com/stevengj/nlopt.

Kang, H., Agus, T.R., Pressnitzer, D., 2017. Auditory memory for random time patterns. J. Acoust. Soc. Am. 142, 2219–2232. doi:10.1121/1.5007730

Kern, P., Heilbron, M., Lange, F.P. De, Spaak, E., 2022. Cortical activity during naturalistic music listening reflects short- range predictions based on long- term experience. Elife 11:e80935.

Kessels, R.P.C., Van Zandvoort, M.J.E., Postma, A., Kappelle, L.J., De Haan, E.H.F., 2000. The Corsi Block-Tapping Task: Standardization and normative data. Appl. Neuropsychol. 7, 252–258. doi:10.1207/S15324826AN0704_8

Kiang, M., Braff, D.L., Sprock, J., Light, G.A., 2009. The relationship between preattentive sensory processing deficits and age in schizophrenia patients. Clin. Neurophysiol. 120, 1949–1957. doi:10.1016/j.clinph.2009.08.019

Kobor, A., Janacsek, K., Takacs, A., Nemeth, D., 2017. Statistical learning leads to persistent memory: Evidence for one-year consolidation. Sci. Rep. 7, 1–10. doi:10.1038/s41598-017-00807-3

Koen, J.D., Yonelinas, A.P., 2016. Recollection, not familiarity, decreases in healthy ageing: Converging evidence from four estimation methods. Memory 24, 75–88. doi:10.1080/09658211.2014.985590

Koen, J.D., Yonelinas, A.P., 2014. The Effects of Healthy Aging, Amnestic Mild Cognitive Impairment, and Alzheimer's Disease on Recollection and Familiarity: A Meta-Analytic Review. Neuropsychol. Rev. 24, 332–354. doi:10.1007/s11065-014-9266-5

Kumar, S., Bonnici, H.M., Teki, S., Agus, T.R., Pressnitzer, D., Maguire, E.A., Griffiths, T.D., 2014. Representations of specific acoustic patterns in the auditory cortex and hippocampus. Proc. R. Soc. B Biol. Sci. 281, 20141000. doi:10.1098/rspb.2014.1000

Leek, M.R., Watson, C.S., 1988. Auditory perceptual learning of tonal patterns. Percept. Psychophys. 43, 389–394. doi:10.3758/BF03208810

Lewandowsky, S., Oberauer, K., Brown, G.D.A.A., 2009. No temporal decay in verbal short-term memory. Trends Cogn. Sci. 13, 120–126. doi:10.1016/j.tics.2008.12.003

Lu, K., Vicario, D.S., 2014. Statistical learning of recurring sound patterns encodes auditory objects in songbird forebrain. Proc. Natl. Acad. Sci. 111, 14553–14558. doi:10.1073/pnas.1412109111

Lukács, Á., Kemény, F., 2015. Development of different forms of skill learning throughout the lifespan. Cogn. Sci. 39, 383–404. doi:10.1111/cogs.12143

Mander, B.A., Rao, V., Lu, B., Saletin, J.M., Lindquist, J.R., Ancoli-Israel, S., Jagust, W.,

1138   Walker, M.P., 2013. Prefrontal atrophy, disrupted NREM slow waves and impaired
1139   hippocampal-dependent memory in aging. Nat. Neurosci. 16, 357–364.
1140   doi:10.1038/nn.3324
1141 Manes, F., Serrano, C., Calcagno, M.L., Cardozo, J., Hodges, J., 2008. Accelerated
1142   forgetting in subjects with memory complaints A new form of Mild Cognitive
1143   Impairment? J Neurol 255, 1067–1070. doi:10.1007/s00415-008-0850-6
1144 Manly, T., Davison, B., Heutink, J., Galloway, M., Robertson, I.H., 2000. Not enough time or
1145   not enough attention ? Speed , error and self-maintained control in the Sustained
1146   Attention to Response Test ( SART ). Clin. Neuropsychol. Assess. 3, 167–177.
1147 Mary, A., Schreiner, S., Peigneux, P., 2013. Accelerated long-term forgetting in aging and
1148   intra-sleep awakenings. Front. Psychol. 4, 1–11. doi:10.3389/fpsyg.2013.00750
1149 Massaro, D.W., Cohen, M.M., 1975. Preperceptual Auditory Storage in Speech Recognition.
1150   Springer, Berlin, Heidelberg, pp. 226–245. doi:10.1007/978-3-642-81000-8_14
1151 McDermott, J.H., Wrobleski, D., Oxenham, A.J., 2011. Recovering sound sources from
1152   embedded repetition. Proc. Natl. Acad. Sci. 108, 1188–1193.
1153   doi:10.1073/pnas.1004765108
1154 McGibbon, T., Jansari, A., Demirjian, J., Nemes, A., Opre, A., 2022. Accelerated forgetting
1155   in healthy older samples: {Implications} for methodology, future ageing studies, and
1156   early identification of risk of dementia. Q. J. Exp. Psychol.
1157   doi:10.1177/17470218221113412
1158 Michael Lawrence, M.A., 2016. Package "ez": Easy Analysis and Visualization of Factorial
1159   Experiments 4–5. doi:10.1515/zava.93.1.1
1160 Milne, A., Bianco, R., Poole, K., Zhao, S., Oxenham, A., Billig, A., Chait, M., 2020. An online
1161   headphone screening test based on dichotic pitch. Behav. Res. Methods.
1162   doi:10.1101/2020.07.21.214395
1163 Murphy, D.R., Craik, F.I.M.M., Li, K.Z.H.H., Schneider, B.A., 2000. Comparing the effects of
1164   aging and background noise on short-term memory performance. Psychol. Aging 15,
1165   323–334. doi:10.1037/0882-7974.15.2.323
1166 Näätänen, R., Kujala, T., Escera, C., Baldeweg, T., Kreegipuu, K., Carlson, S., Ponton, C.,
1167   2012. The mismatch negativity (MMN) - A unique window to disturbed central auditory
1168   processing in ageing and different clinical conditions. Clin. Neurophysiol. 123, 424–458.
1169   doi:10.1016/j.clinph.2011.09.020
1170 Näätänen, R., Paavilainen, P., Rinne, T., Alho, K., 2007. The mismatch negativity (MMN) in
1171   basic research of central auditory processing: A review. Clin. Neurophysiol. 118, 2544–
1172   2590. doi:10.1016/j.clinph.2007.04.026
1173 Narmour, E., 1990. The analysis and cognition of basic melodic structures: The implication-
1174   realization model., University. ed.
1175 Nees, M.A., 2016. Have we forgotten auditory sensory memory? Retention intervals in
1176   studies of nonverbal auditory working memory. Front. Psychol. 7, 1–6.
1177   doi:10.3389/fpsyg.2016.01892
1178 Nemeth, D., Janacsek, K., 2011. The dynamics of implicit skill consolidation in young and
1179   elderly adults. Journals Gerontol. - Ser. B Psychol. Sci. Soc. Sci. 66 B, 15–22.
1180   doi:10.1093/geronb/gbq063
1181 Nyberg, L., Lövdén, M., Riklund, K., Lindenberger, U., Bäckman, L., 2012. Memory aging
1182   and brain maintenance. Trends Cogn. Sci. 16, 292–305. doi:10.1016/j.tics.2012.04.005
1183 Nyberg, L., Pudas, S., 2019. Successful Memory Aging. Annu. Rev. Psychol. 70, 219–243.
1184   doi:10.1146/annurev-psych-010418-103052
1185 Ohno, T., Hasegawa, T., Tsuruoka, T., Terabe, K., Gimzewski, J.K., Aono, M., 2011. Short-
1186   term plasticity and long-term potentiation mimicked in single inorganic synapses. Nat.
1187   Mater. 10, 591–595. doi:10.1038/nmat3054
1188 Omigie, D., Pearce, M., Lehongre, K., Hasboun, D., Navarro, V., Adam, C., Samson, S.,
1189   2019. Intracranial Recordings and Computational Modeling of Music Reveal the Time
1190   Course of Prediction Error Signaling in Frontal and Temporal Cortices. J. Cogn.
1191   Neurosci. 31, 855–873. doi:10.1162/jocn_a_01388
1192 Palmer, S.D., Hutson, J., Mattys, S.L., 2018. Statistical learning for speech segmentation:

Age-related changes and underlying mechanisms. Psychol. Aging 33, 1035–1044.
doi:10.1037/pag0000292

Pearce, M.T.M.T., Ruiz, M.H., Kapasi, S., Wiggins, G. a, Bhattacharya, J., 2010.
Unsupervised statistical learning underpins computational, behavioural, and neural
manifestations of musical expectation. Neuroimage 50, 302–13.
doi:10.1016/j.neuroimage.2009.12.019

Pekkonen, E., Rinne, T., Reinikainen, K., Kujala, T., Alho, K., Näätänen, R., 1996. Aging
Effects on Auditory Processing: An Event-Related Potential Study. Exp. Aging Res. 22,
171–184. doi:10.1080/03610739608254005

Pekkonen, Hirvonen, J., Jääskeläinen, I.P., Kaakkola, S., Huttunen, J., 2001. Auditory
sensory memory and the cholinergic system: Implications for Alzheimer's disease.
Neuroimage 14, 376–382. doi:10.1006/nimg.2001.0805

Petkov, C.I., ten Cate, C., 2020. Structured Sequence Learning: Animal Abilities, Cognitive
Operations, and Language Evolution. Top. Cogn. Sci. 12, 828–842.
doi:10.1111/tops.12444

Poeppel, D., Idsardi, W., 2022. We don't know how the brain stores anything, let alone
words. Trends Cogn. Sci. 26, 1054–1055. doi:10.1016/j.tics.2022.08.010

Powers, D.M.W., 2007. Evaluation: From Precision, Recall and F-Factor to ROC,
Informedness, Markedness & Correlation. Tech. Rep. SIE-07-001.

Quiroga-Martinez, D.R., Hansen, N.C., Højlund, A., Pearce, M., Brattico, E., Vuust, P., 2020.
Decomposing neural responses to melodic surprise in musicians and non-musicians:
Evidence for a hierarchy of predictions in the auditory system. Neuroimage 215.
doi:10.1016/j.neuroimage.2020.116816

Raz, N., Lindenberger, U., 2011. Only Time will Tell: Cross-sectional Studies Offer no
Solution to the Age-Brain-Cognition Triangle—Comment on Salthouse (2011). Psychol
Bull 137, 790–795. doi:doi:10.1037/a0024503

Reber, A.S., 1989. Implicit Learning and Tacit Knowledge. J. Exp. Psychol. Gen. 118, 219–
235. doi:10.1037/0096-3445.118.3.219

Rieckmann, A., Bäckman, L., 2009. Implicit learning in aging: Extant patterns and new
directions. Neuropsychol. Rev. 19, 490–503. doi:10.1007/s11065-009-9117-y

Rimmele, J., Sussman, E., Keitel, C., Jacobsen, T., Schröger, E., 2012. Electrophysiological
evidence for age effects on sensory memory processing of tonal patterns. Psychol
Aging 27, 384–398. doi:10.1037/a0024866

Rimmele, J.M., Sussman, E., Poeppel, D., 2015. The role of temporal structure in the
investigation of sensory memory, auditory scene analysis, and speech perception: A
healthy-aging perspective. Int. J. Psychophysiol. 95, 175–183.
doi:10.1016/j.ijpsycho.2014.06.010

Ringer, H., Schröger, E., Grimm, S., 2022. Perceptual Learning and Recognition of Random
Acoustic Patterns. Audit. Percept. Cogn. 00, 1–23.
doi:10.1080/25742442.2022.2082827

Román-Caballero, R., Arnedo, M., Triviño, M., Lupiáñez, J., 2018. Musical practice as an
enhancer of cognitive function in healthy aging - A systematic review and meta-
analysis. PLoS One 13, 1–23. doi:10.1371/journal.pone.0207957

Rosen, S., 1992. Temporal information in speech: acoustic, auditory and linguistic aspects.
Philos. Trans. R. Soc. Lond. B. Biol. Sci. doi:10.1098/rstb.1992.0070

Ross, G.J., Tasoulis, D.K., Adams, N.M., 2011. Nonparametric monitoring of data streams
for changes in location and scale. Technometrics 53, 379–389.
doi:10.1198/TECH.2011.10069

Rowan, T., 1990. Functional stability analysis of numerical algorithms.

Ryan, T.J., Frankland, P.W., 2022. Forgetting as a form of adaptive engram cell plasticity.
Nat. Rev. Neurosci. 1–14. doi:10.1038/s41583-021-00548-3

Saffran, J.R., Aslin, R.N., Newport, E.L., 1996. Statistical Learning by 8-Month-Old Infants.
Science (80-. ). 274, 1926–1928.

Salthouse, T.A., 2011. Neuroanatomical substrates of age-related cognitive decline.
Psychol. Bull. 137, 753–784. doi:10.1037/a0023262

Salthouse, T.A., 1996. The processing-speed theory of adult age differences in cognition.
Psychol. Rev. 103, 403–428.
Sanchez Lopez, R., Bianchi, F., Fereczkowski, M., Santurette, S., Dau, T., 2018. Data-
Driven Approach for Auditory Profiling and Characterization of Individual Hearing Loss.
Trends Hear. 22, 1–12. doi:10.1177/2331216518807400
Santolin, C., Saffran, J.R., 2018. Constraints on Statistical Learning Across Species. Trends
Cogn. Sci. 22, 52–63. doi:10.1016/j.tics.2017.10.003
Schapiro, A.C., Gregory, E., Landau, B., McCloskey, M., Turk-Browne, N.B., 2014. The
necessity of the medial temporal lobe for statistical learning. J. Cogn. Neurosci. 26,
1736–1747. doi:10.1162/jocn
Schevenels, K., Altvater-Mackensen, N., Zink, I., De Smedt, B., Vandermosten, M., 2021.
Aging effects and feasibility of statistical learning tasks across modalities. Aging,
Neuropsychol. Cogn. 00, 1–30. doi:10.1080/13825585.2021.2007213
Schmithorst, V.J., 2005. Separate cortical networks involved in music perception:
Preliminary functional MRI evidence for modularity of music processing. Neuroimage
25, 444–451. doi:10.1016/j.neuroimage.2004.12.006
Schneider, B.A., Daneman, M., Pichora-Fuller, M.K., 2002. Listening in aging adults: From
discourse comprehension to psychoacoustics. Can. J. Exp. Psychol. 56, 139–152.
doi:10.1037/h0087392
Schneider, B.A., Pichora-Fuller, M.K., 2000. Implications of perceptual deterioration for
cognitive aging research, in: The Handbook of Aging and Cognition. pp. 155–219.
Skerritt-Davis, B., Elhilali, M., 2021. Computational framework for investigating predictive
processing in auditory perception. J. Neurosci. Methods 360, 109177.
doi:10.1016/j.jneumeth.2021.109177
Sluming, V., Barrick, T., Howard, M., Cezayirli, E., Mayes, A., Roberts, N., 2002. Voxel-
Based Morphometry Reveals Increased Gray Matter Density in Broca's Area in Male
Symphony Orchestra Musicians. Neuroimage 17, 1613–1622.
doi:10.1006/nimg.2002.1288
Smalle, E.H.M.M., Page, M.P.A.A., Duyck, W., Edwards, M., Szmalec, A., 2018. Children
retain implicitly learned phonological sequences better than adults: a longitudinal study.
Dev. Sci. 21. doi:10.1111/desc.12634
Sofi, F., Valecchi, D., Bacci, D., Abbate, R., Gensini, G.F., Casini, A., Macchi, C., 2011.
Physical activity and risk of cognitive decline: {A} meta-analysis of prospective studies.
J. Intern. Med. 269, 107–117. doi:10.1111/j.1365-2796.2010.02281.x
Soyman, E., Vicario, D.S., 2019. Rapid and long-lasting improvements in neural
discrimination of acoustic signals with passive familiarization, PLoS ONE.
doi:10.1371/journal.pone.0221819
Squires, K.C., Wickens, C., Squires, N.K., Donchin, E., 1976. The effect of stimulus
sequence on the waveform of the cortical event-related potential. Science (80-. ). 193,
1142–1146. doi:10.1126/science.959831
Swords, G.M., Nguyen, L.T., Mudar, R.A., Llano, D.A., 2018. Auditory system dysfunction in
Alzheimer disease and its prodromal states: A review. Ageing Res. Rev. 44, 49–59.
doi:10.1016/j.arr.2018.04.001
Tóth-Fábera, E., Nemeth, D., Janacsek, K., 2023. Lifespan developmental invariance in
memory consolidation: evidence from procedural memory. PNAS Nexus pgad037.
Wais, P.E., Gazzaley, A., 2014. Distractibility during retrieval of long-term memory:
{Domain}-general interference, neural networks and increased susceptibility in normal
aging. Front. Psychol. 5, 1–12. doi:10.3389/fpsyg.2014.00280
Warren, B.M., Ackroff, J.M., 1976. Two types of auditory sequence perception. Percept.
Psychophys. 20, 387–394. doi:10.3758/BF03199420
Watanabe, T., Yagishita, S., Kikyo, H., 2008. Memory of music: Roles of right hippocampus
and left inferior frontal gyrus. Neuroimage 39, 483–491.
doi:10.1016/j.neuroimage.2007.08.024
Wayne, R. V., Johnsrude, I.S., 2015. A review of causal mechanisms underlying the link
between age-related hearing loss and cognitive decline. Ageing Res. Rev. 23, 154–

166. doi:10.1016/j.arr.2015.06.002
Wearn, A.R., Saunders-Jennings, E., Nurdal, V., Hadley, E., Knight, M.J., Newson, M.,
Kauppinen, R.A., Coulthard, E.J., 2020. Accelerated long-term forgetting in healthy
older adults predicts cognitive decline over 1 year. Alzheimer's Res. Ther. 12, 1–9.
doi:10.1186/s13195-020-00693-4
Weston, P.S.J.J., Nicholas, J.M., Henley, S.M.D.D., Liang, Y., Macpherson, K., Donnachie,
E., Schott, J.M., Rossor, M.N., Crutch, S.J., Butler, C.R., Zeman, A.Z., Fox, N.C., 2018.
Accelerated long-term forgetting in presymptomatic autosomal dominant Alzheimer's
disease: a cross-sectional study. Lancet Neurol. 17, 123. doi:10.1016/S1474-
4422(17)30434-9
Wilson, B., Slater, H., Kikuchi, Y., Milne, A.E., Marslen-Wilson, W.D., Smith, K., Petkov, C.I.,
2013. Auditory artificial grammar learning in Macaque and Marmoset monkeys. J.
Neurosci. 33, 18825–18835. doi:10.1523/JNEUROSCI.2414-13.2013
Winkler, I., Cowan, N., 2005. From sensory to long-term memory: Evidence from auditory
memory reactivation studies. Exp. Psychol. 52, 3–20. doi:10.1027/1618-3169.52.1.3
Winkler, I., Denham, S.L., Nelken, I., 2009. Modeling the auditory scene: predictive regularity
representations and perceptual objects. Trends Cogn. Sci. 13, 532–540.
doi:10.1016/j.tics.2009.09.003
Woods, K.J.P., Mcdermott, J.H., 2018. Schema learning for the cocktail party problem.
PNAS 115, E3313–E3322. doi:10.1073/pnas.1801614115
Zajac, I.T., Nettelbeck, T., 2018. Auditory speed tasks as potential candidates for the study
of cognitive ageing. Aging, Neuropsychol. Cogn. 25, 167–185.
doi:10.1080/13825585.2016.1272671