Cellular iron governs the host response to malaria

Sarah K. Wideman[1], Joe N. Frost[1,6], Felix C. Richter[2,6], Caitlin Naylor[1], José M. Lopes[3,4], Nicole Viveiros[4], Megan R. Teh[1], Alexandra E. Preston[1], Natasha White[1], Shamsideen Yusuf[1], Simon J. Draper[5], Andrew E. Armitage[1], Tiago L. Duarte[3,4], Hal Drakesmith[1,7].

Affiliations

[1] MRC Human Immunology Unit, MRC Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford, United Kingdom.

[2] Kennedy Institute of Rheumatology, Roosevelt Drive, OX3 7FY, Oxford, United Kingdom.

[3] Faculty of Medicine (FMUP) and Institute of Molecular Pathology, Immunology (IPATIMUP), University of Porto, Porto, Portugal.

[4] Instituto de Biologia Molecular e Celular & Instituto de Investigação e Inovação em Saúde (i3S), University of Porto, Porto, Portugal.

[5] Department of Biochemistry, University of Oxford, South Parks Road, Oxford, OX1 3QU, UK.

[6] Authors contributed equally.

[7] Corresponding author: alexander.drakesmith@ndm.ox.ac.uk.

ABSTRACT

Malaria and iron deficiency are major global health problems with extensive epidemiological overlap.

Iron deficiency-induced anaemia can protect the host from malaria by limiting parasite growth. On the

other hand, iron deficiency can significantly disrupt immune cell function. However, the impact of host

cell iron scarcity beyond anaemia remains elusive in malaria. To address this, we employed a transgenic

mouse model carrying a mutation in the transferrin receptor ($Tfrc^{Y20H/Y20H}$), which limits the ability of

cells to internalise iron from plasma. At homeostasis $Tfrc^{Y20H/Y20H}$ mice appear healthy and are not

anaemic. However, $Tfrc^{Y20H/Y20H}$ mice infected with *Plasmodium chabaudi chabaudi AS* showed

significantly higher peak parasitaemia and body weight loss. We found that $Tfrc^{Y20H/Y20H}$ mice displayed

a similar trajectory of malaria-induced anaemia as wild-type mice, and elevated circulating iron did not

increase peak parasitaemia. Instead, *P. chabaudi* infected $Tfrc^{Y20H/Y20H}$ mice had an impaired innate and

adaptive immune response, marked by decreased cell proliferation and cytokine production.

Moreover, we demonstrated that these immune cell impairments were cell-intrinsic, as *ex vivo* iron

supplementation fully recovered CD4 T cell and B cell function. Despite the inhibited immune response

and increased parasitaemia, $Tfrc^{Y20H/Y20H}$ mice displayed mitigated liver damage, characterised by

decreased parasite sequestration in the liver and an attenuated hepatic immune response. Together, these

results show that host cell iron scarcity inhibits the immune response but prevents excessive hepatic

tissue damage during malaria infection. These divergent effects shed light on the role of iron in the

complex balance between protection and pathology in malaria.

INTRODUCTION

Malaria is a major global health problem that causes significant morbidity and mortality worldwide (1). It is caused by *Plasmodium* species parasites, which have a complex life cycle and are transmitted between humans by *Anopheles* mosquitos. In the human host, multiple cycles of asexual parasite replication inside red blood cells (RBC) result in extensive RBC destruction, immune activation, and microvascular obstruction (2). This blood stage of infection gives rise to symptoms such as fever, chills, headache, and malaise. In severe cases, it can also cause life-threatening complications such as acute anaemia, coma, respiratory distress, and organ failure (2).

There is a complex relationship between host iron status and malaria. Iron is an essential micronutrient that is required by most living organisms to maintain physiological and biochemical processes, such as oxygen transport and storage, cellular metabolism, and reduction-oxidation reactions (3,4). Despite the importance of iron, iron deficiency is exceedingly common in humans, and iron deficiency anaemia is estimated to affect a sixth of the world's population (5,6). In the context of human malaria infection, iron deficiency can decrease the risk of disease, severe disease, and mortality (7–9). The protective effect of iron deficiency is at least partly mediated by anaemia, as RBCs isolated from anaemic individuals are less amenable to malaria parasite growth (10).

Meanwhile, oral iron supplementation is a risk factor for malaria in areas with limited access to preventative measures and treatment (11,12). This effect can to some extent be explained by iron supplementation stimulating erythropoiesis and increasing the proportion of reticulocytes and young erythrocytes, which are preferred targets for invasion by *P. falciparum* parasites (10). Malaria and iron deficiency also often disproportionally affect the same populations (e.g. young children in the WHO African Region) (1,6), in part, because malaria causes iron deficiency (13).

Anaemia is the primary and most profound consequence of iron deficiency. However, iron deficiency can also have other negative impacts on human health. Immune cells with high proliferative and anabolic capacities appear to be particularly sensitive to iron deficiency. As such, decreased iron availability can impair the proliferation and maturation of lymphocytes and neutrophils (14–16).

Neutrophils and macrophages also require iron for enzymes involved in microbial killing (16–19). In

animal models of iron deficiency, lymphocyte function is severely impaired, and the immune response

to immunisation and viral infection is inhibited (20,21). Similarly, iron deficiency decreases

inflammation and improves outcomes in mouse models of autoimmune disease (22–25). In humans,

associations between iron deficiency and attenuated responses to some vaccines have been observed

(20,21,26–28). Moreover, patients with a rare mutation in transferrin receptor-1 (TfR1), the primary

receptor for iron uptake in cells, present with lymphocyte dysfunction and combined immunodeficiency

(29,30).

Controlling a malaria infection requires two distinct but complementary immune responses. An early

cell-mediated response, primarily driven by interferon-γ (IFN-γ) producing CD4$^+$ T cells, prevents

uncontrolled exponential parasite growth (31–35). Meanwhile, a humoral response is required to

prevent recrudescence and to clear the infection (36,37). Excessive production of pro-inflammatory

immune cells and cytokines can lead to sepsis-like complications and cause collateral damage to tissues

and organs (38,39). Thus, the pro-inflammatory anti-parasite response must be balanced by

immunoregulatory and tissue-protective responses to prevent immunopathology (40–43).

Although it is known that host iron deficiency influences malaria infection, the mechanisms that affect

host health or *Plasmodium* virulence remain largely unknown. In particular, the effects of iron

deficiency aside from anaemia, have scarcely been explored. Moreover, any effects on malaria

immunity have not been investigated beyond a few observational studies that found associations

between iron deficiency and attenuated antibody responses to malaria in children (7,44,45).

In this study, we aspired to deepen our understanding of how malaria infection is affected by host iron

deficiency. To this end, we employed a genetic mouse model of cellular iron deficiency based on a rare

mutation in TfR1 (*Tfrc$^{Y20H/Y20H}$*), which causes combined immunodeficiency in humans (29,30). We

found that decreasing host cellular iron levels increased peak malaria parasitaemia in mice infected with

P. chabaudi. While *P. chabaudi*-induced anaemia and RBC invasion remained unaffected, the immune

response to *P. chabaudi* was drastically inhibited. Interestingly, mice with cellular iron deficiency also

had attenuated *P. chabaudi*-induced liver damage, suggesting reduced immunopathology. Hence, host

cellular iron deficiency attenuated the immune response to malaria, leading to increased pathogen

burden and mitigated liver pathology.

RESULTS

Decreased cellular iron uptake increases *P. chabaudi* pathogen burden

To investigate the effects of cellular iron availability on the host's response to malaria, we utilised a

transgenic mouse with a mutation in the cellular iron transporter TfR1. The $Tfrc^{Y20H/Y20H}$ mutation

decreases receptor internalisation by approximately 50%, resulting in decreased cellular iron uptake

(29). The effects of the $Tfrc^{Y20H/Y20H}$ mutation in erythroid cells are minimised due to a STEAP3-

mediated compensatory mechanism (29). At homeostasis, adult $Tfrc^{Y20H/Y20H}$ mice are healthy, normal-

sized, and not anaemic (Figure S1A-B). However, they have microcytic RBCs, compensated for by an

increase in RBCs (Figure S1C-D), and mildly suppressed liver and serum iron levels (Figure S1E-F).

$Tfrc^{Y20H/Y20H}$ and wild-type mice were infected with a recently mosquito-transmitted rodent malaria

strain, *P. chabaudi chabaudi* AS, which constitutively expresses GFP (hereafter referred to as *P.*

chabaudi) (46,47) (Figure 1A). Recently mosquito-transmitted parasites were used to mimic a natural

infection more closely, as vector transmission is known to regulate *Plasmodium* virulence and alter the

host's immune response (48-50). Consequently, parasitaemia is expected to be significantly lower upon

infection with recently mosquito-transmitted parasites, compared to infection with serially blood-

passaged parasites that are more virulent (47,48).

Strikingly, mice with decreased cellular iron uptake had significantly higher peak parasitaemia and

higher peak infected red blood cell (iRBC) counts (Figure 1B-C). The higher pathogen burden coincided

with more severe weight loss than wild-type mice (Figure 1D). This phenotype contrasts previous

studies, in which nutritional iron deficiency resulted in lower parasitaemia and increased survival of

malaria infected mice (49,50). Hence, our findings highlight a distinct role for cellular iron in malaria

pathology, which acts inversely to the protective effect of anaemia. This prompted us to investigate the

cause of the higher parasite burden observed in our model.

$Tfrc^{Y20H/Y20H}$ and wild-type mice have comparable malaria-induced RBC loss and anaemia

Anaemia-associated alterations of RBC physiology can affect malaria infection and have been put

forward as the main cause of both the protective effect of iron deficiency and the increased risk

associated with iron supplementation (10). We therefore monitored RBCs in wild-type and $Tfrc^{Y20H/Y20H}$

mice infected with *P. chabaudi.* Both genotypes displayed similar levels of malaria-induced RBC loss

and RBC recovery (Figure 1E). Moreover, $Tfrc^{Y20H/Y20H}$ and wild-type mice were equally severely

anaemic at the nadir of RBC loss, eight days post infection (dpi) (Figure 1F). At the chronic stage of

infection (20 dpi), however, wild-type mice showed improved recovery from anaemia compared to

$Tfrc^{Y20H/Y20H}$ mice (Figure 1G), consistent with a decreased ability of the $Tfrc^{Y20H/Y20H}$ cells to incorporate

iron.

While anaemia and RBC counts were comparable between both genotypes during infection, it was

nevertheless possible that differences in RBC physiology could alter the course of infection.

Consequently, we performed an *in vitro* invasion assay to determine whether $Tfrc^{Y20H/Y20H}$ RBCs were

more susceptible to *P. chabaudi* invasion. Fluorescently labelled wild-type or $Tfrc^{Y20H/Y20H}$ RBCs were

incubated *in vitro* with RBCs from a *P. chabaudi* infected wild-type mouse. Upon completion of one

asexual replication cycle, invasion was assessed, and the susceptibility index was calculated (Figure

1H). The RBC susceptibility indices of both genotypes were comparable (Figure 1I), thus indicating

that the higher parasite burden in $Tfrc^{Y20H/Y20H}$ mice was not due to a higher susceptibility of their RBCs

to *P. chabaudi* invasion.

Hyperferremia does not substantially alter *P. chabaudi* infection

In addition to anaemia, it has been suggested that that variations in host iron levels could affect blood-

stage *Plasmodium* parasite growth (51,52). Consequently, non-haem liver iron and serum iron was

measured in wild-type and $Tfrc^{Y20H/Y20H}$ mice upon *P. chabaudi* infection. At the peak of infection, both

genotypes had elevated liver and serum iron levels compared to homeostasis (Figure S1E-F & Figure

1J-K). Infected wild-type and $Tfrc^{Y20H/Y20H}$ mice had equivalent liver iron levels (Figure 1J), but serum

iron levels were higher in $Tfrc^{Y20H/Y20H}$ mice (Figure 1K).

The elevated serum iron observed in infected $Tfrc^{Y20H/Y20H}$ mice was consistent with their restricted

capacity to take up circulating transferrin-bound iron into tissues. However, we decided to investigate

whether this supraphysiological serum iron (i.e., hyperferremia) could alter *P. chabaudi* parasite

growth. To do this, we treated wild-type mice with a recombinant monoclonal anti-BMP6 IgG antibody

(αBMP6) or an isotype control (Figure S2A). αBMP6 treatment suppresses hepcidin expression and

elevates serum iron, as a consequence of unregulated release of iron from cellular stores (53) (Figure

S2A). *P. chabaudi* infected mice treated with αBMP6 had higher serum iron than isotype control-treated

mice on days 9 and 21 after infection (Figure S2B). Nevertheless, mice treated with αBMP6 and isotype

had comparable peak parasitaemia and peak iRBC counts, although αBMP6 treated mice appeared to

clear the parasites slightly more efficiently (Figure S2C-D). In addition, αBMP6 treatment did not

significantly alter weight loss (Figure S2E). Taken together, this data indicates that hyperferremia, as

observed in infected $Tfrc^{Y20H/Y20H}$ mice, does not increase peak parasitaemia. Accordingly, these findings

further indicate that iron uptake by non-erythropoietic cells is decisive in the host response to malaria.

Decreased cellular iron uptake attenuates the immune response to *P. chabaudi*

The immune response to malaria exerts control of parasitaemia, and the spleen is the main site of the

immune response to blood-stage malaria (39,54). Therefore, we assessed the splenic immune response

to *P. chabaudi* during the acute stage of infection (8 dpi). Interestingly, $Tfrc^{Y20H/Y20H}$ mice had attenuated

splenomegaly during acute *P. chabaudi* infection (Figure 2A-B), suggesting a disrupted splenic

response.

Malaria infection leads to an influx of mononuclear phagocytes (MNP) into the spleen, where they are

involved in cytokine production, antigen presentation, and phagocytosis of iRBCs (34,35,43). Upon *P.*

chabaudi infection, fewer MNPs were detected in the spleen of $Tfrc^{Y20H/Y20H}$ mice (Figure 2C). This

applied both to CD11b$^+$ Ly6C$^+$ MNPs (resembling inflammatory monocytes and/or monocyte-derived

macrophages) and to CD11c$^+$ MHCII$^+$ MNPs (resembling dendritic cells) (Figure 2D-E & Figure S3A).

In malaria infection, some MNPs can produce IFNγ that facilitates naïve CD4$^+$ T cell activation and

polarisation (34). Consequently, splenocytes from infected mice were cultured *ex vivo* with a protein

transport inhibitor, and intracellular cytokine staining was performed. Interestingly, fewer MNPs from

$Tfrc^{Y20H/Y20H}$ mice produced IFNγ compared to MNPs from wild-type mice (Figure 2F-G). Infected wild-

type and $Tfrc^{Y20H/Y20H}$ mice had comparable splenic neutrophil, eosinophil and NK cell numbers during

acute infection (8 dpi) (Figure S3B-D). Thus, mice with decreased cellular iron uptake had an attenuated

MNP response to *P. chabaudi* infection.

Cellular iron deficiency impairs the CD4+ T cell response to *P. chabaudi*

T cells, particularly CD4+ T cells, are a critical component of the immune response to blood-stage

malaria (55). Therefore, we assessed the splenic T cell response to acute *P. chabaudi* infection. The

total splenic CD4+ T cell count was comparable in both genotypes eight days after infection (Figure

3A). However, mice with decreased cellular iron uptake had a decreased proportion of effector CD4+ T

cells (Figure 3B), and, consequently, fewer total splenic effector CD4+ T cells than wild-type mice

(Figure 3C). In addition, the proportion of antigen-experienced CD44+ and PD1+CD4+ T cells was also

reduced in $Tfrc^{Y20H/Y20H}$ mice, re-enforcing their less activated state (Figure 3D-E). Moreover, fewer

$Tfrc^{Y20H/Y20H}$ CD4+ T cells were actively dividing, based on the proliferation marker KI-67 (Figure 3F).

This suggests a functional impairment of the CD4+ T cell response to *P. chabaudi* in mice with

decreased cellular iron uptake.

Similarly, the total CD8+ T cell count did not differ between genotypes (Figure S4A), but *P. chabaudi*

infected $Tfrc^{Y20H/Y20H}$ mice had fewer effector CD8+ T cells eight days after infection (Figure S4B-C).

However, there was no difference in the percentage of antigen-experienced (CD44+ or PD-1+) (Figure

S4D-E), proliferating (KI-67+) (Figure S4F) or IFNγ producing (Figure S4G) CD8+ T cells. Hence the

CD8+ T cell response to *P. chabaudi* infection was also attenuated, albeit to a lesser degree than CD4+

T cells.

T helper 1 (Th1) cells and other T helper subsets that express IFNγ are particularly important for malaria

immunity (55). Interestingly, the proportion of CD4+ T cells that expressed the Th1 transcription factor

T-BET was lower in mice with decreased cellular iron uptake (Figure 3G). Furthermore, fewer CD4 T

cells from $Tfrc^{Y20H/Y20H}$ mice produced IFNγ upon *ex vivo* restimulation (Figure 3H-I). Thus, further

strengthening the evidence of functional CD4+ T cell impairment in $Tfrc^{Y20H/Y20H}$ mice during *P.*

chabaudi infection.

To determine whether these impairments were T cell intrinsic and iron-dependent, we utilized naïve

CD4[+] T cells isolated from uninfected wild-type and *Tfrc*[Y20H/Y20H] mice. The cells were cultured *in vitro*

under Th1 polarising conditions for four days, in standard or iron-supplemented culture media (Figure

4A). *Tfrc*[Y20H/Y20H] lymphocytes can acquire iron under conditions where transferrin is hyper-saturated

and sufficient quantities of free iron are likely to be generated (29,56). Proliferation was significantly

impaired in *Tfrc*[Y20H/Y20H] CD4[+] T cells but could be rescued in a dose-dependent manner by iron

supplementation (Figure 4B-C). In addition, very few *Tfrc*[Y20H/Y20H] CD4[+] T cells cultured in standard

media produced IFNγ. However, iron supplementation completely rescued IFNγ production (Figure

4D-F). Hence, the CD4[+] T cell deficiencies observed in *Tfrc*[Y20H/Y20H] mice during *P. chabaudi* infection

were replicated *in vitro* and could be rescued by iron supplementation. These observations confirm that

host cell iron scarcity disrupts CD4[+] T cell function, leading to an inhibited CD4[+] T cell response to *P.*

chabaudi infection.

Decreased cellular iron uptake disrupts the germinal centre response to *P. chabaudi*

An efficient germinal centre (GC) response is required to generate high-affinity antibodies that enable

malaria clearance (36,37). In light of the impaired CD4[+] T cell response to *P. chabaudi* in *Tfrc*[Y20H/Y20H]

mice, we further examined the B cell supporting T follicular helper cell (Tfh) response. During the acute

stage of infection, a smaller proportion of CD4[+] T cells from *Tfrc*[Y20H/Y20H] mice expressed B cell co-

stimulation receptor ICOS (Figure 5A). ICOS is essential in malaria infection, as it is required to

maintain the Tfh cell response and sustain antibody production (57). In line with this, *Tfrc*[Y20H/Y20H] mice

had fewer Tfh cells, both during the acute (8 dpi) and chronic (20 dpi) stages of infection (Figure 5B-

C). Tfh cells support the activation, differentiation, and selection of high-affinity GC B cells, and are

an essential component of the humoral immune response to malaria (37). Therefore, we next sought to

assess the B cell response to *P. chabaudi* infection in *Tfrc*[Y20H/Y20H] and wild-type mice.

We observed no difference between genotypes in the total number of splenic B cells at the acute stage

of infection (8 dpi) (Figure 5D). However, mice with decreased cellular iron uptake had severely

impaired B cell activation and fewer antibody-secreting effector B cells (Figure 5E-F). Additionally,

Tfrc[Y20H/Y20H] mice had fewer GC B cells during acute infection (8 dpi) (Figure 5G). This effect remained

in the chronic stage of infection (20 dpi) (Figure 5H-I), indicating a prolonged immune inhibition caused

by restricted cellular iron availability.

Cellular iron deficiency impairs B cell function

To determine if the $Tfrc^{Y20H/Y20H}$ mutation also had cell-intrinsic and iron-dependent effects on B cells,

their functionality was further investigated *in vitro.* B cells were isolated from uninfected $Tfrc^{Y20H/Y20H}$

and wild-type mice, activated, and cultured in standard or iron-supplemented media for three days

(Figure 6A). Expression of the B cell activation marker LAT-1 was lower on $Tfrc^{Y20H/Y20H}$ B cells than

wild-type (Figure 6B). However, LAT-1 expression was rescued by iron supplementation, indicating

improved B cell activation (Figure 6B). $Tfrc^{Y20H/Y20H}$ B cell proliferation was also severely impaired

compared to wild-type cells, but was rescued by iron supplementation in a dose-dependent manner

(Figure 6C-D). Iron scarcity also inhibited the potential of $Tfrc^{Y20H/Y20H}$ B cells to differentiate into

antibody-secreting and class-switched cells (Figure 6E-G). This impairment was fully restored upon

iron supplementation (Figure 6E-G). Overall, our data clearly show that the activation, proliferation,

and differentiation of $Tfrc^{Y20H/Y20H}$ B cells were impaired, demonstrating that cellular iron deficiency

causes cell-intrinsic B cell dysfunction.

Decreased cellular iron uptake ameliorates *P. chabaudi*-induced liver pathology

$Tfrc^{Y20H/Y20H}$ mice experienced higher *P. chabaudi* parasitaemia and an inhibited immune response.

However, the precise consequences of this disease phenotype remained unclear. Aspects of the immune

response, such as the cytokine profile and the balance between pro-inflammatory and

immunoregulatory responses, can tip the scales toward protection or pathology in malaria (39). Hence,

an attenuated immune response could cause hyperparasitaemia, but it may also be crucial in limiting

immunopathology. We therefore set out to characterise key indicators of malaria disease severity.

We first measured circulating levels of angiopoietin-2 (ANG-2) and alanine transferase (ALT). ANG-

2 is a marker of endothelial activation that correlates with malaria disease severity and mortality in

humans (58,59). Liver damage is also indicative of severe malaria (60), and ALT is a standard marker

of liver damage. There was a trend towards lower ANG-2 and significantly decreased ALT in

$Tfrc^{Y20H/Y20H}$ mice eight days after *P. chabaudi* infection, suggesting milder pathology (Figure 7A-B).

Considering the substantial difference in serum ALT between genotypes, we further examined the

malaria induced liver pathology. $Tfrc^{Y20H/Y20H}$ mice had lower expression of the tissue-damage and

inflammation-induced acute phase protein genes *Saa1* and *Fga* (Figure S5A-B). Furthermore, while

both genotypes developed malaria-induced hepatomegaly, there was a trend toward less severe

hepatomegaly in $Tfrc^{Y20H/Y20H}$ mice (Figure S5C).

Histological analysis revealed hepatic pathology in all *P. chabaudi* infected mice, characterised by

hepatocellular necrosis, sinusoidal dilatation, glycogen depletion, and infiltration by mononuclear

immune cells (Figure 7C-D & Figure S5D-E). Interestingly, no polymorphonuclear immune cell

infiltration was observed. All infected wild-type mice developed confluent necrosis (areas of lobular

disarray, eosinophilia, and loss of glycogen deposits, score ≥3), and most individuals (8 out of 11) also

displayed bridging necrosis (areas of confluent necrosis extending across multiple lobules, score=4)

(Figure 7E & Figure S5F). In contrast, severe focal necrosis or confluent necrosis (score ≥3) was

detected in just over half (6 out of 10) infected $Tfrc^{Y20H/Y20H}$ mice, and only four individuals developed

bridging necrosis (Figure 7E & Figure S5F). Hence, the proportion of mice that developed severe

hepatic necro-inflammation (score ≥3) upon *P. chabaudi* infection was significantly smaller in

$Tfrc^{Y20H/Y20H}$ than in wild-type mice (Figure 7E).

Excess reactive liver iron and haem are known to cause liver damage in malaria (61,62). However, we

observed no differences in total non-haem liver iron (Figure 1I) or liver lipid peroxidation, which

correlates with ROS levels (Figure S5G). Hence, it is unlikely that tissue level variations in hepatic

reactive iron or haem can explain the difference in liver damage. In addition, we measured the

expression of two genes that are known to have a hepatoprotective effect in the context of iron loading

in malaria: *Hmox1* (encodes haemoxygenase-1 (HO-1)) and *Fth1* (encodes ferritin heavy chain). Liver

gene expression of *Hmox1* was higher in $Tfrc^{Y20H/Y20H}$ mice, while the expression of *Fth1* did not differ

between genotypes, eight days after infection (Figure S5H-I). Thus, the higher expression of *Hmox1*

may have contributed to a hepatoprotective effect in $Tfrc^{Y20H/Y20H}$ mice.

During malaria infection, endothelial activation leads to increased adhesion and sequestration of iRBCs,

resulting in hepatic vascular occlusions and hypoxia that cause damage (2,63). Fewer sequestration,

rosetting, and vascular occlusion events were detected in liver sections from $Tfrc^{Y20H/Y20H}$ mice eight

days after *P. chabaudi* infection (Figure 7F). Together with the trend toward lower ANG-2 levels in

$Tfrc^{Y20H/Y20H}$ mice (Figure 7A), this indicates that decreased endothelial activation and iRBC

sequestration contributed to the attenuated liver pathology observed in $Tfrc^{Y20H/Y20H}$ mice.

Inflammation also causes severe disease and liver pathology in malaria (39,61,64). Hence, hepatic

inflammation was approximated by measuring the expression of genes encoding pro-inflammatory

cytokines IFNγ, TNFα, and IL-1β. We observed no difference in the expression of *Ifng* or *Tnf,* but *Il1b*

expression was lower in $Tfrc^{Y20H/Y20H}$ mice eight days after *P. chabaudi* infection (Figure S5J-L).

Moreover, immunohistochemistry staining showed reduced infiltration of leukocytes (CD45[+] cells) in

livers of $Tfrc^{Y20H/Y20H}$ mice (Figure 7G-H). Additionally, a smaller proportion of liver leukocytes

(CD45[+]) were effector immune cells such as dendritic cells, CD44[+] CD4[+] T cells, and CD44[+] CD8[+] T

cells (Figure 7I-L). Taken together, this data shows that host cell iron scarcity leads to an attenuated

hepatic immune response during *P. chabaudi* infection.

DISCUSSION

Iron deficiency impacts malaria infection in humans (7–9), but beyond the effects of anaemia (10), little

is known about how host iron deficiency influences malaria infection. Here we investigated how

restricted cellular iron acquisition influenced *P. chabaudi* infection in mice. $Tfrc^{Y20H/Y20H}$ mice developed

comparable malaria-induced anaemia to wild-type mice, and RBC susceptibility to parasite invasion

did not differ between genotypes. This therefore allowed us to largely decouple the effects of anaemia

from other effects of iron on the host response to malaria. Strikingly, $Tfrc^{Y20H/Y20H}$ mice displayed an

attenuated *P. chabaudi* induced splenic and hepatic immune response. This immune inhibition was

associated with increased parasitaemia and mitigated liver pathology. Hence, for the first time, we show

a role for host cellular iron acquisition via TfR1 in modulating the immune response to malaria, with

downstream effects on both pathogen control and host fitness.

On first inspection, the higher parasite burden observed in $Tfrc^{Y20H/Y20H}$ mice may appear to be a severe

consequence of cellular iron deficiency. In humans, however, high parasitaemia is not sufficient to

cause severe disease (65). Moreover, the risk of severe malarial disease decreases significantly after

only one or two exposures, whereas anti-parasite immunity is only acquired after numerous repeated

exposures (2,66). It follows that mitigating immunopathology may be more important than restricting

parasite growth for host survival. As previously noted, the $Tfrc^{Y20H/Y20H}$ mutation has relatively mild

consequences for erythropoietic parameters compared to other haematopoietic lineages (29,30).

However, in humans with normal TfR1-mediated iron uptake, iron deficiency sufficient to cause

immune cell iron scarcity also normally causes anaemia (67). In such circumstances, parasite growth

would likely be limited by anaemia, with the final result that iron deficiency may be protective overall,

if it also minimises aspects of immunopathology.

Previous work has demonstrated the importance of regulating tissue haem and iron levels to prevent

organ damage in malaria (61,62,68,69). For example, HO-1 plays an important role in detoxifying free

haem that occurs as a result of haemolysis during malaria infection, thus preventing liver damage due

to tissue iron overload, ROS and inflammation (61). Interestingly, infected $Tfrc^{Y20H/Y20H}$ mice had higher

expression of $Hmox1$, but levels of liver iron and ROS comparable to that of wild-type mice.

Consequently, this may be indicative of increased haem processing that could have a tissue protective

effect. In humans, there is a correlation between transferrin saturation and ALT levels in patients with

symptomatic malaria (62,70), suggesting that iron status may be linked to malaria-induced liver

pathology in humans. However, it can be difficult to interpret measures of iron status in malaria infected

individuals, since those parameters can be altered by inflammation and RBC destruction. Our findings

reveal additional dimensions through which host iron status impacts malaria-induced tissue damage.

The mitigated liver damage that we observed in $P.$ $chabaudi$ infected $Tfrc^{Y20H/Y20H}$ mice can likely be

explained by a combination of factors; increased expression of hepatoprotective HO-1, decreased

immune mediated endothelial activation, iRBC sequestration, and hepatic vascular occlusion, as well

as, inhibited hepatic inflammation.

The pro-inflammatory immune response to malaria has downstream effects on cytoadherence, as pro-

inflammatory cytokines activate endothelial cells, leading to higher expression of receptors for

cytoadherence (2). As a consequence, $P.$ $chabaudi$ infected mice that lack adaptive immunity or IFNγ-

receptor signalling, have substantially decreased sequestration of iRBCs in the liver, and no detectable

liver damage (as measured by ALT) (63). Endothelial cells can also be activated by direct interactions

with iRBCs (2), and in humans, ANG-2 correlates with estimated parasite biomass (59). However,

although *P. chabaudi* infected $Tfrc^{Y20H/Y20H}$ mice had higher peak parasitaemia, they had fewer hepatic

sequestration, rosetting, and vascular occlusion events and lower ANG-2 levels. The attenuated innate

and adaptive immune response is the most probable cause of decreased endothelial activation and

hepatic microvascular obstruction in $Tfrc^{Y20H/Y20H}$ mice. This, in turn, likely contributed to the clearly

mitigated liver pathology, in spite of the higher parasitaemia. Upon *P. chabaudi* infection, we observed

extensive infiltration of mononuclear leukocytes into the liver, but this response was repressed in

$Tfrc^{Y20H/Y20H}$ mice. Specifically, infected $Tfrc^{Y20H/Y20H}$ mice had fewer effector-like immune cells in the

liver. Hepatic immune cells can contribute to liver damage in malaria, for example, by producing pro-

inflammatory cytokines or through bystander killing of hepatocytes (71). Consequently, a weaker

hepatic pro-inflammatory immune response likely limited immunopathology and ameliorated malaria-

induced liver damage in mice with cellular iron deficiency.

We have previously shown that hepcidin mediated hypoferremia inhibits the immune response to

influenza infection in mice (21). In influenza, cellular iron scarcity exacerbated pulmonary tissue

damage, because failed adaptive immunity led to an exacerbated inflammatory response and poor

pathogen control (21). In contrast, we observed that decreased cellular iron acquisition inhibited both

the innate and adaptive immune response to malaria, ultimately mitigating malaria-induced hepatic

tissue damage and inflammation. This highlights the complex effects of iron deficiency on the immune

system and underscores the need to consider its effect on different infectious diseases in a pathogen-

specific manner. A better understanding of how host iron status affects immunity to infection could

benefit the development of improved antimicrobial therapies and increase the safety of iron deficiency

therapies.

The inhibited innate immune response to *P. chabaudi* in $Tfrc^{Y20H/Y20H}$ mice likely contributed to both the

increased pathogen burden and the decreased liver pathology. Splenic MNPs are important for

controlling parasitaemia (34,35,72), but MNPs are also vital for maintaining tissue homeostasis and

preventing tissue damage in malaria (43,73). Although other innate cells, such as neutrophils, NK cells

and γδT cells are an important part of the immune response to malaria, only the MNP response was

distinctly impaired in *Tfrc*$^{Y20H/Y20H}$ mice. Notably, neutrophils are known to be sensitive to iron

deficiency (16,74) and to affect both immunity and pathology in malaria (75,76). However, in the

context of recently mosquito-transmitted *P. chabaudi* it appears that monocytes and macrophages,

rather than granulocytes, may be particularly important for parasite control and tissue homeostasis

(43,72).

CD4$^+$ T cells and B cells become cell intrinsically dysfunctional during iron scarcity, as we have

demonstrated *in vitro*. However, such cell-intrinsic effects are likely further aggravated by interactions

with other iron-depleted cells *in vivo*. For example, CD4$^+$ T cells support the B cell response to malaria

(37,77), and the repressed CD4$^+$ T cell response to *P. chabaudi* in *Tfrc*$^{Y20H/Y20H}$ mice presumably further

constrained the B cell response. Proliferation is an aspect of immune cell function that appears to be

particularly sensitive to iron deficiency (14,20,21). Unsurprisingly, we also see the most significant

inhibitory effect on immune cell populations that expand greatly during *P. chabaudi* infection. In

addition, proliferation is often required for lymphocyte differentiation and effector function (78), and

the differentiation of Tfh and Th1 cells in malaria depends on a highly proliferative precursor CD4$^+$ T

cell subset (79). T cells from *Tfrc*$^{Y20H/Y20H}$ mice also had decreased KI-67 expression, further confirming

impaired proliferation as a critical mechanism of immune inhibition under conditions of cellular iron

scarcity. CD4$^+$ T cells that produce pro-inflammatory cytokine are also sensitive to iron restriction, as

we have shown for IFNγ, and as has been shown previously for IL-2 and IL-17 (80,81). Interestingly,

iron overload can also alter CD4$^+$ T cell cytokine production, and excess iron can have an inhibitory

effect on IFNγ production (22,82). These observations underline that iron imbalance at either extreme

can disturb immune cell function.

Despite the higher peak parasitaemia in *Tfrc*$^{Y20H/Y20H}$ mice, both genotypes were able to clear *P. chabaudi*

parasites at a comparable rate and prevent recrudescence. It follows that even a weakened humoral

immune response appears to be sufficient to control *P. chabaudi* infection. However, our study did not

investigate the effects of immune cell iron deficiency on the formation of long-term immunity, which

may have been more severely affected. The impaired GC response, in particular, suggests that iron

deficiency could counteract the formation of efficient immune memory to subsequent malaria

infections. This is in line with human observational studies that have found a link between iron

deficiency and weak antibody responses to *P. falciparum* (7,44,45). In humans, anti-parasite immunity

forms very slowly and only after numerous repeated exposures to malaria infection (2). Some have

suggested that this effect could be explained by impaired immune cell function in malaria (83,84), and

future studies should consider whether inhibited immunity as a result of iron deficiency could contribute

to this phenomenon. Moreover, the extensive geographical and epidemiological overlap of iron

deficiency and malaria (1,6,13) makes this concept particularly relevant for further research.

It remains to be seen what the broader importance of cellular iron is in human malaria infection, in

particular within the diverse genetic context of both humans and parasites, found in malaria endemic

regions. Murine models of malaria are useful in providing hypothesis-generating results, but such

findings ultimately ought to be confirmed and developed further through studies in human populations.

This study revealed that decreased host cell iron acquisition inhibits the immune response to malaria

and ameliorates hepatic damage, despite a higher parasite load and similar degree of anaemia, in mice.

Altogether, our data highlight a previously underappreciated role for host cell iron in the trade-off

between pathogen control and immunopathology, and add to our understanding of the complex

interactions between iron deficiency and malaria. Hence, these findings have important implications for

these two widespread and urgent global health problems.

METHODS

Mice

Tfrc$^{Y20H/Y20H}$ mice were initially provided by Professor Raif Geha, Boston Children's Hospital/Harvard

Medical School (29), and they were subsequently bred in-house at the University of Oxford. Control

wild-type C57BL/6JOlaHsd mice were purchased from Envigo and co-housed with *Tfrc$^{Y20H/Y20H}$* mice

for 2-3 weeks prior to *P. chabaudi* infection. All mice were housed in individually ventilated specific-

pathogen-free cages under normal light conditions (light 07.00-19.00, dark 19.00-07.00) and fed

standard chow containing 188 ppm iron (SDS Dietex Services, diet 801161) ad-libitum. Age-matched,

8-13 week-old female mice were used for experiments. Females were exclusively utilised to prevent

loss of animals due to fighting, and to minimise the risk of severe adverse events from *P. chabaudi*

infection, which is higher in males (85). Euthanasia was performed through suffocation by rising CO_2

concentrations, and death was confirmed by cervical dislocation.

Ethics

413 All animal experiments were approved by the University of Oxford Animal Welfare and Ethical Review

414 Board and performed following the U.K. Animals (Scientific Procedures) Act 1986, under project

415 licence P5AC0E8C9.

Parasites and infection

Transgenic recently mosquito-transmitted *P. chabaudi chabaudi* AS parasites expressing GFP (46,47)

were obtained from the European Malaria Reagent Repository at the University of Edinburgh. To

generate iRBCs for blood-stage *P. chabaudi* infections, frozen parasite stocks were rapidly thawed by

hand and injected intraperitoneally (i.p.) into a single wild-type mouse. Once ascending parasitaemia

reached 0.5-2%, the animal was euthanised and exsanguinated through cardiac puncture. Subsequent

experimental infections were immediately initiated from the collected blood, by intravenously (i.v.)

injecting 10^5 iRBCs in 100 uL Alsever's solution. Uninfected control mice received Alsever's solution

only.

To monitor *P. chabaudi* infection, blood was collected through micro-sampling from the tail vein of

infected mice. Parasitaemia, iRBC count and RBC count was measured by flow cytometry, as

previously described (46). Briefly, 2 μL of blood was diluted in 500 μL Alsever's solution immediately

after collection. The solution was further diluted 1:10 in PBS before acquisition on an Attune NxT Flow

Cytometer (Thermo Fisher Scientific). A fixed volume of each sample was acquired, thus allowing for

the enumeration of total RBCs and iRBCs per μL of blood.

αBMP6 treatment

432 In order to experimentally raise serum iron levels, an αBMP6 human IgG monoclonal blocking antibody

433 that cross-reacts with murine BMP6 (53) was administered. Control mice received a human IgG4

434   isotype control antibody. Both antibodies were diluted in 100 µL PBS and injected i.p at a dose of

435   approximately 10 mg/kg body weight.

Tissue processing

Organs and tissues were harvested shortly after euthanasia and kept cold until further analysis could be

performed. Liver and spleen indices were calculated as the mass of the respective organs relative to

mouse body weight. Blood was collected into appropriate blood collection tubes (BD Microtainer

K2EDTA for whole blood or BD Microtainer SST/Sarstedt Microvette 100 Serum for serum), either by

tail vein sampling or by cardiac puncture after euthanasia. Serum was prepared by centrifugation of the

collection tubes at 10,000 x g for 5 min, and stored at -80° C.

Blood analysis

RBC count, haemoglobin, and mean cell volume was measured from whole blood using an automatic

KX-21N Haematology Analyser (Sysmex). Serum levels of ANG-2 and ALT were measured according

to the producers' instructions, using the Mouse ALT ELISA Kit (ab282882, Abcam) and the Mouse/Rat

Angiopoietin-2 Quantikine ELISA Kit (MANG20, R&D Systems), respectively. Serum cytokines were

measured using the LEGENDplex Mouse Inflammation Panel (740446, BioLegend) bead-based

immunoassay. The assay was performed according to the manufacturer's instructions, except that the

protocol was adapted to use half-volumes.

***In vitro P. chabaudi* invasion assay**

To assess the susceptibility of wild-type and $Tfrc^{Y20H/Y20H}$ RBCs to *P. chabaudi* invasion, blood was

collected from a *P. chabaudi* infected wild-type mouse during ascending parasitaemia (donor RBCs/Y),

and from uninfected wild-type and $Tfrc^{Y20H/Y20H}$ mice (target RBCs/X). To remove leukocytes, the blood

was passed through a cellulose (C6288, Merck) packed column, as previously described (86). The target

RBCs were fluorescently labelled with 1 µM CellTrace Far Red (C34572, Thermo Fisher Scientific) in

PBS, by diluting blood 1:10 with CellTrace solution and incubating in the dark for 15 min at 37° C,

mixing the samples every 5 min. Afterward, the cells were washed twice in R10 media (RPMI-1640

with 10% FBS, 2 mM glutamine (G7513, Merck), 1% penicillin-streptomycin (P0781, Merck), 50 µM

2-Mercaptoethanol (31350, Thermo Fisher Scientific)) and resuspended in R10 media supplemented

with 0.5 mM sodium pyruvate (1136007050, Thermo Fisher Scientific). 2×10^7 donor RBCs and $2 \times$

10^7 fluorescently labelled target RBCs were plated in the same well of a 96-well plate, and incubated

overnight (~16 h) in a candle jar at 37° C, to allow sufficient time for schizonts to develop and release

merozoites. Invasion was measured as GFP^+ RBCs and compared by calculating the susceptibility

index, as previously described (87).

$$SI = \frac{\frac{X \text{ RBC}}{X \text{ iRBC}}}{\frac{Y \text{ RBC}}{Y \text{ iRBC}}}$$

X = fluorescently labelled target wild-type or $Tfrc^{Y20H/Y20H}$ RBCs

Y = donor derived wild-type RBCs

Iron measurements

Serum iron measurements were performed on an Abbott Architect c16000 automated analyser by

Oxford University Hospitals Clinical Biochemistry staff using the MULTIGENT Iron Kit (Abbott), or

using a Pentra C400 automated analyser with the Iron CP ABX Pentra Kit (HORIBA Medical).

Non-haem liver iron measurements were performed as previously described (88). In short, pieces of

liver tissue were collected, snap-frozen, and stored at -80° C. The tissue was dried at 100º C for ~6 h,

weighed, and then digested in 10% trichloroacetic acid / 30% hydrochloric acid in water for ~20 hours

at 65ºC. Subsequently, a chromogen reagent containing 0.1% bathophenanthrolinedisulphonic acid

(Sigma, 146617) / 0.8% thioglycolic acid (Sigma, 88652) / 11% sodium acetate in water was added,

and the absorbance at 535 nm measured. The iron content was determined by comparing the samples

against a standard curve of serially diluted ammonium ferric citrate (F5879, Merck).

Flow cytometry

Single cell suspensions for flow cytometry were prepared through mechanical and enzymatic

dissociation. Spleens were passed through 70 µM cell strainers, incubated with 120 Kunitz U/mL

deoxyribonuclease I (DN25, Merck) in R10 for 15 min with agitation, and passed through 40 µM cell

strainers. Livers were perfused with PBS with 10% FBS prior to harvest. To prepare single cell

suspensions, the livers were disrupted with scissors, incubated with 0.5 mg/mL collagenase IV (C5138,

Merck) and 120 Kunitz U/mL DNAse I in R10 for 45 min with agitation, and passed through 70 μM

cell strainers. RBC lysis was subsequently performed by resuspending pelleted cells in tris-buffered

ammonium chloride buffer (0.017 M Tris / 0.14 M NH_4Cl, adjusted to pH 7.2 with HCl) and incubating

for ~5 min on ice before washing with R10.

Immune cells were isolated from livers by Percoll (17-08-91, GE Healthcare) separation. Single-cell

suspensions were gently overlayed onto 33% Percoll and centrifuged for 25 min at 800 x g. After

centrifugation, the supernatant was discarded and the remaining leukocytes were washed twice with

R10.

For intracellular cytokine staining, splenocytes were cultured *ex vivo* in R10 at 5-2 x 10^5 cells/mL, in

round-bottom tissue culture treated 96-well plates, with protein transport inhibitor Brefeldin A for 4-6

h at 37° C, 5% CO^2. To activate T cells, 0.5 μg/mL anti-mouse CD3 (100201, BioLegend) was added

to splenocytes from *P. chabaudi* infected mice.

Cells were counted using a CASY Cell Counter and Analyser (BOKE), and 1-5 x 10^6 cells were stained

for flow cytometry. The cells were washed in PBS, blocked with TruStain FcX (101319, BioLegend),

and stained with a viability dye (NIR Fixable Viability Kit (42301/5, BioLegend) or LIVE/DEAD

Fixable Near-IR Dead Cell Stain Kit (L34975, Thermo Fisher Scientific)) for ~10 min at 4° C in the

dark. Next, fluorophore-conjugated antibodies were added to the cells and incubated for ~20 min. The

cells were washed twice in PBS and fixed by incubating with Fixation Buffer (420801, BioLegend) for

~10 min at 4° C in the dark. Alternatively, the cells were fixed and permeabilised using eBioscience

FOXP3/Transcription Factor Staining Buffer Set (00-5523-00, Thermo Fisher Scientific), and

transcription factor staining was performed, according to the manufacturer's instructions. Intracellular

cytokine staining was performed after permeabilization with Intracellular Staining Permeabilization

Wash Buffer (421002, BioLegend) for ~30 min, according to the manufacturer's protocol. The samples

were acquired on an Attune NxT or BD LSR Fortessa X-20 (BD) flow cytometer.

***In vitro* culture of primary immune cells**

Naïve CD4[+] T cells and B cells were isolated according to the manufacturer's instructions from mixed

splenocyte and lymph node single-cell suspensions using the EasySep Mouse Naïve CD4[+] T Cell

Isolation Kit (19765, STEMCELL), or from splenocyte single-cell suspensions using the EasySep

Mouse B Cell Isolation Kit (19854, STEMCELL). The isolated cells were fluorescently labelled with 5

μM CellTrace Violet (C34571, Thermo Fisher Scientific) in PBS for 8 min at 37° C and washed twice

in R10 media. Cell counting was performed with a CASY Cell Counter and Analyser.

For CD4[+] T cells, flat-bottom tissue culture treated 96-well plates were pre-coated with 5 μg/mL anti-

mouse CD3 and the cells were seeded at 5×10^5 cells/mL. They were cultured in Th1-polarising media

consisting of R10 with 1 μg/mL anti-mouse CD28 (102101, BioLegend), 5 μg/mL anti-mouse IL-4

(504102, BioLegend), 10 ng/mL IL-12 (505201, BioLegend), 25 U/mL IL-2 (575404, BioLegend) and

50 μM 2-Mercaptoethanol. The media was replaced after 48 h of culture. To iron supplement the culture

medium, iron sulphate heptahydrate (F8633, Merck) was added at the previously specified

concentrations.

B cells were cultured at 7.5×10^5 cells/mL in flat-bottom tissue culture treated 96-well plates, in R10

media with 1% MEM amino acids (11130, Thermo Fisher Scientific), 2 μg/mL LPS (tlrl-peklps,

InvivoGen), 10 ng/mL IL-4 (574302, BioLegend), 10 ng/mL IL-5 (581502, BioLegend) and 50 μM 2-

Mercaptoethanol. Ammonium ferric citrate was added at the specified concentrations to iron

supplement the media.

CD4[+] T cells were cultured for 96 h and B cells for 72 h at 37° C, 5% CO2, before flow cytometry

staining. The type of iron used to supplement the culture media was chosen to optimise cell viability.

Gene expression analysis

Gene expression analysis by quantitative real-time PCR, was performed on liver samples preserved in

RNAlater Stabilization Solution and stored at -80° C (AM7020, Thermo Fisher Scientific). The tissue

was homogenised with a TissueRuptor II (9002725, QIAGEN) before total RNA was extracted using

the RNeasy Plus Mini Kit (74136, QIAGEN), according to the manufacturer's protocols. cDNA was

synthesised using the High-Capacity RNA-to-cDNA Kit (4387406, Thermo Fisher Scientific) and subsequent gene expression analysis was performed on 1-5 ng/mL cDNA, using TaqMan Gene Expression Master Mix (4369016, Thermo Fisher Scientific) and the TaqMan Gene Expression Assays (Thermo Fisher Scientific) listed in Table 1, all according to the manufacturers' instructions. An Applied Biosystems 6500 Fast Real-Time PCR System (Thermo Fisher Scientific) instrument was used to run the samples, and the relative gene expression was calculated through the $2^{-\Delta CT}$ method (89).

Table 1. **List of TaqMan Gene Expression Assays.**

Protein	Gene	Assay code
Fibrinogen alpha chain	*Fga*	Mm00802584_m1
Hypoxanthine-guanine phosphoribosyltransferase	*Hprt*	Mm01545399_m1
Interferon γ	*Ifng*	Mm01168134_m1
Interleukin 1β	*Il1b*	Mm00434228_m1
Serum amyloid A1	*Saa1*	Mm00656927_g1
Tumour necrosis factor α	*Tnf*	Mm00443258_m1

Liver histology

Liver samples were fixed with 4% paraformaldehyde in PBS and embedded in paraffin. Following deparaffinization with xylene and hydration by a passage through a grade of alcohols, 3 µm-thick sections were stained with haematoxylin-eosin, and Periodic Acid-Schiff, before and after diastase digestion, at IPATIMUP Diagnostics, Portugal, using standard procedures.

Histopathology scores for lobular necro-inflammatory activity were assigned using the criteria of Scheuer (90) for the grading of chronic hepatitis. In short, the scores were assigned as follows, 0 = inflammation absent, 1 = inflammation but no hepatocellular death, 2 = focal necrosis (one or a few necrotic hepatocytes/acidophil bodies), 3 = severe focal death, confluent necrosis without bridging, and 4 = damage includes bridging necrosis. Sections were scored independently by two investigators with experience in liver histopathology who were blinded to the experimental groups. The total numbers of RBC endothelial cytoadherence (sequestration), rosetting and vascular occlusion events were counted

blindly in random high-power (×400 magnification) fields of liver sections. Images were captured using

an Olympus BX50 photomicroscope.

For the immunohistochemical detection of $CD45^+$ cells, liver sections were subjected to antigen

retrieval with citrate buffer, endogenous peroxidases were blocked with 0.6% H_2O_2 and non-specific

antigens were blocked with 5 % bovine serum albumin. Samples were incubated with goat anti-mouse

CD45 antibody (1:50, AF114, R&D Systems, MN, USA) followed by horseradish peroxidase-

conjugated rabbit anti-goat IgG (1:250, R-21459, ThermoFisher Scientific). Immunoreactivity was

visualized using 3,3'-diaminobenzidine. Quantification was performed by counting positive cells in 5

random fields per liver at 200× magnification using QuPath Open Software for Bioimage Analysis

(version 0.4.0).

Thiobarbaturic acid reactive substances assay

Liver ROS/lipid peroxidation was appreciated by quantifying malondialdehyde, using the TBARS

Assay Kit (700870, Cayman Chemical) as described by the manufacturer. Briefly, tissue homogenates

were prepared from snap-frozen liver tissue by adding 1 mL RIPA buffer per 100 mg of tissue, and

lysing using Precellys soft tissue homogenising tubes (KT03961-1-003.2, Bertin Instruments)

according to manufacturers instruction. The lysates were allowed to react with thiobarbaturic acid at

95° C for 1 h, cooled on ice, and centrifuged for 10 min at 1,600 x g at 4° C. Subsequently, the

absorbance of the lysates at 530 nm was measured.

Software and statistical analysis

All flow cytometry data analysis was performed using FlowJo analysis software (BD). Graphs were

generated using GraphPad Prism (GraphPad Software).

Statistical analysis was also performed in GraphPad Prism and differences were considered statistically

different when $p<0.05$ (* $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$). The D'Agostino-Pearson

omnibus normality test was used to determine normality/lognormality. Parametric statistical tests (e.g.

Welch's t-test) were used for normally distributed data. For lognormal distributions, the data was log-

transformed prior to statistical analysis. Where data did not have a normal or lognormal distribution, or

<mark>too few data points were available for normality testing, a nonparametric test (e.g. Mann-Whitney test)</mark>

<mark>was applied.</mark> A t-test (or a comparable nonparametric test) was used to compare the means of two

groups. As a rule, t-tests were performed with Welch's correction, as it corrects for unequal standard

deviations but does not introduce error when standard deviations are equal. Two-way ANOVA was

used for analysis with two categorical variables and one continuous variable. The applied statistical test

and sample size (n) is indicated in each figure legend.

ACKNOWLEDGEMENTS

The authors thank the staff of the University of Oxford Department of Biomedical Services for

assistance with animal husbandry and procedures, and the Weatherall Institute of Molecular Medicine

Flow Cytometry Facility for technical assistance. We are also grateful to the Clinical Biochemistry Unit

(Oxford University Hospitals NHS Foundation Trust), and Samira Lakhal-Littleton and Goran

Mohammad at the Oxford University Department of Physiology, Anatomy & Genetics for assistance

with biochemical measurements. Additionally, we would like to sincerely thank Wiebke Nahrendorf,

Philip Spence and Joanne Thompson at the University of Edinburgh for helpful scientific discussions

and for providing us with *P. chabaudi* parasites.

FUNDING

This work was supported by:

Wellcome Trust Infection, Immunology & Translational Medicine doctoral programme grant (awarded

to SKW, grant no. 108869/Z/15/Z)

UK Medical Research Council (MRC Human Immunology Unit core funding awarded to HD, grant no.

MC_UU_12010/10)

Portuguese National Funds through FCT—Fundação para a Ciência e a Tecnologia, I.P., under the

project UIDB/04293/2020

Wellcome Trust Infection, Immunology & Translational Medicine doctoral programme grant (awarded

to FCR, grant no. 203803/Z16/Z)

CONFLICT OF INTERESTS

The authors declare that they have no conflict of interest.

REFERENCES

1.   World Malaria Report 2022. Geneva: World Health Organisation.

2.   Cowman AF, Healer J, Marapana D, Marsh K. Malaria: Biology and Disease. Cell. 2016 Oct

20;167(3):610–24.

3. Coffey R, Ganz T. Iron homeostasis: An anthropocentric perspective. J Biol Chem.
2017;292(31):12727–34.

4. Drakesmith H, Prentice AM. Hepcidin and the Iron-Infection Axis. Science (80-). 2012 Nov
9;338(6108):768–72.

5. Pasricha SR, Tye-Din J, Muckenthaler MU, Swinkels DW. Iron deficiency. Lancet.
2021;397(10270):233–48.

6. Kassebaum NJ, Jasrasaria R, Naghavi M, Wulf SK, Johns N, Lozano R, et al. A systematic
analysis of global anemia burden from 1990 to 2010. Blood. 2014 Jan 30;123(5):615–24.

7. Nyakeriga AM, Troye-Blomberg M, Dorfman JR, Alexander ND, Bäck R, Kortok M, et al. Iron
deficiency and malaria among children living on the coast of Kenya. J Infect Dis.
2004;190(3):439–47.

8. Jonker FAM, Calis JCJ, van Hensbroek MB, Phiri K, Geskus RB, Brabin BJ, et al. Iron status
predicts malaria risk in Malawian preschool children. PLoS One. 2012;7(8):1–8.

9. Gwamaka M, Kurtis JD, Sorensen BE, Holte S, Morrison R, Mutabingwa TK, et al. Iron
deficiency protects against severe plasmodium falciparum malaria and death in young children.
Clin Infect Dis. 2012;54(8):1137–44.

10. Clark MA, Goheen MM, Fulford A, Prentice AM, Elnagheeb MA, Patel J, et al. Host iron status
and iron supplementation mediate susceptibility to erythrocytic stage Plasmodium falciparum.
Nat Commun. 2014 Dec 25;5(1):4446.

11. Sazawal S, Black RE, Ramsan M, Chwaya HM, Stoltzfus RJ, Dutta A, et al. Effects of routine
prophylactic supplementation with iron and folic acid on admission to hospital and mortality in
preschool children in a high malaria transmission setting: community-based, randomised,
placebo-controlled trial. Lancet. 2006 Jan;367(9505):133–43.

12.   Neuberger A, Okebe J, Yahav D, Paul M. Oral iron supplements for children in malaria-endemic

areas. Cochrane Database Syst Rev. 2016 Feb 27;2016(2).

13.   Muriuki JM, Mentzer AJ, Mitchell R, Webb EL, Etyang AO, Kyobutungi C, et al. Malaria is a

cause of iron deficiency in African children. Nat Med. 2021;27(4):653–8.

14.   Brekelmans P, Van Soest P, Leenen PJM, van Ewijk W. Inhibition of proliferation and

differentiation during early T cell development by anti-transferrin receptor antibody. Eur J

Immunol. 1994;24(11):2896–902.

15.   Ned RM, Swat W, Andrews NC. Transferrin receptor 1 is differentially required in lymphocyte

development. Blood. 2003;102(10):3711–8.

16.   Frost JN, Wideman SK, Preston AE, Teh MR, Ai Z, Wang L, et al. Plasma iron controls

neutrophil production and function. Sci Adv. 2022 Oct 7;8(40).

17.   West AP, Brodsky IE, Rahner C, Woo DK, Erdjument-Bromage H, Tempst P, et al. TLR

signalling augments macrophage bactericidal activity through mitochondrial ROS. Nature. 2011

Apr 27;472(7344):476–80.

18.   Murakawa H, Bland CE, Willis WT, Dallman PR. Iron deficiency and neutrophil function:

different rates of correction of the depressions in oxidative burst and myeloperoxidase activity

after iron treatment. Blood. 1987 May;69(5):1464–8.

19.   Hassan TH, Badr MA, Karam NA, Zkaria M, El Saadany HF, Rahman DMA, et al. Impact of

iron deficiency anemia on the function of the immune system in children. Medicine (Baltimore).

2016 Nov;95(47):e5395.

20.   Jiang Y, Li C, Wu Q, An P, Huang L, Wang J, et al. Iron-dependent histone 3 lysine 9

demethylation controls B cell proliferation and humoral immune responses. Nat Commun.

2019;10(1):2935.

21.   Frost JN, Tan TK, Abbas M, Wideman SK, Bonadonna M, Stoffel NU, et al. Hepcidin-Mediated

Hypoferremia Disrupts Immune Responses to Vaccination and Infection. Med. 2021

Feb;2(2):164-179.e12.

22.   Wang Z, Yin W, Zhu L, Li J, Yao Y, Chen F, et al. Iron Drives T Helper Cell Pathogenicity by
Promoting RNA-Binding Protein PCBP1-Mediated Proinflammatory Cytokine Production.
Immunity. 2018 Jul;49(1):80-92.e7.

23.   Zhao M, Li M ying, Gao X fei, Jia S jie, Gao K qin, Zhou Y, et al. Downregulation of BDH2
modulates iron homeostasis and promotes DNA demethylation in CD4+ T cells of systemic
lupus erythematosus. Clin Immunol. 2018;187:113–21.

24.   Gao X, Song Y, Lu S, Hu L, Zheng M, Jia S, et al. Insufficient Iron Improves Pristane-Induced
Lupus by Promoting Treg Cell Expansion. Front Immunol. 2022;13(February):1–10.

25.   Voss K, Sewell AE, Krystofiak ES, Gibson-Corley KN, Young AC, Basham JH, et al. Elevated
transferrin receptor impairs T cell metabolism and function in systemic lupus erythematosus.
Sci Immunol. 2023 Jan 20;8(79).

26.   Preston AE, Drakesmith H, Frost JN. Adaptive immunity and vaccination – iron in the spotlight.
Immunother Adv. 2021;1(1):1–11.

27.   Stoffel NU, Uyoga MA, Mutuku FM, Frost JN, Mwasi E, Paganini D, et al. Iron Deficiency
Anemia at Time of Vaccination Predicts Decreased Vaccine Response and Iron Supplementation
at Time of Vaccination Increases Humoral Vaccine Response: A Birth Cohort Study and a
Randomized Trial Follow-Up Study in Kenyan Infants. Front Immunol. 2020;11(July).

28.   Brussow H, Sidoti J, Dirren H, Freire WB, Brüssow H, Sidoti J, et al. Effect of malnutrition in
Ecuadorian children on titers of serum antibodies to various microbial antigens. Clin Diagn Lab
Immunol. 1995;2(1):62–8.

29.   Jabara HH, Boyden SE, Chou J, Ramesh N, Massaad MJ, Benson H, et al. A missense mutation
in TFRC, encoding transferrin receptor 1, causes combined immunodeficiency. Nat Genet. 2016
Jan 7;48(1):74–8.

30.   Aljohani AH, Al-Mousa H, Arnaout R, Al-Dhekri H, Mohammed R, Alsum Z, et al. Clinical

and Immunological Characterization of Combined Immunodeficiency Due to TFRC Mutation

in Eight Patients. J Clin Immunol. 2020;40(8):1103–10.

31.   Suss G, Eichmann K, Kury E, Linke A, Langhorne J. Roles of CD4- and CD8-bearing T

689   lymphocytes in the immune response to the erythrocytic stages of Plasmodium chabaudi. Infect

690   Immun. 1988;56(12):3081–8.

691   32.   Stevenson MM, Tam MF, Belosevic M, Van Der Meide PH, Podoba JE. Role of endogenous

692   gamma interferon in host response to infection with blood-stage Plasmodium chabaudi AS.

693   Infect Immun. 1990;58(10):3225–32.

694   33.   Meding SJ, Cheng SC, Simon-Haarhaus B, Langhorne J. Role of gamma interferon during

695   infection with Plasmodium chabaudi chabaudi. Infect Immun. 1990 Nov;58(11):3671–8.

696   34.   Leisewitz AL, Rockett KA, Gumede B, Jones M, Urban B, Kwiatkowski DP. Response of the

697   splenic dendritic cell population to malaria infection. Infect Immun. 2004;72(7):4233–9.

698   35.   Sponaas AM, Do Rosario APF, Voisine C, Mastelic B, Thompson J, Koernig S, et al. Migrating

699   monocytes recruited to the spleen play an important role in control of blood stage malaria. Blood.

700   2009;114(27):5522–31.

701   36.   von der Weid T, Honarvar N, Langhorne J. Gene-targeted mice lacking B cells are unable to

702   eliminate a blood stage malaria infection. J Immunol. 1996 Apr 1;156(7):2510–6.

703   37.   Pérez-Mazliah D, Nguyen MP, Hosking C, McLaughlin S, Lewis MD, Tumwine I, et al.

704   Follicular Helper T Cells are Essential for the Elimination of Plasmodium Infection.

705   EBioMedicine. 2017;24:216–30.

706   38.   Grau GE, Taylor TE, Molyneux ME, Wirima JJ, Vassalli P, Hommel M, et al. Tumor necrosis

707   factor and disease severity in children with falciparum malaria. N Engl J Med. 1989 Jun

708   15;320(24):1586–91.

709   39.   Deroost K, Pham TT, Opdenakker G, Van den Steen PE. The immunological balance between

710   host and parasite in malaria. Vol. 40, FEMS Microbiology Reviews. 2016. 208–257 p.

40. Li C, Corraliza I, Langhorne J. A defect in interleukin-10 leads to enhanced malarial disease in
Plasmodium chabaudi chabaudi infection in mice. Infect Immun. 1999;67(9):4435–42.

41. Kobayashi F, Ishida H, Matsui T, Tsuji M. Effects of in vivo administration of anti-IL-10 or
anti-IFN-gamma monoclonal antibody on the host defense mechanism against Plasmodium
yoelii yoelii infection. J Vet Med Sci. 2000 Jun;62(6):583–7.

42. Omer FM, Riley EM. Transforming Growth Factor β Production Is Inversely Correlated with
Severity of Murine Malaria Infection. J Exp Med. 1998 Jul 1;188(1):39–48.

43. Nahrendorf W, Ivens A, Spence PJ. Inducible mechanisms of disease tolerance provide an
alternative strategy of acquired immunity to malaria. Elife. 2021;10:1–33.

44. Bundi CK, Nalwoga A, Lubyayi L, Muriuki JM, Mogire RM, Opi H, et al. Iron Deficiency Is
Associated with Reduced Levels of Plasmodium falciparum-specific Antibodies in African
Children. Clin Infect Dis. 2021;73(1):43–9.

45. Tchum SK, Sakyi SA, Adu B, Arthur F, Oppong FB, Dzabeng F, et al. Impact of IgG response
to malaria-specific antigens and immunity against malaria in pre-school children in Ghana. A
cluster randomized, placebo-controlled trial. PLoS One. 2021;16(7 July):1–13.

46. Bushell ES, Marr EJ, Milne RM, Anar B, Girling G, Schwach F, et al. An enhanced toolkit for
the generation of knockout and marker-free fluorescent Plasmodium chabaudi. Wellcome Open
Res. 2020;5:1–20.

47. Spence PJ, Jarra W, Lévy P, Nahrendorf W, Langhorne J. Mosquito transmission of the rodent
malaria parasite Plasmodium chabaudi. Malar J. 2012;11:1–7.

48. Spence PJ, Jarra W, Lévy P, Reid AJ, Chappell L, Brugat T, et al. Vector transmission regulates
immune control of Plasmodium virulence. Nature. 2013 Jun 29;498(7453):228–31.

49. Harvey PW, Bell RG, Nesheim MC. Iron deficiency protects inbred mice against infection with
Plasmodium chabaudi. Infect Immun. 1985 Dec;50(3):932–4.

50.   Koka S, Föller M, Lamprecht G, Boini KM, Lang C, Huber SM, et al. Iron deficiency influences

736   the course of malaria in Plasmodium berghei infected mice. Biochem Biophys Res Commun.

737   2007;357(3):608–14.

738   51.   Clark M, Fisher NC, Kasthuri R, Cerami Hand C. Parasite maturation and host serum iron

739   influence the labile iron pool of erythrocyte stage Plasmodium falciparum. Br J Haematol.

740   2013;161(2):262–9.

741   52.   Pollack S, Fleming J. Plasmodium falciparum takes up iron from transferrin. Br J Haematol.

742   1984 Oct;58(2):289–93.

743   53.   Petzer V, Tymoszuk P, Asshoff M, Carvalho J, Papworth J, Deantonio C, et al. A fully human

744   anti-BMP6 antibody reduces the need for erythropoietin in rodent models of the anemia of

745   chronic disease. Blood. 2020;136(9):1080–90.

746   54.   del Portillo HA, Ferrer M, Brugat T, Martin-Jaular L, Langhorne J, Lacerda MVG. The role of

747   the spleen in malaria. Cell Microbiol. 2012;14(3):343–55.

748   55.   Kurup SP, Butler NS, Harty JT. T cell-mediated immunity to malaria. Nat Rev Immunol.

749   2019;19(7):457–71.

750   56.   Arezes J, Costa M, Vieira I, Dias V, Kong XL, Fernandes R, et al. Non-transferrin-bound iron

751   (NTBI) uptake by T lymphocytes: Evidence for the selective acquisition of oligomeric ferric

752   citrate species. PLoS One. 2013;8(11).

753   57.   Wikenheiser DJ, Ghosh D, Kennedy B, Stumhofer JS. The Costimulatory Molecule ICOS

754   Regulates Host Th1 and Follicular Th Cell Differentiation in Response to Plasmodium chabaudi

755   chabaudi AS Infection . J Immunol. 2016;196(2):778–91.

756   58.   Yeo TW, Lampah DA, Gitawat R, Tjitra E, Kenangalem E, Piera K, et al. Angiopoietin-2 is

757   associated with decreased endothelial nitric oxide and poor clinical outcome in severe

758   falciparum malaria. Proc Natl Acad Sci U S A. 2008;105(44):17097–102.

59.   Hanson J, Lee SJ, Hossain AM, Anstey NM, Charunwatthana P, Maude RJ, et al. Microvascular

obstruction and endothelial activation are independently associated with the clinical

manifestations of severe falciparum malaria in adults: An observational study. BMC Med.

2015;13(1):1–11.

60.   Jain A, Kaushik R, Kaushik RM. Malarial hepatopathy: Clinical profile and association with

other malarial complications. Acta Trop. 2016 Jul;159:95–105.

61.   Seixas E, Gozzelino R, Chora Â, Ferreira A, Silva G, Larsen R, et al. Heme oxygenase-1 affords

protection against noncerebral forms of severe malaria. Proc Natl Acad Sci U S A.

2009;106(37):15837–42.

62.   Gozzelino R, Andrade BB, Larsen R, Luz NF, Vanoaica L, Seixas E, et al. Metabolic adaptation

to tissue iron overload confers tolerance to malaria. Cell Host Microbe. 2012;12(5):693–704.

63.   Brugat T, Cunningham D, Sodenkamp J, Coomes S, Wilson M, Spence PJ, et al. Sequestration

and histopathology in Plasmodium chabaudi malaria are influenced by the immune response in

an organ-specific manner. Cell Microbiol. 2014;16(5):687–700.

64.   de Menezes MN, Salles ÉM, Vieira F, Amaral EP, Zuzarte-Luís V, Cassado A, et al. IL-1α

promotes liver inflammation and necrosis during blood-stage Plasmodium chabaudi malaria. Sci

Rep. 2019;9(1):1–12.

65.   Gonçalves BP, Huang C-Y, Morrison R, Holte S, Kabyemela E, Prevots DR, et al. Parasite

Burden and Severity of Malaria in Tanzanian Children. N Engl J Med. 2014 May

8;370(19):1799–808.

66.   Gupta S, Snow RW, Donnelly CA, Marsh K, Newbold C. Immunity to non-cerebral severe

malaria is acquired after one or two infections. Nat Med. 1999;5(3):340–3.

67.   Armitage AE, Agbla SC, Betts M, Sise EA, Jallow MW, Sambou E, et al. Rapid growth is a

dominant predictor of hepcidin suppression and declining ferritin in Gambian infants.

Haematologica. 2019;104(8):1542–53.

68.   Ramos S, Carlos AR, Sundaram B, Jeney V, Ribeiro A, Gozzelino R, et al. Renal control of

disease tolerance to malaria. Proc Natl Acad Sci U S A. 2019;116(12):5681–6.

69.   Pamplona A, Ferreira A, Balla J, Jeney V, Balla G, Epiphanio S, et al. Heme oxygenase-1 and

carbon monoxide suppress the pathogenesis of experimental cerebral malaria. Nat Med.

2007;13(6):703–10.

70.   Das I, Saha K, Mukhopadhyay D, Roy S, Raychaudhuri G, Chatterjee M, et al. Impact of iron

deficiency anemia on cell-mediated and humoral immunity in children: A case control study. J

Nat Sci Biol Med. 2014;5(1):158.

71.   Wunderlich F, Al-Quraishy S, Dkhil MA. Liver-inherent immune system: its role in blood-stage

malaria. Front Microbiol. 2014 Nov 4;5.

72.   Deroost K, Alder C, Hosking C, McLaughlin S, Lin J-W, Lewis MD, et al. Tissue macrophages

and interferon-gamma signalling control blood-stage Plasmodium chabaudi infections derived

from mosquito-transmitted parasites. Curr Res Immunol. 2021;2(July):104–19.

73.   Gupta P, Lai SM, Sheng J, Renia L, Karjalainen K, Ruedl C, et al. Crucial Front Line against

Plasmodium Infection Article a Crucial Front Line against Plasmodium Infection. CellReports.

2016;16(6):1749–61.

74.   Hoffmann A, Haschka D, Valente L, Souza D, Tymoszuk P, Seifert M, et al. EBioMedicine

Baseline iron status and presence of anaemia determine the course of systemic Salmonella

infection following oral iron supplementation in mice. 2021;71.

75.   Knackstedt SL, Georgiadou A, Apel F, Abu-Abed U, Moxon CA, Cunnington AJ, et al.

Neutrophil extracellular traps drive inflammatory pathogenesis in malaria. Sci Immunol.

2019;4(40):1–18.

76.   Lin JW, Sodenkamp J, Cunningham D, Deroost K, Tshitenge TC, McLaughlin S, et al.

Signatures of malaria-associated pathology revealed by high-resolution whole-blood

transcriptomics in a rodent model of malaria. Sci Rep. 2017;7(December 2016):1–13.

77.   Zander RA, Vijay R, Pack AD, Guthmiller JJ, Graham AC, Lindner SE, et al. Th1-like

Plasmodium-Specific Memory CD4+ T Cells Support Humoral Immunity. Cell Rep.
2017;21(7):1839–52.

78. Scharer CD, Barwick BG, Guo M, Bally APR, Boss JM. Plasma cell differentiation is controlled
by multiple cell division-coupled epigenetic programs. Nat Commun. 2018;9(1).

79. Lönnberg T, Svensson V, James KR, Fernandez-Ruiz D, Sebina I, Montandon R, et al. Single-
cell RNA-seq and computational analysis using temporal mixture modeling resolves TH1/TFH
fate bifurcation in malaria. Sci Immunol. 2017;2(9):1–12.

80. Yarosz EL, Ye C, Kumar A, Black C, Choi E-K, Seo Y-A, et al. Cutting Edge: Activation-
Induced Iron Flux Controls CD4 T Cell Proliferation by Promoting Proper IL-2R Signaling and
Mitochondrial Function. J Immunol. 2020 Apr 1;204(7):1708–13.

81. Teh MR, Frost JN, Armitage AE, Drakesmith H. Analysis of Iron and Iron-Interacting Protein
Dynamics During T-Cell Activation. Front Immunol. 2021;12(August):1–18.

82. Pfeifhofer-Obermair C, Tymoszuk P, Nairz M, Schroll A, Klais G, Demetz E, et al. Regulation
of Th1 T Cell Differentiation by Iron via Upregulation of T Cell Immunoglobulin and Mucin
Containing Protein-3 (TIM-3). Front Immunol. 2021;12(May):1–11.

83. Ly A, Hansen DS. Development of B cell memory in malaria. Front Immunol. 2019;10(APR):1–
11.

84. Portugal S, Obeng-Adjei N, Moir S, Crompton PD, Pierce SK. Atypical memory B cells in
human chronic infectious diseases: An interim report. Cell Immunol. 2017;321(June):18–25.

85. Cernetich A, Garver LS, Jedlicka AE, Klein PW, Kumar N, Scott AL, et al. Involvement of
gonadal steroids and gamma interferon in sex differences in response to blood-stage malaria
infection. Infect Immun. 2006;74(6):3190–203.

86. MalariaGEN Resource Centre. Leucocyte depletion of 2.0mL of Plasmodium-infected whole
blood using MN2100ff cellulose columns [Internet]. MalariaGEN Protocols. 2016. Available
from: https://www.malariagen.net/

87.   Clark MA, Goheen MM, Spidale NA, Kasthuri RS, Fulford A, Cerami C. RBC barcoding allows

for the study of erythrocyte population dynamics and P. falciparum merozoite invasion. PLoS

One. 2014;9(7):e101041.

88.   Duarte TL, Neves J V. Measurement of Tissue Non-Heme Iron Content using a

Bathophenanthroline-Based Colorimetric Assay. J Vis Exp. 2022 Jan 31;(179).

89.   Yuan JS, Reed A, Chen F, Stewart CN. Statistical analysis of real-time PCR data. BMC

Bioinformatics. 2006;7:85.

90.   Scheuer PJ. Classification of chronic viral hepatitis: a need for reassessment. J Hepatol. 1991

Nov;13(3):372–4.

FIGURE LEGENDS

Figure 1: **Decreased cellular iron uptake increases the *P. chabaudi* pathogen burden.**

A) C57BL/6 (WT) and *Tfrc$^{Y20H/Y20H}$* (TfR) mice were infected by intravenous (i.v.) injection of 10^5

recently mosquito-transmitted *P. chabaudi* infected red blood cells (iRBC).

B-E) Parasitaemia (B), iRBC count (C), body weight change (D) and RBC count (E) measured

throughout the course of infection. Mean ± SEM, mixed-effects analysis (B, C, E) or repeated measures

two-way ANOVA (D), with Sidak's multiple comparisons test, n=7-9.

F-G) Haemoglobin measured 8 (F) and 20 (G) days after infection. Welch's t-test, n=6-9.

H-I) A mix of unlabelled WT RBC and iRBC were incubated with fluorescently labelled WT or TfR

RBC and the invasion susceptibility index (SI) was determined after completion of a new invasion

cycle. Mean, Welch's t-test, n=3.

J-K) Liver iron and serum iron levels measured 8 days after infection. Mean, Welch's t-test, n=9.

Figure 2: **Decreased cellular iron uptake impairs the splenic MNP response to *P. chabaudi*.** Splenic

immune response to *P.chabaudi* in C57BL/6 (WT) and *Tfrc$^{Y20H/Y20H}$* (TfR) mice at 8 dpi.

A) Representative picture of spleens from naïve and *P. chabaudi* infected mice.

B) Spleen index of spleens from naïve and *P. chabaudi* infected mice. Mean, Welch's t-test n=9.

C-E) Absolute numbers of CD11b$^+$ CD11c$^+$ mononuclear phagocytes (MNPs) (C), Ly6C^{hi} CD11b+ monocytes/macrophages (D) and MHCII$^+$ CD11c$^+$ dendritic cells (E) in spleens from naïve and *P. chabaudi* infected mice . Mean, Welch's t-test on untransformed (C) or log transformed data (D, E) n=9-11.

F) Representative flow cytometry plot of interferon-γ (IFNγ) production of CD11b$^+$ CD11c$^+$ MNPs.

G) Proportion of IFNγ-producing MNPs, detected by intracellular cytokine staining. Mean, Welch's t-test n=9-11.

Dotted line represents uninfected mice.

Figure 3:* Decreased cellular iron uptake disrupts the effector CD4$^+$ T cell response to *P. chabaudi. Conventional CD4$^+$ T cells (FOXP3$^-$) in the spleen of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc$^{Y20H/Y20H}$* (TfR) mice, 8 dpi.

A) Absolute number of CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test, n=9-11.

B) Proportions of naïve (CD44$^-$ CD62L$^+$), effector (CD62L$^-$ CD127$^-$) and memory (CD44$^+$ CD127$^+$) CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, two-way ANOVA with Sidak's multiple comparisons test, n=9-11.

C) Absolute number of effector CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Mann-Whitney test, n=9-11.

D-E) Proportions of CD4$^+$ T cells expressing markers of antigen experience CD44$^+$ (D) and PD-1$^+$ (E) in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=9-11

F) Proportion of proliferating (KI-67$^+$) CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=9-11

G) Proportion of T helper 1 (TBET$^+$) CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR

mice. Mean, Welch's t-test n=9-11

H) Representative flow cytometry plot of IFNγ producing CD4$^+$ T cells in the spleen of *P. chabaudi*

infected WT and TfR mice.

I) Proportion of IFNγ producing CD4$^+$ T cells, detected by intracellular cytokine staining, in the spleen

of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=10-11.

Dotted line represents uninfected mice.

***Figure 4. In vitro* T helper 1 (Th1) polarised *Tfrc$^{Y20H/Y20H}$* CD4$^+$ T cells have impaired proliferation**

and effector function, which can be rescued by iron supplementation.

A) Naïve CD4$^+$ T cells were isolated from uninfected C57BL/6 (WT) and *Tfrc$^{Y20H/Y20H}$* (TfR) mice, and

cultured for 96 h in Th1 polarising media, with varying concentrations of iron sulfate (FeSO$_4$).

B) Representative flow cytometry plot of CD4$^+$ T cell proliferation, quantified using CellTrace Violet.

C) Proportion of CD4$^+$ T cells that have divided more than two times (> 2X). Mean, two-way ANOVA,

Sidak's multiple comparisons test, n=3.

D) Representative flow cytometry plot of IFNγ producing CD4$^+$ T cell in the absence or presence of

FeSO$_4$.

E-F) Proportion of IFNγ producing CD4 T cells (E) and IFNγ production per cell (F). Mean, two-way

ANOVA, Sidak's multiple comparisons test, n=3.

Figure 5. **Decreased cellular iron uptake disrupts the germinal centre response to *P. chabaudi.***

Splenic immune response of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc$^{Y20H/Y20H}$* (TfR) mice.

A) Proportion of CD4$^+$ T cells expressing B cell co-stimulatory receptor ICOS in the spleen of of *P.*

chabaudi infected WT and TfR mice, 8 dpi. Mean, Welch's t-test, n=10-11.

B) Proportion of T follicular helper (Tfh) cells in the spleen of of *P. chabaudi* infected WT and TfR

mice, 8 dpi. Mean, Welch's t-test, n=9.

C) Proportion of Tfh cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 20 dpi. Mean,

Welch's t-test, n=6-7.

D-F) Absolute total number of splenic B cells (D) and proportion of activated (E) and antibody secreting

(F) splenic B cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 8 dpi. Mean, Welch's t-

test, n=9.

G) Proportion of germinal centre B cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 8

dpi. Mean, Welch's t-test, n=9.

H) Representative flow cytometry plot of germinal centre B cells in the spleen of of *P. chabaudi*

infected WT and TfR mice, 20 dpi.

I) Proportion of germinal centre B cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 20

dpi. Mean, Welch's t-test on log transformed data, n=6-9.

Dotted line represents uninfected mice.

Figure 6. In vitro **cultured** *Tfrc^{Y20H/Y20H}* **B cells display impaired activation, proliferation and**

differentiation, which can be rescued by iron supplementation.

A) B cells were isolated from uninfected C57BL/6 (WT) and *Tfrc^{Y20H/Y20H}* (TfR) mice and cultured for

96 h in B cell activating media, with varying concentrations of ammonium ferric citrate (AFeC).

B) Large neutral amino acid transporter-1 (LAT-1/CD98) expression on divided B cells. Mean, two-

way ANOVA, Sidak's multiple comparisons test, n=3.

C) Representative flow cytometry plot of proliferating B cells, quantified using CellTrace Violet.

D) Proportion of proliferating B cells (CTV^{low}). Mean, two-way ANOVA, Sidak's multiple

comparisons test, n=3.

E) Representative flow cytometry plots of antibody secreting (CD138^+) and class-switched (IgG^+)

divided B cells.

F-G) Proportion of antibody secreting (F) and class-switched (G) divided B cells. Mean, two-way ANOVA, Sidak's multiple comparisons test, n=3.

Figure 7. **Decreased cellular iron uptake mitigates *P. chabaudi* liver pathology.** Liver pathology of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc*$^{Y20H/Y20H}$ (TfR) mice, 8 dpi.

A-B) Serum levels of angiopoietin-2 (A) and alanine transaminase (B). Mean, Welch's t-test, n=15-16. Dotted line represents uninfected mice

C-D) Haematoxylin and eosin (C), and periodic acid–Schiff (D) staining of representative liver sections. Labels indicate central veins (CV), portal triads (PT), and areas of focal (black arrows) and bridging (white arrows) necrosis. Original magnification 40X, scale bar 100 µm.

E) Quantification of severe hepatic necrosis (score ≧ 3) as measured by histological scoring. Count, Fisher's exact test, n=10-11.

F) Number of hepatic red blood cell sequestration, rosetting and vascular occlusion events per randomly imaged high-power field (HPF). Mean, Welch's t-test, n=10-11.

G) Immunohistochemistry staining of liver leukocytes (CD45+) in representative liver sections. Original magnification 20X, scale bar 100 µm.

H) Quantification of CD45$^+$ leukocytes in liver sections identified by immunohistochemistry staining. n= 9-11

I-L) Hepatic monocytes/macrophages (I), dendritic cells (J), CD44+ CD4+ T cells (K) and CD44+ CD8+ T cells (L). Mean, Welch's t-test, n=7-12.

Figure S1: Tfrc$^{Y20H/Y20H}$ **mice have mild microcytosis and decreased iron levels at homeostasis.** Uninfected 8–12-week-old C57BL/6 (WT) and *Tfrc*$^{Y20H/Y20H}$ (TfR) mice were used for characterization.

A) Body weight at homeostasis. Mean, Welch's t-test, n=9-10.

B-D) Haemoglobin (B), mean RBC volume (C) and RBC count (D) at homeostasis. Mean, Welch's t-test, n=7.

E-F) Liver iron (E) and serum iron (F) at homeostasis. Mean, Welch's t-test, n=8-10.

Figure S2. **Hyperferremia does not increase *P. chabaudi* parasitaemia.**

A) C57BL/6 mice were infected by intravenous (i.v.) injection of 10^5 *P. chabaudi* infected red blood

cells (iRBC). A monoclonal anti-BMP-6 antibody (αBMP6) or an isotype control antibody (Iso) was

administered 2, 12 and 16 days after infection.

B) Serum iron measured 9 and 21 dpi in mice treated with anti-BMP6 or Iso. At day 9 post-infection,

serum samples, collected through tail bleeding, were pooled for each experimental group to obtain

sufficient sample for the quantification. At day 21 post-infection, mice were sacrificed, and serum

samples collected through cardiac puncture. Mean, Welch's t-test, n=6-8.

C-E) Parasitaemia (C), iRBC count (D) and relative change in body weight (E) were measured

throughout the course of infection. Mean ± SEM, two-way ANOVA with Sidak's multiple comparisons

test, n=6-8.

Figure S3: **Mononuclear phagocyte gating scheme and innate immune response to *P. chabaudi***

infection. Splenic immune response of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc*$^{Y20H/Y20H}$ (TfR)

mice, 8 dpi.

A) Gating strategy for mononuclear phagocytes (MNP), monocytes/macrophages (Mo/Mac) and

dendritic cells (DC).

B-D) Absolute number of splenic neutrophils (B), eosinophils (C) and NK cells (D) of WT and TfR

mice at 8dpi. Mean, Welch's t-test, n = 6-8.

Figure S4: **Decreased cellular iron uptake attenuates the effector CD8$^+$ T cell response to *P.***

***chabaudi*.** CD8$^+$ T cells in the spleen of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc*$^{Y20H/Y20H}$ (TfR)

mice, 8 dpi.

A) Absolute numbers of splenic CD8$^+$ T cells of *P. chabaudi* infected WT and TfR mice. Mean, Welch's

t-test, n=9-10.

B) Proportion of naïve (CD44$^-$ CD62L$^+$), effector (CD62L$^-$ CD127$^-$) and memory (CD44$^+$ CD127$^+$)

splenic CD8$^+$ T cells of *P. chabaudi* infected WT and TfR mice. Mean, two-way ANOVA with Sidak's

multiple comparisons test, n=9-11.

C) Absolute number of effector CD8$^+$ T cells of spleens from *P. chabaudi* infected WT and TfR mice.

Mean, Mann-Whitney test, n=9-11.

D-E) Proportion of splenic CD8$^+$ T cells expressing markers of antigen experience CD44$^+$ (D) and PD-

1$^+$ (E) of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=10

F) Proportion of proliferating (KI-67$^+$) splenic CD8$^+$ T cells of *P. chabaudi* infected WT and TfR mice.

Mean, Welch's t-test n=9-11

G) Proportion of IFNγ producing splenic CD8$^+$ T cells, detected by intracellular cytokine staining of *P.*

chabaudi infected WT and TfR mice. Mean, Welch's t-test n=10-11.

Dotted line represents uninfected mice.

Figure S5. **Decreased cellular iron uptake attenuates *P. chabaudi* induced liver damage.** Hepatic

response of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc$^{Y20H/Y20H}$* (TfR) mice, 8 dpi.

A-B) Liver gene expression of *Saa1* (A) and *Fga* (B) of *P. chabaudi* infected WT and TfR mice. Mean,

Welch's t-test, n=12.

C) Liver index of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test, n=10-11.

D-E) Higher magnification depiction of H&E (D) and PAS (E) stained liver sections from a

representative *P. chabaudi* infected WT mouse. The arrowheads indicate areas of confluent necrosis,

featuring lobular disarray, lympho-histiocytic inflammation, acidophil body formation, and glycogen

depletion. Original magnification 200×, scale bar 20µm.

F) Blinded scoring of lobular necro-inflammatory activity. Mann-Whitney test, n=10-11.

G) Hepatic malondialdehyde (MDA), quantified as an indirect measurement or ROS, using a

thiobarbituric acid reactive substances assay in *P. chabaudi* infected WT and TfR mice. Mean, Welch's

t-test, n=10-12.

H-J) Liver gene expression of *Tnf* (H), *Ifn* (I) and *Il1b* (J). Mean, Welch's t-test on untransformed

(H&J) or log transformed data (I), n=12.

1005

Figure 1: **Decreased cellular iron uptake increases the *P. chabaudi* pathogen burden.**

A) C57BL/6 (WT) and *Tfrc^Y20H/Y20H* (TfR) mice were infected by intravenous (i.v.) injection of 10^5 recently mosquito transmitted *P. chabaudi* infected red blood cells (iRBC).

B-E) Parasitaemia (B), iRBC count (C), body weight change (D) and RBC count (E) measured throughout the course of infection. Mean ± SEM, mixed-effects analysis (B, C, E) or repeated measures two-way ANOVA (D), with Sidak's multiple comparisons test, n=7-9.

F-G) Haemoglobin measured 8 (F) and 20 (G) days after infection. Welch's t-test, n=6-9.

H-I) A mix of unlabelled WT RBC and iRBC were incubated with fluorescently labelled WT or TfR RBC and the invasion susceptibility index (SI) was determined after completion of a new invasion cycle. Mean, Welch's t-test, n=3.

J-K) Liver iron and serum iron levels measured 8 days after infection. Mean, Welch's t-test, n=9.

Figure 2: **Decreased cellular iron uptake impairs the splenic MNP response to *P. chabaudi*.** Splenic immune response to *P.chabaudi* in C57BL/6 (WT) and *Tfrc^{Y20H/Y20H}* (TfR) mice at 8 dpi.

A) Representative picture of spleens from naïve and *P. chabaudi* infected mice.

B) Spleen index of spleens from naïve and *P. chabaudi* infected mice. Mean, Welch's t-test n=9.

C-E) Absolute numbers of CD11b$^+$ CD11c$^+$ mononuclear phagocytes (MNPs) (C), Ly6C^{hi} CD11b+ monocytes/macrophages (D) and MHCII$^+$ CD11c$^+$ dendritic cells (E) in spleens from naïve and *P. chabaudi* infected mice . Mean, Welch's t-test on untransformed (C) or log transformed data (D, E) n=9-11.

F) Representative flow cytometry plot of interferon-γ (IFNγ) production of CD11b$^+$ CD11c$^+$ MNPs.

G) Proportion of IFNγ-producing MNPs, detected by intracellular cytokine staining. Mean, Welch's t-test n=9-11.

Dotted line represents uninfected mice.

Figure 3: **Decreased cellular iron uptake disrupts the effector CD4$^+$ T cell response to *P. chabaudi*.** Conventional CD4$^+$ T cells (FOXP3$^-$) in the spleen of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc$^{Y20H/Y20H}$* (TfR) mice, 8 dpi.

A) Absolute number of CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test, n=9-11.

B) Proportions of naïve (CD44$^-$ CD62L$^+$), effector (CD62L$^-$ CD127$^-$) and memory (CD44$^+$ CD127$^+$) CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, two-way ANOVA with Sidak's multiple comparisons test, n=9-11.

C) Absolute number of effector CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Mann-Whitney test, n=9-11.

D-E) Proportions of CD4$^+$ T cells expressing markers of antigen experience CD44$^+$ (D) and PD-1$^+$ (E) in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=9-11

F) Proportion of proliferating (KI-67$^+$) CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=9-11

G) Proportion of T helper 1 (TBET$^+$) CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=9-11

H) Representative flow cytometry plot of IFNγ producing CD4$^+$ T cells in the spleen of *P. chabaudi* infected WT and TfR mice.

I) Proportion of IFNγ producing CD4$^+$ T cells, detected by intracellular cytokine staining, in the spleen of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=10-11.

Dotted line represents uninfected mice.

***Figure 4. In vitro* T helper 1 (Th1) polarised *Tfrc^{Y20H/Y20H}* CD4^+ T cells have impaired proliferation and effector function, which can be rescued by iron supplementation.**

A) Naïve CD4^+ T cells were isolated from uninfected C57BL/6 (WT) and *Tfrc^{Y20H/Y20H}* (TfR) mice, and cultured for 96 h in Th1 polarising media, with varying concentrations of iron sulfate (FeSO₄).

B) Representative flow cytometry plot of CD4^+ T cell proliferation, quantified using CellTrace Violet.

C) Proportion of CD4^+ T cells that have divided more than two times (> 2X). Mean, two-way ANOVA, Sidak's multiple comparisons test, n=3.

D) Representative flow cytometry plot of IFNγ producing CD4^+ T cell in the absence or presence of FeSO₄.

E-F) Proportion of IFNγ producing CD4 T cells (E) and IFNγ production per cell (F). Mean, two-way ANOVA, Sidak's multiple comparisons test, n=3.

***Figure 5*. Decreased cellular iron uptake disrupts the germinal centre response to *P. chabaudi*.** Splenic immune response of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc^{Y20H/Y20H}* (TfR) mice.

A) Proportion of CD4[+] T cells expressing B cell co-stimulatory receptor ICOS in the spleen of of *P. chabaudi* infected WT and TfR mice, 8 dpi. Mean, Welch's t-test, n=10-11.

B) Proportion of T follicular helper (Tfh) cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 8 dpi. Mean, Welch's t-test, n=9.

C) Proportion of Tfh cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 20 dpi. Mean, Welch's t-test, n=6-7.

D-F) Absolute total number of splenic B cells (D) and proportion of activated (E) and antibody secreting (F) splenic B cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 8 dpi. Mean, Welch's t-test, n=9.

G) Proportion of germinal centre B cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 8 dpi. Mean, Welch's t-test, n=9.

H) Representative flow cytometry plot of germinal centre B cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 20 dpi.

I) Proportion of germinal centre B cells in the spleen of of *P. chabaudi* infected WT and TfR mice, 20 dpi. Mean, Welch's t-test on log transformed data, n=6-9.

Dotted line represents uninfected mice.

Figure 6. *In vitro* cultured *Tfrc^Y20H/Y20H* B cells display impaired activation, proliferation and differentiation, which can be rescued by iron supplementation.

A) B cells were isolated from uninfected C57BL/6 (WT) and *Tfrc^Y20H/Y20H* (TfR) mice and cultured for 96 h in B cell activating media, with varying concentrations of ammonium ferric citrate (AFeC).

B) Large neutral amino acid transporter-1 (LAT-1/CD98) expression on divided B cells. Mean, two-way ANOVA, Sidak's multiple comparisons test, n=3.

C) Representative flow cytometry plot of proliferating B cells, quantified using CellTrace Violet.

D) Proportion of proliferating B cells (CTV^low). Mean, two-way ANOVA, Sidak's multiple comparisons test, n=3.

E) Representative flow cytometry plots of antibody secreting (CD138+) and class-switched (IgG+) divided B cells.

F-G) Proportion of antibody secreting (F) and class-switched (G) divided B cells. Mean, two-way ANOVA, Sidak's multiple comparisons test, n=3.

Figure 7. **Decreased cellular iron uptake mitigates *P. chabaudi* liver pathology.** Liver pathology of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc^Y20H/Y20H* (TfR) mice, 8 dpi.

A-B) Serum levels of angiopoietin-2 (A) and alanine transaminase (B). Mean, Welch's t-test, n=15-16. Dotted line represents uninfected mice

C-D) Haematoxylin and eosin (C), and periodic acid–Schiff (D) staining of representative liver sections. Labels indicate central veins (CV), portal triads (PT), and areas of focal (black arrows) and bridging (white arrows) necrosis. Original magnification 40X, scale bar 100 μm.

E) Quantification of severe hepatic necrosis (score ≧ 3) as measured by histological scoring. Count, Fisher's exact test, n=10-11.

F) Number of hepatic red blood cell sequestration, rosetting and vascular occlusion events per randomly imaged high-power field (HPF). Mean, Welch's t-test, n=10-11.

G) Immunohistochemistry staining of liver leukocytes (CD45+) in representative liver sections. Original magnification 20X, scale bar 100 μm.

H) Quantification of CD45+ leukocytes in liver sections identified by immunohistochemistry staining. n= 9-11

I-L) Hepatic monocytes/macrophages (I), dendritic cells (J), CD44+ CD4+ T cells (K) and CD44+ CD8+ T cells (L). Mean, Welch's t-test, n=7-12.

Figure S1: Tfrc^{Y20H/Y20H} mice have mild microcytosis and decreased iron levels at homeostasis. Uninfected 8–12-week-old C57BL/6 (WT) and *Tfrc^{Y20H/Y20H}* (TfR) mice were used for characterization.

A) Body weight at homeostasis. Mean, Welch's t-test, n=9-10.

B-D) Haemoglobin (B), mean RBC volume (C) and RBC count (D) at homeostasis. Mean, Welch's t-test, n=7.

E-F) Liver iron (E) and serum iron (F) at homeostasis. Mean, Welch's t-test, n=8-10.

***Figure S2.* Hyperferremia does not increase *P. chabaudi* parasitaemia**.

A) C57BL/6 mice were infected by intravenous (i.v.) injection of 10^5 *P. chabaudi* infected red blood cells (iRBC). A monoclonal anti-BMP-6 antibody (αBMP6) or an isotype control antibody (Iso) was administered 2, 12 and 16 days after infection.

B) Serum iron measured 9 and 21 dpi in mice treated with anti-BMP6 or Iso. At day 9 post-infection, serum samples, collected through tail bleeding, were pooled for each experimental group to obtain sufficient sample for the quantification. At day 21 post-infection, mice were sacrificed, and serum samples collected through cardiac puncture. Mean, Welch's t-test, n=6-8.

C-E) Parasitaemia (C), iRBC count (D) and relative change in body weight (E) were measured throughout the course of infection. Mean ± SEM, two-way ANOVA with Sidak's multiple comparisons test, n=6-8.

Figure S3: **Mononuclear phagocyte gating scheme and innate immune response to *P. chabaudi* infection.** Splenic immune response of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc*[Y20H/Y20H] (TfR) mice, 8 dpi.

A) Gating strategy for mononuclear phagocytes (MNP), monocytes/macrophages (Mo/Mac) and dendritic cells (DC).

B-D) Absolute number of splenic neutrophils (B), eosinophils (C) and NK cells (D) of WT and TfR mice at 8dpi. Mean, Welch's t-test, n = 6-8.

Figure S4: **Decreased cellular iron uptake attenuates the effector CD8+ T cell response to *P. chabaudi*.** CD8+ T cells in the spleen of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc^{Y20H/Y20H}* (TfR) mice, 8 dpi.

A) Absolute numbers of splenic CD8+ T cells of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test, n=9-10.

B) Proportion of naïve (CD44- CD62L+), effector (CD62L- CD127-) and memory (CD44+ CD127+) splenic CD8+ T cells of *P. chabaudi* infected WT and TfR mice. Mean, two-way ANOVA with Sidak's multiple comparisons test, n=9-11.

C) Absolute number of effector CD8+ T cells of spleens from *P. chabaudi* infected WT and TfR mice. Mean, Mann-Whitney test, n=9-11.

D-E) Proportion of splenic CD8+ T cells expressing markers of antigen experience CD44+ (D) and PD-1+ (E) of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=10

F) Proportion of proliferating (KI-67+) splenic CD8+ T cells of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=9-11

G) Proportion of IFNγ producing splenic CD8+ T cells, detected by intracellular cytokine staining of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test n=10-11.

Dotted line represents uninfected mice.

Figure S5. **Decreased cellular iron uptake attenuates *P. chabaudi* induced liver damage.** Hepatic response of *P. chabaudi* infected C57BL/6 (WT) and *Tfrc^{Y20H/Y20H}* (TfR) mice, 8 dpi.

A-B) Liver gene expression of *Saa1* (A) and *Fga* (B) of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test, n=12.

C) Liver index of *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test, n=10-11.

D-E) Higher magnification depiction of H&E (D) and PAS (E) stained liver sections from a representative *P. chabaudi* infected WT mouse. The arrowheads indicate areas of confluent necrosis, featuring lobular disarray, lympho-histiocytic inflammation, acidophil body formation, and glycogen depletion. Original magnification 200×, scale bar 20µm.

F) Blinded scoring of lobular necro-inflammatory activity. Mann-Whitney test, n=10-11.

G) Hepatic malondialdehyde (MDA), quantified as an indirect measurement or ROS, using a thiobarbituric acid reactive substances assay in *P. chabaudi* infected WT and TfR mice. Mean, Welch's t-test, n=10-12.

H-I) Liver gene expression of *Hmox1* (H) and *Fth1* (I). Mean, Welch's t-test on log transformed (H) or untransformed (I) data, n=12.

J-L) Liver gene expression of *Tnf* (H), *Ifn* (I) and *Il1b* (J). Mean, Welch's t-test on untransformed (H&J) or log transformed data (I), n=12.