FGF independent MEK1/2 signalling is essential for male germline development in mice

Rheannon O. Blücher[1], Rachel S. Lim[2], Ellen G. Jarred[1], Matthew E. Ritchie[2], Patrick S. Western[1*]

[1]Centre for Reproductive Health, Hudson Institute of Medical Research and Department of Molecular and Translational Science, Monash University, Clayton, Victoria 3168, Australia
[2]Epigenetics and Development Division, The Walter and Eliza Hall Institute of Medical Research, Parkville, Victoria 3052, Australia

*Corresponding author
patrick.western@hudson.org.au

Abstract

Germline development provides the founding cells for spermatogenesis and oogenesis in males and females, respectively. Disrupted germline differentiation or compromised testis development can lead to subfertility or infertility and are strongly associated with testis cancer in humans. In mice, SRY and SOX9 induce expression of a range of genes, including *Fgf9,* that promote Sertoli cell differentiation and testis development. FGF9 is also thought to promote male germline differentiation but the pathway through which it signals is unknown. As FGFs signal through Mitogen-Activated Protein Kinases (MAPKs) in other tissues, we explored whether FGF9 regulates male germline development through MAPK by inhibiting either FGF or MEK1/2 signalling in fetal testis cultures from embryonic day (E)12.5, immediately after gonadal sex determination and testis cord formation, but prior to male germline commitment. Inhibition of MEK1/2 disrupted mitotic arrest, dysregulated a broad range of male germline development genes and prevented the upregulation of key male germline markers DPPA4 and DNMT3L. In contrast, when FGF signalling was inhibited, the male germline specific transcriptional program and the expression of male germline markers DPPA4 and DNMT3L were unaffected, and germ cells entered mitotic arrest normally. While male germline development was not disrupted by FGF inhibition, some genes were commonly altered after 24h of FGF or MEK1/2 inhibition including genes involved in maintenance, germline stem cells, Nodal signalling, proliferation, and germline cancer. Together, these data demonstrate a novel and essential role for MEK1/2 signalling in male germline differentiation, but a more limited role for FGF signalling. Our data strongly indicate that additional ligands act through MEK1/2 to promote male germline differentiation and highlight a need for further mechanistic understanding of male germline development.

Introduction

Development of functional sperm and oocytes is essential for fertility and transmission of genetic and epigenetic information to offspring. A critical part of germline development involves the commitment of germ cells to male or female development in response to somatic cell signalling (Adams and McLaren, 2002). Germ cells within a developing testis commit to male development at around embryonic day (E)12.5 and enter mitotic arrest between E13.5 and E15.5 (Adams and McLaren, 2002, Western et al., 2008). They then re-enter the cell cycle and establish spermatogonial stem cells before entering meiosis during spermatogenesis in post-natal life. In contrast, commitment to female germline in the ovary is closely followed by entry into meiosis by E15.5 (Western et al., 2008, Adams and McLaren, 2002, Miles et al., 2010).

Male germline differentiation depends on development of an appropriate testicular environment. In XY mice, *Sry* (Sex region, Y-chromosome) and SOX9 (SRY box gene 9) drive pre-supporting cell commitment to Sertoli cell development at E11.5 (Koopman et al., 1991, Kent et al., 1996). Sertoli cells proliferate and form testis cords that enclose germ cells, a process supported by FGF9 (fibroblast growth factor 9), which further induces *Sox9* expression (Kim et al., 2006, Colvin et al., 2001). *Fgf9* expression peaks at E11.5, before declining to low levels by E13.5 (Bowles et al., 2010, Colvin et al., 2001, Kim et al., 2006, Yildirim et al., 2020, Jameson et al., 2012b). Loss of function mutations in *Fgf9* or its receptor, *Fgfr2*, disrupt testis development leading to ovary or ovotestis formation, reduced germ cell numbers and germline sex-reversal (DiNapoli et al., 2006, Jameson et al., 2012a, Yildirim et al., 2020, Colvin et al., 2001, Kim et al., 2006, Bagheri-Fam et al., 2008, Bagheri-Fam et al., 2017).

Evidence suggests that FGF9 also directly promotes male germline development by acting as a meiosis inhibiting factor by repressing *Stra8* (Stimulated by retinoic acid gene 8) and ensuring expression of male germline genes including *Nanos2* and *Dnmt3l* (Bowles et al., 2010, Barrios et al., 2010). However, male germ cells develop in *Wnt4/Fgf9* double null mice (Jameson et al., 2012a), indicating that *Fgf9* is dispensable in this double null background.

Furthermore, the mechanism through which FGF9 promotes male germ cell development in wild type mice remains unknown.

FGF ligands, including FGF9, are also required for the establishment of pluripotent embryonic germ cells (EGCs) from primordial germ cells (PGCs) in culture (Durcova-Hills et al., 2006). Consistent with this, FGF9 is thought to maintain expression of the pluripotency genes *Sox2* and *Oct4* in XY germ cells (Bowles et al., 2010) and exposure of isolated E11.5 or E12.5 XY germ cells to high levels of FGF9 maintained proliferation, but low levels of FGF9 promoted male germ cell differentiation (Ulu et al., 2017). *Fgf9* is downregulated by E13.5 and the potential for germ cells to make EGCs is lost from E12.5, consistent with germ cell entry into mitotic arrest and repression of pluripotency between E13.5 and E15.5 (Durcova-Hills et al., 2008, Western et al., 2010, Bowles et al., 2010).

FGF ligands bind FGF receptors and activate intracellular responses via MAP kinase signalling, including through MEK1/2 and phosphorylation of its target ERK1/2 (Dorey and Amaya, 2010). ERK1/2 phosphorylation is low at E11.5 in XX and XY germ cells, but gradually increases between E12.5 and E14.5 (Sorrenti et al., 2020), highlighting a potential role for MEK1/2 signalling during fetal germ cell development. Moreover, inhibiting ERK1/2 signalling maintains stem cells in ground state pluripotency (Ying et al., 2008), demonstrating a role for ERK1/2 in priming differentiation of pluripotent cells, an activity facilitated by the E26 transformation-specific (ETS) factors, ETV4 and ETV5 (Kalkan et al., 2019, Mulas et al., 2019).

We hypothesised that FGF9 signalling through MEK1/2 promotes male germline development soon after supporting cells commit to Sertoli cell development and testis differentiation. We inhibited FGF receptor or MEK1/2 at E12.5, after commitment to testis development at E11.5 but before germ cells irreversibly commit to male development. Our data demonstrate that germ cells are not irreversibly committed to male germline fate at E12.5 but require FGF-independent MEK1/2 signalling to successfully mediate male germline differentiation.

Results

Inhibition of FGF or MEK1/2 signalling disrupts testis development

To determine the role of FGF and potential downstream target signalling pathways required for male germ cell development, E12.5 XY gonad-mesonephros samples were cultured with vehicle control (DMSO) or a range of small molecule inhibitors of FGF receptors (FGFR1-3), MEK1/2, p38MAPK and PI3K. These included pan-FGFR (FGFR1-3) inhibitor BGJ398/Infigratinib (FGFRi; IC50 ~1.0nM), MEK1/2 inhibitor PD0325901/Mirdametinib (MEKi; IC50 0.33nM), p38MAPK inhibitor PH-797804 (p38i; IC50 26nM) or PI3K inhibitor GSK1059615 (PI3Ki, IC50 5nM; Table 1). Importantly all of these inhibitors have IC50 values in the low nM range and have been extensively validated in clinical trials demonstrating their high specificity, potency and cell tolerance (Table 1). A starting concentration of 500nM was used based on their low IC50 values and our observations that drugs with similar IC50s have observable effects in gonad cultures in the 100-1000nM range. The vehicle control, DMSO, was used at a dilution of ≥1/5,000 in all experiments, a concentration that does not affect gonad or germline development (Miles et al., 2013, Gustin et al., 2016b). Bright field and fluorescence examination of E12.5 testis-mesonephros samples cultured with control (DMSO) or drug for 72h, provided an initial readout of the impact of each drug based on germ cell organisation within testis cords, marked by germ cell specific expression of *Oct4-GFP* (Fig 1A). DMSO controls developed well-defined cords containing germ cells, but FGFRi and MEKi treatment resulted in poor testis cord structure and GFP positive germ cells scattered throughout the testis. p38i and PI3Ki treated gonads were morphologically similar to the DMSO controls, with GFP positive germ cells contained within well-defined testis cords (Fig 1A).

We previously demonstrated that FGF9 induces proliferation of XX somatic cells at rates similar to XY gonads (Gustin et al., 2016b). To confirm that FGFRi and MEKi blocked FGF9 activity, XX E12.5 gonads were cultured for 48h in media containing DMSO, 50ng/mL FGF9, 500nM FGFRi, 500nM MEKi, FGF9+FGFRi or FGF9+MEKi, with 5-ethynyl-2'-deoxyuridine (EdU) added for the final two hours of culture to mark cells in S-phase (Fig 1B-C). Using flow cytometry, mouse vasa homologue (MVH, also known as DDX4) positive germ cells were separated from somatic cells and proliferation assessed using EdU incorporation and

propidium iodide to measure DNA content (Fig S1) (Miles et al., 2013, Gustin et al., 2016b, Gustin et al., 2016a). As expected, FGF9 substantially increased the proliferation of XX gonadal somatic cells compared to XX controls ($P<0.05$, $P<0.0001$; Fig 1B-C). Critically, both FGFRi and MEKi completely neutralised FGF9, with somatic cell proliferation decreased to XX control levels in FGF9 + FGFRi or FGF9 + MEKi treated XX gonads (Fig 1B-C). In contrast, neither p38i nor PI3Ki counteracted FGF9 induced somatic cell proliferation in XX gonads, indicating that neither p38MAPK nor PI3K regulate primary pathways through which FGF drives somatic cell proliferation in developing gonads (Fig S2A).

To assess the effects of each treatment on testis development, we used EdU and flow cytometry to measure SOX9 and MVH positive Sertoli and germ cell proliferation in E12.5 gonad-mesonephric samples cultured for 72h with each inhibitor (Fig 1D-E, S1). As expected, Sertoli cells were highly proliferative in DMSO controls (Fig 1D). However, FGFR or MEK1/2 inhibition reduced Sertoli cell proliferation compared to XY DMSO controls ($P<0.05$ or $P<0.001$, respectively, Fig 1D). As expected, germ cell proliferation was very low in DMSO controls (Fig 1E), confirming that the germ cells had entered mitotic arrest, a key milestone in male germline differentiation (Western et al., 2008). In contrast, germ cell proliferation was substantially higher in MEKi treated gonads ($P<0.05$), demonstrating that germ cell mitotic arrest was disrupted (Fig 1E). However, the percentage of germ cells incorporating EdU in FGFRi-treated testes was similar to DMSO controls, demonstrating that the germ cells had entered mitotic arrest (Fig 1E). This outcome was confirmed using another potent FGFR inhibitor (AZD4547; Table 1), which also resulted in reduced Sertoli cell proliferation but did not disrupt germ cell mitotic arrest (Fig S2B-C). Neither p38i (PH-797804) nor PI3Ki (GSK1059615) affected Sertoli cell proliferation or germ cell mitotic arrest (Fig 1D-E), an outcome confirmed using independent, p38 and PI3K inhibitors, ralimetanib dimesylate (LY2228820) and PF-04691502 (Table 1; Fig S2D-E).

FGF and MEK1/2 signalling is required for Sertoli cell proliferation, but only MEK1/2 is required for germ cell mitotic arrest

To determine the dose response to MEK1/2 and FGFR inhibition, E12.5 XY gonad-mesonephros samples were cultured with MEKi or FGFRi at 0 (DMSO diluted ≥1/5,000), 125, 250, 500 and 1000nM and assessed using flow cytometry (Fig S3). Compared to DMSO

control, Sertoli cell proliferation was reduced by all doses of FGFRi ($P<0.0001$), with the effect plateauing at ≥125nM (Fig 2Ai) and from ≥250nM MEKi ($P<0.0001$, Fig 2Aii). Consistent with our initial observations (Fig 1E), germ cell mitotic arrest remained unaffected by FGFRi, even with doses of 1000nM (~750-1000x the drug IC50; Fig 2Bi) or 2500nM (Fig S2F-G). Similarly, while SU5402 , an FGFR inhibitor used in a previous study (Bowles et al., 2010), reduced Sertoli cell proliferation to a similar extent as FGFRi, it did not affect germ cell mitotic arrest (Table 1; S2F-G). In contrast, MEKi potently disrupted germ cell mitotic arrest even when used at 125nM, with increasingly high proportions of EdU positive proliferative germ cells as MEKi concentration increased ($P<0.0001$, Fig 2Bii).

As FGF9 expression peaks at E11.5 (Jameson et al., 2012b), a potential explanation for the inability of FGFRi to disrupt germ cell mitotic arrest may have been that FGF9 was inhibited too late in E12.5 cultures. XY E11.5 gonad-mesonephros samples were therefore cultured for 96h with DMSO or 500nM FGFRi or MEKi (Fig 2C-D). As in E12.5 cultures, FGFRi or MEKi significantly reduced Sertoli cell proliferation ($P<0.01$ and $P<0.001$ respectively, Fig 2C). Germ cell proliferation was moderately higher in FGFRi treated gonads compared to DMSO control (p<0.05, Fig 2D). However, MEKi resulted in much more robust maintenance of germ cell proliferation $(P<0.0001$, Fig 2D), demonstrating a potent ability of MEKi, but a substantially lower ability of FGFRi, to disrupt mitotic arrest.

To determine if MEK1/2 inhibition affected Sertoli cell proliferation and germ cell mitotic arrest after E13.5, XY E13.5 gonad-mesonephros samples were cultured for 48h with DMSO or 500nM MEKi. MEKi treatment from E13.5 significantly reduced Sertoli ($P<0.01$, Fig 2E) and increased germ cell proliferation ($P<0.01$, Fig 2F). However, the effect of MEKi on germ cell mitotic arrest was diminished compared to E12.5 and E11.5 (E13.5: EdU 6%, E12.5: 15%, E11.5: 25%, Fig 2Bii,D,F), demonstrating that the ability of MEKi to disrupt mitotic arrest decreased in E11.5, E12.5 and E13.5 gonads, despite similar impacts on Sertoli cell proliferation at all ages.

FGF and MEK1/2 signalling is required to ensure all germ cells localise inside testis cords
As gonad wholemount images indicated testis cords were disrupted by FGFRi or MEKi (Fig 1A), we used immunofluorescence (IF) to investigate MVH positive germ cells relative to SOX9-expressing Sertoli cells, SMA-expressing peritubular myoid cells or laminin, which delineate

testis cords. In DMSO controls the majority of Sertoli cells were organised in a single layer at the outer edge of the testis cords, with germ cells very rarely found outside the cords (Fig 3). In FGFRi and MEKi treated samples some Sertoli cells localised to the outer edge of testis cords, but gaps were evident between the Sertoli cells, and many Sertoli cells remained dispersed throughout the interior of the testis cords (Fig 3A). Furthermore, germ cells were occasionally present in the gaps between Sertoli cells at the testis cord basement membrane (Fig 3A) and were mis-localised outside testis cords in FGFRi treated cultures, although this was more common in MEKi treated gonads (Fig 3B).

MEK1/2 signalling is required for male germline differentiation

As MEKi prevented germ cell mitotic arrest, the expression of male germline differentiation markers was assessed using IF and flow cytometry in E12.5 XY gonad-mesonephros samples cultured for 72h with DMSO, 125nM or 500nM FGFRi or MEKi. DPPA4 is expressed in XX and XY germ cells at E12.5, but is upregulated in XY germ cells and repressed in XX germ cells as they differentiate (Gustin et al., 2016b). As expected, DPPA4 was not detected in germ cells of XX gonads but was detected in germ cells of XY E12.5 gonads cultured for 72h with DMSO, and fluorescence appeared more intense in XY E12.5+72h than in E12.5 XY germ cells (Fig 4A, S4A). While DPPA4 germ cell levels were similar in XY E12.5+72h DMSO and FGFRi cultures, DPPA4 intensity appeared lower in MEKi treated samples and comparable to E12.5 XY germ cells (Fig 4A, S4A). Flow cytometry revealed that the relative DPPA4 germ cell intensity was 2x higher in XY E12.5+72h DMSO and FGFRi cultures than in E12.5 XY germ cells ($P<0.0001$, Fig 4B), but comparable to E12.5 XY germ cells in MEKi-treated gonads (Fig 4B).

As the germ cell proliferation and DPPA4 levels indicated that male germ cells failed to properly differentiate in MEKi-treated XY gonads, we examined two additional male germline markers, DNMT3L and PIWIL2. IF and flow cytometry revealed that the majority of germ cells were DNMT3L positive in the DMSO control and FGFRi treated gonads (Fig 4C-E, S4B). In contrast, very few germ cells were DNMT3L positive in MEKi-treated gonads, and the DNMT3L staining intensity was significantly lower than in the DMSO- or FGFRi-treated samples ($P<0.01$, $P<0.0001$; Fig 4C-E, S4B). Similarly, PIWIL2 was expressed at similar levels in germ cells of DMSO- and FGFRi-treated samples but was variable in MEKi-treated samples, with some germ

cells staining strongly for PIWIL2 and others negative (Fig 4F, S4C). This was not possible to confirm using flow cytometry because a reliable PIWIL2 flow assay could not be developed.

MEK1/2 inhibition increased STRA8, but failed to promote widespread meiosis in XY germ cells

Since MEKi inhibited mitotic arrest and male germline differentiation, the expression of female germline markers was investigated to determine if FGFRi or MEKi induced female development in XY germ cells. As expected, the pre-meiotic marker STRA8 was detected in the germ cells of XX E12.5+72h DMSO-treated gonads but was not detected in XY DMSO controls (Fig 5A, S5A). While some germ cells appeared very weakly positive for STRA8 in FGFRi treated samples, STRA8 positive germ cells were commonly found in MEKi-treated gonads (Fig 5A, S5A), particularly in germ cells close to the mesonephric-gonadal boundary (Fig S5A). However, while STRA8 staining was localised in the germ cell nucleus in XX controls, it was detected in the germ cell cytoplasm and nucleus in MEKi treatments (Fig 5A, S5A). Flow cytometry demonstrated that 74% of germ cells were STRA8 positive in XX DMSO controls, but only 5% and 7% were STRA8 positive in XY control and FGFRi treated samples (Fig 5B). The proportion of STRA8 positive germ cells in MEKi-treated gonads was 38%, significantly higher than XY controls ($P<0.0001$), but lower than XX controls ($P<0.0001$, Fig 5B).

To determine whether germ cells in FGFRi- or MEKi-treated XY gonads had entered meiosis, gonad sections were triple stained using antibodies specific for SYCP3, phospho-γH2AX (p-γH2AX) and MVH (Fig 5C, S5B). SYCP3 and p-γH2AX were detected in most germ cells in 72h XX DMSO controls but not in XY DMSO- or FGFRi-treated gonads (Fig 5C, S5B). A small number of germ cells were positive for SYCP3 in MEKi-treated gonads and a subset also stained for p-γH2AX (Fig 5C, S5B). In addition, rare cells positive for p-γH2AX were detected in XY control, FGFRi and MEKi treated gonads (Fig S5B), however, most did not express MVH and were likely to be apoptotic somatic cells in which p-γH2AX also marks double strand DNA breaks.

To test the possibility that meiotic entry of germ cells in FGFRi- or MEKi-treated gonads was delayed, we cultured E12.5 XX and XY gonad-mesonephros for 96h with DMSO or 500nM FGFRi or MEKi. IF staining revealed that most XX DMSO germ cells were p-γH2AX positive, indicating they had entered meiosis (Fig 5D, S5C). p-γH2AX positive germ cells were rarely

detected in XY DMSO- or FGFRi-treated gonads but were more common in MEKi-treated samples (Fig 5D, S5C). Quantification using flow cytometry revealed that 88% of germ cells were p-γH2AX positive in XX DMSO samples while only 2%, 4% and 19% were p-γH2AX positive in XY control, FGFRi and MEKi treated samples, respectively (Fig 5E).

Cell cycle analysis of germ cells from the same gonads using EdU (S-phase) to quantify DNA synthesis and propidium iodide to measure DNA content demonstrated that the majority of germ cells in E12.5+96h cultures were in G2/M in XX controls, but were in G0/G1 in XY DMSO- and FGFRi treated gonads (Fig 5F). Significantly more germ cells were in G2/M in MEKi than in XY control or FGFRi cultures ($P<0.0001$), but remained less than in XX controls ($P<0.0001$, Fig 5F). Therefore, while MEKi treatment resulted in a significantly greater percentage of p-γH2AX expressing germ cells that were in G2/M, this proportion was substantially lower than in XX controls indicating that meiosis was not properly induced within the normal temporal window following MEK inhibition.

The majority of transcriptional divergence occurs after E12.5 in XY and XX germ cells

We next used RNA sequencing to gain greater insight into genome-wide transcriptional changes in FACS isolated *Oct4*-GFP positive germ cells of E12.5 XX and XY gonads (Time 0 controls) and gonads cultured for 24 and 72h with DMSO, FGFRi and MEKi (Fig 6A). Differential expression analysis identified 183 and 234 genes that were expressed higher in XY and XX germ cells at E12.5, respectively (time 0; FDR<0.05; absolute Fold-Change (FC) $\geq$ 1.5, absolute Log-FC $\geq$ 0.585; Fig 6B, Table S1.1-1.2). Included in the differentially expressed genes (DEGs) that were higher in E12.5 XY germ cells were a range of Nodal signalling associated genes, including *Nodal, Tdgf1 (Cripto), Lefty1*, *Lefty2, Pitx2* and *Otx2*, which are known to be high in XY germ cells at this time point (Spiller et al., 2012, Miles et al., 2013, Wu et al., 2015, Mayère et al., 2021) (Table S1.1). E12.5 XX germ cells expressed higher levels of BMP target genes, including *Msx1, Msx2, Id1, Id2, Id3, Stra8* and *Gata2,* consistent with observations that BMP2 promotes female germline development (Nagaoka et al., 2020), (Mayère et al., 2021) (Table S1.2). However, despite these sex-specific transcriptional differences in E12.5 germ cells, our functional data strongly indicated that these differences were insufficient to ensure male germline commitment as MEK1/2 inhibition at E12.5 substantially disrupted male germline differentiation (Figs 1-4).

To identify male and female transcriptional changes that occurred as a normal part of sex-specific germline differentiation, XX and XY germ cells of DMSO control samples were compared after 24 and 72h of culture. Multidimensional scaling (MDS) revealed that although time 0 XY and XX E12.5 samples were different, they diverged substantially more after 24 and 72h of gonad culture (Fig 6C). Differential gene expression analysis revealed that 211 and 3437 genes were increased, and 217 and 2160 genes were decreased in XY germ cells after 24 and 72h compared to E12.5 (time 0) XY germ cells, respectively (Fig 6B, Table S1.3-1.4). Similarly, 340 and 3214 genes were increased, and 157 and 2592 genes were decreased in XX germ cells compared to E12.5 XX germ cells after 24 and 72h (Fig 6B, Table S1.5-1.6). Together, these data indicated that while male and female germline differentiation progressed in the first 24h, the greatest transcriptional change occurred between 24 and 72h.

To identify genes specifically associated with male and female germline differentiation we compared XY with XX germ cells from DMSO controls. This revealed 589 and 3548 genes higher in XY than XX germ cells after 24 and 72h, (XY germline genes; Fig 6B, Table S1.7-1.8), including male germline genes *Dnmt3l, Dppa4, Nanos2, Piwil1, Piwil2, Piwil4, Tdrd1, Tdrd5* and *Tdrd9*. By comparison, 978 and 2898 genes were higher in XX than XY germ cells after 24 and 72h (XX germline genes; Fig 6B, Table S1.9-1.10), including female germline and meiosis markers *Atr, Atm, Chek1, Ccnb3, H2ax, Irx3, Irx5, Msx1, Id1, Id2, Id3, Sycp2, Sycp3* and *Stra8*, confirming that E12.5 germ cells diverged in the expected sex-specific manner over time.

FGF and MEK1/2 commonly regulate a subset of genes involved in germ cell tumours, stem cell biology and proliferation

To determine the initial impacts of MEKi and FGFRi on germline development we compared outcomes in XY germ cells cultured with DMSO or 500nM FGFRi or MEKi for 24h. MDS indicated that germ cells from XX E12.5+24h controls were transcriptionally distinct from all XY groups. XY E12.5+24h DMSO, FGFRi or MEKi cultures were also transcriptionally distinct (Fig 6D) and comparison of XY germ cells from MEKi and FGFRi cultures with DMSO controls revealed 43 and 25 DEGs, respectively (FDR<0.05, |FC|≥1.5; Fig 6E, Table S2.1-2.2). These included 23 and 19 genes that were lower than expected (i.e. for which transcription was not properly activated) and 20 and six genes that were higher than expected (i.e. not properly

repressed, or unexpectedly derepressed) in FGFRi or MEKi treatments. Nine genes were commonly dysregulated by MEKi and FGFRi implying that they depend on FGF signalling via MEK1/2 (Fig 6Fi, Table S2.3). Importantly, the direction of change (up- or downregulated) was concordant (roast test *P*=0.00025) for each of the nine common DEGs, with eight expressed lower, and one higher than expected (Fig 6Fii). The simplest interpretation is that the DEGs that were lower than expected were FGF responsive genes that depended on MEK1/2 signalling for their upregulation, and the DEGs that were higher than expected depended on FGF-MEK1/2 for their repression.

Further examination of the genes that were lower than expected revealed six FGFRi-MEKi DEGs associated with stem cell differentiation, cell self-renewal, cancer, cell proliferation and survival and germ cell tumours, including *Etv1* and *Etv4* (Akagi et al., 2015, Gashaw et al., 2005, Oh et al., 2012, Cai et al., 2007), *Pdk1* (Anwar et al., 2021), *Ret* (Miles et al., 2012, Naughton et al., 2006), *Tdgf1* (*Cripto*) (Spiller et al., 2016) and *Calcoco2* (Cui et al., 2021). Few genes commonly associated with sex-specific germline differentiation were represented in the FGFRi and MEKi DEG lists after 24h of culture. However, although unaffected by FGFRi, *Nanos2,* which regulates male germline differentiation (Suzuki and Saga, 2008), was 9.12-fold lower than control in 24h MEKi samples (Table S2.2), consistent with a requirement for MEK1/2 signalling for its upregulation.

Male specific germline differentiation depends on MEK1/2, but FGF signalling is dispensable
To determine the impacts of FGFRi and MEKi on later stages of male germline differentiation we analysed samples after 72h of culture. Surprisingly, germ cells from 72h XY DMSO and FGFRi cultures were transcriptionally similar (Fig 6G). Although 43 genes were differentially expressed after 24h of FGFR inhibition (FDR<0.05, |FC|≥1.5; Fig 6Ei; Table S2.1), only one DEG was identified after 72h, and was higher than expected (FDR<0.05, |FC|≥1.5; Fig 6Hi; Table S2.4). In contrast, MEKi samples were transcriptionally distinct from XX and XY DMSO controls at 72h (Fig 6G), with 817 genes lower and 586 higher than controls (FDR<0.05, |FC|≥1.5; Fig 6Hii, Table S2.5). Of the 817 genes that were lower, 686 were male germline genes (i.e. normally upregulated in the germ cells of XY vs XX E12.5+72h DMSO cultures defined in Fig 6B) and were therefore defined as MEK1/2-dependent male germline genes (Fig 6Ii, Table S2.6). Genes lower than expected included key male germline markers *Dppa4*, *Nanos2*, *Dnd1*,

Nxf2 and *Brdt* (Maldonado-Saldivia et al., 2007, Suzuki and Saga, 2008). Interestingly, the transcriptional levels of *Dppa4* and the germline teratoma gene *Dnd1* (Youngren et al., 2005, Cook et al., 2009) was similar to E12.5 XY germ cells (Fig 6J), consistent with our observations of DPPA4 protein expression in MEKi treated samples (Fig 4A-B). In addition, DNA methylation and piRNA associated genes, including *Dnmt3a*, *Dnmt3l*, *Tdrd1*, *Tdrd5*, *Tdrd9*, *Spocd1*, *Piwil1*, *Piwil4* and *Henmt1* (Sakai et al., 2004, Aravin et al., 2008, Shoji et al., 2009), were not properly upregulated in MEKi treated samples, with levels remaining substantially lower than in XY DMSO-treated controls (Fig 6J). Consistent with persistent germ cell proliferation after MEK1/2 inhibition (Fig 2Cii), genes regulating the G1-S transition, DNA synthesis and germ cell proliferation, including *Mki67, Pcna*, *Ccne1*, *Ccne2* and *Ccnd1*, remained high after 72h of MEKi (Fig 6J).

MEK1/2 signalling inhibition did not effectively promote female germline differentiation

Of the 586 genes that were higher than control in XY germ cells after MEKi (i.e. genes derepressed or not properly repressed via MEK1/2 signalling), 382 were normally expressed at higher levels in differentiating XX than XY germ cells (Fig 6B), suggesting feminisation of the germline (Fig 6Iii; Table S2.7). However, of these 382 genes, 218 were expressed at similar levels in E12.5 germ cells, consistent with MEKi blocking germline differentiation rather than increasing feminisation (Fig S6). These included genes marking germ cell proliferation such as *Ccne1, Pcna* and *Ki67* (Miles et al., 2010, Western et al., 2008) (Fig 6J). As MEKi blocked mitotic arrest but most germ cells did not enter meiosis (Fig 5E, F), the simplest explanation is that these genes remained high due to continued germ cell proliferation, rather than female differentiation.

MEKi upregulated *Stra8*, *Meioc*, *Irx3* and *Irx5* (Soh et al., 2017, Bowles et al., 2006, Miles et al., 2010, Fu et al., 2018) and some genes that normally increase during female germline development (genes in red box in S6). However, genes marking meiosis initiation or progression including *Msx1, Sycp2, Sycp3*, *Ccnb3* (Cyclin B3)*, Atr and Atm* (Miles et al., 2010, Le Bouffant et al., 2011) were not increased by MEKi (Fig 6J). Moreover, *Ccnb1 and Msx1,* which are repressed in differentiating XX germ cells (Miles et al., 2010), remained high in MEKi cultures (Fig 6J). Other well-known meiosis markers including *Check1* and *H2ax* were transcribed higher in both MEKi and E12.5 XY controls than DMSO controls, so were not

informative (Fig 6J). Together, despite higher levels of some female germline markers including *Stra8 and Meioc,* the low expression of many meiosis markers such as *Atr*, *Atm* and *Msx1* (Fig 6J) was consistent with the limited entry of germ cells into meiosis indicated by analyses of cell cycle and p-γH2AX following MEK inhibition (Fig 5D-F).

MEK1/2 inhibition retained germ cells in a relatively undifferentiated state

MEK1/2 inhibition maintained XY germline proliferation and genes including *Dppa4, Dnd1, Ccne1, Ccne2, Ki67, Pcna, H2ax, Chek1* and *Ccnb1* remained at similar levels in E12.5+72h MEKi samples as in E12.5 XY germ cells, indicating that MEKi blocked male germline differentiation. Consistent with this, the *Nodal* regulatory genes *Lefty1* and *Lefty2* were maintained at higher levels, although *Nodal* and *Tdgf1* were not affected (Fig 6J). While *Nodal* and *Tdgf1* were not affected by MEKi at 72h, the Nodal signalling target *Pitx2* was also upregulated by MEKi (Fig 6J).

Key regulators of pluripotency *Sox2* and *Nanog* are expressed in E12.5 XX and XY germ cells but repressed after E13.5 (Western et al., 2010, Maldonado-Saldivia et al., 2007, Bullejos and Koopman, 2004). MEKi maintained *Nanog*, but not *Sox2* transcription in germ cells. *Oct4* remained unchanged, but as *Oct4* transcription is maintained between E12.5 and E15.5 in XY germ cells (Western et al., 2010), this was not informative. As expected, IF analysis revealed that E13.5 XY germ cells expressed OCT4 and SOX2 protein, but germ cells in XX DMSO+72h cultures were negative (Fig S7B-C). Although OCT4 and SOX2 staining intensity was significantly lower than in E13.5 XY gonads, the proportion of germ cells remaining positive and the staining intensity of these proteins was similar in XY gonads treated with DMSO, FGFRi or MEKi for 72h (Fig S7B-E). In addition, 72h MEKi altered genes associated with germ cell or other tumours (Kanetsky et al., 2011, Taguchi et al., 2021), or spermatogonial stem cell function (Raverot et al., 2005, Song et al., 2016) including higher *Dmrt1* and *Sox3* and lower *Rhox10* expression (Fig 6J). Moreover, rather than being down-regulated as normally occurs in differentiating male germ cells, *Etv3* was maintained at levels higher than or similar to E12.5 germ cells (Fig 6J).

Discussion

We have identified a novel and essential role for MEK1/2 signalling in male germ cell differentiation. Our data demonstrate that MEK1/2 signalling is required for XY germ cells to enter mitotic arrest, upregulate a wide range of genes that mark male germline differentiation and repress a range of female germline genes. However, although FGF signalling has been implicated in directly promoting male germline differentiation, our data indicates that FGF signalling is dispensable for germ cell mitotic arrest and male germline differentiation. Instead, our data support a role for FGF-MEK1/2 signalling in regulating genes associated with Nodal signalling and stem cell characteristics in E12.5 XY germ cells, consistent with FGF and Nodal inducing pluripotency in XY germ cells, germ cell tumours and EGCs and the well-defined role for FGF in the derivation of pluripotent EG cells (Durcova-Hills et al., 2006, Western, 2009, Bowles et al., 2010, Spiller et al., 2012, Vallier et al., 2009, Mesnard et al., 2006).

MEK1/2 inhibition from E12.5 disrupted germ cell mitotic arrest, maintained expression of genes associated with germ cell proliferation and prevented appropriate upregulation of 686 male germline development genes. A prominent germline signature of MEKi was the deficient upregulation of *de novo* DNA methylation genes and a range of genes involved in piRNA biology that are required for silencing repetitive elements. This implies that the DNA methylation pathway either responds directly to MEK1/2 signalling or is not properly activated due to poor male germ cell differentiation when MEK1/2 is inhibited. Either way, germline DNA methylation is likely to remain low if MEK1/2 signalling is compromised in the germline, potentially allowing de-repression of repetitive elements.

In addition, several genes associated with germline tumours, including members of the Nodal signalling pathway and genes associated with germline tumours in human GWAS and other studies, including *DMRT1*, were expressed in germ cells of MEK1/2 inhibited samples. While we did not observe increased OCT4 or SOX2 protein levels, *Nanog* transcription remained high, germ cell proliferation was sustained, and germline differentiation was inhibited in the absence of MEK1/2 signalling. Combined with low DNA methylation, these factors could

render germ cells more susceptible to germline tumours, a possibility that requires further investigation.

A previous report indicated that FGF9 is required for transiently maintaining pluripotency in male germ cells and for male germ cell differentiation (Bowles et al., 2010). In this study, we observed a requirement for FGF signalling to regulate 43 genes and MEK1/2 to regulate 24 genes in E12.5 germ cells after 24h of culture. Nine genes were commonly dysregulated by FGFRi and MEKi, suggesting that these genes are regulated by FGF-MEK-ERK signalling. Consistent with a role for FGF-MEK-ERK in regulating stem cell characteristics in E12.5 germ cells, six of these genes have been associated with germ cell tumours, stem cells, cell self-renewal, cancer, cell proliferation and cell survival.

Although inhibition of FGFR for 24h from E12.5 resulted in dysregulation of 43 genes, only one gene was dysregulated after 72h of FGFRi treatment, while 1403 genes were dysregulated by MEKi. Moreover, despite inhibition of FGF signalling at E12.5, germ cells progressed normally into mitotic arrest. Together, these data indicate that although some genes associated with stem cell characteristics were disrupted by FGFR inhibition at E12.5, the germ cells were able to enter the male fate normally. Our observations support a role for FGF signalling in priming stemness in male germ cells but indicate that FGF signalling is dispensable for male germline differentiation. This is consistent with previous observations that FGF9 can be used for deriving EG cells (Durcova-Hills et al., 2006). While it remains possible that a low level of residual FGF activity could explain the ability of E12.5 germ cells to enter male differentiation in our study, our conclusion is consistent with male germline development in mice lacking *Wnt4* and *Fgf9* (Jameson et al., 2012a). Together these findings demonstrate that germ cells require MEK-ERK signalling to properly establish male identity in an FGF9-FGFR independent manner (Fig 7).

While our data strongly indicate that FGF signalling is dispensable for male germline differentiation it remains possible that FGFR inhibition was too late to disrupt male germline development. However, this seems unlikely as FGF-MEK signalling is very rapid and MEKi profoundly disrupted male germline differentiation at E12.5 but FGFRi did not. A second possibility is that FGFR inhibition was insufficient and residual FGF activity promoted male

germline development. However, FGFRi reduced Sertoli cell proliferation to a similar extent in E11.5, E12.5 and E13.5 XY gonads and completely abrogated FGF9 induced somatic cell proliferation in XX gonads. Moreover, two independent, highly potent and specific FGFR inhibitors provided similar outcomes. Indeed, even when E12.5 gonads were treated with 2500nM FGFRi, a dose ~2000-2500x the drug IC50 or a dose of 5000nM of SU5402, which was previously used to inhibit FGF signalling (Bowles et al., 2010), we still did not observe any impact on germ cell mitotic arrest and the impact on Sertoli cell proliferation was not augmented. Given these data, the simplest explanation is that direct FGF9 signalling through germ cells is dispensable for male germline differentiation and the modest impact on germ cell mitotic arrest achieved using FGFRi at E11.5 was likely to be due to disruption of testis development rather than a direct effect on germ cells.

Deletion of FGF9 causes male to female gonadal sex reversal and germ cells enter meiosis (DiNapoli et al., 2006), the latter which may indirectly result from somatic cell sex reversal or be a direct effect of FGF9 loss. While FGF9 treatment of isolated E11.5 germ cells demonstrated that FGF9 can reduce retinoic acid (RA) induced *Stra8* expression, exposure of germ cells to FGF9 in the absence of RA did not significantly affect *Stra8* (Bowles et al., 2010). Moreover, FGFR inhibition in cultured E11.5 XX or XY gonads did not affect *Stra8* expression, but inhibition CYP26B1 led to robust *Stra8* transcription, presumably due to greater RA availability (Bowles et al., 2010). This effect was enhanced in XY gonads but not XX gonads by inhibition of FGF9. While this suggests that FGF9 and CYP26B1 act in concert to inhibit *Stra8* transcription, the effect of FGF9 in this context may be due to an indirect effect on somatic cells and *Cyp26b1* transcription in E11.5 gonads rather than a direct effect on germ cells. In our study *Stra8* expression was not affected by FGFR inhibition from E12.5, perhaps because FGFRi did not reduce *Cyp26b1* transcription and CYP26B1 mediated RA degradation. Combined, it seems plausible that the primary role of FGF9 in *Stra8* repression is mediated by ensuring an appropriate testicular environment and consequent *Cyp26b1* expression and *Stra8* repression, rather than a direct effect of FGF9 on germ cells.

Together our study reveals a novel, essential role of MEK1/2 signalling in fetal male germline development, particularly in the promotion of germ cell mitotic arrest, the expression of an appropriate male germline transcriptional program and the activation of *de novo* DNA

methylation. While FGF9 may be involved in priming male germline development, it appears to be dispensable for commitment of germ cells to male differentiation. Moreover, our data suggest that unknown ligand(s) activate MEK1/2 signalling and promote germline differentiation through an indirect mechanism, but further work is required to determine how this may occur.

Materials and Methods

Mouse strains, animal housing, breeding and ethics

Mice were housed at Monash Medical Centre Animal Facility with controlled temperature and humidity, a 12 hours (h) light-dark cycle and food and water available *ad libitum*. Mouse embryos were obtained from inbred 129T2svJ *Oct4*-eGFP males crossed with Swiss females. Females were checked daily for vaginal plugs, with detection of a plug noted as E0.5. Animal work was undertaken in accordance with Monash Medical Centre Animal Facility Animal Ethics Committee approval.

Organ culture

E11.5 gonads were genetically sexed using PCR as previously described (McFarlane et al., 2013). E12.5 and E13.5 embryos were sexed visually by the presence (male) or absence (female) of testis cords in the gonad. Gonad-mesonephros samples were cultured on 30mm Millicell Biopore membranes with 0.4µM pores (Merck Millipore; PICM03050) in 6-well plates, with each well containing 1400µL media (15mM Hepes, 0.1mM non-essential amino acids, 1mg/mL N-acetylcysteine, 1X penicillin/streptomycin, 55nM beta-mercaptoethanol and 10% fetal calf serum in DMEM/F12 with Glutamax). PBS was placed in between the wells to maintain humidity. Gonads were cultured in media containing DMSO (vehicle control, was used at a dilution of ≥1/5,000, as appropriate for the concentration of each drug dilution), 125, 250, 500 1000nM of BGJ398 (FGFRi; SelleckChem, HY-13241), PD0325901 (MEKi; SelleckChem, S1036), GSK1059615 (PI3Ki; MedChem Express, HY-12036), PH797804 (p38i; MedChem Express, HY-10403). Additional inhibitors for FGFR (AZD4547; SelleckChem, S2801 and SU5402; MedChem Express, HY-10407), p38 (Ralimetinib dimesylate; MedChem Express, HY-13241) and PI3K (PF-04691502; MedChem Express, HY-15177) signalling were also used to ensure consistency of outcomes with each of the primary inhibitors used. All inhibitors

were selected based on their high-specificity, potency and advancement in clinical trials (Table 1). Gonad-mesonephric complexes were randomly allocated to each culture treatment condition and cultured for 24, 48, 72 or 96h in 5% CO_2 at 37°C, with media refreshed daily. To facilitate analysis of cell proliferation, 5-ethynyl-2'-deoxyuridine (EdU) was added to each sample for the final two hours of culture at a final concentration of 20μM. After culture, gonads were photographed under brightfield and fluorescence optics, then processed for flow cytometry, IF or fluorescent activated cell sorting (FACS). For gonad only flow cytometric experiments and all experiments involving FACS purification of germ and somatic cells, gonads were dissected away from mesonephros at the end of the culture period.

Table 1: Summary of treatments and doses used in gonad cultures

Treatment	Dose	IC50 and target	Clinical trial phase, clinicaltrials.gov identifier	Supplier, catalogue number	References
DMSO (vehicle control)	Equal to DMSO in drug treatments (≥1/5000)	NA	NA	Thermo-Fisher, D12345	
BGJ398/ Infigratinib (FGFRi)	125, 250, 500, 1000 and 2500 nM	IC50 – FGFR1-3: 0.9-1.4 nM, FGFR4: 60 nM	Phase II, NCT02150967	SelleckChem, S2183	(Javle et al., 2018)
PD0325901/ Mirdametinib (MEKi)	125, 250, 500 and 1000 nM	IC50 – MEK: 0.33 nM	Phase II, NCT03962543	SelleckChem or MedChem Express, S1036 or HY-10254	(Weiss et al., 2021)
PH-797804 (p38i)	500 nM	IC50 – p38α: 26 nM; p38β 102 nM	Phase II, NCT00559910	MedChem Express, HY-10403	(MacNee et al., 2013)
GSK1059615 (PI3Ki)	500 nM	IC50 – PI3Kα/ β/δ/γ: 0.4-5 nM, mTOR: 12 nM	Phase I, NCT00695448	MedChem Express, HY-12036	
AZD4547 (second FGFR inhibitor)	125, 250 and 500 nM	IC50 – FGFR1-3: 0.2-2.5 nM; FGFR4: 165 nM; VEGFR2: 24 nM	Phase II, NCT02465060	SelleckChem, S2801	(Chae et al., 2020)
SU5402 (third FGFR inhibitor)	5000nM	IC50 VEGFR: 20nM; FGFR1: 30nM	N/A	MedChem Express, HY-10407	(Sun et al., 1999)
Ralimetinib dimesylate/ LY2228820 (second p38 inhibitor)	500 nM	IC50 – p38α: 5.3 nM; p38β: 5.3 nM	Phase I, NCT01663857	MedChem Express, HY-13241	(Patnaik et al., 2016)
PF-04691502 (second PI3K inhibitor)	500 nM	IC50 – PI3Kα/ β/δ/γ: 1.6-2.1 nM; mTOR: 16 nM	Phase II, NCT01420081	MedChem Express, HY-15177	(del Campo et al., 2016)

Flow cytometry

Gonad collection, dissociation, fixation, staining and flow cytometry were performed essentially as described previously (Wakeling et al., 2013), using eGFP or antibodies specific for MVH, DPPA4, SOX9, DNMT3L, STRA8 or H2AX. Mesonephros or limb samples were used as germ cell negative controls to set gates for eGFP, MVH or DPPA4 and E12.5 female gonads were used as a negative control to set gates for SOX9 and DNMT3L. Cultured male gonads were used as a negative control to set gates for STRA8 and H2AX. Representative gating and negative control gates can be found in S1. A rabbit IgG antibody was used as a negative control for determining staining intensities using specific rabbit antibodies. Primary antibodies used are listed in Table 2. Secondary antibodies used include Alexa Fluor Donkey anti Goat 488 (Thermo-Fisher, A11055), Alexa Fluor Donkey anti Goat 647 (Thermo-Fisher, A31573), Biotin Donkey anti Rabbit (Thermo-Fisher, A16027) and Biotin Donkey anti Goat (Thermo-Fisher, A16009). Cell cycle analysis was performed as previously described (Wakeling et al., 2013), with germ cells or Sertoli cells identified by their expression of MVH or SOX9, respectively. Cells were stained with 20µg/mL of propidium iodide, enabling quantitation of cellular DNA content. Proliferation was measured by gating EdU positive cells to identify cells in S-phase, while cells in G0/G1 or G2/M were respectively identified by DNA contents estimated as 2n or 4n in the EdU negative population compared to DMSO controls. All flow cytometry was performed on a BD FACS Canto II analyser (BD, Biosciences) and data analysed with FlowJo (v10.7.2) and Graphpad Prism (v9.2.0). Relative antibody staining intensities were calculated by removing background staining in negative control samples and normalising levels to the XY DMSO control set at 1.0. Data represents 3-16 biological replicates and statistical significance was determined using one-way ANOVA with Tukey's multiple comparisons or unpaired two-tailed t-test, where appropriate. If variance was unequal, a non-parametric Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons was used. P values <0.05 were considered significant.

Tissue fixation, embedding, immunofluorescence and image analysis

Gonads were fixed in 4% paraformaldehyde (PFA) in PBS overnight at 4°C. Samples were washed three times in PBS before 70% ethanol processing, and embedded in paraffin. 4µm sections were cut in a compound series (typically four slides prepared per gonad sample with

around 6 sections collected/slide), mounted on Superfrost Plus slides and dried at least overnight before antibody incubation. Antigen retrieval was conducted using Dako Citrate buffer (pH 6.0) for 30 minutes at 98°C in a PT Link rinse station (Dako). Tissue sections were blocked in PBTx containing 5% BSA (Merck, A9647) and 10% donkey serum (Sigma; D9663) for 1 hour at RT. Primary antibody (Table 2) diluted in PBTx containing 1% BSA was left to incubate overnight at 4°C or for 2h room temperature (RT). Slides were incubated for 1h at RT in the dark in secondary antibody (Alexa Fluor, Thermo-Fisher, Donkey anti Goat 488 A11055 or Donkey anti Rabbit 488 A21206; Donkey anti Rabbit 647 A31573 or Donkey anti Mouse 647 A31571; Donkey anti Mouse 594 A21203; Donkey anti Goat 594 A11058 or Donkey anti Mouse 555 A31570) diluted at 1/300 in PBTx containing 1% BSA. Slides were mounted in ProLong Gold containing DAPI (Thermo-Fisher, P36931). Confocal images were taken using a Nikon C1 Confocal microscope, with images taken using either 10x lens or 40x oil immersion lens, or slides were scanned using a VS120 Virtual Slide microscope (Olympus), collecting single optical sections of the whole area for each section on the slide. Image analysis was conducted with QuPath (v0.3.0) (Bankhead et al., 2017). Data from at least three representative tissue sections from four biological replicates were averaged. Relative antibody staining intensity was calculated by normalising levels to the XY DMSO control set at 1.0. Data represents four biological replicates and statistical significance was determined with Graphpad Prism (v9.2.0) using one-way ANOVA with Tukey's multiple comparisons. If variance was unequal, a non-parametric Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons was used. *P* values <0.05 were considered significant.

Table 2: Antibodies for flow cytometry (F) and immunofluorescence (IF)

Protein	Source and catalogue #	Species	Dilution
Mouse Vasa Homologue (MVH)	R&D Systems, AF2030	Goat	F: 1/100 IF: 1/400
DPPA4	R&D Systems, AF3730	Goat	F: 1/100 IF: 1/400
OCT4	Santa Cruz, sc8628	Goat	IF: 1/400
MVH	Cell Signalling Technology, 8761S	Rabbit	IF: 1/400
SOX9	Sigma-Aldrich, AB5535	Rabbit	F: 1/200 IF: 1/1000
DNMT3L	Cell Signalling Technology, 13451S	Rabbit	F: 1/100 IF: 1/100
STRA8	Abcam, ab49405	Rabbit	F: 1/500 IF: 1/400
Phospho-γH2AX	Cell Signalling Technology, 9718S	Rabbit	F: 1/100 IF: 1/800

IgG	Cell Signalling Technology, 3900S	Rabbit	Equal to primary antibody used
Laminin	R&D Systems, L9393	Rabbit	IF: 1/200
PIWIL2	Cell Signalling Technology, 5940S	Rabbit	IF: 1/200
SYCP3	Abcam, ab97672	Mouse	IF: 1/200
Smooth Muscle Actin (SMA)	Sigma-Aldrich, A2547	Mouse	IF: 1/1000
NR2F2	R&D Systems, PP-H7147-00	Mouse	IF: 1/400
SOX2	Cell Signalling Technologies, 4900S	Mouse	IF: 1/100

Fluorescent activated cell sorting (FACS) of germ and somatic cells

After organ culture, the mesonephros was dissected from the gonads. Six to 15 gonads were pooled for each sample and were dissociated in trypsin. Germ and somatic cell populations were isolated as previously described (Western et al., 2008, Wakeling et al., 2013) using the BD FACSAria™ Fusion cell sorter. GFP positive germ and GFP negative somatic cell populations were isolated from E12.5 XX and XY control gonads cultured for 24 or 72h in DMSO or XY E12.5 gonads cultured for 24 or 72h in DMSO, FGFRi or MEKi (n≥4 for each treatment/group). Germ cells were defined as GFP positive, with dead propidium iodide positive germ cells excluded.

RNA-seq library construction and sequencing

RNA was isolated from $3\text{-}45 \times 10^4$ FACS-sorted germ cells using Macherey-Nagel NucleoSpin® RNA XS extraction kit (Scientifix, 740902.50) following manufacturer instructions. RNA quantity and RNA integrity (RIN) were assessed using Qubit and Bioanalyzer (Agilent Technologies). Libraries were prepared with 30ng of RNA from germ cells with RIN values greater than 7. The library was constructed as previously described by the MHTP Medical Genomics Facility (Grubman et al., 2021). Briefly, during initial poly(A) tail priming, an 8bp sample index along with a 10bp unique molecular identifier (UMI) was added. Samples were pooled and amplified with a template switching oligonucleotide. The Illumina P5 and P7 were added by PCR and Nextera transposase, respectively. The forward read (R1) utilises a custom primer to sequence into the index while the reverse read (R2) uses a standard R2 primer to sequence the cDNA in the sense direction. I7 indexes were added to enable parsing of sample sets. Sequencing was performed on NextSeq2000 (Illumina) following Illumina protocol 1000000109376 v3.

Data pre-processing

FASTQ files were demultiplexed and mapped using scPipe (Tian et al., 2018) and Rsubread (Liao et al., 2019) (Bioconductor) packages in R studio. Briefly, FASTQ files were reformatted with *sc_trim_barcode* to incorporate barcode information from read 1 into read 2 into the header. Reads were aligned to a reference mouse genome (GENECODE GRCm39) using Rsubread. Reads were assigned to annotated exons with sc_*exon_mapping*, data were demultiplexed using *sc_demultiplex* and a gene count matrix was generated with UMI deduplication using *sc_gene_counting*. Gene count matrices from each set were combined into a DGEList object for analysis.

Downstream analysis of RNA-sequencing data

Differential gene expression was assessed using the Limma (Ritchie et al., 2015), Glimma (Su et al., 2017) and edgeR (Robinson et al., 2009) Bioconductor packages following a previously established workflow (Law et al., 2016). Briefly, gene count data was loaded into R studio and genes were annotated with any duplicates removed. Raw counts were transformed into counts per million (CPM). Lowly expressed genes were removed using the *filterByExpr* function in edgeR, and gene expression distributions were normalised using trimmed mean of M-values (TMM) method (Robinson and Oshlack, 2010). Multidimensional scaling (MDS) plots were generated to visualise sample clustering. Since there were many samples present, groups were subdivided into smaller groups to visualise clustering between different culture treatments and culture periods more clearly. Heteroscedasticity was removed from the count data with *voomWithQualityWeights* (Liu et al., 2015). Linear modelling and empirical Bayes moderations was used to test for differential expression. As there were many differentially expressed genes (DEGs) identified following empirical Bayes moderations, to identify DEGs of biological relevance, a $\log_2$fold-change cut-off was set at greater/less than 0.585 (equivalent to a FC of 1.5) using *treat* (McCarthy and Smyth, 2009). Genes were considered differentially expressed if they met the logFC cut-off and had a false discovery rate (FDR) adjusted p-value less than 0.05. DEGs identified in XY E12.5+24h FGFRi vs XY E12.5+24h DMSO (24h FGFRi DEGs) and XY E12.5+24h MEKi vs XY E12.5+24h DMSO (24h MEKi DEGs) were compared, with genes commonly affected by FGFRi and MEKi assessed using Roast (Wu et al., 2010). Heatmaps were generated using ClustVis (Metsalu and Vilo, 2015).

Declarations

Competing interests:
The authors declare that they have no competing interests that affect this work

Funding:
This work was supported by grants and research funding from:
Australian Research Council Discovery Project DP210102342 (PSW, MER)
Hudson Institute of Medical Research (PSW)
Victorian Government's Operational Infrastructure Support Program
Monash University Postgraduate Student Award to ROB

Author contributions:
Conceived and designed experiments: ROB, PSW
Performed experiments and/or analysed data: ROB, EGJ, PSW
Bioinformatic analyses: ROB, RSL
Writing - original draft: ROB, PSW
Writing - review & editing: ROB, PSW, MER
All authors critically read and approved the manuscript.

Ethics approval and consent to participate:
All animal work was undertaken in accordance with Monash University Animal Ethics
Committee (AEC) approvals.

Data and materials availability: With exception of the RNA sequencing data generated in
this study, all data are available in the main text and the Supplementary Materials. The RNA
sequencing data have been deposited in the Gene Expression Omnibus (GEO) and are
publicly available via accession number GSE221453.

Acknowledgments:
We thank Associate Professor Dagmar Wilhelm and Associate Professor Craig Smith for critical
comments on the manuscript, Monash Animal Research Platform staff for assistance with
mouse care, Monash Histology Platform for assistance with slide scanning, the Monash Micro
Imaging Facility and MHTP Medical Genomics Facilities for assistance and technical advice
and Peter Hickey for advice regarding bioinformatic analysis.

References:

ADAMS, I. R. & MCLAREN, A. 2002. Sexually dimorphic development of mouse primordial
    germ cells: switching from oogenesis to spermatogenesis. *Development,* 129**,** 1155-
    1164.
AKAGI, T., KUURE, S., URANISHI, K., KOIDE, H., COSTANTINI, F. & YOKOTA, T. 2015. ETS-
    related transcription factors ETV4 and ETV5 are involved in proliferation and
    induction of differentiation-associated genes in embryonic stem (ES) cells. *The
    Journal of Biological Chemistry,* 290**,** 22460-22473.

ANWAR, S., SHAMSI, A., MOHAMMAD, T., ISLAM, A. & HASSAN, M. I. 2021. Targeting pyruvate dehydrogenase kinase signaling in the development of effective cancer therapy. *Biochimica et Biophysica Acta - Reviews on Cancer,* 1876**,** 188568.

ARAVIN, A. A., SACHIDANANDAM, R., BOURC'HIS, D., SCHAEFER, C., PEZIC, D., TOTH, K. F., BESTOR, T. & HANNON, G. J. 2008. A piRNA Pathway Primed by Individual Transposons Is Linked to De Novo DNA Methylation in Mice. *Molecular Cell,* 31**,** 785-799.

BAGHERI-FAM, S., BIRD, A. D., ZHAO, L., RYAN, J. M., YONG, M., WILHELM, D., KOOPMAN, P., ESWARAKUMAR, V. P. & HARLEY, V. R. 2017. Testis Determination Requires a Specific FGFR2 Isoform to Repress FOXL2. *Endocrinology,* 158**,** 3832-3843.

BAGHERI-FAM, S., SIM, H., BERNARD, P., JAYAKODY, I., TAKETO, M. M., SCHERER, G. & HARLEY, V. R. 2008. Loss of Fgfr2 leads to partial XY sex reversal. *Developmental Biology,* 314**,** 71-83.

BANKHEAD, P., LOUGHREY, M. B., FERNÁNDEZ, J. A., DOMBROWSKI, Y., MCART, D. G., DUNNE, P. D., MCQUAID, S., GRAY, R. T., MURRAY, L. J. & COLEMAN, H. G. 2017. QuPath: Open source software for digital pathology image analysis. *Scientific Reports,* 7**,** 1-7.

BARRIOS, F., FILIPPONI, D., PELLEGRINI, M., PARONETTO, M. P., DI SIENA, S., GEREMIA, R., ROSSI, P., DE FELICI, M., JANNINI, E. A. & DOLCI, S. 2010. Opposing effects of retinoic acid and FGF9 on Nanos2 expression and meiotic entry of mouse germ cells. *Journal of Cell Science,* 123**,** 871-880.

BOWLES, J., FENG, C. W., SPILLER, C., DAVIDSON, T. L., JACKSON, A. & KOOPMAN, P. 2010. FGF9 Suppresses Meiosis and Promotes Male Germ Cell Fate in Mice. *Developmental Cell,* 19**,** 440-449.

BOWLES, J., KNIGHT, D., SMITH, C., WILHELM, D., RICHMAN, J., MAMIYA, S., YASHIRO, K., CHAWENGSAKSOPHAK, K., WILSON, M. J., ROSSANT, J., HAMADA, H. & KOOPMAN, P. 2006. Retinoid signaling determines germ cell fate in mice. *Science,* 312**,** 596-600.

BULLEJOS, M. & KOOPMAN, P. 2004. Germ cells enter meiosis in a rostro-caudal wave during development of the mouse ovary. *Molecular Reproduction and Development,* 68**,** 422-428.

CAI, C., HSIEH, C.-L., OMWANCHA, J., ZHENG, Z., CHEN, S.-Y., BAERT, J.-L. & SHEMSHEDINI, L. 2007. ETV1 Is a Novel Androgen Receptor-Regulated Gene that Mediates Prostate Cancer Cell Invasion. *Molecular Endocrinology,* 21**,** 1835-1846.

CHAE, Y. K., HONG, F., VAKLAVAS, C., CHENG, H. H., HAMMERMAN, P., MITCHELL, E. P., ZWIEBEL, J. A., IVY, S. P., GRAY, R. J., LI, S., MCSHANE, L. M., RUBINSTEIN, L. V., PATTON, D., WILLIAMS, P. M., HAMILTON, S. R., MANSFIELD, A., CONLEY, B. A., ARTEAGA, C. L., HARRIS, L. N., O'DWYER, P. J., CHEN, A. P. & FLAHERTY, K. T. 2020. Phase II Study of AZD4547 in Patients With Tumors Harboring Aberrations in the FGFR Pathway: Results From the NCI-MATCH Trial (EAY131) Subprotocol W. *The Journal of Clinical Oncology,* 38**,** 2407-2417.

COLVIN, J. S., GREEN, R. P., SCHMAHL, J., CAPEL, B. & ORNITZ, D. M. 2001. Male-to-Female Sex Reversal in Mice Lacking Fibroblast Growth Factor 9. *Cell,* 104**,** 875-889.

COOK, M. S., COVENEY, D., BATCHVAROV, I., NADEAU, J. H. & CAPEL, B. 2009. BAX-mediated cell death affects early germ cell loss and incidence of testicular teratomas in Dnd1(Ter/Ter) mice. *Developmental Biology,* 328**,** 377-83.

CUI, F., WANG, S., TAN, J., TANG, H., FAN, Y. & HU, J. 2021. Calcium-binding and coiled-coil domain 2 promotes the proliferation and suppresses apoptosis of prostate cancer cells. *Experimental and Therapeutic Medicine,* 21**,** 405-405.

DEL CAMPO, J. M., BIRRER, M., DAVIS, C., FUJIWARA, K., GOLLERKERI, A., GORE, M., HOUK, B., LAU, S., POVEDA, A., GONZÁLEZ-MARTÍN, A., MULLER, C., MURO, K., PIERCE, K., SUZUKI, M., VERMETTE, J. & OZA, A. 2016. A randomized phase II non-comparative study of PF-04691502 and gedatolisib (PF-05212384) in patients with recurrent endometrial cancer. *Gynecologic Oncology,* 142**,** 62-69.

DINAPOLI, L., BATCHVAROV, J. & CAPEL, B. 2006. FGF9 promotes survival of germ cells in the fetal testis. *Development,* 133**,** 1519-1527.

DOREY, K. & AMAYA, E. 2010. FGF signalling: diverse roles during early vertebrate embryogenesis. *Development,* 137**,** 3731-42.

DURCOVA-HILLS, G., ADAMS, I. R., BARTON, S. C., SURANI, M. A. & MCLAREN, A. 2006. The Role of Exogenous Fibroblast Growth Factor-2 on the Reprogramming of Primordial Germ Cells into Pluripotent Stem Cells. *Stem Cells,* 24**,** 1441-1449.

DURCOVA-HILLS, G., TANG, F., DOODY, G., TOOZE, R. & SURANI, M. A. 2008. Reprogramming Primordial Germ Cells into Pluripotent Stem Cells. *PLOS One,* 3**,** e3531.

FU, A., OBERHOLTZER, S. M., BAGHERI-FAM, S., RASTETTER, R. H., HOLDREITH, C., CACERES, V. L., JOHN, S. V., SHAW, S. A., KRENTZ, K. J., ZHANG, X., HUI, C.-C., WILHELM, D. & JORGENSEN, J. S. 2018. Dynamic expression patterns of Irx3 and Irx5 during germline nest breakdown and primordial follicle formation promote follicle survival in mouse ovaries. *PLOS Genetics,* 14**,** e1007488.

GASHAW, I., GRÜMMER, R., KLEIN-HITPASS, L., DUSHAJ, O., BERGMANN, M., BREHM, R., GROBHOLZ, R., KLIESCH, S., NEUVIANS, T. P. & SCHMID, K. W. 2005. Gene signatures of testicular seminoma with emphasis on expression of ets variant gene 4. *Cellular and Molecular Life Sciences CMLS,* 62**,** 2359-2368.

GRUBMAN, A., CHOO, X. Y., CHEW, G., OUYANG, J. F., SUN, G., CROFT, N. P., ROSSELLO, F. J., SIMMONS, R., BUCKBERRY, S., LANDIN, D. V., PFLUEGER, J., VANDEKOLK, T. H., ABAY, Z., ZHOU, Y., LIU, X., CHEN, J., LARCOMBE, M., HAYNES, J. M., MCLEAN, C., WILLIAMS, S., CHAI, S. Y., WILSON, T., LISTER, R., POUTON, C. W., PURCELL, A. W., RACKHAM, O. J. L., PETRETTO, E. & POLO, J. M. 2021. Transcriptional signature in microglia associated with Aβ plaque phagocytosis. *Nature Communications,* 12**,** 3015.

GUSTIN, S. E., HOGG, K., STRINGER, J. M., RASTETTER, R. H., PELOSI, E., MILES, D. C., SINCLAIR, A. H., WILHELM, D. & WESTERN, P. S. 2016a. WNT/β-catenin and p27/FOXL2 differentially regulate supporting cell proliferation in the developing ovary. *Developmental Biology,* 412**,** 250-260.

GUSTIN, S. E., STRINGER, J. M., HOGG, K., SINCLAIR, A. H. & WESTERN, P. S. 2016b. FGF9, activin and TGFbeta promote testicular characteristics in an XX gonad organ culture model. *Reproduction,* 152**,** 529-543.

JAMESON, S. A., LIN, Y. T. & CAPEL, B. 2012a. Testis development requires the repression of Wnt4 by Fgf signaling. *Developmental Biology,* 370**,** 24-32.

JAMESON, S. A., NATARAJAN, A., COOL, J., DEFALCO, T., MAATOUK, D. M., MORK, L., MUNGER, S. C. & CAPEL, B. 2012b. Temporal Transcriptional Profiling of Somatic and Germ Cells Reveals Biased Lineage Priming of Sexual Fate in the Fetal Mouse Gonad. *PLOS Genetics,* 8**,** e1002575.

JAVLE, M., LOWERY, M., SHROFF, R. T., WEISS, K. H., SPRINGFELD, C., BORAD, M. J., RAMANATHAN, R. K., GOYAL, L., SADEGHI, S. & MACARULLA, T. 2018. Phase II study of BGJ398 in patients with FGFR-altered advanced cholangiocarcinoma. *Journal of Clinical Oncology,* 36**,** 276.

KALKAN, T., BORNELÖV, S., MULAS, C., DIAMANTI, E., LOHOFF, T., RALSER, M., MIDDELKAMP, S., LOMBARD, P., NICHOLS, J. & SMITH, A. 2019. Complementary Activity of ETV5, RBPJ, and TCF3 Drives Formative Transition from Naive Pluripotency. *Cell Stem Cell,* 24**,** 785-801.e7.

KANETSKY, P. A., MITRA, N., VARDHANABHUTI, S., VAUGHN, D. J., LI, M., CIOSEK, S. L., LETRERO, R., D'ANDREA, K., VADDI, M., DOODY, D. R., WEAVER, J., CHEN, C., STARR, J. R., HÅKONARSON, H., RADER, D. J., GODWIN, A. K., REILLY, M. P., SCHWARTZ, S. M. & NATHANSON, K. L. 2011. A second independent locus within DMRT1 is associated with testicular germ cell tumor susceptibility. *Human Molecular Genetics,* 20**,** 3109-3117.

KENT, J., WHEATLEY, S. C., ANDREWS, J. E., SINCLAIR, A. H. & KOOPMAN, P. 1996. A male-specific role for SOX9 in vertebrate sex determination. *Development,* 122**,** 2813-2822.

KIM, Y., KOBAYASHI, A., SEKIDO, R., DINAPOLI, L., BRENNAN, J., CHABOISSIER, M. C., POULAT, F., BEHRINGER, R. R., LOVELL-BADGE, R. & CAPEL, B. 2006. Fgf9 and Wnt4 act as antagonistic signals to regulate mammalian sex determination. *PLoS Biology,* 4**,** e187.

KOOPMAN, P., GUBBAY, J., VIVIAN, N., GOODFELLOW, P. & LOVELL-BADGE, R. 1991. Male development of chromosomally female mice transgenic for Sry. *Nature,* 351**,** 117-121.

LAW, C. W., ALHAMDOOSH, M., SU, S., DONG, X., TIAN, L., SMYTH, G. K. & RITCHIE, M. E. 2016. RNA-seq analysis is easy as 1-2-3 with limma, Glimma and edgeR. *F1000Research,* 5.

LE BOUFFANT, R., SOUQUET, B., DUVAL, N., DUQUENNE, C., HERVÉ, R., FRYDMAN, N., ROBERT, B., HABERT, R. & LIVERA, G. 2011. Msx1 and Msx2 promote meiosis initiation. *Development,* 138**,** 5393-5402.

LIAO, Y., SMYTH, G. K. & SHI, W. 2019. The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Research,* 47**,** e47-e47.

LIU, R., HOLIK, A. Z., SU, S., JANSZ, N., CHEN, K., LEONG, H. S., BLEWITT, M. E., ASSELIN-LABAT, M.-L., SMYTH, G. K. & RITCHIE, M. E. 2015. Why weight? Modelling sample and observational level variability improves power in RNA-seq analyses. *Nucleic Acids Research,* 43**,** e97-e97.

MACNEE, W., ALLAN, R. J., JONES, I., DE SALVO, M. C. & TAN, L. F. 2013. Efficacy and safety of the oral p38 inhibitor PH-797804 in chronic obstructive pulmonary disease: a randomised clinical trial. *Thorax,* 68**,** 738-45.

MALDONADO-SALDIVIA, J., VAN DEN BERGEN, J., KROUSKOS, M., GILCHRIST, M., LEE, C., LI, R., SINCLAIR, A. H., SURANI, M. A. & WESTERN, P. S. 2007. Dppa2 and Dppa4 are closely linked SAP motif genes restricted to pluripotent cells and the germ line. *Stem Cells,* 25**,** 19-28.

MAYÈRE, C., NEIRIJNCK, Y., SARAROLS, P., RANDS, C. M., STÉVANT, I., KÜHNE, F., CHASSOT, A. A., CHABOISSIER, M. C., DERMITZAKIS, E. T. & NEF, S. 2021. Single-cell

transcriptomics reveal temporal dynamics of critical regulators of germ cell fate during mouse sex determination. *The FASEB Journal,* 35**,** e21452.

MCCARTHY, D. J. & SMYTH, G. K. 2009. Testing significance relative to a fold-change threshold is a TREAT. *Bioinformatics,* 25**,** 765-771.

MCFARLANE, L., TRUONG, V., PALMER, J. S. & WILHELM, D. 2013. Novel PCR Assay for Determining the Genetic Sex of Mice. *Sexual Development,* 7**,** 207-211.

MESNARD, D., GUZMAN-AYALA, M. & CONSTAM, D. B. 2006. Nodal specifies embryonic visceral endoderm and sustains pluripotent cells in the epiblast before overt axial patterning. *Development,* 133**,** 2497-2505.

METSALU, T. & VILO, J. 2015. ClustVis: a web tool for visualizing clustering of multivariate data using Principal Component Analysis and heatmap. *Nucleic Acids Research,* 43**,** W566-W570.

MILES, D. C., VAN DEN BERGEN, J. A., SINCLAIR, A. H. & WESTERN, P. 2010. Regulation of the female mouse germ cell cycle during entry into meiosis. *Cell Cycle,* 9**,** 408-418.

MILES, D. C., VAN DEN BERGEN, J. A., WAKELING, S. I., ANDERSON, R. B., SINCLAIR, A. H. & WESTERN, P. S. 2012. The proto-oncogene Ret is required for male foetal germ cell survival. *Developmental Biology,* 365**,** 101-109.

MILES, D. C., WAKELING, S. I., STRINGER, J. M., VAN DEN BERGEN, J. A., WILHELM, D., SINCLAIR, A. H. & WESTERN, P. S. 2013. Signaling through the TGF beta-activin receptors ALK4/5/7 regulates testis formation and male germ cell development. *PLOS One,* 8**,** e54606.

MULAS, C., KALKAN, T., VON MEYENN, F., LEITCH, H. G., NICHOLS, J. & SMITH, A. 2019. Defined conditions for propagation and manipulation of mouse embryonic stem cells. *Development,* 146.

NAGAOKA, S. I., NAKAKI, F., MIYAUCHI, H., NOSAKA, Y., OHTA, H., YABUTA, Y., KURIMOTO, K., HAYASHI, K., NAKAMURA, T., YAMAMOTO, T. & SAITOU, M. 2020. ZGLP1 is a determinant for the oogenic fate in mice. *Science,* 367**,** eaaw4115.

NAUGHTON, C. K., JAIN, S., STRICKLAND, A. M., GUPTA, A. & MILBRANDT, J. 2006. Glial Cell-Line Derived Neurotrophic Factor-Mediated RET Signaling Regulates Spermatogonial Stem Cell Fate1. *Biology of Reproduction,* 74**,** 314-321.

OH, S., SHIN, S. & JANKNECHT, R. 2012. ETV1, 4 and 5: An oncogenic subfamily of ETS transcription factors. *Biochimica et Biophysica Acta - Reviews on Cancer,* 1826**,** 1-12.

PATNAIK, A., HALUSKA, P., TOLCHER, A. W., ERLICHMAN, C., PAPADOPOULOS, K. P., LENSING, J. L., BEERAM, M., MOLINA, J. R., RASCO, D. W., ARCOS, R. R., KELLY, C. S., WIJAYAWARDANA, S. R., ZHANG, X., STANCATO, L. F., BELL, R., SHI, P., KULANTHAIVEL, P., PITOU, C., MULLE, L. B., FARRINGTON, D. L., CHAN, E. M. & GOETZ, M. P. 2016. A First-in-Human Phase I Study of the Oral p38 MAPK Inhibitor, Ralimetinib (LY2228820 Dimesylate), in Patients with Advanced Cancer. *Clinical Cancer Research,* 22**,** 1095-1102.

RAVEROT, G., WEISS, J., PARK, S. Y., HURLEY, L. & JAMESON, J. L. 2005. Sox3 expression in undifferentiated spermatogonia is required for the progression of spermatogenesis. *Developmental Biology,* 283**,** 215-225.

RITCHIE, M. E., PHIPSON, B., WU, D., HU, Y., LAW, C. W., SHI, W. & SMYTH, G. K. 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research,* 43**,** e47-e47.

ROBINSON, M. D., MCCARTHY, D. J. & SMYTH, G. K. 2009. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics,* 26**,** 139-140.

ROBINSON, M. D. & OSHLACK, A. 2010. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biology,* 11**,** R25.

SAKAI, Y., SUETAKE, I., SHINOZAKI, F., YAMASHINA, S. & TAJIMA, S. 2004. Co-expression of de novo DNA methyltransferases Dnmt3a2 and Dnmt3L in gonocytes of mouse embryos. *Gene Expression Patterns,* 5**,** 231-237.

SHOJI, M., TANAKA, T., HOSOKAWA, M., REUTER, M., STARK, A., KATO, Y., KONDOH, G., OKAWA, K., CHUJO, T., SUZUKI, T., HATA, K., MARTIN, S. L., NOCE, T., KURAMOCHI-MIYAGAWA, S., NAKANO, T., SASAKI, H., PILLAI, R. S., NAKATSUJI, N. & CHUMA, S. 2009. The TDRD9-MIWI2 Complex Is Essential for piRNA-Mediated Retrotransposon Silencing in the Mouse Male Germline. *Developmental Cell,* 17**,** 775-787.

SOH, Y. Q. S., MIKEDIS, M. M., KOJIMA, M., GODFREY, A. K., DE ROOIJ, D. G. & PAGE, D. C. 2017. Meioc maintains an extended meiotic prophase I in mice. *PLOS Genetics,* 13**,** e1006704.

SONG, H.-W., BETTEGOWDA, A., LAKE, BLUE B., ZHAO, ADRIENNE H., SKARBREVIK, D., BABAJANIAN, E., SUKHWANI, M., SHUM, ELEEN Y., PHAN, MIMI H., PLANK, T.-DAWN M., RICHARDSON, MARCY E., RAMAIAH, M., SRIDHAR, V., DE ROOIJ, DIRK G., ORWIG, KYLE E., ZHANG, K. & WILKINSON, MILES F. 2016. The Homeobox Transcription Factor RHOX10 Drives Mouse Spermatogonial Stem Cell Establishment. *Cell Reports,* 17**,** 149-164.

SORRENTI, M., KLINGER, F. G., IONA, S., ROSSI, V., MARCOZZI, S. & DE FELICI, M. 2020. Expression and possible roles of extracellular signal-related kinases 1-2 (ERK1-2) in mouse primordial germ cell development. *The Journal of Reproduction and Development,* 66**,** 399-409.

SPILLER, C. M., FENG, C. W., JACKSON, A., GILLIS, A. J., ROLLAND, A. D., LOOIJENGA, L. H., KOOPMAN, P. & BOWLES, J. 2012. Endogenous Nodal signaling regulates germ cell potency during mammalian testis development. *Development* 139**,** 4123-4132.

SPILLER, C. M., GILLIS, A. J. M., BURNET, G., STOOP, H., KOOPMAN, P., BOWLES, J. & LOOIJENGA, L. H. J. 2016. Cripto: Expression, epigenetic regulation and potential diagnostic use in testicular germ cell tumors. *Molecular Oncology,* 10**,** 526-537.

SU, S., LAW, C. W., AH-CANN, C., ASSELIN-LABAT, M.-L., BLEWITT, M. E. & RITCHIE, M. E. 2017. Glimma: interactive graphics for gene expression analysis. *Bioinformatics,* 33**,** 2050-2052.

SUN, L., TRAN, N., LIANG, C., TANG, F., RICE, A., SCHRECK, R., WALTZ, K., SHAWVER, L. K., MCMAHON, G. & TANG, C. 1999. Design, synthesis, and evaluations of substituted 3-[(3- or 4-carboxyethylpyrrol-2-yl)methylidenyl]indolin-2-ones as inhibitors of VEGF, FGF, and PDGF receptor tyrosine kinases. *The Journal of Medicinal Chemistry,* 42**,** 5120-30.

SUZUKI, A. & SAGA, Y. 2008. Nanos2 suppresses meiosis and promotes male germ cell differentiation. *Genes and Development,* 22**,** 430-435.

TAGUCHI, J., SHIBATA, H., KABATA, M., KATO, M., FUKUDA, K., TANAKA, A., OHTA, S., UKAI, T., MITSUNAGA, K., YAMADA, Y., NAGAOKA, S. I., YAMAZAWA, S., OHNISHI, K., WOLTJEN, K., USHIKU, T., OZAWA, M., SAITOU, M., SHINKAI, Y., YAMAMOTO, T. & YAMADA, Y. 2021. DMRT1-mediated reprogramming drives development of cancer

resembling human germ cell tumors with features of totipotency. *Nature Communications,* 12**,** 5041.

TIAN, L., SU, S., DONG, X., AMANN-ZALCENSTEIN, D., BIBEN, C., SEIDI, A., HILTON, D. J., NAIK, S. H. & RITCHIE, M. E. 2018. scPipe: A flexible R/Bioconductor preprocessing pipeline for single-cell RNA-sequencing data. *PLOS Computational Biology,* 14**,** e1006361.

ULU, F., KIM, S.-M., YOKOYAMA, T. & YAMAZAKI, Y. 2017. Dose-dependent functions of fibroblast growth factor 9 regulate the fate of murine XY primordial germ cells†. *Biology of Reproduction,* 96**,** 122-133.

VALLIER, L., MENDJAN, S., BROWN, S., CHNG, Z., TEO, A., SMITHERS, L. E., TROTTER, M. W. B., CHO, C. H. H., MARTINEZ, A., RUGG-GUNN, P., BRONS, G. & PEDERSEN, R. A. 2009. Activin/Nodal signalling maintains pluripotency by controlling Nanog expression. *Development,* 136**,** 1339-1349.

WAKELING, S. I., MILES, D. C. & WESTERN, P. S. 2013. Identifying disruptors of male germ cell development by small molecule screening in ex vivo gonad cultures. *BMC Research Notes,* 6**,** 168.

WEISS, B. D., WOLTERS, P. L., PLOTKIN, S. R., WIDEMANN, B. C., TONSGARD, J. H., BLAKELEY, J., ALLEN, J. C., SCHORRY, E., KORF, B., ROBISON, N. J., GOLDMAN, S., VINKS, A. A., EMOTO, C., FUKUDA, T., ROBINSON, C. T., CUTTER, G., EDWARDS, L., DOMBI, E., RATNER, N., PACKER, R. & FISHER, M. J. 2021. NF106: A Neurofibromatosis Clinical Trials Consortium Phase II Trial of the MEK Inhibitor Mirdametinib (PD-0325901) in Adolescents and Adults With NF1-Related Plexiform Neurofibromas. *The Journal of Clinical Oncology,* 39**,** 797-806.

WESTERN, P. 2009. Foetal germ cells: striking the balance between pluripotency and differentiation. *International Journal of Developmental Biology,* 53**,** 393-409.

WESTERN, P. S., MILES, D. C., VAN DEN BERGEN, J. A., BURTON, M. & SINCLAIR, A. H. 2008. Dynamic regulation of mitotic arrest in fetal male germ cells. *Stem Cells,* 26**,** 339-347.

WESTERN, P. S., VAN DEN BERGEN, J. A., MILES, D. C. & SINCLAIR, A. H. 2010. Male fetal germ cell differentiation involves complex repression of the regulatory network controlling pluripotency. *The FASEB Journal,* 24**,** 3026-3035.

WU, D., LIM, E., VAILLANT, F., ASSELIN-LABAT, M.-L., VISVADER, J. E. & SMYTH, G. K. 2010. ROAST: rotation gene set tests for complex microarray experiments. *Bioinformatics,* 26**,** 2176-2182.

WU, Q., FUKUDA, K., WEINSTEIN, M., GRAFF, J. M. & SAGA, Y. 2015. SMAD2 and p38 signaling pathways act in concert to determine XY primordial germ cell fate in mice. *Development,* 142**,** 575-586.

YILDIRIM, E., AKSOY, S., ONEL, T. & YABA, A. 2020. Gonadal development and sex determination in mouse. *Reproductive Biology,* 20**,** 115-126.

YING, Q.-L., WRAY, J., NICHOLS, J., BATLLE-MORERA, L., DOBLE, B., WOODGETT, J., COHEN, P. & SMITH, A. 2008. The ground state of embryonic stem cell self-renewal. *Nature,* 453**,** 519-523.

YOUNGREN, K. K., COVENEY, D., PENG, X., BHATTACHARYA, C., SCHMIDT, L. S., NICKERSON, M. L., LAMB, B. T., DENG, J. M., BEHRINGER, R. R., CAPEL, B., RUBIN, E. M., NADEAU, J. H. & MATIN, A. 2005. The Ter mutation in the dead end gene causes germ cell loss and testicular germ cell tumours. *Nature,* 435**,** 360-364.

A

E12.5+72h

| | DMSO | FGFRi 500nM | MEKi 500nM | p38i 500nM | PI3Ki 500nM |

Bright-field

GFP

B **XX E12.5+48h**

Somatic cell proliferation

C **XX E12.5+48h**

Somatic cell proliferation

D **XY E12.5+72h**

Sertoli cell proliferation

E **XY E12.5+72h**

Germ cell proliferation

Figure 1

Figure 1: FGF and MEK1/2 signalling inhibition disrupts fetal testis development. **A)** Bright-field and GFP images of E12.5 XY gonad-mesonephros tissue cultured for 72h with DMSO, or 500nM of FGFRi, MEKi, p38i or PI3Ki. Scale bar: 100μm. Dotted lines highlight the gonad. **B-C)** Flow cytometric analysis of gonadal somatic cell proliferation based on EdU incorporation in XX E12.5 gonads-mesonephros tissue cultured for 48h with DMSO, FGF9 (50ng/mL), 500nM FGFRi or FGF9+FGFRi (B), or DMSO, FGF9 (50ng/mL), 500nM MEKi or FGF9+MEKi (C). **D-E)** Flow cytometric analysis of Sertoli (D) or germ (E) cell proliferation based on EdU incorporation in XY E12.5 gonad-mesonephros tissue cultured for 72h with DMSO or 500nM FGFRi, 500nM MEKi, 500nM p38i or 500nM PI3Ki. Replicates: n≥5; Statistics: B&E; Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons. C&D; Ordinary One-way ANOVA with Tukey's multiple comparisons; Error bars: Mean±SEM; Significance between treatments and controls; *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$.

Figure 2

Figure 2: FGF and MEK1/2 signalling inhibition reduce Sertoli cell proliferation, but only MEK1/2 inhibition disrupts germ cell mitotic arrest. A-B) Flow cytometric analysis of XY E12.5 gonad-mesonephros tissue cultured with DMSO, 125, 250, 500 or 1000nM of FGFRi or MEKi for 72h showing the percentage EdU incorporation in Sertoli (A) or germ (B) cells. **C-D)** Percentage EdU incorporation in Sertoli (C) or germ (D) cells from E11.5 XY gonad-mesonephros tissue cultured with DMSO or 500nM of FGFRi or MEKi for 96h. **E-F)** Percentage EdU incorporation in Sertoli (E) or germ (F) cells from E13.5 XY gonad-mesonephros tissue cultured with DMSO or 500nM MEKi for 48h. Replicates: n≥3. Statistics: A&B; Ordinary One-way ANOVA with Tukey's multiple comparisons. C&D; Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons. E&F; Unpaired two-tailed t-test. Error bars: mean±SEM. Significance between treatments and controls: *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$.

Figure 3

Figure 3: FGF and MEK1/2 signalling is required for organisation of Sertoli cells within testis cords. Immunofluorescent images of XY E12.5 gonad-mesonephros tissue cultured with DMSO, 500nM FGFRi or 500nM MEKi for 72h showing Sertoli (A) and germ (A and B) cell localisation. DAPI (blue; A&B), MVH (green; A&B), SMA (red; A), SOX9 (cyan; A) and Laminin (red; B). Scale bars: top panel 100µm, bottom panel 50µm. Replicates: n≥3. A: White arrows identify gaps in the Sertoli cell layer at the testis cord basement; white asterisks identify Sertoli cells dispersed within the inner area of the testis cords. B: White arrows identify germ cells localised outside testis cords.

Figure 4

Figure 4: MEK1/2 signalling is required for male germline differentiation. Analysis of E12.5 XY or XX gonad-mesonephros cultured for 72h with DMSO, 125, or 500nM FGFRi or MEKi. **A)** Immunofluorescent images demonstrating DPPA4 localisation. Top panel: DAPI (blue), MVH (green), DPPA4 (red), SMA (cyan). Bottom panel: DPPA4 (white). **B)** DPPA4 staining intensity in germ cells determined by flow cytometry. **C)** Immunofluorescent images demonstrating DNMT3L localisation. Top panel: DAPI (blue), MVH (green), NR2F2 (red), DNMT3L (cyan). Bottom panel: DNMT3L (white). **D-E)** Percentage DNMT3L positive germ cells (D) and DNMT3L staining intensity (E) determined by flow cytometry. **F)** Immunofluorescent images demonstrating PIWIL2 localisation. Top panel: DAPI (blue), MVH (green), SMA (red), PIWIL2 (cyan). Bottom panel: PIWIL2 (white). Scale bars: 50µm. Replicates: n≥3. Statistics: In B; Brown-Forsythe and Welch ANOVA with Dunnett's multiple comparisons. In D&E; Ordinary One-way ANOVA with Tukey's multiple comparisons. In B&E; Intensity is relative to E12.5+72h XY DMSO control sample set at 1.0. Error bars: mean±SEM. Significance between treatments and controls: **P<0.01, ***P<0.001, ****P<0.0001.

Figure 5

Figure 5: MEK1/2 signalling inhibition permitted *Stra8* expression but failed to effectively induce meiosis in XY germ cells. Analysis of XY or XX E12.5 gonad-mesonephros tissue cultured with DMSO or 500nM FGFRi or MEKi for 72h (A-C) or 96h (D-F). **A)** Immunofluorescent images demonstrating STRA8 localisation. Top panel: DAPI (blue), MVH (green), STRA8 (red), SMA (cyan). Bottom panel: STRA8 (white). **B)** Percentage STRA8 positive germ cells determined by flow cytometry. **C-D)** Immunofluorescent images demonstrating SYCP3 (C) and phospho-γH2AX (p-γH2AX) localisation. Left panel: DAPI (blue), MVH (green), SYCP3 (red; C), SMA (red; D), p-γH2AX (cyan). Middle panel: SCP3 (white; C). Right panel: p-γH2AX (white). **E)** Percentage p-γH2AX positive germ cells determined by flow cytometry. **F)** Flow cytometric cell cycle analysis of G0/G1, S-phase and G2/M based on the incorporation of EdU (S-phase) and propidium iodide (DNA content). A, C&D; scale bar: 50µm. Replicates: n≥3. B, E&F; Replicates: n≥8. Statistics: B; Ordinary One-way ANOVA with Tukey's multiple comparisons. E; Brown-Forsythe and Welch ANOVA with Dunnett's multiple comparisons. F; Repeated measure two-way ANOVA with Geisser-Greenhouse correction and Tukey's multiple comparisons. Error bars: mean±SEM. Significance between treatments and controls: **P<0.05, ***P<0.005, ****P<0.0001.

Figure 6

Figure 6: FGF-MEK signalling supports expression of stem cell associated genes in early germ cells, but only MEK1/2 signalling is required for male germline differentiation. RNA sequencing analysis of germ cells from XX or XY E12.5 gonads, or XX or XY E12.5 gonads cultured for 24 or 72h with DMSO or 500nM FGFRi or MEKi. **A)** Example of FACS scatterplot depicting GFP positive germ cell isolation. **B)** Number of differentially expressed genes (DEGs) between XX or XY E12.5 (time 0) and XX or XY DMSO controls from 24h and 72h cultures. **C)** Multidimensional scaling (MDS) of all control conditions. **D)** MDS of XX and XY gonads cultured for 24h. **E)** Differential gene expression analysis of XY E12.5+24h FGFRi XY (i) or XY E12.5+24h MEKi (ii) vs XY E12.5+24h DMSO. **F)** Venn-diagram of 24h FGFRi and MEKi DEGs (i) and heatmap of common DEGs (ii). Asterisks represent genes associated with germ cell tumours, cancer/stem cell biology and/or proliferation. **G)** MDS of XX and XY gonads cultured for 72h. **H)** Differential gene expression analysis of XY E12.5+72h FGFRi (i) or XY E12.5+72h MEKi (ii) vs XY E12.5+72h DMSO. **I)** Venn-diagram comparing MEKi 72h culture DEGs expressed lower (i) or higher (ii) than expected with XX or XY specific genes identified in B. **J)** Heatmap of DEGs identified in XY E12.5+72h MEKi vs XY E12.5+72h DMSO associated with male germline differentiation, DNA methylation/piRNA pathway, cell cycle, female germline differentiation/meiosis, Nodal signalling, pluripotency and spermatogonial stem cells (SSCs). Asterisks highlight genes associated with cell cycle, meiosis and pluripotency but are not differentially expressed. For all comparisons, genes with FDR <0.05 and |logFC|>0.585 (equivalent to |FC|>1.5) were considered differentially expressed.

Normal testis development

Male germline differentiation:
Mitotic arrest
Dnmt3L
Nanos2
Dppa4

Testis cord development:
Sertoli cell proliferation
Sertoli cell organisation

Erk1/2
Mek1/2
FGFR
Mek1/2
Erk1/2
Fgf9
Sox9
PMC
GC
SC
BM
Erk1/2
Mek1/2
Unknown ligand

FGFR inhibition

Male germline differentiation:
Mitotic arrest
Dnmt3L
Nanos2
Dppa4

Testis cord development:
Reduced *Sertoli cell proliferation*
Poor *Sertoli cell organisation*

FGFR
FGFR
Fgf9
Sox9
Erk1/2
Mek1/2
Unknown ligand

MEK1/2 inhibition

Male germline differentiation:
Failed *mitotic arrest*
Dnmt3L
Nanos2 } **Failed** *upregulatation*
Dppa4
Stra8 - **Increased** *expression*
Nanog - **Failed** *repression*

Testis cord development:
Reduced *Sertoli cell proliferation*
Poor *Sertoli cell organisation*

Mek1/2
FGFR
FGFR
Mek1/2
Fgf9
Sox9
Mek1/2
Unknown ligand

Figure 7

Figure 7: MEK1/2 and FGF signalling in male germline development. MEK1/2 signalling inhibition results in failed germ cell mitotic arrest, failed upregulation of male germline markers including *Nanos2,* DPPA4 and DNMT3L and aberrant expression of the pluripotency marker, *Nanog* and some female germline markers including STRA8*.* We propose that an unknown ligand activates MEK1/2 signalling to promote male germline differentiation including mitotic arrest and expression of male germline and DNA methylation markers such as *Nanos2, Dppa4* and *Dnmt3L.* MEK1/2 signalling is also essential for normal testis cord development, including Sertoli cell proliferation and organisation. FGFR is dispensable for male germline differentiation, but compromises Sertoli cell proliferation and organisation.

Supplementary Tables and Figures

Supplementary Tables 1 and 2

Supplementary Tables 1 and 2 contain gene lists generated from RNA sequencing analyses performed in this study.

Supplementary Figure 1: Representative plots depicting gating for antibodies used in flow cytometric analysis. A = Area, W = Width. **A)** Gates used to separate cells from debris (left) and to isolate single cells based on propidium iodide (PI) staining (right). **B)** Germ and somatic cells populations were identified by detecting Oct4GFP transgene (left). E12.5 mouse limb or mesonephros cells were used as a negative control (right). **C)** Germ and somatic cells populations were identified by detecting MVH staining (left). E12.5 mouse limb or mesonephros were used as a negative control (right). **D)** Sertoli cells were identified based on SOX9 staining (left). XX somatic cells were used as a negative control (right). **E-F)** Incorporation of EdU was used to identify proliferating Sertoli cells (E) or germ cells (F), with PI incorporation used to determine individual cell DNA content (left). E12.5 limb or mesonephros (E) or E12.5 XX germ cells not exposed to EdU (F) were used as a negative control for EdU (right). **G)** DNMT3L+ germ cells identified with DNMT3L staining (left). XX germ cells were used as a negative control (right). **H)** DPPA4+ germ cells were identified with DPPA4 staining (left). Cells not stained for DPPA4 were used as a negative control (right). **I-J)** E12.5+72h XY DMSO germ cells were used as a negative control for STRA8 (I) or p-γH2AX (J) staining (left). E12.5+72h XX DMSO germ cells were used as a positive control (right).

A

XX E12.5+48h
Somatic cell proliferation

B

XY E12.5+72h
Sertoli cell proliferation

AZD4547 concentration

C

XY E12.5+72h
Germ cell proliferation

AZD4547 concentration

D

XY E12.5+72h
Sertoli cell proliferation

E

XY E12.5+72h
Germ cell proliferation

F

XY E12.5+72h
Sertoli cell proliferation

G

XY E12.5+72h
Germ cell proliferation

Supplementary Figure 2

Supplementary Figure 2: Supplementary material relating to Figure 1 and 2. A) Flow cytometric analysis of gonadal somatic cell proliferation identified by EdU incorporation in E12.5 XX gonads/mesonephros tissue cultured for 48h with DMSO, FGF9 (50ng/mL), p38i (500nM), FGF9 + p38i, PI3Ki (500nM) or FGF9 + PI3Ki. **B-E)** A secondary inhibitor to target FGFR signalling (AZD4547; B-C), p38 signalling (ralimetinib dimesylate; D-E) or PI3K signalling (PF-04691502; D-E) was used to assess Sertoli (B&D) and germ cell (C&E) proliferation in E12.5 XY gonad-mesonephros cultured with DMSO or inhibitors for 72h. **F-G)** Sertoli (F) and germ cell (G) proliferation in E12.5 XY gonad-mesonephros cultured with DMSO, 500nM of MEKi, 2500nM of FGFRi or 5000nM of a third FGFR inhibitor (SU5402) for 72h. Replicates: n≥4. Statistics: Ordinary One-way ANOVA with Tukey's multiple comparisons. Error bars: mean±SEM. Significance between treatments and controls: $*P<0.05$, $***P<0.001$, $****P<0.0001$.

A

Sertoli cell proliferation

B

Germ cell proliferation

Supplementary Figure 3

Supplementary Figure 3: Flow cytometric scatterplot depicting Sertoli and germ cell proliferation. Flow cytometric scatterplots of XY E12.5 gonad-mesonephros tissue cultured in DMSO, 125, 250, 500 or 1000nM of FGFRi or MEKi for 72h showing the percentage EdU incorporation in Sertoli (A) or germ (B) cells. Percentage in top left corner of each graph represents the average proportion of Sertoli (A) or germ (B) cells in each treatment.

Supplementary Figure 4: Whole view of images displayed in Figure 4. Immunofluorescent images of XY E12.5 gonad-mesonephros or XY or XX E12.5 gonad-mesonephros tissue cultured with DMSO or 500nM of FGFRi or MEKi for 72h. **A)** Immunofluorescent images demonstrating DPPA4 localisation. Top panel: DAPI (blue), MVH (green), DPPA4 (red) and SMA (cyan). Bottom panel: DPPA4 (white). **B)** Immunofluorescent images demonstrating DNMT3L localisation. Left panel: DAPI (blue), MVH (green), NR2F2 (red) and DNMT3L (cyan). Right panel: DNMT3L (white). **C)** Immunofluorescent images demonstrating PIWIL2 localisation Left panel: DAPI (blue), MVH (green), SMA (red) and PIWIL2 (cyan). Right panel: PIWIL2 (white). Replicates: n≥3. Scale bar: 100μm.

Supplementary Figure 5: Whole view of images displayed in Figure 5. Immunofluorescent images of XY or XX E12.5 gonad-mesonephros tissue cultured with DMSO, FGFRi or MEKi for 72h (A-B) or 96h (C). **A)** Immunofluorescent images demonstrating STRA8 localisation. Top panel: DAPI (blue), MVH (green), STRA8 (red) and SMA (cyan). Bottom panel: Stra8 (white). **B-C)** Immunofluorescent images demonstrating SCP3 and phospho-γH2AX (p-γH2AX) localisation. Left panel: DAPI (blue), MVH (green), SCP3 (red; B) or SMA (red; C) and phospho-γH2AX (cyan). Middle panel: SCP3 (white; B). Right panel: p-γH2AX (white). Replicates: n≥3. Scale bar: 100µm.

Supplementary Figure 6

Supplementary Figure 6: Heatmap of 382 72h MEK1/2 dependent genes expressed higher than expected common in 72h XX germline specific genes. Genes which were expressed higher than expected in XY E12.5+72h MEKi vs XY E12.5+72h DMSO and were present in the 72h XX germline specific genes dataset (identified as genes which were upregulated in XX E12.5+72h DMSO vs XY E12.5+72h DMSO) were assessed. Of these 382 genes, 218 genes were highly expressed in XY E12.5 germ cells and XX E12.5+72h DMSO germ cells and were therefore not considered informative. 164 genes were not or were lowly expressed in XY E12.5 germ cells compared to XX E12.5+72h DMSO germ cells and were therefore considered more reliable female germline differentiation genes (identified by red box). Genes with an FDR <0.05 and |logFC| >0.585 (equivalent to |FC| >1.5) were considered differentially expressed.

Supplementary Figure 7: FGF and MEK1/2 inhibition does not result in abnormal maintenance of pluripotency markers. Immunofluorescent analysis of XY E13.5 gonad-mesonephros tissue or XY or XX E12.5 gonad-mesonephros tissue cultured with DMSO or 500nM of FGFRi or MEKi for 72h. **A)** Whole view immunofluorescent images demonstrating OCT4 and SOX2 localisation. Left panel: DAPI (blue), MVH (green), OCT4 (red) and SOX2 (cyan). Middle panel: OCT4 (white). Right panel: SOX2 (white). Scale bar: 100μm. **B-C)** Percentage of OCT4+ (B) or SOX2+ (C) germ cells calculated from immunofluorescent images. **D-E)** OCT4 (C) or SOX2 (E) intensity in germ cells relative to XY DMSO control set at 1.0, calculated from immunofluorescent images. Replicates: n=4. Statistics: Brown-Forsythe and Welch ANOVA with Dunnett's multiple comparisons. Error bars: mean±SEM. Significance between treatments and controls: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.