Overcoming clinical resistance to EZH2 inhibition using rational epigenetic combination therapy

Yaniv Kazansky[1,2], Daniel Cameron[1,2], Helen Mueller[1,2], Phillip Demarest[1,2], Nadia Zaffaroni[3], Noemi Arrighetti[3], Valentina Zuco[3], Yasumichi Kuwahara[4], Romel Somwar[5], Marc Ladanyi[5], Rui Qu[6], Elisa De Stanchina[6], Filemon Dela Cruz[7], Andrew Kung[7], Mrinal Gounder[8], Alex Kentsis[1,2,9]*

[1] Molecular Pharmacology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[2] Tow Center for Developmental Oncology, Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[3] Molecular Pharmacology Unit, Department of Experimental Oncology, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Milan, Italy

[4] Department of Biochemistry and Molecular Biology, Kyoto Prefectural University of Medicine

[5] Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[6] Antitumor Assessment Core, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[7] Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[8] Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[9] Departments of Pediatrics, Pharmacology, and Physiology & Biophysics, Weill Medical College of Cornell University, New York, NY, USA

* Correspondence: Alex Kentsis, MD, PhD, email kentsisresearchgroup@gmail.com, telephone +1-646-888-2593

Running title: Overcoming resistance to EZH2 inhibition

Keywords: epigenetics, EZH2 inhibition, BAF, combination therapy, PRC2

Abstract

Essential epigenetic dependencies have become evident in many cancers. Based on the functional antagonism between BAF/SWI/SNF and PRC2 in *SMARCB1*-deficient sarcomas, we and colleagues recently completed the clinical trial of the EZH2 inhibitor tazemetostat. However, the principles of tumor response to epigenetic therapy in general, and tazemetostat in particular, remain unknown. Using functional genomics of patient tumors and diverse experimental models, we sought to define molecular mechanisms of tazemetostat resistance in *SMARCB1*-deficient sarcomas and rhabdoid tumors. We found distinct classes of acquired mutations that converge on the RB1/E2F axis and decouple EZH2-dependent differentiation and cell cycle control. This allows tumor cells to escape tazemetostat-induced G1 arrest despite EZH2 inhibition, and suggests a general mechanism for effective EZH2 therapy. This also enables us to develop combination strategies to circumvent tazemetostat resistance using cell cycle bypass targeting via AURKB, and synthetic lethal targeting of PGBD5-dependent DNA damage repair via ATR. This reveals prospective biomarkers for therapy stratification, including PRICKLE1 associated with tazemetostat resistance. In all, this work offers a paradigm for rational epigenetic combination therapy suitable for immediate translation to clinical trials for epithelioid sarcomas, rhabdoid tumors, and other epigenetically dysregulated cancers.

Significance: Genomic studies of patient epithelioid sarcomas, rhabdoid tumors, and their cell lines identify mutations converging on a common pathway that is essential for response to EZH2 inhibition. Resistance mutations decouple drug-induced differentiation from cell cycle control. We identify complementary epigenetic combination strategies to overcome resistance and improve durability of response, supporting their investigation in clinical trials.

Introduction

Unlike conventional cytotoxic chemotherapy, epigenetic therapy offers the ability to target cancer-specific dependencies with increased specificity and reduced toxicity. This is especially true for cancers caused by genetic mutations of transcriptional regulators, such as the chromatin remodeling BAF/SWI/SNF (Brg/Brahma-associated factors) complex that is mutated in more than 20% of human cancers (1). For example, malignant rhabdoid tumors (MRT) and epithelioid sarcomas (ES) are lethal tumors of children and young adults caused by inactivating mutations of *SMARCB1/SNF5/INI1/BAF47*, one of the core BAF/SWI/SNF complex subunits (2-4). Loss of normal BAF function can confer a dependency on the Polycomb Repressive Complex 2 (PRC2) and its methyltransferase EZH2. This dependency is thought to result from epigenetic antagonism between the two complexes, in which normal BAF activity evicts PRC2 from tumor suppressor loci (5, 6).

High *EZH2* expression in many cancer types and gain-of-function *EZH2* mutations in lymphomas (7-10) have led to the development of multiple inhibitors of EZH2 (11-13), several of which have now entered clinical trials (14). This includes clinical trials for *SMARCB1*-deficient tumors without *EZH2* mutations (15). However, despite the potential of such targeted epigenetic therapies, the principles governing their effective application remain unknown. Intrinsic and acquired resistance limits their use as monotherapies (16, 17).

A recent clinical trial led to the FDA approval of the EZH2 methyltransferase inhibitor tazemetostat (TAZ) as the first targeted therapy for *SMARCB1*-deficient epithelioid sarcomas (15). However, only 15% of patients showed objective clinical responses, with most epithelioid sarcomas being resistant to TAZ. Ongoing clinical trials in patients with rhabdoid tumors have shown similar results, with only a subset of brain atypical teratoid rhabdoid tumor (ATRT) patients exhibiting objective responses to TAZ, while extracranial rhabdoid tumors appear to be uniformly resistant (18). Recent studies have nominated the histone methyltransferase NSD1 as a regulator of TAZ susceptibility in rhabdoid tumor cells, but the clinical relevance of this mechanism is currently not known (19). In all, the outcomes for most patients with MRT and ES remain dismal, and the mechanisms of clinical response and resistance to EZH2 inhibition remain unknown. This hinders our ability to stratify treatment using prospective biomarkers to identify patients who may benefit from TAZ and to develop effective combination therapies with improved and durable benefits for patients.

Here, we define the key requirements for effective epigenetic therapy in diverse *SMARCB1-*
deficient epithelioid sarcomas and rhabdoid tumors *in vivo*. Using comparative genomic
analyses of clinical trial patients treated with TAZ, we identify multiple acquired mutations that
cause therapy resistance. Using functional genomic approaches, we show that resistance
mechanisms converge on a common RB1/E2F axis that integrates control of tumor cell division
and differentiation. This organizes patient resistance mutations into a general framework,
allowing us to develop rational combination therapies to circumvent TAZ resistance. Using
diverse patient-derived ES and MRT cell lines and mouse xenografts, we demonstrate cell cycle
bypass and synthetic lethal treatment strategies suitable for immediate translation to
combination clinical trials for patients.

Results

Patient tumor sequencing reveals diverse resistance mutations

To identify mutations associated with clinical resistance to TAZ, we performed targeted gene
sequencing of patient tumors using MSK-IMPACT, which is based on a panel of over 500 genes
recurrently mutated in diverse cancer types (20). We analyzed 33 tumor specimens from 20
patients treated as part of the recent TAZ clinical trial (15), and identified somatic tumor
mutations in matched pre- and post-treatment specimens (**Supplementary Tables 1 and 2**).
We found distinct sets of somatic mutations in responding and non-responding tumors, with
nearly all mutations, apart from *SMARCB1* loss itself, being exclusive to either TAZ-responsive
or TAZ-resistant tumors (**Figure 1A, top panel**). Strikingly, we observed two tumors which
initially responded to TAZ based on radiographic imaging but later developed clinical resistance
(**Figure 1A, bottom panel, 1B**). Targeted sequencing of the resistant tumors revealed two
newly acquired somatic mutations: heterozygous missense mutation of *EZH2* (*EZH2^{Y666N}*) in the
Patient 3 tumor specimen, and biallelic loss of function mutation of *RB1*, including a hemizygous
deletion and a frame shift mutation (*RB1^{del}*) in the remaining allele, in the Patient 15 tumor
specimen. We confirmed both mutations using RNA-seq of the respective tumor specimens
(**Figure S1A-B**). Since one mutation affected *EZH2* directly, and the other involved the known
EZH2 target *RB1* (21), we hypothesized that both mutations were responsible for TAZ
resistance in their respective patients.

First, we investigated the $EZH2^{Y666N}$ mutation. Past studies using forward genetic screens in lymphoma cell lines have identified putative resistance mutations within both the N-terminal D1 domain and the catalytic SET domain of EZH2, both of which are predicted to interact with S-adenosyl methionine (SAM)-competitive EZH2 inhibitors such as TAZ (22, 23). One such SET domain mutation previously identified in cell lines is $EZH2^{Y661D}$. Y661 corresponds to Y666 in isoform 2 of $EZH2$, the isoform referred to in this study, and thus is concordant with the mutation we observed in the Patient 3 tumor (22).

Based on the atomic resolution structure of a pyridone-based EZH2 inhibitor bound to the $Anolis$ $carolinensis$ EZH2 (PDB 5IJ7) (24), we reasoned that residue Y666 in human EZH2 may form a critical part of the TAZ binding site and that its mutation can prevent TAZ from binding to the SET domain (**Figure 1C**). To test $EZH2^{Y666N}$ as a resistance allele, we expressed doxycycline-inducible $EZH2^{Y666N}$ in $SMARCB1$-deficient G401 rhabdoid tumor cells. We observed that $EZH2^{Y666N}$ expressing clones are resistant to TAZ as compared to cells expressing equal levels of wild-type $EZH2$ by assessing both cell viability (**Figure 1D**) and cell morphology (**Figure S1C**). The resistance phenotype of $EZH2^{Y666N}$-expressing cells depends on the intact catalytic activity of the SET domain, as the compound mutant combining the catalytically inactive triple mutant $EZH2^{F672I,H694A,R732K}$ ($EZH2^{CatMut}$) with the Y666N mutation, termed $EZH2^{QuadMut}$, did not confer resistance to TAZ (**Figure S1D**). This indicates that the $EZH2^{Y666N}$ mutation does not significantly interfere with enzymatic activity. We also observed that $EZH2^{Y666N}$ confers resistance to the dual EZH1/2 inhibitor valemetostat (25), consistent with putative resistance to SAM-competitive, pyridone-based EZH2 inhibitors (**Figure S1E**). This suggests that combined inhibition of EZH2 and EZH1 may not overcome this type of acquired EZH2 inhibitor resistance.

Previous studies found that lymphoma cells resistant to EZH2 inhibitors remained susceptible to the inhibition of the non-enzymatic PRC2 subunit EED (26), including those with mutations in the $EZH2$ SET domain ($EZH2^{C663Y}$ and $EZH2^{Y726F}$) and the D1 domain (23, 27). We therefore hypothesized that TAZ resistance conferred by $EZH2^{Y666N}$ could be overcome by PRC2 inhibitors that do not bind to EZH2. Indeed, we found that the allosteric EED inhibitor MAK683 overcomes $EZH2^{Y666N}$-mediated resistance (28), demonstrating that these cells remain generally susceptible to PRC2 inhibition (**Figure 1D**). EED inhibition may thus be an effective strategy to overcome acquired TAZ resistance mutations in $EZH2$.

155 ***RB1* loss allows escape from cell cycle arrest despite effective EZH2 inhibition**

156

157 Past work has shown that *EZH2* is a direct target of repression by RB1/E2F (21, 29). This

158 suggests that acquired *RB1* loss may confer resistance to EZH2 inhibition by increasing EZH2

159 expression. To test *RB1^{del}* as a TAZ resistance allele, we used CRISPR/Cas9 genome editing to

160 generate biallelic *RB1^{del}* mutations in G401 cells, as compared to the isogenic *RB1*-wild type

161 control cells, engineered by targeting the safe harbor locus *AAVS1*. We confirmed absence of

162 RB1 protein expression in two independent clones using Western blotting, and found that *RB1^{del}*

163 cells were indeed resistant to TAZ (**Figure 1E**). Even when the TAZ dosage is increased as

164 high as 10 μM, this resistance persists unabated.

165

166 Despite *EZH2* being a known target gene of RB1/E2F, we were surprised to observe that *RB1^{del}*

167 G401 cells showed similar morphological changes upon TAZ treatment (**Figure S2A**) to those

168 previously reported for TAZ-treated *RB1^{WT}* G401 cells (11). To define the effects of TAZ on

169 *RB1^{del}* cells more precisely, we performed RNA-seq of isogenic G401 *RB1^{del}* and wildtype

170 *AAVS1*-control cells, treated with either 10 μM TAZ or DMSO control for 11 days, based on an

171 established treatment regimen to model EZH2 inhibition *in vitro* (11). As predicted, we observed

172 that *EZH2* mRNA and protein levels remained high in TAZ-treated *RB1^{del}* cells, unlike in *RB1^{WT}*

173 cells (**Figure 2A-C, S2B**). However, EZH2 inhibition induced significant upregulation of

174 hundreds of genes in both *RB1^{WT}* and *RB1^{del}* cells, including upregulation of known PRC2 target

175 genes (**Figure 2A-F**).

176

177 Importantly, trimethylation of the EZH2 substrate H3K27 was substantially reduced by TAZ

178 regardless of *RB1* status (**Figure S2B**). This indicates that despite persistent EZH2 expression,

179 EZH2 methyltransferase activity is effectively inhibited by TAZ regardless of *RB1* loss. Gene set

180 enrichment analysis (GSEA) showed that all three clones significantly upregulated multiple gene

181 sets upon TAZ treatment (**Supplementary Table S3**, **Figure S2C**). This included gene sets

182 associated with mesenchymal cell differentiation such as epithelial-to-mesenchymal transition

183 (EMT; **Figure S2D**), as well as specific markers of differentiation such as *MMP2* (**Figure S3A**),

184 used previously as a mesenchymal marker induced by *SMARCB1* re-expression in rhabdoid

185 tumor organoids (30). This is reminiscent of recent observations that re-expression of

186 *SMARCB1* in G401 cells can lead to a mesenchymal chromatin state (31), and is consistent

187 with the idea that PRC2 inhibition may allow BAF to re-activate a more developmentally normal

188 gene expression state. While MMP2 protein did not appear to be upregulated by TAZ (**Figure**

S3B-C), the induction of mesenchymal genes suggests that TAZ can induce a transcriptional

differentiation program regardless of *RB1* status. Taken together, these findings indicate that

RB1 loss-induced TAZ resistance is independent of EZH2.

In addition to *EZH2*, other RB1/E2F target genes were also upregulated in TAZ-treated *RB1^{del}*

cells compared to TAZ-treated *RB1^{WT}* control cells (**Figure S4A-B**). Given the function of RB1 in

the regulation of the G1/S cell cycle checkpoint, we hypothesized that *RB1* loss could allow cells

to escape TAZ-induced cell cycle arrest. Flow cytometry cell cycle analysis showed that G401

cells treated with TAZ doses as low as 1 µM arrest at the G1/S checkpoint, as reported

previously (11). However, *RB1^{del}* cells exhibited a significant reduction in the proportion of cells

in G1 phase upon TAZ treatment (50% of TAZ-treated *RB1^{WT}* cells versus 31% and 28% for

RB1^{del} E1 and F2 clones, respectively; Student's t-test p = 1.4E-6 and 1.1E-6 for *RB1^{WT}* versus

RB1^{del} E1 and F2, respectively), with a corresponding increase of the proportion of cells

remaining in S and G2/M phases (**Figure 2G**). In agreement with these findings, we observed

persistent mRNA expression of S/G2/M-phase-associated *CCNA2*, *CDK2*, and *AURKB* genes in

RB1^{del} cells upon TAZ treatment (**Figure 2A-C**), despite the upregulation of the CDK4/6 inhibitor

CDKN2A (p16), a known PRC2 target in MRT (5, 32, 33). We confirmed persistent maintenance

of the S-phase cyclin A2 (CCNA2) protein levels despite p16 upregulation using Western

blotting (**Figure 2H**). Together, these results show that *RB1* loss is sufficient to evade TAZ-

induced cell cycle arrest at the G1/S restriction point despite maintaining the expected global

transcriptional response to EZH2 inhibition, including upregulation of cell cycle inhibitor genes.

Intact RB1/E2F axis is required for TAZ susceptibility

The requirement for *RB1* expression in the therapeutic response to TAZ suggests that an intact

RB1/E2F axis may be a general requirement for effective EZH2 inhibitor therapy. This would

predict that other genetic and epigenetic perturbations to the RB1/E2F axis, beyond *RB1* loss

itself, would similarly confer escape from TAZ-induced cell cycle arrest. Analysis of our TAZ

clinical trial treatment cohort revealed one patient tumor specimen with primary resistance to

TAZ with intact *RB1* but inactivating mutations of both *CDKN2A* and *CDKN2B* (**Figure 1A**,

Supplementary Tables 1-2), both of which are known to inhibit CDK4/6-mediated

phosphorylation of RB1. Two additional specimens had missense mutations in *ANKRD11*: One

tumor with primary resistance to TAZ and another which initially responded but later progressed

on treatment, at which point a newly acquired *ANKRD11* mutation was detected (**Figure 1A**-

Patients 9 and 16, respectively). *ANKRD11* is a known TP53 cofactor and putative tumor
suppressor that contributes to TP53-mediated expression of pan-CDK inhibitor *CDKN1A* (34-
36). *CDKN1A* itself is also known to be a PRC2 target in tumors (37-39), although its role in the
response to EZH2 inhibition in *SMARCB1*-deficient sarcomas is currently unknown. These
results converge on the dysregulation of the RB1/E2F axis as a clinical mechanism of evasion
of TAZ-induced cell cycle arrest.

We hypothesized that epithelioid and rhabdoid tumor cell lines could also harbor mutations of
the RB1/E2F axis. To investigate the functional determinants of tumor cell response to TAZ, we
first analyzed the response to TAZ of seven MRT and four ES cell lines in which we confirmed
loss of SMARCB1 protein expression in all ES and MRT cell lines using Western blotting, as
compared to SMARCB1-expressing HEK293T cells (**Figure S5A**). We classified each line as
sensitive or resistant based on the area under the curve (AUC) of their TAZ dose responses
(AUC > 0.3 for sensitive G401, KP-MRT-NS, TTC642, A204, TM8716, KP-MRT-RY cell lines
and AUC ≤ 0.3 for resistant ES1, VAESBJ, ES2, EPI544, MP-MRT-AN cell lines; **Figure S5B**).
Given that TAZ treatment requires at least 4 days for the cellular reduction of methylated EZH2
substrates and at least 7 days for apparent antiproliferative effects in the rapidly-dividing G401
cell line (11), we confirmed that the apparent TAZ susceptibilities of these MRT and ES cell
lines are not correlated with their proliferation rates (Pearson's r = -0.073 and p = 0.81; **Figure
S5C**).

In agreement with somatic mutations affecting the RB1/E2F axis associated with TAZ resistance
in clinical tumor specimens (**Figure 1A**), we found mutations of *CDKN2A* in 4 out of 5 TAZ-
resistant cell lines, *CDKN2B* in 2 out of 5 resistant cell lines, *CDKN1A* in 1 out of 5, and
ANKRD11 in 3 out of 5*,* as compared to no such mutations in any TAZ-sensitive MRT and ES
cell lines (**Figure 3A**, **Supplementary Table 4**). While our analysis detected a reduction in copy
number of *RB1* in TAZ-sensitive KP-MRT-RY cells, manual inspection of sequencing reads
within the *RB1* gene revealed lack of homozygous deletion, with presumed retention of *RB1*
expression (**Figure S5D**). Thus, mutations associated with the RB1/E2F axis are associated
with resistance to TAZ in *SMARCB1*-deficient cell lines and patient tumors.

We confirmed that genetic mutation of *CDKN2A* correlates with lack of p16 protein induction
upon TAZ treatment despite effective EZH2 inhibition by Western blotting of p16 and
H3K27me3 in four TAZ-sensitive and four TAZ-resistant cell lines (**Figure S6A**). Consistent with

the genomic data and observations in $RB1^{del}$ cells, TAZ-resistant cell lines showed no p16
induction despite complete loss of H3K27me3. Furthermore, TAZ reduced protein expression of
E2F targets EZH2 and CCNA2 only in TAZ-sensitive cell lines (**Figure S6A**).

To test whether *CDKN2A* and *CDKN1A* mutations can confer resistance to TAZ, we engineered
CRISPR/Cas9-mediated biallelic knockouts of *CDKN2A* or *CDKN1A* in G401 cells, generating
single-cell clones with respective mutant alleles (**Figure S6B-D**). We also attempted to knock
out the *ANKRD11* gene, but were unable to obtain viable biallelic mutant clones. We found that
deletion of either *CDKN1A* or *CDKN2A* was sufficient to confer resistance to TAZ (**Figure S6E,**
3B). Thus, perturbations of the RB1/E2F axis beyond loss of *RB1* itself are sufficient to allow
rhabdoid tumor cells to proliferate in spite of EZH2 inhibition.

We reasoned that the analysis of gene expression dynamics of TAZ-sensitive and TAZ-resistant
tumor cells may be used to identify functional markers of response and resistance. Thus, we
assessed the apparent transcriptional activity of the RB1/E2F axis in patient tumors using
quantitative gene expression RNA-seq analysis of TAZ responding and non-responding tumors
biopsied before and after TAZ treatment (**Supplementary Table S1**). Pre-treatment TAZ-
resistant tumors exhibited significant enrichment of multiple Gene Ontology (GO) terms
associated with the cell cycle and in particular with the S and G2/M phases (**Figure S7A**).
Similarly, post-treatment tumors that progressed on TAZ showed increased gene expression of
GO terms associated with mitosis, as compared to TAZ-responsive tumors (**Figure S7B**).
Indeed, TAZ-resistant tumors exhibited consistently higher expression of S/G2/M-phase-
associated genes prior to treatment (**Figure S7C-D**). These findings suggest that in addition to
the mutations affecting the RB1/E2F axis associated with TAZ resistance, additional mutations
not captured by MSK-IMPACT targeted gene sequencing and/or epigenetic dysregulation, such
as putative silencing of tumor suppressor genes like *CDKN1A* or *CDKN2A*, may contribute to
the decoupling of RB1/E2F-mediated proliferation and PRC2-regulated differentiation, and
consequent TAZ resistance.

Since TAZ-resistant MRT and ES cell lines and patient tumors show distinct mutations and gene
expression changes that converge on the RB1/E2F axis, we inquired whether these
perturbations would similarly converge on common prognostic biomarkers of TAZ resistance.
Comparative gene expression analysis of untreated $RB1^{del}$ cells versus $RB1^{WT}$ G401 cells
showed a small set of consistently and significantly up- and down-regulated genes in two

independent clones (**Figure S8A**). The most substantially and significantly upregulated gene associated with *RB1* loss was *PRICKLE1* (**Figure 3C-D**), which we confirmed to be overexpressed at the protein level in both *RB1^{del}* clones using Western blotting (**Figure 3E**).

We also found that *PRICKLE1* was among the most differentially expressed genes between 10 pre-treatment patient tumors with response (6) and resistance (4) to TAZ, with *PRICKLE1* expression being higher in TAZ-resistant tumors (mean normalized read counts = 10,237 and 300 for resistant and responsive tumors, respectively; Student's t-test *p* = 0.013; **Figure 3F, S8B**). *PRICKLE1* can control planar cell polarity (PCP), a key cell differentiation pathway, and has previously been implicated as a prognostic biomarker of poor prognosis in breast cancer (40, 41), acute myeloid leukemia (42), and gastric cancer (43, 44). Several other genes encoding PCP pathway factors were also upregulated in TAZ-resistant tumors compared to TAZ-sensitive tumors (**Figure S8B**). We also examined differentially expressed genes in TAZ-sensitive tumors as potential markers of sensitivity. These included the transcription factor *KLF4* which can control the G1/S transition by regulating *CDKN1A* expression (45, 46) (mean normalized read counts = 200 and 2,489 for resistant and responsive tumors, respectively; Student's-test *p* = 0.03; **Figure S8C, 3G**). Thus, PRICKLE1 and possibly additional factors controlling PCP and integration of RB1/E2F cell cycle and differentiation are potential prognostic pre-treatment biomarkers to identify clinical TAZ resistance and susceptibility of *SMARCB1*-deficient tumors.

Synthetic lethal and cell cycle bypass epigenetic combination strategies overcome tazemetostat resistance

Given that *RB1^{del}* cells are able to bypass cell cycle arrest at the G1/S checkpoint, we reasoned that inhibiting cell cycle kinases downstream of this checkpoint could overcome TAZ resistance. We focused on the cell cycle kinases *CDK2* and *AURKB*, which are downregulated by EZH2 inhibition in TAZ-sensitive cells but persistently expressed in TAZ-resistant cells (**Figure 2A-C**), as potential cell cycle bypass targets. We found that the CDK2 inhibitor seleciclib (47), the mitotic kinase Aurora A inhibitor alisertib (48), and the Aurora B inhibitor barasertib (49) were able to overcome TAZ resistance in *RB1^{del}* G401 cells (**Figure 4A, Figure S9A-B**). Consistent with the function of CDK4/6 kinases upstream of RB1/E2F, sensitivity to CDK4/6 inhibitors palbociclib and abemaciclib was reduced by *RB1^{del}* mutation (**Figure S9C-D**). In support of the cell cycle bypass TAZ combination strategy, we observed that patient tumors which progressed

on TAZ showed higher expression of *AURKB* mRNA as compared to those that responded (**Figure 3H**). Combined with the high sensitivity of G401 cells to barasertib (**Figure 4A**; half-maximal effective concentration of 6.5 ± 0.5 nM, 5.5 ± 0.6 nM, and 5.9 ± 0.9 nM for $RB1^{WT}$, $RB1^{del}$ E1, and F2 clones, respectively), these findings suggest that the cell cycle bypass strategy may be used to overcome TAZ resistance.

We therefore asked whether the combination of TAZ and barasertib would have activity against both TAZ-responsive and TAZ-resistant *SMARCB1*-deficient MRT and ES cell lines, using RPE cells as a *SMARCB1*-proficient control. The effects of the combination of TAZ and barasertib on cell viability did not substantially exceed the effect of barasertib alone at the doses tested (**Figure 4B**; 200 nM TAZ, 8 nM barasertib). However, nearly all cell lines tested, including those resistant to TAZ monotherapy, showed substantial susceptibility to barasertib, with RPE cells displaying the lowest sensitivity (**Figure 4B**). Cell cycle analysis on G401 cells treated with this combination showed that the combined TAZ and barasertib treatment caused a greater degree of cell cycle arrest and reduced entry into S phase than either drug alone, as measured by EdU incorporation (**Figure S10A**). This was the case in both $RB1^{WT}$ and $RB1^{del}$ cells, consistent with the prediction that cell cycle inhibition downstream of the G1/S checkpoint would cause cell cycle arrest even in cells with *RB1* loss (**Figure S8A**). Furthermore, the S phase reduction was not due to apoptosis, as measured by Western blot and immunofluorescence analysis of Caspase 3 cleavage (**Figure S10B-C**). Finally, this effect was associated with the induction of large multi-nucleated cells (**Figure S10C**), consistent with observations of postmitotic endoreduplication upon AURKB inhibition (50, 51).

To test the effects of combined TAZ and barasertib therapy *in vivo*, we treated a panel of five patient-derived rhabdoid tumor and epithelioid sarcoma xenografts (PDX) engrafted in immunodeficient mice (**Supplementary Table S5**). In contrast to the modest reduction of tumor growth and extension of survival of mice with tumors <1,000 mm^3 with TAZ or barasertib treatment alone, PDX mice treated with the combination of TAZ and barasertib showed significant reductions in tumor growth (Vardi *U*-test *p* = 4.0E-4 and 2.0E-4 for combination versus barasertib or TAZ, respectively; **Figure 4C**) (52). In two of the PDX models, this combination led to tumor regressions (**Figures 4E-F & S10D**). Consistent with this benefit, the combination was also found to significantly increase mean tumor-free animal survival from 65 days (95% confidence interval (CI) = 52-78 days) for barasertib and 68 days (95% CI = 53-82 days) for TAZ to 98 days (95% CI = 84-112 days) for the combination (log-rank test *p* = 3.3E-3

and 7.4E-3 for combination versus barasertib or TAZ, respectively; **Figure 4D**). These results

indicate that the combination of TAZ with a downstream cell cycle inhibitor such as barasertib

can improve response and overcome resistance to TAZ in diverse rhabdoid tumors and

epithelioid sarcomas *in vivo*.

Our studies with MRT and ES cell lines showed that p16 induction, rather than inhibition of

EZH2 and H3K27me3 loss, correlates with TAZ response. To determine whether this

relationship held *in vivo*, we measured H3K27me3 and p16 levels in three PDX models for

which sufficient mice were engrafted and treated. One model, HYMAD_EPIS_X0004aS1, did

not respond to TAZ monotherapy (log-rank test p = 0.63), while the other two,

SOMWR_EPIS_X00013aS1 and KUNGA_MRT_X0002aS1 showed significant anti-tumor

activity of the TAZ alone (log-rank test p = 7.1E-6 and 8.0E-3 respectively; **Figure S11A**). While

all three models showed reductions of H3K27me3 upon TAZ treatment, only TAZ-sensitive

tumors showed upregulation of p16 and reduction of the cell proliferation marker Ki67 (**Figure**

S11B-C). In addition, TAZ did not appear to induce apoptosis *in vivo* as evidenced by the

absence of cleaved caspase 3 staining (**Figure S11B-C**), consistent with the results *in vitro*

(**Figure S10B-C**).

In addition to the distinct cell cycle dynamics of TAZ resistance, we observed that TAZ

treatment also caused significant increase in the expression of the *PiggyBac transposable*

element derived 5 (*PGBD5)* regardless of *RB1* mutation (**Figure 2A-C**). *PGBD5* is a

transposase-derived gene with retained nuclease activity in human cells (53-55), which requires

active end-joining DNA repair, including the DNA damage repair signaling kinase ATR (56). In

rhabdoid tumors and medulloblastomas, PGBD5 has been implicated as a developmental

mutator and effector of somatic deletions of *SMARCB1* itself, and has been found to induce

double-strand DNA (dsDNA) breaks and DNA rearrangements (53, 56-58).

Since *PGBD5* is both necessary and sufficient to confer a dependency on ATR kinase signaling

in tumor cells (56), we reasoned that TAZ-induced upregulation of *PGBD5* expression may

potentiate the synthetic lethal dependency between PGBD5 and ATR-dependent DNA damage

repair. To test this idea, we used the ATR-selective kinase inhibitor elimusertib, which is

currently undergoing clinical trials in patients with solid tumors, including patients with PGBD5-

expressing tumors such as MRT and ES (Clinical Trials Identifier NCT05071209). We found that

elimusertib exhibited low-nanomolar potency against $RB1^{WT}$ and $RB1^{del}$ G401 cells *in vitro* (half-

maximal effective concentration of 18 ± 1.6 nM, 19 ± 3.8 nM, and 27 ± 3.2 nM for $RB1^{WT}$, $RB1^{del}$

E1 and F2 clones, respectively; **Figure 5A**). We also found that the combination of TAZ and

elimusertib exerted greater antitumor effects than either drug alone against diverse MRT and

ES cell lines (**Figure 5B**), exhibiting synergy in a subset of the cell lines (**Figure S12A-B**). To

determine whether the synergistic combination of elimusertib and TAZ antitumor effects were

due to increased DNA damage, we used confocal immunofluorescence microscopy to quantify

γH2AX, as a specific marker of DNA damage (59). In agreement with prior studies (56),

untreated G401 cells showed measurable γH2AX associated with baseline *PGBD5* expression

(**Figure 5D**). Consistent with TAZ-mediated induction of *PGBD5* expression (**Figure 2A-C**), we

found that TAZ treatment alone significantly increased nuclear γH2AX (mean normalized level =

0.42 versus 0.29 for TAZ and DMSO, respectively; *t*-test p = 9.3E-3; **Figures 5C-D, S12C**), and

the combination of TAZ and elimusertib induced additional increases in γH2AX than either drug

alone (mean normalized levels = 1.8 for the combination versus 0.42 and 0.85 for TAZ and

elimusertib, respectively; *t*-test p = 6.2E-14 and 1.5E-29 for combination versus elimusertib and

TAZ, respectively; **Figure 5D, S12C**).

To determine whether TAZ-mediated induction of DNA damage, associated with increased

expression of *PGBD5*, is indeed PGBD5-dependent, we engineered short hairpin RNA-

mediated knockdown of *PGBD5* in G401 cells using two specific shRNA constructs (shPGBD5),

as compared to a non-targeting control GFP targeting shRNA (shGFP; **Figure S13A**). We found

that while the combination of TAZ and elimusertib induced DNA damage in shPGBD5 cells, this

effect was significantly reduced compared to control shGFP cells (Student's t-test, p = 3.9E-8

and 4.4E-8 for shGFP versus shPGBD5-1 and shPGBD5-3, respectively; **Figure S13B**). This

was the case both when examining all nuclei with γH2AX (**Figure S13B**), as well as analyzing

nuclei with punctate γH2AX staining (**Figure S13C**), which have more localized DNA damage

versus those with pan-nuclear γH2AX staining (**Figure S13D**), which are cells with genome-

wide unrepaired DNA damage including cells undergoing apoptosis. Thus, the induction of DNA

damage by the combination of TAZ and elimusertib is PGBD5-dependent, as explained by the

synthetic lethal dependency between PGBD5 and end-joining DNA repair and the TAZ-

mediated increase in PGBD5 expression and its activity as a DNA nuclease that induces DNA

damage.

This synthetic lethal targeting strategy is specific, because the combination of TAZ with the

SRA737 inhibitor of another DNA damage repair signaling kinase CHK1 showed no increased

activity as compared to either drug alone (**Figure S14A-B**). Indeed, TAZ treatment did not
induce apparent replication stress, as measured by RPA phosphorylation (**Figure S14B**), which
was also not potentiated by the combined CHK1 inhibition with SRA737, despite effective
suppression of CHK1 auto-phosphorylation (**Figure S14B**). Unlike T-ALL, where EZH2
suppression induces MYCN protein expression and replication stress (60), TAZ treatment of
G401 rhabdoid tumor cells failed to increase MYCN protein levels (**Figure S14C**), in spite of
significant upregulation of *MYCN* mRNA (**Figure 2A-C**).

Encouraged by the potent and specific antitumor activity of synthetic lethal combination TAZ
therapy *in vitro*, we tested the antitumor activity of elimusertib and TAZ combination using a
diverse cohort of MRT and ES PDX mice *in vivo* (**Supplementary Table S5**). The combination
of TAZ and elimusertib exceeded the effect of treatment with either drug alone when assessed
by tumor measurements (Vardi *U*-test p = 2.33E-2 and 0.20 for combination versus elimusertib
or TAZ, respectively; **Figure 5E**) and significantly extended tumor-free survival from 51 days
(95% CI = 42-60 days) for elimusertib and 68 days (95% CI = 53-82 days) for TAZ to 100 days
(95% CI = 76-124 days) for the combination (log-rank test p = 5.8E-4 and 3.8E-2 for
combination versus elimusertib or TAZ, respectively; **Figure 5F**). This was most pronounced for
the HYMAD_EPIS_X0004aS1 and SOMWR_EPIS_X00013aS1 PDX models (**Figures 5G-H &**
S15A-D), despite the former exhibiting a relatively poor response to TAZ monotherapy, when
assessed by tumor growth measurements (Vardi *U*-test p = 2.0E-4 for combination versus
elimusertib or TAZ for HYMAD_EPIS_X0003aS1; **Figure 5G**; p = 1.0E-3 and 6.0E-2 for
combination versus elimusertib and TAZ, respectively for SOMWR_EPIS_X00013aS1; **Figure**
S15C) and tumor-free survival (log-rank test p = 6.2E-3 and 6.3E-5 for combination versus
elimusertib or TAZ, respectively for HYMAD_EPIS_X0003aS1; **Figure 5H**; p = 9.5E-5 and 6.0E-
2 for combination versus elimusertib or TAZ, respectively for SOMWR_EPIS_X00013aS1;
Figure S15D). Thus, the combination of EZH2 and ATR inhibition constitutes a synthetic lethal
rational combination strategy to improve TAZ clinical response and overcome resistance.

Finally, we assessed the impact of TAZ treatment on immune-related gene sets, as recent
studies have reported that EZH2 inhibition can increase tumor immunogenicity (61-63). We
found that TAZ treatment upregulated several immune-related gene sets in both $RB1^{WT}$ and
$RB1^{del}$ G401 cells, with 8 of the top 25 upregulated gene sets in all cell clones involving the
immune response (**Supplementary Table S3, Figure S16A-C**). We also found that post-
treatment patient tumor samples which responded to TAZ expressed higher levels of genes

associated with immune-related GO terms (**Figure S16D**). Immune genes upregulated by TAZ
in G401 cells included both MHC-I and MHC-II genes as well as their respective regulators
NLRC5 and CIITA (**Figure S16E-F**), interferon-γ and its receptor (**Figure S16G**), and *CD274*
encoding PD-L1 (**Figure S16G**). Similarly, we observed TAZ induction of multiple endogenous
retroelements (**Figure S16H-I**, **Supplementary Table S6**), and the dsRNA sensor *IFIH1/MDA5*
(**Figure S16G**). Further investigation of EZH2-mediated and TAZ induced antigen presentation
and immune responses of *SMARCB1*-deficient sarcomas is warranted, including the
investigation of the combination of immune checkpoint blockade with epigenetic TAZ
combination therapies.
470
471
472 **Discussion**
473
What defines effective epigenetic EZH2 inhibition therapy for *SMARCB1*-deficient epithelioid
sarcomas and rhabdoid tumors? Our studies of ES patients treated with TAZ clinically
demonstrate that effective inhibition of PRC2 enzymatic activity is necessary but not sufficient
for durable antitumor effects. Based on clinical genomics and transcriptomics, combined with
functional genetic studies of more than 15 diverse MRT and ES cell lines and patient-derived
tumors *in vitro* and *in vivo*, we propose a general molecular model for effective epigenetic TAZ
therapy (**Figure 6**). This model places validated *RB1* and *EZH2* TAZ resistance alleles within
the context of a molecular sequence of events required for clinical TAZ response. This model
also explains additional mutations associated with TAZ resistance based on their perturbation of
each stage of this sequence and provides candidate prognostic biomarkers and therapeutic
combination strategies. We discuss the evidence for this model below, and summarize its novel
predictions and implications.
486
487 First, TAZ must be able to bind and enzymatically inhibit the EZH2 SET domain (**Figure 6; Step**
488 **1**). This inhibition can be blocked by gatekeeper mutations of the EZH2 drug binding site, as
489 observed in lymphoma cell lines (22), and demonstrated for the first time here in an epithelioid
490 sarcoma with clinically acquired $EZH2^{Y666N}$ mutation. Such resistance mutations can be
491 overcome by targeting EED, a non-enzymatic PRC2 subunit.
492
493 For effective epigenetic TAZ therapy, chromatin remodeling complexes must act on tumor
494 suppressor loci that were aberrantly repressed by PRC2 (**Figure 6; Step 2**). The canonical BAF

complex is thought to oppose the activity of the PRC, associated with its chromatin eviction (5, 6). However, the precise mechanism of eviction of TAZ-inhibited PRC2 in *SMARCB1*-deleted tumors is not fully defined. This may involve TAZ-induced remodeling of BAF and/or PRC2. For example, a recent study of *SMARCA4*-deficient cell lines found that upregulation of the expression of BAF helicase *SMARCA2*, which is under PRC2 control in these cells, is necessary for response to EZH2 inhibition (64). In our study, we observed *RB1*-independent upregulation of the expression of BAF subunits *SMARCA2* and *DPF3* upon TAZ treatment in G401 cells (**Figure S17A**). This suggests that TAZ may impact BAF complex assembly as part of its therapeutic mechanism in *SMARCB1*-deficient tumor cells. However, this also indicates that *SMARCA2* re-expression upon EZH2 inhibition is not sufficient to induce cell cycle arrest in the absence of *RB1* expression. After assessing TAZ-induced gene expression changes in PRC2 subunits, we also observed *RB1*-independent TAZ-induced upregulation of PRC2 subunit *JARID2*, and *RB1*-dependent downregulation of *PHF19*, suggesting that PRC2 composition itself may be affected by TAZ treatment (**Figure S10A**). Additionally, it is unknown whether PRC2 eviction in TAZ-treated cells requires a specific form of the BAF complex, such as the non-canonical *SMARCB1*-deficient ncBAF or GBAF complex described previously (65, 66).

If the activity of specific BAF subtypes is indeed needed to evict PRC2 from chromatin upon EZH2 inhibition, then genetic perturbation of specific BAF subunits may impact tumor response to TAZ. For example, in our patient cohort, we observed one TAZ-sensitive tumor with a truncation in the canonical BAF-specific subunit *ARID1B*, while a TAZ-resistant tumor had a missense mutation in PBAF-specific subunit *ARID2* (**Figure 1A**). We also found *ARID1B* to be mutated in 2 out of 5 TAZ-responsive cell lines (0 out of 5 TAZ-sensitive cell lines) and *ARID2* to be mutated in 2 out of 5 TAZ-resistant cell lines, though also in 1 out of 5 TAZ-sensitive lines (**Figure 3B**). It is not known whether these mutations affect tumor response to TAZ, and further work will be needed to elucidate the specific mechanism of chromatin de-repression in TAZ-treated tumor cells, as well as the effects of TAZ on BAF and PRC2 complex assembly.

Effective epigenetic TAZ therapy must also upregulate PRC2-repressed tumor suppressor loci (**Figure 6; Step 3**). In our patient cohort, one TAZ-resistant tumor had deletions of both *CDKN2A* and *CDKN2B* (**Figure 1A**), which can inhibit CDK4/6 from phosphorylating RB1 and are known to be de-repressed by TAZ treatment (11). Our genomic analysis of MRT and ES cell lines also showed that 4 out of 5 TAZ-resistant cell lines tested had apparent loss of *CDKN2A*, with two also having loss of *CDKN2B*, and one having *CDKN1A* loss as well (**Figure 3B**). These

mutations may phenocopy *RB1* loss, suggesting that upregulation of these cell cycle inhibitors

may be necessary for effective TAZ therapy. Indeed, loss of *CDKN1A* or *CDKN2A* is sufficient

to confer resistance to TAZ, similar to loss of *RB1.*

Our findings establishing *CDKN2A* as a TAZ resistance mutation are particularly significant

given the recent genomic studies of epithelioid sarcomas (67, 68), which found recurrent

CDKN2A mutations with loss of p16 expression in more than one third of tumors (67). This also

suggests that epigenetic dysregulation of p16 expression, for example due to repression by

non-EZH2 regulators, may explain the prevalence of TAZ resistance in other *SMARCB1*-

deficient tumors, such as brain ATRT and extracranial rhabdoid tumors.

In addition to maintenance of the expression of these tumor suppressor loci, transcription

factors and coactivators that upregulate their expression must also be intact and expressed for

tumor cells to effectively respond to TAZ (**Figure 6; Step 3**). In our patient cohort, two TAZ-

resistant patient tumors had missense mutations in *ANKRD11* (**Figure 1A**), as did 3 out of 5

TAZ-resistant cell lines. *ANKRD11* is a putative tumor suppressor that exhibits loss of

heterozygosity in breast cancer (69), and is recurrently mutated in other cancers (70, 71).

ANRKD11 can cooperate with p53 to upregulate *CDKN1A*, and previous reports have

suggested a possible risk of cancer development in patients with a constitutional loss of

ANKRD11 (72, 73). We should note that our inability to obtain viable clones with biallelic

ANKRD11 loss, together with the presence of only missense *ANKRD11* mutations in our

samples, suggests a pleiotropic function of this gene. Further work to elucidate the role of

ANKRD11 in TAZ response will require the introduction of specific missense mutations detected

in cell lines and tumors that can separate the cell viability functions from any effects on TAZ

resistance. Interestingly, a recent report described a patient with KBG syndrome, a

developmental condition caused by mutation of *ANKRD11*, who also developed a rhabdoid

tumor (73). The co-occurrence of these two exceedingly rare conditions in the same patient is

consistent with potential functional involvement of *ANKRD11* in rhabdoid tumor development

and susceptibility to EZH2 inhibition.

We also identified *KLF4* as a putative marker of susceptibility to TAZ in patient tumors (**Figure**

3G), and found *KLF4* to be mutated in 1 out of 5 TAZ-resistant cell lines (**Figure 3B**). Although

KLF4 is a transcription factor with both activating and repressing functions, its role in

SMARCB1-deleted tumors is not known. Its expression has long been known to upregulate

CDKN1A, causing cell cycle arrest at the G1/S checkpoint (45, 74). This suggests that KLF4 may regulate the induction of tumor suppressor genes in response to EZH2 inhibition, such as its bona fide target *CDKN1A*. This may occur through recruitment of BAF to tumor suppressor loci; KLF4 can recruit BAF to target genes to upregulate them (75). Like *ANKRD11*, *KLF4* may thus be a key activator of PRC2-repressed genes.

Finally, effective TAZ therapy also requires the function of downstream cell cycle effectors of the relevant tumor suppressor loci (**Figure 6; Step 4**). As we have demonstrated in this study, loss of *RB1* leads to the evasion of TAZ-induced cell cycle arrest, despite effective inhibition of EZH2 activity and sustained transcriptional response to TAZ. A recent genome-wide CRISPR screen also found *RB1* as a top mediator of TAZ resistance (19). This is reminiscent of the necessity of intact *RB1* for therapeutic response to clinical inhibitors of CDK4/6 (76-78). The transcriptional upregulation of hundreds of genes by TAZ in $RB1^{del}$ tumors, including EMT gene sets (**Figure S2C&F**), suggests that these cells are undergoing forced differentiation, even while maintaining proliferation (**Figure 2G-H**). This upregulation of mesenchymal gene sets is consistent with previous work that has shown that PRC2 inhibition, similar to *SMARCB1* re-expression, can drive *SMARCB1*-deficient tumors into a terminally differentiated, mesenchymal-like state (11, 30, 79), possibly recapitulating the normal developmental trajectory of their cells of origin (30). In this way, *RB1* loss and dysregulation of the RB1/E2F axis appear to decouple the regulation of cell fate and identity from its control of cell cycle progression. Although in our patient tumor cohort, only a single *RB1* mutation was detected, recent genomic studies found *RB1* loss to be one of the few co-occurring mutations in primary and metastatic or recurrent epithelioid sarcomas (68). Together with the mutational deletion of CDKN2A, mutation of *ANKRD11*, and epigenetic silencing of p16, this mechanism therefore represents a frequent perturbation of the G1/S checkpoint that accounts for TAZ resistance in 5 out of 16 TAZ-resistant tumor specimens studied, as well as in 4 of 5 TAZ-resistant cell lines. Further work will be needed not only to define the contribution of other mutations and their place within the model we propose here (**Figure 6**), but also on the non-cell autonomous effects of EZH2 inhibition *in vivo*, which may include increased immunogenicity stimulated by TAZ treatment and consequent immunologic tumor clearance *in vivo* (**Figure S16**).

In our search for predictive biomarkers of TAZ response, we identified increased expression of the PCP gene *PRICKLE1* to be associated with a deficient RB1/E2F axis and TAZ resistance (**Figure 3C-F**). The molecular mechanism connecting G1/S dysregulation and *PRICKLE1*

expression is currently unknown, but the PCP pathway is known to be under cell cycle control (80, 81). This is mediated at least in part by PLK1, which we found to be one of the top upregulated genes in TAZ-resistant tumors (**Figure S5B**). Likewise, deletion of the *Drosophila* *RB1* homologue *Rbf1* results in the upregulation of several PCP genes, including the *PRICKLE1* homologue *pk* (82). This suggests that dysregulation of the RB1/E2F axis may lead to upregulation of *PRICKLE1* through dysregulation of normal cell cycle control of PCP. Further work will be needed to define this mechanism, as well as to investigate PRICKLE1 as a clinical biomarker for TAZ resistance.

Finally, these findings enabled two strategies to circumvent clinical TAZ resistance. First, since dysregulation of the RB1/E2F axis mediates escape from cell cycle arrest at the G1/S checkpoint, cell cycle kinases that function downstream of this checkpoint should remain viable therapeutic targets. This cell cycle bypass strategy is supported by previous work showing that loss of *RB1* can sensitize cancer cells to Aurora kinase inhibition through a primed spindle assembly checkpoint (83). As predicted, $RB1^{del}$ cells remain sensitive to CDK2, AURKA, and AURKB inhibition (**Figures 4A, S6A-B**). We found that MRT and ES cell lines resistant to TAZ, including those with *ANKRD11* and *CDKN1A/2A/2B* mutations were sensitive to barasertib. Most compellingly, we found that the combination of TAZ and barasertib exhibits improved antitumor activity *in vivo* compared to either drug alone.

This is reminiscent of the therapeutic combination mechanism proposed by Sorger and Palmer (84), in which inter-patient and inter-tumor variability in response to individual drugs rather than their pharmacological interactions lead to apparent combined effects. We observed substantial benefit of TAZ and barasertib combination within individual PDX models. This suggests that the improved efficacy of this combination may also result from intra-tumor heterogeneity and tumor evolution *in vivo*, as proposed for combination therapy more than fifty years ago (85, 86). Dual targeting of two different parts of the cell cycle can prevent tumors from evading cell cycle arrest through the presence or acquisition of mutations in cell cycle control genes. For example, cells with a defective G1/S checkpoint that continue to transit through the cell cycle remain sensitive to AURKB inhibition. Further work will be needed to elucidate whether barasertib causes postmitotic endoreduplication due to its disruption of the spindle assembly and/or direct inhibition of the mitotic checkpoint (87, 88).

Our findings also advance a synthetic lethal strategy for rational epigenetic TAZ combination
therapy due to TAZ-induced expression of *PGBD5*, the putative developmental mutator in
rhabdoid and other young-onset solid tumors, through its induction of DNA damage and
requirement for ongoing DNA damage repair signaling (57). We found that the DNA damage
repair ATR kinase inhibitor elimusertib not only can overcome RB1/E2F axis-mediated
resistance, but in combination with TAZ, can also exert synergistic anti-tumor effects *in vitro* and
in vivo. Indeed, PGBD5 is required for the induction of DNA damage by the combination of TAZ
and elimusertib, substantiating the specific nature of the synthetic lethal dependency and
therapeutic vulnerability created by TAZ-mediated induction of PGBD5 and its DNA nuclease
activity.

It is possible that the enhanced sensitivity to ATR inhibition due to the increased requirements
for DNA repair also depends on the intrinsic variation in DNA damage repair signaling among
different tumor subtypes. This may also be due to the variation in the expression and activity of
PGBD5 nuclease activity among tumors, both of which may be associated with the recently
described molecular subtypes of ES and MRT tumors (56). Finally, additional synthetic lethal
strategies may also be developed based on the immunologic effects of EZH2 inhibitors,
particularly as combined with epigenetic and synthetic lethal therapies, both of which can
promote tumor immunogenicity. In all, this study develops a paradigm for rational epigenetic
combination therapy, including candidate prognostic biomarkers, all of which should be
incorporated into future clinical trials for patients.

Acknowledgements

This work is dedicated to Maggie Schmidt and her family, and many other patients and their
advocates who inspire and support our research. We thank Richard Koche, Nicholas Socci, and
Mithat Gönen for technical advice, Alejandro Gutierrez and Stuart Orkin for *EZH2* expression
vectors, members of our labs for critical advice and manuscript comments, and Epizyme (now
Ipsen), Bayer and AstraZeneca for supplying tazemetostat, elimusertib, and barasertib,
respectively. This work was supported by the MSK Integrated Genomics Operation Core, Anti-
Tumor Assessment Core, Bioinformatics Core, Molecular Diagnostics Service and the
Department of Pathology, the Marie-Josée and Henry R. Kravis Center for Molecular Oncology,
NIH R01 CA214812, P30 CA08748, T32 GM007739, Burroughs Wellcome Fund, Rita Allen
Foundation, Pershing Square Sohn Cancer Research Alliance and the G. Harold and Leila Y.

Mathers Foundation, Cycle for Survival, MSK Sarcoma Center, the Starr Cancer Consortium,

and Maggie's Mission. Yaniv Kazansky was supported by a Medical Scientist Training Program

grant from the National Institute of General Medical Sciences of the National Institutes of Health

under award number T32 GM007739 to the Weill Cornell/Rockefeller/Sloan Kettering Tri-

Institutional MD-PhD Program. AK is a Scholar of the Leukemia & Lymphoma Society.

Conflict of Interest

The authors have no competing financial interests. AK is a consultant for Novartis, Rgenta,
Blueprint, and Syndax.

Methods

EZH2 mutant plasmids

The $EZH2^{Y666N}$ mutation detected in the clinical trial patient refers to amino acid numbering in

isoform 2 of the protein. For consistency of nomenclature, all engineered mutations use

numbering referring to isoform 2 (Uniprot ID: Q15190-2), although isoform 1 was expressed in

cells for this study. Plasmids containing wild-type EZH2 ($EZH2^{WT}$) and catalytically inactive triple

mutant (F672I, H694A, R732K, referred to as $EZH2^{CatMut}$) plasmids were kindly provided by

Alejandro Gutierrez in the doxycycline-inducible pINDUCER20 vector (89). The plasmids

contain human *EZH2* tagged N-terminally with a FLAG-Avi tag.

The Y666N mutation was engineered in both the $EZH2^{WT}$ and $EZH2^{CatMut}$ plasmids to yield

$EZH2^{Y666N}$ and what we termed $EZH2^{QuadMut}$. The mutation was introduced by site-directed

mutagenesis using the QuikChange Lightning Kit (Agilent) per manufacturer's instructions

(mutagenesis primers: 5'-GCAAAGTGTACGACAAGAACATGTGCAGCTTTCTG-3' and 5'-

CAGAAAGCTGCACATGTTCTTGTCGTACACTTTGC-3') to engineer a TAC to AAC codon

change. After mutagenesis, PCR products were transformed into Stbl3 *E. coli* cells and

expanded at 30°C. Correct plasmid sequences were confirmed by Sanger sequencing (Eton).

See oligonucleotide table below for all sequencing primers used.

Generating CRISPR knockout cells

G401 cells with *RB1* mutations were engineered by Synthego. Briefly, a single guide RNA

targeting exon 2 of *RB1* was used (see oligonucleotide table below for gRNA and primer

sequences). Ribonucleoprotein containing spCas9 and sgRNA was transfected into G401 cells

by electroporation. The target site was then PCR-amplified and Sanger sequenced to ensure

homozygous indels. The cells were then single-cell cloned and re-verified by Sanger

sequencing. Loss of the RB1 protein was confirmed by Western blot.

G401 cells with *CDKN2A* and *CDKN1A* KO were generated in-house, using crRNAs against

exon 1 of *CDKN2A* and exon 2 of *CDKN1A*. Ribonucleoproteins were prepared using crRNA,

ATTO550-labeled tracrRNA, and Alt-R S.p. Cas9 (IDT). Cells were transfected by

electroporation using the Neon transfection system (Invitrogen) with parameters set at 1300 V,

20 ms, 1 pulse Transfected cells were selected by flow sorting and single-cell cloned.

Homozygous indels were confirmed by Sanger sequencing (see table below for gRNA and

primers sequences).

<u>Generating shPGBD5 cells</u>

For shRNA cells, pLKO.1 shRNA vectors targeting *PGBD5* (TRCN0000138412,

TRCN0000135121) and control shGFP were obtained from the RNAi Consortium (Broad

Institute). G401 cells were transduced at an MOI ~1.5 and selected with puromycin at 2 µg/mL

for 72 hours. Knockdown was confirmed by quantitative RT-PCR (see Supplementary Methods)

using primers specified in the oligonucleotide table below.

Oligonucleotides:

gRNAs	
RB1 crRNA	5'- AGAGAGAGCUUGGUUAACUU -3'
CDKN2A crRNA	5'- GACCCTCTACCCACCTGGAT -3'
CDKN1A crRNA	5'- CTTTGTCACCGAGACACCAC -3'
Primers for amplification	
RB1 forward	5'- CACTGTGTGGTATCCTTATTTTGGA -3'
RB1 reverse	5'- AGGTAAATTTCCTCTGGGTAATGGA -3'
CDKN2A forward	5'- ATGGAGCCTTCGGCTGACT -3'
CDKN2A reverse	5'- CCTTCAGATCTTCTCAGCATTCG -3'
CDKN1A forward	5'- GCTGATAATACCTGGTGGCAAA -3'
CDKN1A reverse	5'- CACATGTCCGCACCTGTCAT -3'
PGBD5 forward	5'- CTCTGGGTCAGACAATGTTCTTC -3'
PGBD5 reverse	5'- GCTTATTCTTCAGCGCATCCA -3'
GAPDH forward	5'- GATCATCAGCAATGCCTCCT -3'
GAPDH reverse	5'- GTCATGAGTCCTTCCACGATAC -3'
Primers for sequencing	
EZH2 forward 1	5'- GGACAGCAGAGATCCAGTTTG -3'
EZH2 reverse 1	5'- GGTCCGTTCCAGGATCTTCT -3'
EZH2 forward 2	5'- TCCAGTGTGGTGGAATTCTG -3'
EZH2 reverse 2	5'- TATCGCTGGGGAACTTTCTG -3'
EZH2 forward 3	5'- TGCTGCACAACATCCCTTAC -3'
EZH2 reverse 3	5'- TGCTGGTTTCGTCCTTCTTT -3'
EZH2 forward 4	5'- CCTACAAGCGGAAGAACACC -3'
EZH2 reverse 4	5'- GTTCTTGCTGTCCCAGTGGT -3'

EZH2 forward 5	5'- CTGAAGAAGGATGGCAGCTC -3'
EZH2 reverse 5	5'- CTTGGGTGGGTTACTCCAGA -3'
RB1	5'- CACTGTGTGGTATCCTTATTTTGGA -3'
CDKN2A	5'- GTGGCTCCTCATTCCTCTTCCTTG -3'
CDKN1A	5'- ACCTGAGGTGACACAGCAAAGCCC -3'

Cell cycle analysis

G401 cells were plated on Day 0 and treated for 10 days with 1 µm tazemetostat or equivalent

volume of DMSO, with drug and media replaced on Days 4 and 7. On Day 11, cells were pulsed

with EdU for 1 hour. Cells were then harvested and processed for flow cytometry using the

manufacturer's protocol (Click-iT, Invitrogen). Briefly, cells were washed with PBS with 1% BSA,

permeabilized, and incubated with AlexaFluor647 for 30 minutes. DNA content was measured

using propidium iodide (0.05 µg/µL). Cells were analyzed on a CytoFLEX LX (Beckman

Coulter).

Patient tumor samples

Patient tumor and matched normal blood samples were obtained from patients at Memorial

Sloan Kettering Cancer Center (MSKCC) enrolled in the TAZ clinical trial (15). All patients

provided informed consent for this study under the Institutional Review Board approved

research protocol 12-245. Patient tumors were classified into "Response" or "Progression"

groups based on RECIST 1.1 criteria (90). "Response" included tumors exhibiting a complete

response, partial response, or stable disease. All other tumors were classified under the

"Progression" group. The complete list of tumor samples used and corresponding clinical data

may be found in Supplementary Table 1. This cohort includes both tumor samples that

underwent targeted sequencing with MSK-IMPACT (20) as part of their clinical care at MSKCC

as well as archived tumors that were analyzed for this study. For genomic analysis, DNA was

extracted from either flash frozen tumor samples or formalin-fixed, paraffin-embedded (FFPE)

blocks or slides and samples were processed using the IMPACT468 or IMPACT505 panels

depending on the time of their sequencing (20). The detected mutations and copy number

alterations were obtained from cBioPortal (91, 92) and can be found in Supplementary Table 2.

Oncoprints were generated using Oncoprinter (cBioPortal).

For transcriptomic analysis, archived frozen tumor samples were weighed and up to 20-30mg

were homogenized in RLT buffer, followed by extraction using the AllPrep DNA/RNA Mini Kit

(QIAGEN catalog 80204) according to the manufacturer's instructions. RNA was eluted in 13 µL

nuclease-free water. After RiboGreen quantification and quality control by Agilent BioAnalyzer, 1

µg of total RNA with DV200 percentages varying from 78% to 100% underwent ribosomal

depletion and library preparation using the TruSeq Stranded Total RNA LT Kit (Illumina catalog

RS-122-1202) according to instructions provided by the manufacturer with 8 cycles of PCR.

Samples were barcoded and sequenced using NovaSeq 6000 in a PE150 mode, with the

NovaSeq 6000 S4 Reagent Kit (Illumina). On average, 84 million paired reads were generated

per sample and 70% of the data mapped to mRNA.

Targeted sequencing of cell lines

To assess for the presence of somatic mutations in MRT and ES cell lines, DNA was extracted

using the PureLink Genomic DNA Minikit (Invitrogen) and processed using the IMPACT505

panel as above. Due to the lack of matched normal tissue for cell lines, copy number alterations

were detected using a custom algorithm using circular binary segmentation (93) implemented by

the MSK Bioinformatics core. Code is available on github at:

https://github.com/kentsisresearchgroup/seqCNA_tazemetostat_resistance.

Additional methods may be found in the Supplementary Methods and Figures file.

Data Availability

RNA-seq data from G401 cells can be found at the Gene Expression Omnibus (GEO)

repository, accession number GSE213845. RNA-seq data from patient tumor samples has been

deposited to the Database of Genotypes and Phenotypes (dbGaP), accession number

phs003188.v1.p1. Additional supplementary data files and computational analysis scripts are

available at Zenodo (https://doi.org/10.5281/zenodo.7595037).

References

1. Kadoch C, Crabtree GR. Mammalian SWI/SNF chromatin remodeling complexes and cancer: Mechanistic insights gained from human genomics. Sci Adv. 2015;1(5):e1500447.

2. Versteege I, Sevenet N, Lange J, Rousseau-Merck MF, Ambros P, Handgretinger R, et al. Truncating mutations of hSNF5/INI1 in aggressive paediatric cancer. Nature. 1998;394(6689):203-6.

3. Kim KH, Kim W, Howard TP, Vazquez F, Tsherniak A, Wu JN, et al. SWI/SNF-mutant cancers depend on catalytic and non-catalytic activity of EZH2. Nat Med. 2015;21(12):1491-6.

4. McKenna ES, Sansam CG, Cho YJ, Greulich H, Evans JA, Thom CS, et al. Loss of the epigenetic tumor suppressor SNF5 leads to cancer without genomic instability. Mol Cell Biol. 2008;28(20):6223-33.

5. Wilson BG, Wang X, Shen X, McKenna ES, Lemieux ME, Cho YJ, et al. Epigenetic antagonism between polycomb and SWI/SNF complexes during oncogenic transformation. Cancer Cell. 2010;18(4):316-28.

6. Kadoch C, Williams RT, Calarco JP, Miller EL, Weber CM, Braun SM, et al. Dynamics of BAF-Polycomb complex opposition on heterochromatin in normal and oncogenic states. Nat Genet. 2017;49(2):213-22.

7. Kim KH, Roberts CW. Targeting EZH2 in cancer. Nat Med. 2016;22(2):128-34.

8. Yap DB, Chu J, Berg T, Schapira M, Cheng SW, Moradian A, et al. Somatic mutations at EZH2 Y641 act dominantly through a mechanism of selectively altered PRC2 catalytic activity, to increase H3K27 trimethylation. Blood. 2011;117(8):2451-9.

9. McCabe MT, Graves AP, Ganji G, Diaz E, Halsey WS, Jiang Y, et al. Mutation of A677 in histone methyltransferase EZH2 in human B-cell lymphoma promotes hypertrimethylation of histone H3 on lysine 27 (H3K27). Proc Natl Acad Sci U S A. 2012;109(8):2989-94.

10. Majer CR, Jin L, Scott MP, Knutson SK, Kuntz KW, Keilhack H, et al. A687V EZH2 is a gain-of-function mutation found in lymphoma patients. FEBS Lett. 2012;586(19):3448-51.

11. Knutson SK, Warholic NM, Wigle TJ, Klaus CR, Allain CJ, Raimondi A, et al. Durable tumor regression in genetically altered malignant rhabdoid tumors by inhibition of methyltransferase EZH2. Proc Natl Acad Sci U S A. 2013;110(19):7922-7.

12. McCabe MT, Ott HM, Ganji G, Korenchuk S, Thompson C, Van Aller GS, et al. EZH2 inhibition as a therapeutic strategy for lymphoma with EZH2-activating mutations. Nature. 2012;492(7427):108-12.

13. Konze KD, Ma A, Li F, Barsyte-Lovejoy D, Parton T, Macnevin CJ, et al. An orally bioavailable chemical probe of the Lysine Methyltransferases EZH2 and EZH1. ACS Chem Biol. 2013;8(6):1324-34.

14. Li C, Wang Y, Gong Y, Zhang T, Huang J, Tan Z, et al. Finding an easy way to harmonize: a review of advances in clinical research and combination strategies of EZH2 inhibitors. Clin Epigenetics. 2021;13(1):62.

15. Gounder M, Schöffski P, Jones RL, Agulnik M, Cote GM, Villalobos VM, et al. Tazemetostat in advanced epithelioid sarcoma with loss of INI1/SMARCB1: an international, open-label, phase 2 basket study. Lancet Oncol. 2020;21(11):1423-32.

16. Guerra S, Cichowski K. Targeting Cancer at the Intersection of Signaling and Epigenetics. Annual Review of Cancer Biology. 2019;3(1):365-84.

17. Baylin SB, Jones PA. Epigenetic Determinants of Cancer. Cold Spring Harb Perspect Biol. 2016;8(9).

18. Chi SN, Bourdeaut F, Casanova M, Kilburn LB, Hargrave DR, McCowage GB, et al. Update on phase 1 study of tazemetostat, an enhancer of zeste homolog 2 inhibitor, in pediatric patients with relapsed or refractory integrase interactor 1–negative tumors. Journal of Clinical Oncology. 2022;40(16_suppl):10040-.

19. Drosos Y, Myers JA, Xu B, Mathias KM, Beane EC, Radko-Juettner S, et al. NSD1 mediates
antagonism between SWI/SNF and polycomb complexes and is required for transcriptional
activation upon EZH2 inhibition. Mol Cell. 2022;82(13):2472-89.e8.
20. Cheng DT, Mitchell TN, Zehir A, Shah RH, Benayed R, Syed A, et al. Memorial Sloan
Kettering-Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT): A
Hybridization Capture-Based Next-Generation Sequencing Clinical Assay for Solid Tumor
Molecular Oncology. J Mol Diagn. 2015;17(3):251-64.
21. Bracken AP, Pasini D, Capra M, Prosperini E, Colli E, Helin K. EZH2 is downstream of the
pRB-E2F pathway, essential for proliferation and amplified in cancer. Embo j.
2003;22(20):5323-35.
22. Gibaja V, Shen F, Harari J, Korn J, Ruddy D, Saenz-Vash V, et al. Development of
secondary mutations in wild-type and mutant EZH2 alleles cooperates to confer resistance to
EZH2 inhibitors. Oncogene. 2016;35(5):558-66.
23. Bisserier M, Wajapeyee N. Mechanisms of resistance to EZH2 inhibitors in diffuse large B-
cell lymphomas. Blood. 2018;131(19):2125-37.
24. Brooun A, Gajiwala KS, Deng Y-L, Liu W, Bolaños B, Bingham P, et al. Polycomb
repressive complex 2 structure with inhibitor reveals a mechanism of activation and drug
resistance.
25. Honma D, Kanno O, Watanabe J, Kinoshita J, Hirasawa M, Nosaka E, et al. Novel orally
bioavailable EZH1/2 dual inhibitors with greater antitumor efficacy than an EZH2 selective
inhibitor. Cancer Sci. 2017;108(10):2069-78.
26. He Y, Selvaraju S, Curtin ML, Jakob CG, Zhu H, Comess KM, et al. The EED protein–
protein interaction inhibitor A-395 inactivates the PRC2 complex. Nature Chemical Biology.
2017;13(4):389-95.
27. Qi W, Zhao K, Gu J, Huang Y, Wang Y, Zhang H, et al. An allosteric PRC2 inhibitor
targeting the H3K27me3 binding pocket of EED. Nat Chem Biol. 2017;13(4):381-8.
28. Tomassi S, Romanelli A, Zwergel C, Valente S, Mai A. Polycomb Repressive Complex 2
Modulation through the Development of EZH2–EED Interaction Inhibitors and EED Binders.
Journal of Medicinal Chemistry. 2021;64(16):11774-97.
29. Coe BP, Thu KL, Aviel-Ronen S, Vucic EA, Gazdar AF, Lam S, et al. Genomic deregulation
of the E2F/Rb pathway leads to activation of the oncogene EZH2 in small cell lung cancer.
PLoS One. 2013;8(8):e71670.
30. Custers L, Khabirova E, Coorens THH, Oliver TRW, Calandrini C, Young MD, et al. Somatic
mutations and single-cell transcriptomes reveal the root of malignant rhabdoid tumours. Nat
Commun. 2021;12(1):1407.
31. Kenny C, O'Meara E, Ulaş M, Hokamp K, O'Sullivan MJ. Global Chromatin Changes
Resulting from Single-Gene Inactivation-The Role of SMARCB1 in Malignant Rhabdoid Tumor.
Cancers (Basel). 2021;13(11).
32. Kia SK, Gorski MM, Giannakopoulos S, Verrijzer CP. SWI/SNF mediates polycomb eviction
and epigenetic reprogramming of the INK4b-ARF-INK4a locus. Mol Cell Biol. 2008;28(10):3457-
64.
33. Kotake Y, Cao R, Viatour P, Sage J, Zhang Y, Xiong Y. pRB family proteins are required for
H3K27 trimethylation and Polycomb repression complexes binding to and silencing
p16INK4alpha tumor suppressor gene. Genes Dev. 2007;21(1):49-54.
34. Gallagher D, Voronova A, Zander MA, Cancino GI, Bramall A, Krause MP, et al. Ankrd11 is
a chromatin regulator involved in autism that is essential for neural development. Dev Cell.
2015;32(1):31-42.
35. Neilsen PM, Cheney KM, Li CW, Chen JD, Cawrse JE, Schulz RB, et al. Identification of
ANKRD11 as a p53 coactivator. J Cell Sci. 2008;121(Pt 21):3541-52.

36. Noll JE, Jeffery J, Al-Ejeh F, Kumar R, Khanna KK, Callen DF, et al. Mutant p53 drives
multinucleation and invasion through a process that is suppressed by ANKRD11. Oncogene.
2012;31(23):2836-48.
37. Béguelin W, Rivas MA, Calvo Fernández MT, Teater M, Purwada A, Redmond D, et al.
EZH2 enables germinal centre formation through epigenetic silencing of CDKN1A and an Rb-
E2F1 feedback loop. Nat Commun. 2017;8(1):877.
38. Fan T, Jiang S, Chung N, Alikhan A, Ni C, Lee CC, et al. EZH2-dependent suppression of a
cellular senescence phenotype in melanoma cells by inhibition of p21/CDKN1A expression. Mol
Cancer Res. 2011;9(4):418-29.
39. Negishi M, Wongpalee SP, Sarkar S, Park J, Lee KY, Shibata Y, et al. A new lncRNA,
APTR, associates with and represses the CDKN1A/p21 promoter by recruiting polycomb
proteins. PLoS One. 2014;9(4):e95216.
40. Daulat AM, Finetti P, Revinski D, Silveira Wagner M, Camoin L, Audebert S, et al. ECT2
associated to PRICKLE1 are poor-prognosis markers in triple-negative breast cancer. Br J
Cancer. 2019;120(9):931-40.
41. Daulat AM, Bertucci F, Audebert S, Sergé A, Finetti P, Josselin E, et al. PRICKLE1
Contributes to Cancer Cell Dissemination through Its Interaction with mTORC2. Dev Cell.
2016;37(4):311-25.
42. Jiang D, He Y, Mo Q, Liu E, Li X, Huang L, et al. PRICKLE1, a Wnt/PCP signaling
component, is overexpressed and associated with inferior prognosis in acute myeloid leukemia.
J Transl Med. 2021;19(1):211.
43. Zhou R, Tang Z, Li H, Wang X, Sun Y. PRICKLE1 promotes gastric cancer metastasis by
activating mTOR signaling. Am J Transl Res. 2021;13(5):4266-80.
44. Hu J, Yang Y, Ma Y, Ning Y, Chen G, Liu Y. Proliferation Cycle Transcriptomic Signatures
are Strongly associated With Gastric Cancer Patient Survival. Front Cell Dev Biol.
2021;9:770994.
45. Chen X, Johns DC, Geiman DE, Marban E, Dang DT, Hamlin G, et al. Krüppel-like factor 4
(gut-enriched Krüppel-like factor) inhibits cell proliferation by blocking G1/S progression of the
cell cycle. J Biol Chem. 2001;276(32):30423-8.
46. Chen X, Whitney EM, Gao SY, Yang VW. Transcriptional profiling of Krüppel-like factor 4
reveals a function in cell cycle regulation and epithelial differentiation. J Mol Biol.
2003;326(3):665-77.
47. De Azevedo WF, Leclerc S, Meijer L, Havlicek L, Strnad M, Kim SH. Inhibition of cyclin-
dependent kinases by purine analogues: crystal structure of human cdk2 complexed with
roscovitine. Eur J Biochem. 1997;243(1-2):518-26.
48. Manfredi MG, Ecsedy JA, Chakravarty A, Silverman L, Zhang M, Hoar KM, et al.
Characterization of Alisertib (MLN8237), an investigational small-molecule inhibitor of aurora A
kinase using novel in vivo pharmacodynamic assays. Clin Cancer Res. 2011;17(24):7614-24.
49. Wilkinson RW, Odedra R, Heaton SP, Wedge SR, Keen NJ, Crafter C, et al. AZD1152, a
selective inhibitor of Aurora B kinase, inhibits human tumor xenograft growth by inducing
apoptosis. Clin Cancer Res. 2007;13(12):3682-8.
50. Yang J, Ikezoe T, Nishioka C, Tasaka T, Taniguchi A, Kuwayama Y, et al. AZD1152, a
novel and selective aurora B kinase inhibitor, induces growth arrest, apoptosis, and sensitization
for tubulin depolymerizing agent or topoisomerase II inhibitor in human acute leukemia cells in
vitro and in vivo. Blood. 2007;110(6):2034-40.
51. Nair JS, Ho AL, Tse AN, Coward J, Cheema H, Ambrosini G, et al. Aurora B kinase
regulates the postmitotic endoreduplication checkpoint via phosphorylation of the
retinoblastoma protein at serine 780. Mol Biol Cell. 2009;20(8):2218-28.
52. Vardi Y, Ying Z, Zhang CH. Two-sample tests for growth curves under dependent right
censoring. Biometrika. 2001;88(4):949-60.

53. Henssen AG, Henaff E, Jiang E, Eisenberg AR, Carson JR, Villasante CM, et al. Genomic
DNA transposition induced by human PGBD5. Elife. 2015;4.
54. Helou L, Beauclair L, Dardente H, Piégu B, Tsakou-Ngouafo L, Lecomte T, et al. The
piggyBac-derived protein 5 (PGBD5) transposes both the closely and the distantly related
piggyBac-like elements Tcr-pble and Ifp2. J Mol Biol. 2021;433(7):166839.
55. Bigot Y, Yamada M, Mueller H, Morell V, Alves S, Lecomte T, et al. Analysis of DNA
transposition by DNA transposases in human cells. bioRxiv. 2023:2023.04.26.538406.
56. Henssen AG, Reed C, Jiang E, Garcia HD, von Stebut J, MacArthur IC, et al. Therapeutic
targeting of PGBD5-induced DNA repair dependency in pediatric solid tumors. Science
translational medicine. 2017;9(414).
57. Henssen AG, Koche R, Zhuang J, Jiang E, Reed C, Eisenberg A, et al. PGBD5 promotes
site-specific oncogenic mutations in human tumors. Nat Genet. 2017;49(7):1005-14.
58. Keller R, Yamada M, Cameron D, Suzuki H, Sanghrajka R, Vaynshteyn J, et al. Childhood
cancer mutagenesis caused by a domesticated DNA transposase. bioRxiv.
2023:2022.07.05.498128.
59. Mah LJ, El-Osta A, Karagiannis TC. gammaH2AX: a sensitive molecular marker of DNA
damage and repair. Leukemia. 2010;24(4):679-86.
60. León TE, Rapoz-D'Silva T, Bertoli C, Rahman S, Magnussen M, Philip B, et al. EZH2-
Deficient T-cell Acute Lymphoblastic Leukemia Is Sensitized to CHK1 Inhibition through
Enhanced Replication Stress. Cancer Discov. 2020;10(7):998-1017.
61. Burr ML, Sparbier CE, Chan KL, Chan YC, Kersbergen A, Lam EYN, et al. An Evolutionarily
Conserved Function of Polycomb Silences the MHC Class I Antigen Presentation Pathway and
Enables Immune Evasion in Cancer. Cancer Cell. 2019;36(4):385-401.e8.
62. Mehdipour P, Marhon SA, Ettayebi I, Chakravarthy A, Hosseini A, Wang Y, et al. Epigenetic
therapy induces transcription of inverted SINEs and ADAR1 dependency. Nature.
2020;588(7836):169-73.
63. Ishak CA, Marshall AE, Passos DT, White CR, Kim SJ, Cecchini MJ, et al. An RB-EZH2
Complex Mediates Silencing of Repetitive DNA Sequences. Mol Cell. 2016;64(6):1074-87.
64. Januario T, Ye X, Bainer R, Alicke B, Smith T, Haley B, et al. PRC2-mediated repression of
SMARCA2 predicts EZH2 inhibitor activity in SWI/SNF mutant tumors. Proc Natl Acad Sci U S
A. 2017;114(46):12249-54.
65. Wang X, Wang S, Troisi EC, Howard TP, Haswell JR, Wolf BK, et al. BRD9 defines a
SWI/SNF sub-complex and constitutes a specific vulnerability in malignant rhabdoid tumors. Nat
Commun. 2019;10(1):1881.
66. Michel BC, D'Avino AR, Cassel SH, Mashtalir N, McKenzie ZM, McBride MJ, et al. A non-
canonical SWI/SNF complex is a synthetic lethal target in cancers driven by BAF complex
perturbation. Nat Cell Biol. 2018;20(12):1410-20.
67. Jamshidi F, Bashashati A, Shumansky K, Dickson B, Gokgoz N, Wunder JS, et al. The
genomic landscape of epithelioid sarcoma cell lines and tumours. J Pathol. 2016;238(1):63-73.
68. Dermawan JK, Singer S, Tap WD, Nacev BA, Chi P, Wexler LH, et al. The genetic
landscape of SMARCB1 alterations in SMARCB1-deficient spectrum of mesenchymal
neoplasms. Mod Pathol. 2022;35(12):1900-9.
69. Powell JA, Gardner AE, Bais AJ, Hinze SJ, Baker E, Whitmore S, et al. Sequencing,
transcript identification, and quantitative gene expression profiling in the breast cancer loss of
heterozygosity region 16q24.3 reveal three potential tumor-suppressor genes. Genomics.
2002;80(3):303-10.
70. Vasmatzis G, Johnson SH, Knudson RA, Ketterling RP, Braggio E, Fonseca R, et al.
Genome-wide analysis reveals recurrent structural abnormalities of TP63 and other p53-related
genes in peripheral T-cell lymphomas. Blood. 2012;120(11):2280-9.

71. Gebauer N, Künstner A, Ketzer J, Witte HM, Rausch T, Benes V, et al. Genomic insights
into the pathogenesis of Epstein-Barr virus-associated diffuse large B-cell lymphoma by whole-
genome and targeted amplicon sequencing. Blood Cancer J. 2021;11(5):102.
72. Isrie M, Hendriks Y, Gielissen N, Sistermans EA, Willemsen MH, Peeters H, et al.
Haploinsufficiency of ANKRD11 causes mild cognitive impairment, short stature and minor
dysmorphisms. Eur J Hum Genet. 20. England2012. p. 131-3.
73. Behnert A, Auber B, Steinemann D, Frühwald MC, Huisinga C, Hussein K, et al. KBG
syndrome patient due to 16q24.3 microdeletion presenting with a paratesticular rhabdoid tumor:
Coincidence or cancer predisposition? Am J Med Genet A. 2018;176(6):1449-54.
74. Zhang W, Geiman DE, Shields JM, Dang DT, Mahatan CS, Kaestner KH, et al. The gut-
enriched Kruppel-like factor (Kruppel-like factor 4) mediates the transactivating effect of p53 on
the p21WAF1/Cip1 promoter. J Biol Chem. 2000;275(24):18391-8.
75. Moonen JR, Chappell J, Shi M, Shinohara T, Li D, Mumbach MR, et al. KLF4 recruits
SWI/SNF to increase chromatin accessibility and reprogram the endothelial enhancer landscape
under laminar shear stress. Nat Commun. 2022;13(1):4941.
76. Schoninger SF, Blain SW. The Ongoing Search for Biomarkers of CDK4/6 Inhibitor
Responsiveness in Breast Cancer. Mol Cancer Ther. 2020;19(1):3-12.
77. Finn RS, Dering J, Conklin D, Kalous O, Cohen DJ, Desai AJ, et al. PD 0332991, a selective
cyclin D kinase 4/6 inhibitor, preferentially inhibits proliferation of luminal estrogen receptor-
positive human breast cancer cell lines in vitro. Breast Cancer Res. 2009;11(5):R77.
78. Dean JL, Thangavel C, McClendon AK, Reed CA, Knudsen ES. Therapeutic CDK4/6
inhibition in breast cancer: key mechanisms of response and failure. Oncogene.
2010;29(28):4018-32.
79. Nakayama RT, Pulice JL, Valencia AM, McBride MJ, McKenzie ZM, Gillespie MA, et al.
SMARCB1 is required for widespread BAF complex-mediated activation of enhancers and
bivalent promoters. Nat Genet. 2017;49(11):1613-23.
80. Noatynska A, Tavernier N, Gotta M, Pintard L. Coordinating cell polarity and cell cycle
progression: what can we learn from flies and worms? Open Biol. 2013;3(8):130083.
81. Shrestha R, Little KA, Tamayo JV, Li W, Perlman DH, Devenport D. Mitotic Control of Planar
Cell Polarity by Polo-like Kinase 1. Dev Cell. 2015;33(5):522-34.
82. Payankaulam S, Yeung K, McNeill H, Henry RW, Arnosti DN. Regulation of cell polarity
determinants by the Retinoblastoma tumor suppressor protein. Sci Rep. 2016;6:22879.
83. Gong X, Du J, Parsons SH, Merzoug FF, Webster Y, Iversen PW, et al. Aurora A Kinase
Inhibition Is Synthetic Lethal with Loss of the RB1 Tumor Suppressor Gene. Cancer Discov.
2019;9(2):248-63.
84. Palmer AC, Sorger PK. Combination Cancer Therapy Can Confer Benefit via Patient-to-
Patient Variability without Drug Additivity or Synergy. Cell. 2017;171(7):1678-91.e13.
85. Law LW. Effects of combinations of antileukemic agents on an acute lymphocytic leukemia
of mice. Cancer Res. 1952;12(12):871-8.
86. Frei E, 3rd, Karon M, Levin RH, Freireich EJ, Taylor RJ, Hananian J, et al. The effectiveness
of combinations of antileukemic agents in inducing and maintaining remission in children with
acute leukemia. Blood. 1965;26(5):642-56.
87. Tsuda Y, Iimori M, Nakashima Y, Nakanishi R, Ando K, Ohgaki K, et al. Mitotic slippage and
the subsequent cell fates after inhibition of Aurora B during tubulin-binding agent-induced mitotic
arrest. Sci Rep. 2017;7(1):16762.
88. Hauf S, Cole RW, LaTerra S, Zimmer C, Schnapp G, Walter R, et al. The small molecule
Hesperadin reveals a role for Aurora B in correcting kinetochore-microtubule attachment and in
maintaining the spindle assembly checkpoint. J Cell Biol. 2003;161(2):281-94.
89. Meerbrey KL, Hu G, Kessler JD, Roarty K, Li MZ, Fang JE, et al. The pINDUCER lentiviral
toolkit for inducible RNA interference in vitro and in vivo. Proc Natl Acad Sci U S A.
2011;108(9):3665-70.

90. Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, et al. New
response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). Eur J
Cancer. 2009;45(2):228-47.
91. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer
Discov. 2012;2(5):401-4.
92. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative analysis
of complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal.
2013;6(269):pl1.
93. Olshen AB, Venkatraman ES, Lucito R, Wigler M. Circular binary segmentation for the
analysis of array-based DNA copy number data. Biostatistics. 2004;5(4):557-72.

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044
Figure 1: Patient tumor sequencing reveals diverse tazemetostat resistance mutations.
(**A**) Abridged oncoprint of selected genes from MSK-IMPACT sequencing on patient tumor

samples. Top panel: Tumor samples prior to TAZ treatment. Bottom panel: Matched samples
pre- and post-TAZ or pre- and post-acquisition of resistance. (**B**) Pre- and post- treatment CT
imaging of the indicated patient tumors which acquired *EZH2* and *RB1* mutations. (**C**) Atomic
molecular model of the chimeric *Homo sapiens/Anolis carolinensis* EZH2 bound to pyridone-
based EZH2 inhibitor I (blue), PDB: 5IJ7. Y666 is highlighted in red. (**D**) Top panel: Doxycycline-
inducible *EZH2* is expressed in single-cell G401 clones at near-physiological levels after 3 days
treatment with doxycycline at 1 µg/mL. Numbers indicate clone ID. Bottom panel: Cell viability
measured by CellTiter-Glo after 14 days of treatment with the indicated drug at 10 µM or
equivalent volume of DMSO. n=5 biological replicates per condition. *p = 3.5E-3, **p = 5.2E-3,
***p = 3.8E-5, ****p = 1.1E-5 by two-sided Student's t-test. (**E**) Top panel: *RB1* knockout in two
G401 clones (E1 and F2). Bottom panel: Cell viability measured by CellTiter-Glo after treatment
with 10 µM tazemetostat or DMSO for 14 days. *n*=5 biological replicates per condition. *p =
2.8E-5, **p = 9.0E-5 by two-sided Student's t-test.
1060
1061
1062

Figure 2: *RB1* loss allows escape from cell cycle arrest despite effective EZH2 inhibition.
(**A-C**) Volcano plots of RNA-seq data from TAZ-treated G401 *RB1^WT^* (**A**), *RB1^del^ E1* (**B**), and
RB1^del^ F2 (**C**). Plots show gene expression changes of cells treated with 10 µM TAZ versus
equivalent volume of DMSO for 11 days. n=3 biological replicates per condition. Dots in red
show genes with expression changes of $\log_2$(fold change) > ±1 and *p*-value < 0.01. (**D-F**) GSEA
plots showing the PRC2_EZH2_UP.UP gene set for G401 *RB1^WT^* (**D**), *RB1^del^ E1* (**E**), and *RB1^del^*
F2 (**F**) cells treated in A-C. (**G**) Cell cycle analysis of the indicated G401 clone. Plot shows
triplicate measurements of cells treated with 1 µM TAZ versus equivalent volume of DMSO for
11 days. Y-axis shows the percent difference of the TAZ-treated cells in each cell cycle phase
compared to DMSO-treated cells. *p=2.4E-4, **p=2.3E-4, ***p=5.0E-5, ****p=2.7E-5, #*p*=1.4E-6,
##*p*=1.1E-6. (**H**) Western blot of the indicated G401 cell clones treated with 10 µM TAZ versus
equivalent volume of DMSO for 11 days. Actin serves as loading control.

Figure 3: An intact RB1/E2F axis is a key requirement for response to TAZ: (**A**) Abridged oncoprint of MRT and ES cell lines. Genes included here are BAF subunits and CDK4/6/RB1/E2F axis genes. Full list of mutations is listed in Supplementary Table 4. (**B**) Indicated cell lines treated with TAZ for 11 days. (**C-D**) Volcano plots of RNA-seq data comparing DMSO-treated G401 $RB1^{del}$ E1 (**C**) and F2 (**D**) with $RB1^{WT}$ cells. (**E**) Western blot for PRICKLE1 in untreated G401 cells. (**F-G**) Read counts for *PRICKLE1* (**F**) and *KLF4* (**G**) showing counts normalized by DESeq2 for patient tumor samples collected prior to TAZ treatment. (**H**) Read counts for *AURKB* for patient tumor samples collected after TAZ treatment. *p=0.029, **p=0.027, ***p=0.013.

Figure 4: Cell cycle bypass combination strategy using AURKB inhibition overcomes TAZ resistance and improves response: (**A**) G401 cells treated with barasertib for 6 days (**A**). (**B**) Panel of MRT and ES cell lines ordered left to right by decreasing response to TAZ monotherapy. Cells were treated with the indicated monotherapy or combination for 11 days. Drug concentrations used were: TAZ: 200 nM, barasertib: 8 nM. (**C**) Tumor growth curves showing volumes calculated from caliper measurements for 5 mouse PDXs treated with the indicated drug regimen. n = 20 mice for vehicle and barasertib-treated groups, n = 21 for TAZ and TAZ + barasertib-treated groups. Vardi *U*-test *p* = 4.0E-4 and 2.0E-4 for combination vs. barasertib or TAZ, respectively. (**D**) Kaplan-Meier curves showing tumor-free survival (defined

as tumor volume $\leq$ 1,000 mm^3) for the PDXs in panel C. Mean survival is 65 days (95% CI: 52-
78 days) for barasertib, 68 days (95% CI: 53-82 days) for TAZ, 98 days (95% CI: 84-112 days)
for the combination. Log-rank test p = 3.3E-3 and 7.4E-3 for combination vs. barasertib or TAZ,
respectively. (**E**) Tumor growth curves for the subset of mouse tumors in panel C from the
SOMWR_EPIS_X00013aS1 PDX model. n = 3 mice per treatment group. Vardi U-test p = 0.10
and 9.4E-2 for combination vs. barasertib or TAZ, respectively. (**F**) Tumors from panel E
harvested on Day 135 of treatment.

1105

Figure 5: Synthetic lethal combination strategy using ATR inhibition overcomes TAZ
resistance and improves response: (**A**) G401 cells treated with elimusertib for 4 days. (**B**)
Panel of MRT and ES cell lines ordered left to right by decreasing response to TAZ

monotherapy. Cells were treated with the indicated monotherapy or combination for 11 days.
Drug concentrations used were: TAZ: 200 nM, elimusertib: 8 nM. We selected an elimusertib
dose below its monotherapy IC_{50} (for G401 cells) in order to visualize any additive effects upon
combination with TAZ. *p = 0.14, **p = 3.3E-3, ***p = 1.0E-3, ***p = 2.6E-4, #p = 6.1E-5, ##p =
3.0E-529 by two-sided Student's t-test. All comparisons refer to TAZ + elimusertib vs.
elimusertib conditions, except for G401 cells, in which comparison is for TAZ + elimusertib vs.
TAZ. n = 5 replicates per condition. (**C**) Representative images of G401 cells treated with the
indicated treatment for 7 days. Elimusertib was added on Day 5. Doses used were 500 mM TAZ
and 100 nM elimusertib. (**D**) Quantification of γH2AX fluorescence relative to DAPI
fluorescence. *p = 9.3E-3, **p = 6.2E-14, ***p = 1.5E-29 by two-sided Student's t-test. n = 332
nuclei for DMSO, 404 for TAZ, 400 for elimusertib, 257 for TAZ + elimusertib. (**E**) Tumor growth
curves for 5 mouse PDXs treated with the indicated drug regimen. n = 20 mice for vehicle and
elimusertib-treated groups, n = 21 for TAZ and TAZ + elimusertib-treated groups. Vardi *U*-test p
= 2.3E-2 and 0.20 for combination vs. elimusertib or TAZ, respectively. (**F**) Kaplan-Meier curves
showing tumor-free survival (defined as tumor volume ≤ 1,000 mm^3) for the PDXs in panel C.
Mean survival is 51 days (95% CI: 42-60 days) for elimusertib, 68 days (95% CI: 53-82 days) for
TAZ to 100 days (95% CI: 74-124 days) for the combination. Log-rank test p = 5.8E-4 and 3.8E-
2 for combination versus elimusertib or TAZ, respectively. (**G**) Tumor growth curves for the
HYMAD_EPIS_X0004aS1 PDX model treated with the indicated drug regimen. Vardi *U*-test p =
2.0E-4 for combination versus elimusertib or TAZ. n = 14 mice per treatment group. (**H**) Kaplan-
Meier curves showing tumor-free survival (defined as tumor volume ≤ 1,000 mm^3) for the PDXs
in panel E. Log-rank test p = 6.2E-3 and 6.3E-5 for combination versus elimusertib or TAZ,
respectively.

1132
1133

1134
1135 **Figure 6: Mechanistic schematic for the response of BAF-deficient tumors to effective**
1136 **EZH2 therapy: Step 1**: TAZ inhibits histone methylation activity of PRC2. **Step 2:** Activating

1137 chromatin-bound complexes, such as ncBAF bind to tumor suppressor loci. **Step 3:** Tumor
1138 suppressor loci are activated by their transcription factors and their coactivators. **Step 4:** Tumor
1139 suppressors inhibit cell cycle progression through their downstream effectors, such as RB1/E2F.

1181 **Supplementary Methods and Figures**

1182

1183 **for**

1184

1185 **Kazansky et al**

1186

1187 **Overcoming clinical resistance to EZH2 inhibition using rational epigenetic combination**
1188 **therapy**

1189

1190

1191 This file includes:

1192

1193 Supplementary methods
1194 Supplementary table captions
1195 Supplementary figures S1-S9
1196 Additional references

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

Supplementary methods

<u>Cell culture</u>

All cell lines were obtained from the American Type Culture Collection if not otherwise specified.
ES1 and ES2 cells were generated and kindly provided by Nadia Zaffaroni. EPI544 cells were
obtained from the MD Andersen Cancer Center Cytogenetics and Cell Authentication Core.
Rhabdoid tumor cell lines KP-MRT-NS, KP-MRT-RY, and MP-MRT-AN were kindly provided by
Yasumichi Kuwahara and Hajime Hosoi. The identity of all cell lines was verified by STR
analysis. Absence of *Mycoplasma* contamination was determined using the MycoAlert kit
according to manufacturer's instructions (Lonza). Cell lines were cultured in 5% CO_2 in a
humidified atmosphere in 37°C. All media were obtained from Corning and supplemented with
10% fetal bovine serum (FBS), 1% L-glutamine, and 100 U/mL penicillin and 100 µg/mL
streptomycin (Gibco). RPE, G401, A204, ES1, ES2, and VAESBJ cells were cultured in
Dulbecco's Modified Eagle Medium (DMEM). TTC642, TM8716, MP-MRT-AN, KP-MRT-NS,
and KP-MRT-RY cells were cultured in Roswell Park Memorial Institute (RPMI) medium. EPI544
cells were cultured in DMEM/F12 medium.

1243

<u>Western Blotting</u>

To assess protein expression by Western immunoblotting, pellets of 1 million cells were
prepared and washed once in cold PBS. Cells were resuspended in 100-130 µL of RIPA lysis
buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1.0% NP-40, 0.5% sodium deoxycholate, 0.1%
sodium dodecyl sulfate) and incubated on ice for 10 minutes. Cell suspensions were then
disrupted using a Covaris S220 adaptive focused sonicator for 5 minutes (peak incident power:
35W, duty factor: 10%, 200 cycles/burst) at 4 °C. Lysates were cleared by centrifugation at
18,000 g for 15 min at 4 °C. Protein concentration was assayed using the DC Protein Assay
(Bio-Rad) and 15-35 µg whole cell extract was used per sample. Samples were boiled at 95 °C
in Laemmli buffer (Bio-Rad) with 40 mM DTT and resolved using sodium dodecyl sulfate-
polyacrylamide gel electrophoresis. Proteins were transferred to Immobilon FL PVDF
membranes (Millipore), and membranes were blocked using Intercept Blocking buffer (Li-Cor).
Primary antibodies used were: anti-EZH2 (Cell Signaling Technology, 5246) at 1:1,000, anti-
RB1 (Cell Signaling Technology, 9309) at 1:250, anti-H3K27me3 (Cell Signaling Technology,
9733) at 1:500, anti-p16 (Abcam, ab108349) at 1:500, anti-CCNA2 (Santa Cruz, sc-271682) at
1:100, anti-PRICKLE1 (Santa Cruz, sc-393034) at 1:100, anti-SMARCB1 (BD Biosciences,
612110) at 1:500, anti-RPA32 pT21 (abcam, ab109394) at 1:2,000, anti-RPA32 pS4/pS8
(ThermoFisher, A300-245A) at 1:2,000, anti-pCHK1 S296 (Cell Signaling Technology, 90178) at

1:250, anti-MYCN (Cell Signaling Technology, 9405) at 1:250, anti-MMP2 (Cell Signaling

Technology, 40994S) at 1:250, anti-cleaved Caspase 3 (Cell Signaling Technology, 9661S) at

1:500, anti-Actin (Cell Signaling Technology, 4970 and 3700) at 1:5,000. Blotted membranes

were visualized using goat secondary antibodies conjugated to IRDye 680RD or IRDye 800CW

(Li-Cor, 926-68071 and 926-32210) at 1:15,000 and the Odyssey CLx fluorescence scanner,

according to manufacturer's instructions (Li-Cor). Image analysis was done using the Li-Cor

Image Studio software (version 4).

1269

<u>Lentivirus production</u>

Lentivirus production was carried out as described previously (1). Briefly, HEK293T cells were

transfected using TransIT-LT1 using a 2:1:1 ratio of the lentiviral vector and psPAX2 and

pMD2.G packaging plasmids, according to manufacturer's instructions (Mirus). Viral supernatant

was collected at 48 and 72 hours post-transfection, pooled, filtered and stored in aliquots at -80

°C. G401 cells were transduced at a multiplicity of infection (MOI) of 0.3. Transduced cells were

selected for 7 days with G418 sulfate (ThermoFisher) at 1 mg/mL. Single-cell clones were then

isolated and expanded. Inducible EZH2 expression was confirmed by Western blotting against

EZH2.

1279

1280   <u>Cell Viability Testing</u>

1281   Drugs used for *in vitro* treatment were supplied by Selleckchem (TAZ; S7128, Elimusertib;

1282   S9864, abemaciclib; LY2835219, palbociclib; S1116, seleciclib; S1153, alisertib; S1133,

1283   barasertib; S1147, camptothecin, S1288).

1284

The effects of *RB1* loss or *EZH2* mutation on TAZ susceptibility was assessed over 14 days.

Cells were plated in 96-well microplates at equal densities and treated with 10 µM TAZ or

equivalent volume of DMSO on Day 0. Drug and media were replaced on Days 4, 7, and 11.

CellTiter-Glo assays were performed on Day 14, with luminescence readings taken using an

automated fluorescence plate reader (Tecan). CellTiter-Glo reagent was freshly reconstituted on

the day of measurement and added in a 1:1 proportion to cell media. A similar protocol was

used for all other cell viability experiments, with treatment times indicated in the relevant figure

legends. Cell line doubling time was determined by measuring cell viability every 24 hours over

the course of 4 days, and fitting the cell viability to a two-parameter exponential curve. For

combination treatment with TAZ and elimusertib, we used a two-dimensional dose matrix

design, treating the cells for 9 days. After the addition of cells, drugs were added using a pin tool

(stainless steel pins with 50 nL slots, V&P Scientific) mounted onto a liquid handling robot

(CyBio Well vario, Analytik Jena). For analysis of synergy, we used the synergyFinder package

(2). Outliers due to pinning errors were excluded after manual examination.

1299

Cell line RNA-sequencing

G401 cells with or without *RB1* loss were plated and treated with 10 μM TAZ or equivalent

volume of DMSO on Day 0. Drug and media were replaced on Days 4 and 8. Cells were

harvested on Day 11 and RNA was isolated using RNeasy Mini kit, according to manufacturer's

instructions (Qiagen). After RiboGreen quantification and quality control by Agilent BioAnalyzer,

149-500ng of total RNA underwent Poly(A) selection and TruSeq library preparation according

to instructions provided by Illumina (TruSeq Stranded mRNA LT Kit, catalog RS-122-2102), with

8 cycles of PCR. Samples were barcoded and sequenced using a HiSeq 4000 instrument using

50bp/50bp paired end mode, using the HiSeq 3000/4000 SBS Kit (Illumina). An average of 42

million paired reads was generated per sample. Ribosomal reads represented less than 0.03%

of the total reads generated and proportion of mRNA bases averaged 74%.

1311

Analysis of RNA-seq data

For RNA-seq analysis of G401 cell lines, read adaptors were trimmed and quality filtered using

'trim_galore' (v0.4.4_dev) and mapped to GRCh38/hg19 reference genome using STAR v2.6.0a

with default parameters (3). Read counts tables were generated using HTSeq (4). Normalization

was performed using DESeq2 using the default parameters (5). Differential gene expression

analysis of transposable elements (TEs) was performed using the TEtranscripts (6) package

with default parameters, with GTF files for TE annotations for hg19 generated by the Hammell

lab using UCSC Repeat Masker.

1320

1321

For RNA-seq analysis of patient tumor samples, read adaptors were trimmed and quality filtered

using 'trim_galore' and mapped to GRCh38/hg19 reference genome using STAR v2.7.9 with

default parameters (3). Read count tables were generated using HTSeq v0.11.3 (4). Bam files

were sorted by name using 'samtools' and alignment quality was assessed using 'qualimap'

v2.2.2. Normalization was performed using DESeq2 v1.34.0 using the default parameters (5).

To assess gene expression changes between TAZ-sensitive and TAZ-resistant tumors, samples

in both categories were compared by two-tailed Student's t-test using 'rowttests' in R v4.1.3.

Genes were filtered by $p < 0.05$ and sorted by t-statistic. Heatmaps were then generated using

'pheatmap.' Genome browser tracks were visualized from bam files using Integrated Genomics

Viewer v2.13.1.

1332

1333 Gene ontology analysis

1334 Genes significantly up- or down-regulated in TAZ-sensitive and TAZ-resistant tumors

1335 determined by two-tailed Student's t-test at p-value < 0.5 were searched against the Gene

1336 Ontology database (DOI: 10.5281/zenodo.5725227 Downloaded 2021-11-16).

1337

Microscopy

Bright field microscopy was performed using an Evos FL Auto 2 imager at 10x magnification,

with cells grown on plastic dishes. Immunofluorescence for γH2AX was performed on cells

plated on Millicell EZ Slide glass slides (EMD Millipore), coated for 45 minutes with bovine

plasma fibronectin (Millipore Sigma). After drug treatment, cells were washed once with PBS

and fixed in 4% formaldehyde for 10 minutes at room temperature. Slides were then washed

three times in PBS for 5 minutes, permeabilized for 15 minutes in 0.3% Triton X-100, washed

again in PBS three times, and blocked with 5% goat serum (Millipore Sigma, G9023) in PBS for

1 hour at room temperature. Slides were incubated with mouse anti-γH2A.X primary antibody

(Sigma-Aldrich, 05-636) at 1:500 in blocking buffer for 1 hour, washed three times in PBS, and

incubated with goat anti-mouse secondary antibody conjugated to AlexaFluor555 (Invitrogen, A-

21422) at 1:1,000. Cells were then counterstained with DAPI at 1:1,000 for 10 minutes and

treated with ProLong Diamond Antifade Mountant with DAPI (Invitrogen, P36962) for 48 hours.

For MMP2 and cleaved caspase 3 immunofluorescence, cells were processed as above, using

anti-MMP2 antibody (Cell Signaling Technology, 40994S) at 1:200 or anti-cleaved Caspase 3

antibody (Cell Signaling Technology, 9661S) at 1:300, and Phalloidin conjugated to

AlexaFluor488 (ThermoFisher, A12379) at 1:400 added to the secondary antibody mix.

1355

Images were acquired on a Zeiss LSM880 confocal microscope at 63x magnification. Images

were then processed using a custom pipeline in CellProfiler (7). Per-cell integrated γH2A.X

intensity was normalized against per-cell integrated DAPI intensity. All image analysis used the

same pipeline settings, with the exception of the RescaleIntensity module for the AF555

channel, which used the settings 0.009-0.09 for the images in Figure 5 and 0.005-0.09 for

Figure S13. Overlaid images were prepared using Fiji (8).

1362
1363
1364

<u>Xenografts</u>

All mouse experiments were carried out in accordance with institutionally approved animal use

protocols. To generate PDXs, tumor specimens were collected under approved IRB protocol 14-

091, immediately minced and mixed (50:50) with Matrigel (Corning, New York, NY) and

implanted subcutaneously in the flank of 6-8 weeks-old female NOD.Cg-$Prkdc^{scid}$ $Il2rg^{tm1Wjl}$/Szj

(NSG) mice (Jackson Laboratory, Bar Harbor, ME), as described previously (9). Mice were

monitored daily and PDX samples were serially transplanted three times before being deemed

established. PDX tumor histology was confirmed by review of H&E slides and direct comparison

to the corresponding patient tumor slides. PDX identity was further confirmed by MSK-IMPACT

sequencing analysis.

1375

Therapeutic studies used female and male NSG mice obtained from the Jackson Laboratory.

Xenografts were prepared as single-cell suspensions, resuspended in Matrigel, and implanted

subcutaneously into the right flank of 6-10 week old mice. 100 μL of tumor cell suspension was

used for each mouse. Tumors were allowed to grow until they reached a volume of 100 mm^3, at

which point they were randomized into treatment groups without blinding. Drugs were prepared

using the following formulations: Tazemetostat was dissolved at 25 mg/mL in 5% DMSO, 40%

PEG 300, 5% Tween 80, and 50% water. Elimusertib was dissolved at 5 mg/mL in 10% DMSO,

40% PEG 300, 5% Tween 80, and 45% water using a sonicator. Barasertib was dissolved at 2.5

mg/mL in 5% DMSO, 40% PEG 300, 5% Tween 80, and 50% water. Drugs were reconstituted

daily. The following drug doses and schedules were used: TAZ was dosed at 250 mg/kg twice

daily by oral gavage, 7 days per week. Barasertib was dosed at 25 mg/kg once daily by

intraperitoneal injection using 3 days on and 4 days off cycle. Elimusertib was dosed at 40

mg/kg twice daily by oral gavage using 2 days on and 12 days off cycle. Caliper tumor

measurements were taken twice weekly. Tumor volumes were calculated using the formula

Volume = (π/6) x length x width^2. Tumor growth analysis was performed using the Vardi U-test

(10), as implemented in the clinfun R package using the aucVardiTest function. Tumor-free

survival analysis was calculated using OriginPro (Microcal) by the Kaplan-Meier method, using

the log-rank test.

1394

<u>Immunohistochemistry</u>

The immunohistochemistry detection of Ki67, Cl-caspase3, p-H2AX, p16, p21, and H3K27me3

was performed at Molecular Cytology Core Facility of Memorial Sloan Kettering Cancer Center

using Discovery Ultra processor (Ventana Medical Systems). For all markers with the exception

of H3K27me3, paraffin-embedded tissues were sectioned at 5 μm and baked at 58°C for 1 hr.

Slides were loaded in Ultra autostainer and IHC staining was performed as follows. Samples

were baked and dewaxed followed by pretreatment with epitope retrieval CC1 solution

(Ventana, 950-500) for 48 min at 100°C. The primary antibodies (rabbit monoclonal Ki67

antibody (1ug/ml, Cell Signaling Technology, 9027), rabbit anti- Cleaved-caspase3 antibody

(0.2ug/ml, Cell Signaling Technology, 9661), rabbit anti-p-H2AX (0.2ug/ml, Abcam, ab11174),

rabbit anti p21 (0.8ug/ml, Cell Signaling Technology, 2947), and mouse anti-p16 antibody

(0.2ug/ml,Santa Cruz,sc56330) were incubated for 4h at room temperature. Samples were then

incubated with either OmniMap anti-rabbit HRP (Roche Diagnostics, 760-4311) secondary

antibody or OmniMap anti-mouse HRP (Roche Diagnostics760-4310) for 20 minutes. Discovery

ChromoMap DAB detection kit (Roche Diagnostics) was used according to the manufacturer's

instructions.

For H3K27me3 staining, slides were loaded into Leica Bond RX and dewaxed at 72°C before

being pretreated with EDTA-based epitope retrieval ER2 solution (Leica, AR9640) for 20 mins.

at 100°C. The Rabbit anti-H3K27me3 (0.2ug/ml, Cell Signaling Technology, 9733) were applied

for 60 mins. Samples were then incubated with Leica Bond Polymer (anti-rabbit HRP) (included

in Polymer Refine Detection Kit (Leica, DS9800) for another 8 min. Slides were then incubated

with mixed DAB reagent (Polymer Refine Detection Kit) for 10 mins, followed by hematoxylin

(Refine Detection Kit) counterstaining for 10 mins.

After staining, all sample slides were washed in water, dehydrated using an ethanol gradient

(70%, 90%, 100%), washed three times in HistoClear II (National Diagnostics, HS-202), and

mounted in Permount (Fisher Scientific, SP15).

1421

1422

1423

Supplementary Table S1: List of patient tumor specimens used for RNA-seq and MSK-
IMPACT analysis. *We note that two primary tumors in patients who responded to TAZ
harbored deletions of *RB1* (patient 2, sample ES_02_T_02) in one tumor and *CDKN2A/B* in
another tumor (patient 5, sample ES_05_T_01). However, these primary tumors were fully
resected prior to the initiation of TAZ treatment and did not recur at the primary sites. In the
case of patient 2, a later TAZ-responsive metastasis (ES_2_T_03) did not harbor the *RB1* loss.
In the case of patient 5, a later TAZ-responsive metastasis (ES_05_T_09) did not harbor the
CDKN2A/B loss. This suggests that these mutations were subclonal and were not present in
tumors exposed to TAZ treatment. Thus, the mutations in these tumors were unlikely to have
impacted their response to TAZ.
1443
1444 **Supplementary Table S2:** List of mutations found in all patient tumor specimens in
1445 Supplementary table 1 for which MSK-IMPACT data is available.
1446
1447 **Supplementary Table S3:** List of Hallmark gene sets identified by Gene Set Enrichment
1448 Analysis (GSEA) up- and down-regulated by TAZ treatment. Related to Supplementary Figure
1449 S2C.
1450
1451 **Supplementary Table S4:** List of mutations found in all MRT and ES cell lines used in this
1452 study as determined by targeted MSK-IMPACT sequencing. Related to Figure 3A.
1453
1454 **Supplementary Table S5:** List of PDX models used in this study, with clinical characteristics of
1455 the original tumor specimens, followed by a list of mutations found in all PDX models, as
1456 determined by targeted MSK-IMPACT sequencing.
1457
1458 **Supplementary Table S6:** List of endogenous transposable elements whose expression is up-
1459 or downregulated by TAZ treatment. Related to Supplementary Figure S16

Figure S1: Validation of tumor resistance mutations of *EZH2*: (**A-B**) Integrated Genome Viewer (IGV) tracks of RNA-seq data for post-treatment tumor samples from patients 3 and 15. Purple boxes indicate the mutation sites. (**A**) Aligned reads and coverage plot of exon 16 of *EZH2*, showing expression of mRNA with the T (red) → A (green) mutant allele. (**B**) Aligned reads and coverage plot of exon 3 of *RB1*, showing expression of mRNA with frame shift deletion at I124. (**C**) Phase-contrast microscopy of G401 single-cell clones expressing the indicated form of *EZH2*. Cells were treated with 10 μM tazemetostat or DMSO for 9 days and imaged with an Evos FL Auto 2 imager at 10X magnification. Arrow indicates a refractile, mitotic cell. Arrowhead indicates a post-treatment, morphologically altered cell. (**D**) Cell viability measured by CellTiter-Glo after 14 days of treatment with 10 μM tazemetostat or equivalent volume of DMSO. Data is the same as in Figure 1D with the addition of *EZH2^CatMut^* and *EZH2^QuadMut^* clones. n=5 replicates per condition. (**E**) Cell viability after treatment with 10 μM valemetostat or DMSO for 14 days. n=5 replicates per condition.

1485
1486
Figure S2: *RB1^{del}* cells show morphological and transcriptional responses to TAZ: (**A**) Phase-contrast
microscopy of G401 cells with or without *RB1* expression. Cells were treated with 10 µM tazemetostat or DMSO for 9
days and imaged with an Evos FL Auto 2 imager at 10X magnification. Arrow indicates a refractile, mitotic cell.
Arrowhead indicates a post-treatment, morphologically altered cell. (**B**) Western blot of the indicated G401 cell clones
treated with 10 µM TAZ vs. equivalent volume of DMSO for 11 days. Bulk H3K27me3 levels are reduced in all three
clones despite persistent EZH2 expression in *RB1^{del}* clones. (**C**) Comparison of all Hallmark gene sets upregulated in
G401 cells upon TAZ treatment with significance at FDR < 25%. (**D**) GSEA plots showing the
Hallmark_Epithelial_Mesenchymal_Transition gene set for the indicated TAZ-treated G401 cells compared to DMSO.

1495

1496
1497
1498 **Figure S3: TAZ-treated *RB1^del* cells show evidence of differentiation at the transcript, but not protein level.** (**A**)
1499 RNA-seq data from cells in Figure 2A-F, showing normalized read counts for the *MMP2* gene. (**B**) Indicated cells
1500 treated with 1 µM TAZ for 11 days, stained with MMP2 antibody. U2OS cells are shown as a positive control. (**C**)
1501 MMP2 western blot on cells treated as in (B).

1513
1514
1515 **Figure S4: *RB1^del* cells show increased expression of E2F targets.** (**A-B**) GSEA plots showing the
1516 Hallmark_E2F_Targets gene set comparing TAZ-treated *RB1^del* G401 cells with *RB1^WT* cells.
1517

Figure S5: Characterization of MRT and ES cell lines: (**A**) Confirmed loss of SMARCB1 expression in all MRT and ES cell lines used. (**B**) Dose-response curves of a panel of MRT and ES cell lines treated with TAZ for 13 days. Curves in red correspond to TAZ-responsive cell lines, grey to TAZ-resistant cell lines. *n=3* replicates per condition. (**C**) Plot of area under the curve (AUC) of the dose-response curves from Figure 3A plotted against doubling time. AUC was integrated from 40 nM to 50 µM (**D**) IGV track of reads from MSK-IMPACT sequencing of KP-MRT-RY cells, focusing on the *RB1* gene, shows reads spanning segments across the full gene and indicates the presence of at least one allele.

1530
1531 **Figure S6: Generating and testing *CDKN1A^del^* and *CDKN2A^del^* G401 cells.** (**A**) Indicated cell lines were treated
1532 with 10 μM TAZ or DMSO for 11 days. (**B-C**) Sanger traces of the indicated G401 clones for *CDKN2A^del^* (**B**) and
1533 *CDKN1A^del^* (**C**) showing homozygous editing. Red arrows indicate crRNA sequences. (**D**) Indicated G401 *CDKN2A^del^*
1534 cell clones treated with 10 μM TAZ or DMSO for 7 days, confirming loss of CDKN2A induction. (**E**) Dose response
1535 curves of the indicated cell lines treated with TAZ for 11 days. *n=3* replicates per condition.

1536
1537 **Figure S7: TAZ-resistant patient tumors show upregulation of cell cycle genes:** Top 30 GO terms, sorted by p-
1538 value, enriched in pre-treatment (**A**) and post-treatment (**B**) patient tumor specimens that did not respond to TAZ,
1539 compared to those that did. (**C-D**) DESeq2-normalized read counts of genes from the indicated GO terms comparing
1540 pre-treatment TAZ-responding tumors to pre-treatment non-responding tumors.
1541

1542
1543
Figure S8: Transcriptomic analysis of patient tumors nominates putative biomarkers of TAZ sensitivity and
resistance: (**A**) Venn diagrams of genes up- or down-regulated by *RB1* knockout in the indicated clone. Same data
as in Figure 2. (**B-C**) Heatmaps showing the top 60 upregulated (**B**) and downregulated (**C**) genes in patient tumors
collected prior to TAZ-treatment. Each column indicates a separate tumor. Heatmaps are sorted by t-statistic
calculated using a two-tailed Student's t-test, $p<0.05$. Red boxes indicate planar cell polarity genes *CELSR2*, *PLK1*,
and *PRICKLE1* and *CDKN1A* regulator *KLF4*.

1550

1551

1552

1553

1554

1555

1556

1557

1558

1559 **Figure S9: Downstream cell cycle inhibitors overcome resistance to TAZ:** G401 cells treated with seleciclib (**A**),

1560 alisertib (**B**), abemaciclib (**C**), or palbociclib (**D**) 9 days. n=3 replicates per condition.

1561

50 μm

1562
Figure S10: TAZ + barasertib increases cell cycle arrest without inducing apoptosis. (**A**) EdU incorporation
measured by flow cytometry for the indicated cell lines and treatments. Y-axis shows percent change vs. DMSO
treatment for each cell line. TAZ dose was 1 μM for 11 days, with barasertib added at 12 nM or equivalent volume of
DMSO added for the last 3 days. *p=1.4E-3 , **p=5.3E-6 , ***p=2.2E-9 . (**B-C**) Western blot (**B**) and
immunofluorescence (**C**) of G401 and A204 cells treated as indicated, with drug doses and times as in (**A**).
Staurosporine treatment for 6 h at 0.5 μM was used as a positive control. (**D**) Tumor growth curves for the subset of
mouse tumors in Figure 4C from the HYMAD_EPIS_X0003aS1 PDX model. n = 2 mice for vehicle and barasertib-
treated groups, n = 3 mice for TAZ and TAZ + barasertib-treated groups.
1571

1572
Figure S11: p16 induction correlates with TAZ response *in vivo*. (**A**) Kaplan-Meier curves showing tumor-free
survival (defined as tumor volume ≤ 1,000 mm^3) for the indicated PDXs. HYMAD_EPIS_X0004aS1; mean survival is
34 days (95% CI: 29-40 days) for vehicle and 36 days (95% CI: 32-39 days) for TAZ. Log-rank test *p* = 0.63.

SOMWR_EPIS_X00013aS1; mean survival is 44 days (95% CI: 37-51 days) for vehicle and 84 days (95% CI: 76-92
days) for TAZ. Log-rank p= 7.1E-6. KUNGA_MRT_X0002aS1; mean survival is 38 days (95% CI: 30-46 days) for
vehicle and 58 days (95% CI: 47-69 days) for TAZ. Log-rank p = 8.0E-3. (**B**) Representative IHC images of PDX
tissue after treatment with vehicle vs. TAZ. (**C**) Quantification of IHC images. 3 fields for 3 tumors were quantified for
each condition.

1586
1587
1588 **Figure S12: TAZ induces DNA damage in G401 cells** (**A-B**) Synergy plots for combination treatment with TAZ and
1589 elimusertib for (**A**) G401 and (**B**) ES1 cells. Cells were treated at the indicated doses for 9 days and analyzed for
1590 synergy using the Zero Interaction Potency (ZIP) m**odel.** (**C**) Representative uncropped images of G401 cells treated
1591 as indicated (same treatment as in Figure 5D). 2 fields are shown per condition.
1592

1593
Figure S13: TAZ-induced DNA damage is PGBD5-dependent: (**A**) RT-qPCR showing *PGBD5* expression vs.
GAPDH in G401 cells with the indicated shRNA. *p = 1.6E-2, **p = 3.8E-3. (**B**) Quantification of γH2AX fluorescence
relative to DAPI fluorescence in all nuclei. *p = 8.6E-3, **p = 3.9E-8, ***p = 4.4E-8, ****p = 1.1E-12, # p = 4.9E-34, ##
p = 4.0E-47 by two-sided Student's t-test. (**C**) Quantification of γH2AX fluorescence relative to DAPI fluorescence in
nuclei with punctate γH2AX staining. *p = 3.6E-2, **p = 2.7E-5, ***p = 2.4E-5, ****p = 1.1E-15, #p = 5.6E-51, ##p =
1.7E-81. For B-C, n for shGFP cells is 539 for DMSO, 321 for TAZ, 491 for elimusertib, 418 for TAZ + elimusertib. n
for shPGBD5-1 is 497 for DMSO, 390 for TAZ, 354 for elimusertib, 277 for TAZ + elimusertib. n for shPGBD5-3 is 703
for DMSO, 418 for TAZ, 554 for elimusertib, 317 for TAZ + elimusertib. (**D**) Proportion of nuclei with pan-nuclear
γH2AX staining per field. Each dot represents one field. *p = 9.2E-2, **p = 1.6E-2, ***p = 9.0E-4 n for shGFP cells is
22 fields for DMSO, 22 for TAZ, 22 for elimusertib, 28 for TAZ + elimusertib. n for shPGBD5-1 is 22 for DMSO, 22 for
TAZ, 22 for elimusertib, 20 for TAZ + elimusertib. n for shPGBD5-3 is 22 for DMSO, 22 for TAZ, 20 for elimusertib, 22
for TAZ + elimusertib.
1606
1607
1608

1609
1610
1611
Figure S14: CHK1 inhibition does not induce replication stress or synergize with TAZ: (**A**) Dose-response
curves of G401 cells treated with the CHK1 inhibitor SRA737 for 9 days. (**B**) Western blot assaying replication stress
as measured by RPA phosphorylation at S4/8 and T21. Camptothecin treatment (1.5 µM) for 2 h was used as a
positive control for replication stress. Autophosphorylation of CHK1 at S296 was used to confirm CHK1 inhibition.
Cells were pre-treated with 10 µM TAZ or DMSO for 9 days. Cells were then split and additionally treated with
SRA737 (3 µM) or equivalent volume of DMSO for 2 days. (**C**) Cells treated with 10 µM TAZ or DMSO for 11 days do
not express MYCN protein. *MYCN*-amplified neuroblastoma cell line IMR5 was used as a positive control for MYCN
expression.
1620

1621
1622
Figure S15: Response of individual PDX models to combination therapy: (**A**) Image of representative tumors
extracted from mice in Figure 5G and on Day 52 of treatment (**B**) their corresponding weights. *p = 6.7E-3 by two-
sided Student's t-test. (**C**) Tumor growth curves for SOMWR_EPIS_X00013aS1 PDX model. Vardi U-test p = 1.0E-3
and 6.0E-2 for combination versus elimusertib and TAZ, respectively. n = 8 mice for all groups. (**D**) Kaplan-Meier
curves showing tumor-free survival (defined as tumor volume ≤ 1,000 mm^3) for SOMWR_EPIS_X00013aS1 PDX
model; Mean survival is 90 days (95% CI: 83-97 days) for combination versus 80 days (95% CI: 70-90 days) for TAZ
and 45 days (95% CI: 37-52 days) for elimusertib. Log-rank test p = 9.5E-5 and 6.0E-2 for combination versus
elimusertib and TAZ, respectively.
1631

Figure S16: Induction of immune-related genes and endogenous transposable elements by TAZ: (**A-B**)
Examples of immune-related gene sets induced by TAZ in G401 cells with and without *RB1*. Data is the same as that

in Figure 2. (**C**) Venn diagrams comparing immune-related gene sets in $RB1^{wt}$ and $RB1^{del}$ cells. (**D**) Top GO terms, sorted by p-value, enriched in post-treatment patient tumor specimens that did responded to TAZ, compared to those that did not. (**E-G**) DESeq2-normalized gene expression for the indicated genes in TAZ or DMSO-treated G401 cells. Data is the same as that shown in Figure 2. (**H**) Venn diagrams comparing endogenous transposable elements up- and down-regulated by TAZ in G401 cells.

1642

1643
1644
1645 **Figure S17: TAZ may remodel BAF and PRC2 composition by transcriptional regulation of their subunits:** (**A**)
1646 DESeq2-normalized read counts for all BAF and PRC2 subunits showing significantly altered gene expression
1647 between TAZ and DMSO-treated cells. Same data as in Figure 2.
1648
1649
1650
1651

Supplementary References

1. Kentsis A, Reed C, Rice KL, Sanda T, Rodig SJ, Tholouli E, et al. Autocrine activation of the MET receptor tyrosine kinase in acute myeloid leukemia. Nat Med. 2012;18(7):1118-22.

2. Ianevski A, Giri AK, Aittokallio T. SynergyFinder 2.0: visual analytics of multi-drug combination synergies. Nucleic Acids Res. 2020;48(W1):W488-w93.

3. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.

4. Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput sequencing data. Bioinformatics. 2015;31(2):166-9.

5. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

6. Jin Y, Tam OH, Paniagua E, Hammell M. TEtranscripts: a package for including transposable elements in differential expression analysis of RNA-seq datasets. Bioinformatics. 2015;31(22):3593-9.

7. Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A. CellProfiler 4: improvements in speed, utility and usability. BMC Bioinformatics. 2021;22(1):433.

8. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.

9. Mattar M, McCarthy CR, Kulick AR, Qeriqi B, Guzman S, de Stanchina E. Establishing and Maintaining an Extensive Library of Patient-Derived Xenograft Models. Front Oncol. 2018;8:19.

10. Vardi Y, Ying Z, Zhang CH. Two-sample tests for growth curves under dependent right censoring. Biometrika. 2001;88(4):949-60.