Overcoming clinical resistance to EZH2 inhibition using rational epigenetic combination therapy

Yaniv Kazansky[1,2], Daniel Cameron[1,2], Phillip Demarest[1,2], Nadia Zaffaroni[3], Noemi Arrighetti[3], Valentina Zuco[3], Yasumichi Kuwahara[4], Rui Qu[5], Elisa De Stanchina[5], Filemon Dela Cruz[6], Andrew Kung[6], Mrinal Gounder[7], Alex Kentsis[1,2,8]*

[1] Molecular Pharmacology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[2] Tow Center for Developmental Oncology, Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[3] Molecular Pharmacology Unit, Department of Experimental Oncology, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Milan, Italy

[4] Department of Biochemistry and Molecular Biology, Kyoto Prefectural University of Medicine

[5] Antitumor Assessment Core, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[6] Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[7] Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[8] Departments of Pediatrics, Pharmacology, and Physiology & Biophysics, Weill Medical College of Cornell University, New York, NY, USA

* Correspondence: Alex Kentsis, MD, PhD, email kentsisresearchgroup@gmail.com, telephone +1-646-888-2593

Running title: Overcoming resistance to EZH2 inhibition

Keywords: epigenetics, EZH2 inhibition, BAF, combination therapy, PRC2

Abstract

Essential epigenetic dependencies have become evident in many cancers. Based on functional antagonism between BAF/SWI/SNF and PRC2 in *SMARCB1*-deficient sarcomas, we and colleagues recently completed the clinical trial of the EZH2 inhibitor tazemetostat, leading to its FDA approval. However, the principles of tumor response to epigenetic therapy in general, and tazemetostat in particular, remain unknown. Using functional genomics of patient tumors and diverse experimental models, we define molecular mechanisms of tazemetostat resistance in *SMARCB1*-deficient sarcomas and rhabdoid tumors. We found distinct classes of acquired mutations that converge on the RB1/E2F axis and decouple EZH2-dependent differentiation and cell cycle control. This allows tumor cells to escape tazemetostat-induced G1 arrest despite EZH2 inhibition, and suggests a general mechanism for effective EZH2 therapy. Thus, we develop combination strategies to circumvent tazemetostat resistance using cell cycle bypass and synthetic lethal targeting, and provide prospective biomarkers for therapy stratification. This offers a paradigm for rational epigenetic combination therapy suitable for immediate translation to clinical trials for epithelioid sarcomas, rhabdoid tumors, and other epigenetically dysregulated cancers.

Significance: Genomic studies of patient tumors and cell lines identify mutations converging on a common pathway that is essential for response to EZH2 inhibition. Resistance mutations decouple drug-induced differentiation from cell cycle control. We identify complementary epigenetic combination strategies to overcome resistance and improve response, supporting their investigation in clinical trials.

Introduction

Unlike conventional cytotoxic chemotherapy, epigenetic therapy offers the ability to target cancer-specific dependencies with increased specificity and reduced toxicity. This is especially true for cancers caused by genetic mutations of transcriptional regulators, such as the chromatin remodeling BAF/SWI/SNF (Brg/Brahma-associated factors) complex that is mutated in more than 20% of human cancers (1). For example, malignant rhabdoid tumors (MRT) and epithelioid sarcomas (ES) are lethal tumors of children and young adults caused by inactivating mutations of *SMARCB1/SNF5/INI1/BAF47*, one of the core BAF/SWI/SNF complex subunits (2-4). Loss of normal BAF function can confer a dependency on the Polycomb Repressive Complex 2 (PRC2) and its methyltransferase EZH2. This dependency results from epigenetic antagonism between the two complexes, in which normal BAF activity evicts PRC2 from tumor suppressor loci (5, 6). This discovery has led to the development of multiple inhibitors of EZH2 (7-9), several of which have now entered clinical trials (10). However, despite the potential of such targeted epigenetic therapies, the principles governing their effective application remain unknown. Intrinsic and acquired resistance limits their use as monotherapies (11, 12).

A recent clinical trial led to the FDA approval of the EZH2 methyltransferase inhibitor tazemetostat (TAZ) as the first targeted therapy for *SMARCB1*-deficient epithelioid sarcomas (13). However, only 15% of patients showed objective clinical responses, with most epithelioid sarcomas being resistant to TAZ. Ongoing clinical trials in patients with rhabdoid tumors have shown similar results, with only a subset of brain rhabdoid tumor patients exhibiting objective responses to TAZ, while extracranial rhabdoid tumors appear to be uniformly resistant (14). Recent studies have nominated the histone methyltransferase NSD1 as a regulator of TAZ susceptibility in rhabdoid tumor cells, but the clinical relevance of this mechanism is currently not known (15). In all, the outcomes for most patients with MRT and ES remain dismal, and the mechanisms of clinical response and resistance to EZH2 inhibition remain unknown. This hinders our ability to stratify treatment using prospective biomarkers to identify patients who may benefit from TAZ and to develop effective combination therapies with improved and durable benefits for patients.

Here, we define the key requirements for effective epigenetic therapy in diverse *SMARCB1*-deficient epithelioid sarcomas and rhabdoid tumors *in vivo*. Using comparative genomic analyses of clinical trial patients treated with TAZ, we identify multiple acquired mutations that

cause therapy resistance. Using functional genomic approaches, we show that resistance

mechanisms converge on a common RB1/E2F axis that integrates control of tumor cell division

and differentiation. This organizes patient resistance mutations into a general framework,

allowing us to develop rational combination therapies to circumvent TAZ resistance. Using

diverse patient-derived ES and MRT cell lines and mouse xenografts, we demonstrate cell cycle

bypass and synthetic lethal treatment strategies suitable for immediate translation to

combination clinical trials for patients.

Results

Patient tumor sequencing reveals diverse resistance mutations

To identify mutations associated with clinical resistance to TAZ, we performed targeted gene

sequencing of patient tumors using MSK-IMPACT, which is based on a panel of over 500 genes

recurrently mutated in diverse cancer types (16). We analyzed 33 tumor specimens from 20

patients treated as part of the recent TAZ clinical trial (13), and identified somatic tumor

mutations in matched pre- and post-treatment specimens (**Supplementary Tables 1 and 2**).

We found distinct sets of somatic mutations in responding and non-responding tumors, with

nearly all mutations, apart from *SMARCB1* loss itself, being exclusive to either TAZ-responsive

or TAZ-resistant tumors (**Figure 1A, top panel**). Strikingly, we observed two tumors which

initially responded to TAZ based on radiographic imaging but later developed clinical resistance

(**Figure 1A, bottom panel, 1B**). Targeted sequencing of the resistant tumors revealed two

newly acquired somatic mutations: homozygous missense mutation of *EZH2* ($EZH2^{Y666N}$) in the

Patient 3 tumor specimen, and biallelic loss of function mutation of *RB1*, including a hemizygous

deletion and a frame shift mutation ($RB1^{del}$) in the remaining allele, in the Patient 15 tumor

specimen. We confirmed both mutations using RNA-seq of the respective tumor specimens

(**Figure S1A-B**). Since one mutation affected *EZH2* directly, and the other involved the known

EZH2 target *RB1* (17), we hypothesized that both mutations were responsible for TAZ

resistance in their respective patients.

First, we investigated the $EZH2^{Y666N}$ mutation. Past studies using forward genetic screens in

lymphoma cell lines have identified putative resistance mutations within both the N-terminal D1

domain and the catalytic SET domain of EZH2, both of which are predicted to interact with S-

adenosyl methionine (SAM)-competitive EZH2 inhibitors such as TAZ (18, 19). One such SET

domain mutation previously identified in cell lines is $EZH2^{Y661D}$, with Y661 corresponding to
Y666 in isoform 2 of $EZH2$, the isoform referred to in this study, and thus concordant with the
mutation we observed in the Patient 3 tumor (18).

Based on the atomic resolution structure of a pyridone-based EZH2 inhibitor bound to the *Anolis*
carolinensis EZH2 (PDB 5IJ7) (20), we reasoned that residue Y666 in human EZH2 may form a
critical part of the TAZ binding site and that its mutation can prevent TAZ from binding to the
SET domain (**Figure 1C**). To test $EZH2^{Y666N}$ as a resistance allele, we expressed doxycycline-
inducible $EZH2^{Y666N}$ in $SMARCB1$-deficient G401 rhabdoid tumor cells. We observed that
$EZH2^{Y666N}$ expressing clones are resistant to TAZ as compared to cells expressing equal levels
of wild-type $EZH2$ by assessing both cell viability (**Figure 1D**) and cell morphology (**Figure**
S1C). The resistance phenotype of $EZH2^{Y666N}$-expressing cells depends on the intact catalytic
activity of the SET domain, as a compound mutant combining the catalytically inactive triple
mutant $EZH2^{F672I,H694A,R732K}$ ($EZH2^{CatMut}$) with the Y666N mutation, termed $EZH2^{QuadMut}$, did not
confer resistance to TAZ (**Figure S1D**). We also observed that $EZH2^{Y666N}$ confers resistance to
the dual EZH1/2 inhibitor valemetostat (21), consistent with putative resistance to SAM-
competitive, pyridone-based EZH2 inhibitors (**Figure S1E**). This also suggests that combined
inhibition of EZH2 and EZH1 may not overcome this type of acquired EZH2 inhibitor resistance.

Previous studies found that lymphoma cells resistant to EZH2 inhibitors remained susceptible to
the inhibition of the non-enzymatic PRC2 subunit EED (22), including those with mutations in
the $EZH2$ SET domain ($EZH2^{C663Y}$ and $EZH2^{Y726F}$) and the D1 domain (19, 23). We therefore
hypothesized that TAZ resistance conferred by $EZH2^{Y666N}$ could be overcome by PRC2
inhibitors that do not bind to EZH2. Indeed, we found that the allosteric EED inhibitor MAK683
overcomes $EZH2^{Y666N}$-mediated resistance (24), demonstrating that these cells remain generally
susceptible to PRC2 inhibition (**Figure 1D**). EED inhibition may thus be an effective strategy to
overcome acquired TAZ resistance mutations in $EZH2$.

***RB1* loss allows escape from cell cycle arrest despite effective EZH2 inhibition**

Past work has shown that $EZH2$ is a direct target of repression by RB1/E2F (17, 25). This
suggests that acquired $RB1$ loss observed upon TAZ treatment may confer resistance to EZH2
inhibition by increasing EZH2 expression. To test $RB1^{del}$ as a TAZ resistance allele, we used
CRISPR/Cas9 genome editing to generate biallelic $RB1^{del}$ mutations in G401 cells, as compared

to isogenic *RB1*-wild type control cells produced by targeting the safe harbor locus *AAVS1*. We

confirmed absence of RB1 protein expression in two independent clones using Western blotting,

and found that *RB1^{del}* cells were indeed resistant to TAZ (**Figure 1E**).

Despite *EZH2* being a known target gene of RB1/E2F, we were surprised to observe that *RB1^{del}*

G401 cells showed similar morphological changes upon TAZ treatment (**Figure S2A**) to those

previously reported for TAZ-treated *RB1^{WT}* G401 cells (7). To define the effects of TAZ on *RB1^{del}*

cells more precisely, we performed RNA-seq of isogenic G401 *RB1^{del}* and wildtype *AAVS1*-

control cells, treated with either 10 μM TAZ or DMSO control for 11 days, based on an

established treatment regimen to model EZH2 inhibition *in vitro* (7). As predicted, we observed

that *EZH2* mRNA and protein levels remained high in TAZ-treated *RB1^{del}* cells, unlike in *RB1^{WT}*

cells (**Figure 2A-C, S2B**). However, EZH2 inhibition induced significant upregulation of

hundreds of genes in both *RB1^{WT}* and *RB1^{del}* cells, including upregulation of known PRC2 target

genes (**Figure 2A-F**).

Importantly, trimethylation of the EZH2 substrate H3K27 was substantially reduced by TAZ

regardless of *RB1* status (**Figure S2B**). This indicates that despite persistent EZH2 expression,

EZH2 methyltransferase activity is effectively inhibited by TAZ despite *RB1* loss. Gene set

enrichment analysis (GSEA) showed that all three clones significantly upregulated multiple gene

sets upon TAZ treatment (**Supplementary Table S3**, **Figure S2C**). This included gene sets

associated with cell differentiation such as epithelial-to-mesenchymal transition (EMT; **Figure**

S2F). This is reminiscent of recent observations that re-expression of *SMARCB1* in G401 cells

can lead to a mesenchymal chromatin state (26), and is consistent with the idea that PRC2

inhibition may allow BAF to re-activate a more developmentally normal gene expression state.

Taken together, these findings indicate that *RB1* loss-induced TAZ resistance is independent of

EZH2.

In addition to *EZH2*, additional RB1/E2F target genes were also upregulated in TAZ-treated

RB1^{del} cells compared to TAZ-treated *RB1^{WT}* control cells (**Figure S2D-E**). Given the function of

RB1 in the regulation of the G1/S cell cycle checkpoint, we hypothesized that *RB1* loss could

allow cells to escape TAZ-induced cell cycle arrest. Flow cytometry cell cycle analysis showed

that G401 cells treated with 1 μM TAZ arrest at the G1/S checkpoint, as reported previously (7).

However, *RB1^{del}* cells exhibited a significant reduction in the proportion of cells in G1 phase

upon TAZ treatment (50% of TAZ-treated *RB1^{WT}* cells versus 31% and 28% for *RB1^{del}* E1 and

F2 clones, respectively), with a corresponding increase of the proportion of cells remaining in S and G2/M phases (**Figure 2G**). In agreement with this, we observed persistent mRNA expression of S/G2/M-phase-associated *CCNA2*, *CDK2*, and *AURKB* genes in $RB1^{del}$ cells upon TAZ treatment (**Figure 2A-C**). This is despite upregulation of the CDK4/6 inhibitor *CDKN2A* (p16), a known PRC2 target in MRT (5, 27, 28), upon TAZ treatment in both $RB1^{WT}$ and $RB1^{del}$ cells (**Figure 2A-C**). We confirmed persistent maintenance of S-phase cyclin A2 (CCNA2) protein levels despite p16 upregulation using Western blotting (**Figure 2H**). Together, these results show that *RB1* loss is sufficient to evade TAZ-induced cell cycle arrest at the G1/S restriction point despite maintaining the expected global transcriptional response to EZH2 inhibition, including upregulation of cell cycle inhibitor genes.

Intact RB1/E2F axis is required for TAZ susceptibility

The requirement for *RB1* expression in the therapeutic response to TAZ suggests that an intact RB1/E2F axis may be a general requirement for effective EZH2 inhibitor therapy. This would predict that other genetic and epigenetic perturbations to the RB1/E2F axis, beyond *RB1* loss itself, would similarly confer escape from TAZ-induced cell cycle arrest. Analysis of our TAZ clinical trial treatment cohort revealed one patient tumor specimen with primary resistance to TAZ with intact *RB1* but inactivating mutations of both *CDKN2A* and *CDKN2B* (**Figure 1A**, **Supplementary Tables 1-2**), both of which are known to inhibit CDK4/6-mediated phosphorylation of RB1. Two additional specimens had missense mutations in *ANKRD11*: One tumor with primary resistance to TAZ and another which initially responded but later progressed on treatment, at which point a newly acquired *ANKRD11* mutation was detected (**Figure 1A**-Patients 9 and 16, respectively). *ANKRD11* is a known TP53 cofactor and putative tumor suppressor that contributes to TP53-mediated expression of pan-CDK inhibitor *CDKN1A* (29-31). *CDKN1A* itself is also known to be a PRC2 target in tumors (32-34), although its role in the response to EZH2 inhibition in *SMARCB1*-deficient sarcomas is currently unknown. These results converge on the dysregulation of the RB1/E2F axis as a mechanism of evasion of TAZ-induced cell cycle arrest.

To investigate the functional determinants of tumor cell response to TAZ, we first analyzed the response to TAZ of seven MRT and four ES cell lines in which we confirmed loss of SMARCB1 protein expression in all ES and MRT cell lines using Western blotting, as compared to SMARCB1-expressing HEK293T cells (**Figure S3A**). We classified each line as sensitive or

resistant based on the area under the curve (AUC) of their TAZ dose responses (AUC > 0.3 for sensitive G401, KP-MRT-NS, TTC642, A204, TM8716, KP-MRT-RY cell lines and AUC ≤ 0.3 for resistant ES1, VAESBJ, ES2, EPI544, MP-MRT-AN cell lines; **Figure 3A**). Given that TAZ treatment requires at least 4 days for the cellular reduction of methylated EZH2 substrates and at least 7 days for apparent antiproliferative effects in the rapidly-dividing G401 cell line (7), we confirmed that the apparent TAZ susceptibilities of these MRT and ES cell lines are not correlated with their proliferation rates (Pearson's *r* = -0.073 and *p* = 0.81; **Figure S3B**). In agreement with somatic mutations affecting the RB1/E2F axis associated with TAZ resistance in clinical tumor specimens (**Figure 1A**), we found mutations of *CDKN2A* in 4 out of 5 TAZ-resistant cell lines, *CDKN2B* in 2 out of 5 resistant cell lines, *CDKN1A* in 1 out of 5, and *ANKRD11* in 3 out of 5*,* as compared to no such mutations in any TAZ-sensitive MRT and ES cell lines (**Figure 3B**, **Supplementary Table 4**). While our analysis detected reduction in copy number of *RB1* in TAZ-sensitive KP-MRT-RY cells, manual inspection of sequencing reads within the *RB1* gene revealed lack of homozygous deletion, with presumed retention of *RB1* expression (**Figure S3C**). Thus, mutations associated with the RB1/E2F axis are associated with resistance to TAZ in *SMARCB1*-deficient cell lines and patient tumors.

We next assessed the apparent transcriptional activity of the RB1/E2F axis in patient tumors using quantitative gene expression analysis of TAZ responding and non-responding tumors biopsied before and after TAZ treatment using RNA-seq (**Supplementary Table S1**). Pre-treatment TAZ-resistant tumors exhibited significant enrichment of multiple Gene Ontology (GO) terms associated with the cell cycle and in particular with the S and G2/M phases (**Figure S4A**). Similarly, post-treatment tumors that progressed on TAZ showed increased gene expression of GO terms associated with mitosis, as compared to TAZ-responsive tumors (**Figure S4B**). Indeed, TAZ-resistant tumors exhibited consistently higher expression of S/G2/M-phase-associated genes prior to treatment (**Figure S4C-D**). These findings suggest that in addition to the mutations affecting the RB1/E2F axis associated with TAZ resistance, additional mutations not captured by MSK-IMPACT targeted gene sequencing and/or epigenetic dysregulation, such as putative silencing of tumor suppressor genes like *CDKN1A* or *CDKN2A*, likely contribute to TAZ resistance and the decoupling of RB1/E2F-mediated proliferation and PRC2-regulated differentiation.

Since TAZ-resistant MRT and ES cell lines and patient tumors show distinct mutations and gene expression changes that converge on the RB1/E2F axis, we inquired whether these

perturbations would similarly converge on common prognostic biomarkers of TAZ resistance.
Comparative gene expression analysis of untreated $RB1^{del}$ cells versus $RB1^{WT}$ G401 cells
showed a small set of consistently and significantly up- and down-regulated genes in two
independent clones (**Figure S5A**). The most substantially and significantly upregulated gene
associated with *RB1* loss was *PRICKLE1* (**Figure 3C-D**), which we confirmed to be
overexpressed at the protein level in both $RB1^{del}$ clones using Western blotting (**Figure 3E**).
260
In agreement with this, we also found that *PRICKLE1* was among the most differentially
expressed genes between 10 pre-treatment patient tumors with response and resistance to
TAZ, with *PRICKLE1* expression being higher in TAZ-resistant tumors (mean normalized reads
= 10,237 and 300 for resistant and responsive tumors, respectively; Student's t-test p = 0.013;
Figure 3F, S5A). *PRICKLE1* can control planar cell polarity (PCP), a key cell differentiation
pathway, and has previously been implicated as a prognostic biomarker of poor prognosis in
breast cancer (35, 36), acute myeloid leukemia (37), and gastric cancer (38, 39). Several other
genes encoding PCP pathway factors were also upregulated in TAZ-resistant tumors compared
to TAZ-sensitive tumors (**Figure S5A**). We next looked for differentially expressed genes in
TAZ-sensitive tumors as potential markers of sensitivity. These included the transcription factor
KLF4 which can control the G1/S transition by regulating *CDKN1A* expression (40, 41) (mean
normalized reads = 200 and 2,489 for resistant and responsive tumors, respectively; Student's-
test p = 0.03; **Figure S5B, 3G**). Thus, PRICKLE1 and additional factors controlling PCP and
integration of RB1/E2F cell cycle and differentiation are potential prognostic pre-treatment
biomarkers to identify clinical TAZ resistance and susceptibility of *SMARCB1*-deficient tumors.
276
277 **Synthetic lethal and cell cycle bypass epigenetic combination strategies overcome**
278 **tazemetostat resistance**
279
Given that $RB1^{del}$ cells are able to bypass cell cycle arrest at the G1/S checkpoint, we reasoned
that inhibiting cell cycle kinases downstream of this checkpoint could overcome TAZ resistance.
In particular, cell cycle kinases *CDK2* and *AURKB*, which are downregulated by EZH2 inhibition
in TAZ-sensitive cells but persistently expressed in TAZ-resistant cells, may offer especially
compelling therapeutic targets to overcome TAZ resistance (**Figure 2A-C**). Indeed, we found
that the CDK2 inhibitor seleciclib (42), as well as the mitotic kinase Aurora A inhibitor alisertib
(43), and Aurora B inhibitor barasertib (44), were able to overcome TAZ resistance in $RB1^{del}$
G401 cells (**Figure 4A, Figure S6A-B**). We term this combination strategy cell cycle bypass.

Consistent with the function of CDK4/6 kinases upstream of RB1/E2F, sensitivity to CDK4/6 inhibitors palbociclib and abemaciclib was reduced by $RB1^{del}$ mutation (**Figure S6C-D**). In support of the cell cycle bypass strategy for TAZ combination therapy, we observed that patient tumors which progressed on TAZ showed higher expression of $AURKB$ mRNA as compared to those that responded (**Figure 3H**). Combined with the high sensitivity of G401 cells to barasertib (**Figure 4A**; half-maximal effective concentration of 6.5 ± 0.5 nM, 5.5 ± 0.6 nM, and 5.9 ± 0.9 nM for $RB1^{WT}$, $RB1^{del}$ E1, and F2 clones, respectively), these findings suggest that the cell cycle bypass strategy may effectively overcome TAZ resistance.

We therefore asked whether the combination of TAZ and barasertib would have activity against both TAZ-responsive and TAZ-resistant $SMARCB1$-deficient MRT and ES cell lines, using RPE cells as a $SMARCB1$-proficient control. The effects of the combination of TAZ and barasertib on cell viability did not substantially exceed the effect of barasertib alone at the doses tested (**Figure 4B**; 200 nM TAZ, 8 nM barasertib). However, nearly all cell lines tested, including those resistant to TAZ monotherapy, showed substantial susceptibility to barasertib, with RPE cells displaying the lowest sensitivity (**Figure 4B**).

To test the effects of this combination *in vivo*, we treated a panel of five patient-derived rhabdoid tumor and epithelioid sarcoma xenografts (PDX) in immunodeficient mice (**Supplementary Table S5**), comparing TAZ and barasertib monotherapies with their combination. Importantly, in contrast to the modest reduction of tumor growth and extension of survival of mice with tumors <1,000 mm^3 with TAZ or barasertib alone, PDX mice treated with the combination of TAZ and barasertib showed significant reductions in tumor growth (Vardi U-test p = 4.0E-4 and 2.0E-4 for combination versus barasertib or TAZ, respectively; **Figure 4C**) (45). In two of the PDX models, this combination led to tumor regressions (**Figures 4E-F & S8A**). Consistent with this benefit, the combination was also found to significantly increase mean tumor-free animal survival from 65 days (95% confidence interval (CI) = 51-78 days) for barasertib and 67 days (95% CI = 53-81 days) for TAZ to 98 days (95% CI = 84-112 days) for the combination (log-rank test p = 3.3E-3 and 5.8E-3 for combination versus barasertib or TAZ, respectively; **Figure 4D**). These results indicate that the combination of TAZ with a downstream cell cycle inhibitor such as barasertib can improve response and overcome resistance to TAZ in diverse rhabdoid tumors and epithelioid sarcomas *in vivo*.

In addition to distinct cell cycle dynamics of TAZ resistance, we observed that regardless of *RB1*

status, TAZ treatment also caused significant increase in expression of *PiggyBac transposable*

element derived 5 (*PGBD5)* (**Figure 2A-C**). *PGBD5* is a transposase-derived gene with retained

nuclease activity in human cells, which has been implicated as a somatic mutator and inducer of

double-strand DNA (dsDNA) breaks in childhood solid tumors (46-48). In rhabdoid tumors in

particular, PGBD5 was observed to induce sequence-specific mutations and DNA

rearrangements, including somatic deletions of *SMARCB1* itself (46). In turn, *PGBD5*

expression was both necessary and sufficient to confer a cellular dependency on end-joining

DNA repair and ATR kinase signaling (47).

TAZ-induced upregulation of *PGBD5* expression suggests that TAZ treatment may potentiate

this synthetic lethal dependency. To test this idea, we used the ATR-selective kinase inhibitor

elimusertib, which is currently undergoing clinical trials in patients with solid tumors, including

patients with PGBD5-expressing tumors such as MRT and ES (Clinical Trials Identifier

NCT05071209). We found that elimusertib exhibited low-nanomolar potency against $RB1^{WT}$ and

$RB1^{del}$ G401 cells *in vitro* (half-maximal effective concentration of 17.6 ± 1.6 nM, 19.2 ± 3.8 nM,

and 26.7 ± 3.2 nM for $RB1^{WT}$, $RB1^{del}$ E1 and F2 clones, respectively; **Figure 5A**). We also found

that the combination of TAZ and elimusertib exerted greater antitumor effects than either drug

alone against diverse MRT and ES cell lines (**Figure 5B**), exhibiting synergy in a subset of the

cell lines (**Figure S6E-F**). To determine whether the synergistic elimusertib and TAZ

combination antitumor effects were due to increased DNA damage, we used confocal

immunofluorescence microscopy to quantify γH2AX phosphorylation, a specific marker of

dsDNA breaks (49). In agreement with prior studies (47), untreated G401 cells showed dsDNA

breaks associated with baseline *PGBD5* expression (**Figure S10A**). Consistent with TAZ-

mediated induction of *PGBD5* expression (**Figure 2A-C**), we found that TAZ treatment alone

significantly increased nuclear γH2AX fluorescence (median normalized level = 0.061 versus

0.12, respectively; *t*-test *p* = 1.7E-8; **Figures 5C & S10B**), and the combination of TAZ and

elimusertib induced additional dose-dependent increases in dsDNA break levels than either

drug alone (median normalized level = 0.079 versus 0.12 and 0.20, respectively; *t*-test *p* = 5.1E-

3 and 6.5E-4 for 50 and 100 nM TAZ, respectively; **Figure 5D**). Targeting TAZ-potentiated and

PGBD5-induced DNA damage using the ATR kinase-selective inhibitor elimusertib was specific,

because combination of TAZ with the DNA replication repair CHK1 kinase-selective inhibitor

SRA737 showed no increased activity as compared to either drug alone (**Figure S9A-B**).

Indeed, TAZ treatment did not induce apparent replication stress, as measured by RPA

phosphorylation (**Figure S9B**), which was also not potentiated by combined CHK1 inhibition with SRA737, in spite of effective suppression of CHK1 auto-phosphorylation (**Figure S9B**). Unlike T-ALL, where EZH2 suppression induces MYCN protein expression and replication stress (50), TAZ treatment of G401 rhabdoid tumor cells failed to increase MYCN protein abundance (**Figure S9C**), in spite of significant upregulation of *MYCN* mRNA (**Figure 2A-C**).

Encouraged by the potent and specific antitumor activity of synthetic lethal combination TAZ therapy *in vitro*, we tested the antitumor activity of elimusertib and TAZ combination using a diverse cohort of MRT and ES PDX mice *in vivo* (**Supplementary Table S5**). The combination of TAZ and elimusertib exceeded the effect of treatment with either drug alone when assessed by tumor measurements (Vardi *U*-test p = 2.0E-2 and 0.19 for combination versus elimusertib or TAZ, respectively; **Figure 5E**) and significantly extended tumor-free survival from 51 days (95% CI = 42-60 days) for elimusertib and 67 days (95% CI = 53-81 days) for TAZ to 99 days (95% CI = 74-123 days) for the combination (log-rank test p = 5.8E-4 and 3.9E-2 for combination versus elimusertib or TAZ, respectively; **Figure 5F**). This was most pronounced for the HYMAD_EPIS_X0004aS1 tumor (**Figures 5G-H & S8B-C**), which exhibited a relatively poor response to TAZ monotherapy, when assessed by tumor growth measurements (Vardi *U*-test p = 2.0E-4 for combination versus elimusertib or TAZ; **Figure 5G**) and tumor-free survival (log-rank test p = 6.2E-3 and 6.3E-5 for combination versus elimusertib or TAZ, respectively; **Figure 5H**). Thus, the combination of EZH2 and ATR inhibition constitutes a synthetic lethal rational combination strategy to improve TAZ clinical response and overcome resistance.

Discussion

What defines effective epigenetic EZH2 inhibition therapy for *SMARCB1*-deficient epithelioid sarcomas and rhabdoid tumors? Our studies of ES patients treated with TAZ demonstrate that effective inhibition of PRC2 enzymatic activity is necessary but not sufficient for durable antitumor effects. Using clinical genomics and transcriptomics, combined with functional genetic studies of more than 15 diverse MRT and ES cell lines and patient-derived tumors *in vitro* and *in vivo*, we propose a general molecular model for effective epigenetic TAZ therapy (**Figure 6**). This model places validated *RB1* and *EZH2* TAZ resistance alleles within the context of a molecular sequence of events required for clinical TAZ response. This model also explains additional mutations associated with TAZ resistance based on the perturbation of each stage of this sequence and provides candidate prognostic biomarkers and therapeutic combination

strategies. We discuss the evidence for this model below, and summarize its novel predictions
and implications.

First, TAZ must be able to bind and enzymatically inhibit the EZH2 SET domain (**Figure 6; Step**
1). This inhibition can be blocked by gatekeeper mutations of the EZH2 drug binding site, as
observed in lymphoma cell lines (18), and demonstrated for the first time here in an epithelioid
sarcoma with clinically acquired $EZH2^{Y666N}$ mutation. Such resistance mutations can be
overcome by targeting EED, a non-enzymatic PRC2 subunit.

For effective epigenetic TAZ therapy, chromatin remodeling complexes must act on tumor
suppressor loci that were aberrantly repressed by PRC2 (**Figure 6; Step 2**). The canonical BAF
complex is thought to oppose the activity of the Polycomb Repressive Complex (PRC),
associated with its chromatin eviction (5, 6). However, the precise mechanism of eviction of
TAZ-inhibited PRC2 in *SMARCB1*-deleted tumors is not fully defined. This may involve TAZ-
induced remodeling of BAF and/or PRC2. For example, a recent study of *SMARCA4*-deficient
cell lines found that upregulation of the expression of BAF helicase *SMARCA2*, which is under
PRC2 control in these cells, is necessary for response to EZH2 inhibition (51). In our study, we
observed *RB1*-independent upregulation of the expression of BAF subunits *SMARCA2* and
DPF3 upon TAZ treatment in G401 cells (**Figure S10A**). This suggests that TAZ may impact
BAF complex assembly as part of its therapeutic mechanism in *SMARCB1*-deficient tumor cells.
However, this also indicates that *SMARCA2* re-expression upon EZH2 inhibition is not sufficient
to induce cell cycle arrest in the absence of *RB1* expression. After assessing TAZ-induced gene
expression changes in PRC2 subunits, we also observed *RB1*-independent TAZ-induced
upregulation of PRC2 subunit *JARID2*, and *RB1*-dependent downregulation of *PHF19*,
suggesting that PRC2 composition itself may be affected by TAZ treatment (**Figure S10A**).
Additionally, it is unknown whether PRC2 eviction in TAZ-treated cells requires a specific form
of the BAF complex, such as the non-canonical *SMARCB1*-deficient ncBAF or GBAF complex
described previously (52, 53).

If the activity of specific BAF subtypes is indeed needed to evict PRC2 from chromatin upon
EZH2 inhibition, then genetic perturbation of specific BAF subunits may impact tumor response
to TAZ. For example, in our patient cohort, we observed one TAZ-sensitive tumor with a
truncation in the canonical BAF-specific subunit *ARID1B*, while a TAZ-resistant tumor had a
missense mutation in PBAF-specific subunit *ARID2* (**Figure 1A**). We also found *ARID1B* to be

mutated in 2 out of 5 TAZ-responsive cell lines (0 out of 5 TAZ-sensitive cell lines) and *ARID2* to

be mutated in 2 out of 5 TAZ-resistant cell lines, though also in 1 out of 5 TAZ-sensitive lines

(**Figure 3B**). It is not known whether these mutations affect tumor response to TAZ, and further

work will be needed to elucidate the specific mechanism of chromatin de-repression in TAZ-

treated tumor cells, and its requirement of specific BAF subunits and complexes.

Effective epigenetic TAZ therapy must also upregulate PRC2-repressed tumor suppressor loci

(**Figure 6; Step 3**). In our patient cohort, one TAZ-resistant tumor had deletions of both

CDKN2A and *CDKN2B* (**Figure 1A**), which can inhibit CDK4/6 from phosphorylating RB1 and

are known to be de-repressed by TAZ treatment (7). Our genomic analysis of MRT and ES cell

lines also showed that 4 out of 5 TAZ-resistant cell lines tested had apparent loss of *CDKN2A*,

with two also having loss of *CDKN2B*, and one having *CDKN1A* loss as well (**Figure 3B**). These

mutations may phenocopy *RB1* loss, suggesting that upregulation of these cell cycle inhibitors

may be necessary for effective TAZ therapy.

In addition to the preservation of these tumor suppressor loci, transcription factors and

coactivators that upregulate their expression must also be intact and expressed for tumor cells

to effectively respond to TAZ (**Figure 6; Step 3**). In our patient cohort, 2 TAZ-resistant patient

tumors had missense mutations in *ANKRD11* (**Figure 1A**), as did 3 out of 5 TAZ-resistant cell

lines. *ANKRD11* is a putative tumor suppressor that exhibits loss of heterozygosity in breast

cancer (54), and is recurrently mutated in other cancers (55, 56). *ANRKD11* can cooperate with

p53 to upregulate *CDKN1A*, and previous reports have suggested possible risk of cancer

development in patients with a constitutional loss of *ANKRD11* (57, 58). While its importance in

SMARCB1-deficient sarcomas is unknown, its association with TAZ resistance suggests

potential causality. We note that its proposed tumor suppressor function would require all alleles

to be lost or mutated. Consistent with potential functional significance, EPI544 cells harbor

multiple mutations in *ANKRD11*, two with an apparent allele frequency of 1.0 and MP-MRT-AN

cells harbor two mutations, both with 0.5 allele frequencies (**Supplementary Table S4**).

Interestingly, a recent report described a patient with KBG syndrome, a developmental condition

caused by mutation of *ANKRD11*, who also developed a rhabdoid tumor (58). The co-

occurrence of these two exceedingly rare conditions in the same patient is consistent with

potential functional involvement of *ANKRD11* in rhabdoid tumor development and susceptibility

to EZH2 inhibition.

We also identified *KLF4* as a putative marker of susceptibility to TAZ in patient tumors (**Figure
3G**), and found *KLF4* to be mutated in 1 out of 5 TAZ-resistant cell lines (**Figure 3B**). Although
KLF4 is a transcription factor with both an activating and repressing functions, its role in
SMARCB1-deleted tumors is not known. Its expression has long been known to upregulate
CDKN1A, causing cell cycle arrest at the G1/S checkpoint (40, 59). This suggests that KLF4
may regulate the induction of tumor suppressor genes in response to EZH2 inhibition, such as
its bona fide target *CDKN1A*. This may occur through recruitment of BAF to tumor suppressor
loci; KLF4 can recruit BAF to target genes to upregulate them (60). Like *ANKRD11*, *KLF4* may
thus be a key activator of PRC2-repressed genes (**Figure 5**).

Finally, effective TAZ therapy also requires the function of downstream cell cycle effectors of the
relevant tumor suppressor loci (**Figure 6; Step 4**). As we have demonstrated in this study, loss
of *RB1* leads to the evasion of TAZ-induced cell cycle arrest, despite effective inhibition of EZH2
activity and sustained transcriptional response to TAZ. A recent genome-wide CRISPR screen
also found *RB1* as a top mediator of TAZ resistance (15). This is reminiscent of the necessity of
intact *RB1* for therapeutic response to clinical inhibitors of CDK4/6 (61-63). The transcriptional
upregulation of hundreds of genes by TAZ in $RB1^{del}$ tumors, including EMT gene sets (**Figure
S2C&F**), suggests that these cells are undergoing forced differentiation, even while maintaining
proliferation (**Figure 2G-H**). This upregulation of mesenchymal gene sets is consistent with
previous work that has shown that PRC2 inhibition, similar to *SMARCB1* re-expression, can
drive *SMARCB1*-deficient tumors into a terminally differentiated, mesenchymal-like state (7, 64,
65), possibly recapitulating the normal developmental trajectory of their cells of origin (64). In
this way, *RB1* loss and dysregulation of the RB1/E2F axis appear to decouple the regulation of
cell fate and identity from its control of cell cycle progression.

In our search for predictive biomarkers of TAZ response, we identified increased expression of
the PCP gene *PRICKLE1* to be associated with a deficient RB1/E2F axis and TAZ resistance
(**Figure 3C-F**). The molecular mechanism connecting G1/S dysregulation and *PRICKLE1*
expression is currently unknown, but the PCP pathway is known to be under cell cycle control
(66, 67). This is mediated at least in part by PLK1, which we found to be one of the top
upregulated genes in TAZ-resistant tumors (**Figure S5B**). Likewise, deletion of the *Drosophila*
RB1 homologue *Rbf1* results in the upregulation of several PCP genes, including the *PRICKLE1*
homologue *pk* (68). This suggests that dysregulation of the RB1/E2F axis may lead to
upregulation of *PRICKLE1* through dysregulation of normal cell cycle control of PCP. Further

work will be needed to define this mechanism, as well as to investigate PRICKLE1 as a clinical

biomarker for TAZ resistance.

Finally, our study developed two strategies to circumvent clinical TAZ resistance. First, since

dysregulation of the RB1/E2F axis mediates escape from cell cycle arrest at the G1/S

checkpoint, we reasoned that cell cycle kinases that function downstream of this checkpoint

would remain viable therapeutic targets. This cell cycle bypass strategy is supported by

previous work showing that loss of *RB1* can sensitize cancer cells to Aurora kinase inhibition

through a primed spindle assembly checkpoint (69). As predicted, *RB1^{del}* cells remain sensitive

to CDK2, AURKA, and AURKB inhibition (**Figures 4A, S6A-B**). We found that MRT and ES cell

lines resistant to TAZ, including those with *ANKRD11* and *CDKN1A/2A/2B* mutations were

sensitive to barasertib. Most compellingly, we found that the combination of TAZ and barasertib

exhibits improved antitumor activity *in vivo* compared to either drug alone.

This is reminiscent of the therapeutic combination mechanism proposed by Sorger and Palmer

(70), in which inter-patient and inter-tumor variability in response to individual drugs rather than

their pharmacological interactions lead to apparent combined effects. We observed substantial

benefit of TAZ and barasertib combination within individual PDX models. This suggests that the

improved efficacy of this combination may also result from intra-tumor heterogeneity and tumor

evolution *in vivo*, as proposed for combination therapy more than fifty years ago (71, 72). Dual

targeting of two different parts of the cell cycle can prevent tumors from evading cell cycle arrest

through the presence or acquisition of mutations in cell cycle control genes. For example, cells

with a defective G1/S checkpoint that continue to transit through the cell cycle remain sensitive

to AURKB inhibition. Further work will be needed to elucidate whether barasertib causes

postmitotic endoreduplication due to its disruption of the spindle assembly and/or direct

inhibition of the mitotic checkpoint (73, 74).

Our findings also advance synthetic lethal strategy for rational epigenetic TAZ combination

therapy due to TAZ-induced expression of *PGBD5*, the putative developmental mutator in

rhabdoid and other young-onset solid tumors, due to its induction of dsDNA breaks (46). We

found that the DNA damage repair ATR kinase inhibitor elimusertib not only overcomes

RB1/E2F axis-mediated resistance, but in combination with TAZ, also exerts synergistic anti-

tumor effects *in vitro* and *in vivo*. TAZ-induced upregulation of PGBD5 is associated with the

induction of DNA damage. It is possible that the enhanced sensitivity to ATR inhibition due to

the increased requirements for DNA repair also depends on the intrinsic variation in DNA damage repair signaling among different tumor subtypes. This may also be due to the variation in the expression and activity of PGBD5 nuclease activity among tumors, both of which may be associated with the recently described molecular subtypes of ES and MRT tumors (47). Additional synthetic lethal strategies may also be developed based on the immunologic effects of EZH2 inhibitors, particularly as combined with epigenetic and synthetic lethal therapies, both of which can promote tumor immunogenicity. In all, this study develops a paradigm for rational epigenetic combination therapy, including candidate prognostic biomarkers, all of which should be incorporated into future clinical trials for patients.

Acknowledgements

This work is dedicated to Maggie Schmidt and her family, and many other patients and their advocates who inspire and support our research. We thank Richard Koche, Nicholas Socci, and Mithat Gönen for technical advice, Alejandro Gutierrez and Stuart Orkin for *EZH2* expression vectors, Helen Mueller and other members of our labs for critical advice and manuscript comments, and Epizyme, Bayer and AstraZeneca for supplying tazemetostat, elimusertib, and barasertib, respectively. This work was supported by the MSK Integrated Genomics Operation Core, Anti-Tumor Assessment Core, Bioinformatics Core, Molecular Diagnostics Service and the Department of Pathology, the Marie-Josée and Henry R. Kravis Center for Molecular Oncology, NIH R01 CA214812, P30 CA08748, T32 GM007739, Burroughs Wellcome Fund, Rita Allen Foundation, Pershing Square Sohn Cancer Research Alliance and the G. Harold and Leila Y. Mathers Foundation, Cycle for Survival, MSK Sarcoma Center, the Starr Cancer Consortium, and Maggie's Mission. Yaniv Kazansky was supported by a Medical Scientist Training Program grant from the National Institute of General Medical Sciences of the National Institutes of Health under award number T32 GM007739 to the Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program. AK is a Scholar of the Leukemia & Lymphoma Society.

Conflict of Interest

The authors have no competing financial interests. AK is a consultant for Novartis, Rgenta, and Blueprint.

Methods

EZH2 mutant plasmids

The *EZH2^{Y666N}* mutation detected in the clinical trial patient refers to amino acid numbering in isoform 2 of the protein. For consistency of nomenclature, all engineered mutations use numbering referring to isoform 2 (Uniprot ID: Q15190-2), although isoform 1 was expressed in cells for this study. Plasmids containing wild-type EZH2 (*EZH2^{WT}*) and catalytically inactive triple mutant (F672I, H694A, R732K, referred to as *EZH2^{CatMut}*) plasmids were kindly provided by Alejandro Gutierrez in the doxycycline-inducible pINDUCER20 vector (75). The plasmids contain human *EZH2* tagged N-terminally with a FLAG-Avi tag.

The Y666N mutation was engineered in both the *EZH2^{WT}* and *EZH2^{CatMut}* plasmids to yield *EZH2^{Y666N}* and what we termed *EZH2^{QuadMut}*. The mutation was introduced by site-directed mutagenesis using the QuikChange Lightning Kit (Agilent) per manufacturer's instructions (mutagenesis primers: 5'-GCAAAGTGTACGACAAGAACATGTGCAGCTTTCTG-3' and 5'-CAGAAAGCTGCACATGTTCTTGTCGTACACTTTGC-3') to engineer a TAC to AAC codon change. After mutagenesis, PCR products were transformed into Stbl3 *E. coli* cells and expanded at 30°C. Correct plasmid sequences were confirmed by Sanger sequencing (Eton). The sequencing primers used are:

F1: 5'-GGACAGCAGAGATCCAGTTTG-3'

R1: 5'-GGTCCGTTCCAGGATCTTCT-3'

F2: 5'-TCCAGTGTGGTGGAATTCTG-3'

R2: 5'-TATCGCTGGGGAACTTTCTG-3'

F3: 5'-TGCTGCACAACATCCCTTAC-3'

R3: 5'-TGCTGGTTTCGTCCTTCTTT-3'

F4: 5'-CCTACAAGCGGAAGAACACC-3'

R4: 5'-GTTCTTGCTGTCCCAGTGGT-3'

F5: 5'-CTGAAGAAGGATGGCAGCTC-3'

R5: 5'-CTTGGGTGGGTTACTCCAGA-3'

RB1 Knockout

G401 cells with *RB1* mutations were engineered by Synthego. Briefly, a single guide RNA targeting exon 2 of *RB1* was used (AGAGAGAGCUUGGUUAACUU). Ribonucleoprotein containing spCas9 and sgRNA was transfected into G401 cells by electroporation. The target

site was then PCR-amplified and Sanger sequenced to ensure homozygous indels (PCR and

sequencing primers: Forward- CACTGTGTGGTATCCTTATTTTGGA, Reverse-

AGGTAAATTTCCTCTGGGTAATGGA, with the forward primer used for sequencing). The cells

were then single-cell cloned and re-verified by Sanger sequencing. Loss of the RB1 protein was

confirmed by Western blot.

Cell cycle analysis

G401 cells were plated on Day 0 and treated for 10 days with 1 μm tazemetostat or equivalent

volume of DMSO, with drug and media replaced on Days 4 and 7. On Day 11, cells were pulsed

with EdU for 1 hour. Cells were then harvested and processed for flow cytometry using the

manufacturer's protocol (Click-iT, Invitrogen). Briefly, cells were washed with PBS with 1% BSA,

permeabilized, and incubated with AlexaFluor647 for 30 minutes. DNA content was measured

using propidium iodide (0.05 μg/μL). Cells were analyzed on a CytoFLEX LX (Beckman

Coulter).

Patient tumor samples

Patient tumor and matched normal blood samples were obtained from patients at Memorial

Sloan Kettering Cancer Center (MSKCC) enrolled in the TAZ clinical trial (13). All patients

provided informed consent for this study under the Institutional Review Board approved

research protocol 12-245. Patient tumors were classified into "Response" or "Progression"

groups based on RECIST 1.1 criteria (76). "Response" included tumors exhibiting a complete

response, partial response, or stable disease. All other tumors were classified under the

"Progression" group. The complete list of tumor samples used and corresponding clinical data

may be found in Supplementary Table 1. This cohort includes both tumor samples that

underwent targeted sequencing with MSK-IMPACT (16) as part of their clinical care at MSKCC

as well as archived tumors that were analyzed for this study. For genomic analysis, DNA was

extracted from either flash frozen tumor samples or formalin-fixed, paraffin-embedded (FFPE)

blocks or slides and samples were processed using the IMPACT468 or IMPACT505 panels

depending on the time of their sequencing (16). The detected mutations and copy number

alterations were obtained from cBioPortal (77, 78) and can be found in Supplementary Table 2.

Oncoprints were generated using Oncoprinter (cBioPortal).

For transcriptomic analysis, archived frozen tumor samples were weighed and up to 20-30mg

were homogenized in RLT buffer, followed by extraction using the AllPrep DNA/RNA Mini Kit

(QIAGEN catalog 80204) according to the manufacturer's instructions. RNA was eluted in 13 µL

nuclease-free water. After RiboGreen quantification and quality control by Agilent BioAnalyzer, 1

µg of total RNA with DV200 percentages varying from 78% to 100% underwent ribosomal

depletion and library preparation using the TruSeq Stranded Total RNA LT Kit (Illumina catalog

RS-122-1202) according to instructions provided by the manufacturer with 8 cycles of PCR.

Samples were barcoded and sequenced using NovaSeq 6000 in a PE150 mode, with the

NovaSeq 6000 S4 Reagent Kit (Illumina). On average, 84 million paired reads were generated

per sample and 70% of the data mapped to mRNA.

Targeted sequencing of cell lines

To assess for the presence of somatic mutations in MRT and ES cell lines, DNA was extracted

using the PureLink Genomic DNA Minikit (Invitrogen) and processed using the IMPACT505

panel as above. Due to the lack of matched normal tissue for cell lines, copy number alterations

were detected using a custom algorithm using circular binary segmentation (79) implemented by

the MSK Bioinformatics core. Code is available on github at:

https://github.com/kentsisresearchgroup/seqCNA_tazemetostat_resistance.

Additional methods may be found in the Supplementary Methods and Figures file.

Data Availability

RNA-seq data from G401 cells can be found at the Gene Expression Omnibus (GEO)

repository, accession number GSE213845. RNA-seq data from patient tumor samples has been

deposited to the Database of Genotypes and Phenotypes (dbGaP), accession number

phs003188.v1.p1. Additional supplementary data files and computational analysis scripts are

available at Zenodo (10.5281/zenodo.7595037).

References

1.   Kadoch C, Crabtree GR. Mammalian SWI/SNF chromatin remodeling complexes and cancer: Mechanistic insights gained from human genomics. Sci Adv. 2015;1(5):e1500447.

2.   Versteege I, Sevenet N, Lange J, Rousseau-Merck MF, Ambros P, Handgretinger R, et al. Truncating mutations of hSNF5/INI1 in aggressive paediatric cancer. Nature. 1998;394(6689):203-6.

3.   Kim KH, Kim W, Howard TP, Vazquez F, Tsherniak A, Wu JN, et al. SWI/SNF-mutant cancers depend on catalytic and non-catalytic activity of EZH2. Nat Med. 2015;21(12):1491-6.

4.   McKenna ES, Sansam CG, Cho YJ, Greulich H, Evans JA, Thom CS, et al. Loss of the epigenetic tumor suppressor SNF5 leads to cancer without genomic instability. Mol Cell Biol. 2008;28(20):6223-33.

5.   Wilson BG, Wang X, Shen X, McKenna ES, Lemieux ME, Cho YJ, et al. Epigenetic antagonism between polycomb and SWI/SNF complexes during oncogenic transformation. Cancer Cell. 2010;18(4):316-28.

6.   Kadoch C, Williams RT, Calarco JP, Miller EL, Weber CM, Braun SM, et al. Dynamics of BAF-Polycomb complex opposition on heterochromatin in normal and oncogenic states. Nat Genet. 2017;49(2):213-22.

7.   Knutson SK, Warholic NM, Wigle TJ, Klaus CR, Allain CJ, Raimondi A, et al. Durable tumor regression in genetically altered malignant rhabdoid tumors by inhibition of methyltransferase EZH2. Proc Natl Acad Sci U S A. 2013;110(19):7922-7.

8.   McCabe MT, Ott HM, Ganji G, Korenchuk S, Thompson C, Van Aller GS, et al. EZH2 inhibition as a therapeutic strategy for lymphoma with EZH2-activating mutations. Nature. 2012;492(7427):108-12.

9.   Konze KD, Ma A, Li F, Barsyte-Lovejoy D, Parton T, Macnevin CJ, et al. An orally bioavailable chemical probe of the Lysine Methyltransferases EZH2 and EZH1. ACS Chem Biol. 2013;8(6):1324-34.

10.   Li C, Wang Y, Gong Y, Zhang T, Huang J, Tan Z, et al. Finding an easy way to harmonize: a review of advances in clinical research and combination strategies of EZH2 inhibitors. Clin Epigenetics. 2021;13(1):62.

11.   Guerra S, Cichowski K. Targeting Cancer at the Intersection of Signaling and Epigenetics. Annual Review of Cancer Biology. 2019;3(1):365-84.

12.   Baylin SB, Jones PA. Epigenetic Determinants of Cancer. Cold Spring Harb Perspect Biol. 2016;8(9).

13.   Gounder M, Schöffski P, Jones RL, Agulnik M, Cote GM, Villalobos VM, et al. Tazemetostat in advanced epithelioid sarcoma with loss of INI1/SMARCB1: an international, open-label, phase 2 basket study. Lancet Oncol. 2020;21(11):1423-32.

14.   Chi SN, Bourdeaut F, Casanova M, Kilburn LB, Hargrave DR, McCowage GB, et al. Update on phase 1 study of tazemetostat, an enhancer of zeste homolog 2 inhibitor, in pediatric patients with relapsed or refractory integrase interactor 1–negative tumors. Journal of Clinical Oncology. 2022;40(16_suppl):10040-.

15.   Drosos Y, Myers JA, Xu B, Mathias KM, Beane EC, Radko-Juettner S, et al. NSD1 mediates antagonism between SWI/SNF and polycomb complexes and is required for transcriptional activation upon EZH2 inhibition. Mol Cell. 2022;82(13):2472-89.e8.

16.   Cheng DT, Mitchell TN, Zehir A, Shah RH, Benayed R, Syed A, et al. Memorial Sloan Kettering-Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT): A Hybridization Capture-Based Next-Generation Sequencing Clinical Assay for Solid Tumor Molecular Oncology. J Mol Diagn. 2015;17(3):251-64.

17.   Bracken AP, Pasini D, Capra M, Prosperini E, Colli E, Helin K. EZH2 is downstream of the pRB-E2F pathway, essential for proliferation and amplified in cancer. Embo j. 2003;22(20):5323-35.

18.	Gibaja V, Shen F, Harari J, Korn J, Ruddy D, Saenz-Vash V, et al. Development of secondary mutations in wild-type and mutant EZH2 alleles cooperates to confer resistance to EZH2 inhibitors. Oncogene. 2016;35(5):558-66.

19.	Bisserier M, Wajapeyee N. Mechanisms of resistance to EZH2 inhibitors in diffuse large B-cell lymphomas. Blood. 2018;131(19):2125-37.

20.	Brooun A, Gajiwala KS, Deng Y-L, Liu W, Bolaños B, Bingham P, et al. Polycomb repressive complex 2 structure with inhibitor reveals a mechanism of activation and drug resistance.

21.	Honma D, Kanno O, Watanabe J, Kinoshita J, Hirasawa M, Nosaka E, et al. Novel orally bioavailable EZH1/2 dual inhibitors with greater antitumor efficacy than an EZH2 selective inhibitor. Cancer Sci. 2017;108(10):2069-78.

22.	He Y, Selvaraju S, Curtin ML, Jakob CG, Zhu H, Comess KM, et al. The EED protein–protein interaction inhibitor A-395 inactivates the PRC2 complex. Nature Chemical Biology. 2017;13(4):389-95.

23.	Qi W, Zhao K, Gu J, Huang Y, Wang Y, Zhang H, et al. An allosteric PRC2 inhibitor targeting the H3K27me3 binding pocket of EED. Nat Chem Biol. 2017;13(4):381-8.

24.	Tomassi S, Romanelli A, Zwergel C, Valente S, Mai A. Polycomb Repressive Complex 2 Modulation through the Development of EZH2–EED Interaction Inhibitors and EED Binders. Journal of Medicinal Chemistry. 2021;64(16):11774-97.

25.	Coe BP, Thu KL, Aviel-Ronen S, Vucic EA, Gazdar AF, Lam S, et al. Genomic deregulation of the E2F/Rb pathway leads to activation of the oncogene EZH2 in small cell lung cancer. PLoS One. 2013;8(8):e71670.

26.	Kenny C, O'Meara E, Ulaş M, Hokamp K, O'Sullivan MJ. Global Chromatin Changes Resulting from Single-Gene Inactivation-The Role of SMARCB1 in Malignant Rhabdoid Tumor. Cancers (Basel). 2021;13(11).

27.	Kia SK, Gorski MM, Giannakopoulos S, Verrijzer CP. SWI/SNF mediates polycomb eviction and epigenetic reprogramming of the INK4b-ARF-INK4a locus. Mol Cell Biol. 2008;28(10):3457-64.

28.	Kotake Y, Cao R, Viatour P, Sage J, Zhang Y, Xiong Y. pRB family proteins are required for H3K27 trimethylation and Polycomb repression complexes binding to and silencing p16INK4alpha tumor suppressor gene. Genes Dev. 2007;21(1):49-54.

29.	Gallagher D, Voronova A, Zander MA, Cancino GI, Bramall A, Krause MP, et al. Ankrd11 is a chromatin regulator involved in autism that is essential for neural development. Dev Cell. 2015;32(1):31-42.

30.	Neilsen PM, Cheney KM, Li CW, Chen JD, Cawrse JE, Schulz RB, et al. Identification of ANKRD11 as a p53 coactivator. J Cell Sci. 2008;121(Pt 21):3541-52.

31.	Noll JE, Jeffery J, Al-Ejeh F, Kumar R, Khanna KK, Callen DF, et al. Mutant p53 drives multinucleation and invasion through a process that is suppressed by ANKRD11. Oncogene. 2012;31(23):2836-48.

32.	Béguelin W, Rivas MA, Calvo Fernández MT, Teater M, Purwada A, Redmond D, et al. EZH2 enables germinal centre formation through epigenetic silencing of CDKN1A and an Rb-E2F1 feedback loop. Nat Commun. 2017;8(1):877.

33.	Fan T, Jiang S, Chung N, Alikhan A, Ni C, Lee CC, et al. EZH2-dependent suppression of a cellular senescence phenotype in melanoma cells by inhibition of p21/CDKN1A expression. Mol Cancer Res. 2011;9(4):418-29.

34.	Negishi M, Wongpalee SP, Sarkar S, Park J, Lee KY, Shibata Y, et al. A new lncRNA, APTR, associates with and represses the CDKN1A/p21 promoter by recruiting polycomb proteins. PLoS One. 2014;9(4):e95216.

35.	Daulat AM, Finetti P, Revinski D, Silveira Wagner M, Camoin L, Audebert S, et al. ECT2 associated to PRICKLE1 are poor-prognosis markers in triple-negative breast cancer. Br J Cancer. 2019;120(9):931-40.

36. Daulat AM, Bertucci F, Audebert S, Sergé A, Finetti P, Josselin E, et al. PRICKLE1
Contributes to Cancer Cell Dissemination through Its Interaction with mTORC2. Dev Cell.
2016;37(4):311-25.
37. Jiang D, He Y, Mo Q, Liu E, Li X, Huang L, et al. PRICKLE1, a Wnt/PCP signaling
component, is overexpressed and associated with inferior prognosis in acute myeloid leukemia.
J Transl Med. 2021;19(1):211.
38. Zhou R, Tang Z, Li H, Wang X, Sun Y. PRICKLE1 promotes gastric cancer metastasis
by activating mTOR signaling. Am J Transl Res. 2021;13(5):4266-80.
39. Hu J, Yang Y, Ma Y, Ning Y, Chen G, Liu Y. Proliferation Cycle Transcriptomic
Signatures are Strongly associated With Gastric Cancer Patient Survival. Front Cell Dev Biol.
2021;9:770994.
40. Chen X, Johns DC, Geiman DE, Marban E, Dang DT, Hamlin G, et al. Krüppel-like factor
4 (gut-enriched Krüppel-like factor) inhibits cell proliferation by blocking G1/S progression of the
cell cycle. J Biol Chem. 2001;276(32):30423-8.
41. Chen X, Whitney EM, Gao SY, Yang VW. Transcriptional profiling of Krüppel-like factor
4 reveals a function in cell cycle regulation and epithelial differentiation. J Mol Biol.
2003;326(3):665-77.
42. De Azevedo WF, Leclerc S, Meijer L, Havlicek L, Strnad M, Kim SH. Inhibition of cyclin-
dependent kinases by purine analogues: crystal structure of human cdk2 complexed with
roscovitine. Eur J Biochem. 1997;243(1-2):518-26.
43. Manfredi MG, Ecsedy JA, Chakravarty A, Silverman L, Zhang M, Hoar KM, et al.
Characterization of Alisertib (MLN8237), an investigational small-molecule inhibitor of aurora A
kinase using novel in vivo pharmacodynamic assays. Clin Cancer Res. 2011;17(24):7614-24.
44. Wilkinson RW, Odedra R, Heaton SP, Wedge SR, Keen NJ, Crafter C, et al. AZD1152, a
selective inhibitor of Aurora B kinase, inhibits human tumor xenograft growth by inducing
apoptosis. Clin Cancer Res. 2007;13(12):3682-8.
45. Vardi Y, Ying Z, Zhang CH. Two-sample tests for growth curves under dependent right
censoring. Biometrika. 2001;88(4):949-60.
46. Henssen AG, Koche R, Zhuang J, Jiang E, Reed C, Eisenberg A, et al. PGBD5
promotes site-specific oncogenic mutations in human tumors. Nat Genet. 2017;49(7):1005-14.
47. Henssen AG, Reed C, Jiang E, Garcia HD, von Stebut J, MacArthur IC, et al.
Therapeutic targeting of PGBD5-induced DNA repair dependency in pediatric solid tumors.
Science translational medicine. 2017;9(414).
48. Henssen AG, Henaff E, Jiang E, Eisenberg AR, Carson JR, Villasante CM, et al.
Genomic DNA transposition induced by human PGBD5. Elife. 2015;4.
49. Mah LJ, El-Osta A, Karagiannis TC. gammaH2AX: a sensitive molecular marker of DNA
damage and repair. Leukemia. 2010;24(4):679-86.
50. León TE, Rapoz-D'Silva T, Bertoli C, Rahman S, Magnussen M, Philip B, et al. EZH2-
Deficient T-cell Acute Lymphoblastic Leukemia Is Sensitized to CHK1 Inhibition through
Enhanced Replication Stress. Cancer Discov. 2020;10(7):998-1017.
51. Januario T, Ye X, Bainer R, Alicke B, Smith T, Haley B, et al. PRC2-mediated repression
of SMARCA2 predicts EZH2 inhibitor activity in SWI/SNF mutant tumors. Proc Natl Acad Sci U
S A. 2017;114(46):12249-54.
52. Wang X, Wang S, Troisi EC, Howard TP, Haswell JR, Wolf BK, et al. BRD9 defines a
SWI/SNF sub-complex and constitutes a specific vulnerability in malignant rhabdoid tumors. Nat
Commun. 2019;10(1):1881.
53. Michel BC, D'Avino AR, Cassel SH, Mashtalir N, McKenzie ZM, McBride MJ, et al. A
non-canonical SWI/SNF complex is a synthetic lethal target in cancers driven by BAF complex
perturbation. Nat Cell Biol. 2018;20(12):1410-20.
54. Powell JA, Gardner AE, Bais AJ, Hinze SJ, Baker E, Whitmore S, et al. Sequencing,
transcript identification, and quantitative gene expression profiling in the breast cancer loss of

heterozygosity region 16q24.3 reveal three potential tumor-suppressor genes. Genomics.
2002;80(3):303-10.
55.   Vasmatzis G, Johnson SH, Knudson RA, Ketterling RP, Braggio E, Fonseca R, et al.
Genome-wide analysis reveals recurrent structural abnormalities of TP63 and other p53-related
genes in peripheral T-cell lymphomas. Blood. 2012;120(11):2280-9.
56.   Gebauer N, Künstner A, Ketzer J, Witte HM, Rausch T, Benes V, et al. Genomic insights
into the pathogenesis of Epstein-Barr virus-associated diffuse large B-cell lymphoma by whole-
genome and targeted amplicon sequencing. Blood Cancer J. 2021;11(5):102.
57.   Isrie M, Hendriks Y, Gielissen N, Sistermans EA, Willemsen MH, Peeters H, et al.
Haploinsufficiency of ANKRD11 causes mild cognitive impairment, short stature and minor
dysmorphisms. Eur J Hum Genet. 20. England2012. p. 131-3.
58.   Behnert A, Auber B, Steinemann D, Frühwald MC, Huisinga C, Hussein K, et al. KBG
syndrome patient due to 16q24.3 microdeletion presenting with a paratesticular rhabdoid tumor:
Coincidence or cancer predisposition? Am J Med Genet A. 2018;176(6):1449-54.
59.   Zhang W, Geiman DE, Shields JM, Dang DT, Mahatan CS, Kaestner KH, et al. The gut-
enriched Kruppel-like factor (Kruppel-like factor 4) mediates the transactivating effect of p53 on
the p21WAF1/Cip1 promoter. J Biol Chem. 2000;275(24):18391-8.
60.   Moonen JR, Chappell J, Shi M, Shinohara T, Li D, Mumbach MR, et al. KLF4 recruits
SWI/SNF to increase chromatin accessibility and reprogram the endothelial enhancer landscape
under laminar shear stress. Nat Commun. 2022;13(1):4941.
61.   Schoninger SF, Blain SW. The Ongoing Search for Biomarkers of CDK4/6 Inhibitor
Responsiveness in Breast Cancer. Mol Cancer Ther. 2020;19(1):3-12.
62.   Finn RS, Dering J, Conklin D, Kalous O, Cohen DJ, Desai AJ, et al. PD 0332991, a
selective cyclin D kinase 4/6 inhibitor, preferentially inhibits proliferation of luminal estrogen
receptor-positive human breast cancer cell lines in vitro. Breast Cancer Res. 2009;11(5):R77.
63.   Dean JL, Thangavel C, McClendon AK, Reed CA, Knudsen ES. Therapeutic CDK4/6
inhibition in breast cancer: key mechanisms of response and failure. Oncogene.
2010;29(28):4018-32.
64.   Custers L, Khabirova E, Coorens THH, Oliver TRW, Calandrini C, Young MD, et al.
Somatic mutations and single-cell transcriptomes reveal the root of malignant rhabdoid tumours.
Nat Commun. 2021;12(1):1407.
65.   Nakayama RT, Pulice JL, Valencia AM, McBride MJ, McKenzie ZM, Gillespie MA, et al.
SMARCB1 is required for widespread BAF complex-mediated activation of enhancers and
bivalent promoters. Nat Genet. 2017;49(11):1613-23.
66.   Noatynska A, Tavernier N, Gotta M, Pintard L. Coordinating cell polarity and cell cycle
progression: what can we learn from flies and worms? Open Biol. 2013;3(8):130083.
67.   Shrestha R, Little KA, Tamayo JV, Li W, Perlman DH, Devenport D. Mitotic Control of
Planar Cell Polarity by Polo-like Kinase 1. Dev Cell. 2015;33(5):522-34.
68.   Payankaulam S, Yeung K, McNeill H, Henry RW, Arnosti DN. Regulation of cell polarity
determinants by the Retinoblastoma tumor suppressor protein. Sci Rep. 2016;6:22879.
69.   Gong X, Du J, Parsons SH, Merzoug FF, Webster Y, Iversen PW, et al. Aurora A Kinase
Inhibition Is Synthetic Lethal with Loss of the RB1 Tumor Suppressor Gene. Cancer Discov.
2019;9(2):248-63.
70.   Palmer AC, Sorger PK. Combination Cancer Therapy Can Confer Benefit via Patient-to-
Patient Variability without Drug Additivity or Synergy. Cell. 2017;171(7):1678-91.e13.
71.   Law LW. Effects of combinations of antileukemic agents on an acute lymphocytic
leukemia of mice. Cancer Res. 1952;12(12):871-8.
72.   Frei E, 3rd, Karon M, Levin RH, Freireich EJ, Taylor RJ, Hananian J, et al. The
effectiveness of combinations of antileukemic agents in inducing and maintaining remission in
children with acute leukemia. Blood. 1965;26(5):642-56.

73.   Tsuda Y, Iimori M, Nakashima Y, Nakanishi R, Ando K, Ohgaki K, et al. Mitotic slippage
and the subsequent cell fates after inhibition of Aurora B during tubulin-binding agent-induced
mitotic arrest. Sci Rep. 2017;7(1):16762.
74.   Hauf S, Cole RW, LaTerra S, Zimmer C, Schnapp G, Walter R, et al. The small molecule
Hesperadin reveals a role for Aurora B in correcting kinetochore-microtubule attachment and in
maintaining the spindle assembly checkpoint. J Cell Biol. 2003;161(2):281-94.
75.   Meerbrey KL, Hu G, Kessler JD, Roarty K, Li MZ, Fang JE, et al. The pINDUCER
lentiviral toolkit for inducible RNA interference in vitro and in vivo. Proc Natl Acad Sci U S A.
2011;108(9):3665-70.
76.   Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, et al. New
response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). Eur J
Cancer. 2009;45(2):228-47.
77.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer
Discov. 2012;2(5):401-4.
78.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative
analysis of complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal.
2013;6(269):pl1.
79.   Olshen AB, Venkatraman ES, Lucito R, Wigler M. Circular binary segmentation for the
analysis of array-based DNA copy number data. Biostatistics. 2004;5(4):557-72.

885
Figure 1: Patient tumor sequencing reveals diverse tazemetostat resistance mutations.
(**A**) Abridged oncoprint of selected genes from MSK-IMPACT sequencing on patient tumor

samples. Top panel: Tumor samples prior to TAZ treatment. Bottom panel: Matched samples
pre- and post-TAZ or pre- and post-acquisition of resistance. (**B**) Pre- and post- treatment CT
imaging of the indicated patient tumors which acquired *EZH2* and *RB1* mutations. (**C**) Atomic
molecular model of the chimeric *Homo sapiens/Anolis carolinensis* EZH2 bound to pyridone-
based EZH2 inhibitor I (blue), PDB: 5IJ7. Y666 is highlighted in red. (**D**) Top panel: Doxycycline-
inducible *EZH2* is expressed in single-cell G401 clones at near-physiological levels after 3 days
treatment with doxycycline at 1 μg/mL. Numbers indicate clone ID. Bottom panel: Cell viability
measured by CellTiter-Glo after 14 days of treatment with the indicated drug at 10 μM or
equivalent volume of DMSO. n=5 biological replicates per condition. *p = 3.5E-3, **p = 5.2E-3,
***p = 3.8E-5, ****p = 1.1E-5 by two-sided Student's t-test. (**E**) Top panel: *RB1* knockout in two
G401 clones (E1 and F2). Bottom panel: Cell viability measured by CellTiter-Glo after treatment
with 10 μM tazemetostat or DMSO for 14 days. *n*=5 biological replicates per condition. *p =
2.8E-5, **p = 9.0E-5 by two-sided Student's t-test.
901
902
903

Figure 2: *RB1* loss allows escape from cell cycle arrest despite effective EZH2 inhibition.
(**A-C**) Volcano plots of RNA-seq data from TAZ-treated G401 *RB1^{WT}* (**A**), *RB1^{del} E1* (**B**), and
RB1^{del} F2 (**C**). Plots show gene expression changes of cells treated with 10 μM TAZ versus
equivalent volume of DMSO for 11 days. n=3 biological replicates per condition. Dots in red
show genes with expression changes of log$_2$(fold change) > ±1 and *p*-value < 0.01. (**D-F**) GSEA
plots showing the PRC2_EZH2_UP.UP gene set for G401 *RB1^{WT}* (**D**), *RB1^{del} E1* (**E**), and *RB1^{del}*
F2 (**F**) cells treated in A-C. (**G**) Cell cycle analysis of the indicated G401 clone. Plot shows
triplicate measurements of cells treated with 1 μM TAZ versus equivalent volume of DMSO for
11 days. Y-axis shows the percent difference of the TAZ-treated cells in each cell cycle phase
compared to DMSO-treated cells. (**H**) Western blot of the indicated G401 cell clones treated
with 10 μM TAZ versus equivalent volume of DMSO for 11 days. Actin serves as loading
control.

Figure 3: An intact RB1/E2F axis is a key requirement for response to TAZ: (**A**) Dose-response curves of a panel of MRT and ES cell lines treated with TAZ for 13 days. Curves in red correspond to TAZ-responsive cell lines, grey to TAZ-resistant cell lines. (**B**) Abridged oncoprint of MRT and ES cell lines. Genes included here are BAF subunits and CDK4/6/RB1/E2F axis genes. Full list of mutations is listed in Supplementary Table 4. (**C-D**) Volcano plots of RNA-seq data comparing DMSO-treated G401 $RB1^{del}$ E1 (**C**) and F2 (**D**) with $RB1^{WT}$ cells. (**E**) Western blot for PRICKLE1 in untreated G401 cells. (**F-G**) Read counts for $PRICKLE1$ (**F**) and $KLF4$ (**G**) showing counts normalized by DESeq2 for patient tumor samples collected prior to TAZ treatment. (**H**) Read counts for $AURKB$ for patient tumor samples collected after TAZ treatment. $*p$=0.013, $**p$=0.03.

Figure 4: Cell cycle bypass combination strategy using AURKB inhibition overcomes TAZ resistance and improves response: (**A**) G401 cells treated with barasertib for 6 days (**A**). (**B**) Panel of MRT and ES cell lines ordered left to right by decreasing response to TAZ monotherapy. Cells were treated with the indicated monotherapy or combination for 11 days. Drug concentrations used were: TAZ: 200 nM, barasertib: 8 nM. (**C**) Tumor growth curves showing volumes calculated from caliper measurements for 5 mouse PDXs treated with the indicated drug regimen. n = 20 mice for vehicle and barasertib-treated groups, n = 21 for TAZ and TAZ + barasertib-treated groups. Vardi *U*-test *p* = 4.0E-4 and 2.0E-4 for combination vs. barasertib or TAZ, respectively. (**D**) Kaplan-Meier curves showing tumor-free survival (defined

as tumor volume ≤ 1,000 mm^3) for the PDXs in panel C. Mean survival is 65 days (95% CI: 51-

78 days) for barasertib, 67 days (95% CI: 53-81 days) for TAZ, 98 days (95% CI: 84-112 days)

for the combination. Log-rank test p = 3.3E-3 and 5.8E-3 for combination vs. barasertib or TAZ,

respectively. (**E**) Tumor growth curves for the subset of mouse tumors in panel C from the

SOMWR_EPIS_X00013aS1 PDX model. n = 3 mice per treatment group. Vardi *U*-test p = 0.10

and 9.4E-2 for combination vs. barasertib or TAZ, respectively. (**F**) Tumors from panel E

harvested on Day 135 of treatment.

Figure 5: Synthetic lethal combination strategy using ATR inhibition overcomes TAZ resistance and improves response: (**A, C**) G401 cells treated with elimusertib for 4 days (**C**).

(**B**) Panel of MRT and ES cell lines ordered left to right by decreasing response to TAZ
monotherapy. Cells were treated with the indicated monotherapy or combination for 11 days.
Drug concentrations used were: TAZ: 200 nM, elimusertib: 8 nM. We selected an elimusertib
dose below its monotherapy IC$_{50}$ (for G401 cells) in order to visualize any additive effects upon
combination with TAZ. (**C**) Representative images of G401 cells treated with the indicated
treatment for 7 days. (**D**) Quantification of γH2AX fluorescence relative to DAPI fluorescence. *p
= 0.042, **p = 5.1E-3, ***p = 1.7E-8, ****p = 6.5E-4 by two-sided Student's t-test. n = 548 nuclei
for DMSO, 432 for elimusertib, 696 for 50 nM TAZ, 401 for 50 nM TAZ + elimusertib, 373 for
100 nM TAZ, 63 for 100 nM TAZ + elimusertib. (**E**) Tumor growth curves for 5 mouse PDXs
treated with the indicated drug regimen. n = 20 mice for vehicle and elimusertib-treated groups,
n = 21 for TAZ and TAZ + elimusertib-treated groups. Vardi *U*-test p = 3.2E-2 and 0.23 for
combination vs. elimusertib or TAZ, respectively. (**F**) Kaplan-Meier curves showing tumor-free
survival (defined as tumor volume ≤ 1,000 mm^3) for the PDXs in panel C. Mean survival is 51
days (95% CI: 42-60 days) for elimusertib, 67 days (95% CI: 53-81 days) for TAZ to 99 days
(95% CI: 74-123 days) for the combination. Log-rank test p = 5.8E-4 and 3.9E-2 for combination
versus elimusertib or TAZ, respectively. (**G**) Tumor growth curves for the
HYMAD_EPIS_X0004aS1 PDX model treated with the indicated drug regimen. Vardi *U*-test p =
2.0E-4 for combination versus elimusertib or TAZ. n = 14 mice per treatment group. (**H**) Kaplan-
Meier curves showing tumor-free survival (defined as tumor volume ≤ 1,000 mm^3) for the PDXs
in panel E. Log-rank test p = 6.2E-3 and 6.3E-5 for combination versus elimusertib or TAZ,
respectively.
971
972

973
974 **Figure 6: Mechanistic schematic for the response of BAF-deficient tumors to effective**
975 **EZH2 therapy: Step 1**: TAZ inhibits histone methylation activity of PRC2. **Step 2:** Activating
976 chromatin-bound complexes, such as ncBAF bind to tumor suppressor loci. **Step 3:** Tumor

977   suppressor loci are activated by their transcription factors and their coactivators. **Step 4:** Tumor

978   suppressors inhibit cell cycle progression through their downstream effectors, such as RB1/E2F.