Individual differences in developmental trajectory leave

a male polyphenic signature in bulb mite populations

Jacques A. Deere[1*] & Isabel M. Smallegange[2*]

[1] Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, PO Box 94240, 1090

GE Amsterdam, the Netherlands

[2] School of Natural and Environmental Sciences, Newcastle University, Newcastle upon Tyne, NE1

7RU, UK

*Equal contribution

Corresponding author: Isabel M. Smallegange – Isabel.smallegange@newcastle.ac.uk

Abstract

Developmental plasticity alters phenotypes and can in that way change the response to selection.

When alternative phenotypes show different life history trajectories, developmental plasticity can

also affect, and be affected by, population size-structure in an eco-evolutionary interaction.

Developmental plasticity often functions to anticipate future conditions but it can also mitigate

current stress conditions. Both types of developmental plasticity have evolved under different

selections and this raises the question if they underlie different eco-evolutionary population

dynamics. Here, we tested, in a long-term population experiment using the male polyphenic bulb

mite (*Rhizoglyphus robini*), if the selective harvesting of juveniles of different developmental stages

concurrently alters population size (ecological response) and male adult phenotype expression

(evolutionary response) in line with eco-evolutionary predictions that assume the male polyphenism

is anticipatory or mitigating. We found that the frequency of adult males that expressed costly

(fighter) morphology was lowest under the most severe juvenile harvesting conditions. This

response cannot be explained if we assume that adult male phenotype expression is to anticipate

adult (mating) conditions because, in that case, only the manipulation of adult performance would

have an effect. Instead, we suggest that juveniles mitigate their increased mortality risk by

expediating ontogeny to forego the development of costly morphology and mature quicker but as a

defenceless scrambler. If, like in mammals and birds where early-life stress effects are extensively

studied, we account for such pre-adult viability selection in coldblooded species, it would allow us to

(i) better characterise natural selection on trait development like male polyphenisms, (ii) understand

how it can affect the response to other selections in adulthood, and (iii) understand how such trait

dynamics influence, and are influenced by, population dynamics.

Keywords: alternative male phenotypes, dispersal, eco-evolutionary dynamics, male morph

coexistence, polyphenism

Introduction

Across the animal kingdom, developmental plasticity comprises a switch to alternative

developmental pathways that depends on an individual's condition – the resource budget available

for the production and maintenance of adaptive traits (Hill 2011; Casasa et al. 2020; Nijhout &

McKenna 2018) – and which can lead to food-dependent allometric plasticity and the adaptive

expression of alternative phenotypes in adulthood (polyphenism) (West-Eberhard 2003; Nijhout &

McKenna 2018). For example, many intraspecific, arthropod male dimorphisms comprise large

majors that have weapons for male-male competition, and smaller, weaponless minors that sneak

matings (Moczek 2003; Smallegange 2011; Buzatto & Machado 2014). Importantly, these adaptive,

condition-dependent alternative phenotypes typically have different survival, growth and

reproduction rates because they follow different developmental trajectories (West-Eberhard 2003;

Smallegange 2011; Weir et al. 2016). Differences in individual development, be they plastic or

genetic, will thus affect a population's age or life stage structure, size and growth. Such population

changes can in turn influence individual development through processes like density-dependent

competition over resources (food) and density-dependent selection (Travis et al. 2013), creating eco-

evolutionary feedbacks at the population level (Smallegange & Deere 2014; Croll et al. 2019;

Smallegange 2022). However, accurately predicting the population dynamical responses to

perturbation of developmental trajectories is still difficult (Smallegange et al. 2019; Smallegange

2022).

The way in which a perturbation to individual development can impact the eco-evolutionary

dynamics of populations depends on what selection pressures are assumed to act on adaptive

developmental plasticity (Smallegange 2022). Different selection pressures determine different

types of adaptive developmental plasticity (e.g., Forsman 2015; Nettle and Bateson 2015) and here

we focus on two distinct types: anticipatory and mitigating adaptive developmental plasticity

(Smallegange et al. 2019; Smallegange 2022). Specifically, in case of the minor and major males

described above, whether a male develops into an armed major male or an unarmed minor male

depends on whether it reaches a critical resource budget threshold, assumed to be approximated by

a body size threshold, during development (Plaistow et al. 2005; Sasson et al. 2016; Cotton et al.

2004). If the adaptive developmental plasticity is *anticipatory*, this condition-cued threshold

mechanism can be interpreted as the evolutionary result of disruptive sexual selection on alternative

mating tactics (West-Eberhard 2003; Smallegange et al. 2019): good-condition juvenile males

maximise fitness by becoming majors, anticipating competitive mating success; bad-condition

juveniles anticipate failure at competitive success and salvage fitness by becoming minors (Shuster &

Wade 2003). On the other hand, the condition-cued threshold mechanism could be part of a

heritable developmental system that adaptively mitigates stressful environmental conditions

(Badyaev 2005, Del Giudice et al. 2018; Ellis & Del Giudice 2019). Such *mitigating* adaptive

developmental plasticity is the result of natural selection that favours developmental systems that

tend to construct phenotypes that are successful, relative to other variant systems, at surviving and

reproducing (Griffiths & Gray 1994). This means that juvenile males can mitigate stress during

maturation by refraining from developing costly morphology to maintain fitness and mature and

survive as an adult minor, as opposed to risk dying during maturation when investing in costly major

morphology (Smallegange et al. 2019). To summarise, in both kinds of adaptive plasticity, the fitness

functions of the alternative male phenotypes cross over a condition gradient, but anticipatory

adaptive developmental plasticity assumes that the threshold for alternative male phenotype

expression at the cross-section of the fitness functions is evolutionarily regulated by how adult males

perform, whereas mitigating adaptive developmental plasticity assumes that the threshold is

regulated by the dynamics of population density, food competition and individual resource budgets.

The fitness functions of both types of developmental plasticity are formalised in the

environmental threshold (ET) model (Hazel et al. 1990; 2004). Under the ET model, the alternative

expression of phenotypes has been hypothesized to be an environmentally cued threshold trait. It is

assumed that threshold traits are based on a continuously distributed, polygenic trait, called the

'liability' and a threshold of expression such that individuals that are above this threshold express

one phenotype while those below the threshold express the alternative (Roff 1996). Candidate traits

for the liability are, for example, hormone profiles (Roff 1996). The position of the threshold

depends on a cue that reliably correlates with the status of the environment, which in many taxa

represents condition (Tomkins et al. 2011; Smallegange & Deere 2014). The ET model assumes that,

in response to environment-specific selection, alternative phenotypes have evolved different fitness

functions, through which selection can affect the distribution of individual liabilities. Because

alternative phenotype frequency depends on the distribution of individual liabilities and the cue

distribution, both are taken into account in determining how phenotype fitness influences the

evolution of liabilities and hence the evolution of a threshold trait. The threshold thus evolves if the

fitness functions change in response to changing selection(s), evolutionary impacting alternative

phenotype expression. Furthermore, the cue (here: condition) distribution within the ET model

informs on population size and structure (Smallegange & Coulson 2013). If the cue (here: condition)

distribution, and thus population size and structure, changes, it ecologically impacts what proportion

of individuals expresses which alternative phenotype (e.g., a reduction in overall condition reduces

the frequency of males that can reach a high enough condition to develop into a fighter). Thus, the

ET model gives us the ingredients to create predictions on how changes to the fitness functions and

condition distribution of anticipatory or mitigating developmental plasticity fuel eco-evolutionary

population change.

We conducted a long-term population experiment to test hypotheses on how the selective

removal of juveniles of different condition in populations of the male polyphenic bulb mite

Rhizoglyphus robini impacts eco-evolutionary population responses through anticipatory and

mitigating developmental plasticity, under the assumptions of the ET model. Adult fighter males of

the bulb mite possess a proportionally thickened third leg pair with dagger-like claws that are used

in fights to kill rival males (Radwan et al. 2000). Whereas fighter males metamorphose from good-

condition juveniles that have a large resource budget, juveniles that are in a bad condition (i.e., have

a small resource budget) when metamorphosing into an adult instead express the scrambler

phenotype that does not have such leg modifications (Rhebergen et al. 2022). Note that male adult

phenotype expression does not depend on pheromone cues that inform on the frequency of each

male phenotype in the population (Deere & Smallegange 2014) or mite density (Radwan 1995).

Further, in response to unfavourable environmental conditions (*e.g.*, low food), *R. robini* expresses a

facultative dispersal morph during development, called the deutonymph, which does not feed and is

adapted to phoretic dispersal (Díaz et al. 2000). Deutonymph expression occurs mid-development,

prior to the final instar stage from which juveniles metamorphose into adults (Díaz et al. 2000), and,

thus, prior to when adult male phenotype is determined (which is immediately before

metamorphosis [Rhebergen et al. 2022]). Because deutonymph expression carries developmental

costs, including reduced size at metamorphosis in both sexes and reduced egg production in females

(Deere et al. 2015), our working assumption is that only individuals that are of sufficiently good

condition can develop into one if cued by poor environmental conditions. This assumption is

supported by the fact that in two of our previous studies (Smallegange & Coulson 2011; Deere et al.

2015), male deutonymphs always matured as a fighter, from which we surmise that males always

were in sufficiently good condition to develop costly fighter legs.

In our experiment, we selectively removed individuals of different conditions by removing

deutonymphs, or other juveniles. From the ET model, assuming that the male polyphenism is

anticipatory, we predict that the ecological response of removing deutonymphs from the population

is that it will lower the righthand side of the distribution of individual conditions (because individuals

of relatively good condition are removed), and thus population size (compare blue dotted

distribution size [before harvesting] with red dashed distribution size [after harvesting for several

generations] in Fig. 1c). However, we expect that this will not elicit an evolutionary response

because male fitness is assumed to be unaffected by juvenile performance (it is only affected by

adult male performance). Thus, we expect that the mean threshold for fighter expression, and thus

the proportion of males that are fighters, will track the latter change in the condition distribution

129 **Figure 1. Eco-evolutionary changes due to the**
130 **selective harvesting of deutonymphs and other**
131 **juveniles.** According to the ET model (Hazel et al.
132 1990; 2004), fitness functions of fighters (black lines)
133 and scramblers (grey lines) cross over a condition
134 gradient; the area under each condition distribution
135 curve represents population size. The proportion of
136 individuals adopting alternative phenotypes
137 (contrastingly shaded areas under the curves: light
138 grey for scramblers and dark grey for fighters) is
139 determined by the threshold (t^*) condition value at
140 the intersection of the fitness functions (indicated by
141 a vertical line). Blue dotted lines and red dashed lines
142 represent condition distributions before (**a,b**) and
143 after several generations of harvesting (**c-f**),
144 respectively. **(c)** Although the selective harvesting of
145 deutonymphs will reduce the size of the condition
146 distribution (i.e., population size), anticipatory
147 developmental plasticity predicts that the fitness
148 functions, and, thus, the threshold (t^*) for fighter
149 expression, will track the change in the condition
150 distribution and remain in the same, position relative
151 to the condition distribution. In this example, the
152 threshold tracks the mean of the condition
153 distribution. (**e**) Assuming that the harvesting of
154 juveniles reduces population size but does not

155 change mean condition, the threshold (t^*) for fighter expression remains unaffected in case of anticipatory
156 developmental plasticity. (**d,f**) In contrast, the stress mechanism of a developmental system can facilitate the
157 mitigating response to increased deutonymph or juvenile mortality, because males can mature early as a
158 minor to escape the juvenile stage quickly. This will increase scrambler fitness, fuelling the evolution of
159 scrambler expression to a higher threshold (t^*) for fighter expression. Additionally, because scramblers mature
160 earlier in life (Smallegange 2011), live longer than fighters (Radwan & Bogacz 2000) and sire more offspring
161 than fighters (van den Beuken et al. 2019), an increase in scrambler expression can increase population size
162 (Smallegange & Deere 2014), and thus increase the size of the condition distribution.

163 because relative, and not absolute condition is important in determining alternative phenotype

164 fitness (Tomkins et al. 2011) (compare the proportions of the differently shaded areas under the

165 condition distribution in Fig. 1a [before harvesting] with Fig. 1c [after harvesting for several

166 generations]). Likewise, we predict that the selective removal of juveniles (bar deutonymphs) of

167 random condition has no effect on fighter expression (compare the proportions of the differently

168 shaded areas under the condition distribution in Fig. 1a [before harvesting] with Fig. 1e [after

169 harvesting for several generations]), because fitness functions remain unaltered. The reduction in

170 population size because of the harvesting of juveniles, however, will be reflected in an overall

reduced size of the condition distribution (compare blue dotted distribution size [before harvesting]

with red dashed distribution size [after harvesting for several generations] in Fig. 1e).

In contrast, according to the ET model and assuming that the male polyphenism is

mitigating, the stress mechanism of the developmental system can fuel a mitigating response to

increased deutonymph and juvenile mortality because males can mature early as a minor to escape

the juvenile stage quickly (Ernande et al. 2004), fuelling the evolution of the threshold for scrambler

expression. If scrambler fitness increases relative to that of fighters, the threshold for fighter

expression will evolve to increase, both in response to deutonymph harvesting (compare the

proportions of the differently shaded areas under the condition distribution in Fig. 1b [before

harvesting] with Fig. 1d [after harvesting for several generations]) and in response to the selective

harvesting of other juveniles (compare the proportions of the differently shaded areas under the

condition distribution in Fig. 1b [before harvesting] with Fig. 1f [after harvesting for several

generations]). Further, because scramblers mature earlier in life (Smallegange 2011), live longer than

fighters (Radwan & Bogacz 2000) and sire more offspring than fighters (van den Beuken et al. 2019),

an increase in scrambler expression can increase population size (Smallegange & Deere 2014). This

ecological response will increase the size of the condition distribution (compare blue dotted

distribution size [before harvesting] with red dashed distribution size [after harvesting for several

generations] in Fig. 1d and 1f). But note that the removal of good-condition juveniles when removing

deutonymphs will result in the right-hand side of the distribution to be at lower condition values

(compare blue dotted distribution size [before harvesting] with red dashed distribution size [after

harvesting for several generations] in Fig. 1d).

Methods

Study system

The bulb mite has six life stages: egg, larva, protonymph, deutonymph, tritonymph and adult (Fig. 2). The deutonymph stage is a facultative, non-feeding dispersal stage and mites only develop into this stage during unfavourable conditions (i.e. low food quality and quantity) (Díaz et al 2000). When growing from one stage to the next, mites moult; this stage is known as the quiescent stage which is immobile and non-feeding. Source populations were established from mites collected in 2010 from flower bulbs in storage rooms in North Holland. In our source populations, mites rarely develop into a deutonymph and we previously estimated the percentage of males developing into one after removing food and imposing a dry period at 3% (Smallegange & Coulson 2011). Mites were kept on an oats diet and maintained as described in Smallegange (2011).

Figure 2. Life cycle of the bulb mite *Rhizoglyphus robini*. The bulb mite has six life stages: egg, larva, protonymph, deutonymph, tritonymph and adult. The deutonymph stage is a facultative, non-feeding dispersal stage and mites only develop into this stage during unfavourable conditions (i.e., low food quality and quantity) (Díaz et al 2000). When growing from one stage to the next, mites moult; this stage is known as the quiescent stage which is immobile and non-feeding. Show are ventral and dorsal images for each stage.

Experimental treatments were established from these source populations. Individuals across

populations are highly inbred except for scrambler males in food-limited environments (Stewart et

al. 2019). Inbred fighter males arise due to a combination of mating monopolization and increased

survival, compared to scrambler males, which limit the genetic pool in the resulting fighter offspring

(Stewart et al. 2019). The driver for inbreeding in females is likely driven by intralocus sexual conflict

given the mating monopolization of fighter males (Stewart et al. 2019).

Experimental procedure

The experiment ran between June 2016 and April 2017 (302 days) and comprised five treatments

(Fig. 3): weekly harvesting of (i) 100% of all deutonymphs (D100), (ii) 50% of deutonymphs (D50), (iii)

other juveniles (excluding eggs and deutonymphs) at the same percentage at which deutonymphs

were harvested from all juveniles in D100 (J-D100) to keep the demographic impact of harvesting

comparable over the length of the experiment, (iv) other juveniles (excluding eggs and

deutonymphs) at the same percentage at which deutonymphs were harvested from all juveniles in

D50 (J-D50), again, to keep the demographic impact of harvesting comparable over the length of the

experiment, and (v) no harvesting (C; general control treatment). For the removal of 'other juveniles'

we first calculated the average number of deutonymphs removed, across all three replicates, in a

deutonymph removal treatment (D100/D50). Second, we calculated the average number of

protonymphs and tritonymphs (i.e., other juveniles), across all three replicates, in the same

deutonymph removal treatment. From these values we calculated a deutonymph proportion value:

proportion value = average deutonymphs/(average protonymphs + average tritonymphs). Finally, for

a treatment where other juveniles were removed (J-D100/J-D50), we multiplied the deutonymph

proportion value calculated for a deutonymph removal treatment (D100/D50) to the sum of the

protonymph and tritonymphs within each replicate. In the case of the 'other juveniles' treatments,

juveniles (i.e., protonymphs and tritonymphs) were removed randomly (see further Fig. S6 in the

online appendix). Because deutonymph numbers are generally low in our source populations

(Smallegange & Coulson 2011), one treatment was to remove all deutonymphs. Two levels of

harvesting deutonymphs (D100, D50) and juveniles (J-D100 and J-D50) were chosen to be able to

compare harvesting impacts. Because we expect deutonymph numbers to be low, this means that in

the other juvenile harvesting treatments, we expect to harvest low numbers as well. We anticipate

that these harvesting regimes will still induce population responses. For example, in previous

experiments, we also imposed harvesting

241
Figure 3. Experimental design. The photos show the length measurements of adult females (top left), adult
males (top right), deutonymphs (bottom left) and quiescent individuals (bottom right). The schematic shows
the experimental design and harvesting treatments.

regimes within which we removed only a few individuals on a regular basis and found these

selections to have significant impacts on the threshold for male adult phenotype expression and

population size-structure (Smallegange & Deere 2014; Smallegange & Ens 2018). Treatments were

replicated three times resulting in fifteen populations. The juvenile harvesting populations were

started 8 weeks after the other treatments, but before harvesting commenced (Fig. 4), for logistical

reasons. At the point where the juvenile harvesting populations matched the total population sizes

of the other treatments (Fig. 4), harvesting started. Populations were kept in 20 mm diameter, flat-

bottomed glass tubes with a plaster of Paris and powdered charcoal base, which was kept moist to

avoid desiccation of the mites. Tubes were sealed by a circle of very fine mesh (allowing gaseous

diffusion), which was held in place by the tubes' standard plastic caps with ventilation holes cut into

them. Populations were kept in an unlit climate chamber at 25°C and >70% relative humidity.

Figure 4. Time series of mean total population size, averaged per harvesting treatment. The treatments
were: harvesting of 100% of all deutonymphs (D100: grey line), harvesting of 50% of deutonymphs (D50: black
line), harvesting of other juveniles (excluding deutonymphs) at the same percentage at which deutonymphs
where harvested from all juveniles in D100 (J-D100: purple line), harvesting of other juveniles (excluding
deutonymphs) at the same percentage at which deutonymphs where harvested from all juveniles in D50 (J-
D50: indigo line), and no harvesting (C; control treatment: orange line). The harvesting treatments began in
week 13 of the experiment. No data were collected in week 29 of the experiment due to illness. Vertical grey
lines denote the beginning and end of each time period over which we averaged results (see main text).

Each population was established with 50 randomly selected mites from the source

population, which we refer to as founder mites, which we removed after 15 days to reduce founder

effects (founder mites can be easily distinguished because they are much bigger than the first

generation emerging after 15 days in the experimental population). Each population was fed

approximately 1mg of oats per day (the weight was never less than 0.9g and never exceeded 1.1g).

After population initialization, on each Thursday, mites of each life stage in all population were

counted using a hand counter at 15 X magnification. For three weeks at the start of each time period

(day 85-161, day 162-238 and day 239-302: see statistical analyses below), photos were taken of up

to 5 (average: 4.05 ± 1.46 SD) individual deutonymphs, fighters, scramblers, and females and their

sizes measured as proxies of condition (Rhebergen et al. 2022) to assess any shifts in mean condition

across the different life stages in response to experimental conditions. The body length of mites on

each photo was measured to the nearest 0.1 μm using ZEN software blue edition (ZEISS microscopy,

Carl Zeiss, Germany). We measured body length for adult mites as the ventral distance of the

idiosoma (the white bulbous posterior part of the body; from the mouthparts to the furthest part of

the genitalia); for deutonymphs as the dorsal anterior-posterior length (without mouthparts); and

for quiescent tritonymphs as the lateral body length (without mouthparts) (Fig. 3). Harvesting

treatments started 85 days after populations were initialized, which is a sufficiently long time for

population numbers to stabilise (which typically happens at 40 days after initialisation [Cameron and

Benton 2004; Smallegange and Deere 2014]) (see also Fig. 1). Harvesting was done once a week (on

Thursday), after counting and before feeding. The number of individuals harvested in the control

populations (J-D100 and J-D50) were calculated based on the proportion of deutonymphs removed

from the D100 and D50 populations respectively. The proportion of deutonymphs removed was

calculated relative to the combined number of protonymphs and tritonymphs. This proportion value

was then applied to the combined number of protonymphs and tritonymphs in the relevant control

populations (proportion value for D100 (D50) was applied to J-D100 (J-D50)). The proportion value

applied to the populations calculated the total number of individuals to remove from the J-D100 and

J-D50 populations. Only protonymphs and tritonymphs were removed from the control populations

and this was done randomly. The total protonymphs and deutonymphs removed for each population

for each week can be found in data file in the online appendix (Smallegange and Deere 2023). The

harvesting treatments were applied for a total of 217 days, which in a similar experiment was

sufficient to observe an evolutionary shift in fighter expression (Smallegange & Deere 2014).

However, even if this period of time is too short to observe an evolutionary response to harvesting,

we can still distinguish between our anticipatory and mitigating hypotheses. Specifically, under

juvenile harvesting (J-D100 and J-D50), fighter expression is predicted to remain unaltered if it is

anticipatory (Fig. 1e). However, fighter expression is predicted to decrease if it is mitigating (Fig. 1f)

because juvenile males can respond quickly (within a generation) to increased juvenile mortality by

expediating ontogeny and mature as a scrambler. This plastic (ecological) response, in turn, can

further fuel the evolution towards developmental systems that produce scramblers in response to

the juvenile harvesting selection pressure. What is important to note is that the ET model states that

any change in the threshold for alternative male phenotype expression will affect the proportion of

individuals developing either phenotype because it is expected to track the intersection of the

alternative phenotype fitness functions. Therefore, we can interpret evolutionary shifts in the

threshold from evolutionary changes in fighter expression (the proportion of adult males that are

fighters).

Life-history assays

Following the end of the long-term population experiment we conducted a common garden life

history assay to assess if any differences in body size (which we assume to approximate condition)

and fighter expression between treatments were plastic or genetic. Logistical constraints prevented

us from conducting the time-consuming and labour-intensive assays to measure the threshold for

alternative male phenotype expression itself. For each assay we removed ten adult females from

each population, individually isolated them and gave them ad lib access to oats and allowed them to

lay eggs. Three of the ten females (isolating more individuals was logistically not possible) were then

randomly selected and their offspring individually isolated, given ad lib access to oats, and followed

until they reached maturity. Deutonymphs, quiescent tritonymphs and adults (2-5 days after

maturation) were size-measured as above and their sex and morph scored (we measured a total of

420 individuals). The effects of maternal nutritional conditions on size, age and adult male

phenotype are negligible in this species (the effect size of offspring environment is 15 times larger

than that of maternal environment: Smallegange 2011a), therefore, rearing mites in a common

garden environment condition for one generation is sufficient to eliminate maternal effects. Females

and their offspring were kept in 10 mm diameter tubes with a plaster of Paris and powdered

charcoal base, which were kept in an unlit climate chamber at 25°C and >70% relative humidity.

Statistical analyses

To analyse the results from the population experiment, we only used data from day 85 onwards

(when harvesting commenced) and excluded two population tubes that had fallen over during the

population experiment (one tube in treatment D100 and one in treatment J-D50). We then divided

the experimental period into three periods, day 85-161, day 162-238 and day 239-302 to assess

long-term (instead of transient) temporal changes in the response variables within each treatment

group. To analyse the results of the population experiment, we used a generalised linear mixed

model (GLMM) with Gaussian errors to analyse the effect of the fixed factors harvesting treatment

(C, D100, D50, J-D100, J-D50) and time period (period 1, 2, and 3) and their interaction, including

'population tube' as a random effect (to account for the repeated measures within each

experimental population) on: fighter expression (the number of fighters with the number of adult

males set as an offset so that the model prediction will be proportion of fighter males), and the

mean size of deutonymphs, adult fighters, scramblers and females. The model assumptions of

Gaussian errors and homoscedacity were confirmed by inspecting the probability plots and error

structures. We used a GLMM with Poisson errors to analyse the effects of the fixed factors

harvesting treatment and time period, their interaction, with 'population tube' as a random effect,

on the number of deutonymphs and on total population size (all individuals). In the online appendix

(Smallegange and Deere 2023) we present the analysis of treatment effects on the number of

juveniles and female and male adults.

To analyse the results of the life history assay conducted at the end of the population

experiment, we used a GLMM with Gaussian errors to analyse the effect of the fixed factor

harvesting treatment (C, D100, D50, C-D100, C-D50), including 'population tube' and 'maternal

identity' as random effects on the mean size of adult fighters, scramblers and females. The model

assumptions of Gaussian errors and homoscedacity were again confirmed by inspecting the

probability plots and error structures. We used a GLMM with binomial errors to analyse the effect of

the harvesting treatment on fighter expression ('0' if a male developed into a scrambler, and '1' if in

a fighter), including 'population tube' and 'maternal identity' as random effects.

In each GLMM, the significance of the treatment effects was tested using a model

simplification procedure, whereby the full model was fitted after the least significant term had been

removed (never deleting random terms from the model), if the deletion caused an insignificant

increase in deviance as assessed by a maximum likelihood ratio test (LRT). Pairwise comparisons

among the treatment levels were conducted by creating a reduced model within which two levels

were combined. If the reduced model caused an insignificant increase in deviance as assessed by a

maximum likelihood ratio test ($\alpha = 0.05$), then the two levels were considered to be not significantly

different. All analyses were conducted in R and for all GLMMs we used the *lme4* package (www-r-

project.org).

Results

Population size, fighter expression and number of deutonymphs

Mean total population size slightly increased over time since we started harvesting (Fig. 4).

Indeed, mean population size significantly differed between time periods (χ^2_2 = 262.07, p<0.001)

(non-significant interaction: χ^2_8 = 7.30, p=0.063) and was also significantly affected by harvesting (χ^2_4

= 15.58, p=0.004). Population size was highest when 100% of deutonymphs were harvested (D100),

lower when 50% of deutonymphs were harvested (D50), and lowest in all the other treatments (C, J-

D50 and J-D100) (treatments C and J-D50 did not differ significantly (χ^2_1 = 0.35, p=0.553), and neither

did C and J-D50 differ significantly from J-D100 (χ^2_1 = 0.47, p=0.494)) (Fig. 5A). Irrespective of

harvesting treatment, total population size significantly increased with increasing time period (Fig.

5B). The harvesting treatment significantly affected the proportion of fighters but differently at

different time periods (significant interaction harvesting $\times$ time period: χ^2_8 = 29.92, p<0.001). Within

the interaction, the harvesting treatments C and J-D50 did not significantly differ in their effect on

proportion of fighters in the different time periods (χ^2_3 = 4.22, p=0.239) and were combined into one

level. The next two similar responses to the harvesting treatments J-D100 and D50 did, however,

differ significantly (χ^2_3 = 11.80, p=0.008) (Fig. 5C). In summary, the proportion of fighters was always

relatively the lowest when 100% of deutonymphs were harvested (D100 treatment) (Fig. 5C), but

when other juveniles were harvested at the same percentage as in D100, (J-D100), and when

harvesting 50% of deutonymphs (D50), the proportion of fighters was relatively high in the first time

periods, but lower in the last time period (Fig. 5C). In contrast, under the combined harvesting

treatments of C and J-D50, the proportion of fighters was always high over the whole course of the

experiment (Fig. 5C). In the online appendix we present the number of fighters and scramblers for

each harvesting treatment (Smallegange and Deere 2023).

Figure 5. Population size and structure of the long-term population experiment. Mean total population size differed between different harvesting treatments (a) and between the different time periods (b). Fighter expression varied over time in most but not all harvesting treatments (c). The number of deutonymphs also varied over time in some but not all harvesting treatments (d). Errors in all panels are standard errors. Harvesting treatments were: harvesting of 100% of all deutonymphs (D100: solid black lines), harvesting of 50% of deutonymphs (D50: dashed black lines), harvesting of other juveniles (excluding deutonymphs) at the same percentage at which deutonymphs where harvested from all juveniles in D100 (J-D100: grey dashed line (c) and solid grey line (d)), harvesting of other juveniles (excluding deutonymphs) at the same percentage at which deutonymphs where harvested from all juveniles in D50 (J-D50: solid grey lines), and no harvesting (C; control treatment: solid grey lines). Treatment levels that did not significantly differ from each other are grouped (see main text). Note that y-axes do not start at zero.

The harvesting treatment also significantly affected the number of deutonymphs, but, again, differently at different time periods (significant interaction harvesting × time period: χ_8^2 = 49.70, p<0.001). Within the interaction, the harvesting treatment levels J-D100 and C did not significantly differ (χ_3^2 = 0.40, p=0.940) and were combined into one level. Likewise, the latter combined level did not significantly differ from the J-D50 harvesting treatment (χ_3^2 = 3.08, p=0.380) and all three control levels were combined into one level. These three levels, combined, significantly differed from the D50 harvesting level (χ_3^2 = 12.42, p=0.006) and from the D100 harvesting level (χ_3^2 = 67.71, p<0.001). We therefore find that, counterintuitively, the number of deutonymphs was higher in the last time

period than in the first one when all deutonymphs were harvested (D100), and, to a lesser extent,

when 50% of deutonymphs were harvested (D50) (Fig. 5D). In contrast, in the other treatments

combined, the number of deutonymphs was slightly lower in the last time period than in the first

(Fig. 5D). Furthermore, at any given time period, the number of deutonymphs was always the

highest in the treatment where weekly harvesting removed all deutonymphs (D100) (Fig. 5D). In the

online appendix (Smallegange and Deere 2023), we present interaction plots for the three response

variables.

Deutonymph, fighter, scrambler and adult female mean size

Mean deutonymph size was not significantly affected by harvesting (χ^2_4 = 4.81, p=0.307) but

decreased with each consecutive time period (significant effect of time period: χ^2_2 = 106.31, p<0.001)

(Fig. 6A) (non-significant interaction: χ^2_8 = 7.22, p=0.513). Mean fighter size showed a similar

response but size differed significantly between the different harvesting treatments (significant

interaction harvesting × time period: χ^2_8 = 19.44, p=0.013) (Fig. 6B). Within the interaction, fighter

size in the harvesting treatment levels J-D50 and J-D100 did not significantly differ (χ^2_3 = 2.33,

p=0.507) and were combined into one level (Fig. 6B). Similarly, fighter size in the harvesting

treatment levels C and D50 did not significantly differ (χ^2_3 = 1.83, p=0.609) and were combined into

one level (Fig. 6B). The overall result was that mean fighter size in the combined levels J-D50 and J-

D100 and in the combined levels C and D50 was lower in the last than in the first time period,

whereas mean fighter size in the D100 treatment did not decrease until the last time period (Fig. 6B).

Mean scrambler size was not significantly affected by harvesting (χ^2_4 = 1.17, p=0.883) but was

significantly lower in each consecutive time period (significant effect of time period: χ^2_2 = 96.44,

p<0.001) (Fig. 6C) (non-significant interaction: χ^2_8 = 9.48, p=0.303). Finally, mean adult female size

was also not significantly affected by harvesting (χ^2_4 = 0.62, p=0.961) but significantly decreased with

each consecutive time period (significant effect of time period: χ^2_2 = 67.65, p<0.001) (Fig. 6D) (non-

significant interaction: χ_8^2 = 14.55, p=0.068). In the online appendix (Smallegange and Deere 2023),

we present interaction plots for the four response variables.

Figure 6. Mean body sizes observed in the long-term population experiment. Mean deutonymph size (μm)
(a), mean fighter size (μm) **(b)**, mean scrambler size (μm) **(c)** and mean female size (μm) **(d)** all significantly
decreased over the course of the experiment, but for fighters, this decrease differed between harvesting
treatments (d). Errors in all panels are standard errors. Harvesting treatments were: harvesting of 100% of all
deutonymphs (D100: solid black line), harvesting of 50% of deutonymphs (D50: dashed black line), harvesting
of other juveniles (excluding deutonymphs) at the same percentage at which deutonymphs where harvested
from all juveniles in D100 (J-D100: grey solid line), harvesting of other juveniles (excluding deutonymphs) at
the same percentage at which deutonymphs where harvested from all juveniles in D50 (J-D50: grey solid line),
and no harvesting (C; control treatment: dashed black line). Treatment levels that did not significantly differ
from each other are grouped (see main text). Note that y-axes do not start at zero.

Life history assay

Neither fighter size (χ_4^2 = 6.58, p=0.160, mean size: 439.76 μm $\pm$ 2.39 SE, n = 190) (Fig. 7A), scrambler

size (χ_4^2 = 2.77, p=0.597, mean size: 401.07 μm $\pm$ 7.33 SE, n = 14) (Fig. 7B) or female adult size (χ_4^2 =

3.75, p=0.440, mean size: 334.66 μm $\pm$ 1.65 SE, n = 206) (Fig. 7C) were affected by the harvesting

treatments in the common garden life history assay. Fighter expression was also not significantly

affected by harvesting treatment (χ_4^2 = 0.51, p=0.973) (Fig. 7D); however, only 14 scramblers

emerged in the life history assay (as opposed to 190 fighters), prohibiting any statistical inference.

Figure 7. Life history assay results. Mean fighter size (μm) (**a**), scrambler size (μm) **(b)**, female size (μm) **(c)**,
and fighter expression (proportion of adult males that are fighters) **(d)** for each harvesting treatment. Errors in
all panels are standard errors. Harvesting treatments were: harvesting of 100% of all deutonymphs (D100),
harvesting of 50% of deutonymphs (D50), harvesting of other juveniles (excluding deutonymphs) at the same
percentage at which deutonymphs where harvested from all juveniles in D100 (J-D100), harvesting of other
juveniles (excluding deutonymphs) at the same percentage at which deutonymphs where harvested from all
juveniles in D50 (J-D50), and no harvesting (C; control treatment). Note that y-axes do not start at zero and
differ between panels.

Discussion

We tested two hypotheses on how developmental plasticity impacts the eco-evolutionary response

of male polyphenic bulb mite populations to the selective removal of individuals of different

developmental stages. If alternative male phenotype expression is taken to be anticipatory

(anticipating future mating success), we predicted that the selective harvesting of deutonymphs and

other juveniles would not impact fighter expression. In contrast, if alternative male phenotype

expression is taken to be mitigating, we predicted that the selective harvesting of deutonymphs and

other juveniles would decrease fighter expression both plastically, as male juveniles can expediate

development and escape the juvenile stage quicker by maturing as a scrambler, and evolutionarily,

as the selective removal of juveniles fuels the evolution of developmental systems that produce

scramblers. In line with the mitigating hypothesis, we found that fighter expression was overall

lowest when 100% of all deutonymphs were harvested (D100), and, at the end of the experiments,

had reached similarly low levels in the treatments where 50% of deutonymphs were harvested (D50)

and where the highest number of other juveniles were harvested (J-D100). This 'delayed' response

could be due to that fact that our proportional harvesting treatments became more severe as

population size increased over the course of the experiment, whereas per capita food availability

would also have decreased with increasing population size, potentially reducing fighter expression.

In contrast, fighter expression remained high throughout the experiment in the control treatment

and in the treatment where the lowest number of juveniles were harvested (J-D50), perhaps

because, there, the harvesting impact was the lowest. Our life history assay at the end of the

population experiment failed at showing any sign of evolutionary differentiation between our

treatments, which could be due to low statistical power, no evolution taking place or insufficient

initial genetic diversity. Also, if the overall observed reduction in body size (condition) affected

deutonymph numbers, this plastic response could have reduced the strength of the selection

imposed by our selective harvesting of deutonymphs. However, it is striking that the harvesting

treatments where we observed the lowest levels of fighter expression in the population experiment

are the same as where we observed the lowest fighter expression levels in the life history assay. But

even if this pattern was coincidental and not indicative of an evolutionary change in fighter

expression, our findings still point to a role for mitigating developmental plasticity. That is, we

expected no shift in fighter expression at all in response to the harvesting of juveniles (bar

deutonymphs) under anticipatory developmental plasticity (Fig. 1e), but a reduction in fighter

expression under mitigating developmental plasticity (Fig. 1f) because individuals mature earlier as a

scrambler to escape the risky juvenile stage. The latter reduction is indeed what we observed

towards the end of the experiment in the treatment where we harvested the highest number of

juveniles other than deutonymphs (J-D100). Our results therefore suggest that nutrition-deprived

male juveniles do not 'make the best of a bad job' by anticipating a non-fighting mating tactic

(Dawkins 1980; Eberhard 1982), but instead develop into that male adult phenotype because they

can adaptively mitigate their early-life stress (Rhebergen 2022). Such adaptive early-life stress

responses that have negative effects in adulthood are common in (warm-blooded) mammals and

birds (Lindström 1999; Fisher et al. 2006; Wells 2011; Vickers 2014), likely because their growth

during development is physiologically constrained. However, if we account for such pre-adult

viability selection in coldblooded species that typically show flexible growth, it would allow us to

better characterise natural selection on trait development like male polyphenisms in these species,

how it can affect the response to other selections in adulthood (e.g., sexual selection: Mojica and

Kelly 2010; Lürig & Matthews 2021; Rhebergen 2022), and how such trait dynamics influence, and

are influenced by, population dynamics (Smallegange 2022).

One unexpected finding in our experiment was that the number of deutonymphs was

highest when we harvested all of them, each week. Further, despite being harvested, deutonymph

number increased as the experiment progressed. We can only speculate why this happened.

Perhaps this overcompensatory response was because the per capita food availability increased as

we removed individuals (Verhulst 1838, Schaefer 1954, Hilborn and Walters 1992), although we still

only maximally removed a small percentage of the total population in the deutonymph harvesting

treatments, and we did not see a similar overcompensatory response in the treatment where we

harvested the same percentage of other juveniles. In fact, total population size increased as well so

that per capita food intake rate would decline, cueing more individuals to develop into

deutonymphs. It could also be simply a numerical response because mean total population size

increased over the course of the experiment and thus there were more individuals that could

develop into a deutonymph. Finally, the higher number of deutonymphs we found could also reflect

eco-evolutionary dynamics in which another selection pressure, for example imposed by strong-

density dependence (population size was highest in the 100% deutonymph harvesting treatment),

opposed the effects of direct harvesting (Edeline & Loeuille 2021). Or, perhaps, because twice as

many females emerge from deutonymphs than males (fighters) (Deere et al. 2015; Stewart &

Smallegange in prep), removing deutonymphs reduces females and egg output, reducing density-

dependence. Either way, failure to account for either of these eco-evolutionary processes might

fundamentally hamper our ability to understand the expression of dispersal phenotypes. Another

explanation is that it was the actual lack of deutonymphs and their olfactory cues that triggered

deutonymph development. Olfactory cues are common in acarine mites. All acarines have a sensory

site on their forelegs known as the "foretarsal sensory organ" and olfaction is one of this site's

functions (Carr & Roe 2016). If the harvesting of deutonymphs reduced the concentration of

deutonymph olfactory cues, this could trigger more juveniles to develop into a deutonymph

because, with fewer deutonymphs around, competition for attachment sites on insects is lower

(deutonymphs disperse via phoresy on insects [Díaz et al 2000]). We realise, however, that this is

speculation and olfactory cues produced by deutonymphs have yet to be tested in *R. robini*. Finally,

in our previous studies (Smallegange & Coulson 2011; Deere et al. 2015), male deutonymphs always

matured as a fighter, but here, increases in deutonymph numbers did not coincide with increases in

fighter frequency. Perhaps this was because twice as many females emerge from deutonymphs than

males (fighters) (Deere et al. 2015; Stewart & Smallegange in prep) so that our statistical analyses

were unable to pick up significant differences in the few more fighters that emerged. It does

highlight, however, that we still have some way to go to unravel the intricate links between

deutonymph expression, alternative male phenotype expression and sex ratio and how they

together drive population dynamics.

Because a change in the distribution of a trait will almost always also change the size of the

distribution of that trait, it follows that trait change is accompanied by a change in population size. If

a trait is heritable, this constitutes eco-evolutionary change at the population level (Smallegange &

Coulson 2013). This is especially so in the case of alternative phenotypes where we expect a shift in

phenotype expression to be accompanied by a change in population size because the different

phenotypes often have different demographic rates. For example, in the case of bulb mite males,

scramblers mature earlier in life (Smallegange 2011), live longer than fighters (Radwan & Bogacz

2000) and sire more offspring than fighters (van den Beuken et al. 2019) so that an increase (or

decrease) in scrambler (or fighter) expression will increase population size. Indeed, we found the

highest population size in the harvesting treatment with the lowest level of fighter expression (D100:

100% deutonymph harvesting). We suspect that the increase in population size that we observed

over the course of the experiment is associated with the observed decrease in mean body size over

time in response to the strong, density-dependent food conditions, because smaller organisms

typically reproduce faster and can reach higher equilibrium population sizes than larger ones (e.g.,

Huryn & Benke 2007). Indeed, Gilbert et al. (2022) recently showed in an experiment with the protist

Tetrahymena pyriformis that mean body size decreased in response to increased density-

dependence and that this rapid change in mean body size had a larger effect on population density

than the reverse. Reductions in mean body size have been observed across a range of taxa in

response to adverse environmental change (Gardner et al. 2011; Sheridan & Bickford 2011). Often,

these reductions are associated with increased mortality rates (e.g., in red knots *Calidris canutus*

[van Gils et al., 2016]). What is more, when populations not only suffer from directional

environmental change but also from selective harvesting, the removal of good condition individuals

(of large size) can lead to population extinction (Knell & Martínez-Ruiz 2017). Thus, unravelling the

intertwined ecological and evolutionary dynamics of population size, body size, condition and

condition-dependent (alternative) phenotypes is integral to understanding eco-evolutionary

population responses to change (Edeline & Loeuille 2021). If our experiment would form the basis

for such further studies, we suggest using outbred populations and increasing sample size in the life

history assays to be able to assess any evolutionary shifts in the actual threshold for alternative male

phenotype expression, and to consider other harvesting treatments with potentially stronger

selective effects, like the selective removal of a high proportion of large versus randomly selected

juveniles.

Data availability statement

The raw data, the R scripts that we used to analyse the data, and the additional analyses on

population structure can be found on Figshare (Deere & Smallegange 2023).

References

Badyaev AV. 2005. Stress-induced variation in evolution: from behavioural plasticity to genetic
assimilation. Proc. R. Soc. B 272: 877–886.
Buzatto BA, Machado G. 2014. Male dimorphism and alternative reproductive tactics in harvestmen
(Arachnida: Opiliones). Behav. Processes 109: 2–13.
Carr AL, Roe M. 2016. Acarine attractants: Chemoreception, bioassay, chemistry and control. Pestic
Biochem Physiol 131: 60-79.
Casasa S, Zattara EE, Moczek AP. 2020. Nutrition-responsive gene expression and the developmental
evolution of insect polyphenism. Nature Ecol. Evol. 4: 970–978.
Cotton S, Fowler K, Pomiankowski A. 2004. Condition-dependence of sexual ornament size and
variation in the stalk-eyed fly *Cyrtodiopsis dalmanni* (Diptera: Diopsidae). Evolution 58: 1038–
1046.
Dawkins, R. 1980. Good Strategy or Evolutionarily Stable Strategy? In: Sociobiology: Beyond
Nature/Nurture? (G. W. Barlow & J. Silverberg, eds), pp. 331–367. Westview Press, Boulder, CO
Deere JA, Coulson T, Smallegange IM. 2015. Life history consequences of the facultative expression
of a dispersal life stage in the phoretic bulb mite (*Rhizoglyphus robini*). PLoS ONE 10: e0136872.
Deere JA, Smallegange IM. 2014. Does frequency-dependence determine male morph survival in the
bulb mite, *Rhizoglyphus robini*? Experimental and Applied Acarology 62: 425-436
Deere JA, Smallegange IM. 2023. Data and analysis of "Individual differences in developmental
trajectory leave a male polyphenic signature in bulb mite populations". figshare. Dataset.
https://doi.org/10.6084/m9.figshare.22012589.v1

Del Giudice M, Buck CL, Chaby LE, Gormally BM, Taff CC, Thawley CJ, Vitousek MN, Wada H. 2018.
What Is Stress? A Systems Perspective. Integr. Comp. Biol. 58: 1019-1032.

Díaz A, Okabe K, Eckenrode CJ, Villani MG, Oconnor BM. 2000. Biology, ecology, and management of
the bulb mites of the genus *Rhizoglyphus* (Acari: Acaridae). Exp. Appl. Acarol. 24: 85–113.

Eberhard WG. 1982. Beetle horn dimorphism: making the best of a bad lot. Am. Nat. 119: 420–426.

Edeline E, Loeuille N. 2021. Size-dependent eco-evolutionary feedbacks in harvested systems. Oikos
130: 1636-1649.

Ellis BJ, Del Giudice M. 2019. Developmental Adaptation to Stress: An Evolutionary Perspective.
Annu. Rev. Psychol. 70: 111–139.

Ernande B, Dieckmann U, Heino M. 2004. Adaptive changes in harvested populations: plasticity and
evolution of age and size at maturation. Proc Biol Sci. 271: 415-423.

Fisher MO, Nager RG, Monaghan P. 2006. Compensatory Growth Impairs Adult Cognitive
Performance. PLoS Biology 4: e251.

Forsman, A. 2015. Rethinking phenotypic plasticity and its consequences for individuals, populations
and species. Heredity 115, 276–284

Gardner JL, Peters A, Kearney MR, Joseph L, Heinsohn R. 2011. Declining body size: A third universal
response to warming? Trends Ecol. Evol. 26: 285–291.

Gilbert JP, Han, Z-Y, Wieczynski DJ, Votzke S. 2022. Feedbacks between size and density determine
rapid eco-phenotypic dynamics. Funct. Ecol. 36: 1668-1680.

Griffiths PE, Gray RD. 1994. Developmental Systems and Evolutionary Explanation. J. Phil. 91: 277-
304.

Hazel, W.N., Smock, R., Johnson, M.D. 1990. A polygenic model for the evolution and maintenance
of conditional strategies. Proc. R Soc. Lond. B Biol. Sci. 242, 181–187.

Hazel, W.N., Smock, R., Lively, C.M. 2004. The ecological genetics of conditional strategies. Am. Nat.
163, 888–900.

Hill GE. 2011. Condition-dependent traits as signals of the functionality of vital cellular processes.
Ecol. Lett. 14: 625–634.

Hilborn R. Walters C. 1992. Quantitative fisheries stock assessment: choice, dynamics and
uncertainty. Springer.

Huryn A, Benke A. 2007. Relationship between biomass turnover and body size for stream
communities. In A. Hildrew, D. Raffaelli, & R. Edmonds-Brown (Eds.), Body size: The structure
and function of aquatic ecosystems (pp. 55–76). Cambridge University Press.

Knell RJ, Martínez-Ruiz C. 2017. Selective harvesting focused on sexual signal traits can lead to
extinction under directional environmental change. Proc. R. Soc. B 284: 20171788.

Lindström J. 1999 Early development and fitness in birds and mammals. Trends Ecol. Evol. 14: 343–
348.

Lürig MD, Matthews B. 2021. Dietary-based developmental plasticity affects juvenile survival in an
aquatic detritivore. Proc. R. Soc. B 288: 20203136.

Moczek, AP. 2003. The behavioral ecology of threshold evolution in a polyphenic beetle. Behav. Ecol.
14: 841–854.

Mojica JP, Kelly JK. 2010. Viability selection prior to trait expression is an essential component of
natural selection. Proc. R. Soc. B 277: 2945–2950.

Nijhout HF, McKenna KZ. 2018. The distinct roles of insulin signaling in polyphenic development.
Curr. Opinion Insect Sci. 25:58–64.

Plaistow SJ, Tsuchida K, Tsubaki Y, Setsuda K. 2005 The effect of a seasonal time constraint on
development time, body size, condition, and morph determination in the horned beetle
Allomyrina dichotoma L. (Coleoptera: Scarabaeidae). Ecol. Entomol. 30: 692–699.
Radwan J. 1995. Male morph determination in two species of acarid mites. Heredity 74: 669–673
Radwan J, Bogacz I. 2000. Comparison of life-history traits of the two male morphs of the bulb mite,
Rhizoglyphus robini. Exp. Appl. Acarol. 24: 115–121.
Radwan JW, Czyz M, Konior M, Kołodziejczyk M. 2000. Aggressiveness in two male morphs of the
bulb mite *Rhizoglyphus robini*. Ethology 106: 53-62.
Rhebergen FT. 2022. The ecology of adaptive condition-dependent polyphenism. PhD thesis,
University of Amsterdam, The Netherlands.
Rhebergen FT, Stewart KA, Smallegange IM. 2022. Nutrient-dependent allometric plasticity in a
male-diphenic mite. Ecol. Evol. e9145.
Roff, D.A. 1996. The evolution of threshold traits in animals. Q. Rev. Biol. 71, 3–35
Sasson DA, Munoz PR, Gezan SA, Miller CW. 2016 Resource quality affects weapon and testis size
and the ability of these traits to respond to selection in the leaf-footed cactus bug, *Narnia*
femorata. Ecol. Evol. 6: 2098–2108.
Schaefer MB. 1954. Some aspects of the dynamics of populations important to the management of
commercial marine fisheries. Inter-Am. Trop. Tuna Comm. Bull. 1: 23–56.
Sheridan JA, Bickford D. 2011. Shrinking body size as an ecological response to climate change.
Nature Climate Change 1: 401–406.
Shuster SM, Wade MJ. 2003. Mating Systems and Strategies. Princeton University Press, Princeton,
NJ.
Smallegange IM. 2011. Complex environmental effects on the expression of alternative reproductive
phenotypes in the bulb mite. Evol. Ecol. 25: 857–873.
Smallegange IM. 2022. Integrating developmental plasticity into eco-evolutionary population
dynamics. Trends Ecol. Evol. 37:129-137
Smallegange IM, Coulson T. 2011. The stochastic demography of two coexisting male morphs.
Ecology 92: 755–764.
Smallegange IM, Coulson T. 2013. Towards a general, population-level understanding of eco-
evolutionary change. Trends Ecol. Evol. 28: 143-148.
Smallegange IM, Deere JA. 2014. Eco-evolutionary interactions as a consequence of selection on a
secondary sexual trait. Adv. Ecol. Res. 50: 145–169.
Smallegange IM, Ens HM. 2018. Trait-based predictions and responses from laboratory mite
populations to harvesting in stochastic environments. Journal of Animal Ecology 87: 893-905
Smallegange IM, Rhebergen FT, Stewart KA. 2019. Cross-level considerations for explaining selection
pressures and the maintenance of genetic variation in condition-dependent male morphs. Curr.
Opinion Insect Sci. 36: 66–73.
Stewart KA, Draaijer R, Kolasa MR, Smallegange IM. 2019. The role of genetic diversity in the
evolution and maintenance of environmentally-cued, male alternative reproductive tactics.
BMC Evol Biol 19: 58.
Tomkins JL, Hazel WN, Penrose MA, Radwan JW, Lebas NR. 2011. Habitat complexity drives
experimental evolution of a conditionally expressed secondary sexual trait. Curr. Biol. 21: 569–
573.
Travis J, Leips J, Rodd FH. 2013. Evolution in population parameters: density-dependent selection or
density-dependent fitness? Am. Nat. 181: S9-S20.

Van den Beuken TPG, Duinmeijer CC, Smallegange IM. 2019. Costs of weaponry: unarmed males sire
more offspring than armed males in a male-dimorphic mite. J. Evol. Biol. 32: 153-162.

Van Gils JA, Lisovski S, Lok T, Meissner W, Ozarowska A, de Fouw J, … Klaassen M. (2016). Body
shrinkage due to Arctic warming reduces red knot fitness in tropical wintering range. Science
352: 819–821.

Verhulst PF. 1838. Notice sur la loi que la population suit dans son accroissement. – Corresp.
Mathématique Phys. 10: 113–121.

Vickers MH. 2014 Early life nutrition, epigenetics and programming of later life disease. Nutrients 6:
2165–2178.

Weir LK, Kindsvarter HK, Young KA, Reynolds JD. 2016. Sneaker males affect fighter male body size
and sexual size dimorphism in salmon. Am. Nat. 188: 264-271.

Wells JCK. 2011. The Thrifty Phenotype: An Adaptation in Growth or Metabolism? Am. J. Human Biol.
75: 65–75.

West-Eberhard MJ. 2003. Developmental Plasticity and Evolution. Oxford University Press, Oxford.