A selfish genetic element and its suppressor causes abnormalities to testes in a fly

Authors

Sophie Lyth[1], Andri Manser[2], Tom A. R. Price[1], Gregory D. D. Hurst[1], Rudi L. Verspoor[1].

[1]Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Biosciences Building, Liverpool, UK

[2]Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland

Abstract

Selfish genetic elements (SGEs), specifically X-chromosome meiotic drive (XCMD), create huge conflicts within a host's genome and can have profound effects on fertility. Suppressors are a common evolutionary response to XCMD to negate its costs. However, whether suppressors themselves can cause negative non-target effects remains understudied. Here, we examine whether the intragenomic conflicts created by XCMD and its suppressor affect gonad morphology in *Drosophila subobscura.* We found significant differences in testes, seminal vesicle, and accessory gland size depending on whether a male carried a non-driving X chromosome, an XCMD, and if the XCMD was suppressed. We also found the first evidence of abnormal testes development that is specifically associated with a suppressor of XCMD. Unlike other studies, our evidence suggests that XCMD in *D. subobscura* creates major abnormalities to male gonads. These abnormalities are most frequent if both XCMD and its suppressor are both present. While costs of suppression have importance in theoretical models, they have largely been ignored in empirical XCMD systems. Overall, this study highlights that genetic conflict, created by SGEs and their suppressors, is a potent evolutionary force that can have major impacts on gonad development and gametogenesis.

Introduction

Gamete production, and with this testes and ovary development, are shaped by both natural and sexual selection. For instance, egg size is strongly affected by natural selection (1,2) whereas testis size and sperm number is heavily influenced by sexual selection (3,4), with trade-offs between investment in testes and somatic function. Thus, testes are classic examples of organs that have to trade-off between sexual and natural selection (5). However, gonads are also the site of an additional set of trade-offs; the need to defend against selfish genetic elements (SGEs) (6–8).

Challenging the laws of 'fair' Mendelian inheritance, SGEs can bias their own transmission into the next generation without increasing the fitness of the carrier and often at the cost of the rest of the genome (9). For example, B chromosomes are additional to an organism's normal chromosome complement, that carry superfluous DNA (10,11). They accumulate in the ova rather than polar bodies, allowing them to selfishly accumulate in the germline (10). As a result, they are widely reported in nature and can cause phenotypic differences and reduce fertility (10,12,13). Co-evolutionary responses to intragenomic conflicts created by SGEs in gonads are likely to shape their evolution, alongside natural and sexual selection. Thus, the role of SGEs on gonad evolution should not be overlooked, and could be significant and universal because SGEs are both diverse and near ubiquitous in nature (9).

Conflicts created by SGEs can have enormous consequences on the host's fertility. For instance, P element dysgenesis is a property of transposable elements active in *Drosophila simulans* germlines (14). Strong evidence for the impact of SGEs in evolution comes from observations when SGEs enter a naïve host that it has not coevolved with. The P element in its native host is not associated with any gross defects (15). Yet when this element is horizontally transmitted into naïve hosts, it becomes associated with a number of abnormalities, including sterility and dysgenic ovaries (16). Another example is X chromosome meiotic drive (XCMD), where male carriers have their Y-sperm killed or damaged thus ensuring the transmission of only the selfish X chromosome to the

next generation (17). This reduces a male's fertility and sperm competitive ability (18–20). Thus, a

range of SGEs can negatively impact their hosts and are capable of damaging the gonads and fertility

of both sexes. When faced with a multitude of conflicts caused by SGEs, how do organisms and in

particular their gonads, respond to deal with these challenges?

Here we investigate how the development of the testes responds when faced with XCMD

and a suppressor of this system. XCMD is a type of SGE that creates huge conflicts within the testes

(21). XCMD manipulates the process of meiosis in males, where selfish X chromosomes kill Y-bearing

sperm during spermatogenesis, resulting in the overproduction of daughters (17). The transmission

advantage gained by XCMD can allow it to rapidly spread through populations, distorting sex ratios

and even leading to population extinction owing to a lack of males (22–24). Thus, testes in XCMD

carrying males are a good model to investigate how SGEs impact the evolution of gonads.

Additionally, testes are particularly vulnerable to SGEs due to the asymmetry of gamete production

between males and females. Eggs are large and very costly for a female to produce, so anything that

damages eggs will be extremely costly and very quickly selected against. On the other hand, because

male gametes are generally massively over-produced, damage to sperm may be more readily be

tolerated. In cases where males only produce a reduced amount of sperm, they can still be

successful, making sperm and testes a good target for the activity of SGEs.

In XCMD, male carriers lose up to half of all the sperm that they produce (25). Despite the

lack of sperm produced, drive carrying males do not often show fertility costs in single matings when

compared to wildtype males (18,26–29). Indeed, in stalk eyed flies, the presence of sperm-killing

XCMD appears to have driven the evolution of a compensatory increase in the testes size of XCMD

males (30–32) . This allows them to produce a similar amount of sperm to non-XCMD males. These

studies, that look at how the testes respond to drive, raises the question as to whether drive

systems in general promote changes in the testes.

78   Meiotic drive is deleterious to male function, and selection commonly favours unlinked

suppressor factors that restore Mendelian segregation (21,33). Suppressors, either Y-linked or

autosomal, must have the ability to prevent some killing of Y-bearing sperm when a driving X

chromosome is present (34). Once a suppressor has evolved, and as long as it has no associated

costs, it is expected to rapidly spread to fixation (35,36). Despite this, the impact of suppressors of

XCMD on the testes themselves has been rarely considered in empirical studies, despite being

recognised as potentially important in theoretical studies (37). Ongoing co-evolution between drive

and its suppressor could have different outcomes depending on how a suppressor works, whether it

has associated costs, and if it has non-target effects. Thus, the question remains: does the

emergence of suppressors negate any testes phenotype (and associated selection) produced by

SGEs? Alternatively, the need for testes to evolve could be enhanced if they have to deal with both

XCMD and suppression, which are locked in a spiralling co-evolutionary arms race (37–40).

90   In this paper, we use the XCMD system in *Drosophila subobscura* to look for any effects on

testes morphology. *D. subobscura* harbours an XCMD system called 'SR' (41). The strength of drive is

strong with SR carrying males producing broods that are 85-100% female (41,42). Given this strong

drive, a simple expectation is that SR should rapidly spread to fixation everywhere. However, it does

not. Instead, in North Africa, SR has remained at a stable frequency of approximately 20% for at least

50 years (41–43). Additionally, *D. subobscura* are reported to be strictly monandrous (28,44) so

effects of sperm competition are irrelevant in this species (18,45). We recently discovered

suppressors of the SR system (43) that are capable of fully restoring a normal 50:50 sex ratio to SR

carrying males. Despite the prediction that suppressors against drive should rapidly fix within

populations, this suppressor remains rare at a frequency of 1-5%. The suppressor is also found in

multiple locations across North Africa at these low frequencies. The lack of spread of this suppressor

within populations may indicate that it is associated with its own costs and creates its own burden

upon its host. While we refer to "the suppressor", we have not fully characterised it. Initial crosses of

XCMD into suppressor genotypes results in partial suppression, and several generations of

introgression are required to reach complete drive suppression (Supplementary figure 1). Thus, we

believe "the suppressor" involves multiple loci across at least two chromosomes, and the Y is not

involved. However, in the interests of brevity, we will refer to it as "the suppressor" here, and aim to

dissect the complexity in future work.

We determine whether testes morphology is affected when males carry the selfish SR X

chromosome (henceforth referred to as SR males) in comparison to standard (ST) males that do not

carry SR. We also examine if SR males that also carry the suppressor genes (^{Supp}SR males) can restore

intact testes or whether the suppressor itself can have its own influence on testes morphology. We

used independent SR from multiple locations in Morocco to investigate whether drive in populations

have evolved and responded in the same way when it is paired with suppressors that originated

from populations in Tunisia.

Materials and Methods

Origin and maintenance of fly stocks

Isolines of *Drosophila subobscura* were originally collected from Tabarka, Tunisia (36.95°N 8.74°E) in 2013 (46), Ras el Oud, Morocco (34.15°N 4.00°W), Demnat, Morocco (31.73°N 7.01°W) and Ouzoud, Morocco (32.02°N 6.72°W). Moroccan collections took place in 2019 and all flies were caught using banana, yeast and beer baited traps (47). All isolines were kept in standard *Drosophila* vials on an ASG double yeast medium (10g agar, 85g sucrose, 40g yeast extract, 60g maize, 1000ml H_2O, 25ml 10% nipagin) at 18°C on a 12:12 light/dark cycle. Experimental flies were generated by collecting unmated flies from isolines from each population. All eclosed flies were removed from populations 24 hours prior to ensure flies had not mated. Fly collection was done using CO_2 anaesthesia to separate flies by sex. Flies were stored in new vials on a sugar yeast medium (100g sugar, 100g yeast extract, 20g agar, 1000ml H_2O, 25ml 10% nipagin, 5ml propionic acid) with no more than 10 flies per vial.

Isolation and introgression of SR X chromosomes

We isolated an SR chromosome from a wild caught fly from each of Ras el Oud, Morocco, Demnat, Morocco and Ouzoud, Morocco, where SR occurs at a frequency of approximately 20% (43). SR males were initially identified though a phenotypic scan by counting their offspring sex ratio. Males that produced more than 85% females were deemed SR males and were used to isolate their driving X chromosomes. In order to produce the SR males (that carry an SR chromosome), suppressed SR males (^{Supp}SR), that carry a suppressed version of the SR chromosome, and standard (ST) males (that do not carry the SR chromosome) that were used in this investigation, we introgressed both vulnerable and suppressing ST isolines onto the isolated SR chromosomes. This introgression process meant we were able to produce flies that only differ genetically in carrying drive (males either carried an SR X chromosome or an ST X chromosome) and either carried the suppression genes or genes susceptible to drive. Preliminary experiments have shown that the

Moroccan isolines show no suppression against SR (43), thus these isolines were used to produce the

SR males that produce 100% daughters. Additionally, the ST isoline from Tunisia contains genes that

are capable of suppressing SR. This Tunisian ST isoline was therefore used to produce the ^{Supp}SR

males that produce a normal 50:50 sex ratio despite the male carrying SR (Supplementary figure 1).

We introgressed the SR X chromosomes onto either the vulnerable Moroccan ST isolines or the

suppressing Tunisian ST isoline for 14 to 17 generations (Supplementary figure 2). The introgression

process was done in two crosses per generation. Cross 1: females homozygous for the isolated SR X

chromosome ('SRSR' females) were crossed to a standard ('ST') male from the target isoline. This

cross produces heterozygous females ('SRST' females) that are discarded, and 'SR' males that carry

the SR X chromosome and the target isoline autosomes. Cross 2: simultaneously, SRSR females were

crossed to SR males containing the isolated SR X chromosome. This cross produces only SRSR

females. SR males containing the target autosomes generated from cross 1 are then crossed to the

SRSR females generated from cross 2 to initiate the subsequent generation. ST males are also re-

crossed to the SRSR females from cross 2 to reset cross 1 for the next generation. Over multiple

generations the target autosomes will be introgressed onto the SR X chromosomes (Supplementary

figure 3).

Dissection of male testes

We examined the size of male testes and seminal vesicles (SV's) of SR males, ^{Supp}SR males and ST

males. To look for changes to male testes that may develop over time as they mature or

alternatively be present when just sexually mature and disappear with age, we dissected males

when they were either just sexually mature (at 3-days-old) or when they had reached full sexual

maturity (at 7-days-old) (48). When at either 3- or 7-days-old, males were anesthetised using FlyNap

(Carolina Biological Supply Co., Burlington, USA) and the testes were dissected out of the males in

300µl of phosphate buffered saline (PBS). Once dissected, the testes were transferred onto a

microscope slide containing 80µl of PBS. A photograph of the testes was taken under constant light

conditions with a microscope mounted Nikon D5100 camera at x40 magnification. To control for

body size effects, the left wing of each male was removed using forceps under a microscope

(alternatively, the right wing was taken if the left was damaged) and mounted onto a glass

microscope slide. A photograph of each wing was taken using the microscope mounted camera.

Measuring abnormalities to testes

We first examined photographs for evidence of abnormalities in the testes. Only abnormalities

in the testes themselves was considered. Abnormality was scored as a binary variable by the

presence of physical holes in the orange sheath that coats the testes (Figure 1a). An intact orange

sheath that showed thinning was not classed as abnormal. To avoid bias and ensure consistency, this

process was double blinded and each photo was independently evaluated by three people. Here, the

proportion of males that showed abnormal testes differed by observer. However, when repeating

the analysis just using the data of each observer, we found the same effect that [Supp]SR males

experienced more abnormalities (Analysis 1: GLM: $F_{8,210}$ = 11.227, p <0.001; Analysis 2: GLM: $F_{8,210}$ =

9.821, p <0.001; Analysis 3: GLM: $F_{8,210}$ = 8.818, p <0.001) (Supplementary Figure 4). Thus, a

composite of the three analyses was used to determine a final testis abnormality score for each male

in which testes were assigned as abnormal if two or more evaluators scored them as abnormal.

Measuring testes size, SV size, accessory gland size and wing length

Testes and SV size was measured from images using ImageJ 1.48 (49). A subset of images with

visible accessory glands were also analysed in ImageJ to measure accessory gland size. The area of

each individual testis was measured (Figure 1b) and a mean of the two taken for each male. If only

one testis could be measured for size, it was this single measurement that represented a male's

mean testes size. The same procedure was performed for the SV's and accessory glands. Wing

lengths were also measured in ImageJ to control for body size (Figure 1b), using the method in

Gilchrist *et al.,* 2001 (50). The mean testes, SV and accessory gland size for each male was divided by

wing length to generate testes, SV or accessory gland size relative to wing length. These relative

measurements were used in all statistical analyses.

Figure 1: Analysis and measurement of male reproductive organs and wing size. a) Example images of testes (T), seminal vesicles (SV), accessory glands (AG) and visible abnormalities to testes (D). ai) Image of intact testes with no abnormalities to the orange sheath. aii) Abnormal testes defined by the presence of physical holes in the orange sheath. bi) Measurement of testes (yellow), seminal vesicles (green) and accessory gland (blue) area using ImageJ. bii) Measurement of wing length in ImageJ as defined in Gilchrist *et al.,* 2001 (50).

Statistical analysis

All statistical analysis was performed using the statistical software R, version 3.6.0 (51). A

binomial generalised linear model (GLM) was used to investigate differences in the frequency of

abnormalities to male testes. Testes size, SV size and accessory gland size were analysed by fitting a

GLM with a gaussian error distribution. All models considered the effects of the type of male (SR,

[Supp]SR, ST susceptible or ST suppressed males), and the age of the male when it was dissected (3- or

7- days old). ST males were split into either ST susceptible or ST suppressed to allow comparisons of

the suppressor in the absence of the selfish X chromosome. For testes abnormalities, testes size, and

SV size analysis, the original location of origin of the SR X chromosome was included in the model as

this provided a better model fit. In these analyses, the effect of SR X chromosome location was

significant but idiosyncratic and does not affect the main effects of drive (for individual line effects,

see supplementary figures 7-10). For all GLMs, a fully saturated model was produced and was

systematically reduced to the minimum adequate model through a step-wise simplification process

using Akaike information criterion (AIC). Difference in responses to all GLMs under different

treatments were assessed by analysis of variance (ANOVA) followed by a TukeyHSD comparison. For

simplicity, all graphs presented align SR next to ST susceptible and [Supp]SR next to ST suppressed as

they share genetic backgrounds to make interpretation clearer of what happens when SR is added to

each genetic background.

Results

Wing size

Wing size (a proxy for body size), was analysed using a GLM. The model considered the type

of male and the age of the male and their interaction. The interaction was not significant and was

dropped from the model. Wing size was affected by age (GLM: $F_{1,218}$ = 7.518, p = 0.007), where 7-

day-old males were smaller than 3-day-old males (TukeyHSD, p = 0.007). The type of male also had a

significant effect on wing size (Supplementary figure 6: GLM: $F_{3,218}$ =32.762, p < 0.001). ST

suppressed males were smaller than [Supp]SR (TukeyHSD, p = 0.002), SR (TukeyHSD, p <0.001) and ST

susceptible males (TukeyHSD, p < 0.001). [Supp]SR males were smaller than SR and ST susceptible males

(TukeyHSD, p < 0.001 and p = 0.002, respectively). There was no difference in the wing size of SR and

ST susceptible males (TukeyHSD, p = 0.120).

Testes size

Mean testes size relative to wing length was analysed using a GLM. The model considered

the type of male and the age of the male. A significant age by type of male interaction was found

(Figure 2a: GLM: $F_{3,213}$ =3.663, p = 0.0132). Here, ST males (both suppressed and susceptible) have

significantly reduced testes size as they age from 3-days-old to 7-days-old (TukeyHSD, p < 0.001 for

both ST types). This reduction in testes size was not seen in SR and [Supp]SR males as they aged

(TukeyHSD, p = 0.975, TukeyHSD, p = 0.780, respectively). There was an overall effect of age on

testes size (GLM: $F_{1,213}$ = 15.705, p < 0.001), where 7-day old-males had larger testes (TukeyHSD, p

<0.001).

Seminal vesicle size

Mean SV size relative to wing length was analysed using a GLM. The model considered the

type of male and the age of the male and their interaction. The interaction was not significant, so

was dropped from the model. Overall, SV's were bigger in 7-day-old males than 3-day-old males

(Figure 2b: GLM: $F_{1,213}$ = 24.095, p < 0.001). The type of the male also had a significant effect on

mean SV size (Figure 3b: GLM: $F_{3, 213}$ = 7.384, $p < 0.001$). Here, ST suppressed males have larger SV's

than SR males (TukeyHSD, $p < 0.01$), ST susceptible males (TukeyHSD, $p = 0.005$) and ^{Supp}SR males

(TukeyHSD, $p = 0.026$). However, there was no difference in SV size between ^{Supp}SR and SR and ST

susceptible males.

Figure 2: The mean size of a male's two testes (a) or seminal vesicles (b) of *Drosophila subobscura* males that were either SR, ST susceptible, ST suppressed or [Supp]SR. All measurements were relative to individual body size. The boxplots display the upper and lower quartiles, the median and the range. Individual points represent each measurement obtained.

Abnormal testes

The presence of visible abnormalities to testes was analysed using a binomial GLM. The

model considered the type of male and the age of the male and their interaction. Neither age nor

the age by male type interaction had an effect on visible abnormalities to a male's testes (GLM: $F_{1, 199}$

$= 0.064$, $p = 0.801$, $F_{10,199} = 1.609$, $p = 0.106$, respectively) so they were dropped from the model.

Male type had a significant effect on the likelihood of male having abnormal testes (Figure 3: GLM:

$F_{3, 210} = 9.580$, $p < 0.001$). While most categories differed from one another, the most noticeable

differences were that ^{Supp}SR males had a significantly higher proportion of abnormal testes

compared to ST suppressed males (where by our measure, 49% of ^{Supp}SR males had visibly abnormal

testes in comparison to 1% ST suppressed males) (Figure 3: TukeyHSD, $p < 0.001$). On the other

hand, SR males did not differ significantly in the frequency of abnormal testes in comparison to ST

susceptible (18% of SR males had damage to their testes in comparison to 22% in ST susceptible

males, TukeyHSD, $p = 0.594$). ^{Supp}SR also had a higher amount of damage than SR males (TukeyHSD, p

$= 0.0015$). An example of both normal and abnormal testes can be seen in Figure 1a.

We also dissected the testes from SR, ^{Supp}SR and ST susceptible males when 7-days-old for

preliminary and exploratory analysis of the internal structure of each male type's testes (see

supplementary figure 11 for details). SR and ^{Supp}SR testes showed disorganisation, with gaps

between cells, while ST susceptible males showed normal tightly packed sperm development

(supplementary figure 11). While ^{Supp}SR males showed a higher level of organisation than SR males,

the internal structure of male testes did not resemble that of ST susceptible males.

Figure 3: The mean proportion of males that have visible abnormalities to their testes that were
either SR males, ST susceptible males, ST suppressed males or ^{Supp}SR males, with 95% confidence
intervals. The figure also reports statistical comparisons where NS is non-significant, * is $p < 0.05$, **
is $p < 0.01$ and *** is $p < 0.001$.

Accessory gland size

Accessory glands were measured in any images where they were clearly visible. Mean

accessory gland size relative to wing length was analysed using a GLM. The model considered the

type of male and the age of the male. Models analysing accessory gland size did not consider the

individual location the SR X chromosome originated from due to the smaller sample size and having

little power to detect differences. A significant age by type of male interaction was found (Figure 4:

GLM: $F_{3, 143}$ = 3.689, p = 0.0135). Here, [Supp]SR males' accessory glands increase in size as they age

(TukeyHSD, p = 0.004). This change in accessory gland size was not seen in SR males (TukeyHSD, p =

0.999), ST susceptible males (TukeyHSD, p = 0.999) or ST suppressed males (TukeyHSD, p = 0.933).

Male type also had an effect on accessory gland size (GLM: $F_{3, 143}$ = 3.689, p = 0.0135). Overall, [Supp]SR

males had larger accessory glands than SR and ST susceptible and ST suppressed males (TukeyHSD, p

= 0.046, TukeyHSD, p < 0.001, TukeyHSD, p = 0.037, respectively). SR males also had larger accessory

glands than ST susceptible males (TukeyHSD, p = 0.003). Finally, the age of the male had an effect on

accessory gland size (GLM: $F_{1, 143}$ = 5.713, p = 0.0181). Overall, 7-day-old males had larger accessory

glands than 3-day-old males (TukeyHSD, p = 0.019).

Figure 4: The mean size of the two accessory glands of male *Drosophila subobscura* that were either
SR, ST susceptible, ST suppressed or ^Supp^SR. All measurements were relative to individual body size.
The boxplots display the upper and lower quartiles, the median and the range. Individual points
represent each measurement obtained.

Discussion

XCMD and their suppressors generate potent genetic conflict that may also be a force influencing gonad development, morphology and evolution (31,32). In this study, we examined whether the intragenomic conflicts created by SR and its suppressor affects gonad morphology in *D. subobscura.* We found significant differences in testes, seminal vesicle, and accessory gland size depending on whether a male carried XCMD or not, and also if XCMD was suppressed. We also found the first evidence of extreme whole-organ abnormal development of testes associated with carrying a suppressor of XCMD. While males with SR from independent origins had differences in the morphological traits observed, this variation was idiosyncratic and overall the main effect of SR and its suppressor were the same, suggesting these effects are robust across populations. In conclusion, we found strong evidence that XCMD and its suppressor can have a major impact on gonad development and morphology of its carriers.

Testes and Seminal Vesicle size

We found evidence that the testes of ST (both suppressed and susceptible) males, but not those of SR and ^{Supp}SR males, reduce in size as they age. *Drosophila subobscura* are strictly monandrous (28,44), and males may not benefit from high investment in continually producing large quantities of sperm (52–54). In particular, if *D. subobscura* can partition their ejaculates, like in numerous species across a wide array of taxa (55,56), they may only transfer to a female the sperm she will need. This results in reduced pressure on the testes to continually produce sperm (and the costs of spermatogenesis this entails). In contrast to the reduction in testes size observed in ST males, it appears that SR and ^{Supp}SR testes remain the same size as males age. We propose three mutually non-exclusive reasons for this: (1) compensation, (2) damage or (3) a delay in sperm production.

In stalk-eyed flies, XCMD males have enlarged testes to compensate drive induced sperm mortality, and males produce an equivalent number of sperm to normal males (31). This

compensation of male testes appears to occur from an early development stage (32). Compensation

may also be the effect that we have observed in *D. subobscura*. Indeed, a previous study has

suggested that the loss of Y-bearing sperm in SR males is compensated through cyst doubling in the

testes (57,58). Unlike in stalk-eyed flies, we do not observe this increase in testes size at all ages. At

3-days-old, all male types have equivalent sized testes. If this was compensation, would SR male

carriers not always have enlarged testes? In addition, in a species with no sperm competition

(28,44), loss of sperm number may not need to be compensated.

Contrary to the compensation hypothesis, damage may instead be inflicted upon the testes

resulting in their enlarged size. The internal testes structure of SR and [Supp]SR males showed

disorganisation, with gaps between cells, while ST testes showed normal tightly packed sperm

development (supplementary figure 11). This may indicate that XCMD carriers instead experience

internal testes damage and/or spermatogenic failure, the severity of which may accumulate over

time. Indeed, the gapping between cells and their general disorganisation may explain why SR and

[Supp]SR male have enlarged testes at this age. However, a full investigation of the internal morphology

of testes and associated sperm development is required to explore this further. Currently, little is

known about the interaction of XCMD and genes generally involved in gonad development (59,60).

Finally, if testes are largest simply when the process of spermatogenesis is at its peak, then

sperm production, and the reaching of a full sperm complement, may simply be delayed in SR and

[Supp]SR males. As a result, testes would remain larger for a longer period of time, while they continue

full sperm production to an older age. A detailed temporal breakdown of sperm production and

storage would be required to confirm this hypothesis.

We found no strong effects of carrying XCMD on SV size, as these were equivalent in all male

types, other than ST suppressed. The reasons for this are unclear and do not provide strong support

for any of our above theories. SVs are sperm storage organs in *Drosophila* males, where mature

sperm that develop in the testes move into the SV for storage until mating (61,62). Thus, in unmated

males, SV size is generally related to the amount of sperm stored (63). The equivalent SV size may

indicate that these male types are all producing and storing a similar amount of sperm, providing

some additional support for the compensation theory. Conversely, all the sperm stored in the SVs of

SR and ^{Supp}SR males may not be viable. It is possible that these males are bundling dead or damaged

sperm that is incapable of fertilisation or zygote formation. The disruption to the internal structure

of these testes suggests that spermatogenesis is likely to be affected in these males (supplementary

figure 11). A full examination of the viability of sperm stored within the SVs would be required to

confirm this.

Whole testes abnormalities

Remarkably, we found evidence that, when the suppressor is paired with a driving SR X

chromosome, it causes gross visible abnormalities to testes. Overall, these abnormalities are less in

XCMD males that lack the suppressor which is comparable to that of ST susceptible males. Once

evolved, suppressors against drive are usually expected to remove or minimise the negative impacts

of drive (33,64,65). However, we demonstrate for the first time that suppression, while mitigating

some negative effects of XCMD, can also have its own effects on testes morphology, that likely

imposes costs. In addition, given the scope of the morphological differences we see, suppression

genes may influence general testes development that, when expressed and paired with SR, results in

their abnormal development and potential spermatogenesis failures. In addition, these

abnormalities are not restricted to the external testes. Our preliminary SEM analysis of the internal

structure of male's testes shows that both SR and ^{Supp}SR males have a high level of disorganisation

and gapping between cells, while ST shows normal tightly packed sperm development. Thus, SR

appears to be having its own influence of testes morphology, that becomes enhanced when the

suppressor is also present. This is in support of our theory that SR and its suppressor are inflicting

damage or developmental changes upon a male's testes. To what effect, if any, these abnormalities

then has on a male's fertility remains an open question.

It is critical these abnormalities, and any associated costs is understood as it could affect the

balance between drivers and suppressor that may provide insight into the long-term stability of

drive that we see in many natural systems (43,66–68). Additionally, if suppressors can spread

despite deleterious effects, they have the potential to fix negative mutations within a population,

which is rarely considered in empirical studies. There are many cases of cryptic (or muted) drive

systems that are fixed within populations (64,65,69). Here, drive systems are only revealed through

outcrosses with other populations or species that do not have the suppressor. These factors are

especially important to consider in relation to artificial drive systems (70). What would happen to

artificial drive if a suppressor unexpectedly evolved and fixed deleterious mutations within the

target population? The evidence we have found that major abnormalities are associated with

suppression against XCMD suggests that not only can SGEs carry huge costs, but in addition

suppressors and their interactions with SGEs must be accounted for.

Accessory gland size

The extent of changes to male gonads as a result of XCMD seem not to just be limited to the

testes. We observed changes to the whole of the male gonads, including the accessory glands.

Accessory glands are organs that are responsible for the production of seminal fluid that contains

accessory gland proteins (71). In _Drosophila_, accessory gland proteins influence female behaviour

and a males sperm competitive ability (72,73). In some systems, accessory gland size is also

correlated with male mating rate and success (74,75). In this study, we found that [Supp]SR accessory

glands are the largest, followed by SR which were larger than ST (both suppressed and susceptible)

males. Carrying SR may be associated with compensatory increases in investment in accessory gland

proteins to ensure their mating success rather than transferring an equivalent amount of sperm to

ST males. Additionally, the highest investment in accessory glands was from [Supp]SR. When SR males

are suppressed they also endured the highest frequency of testes abnormalities, which may indicate

males are investing more in gaining a mate. Despite this, it remains unclear as to what the role of

accessory glands is in a monogamous species where the importance of them becomes largely

unknown.

Conclusion

Here, we demonstrate that the presence of XCMD and its suppressor can cause significant

morphological changes to male gonads. This is the second system, that we are aware of, where

XCMD strongly influences male gonad development (31,32). Unlike the XCMD system in stalk-eyed

flies, where morphological changes were deemed a compensation mechanism against drive, our

evidence suggests that XCMD in *D. subobscura* inflicts gross abnormalities on male gonads that

becomes even more severe in the presence of its suppressor. This highlights that SGEs are a strong

evolutionary force that can broadly impact gonad development and gametogenesis. Additionally,

considering drive alone may only reveal half of the story. Suppressors can also have their own severe

effects on their carriers. Overall, the fitness effects of either an SGE or its suppressors may be severe

for individuals that carry them.

Acknowledgements

The authors would like to thank Jola Tanianis for her help with the isolation of the SR chromosomes,

and Kieran Romain for measuring the accessory glands. The authors acknowledge the use of the

biomedical electron microscopy unit provided by Liverpool shared research facilities, Faculty of

Health and Life Sciences, University of Liverpool. We would also like to thank the three anonymous

referees for their helpful comments on the manuscript. This work was funded by the National

Environment Research Council ACCE DTP and the grant no. NE/S001050/1.

References

References

1. Fox CW. Natural selection on seed-beetle egg size in nature and the laboratory: Variation among
environments. Ecology. 2000;81(11):3029–35.

2. Bernardo J. The particular maternal effect of propagule size, especially egg size: Patterns,
models, quality of evidence and interpretations. Am Zool. 1996;36(2):216–36.

3. Parker BYGA. Sperm competition and its evolutionary consequences in the insects. Biol Rev.
1970;45(May):535–67.

4. Simmons LW, García-González F. Evolutionary reduction in testes size and competitive
fertilization success in response to the experimental removal of sexual selection in dung beetles.
Evolution. 2008;62(10):2580–91.

5. Pitnick S, Markow TA. Large-male advantages associated with costs of sperm production in
Drosophila hydei, a species with giant sperm. Proc Natl Acad Sci U S A. 1994;91(20):9277–81.

6. Senti KA, Brennecke J. The piRNA pathway: A fly's perspective on the guardian of the genome.
Trends Genet. 2010;26(12):499–509.

7. Malone CD, Hannon GJ. Small RNAs as Guardians of the Genome. Cell. 2009;136(4):656–68.

8. Zeh JA, Zeh DW. The evolution of polyandry I: Intragenomic conflict and genetic incompatibility.
Proc R Soc B Biol Sci. 1996;263(1377):1711–7.

9. Burt A, Trivers R. Genes in conflict: the biology of selfish genetic elements. Harvard University
Press. 2009.

10. Camacho JPM, Sharbel TF, Beukeboom LW. B-chromosome evolution. Philos Trans R Soc B Biol
Sci. 2000;355(1394):163–78.

11. Houben A, Banaei-Moghaddam AM, Klemme S, Timmis JN. Evolution and biology of
supernumerary B chromosomes. Cell Mol Life Sci. 2014;71(3):467–78.

12. Jones N, Houben A. B chromosomes in plants: Escapees from the A chromosome genome?
Trends Plant Sci. 2003;8(9):417–23.

13. Price TAR, Verspoor RL, Wedell N. Ancient gene drives: an evolutionary paradox. Proc Biol Sci.
2019;286(1917):20192267.

14. Hill T, Schlötterer C, Betancourt AJ. Hybrid Dysgenesis in *Drosophila simulans* Associated with a
Rapid Invasion of the P-Element. PLoS Genet. 2016;12(3):1–17.

15. Mizrokhi LJ, Mazo AM. Evidence for horizontal transmission of the mobile element jockey
between distant *Drosophila* species. Proc Natl Acad Sci U S A. 1990;87(23):9216–20.

16. Kidwell MG, Kidwell JF, Sved JA. Hybrid dysgenesis in *Drosophila melanogaster*: a syndrome of
aberrant traits including mutation, sterility and male recombination. Genetics. 1977;86(4):813–
33.

17. Jaenike J. Sex chromosome meiotic drive. Annu Rev Ecol Syst. 2001;32:25–49.

18. Keais GL, Lu S, Perlman SJ. Autosomal suppression and fitness costs of an old driving X
chromosome in *Drosophila testacea*. J Evol Biol. 2020;

19. Dyer KA, Hall DW. Fitness consequences of a non-recombining sex-ratio drive chromosome can
explain its prevalence in the wild. Proc Biol Sci. 2019;286(1917):20192529.

20. Price TAR, Bretman A, Avent TD, Snook RR, Hurst GDD, Wedell N. Sex ratio distorter reduces
sperm competitive ability in an insect. Evolution. 2008;62(7):1644–52.

21. Lindholm AK, Dyer KA, Firman RC, Fishman L, Forstmeier W, Holman L, et al. The Ecology and
Evolutionary Dynamics of Meiotic Drive. Trends Ecol Evol. 2016;31(4):315–26.

22. Hamilton WD. Extraordinary Sex Ratios. Science. 1967;156:477–88.

23. Price TAR, Hurst GDD, Wedell N. Polyandry Prevents Extinction. Curr Biol. 2010;20(5):471–5.

24. Holman L, Price TAR, Wedell N, Kokko H. Coevolutionary dynamics of polyandry and sex-linked
meiotic drive. Evolution. 2015;69(3):709–20.

25. Policansky D, Ellison J. 'Sex ratio' in *Drosophila pseudoobscura*: spermiogenic failure. Science.
1970;169(3948):888–9.

26. Beckenbach AT. Selection and the 'Sex-Ratio' Polymorphism in Natural Populations of
Drosophila pseudoobscura. Evolution. 1996;50(2):787–94.

27. Price TAR, Lewis Z, Smith DT, Hurst GDD, Wedell N. No evidence of mate discrimination against
males carrying a sex ratio distorter in *Drosophila pseudoobscura*. Behav Ecol Sociobiol.
2012;66(4):561–8.

28. Verspoor RL, Hurst GDD, Price TAR. The ability to gain matings, not sperm competition, reduces
the success of males carrying a selfish genetic element in a fly. Anim Behav. 2016;115:207–15.

29. Wu C i. Virility deficiency and the sex-ratio trait in *Drososphila pseudoobscura*. I. Sperm
displacement and sexual selection. Genetics. 1983;105:651–62.

30. Meade L, Dinneen D, Kad R, Lynch DM, Fowler K, Pomiankowski A. Ejaculate sperm number
compensation in stalk-eyed flies carrying a selfish meiotic drive element. Heredity.
2019;122(6):916–26.

31. Meade L, Finnegan S, Kad R, Fowler K, Pomiankowski A. Maintenance of fertility in the face of
meiotic drive. Am Nat. 2020;

32. Bradshaw SL, Meade L, Tarlton-Weatherall J, Pomiankowski A. Meiotic drive adaptive testes
enlargement during early development in the stalk-eyed fly. Biol Lett. 2022;18(11):20220352.

33. Price TAR, Windbichler N, Unckless RL, Sutter A, Runge JN, Ross PA, et al. Resistance to natural
and synthetic gene drive systems. J Evol Biol. 2020;(June):1–16.

34. Helleu Q, Gerard PR, Montchamp-Moreau C. Sex chromosome drive. Cold Spring Harb Perspect
Biol. 2015;7(2).

35. Bastide H, Gérard PR, Ogereau D, Cazemajor M, Montchamp-Moreau C. Local dynamics of a fast-
evolving sex-ratio system in *Drosophila simulans*. Mol Ecol. 2013;22(21):5352–67.

36. Bastide H, Cazemajor M, Ogereau D, Derome N, Hospital F, Montchamp-Moreau C. Rapid rise
and fall of selfish sex-ratio X chromosomes in *Drosophila simulans*: Spatiotemporal analysis of
phenotypic and molecular data. Mol Biol Evol. 2011;28(9):2461–70.

37. Hall DW. Meiotic drive and sex chromosome cycling. Evolution. 2004;58(5):925–31.

38. Vaz SC, Carvalho AB. Evolution of Autosomal Suppression of the Sex-Ratio Trait in *Drosophila*.
Genetics. 2004;166(1):265–77.

39. Courret C, Chang CH, Wei KHC, Montchamp-Moreau C, Larracuente AM. Meiotic drive
mechanisms: Lessons from *Drosophila*. Proc R Soc B Biol Sci. 2019;286(1913).

40. Searle JB, de Villena FPM. The evolutionary significance of meiotic drive. Heredity.
2022;129(1):44–7.

41. Jungen H. Abnormal sex ratio linked with inverted gene sequence in populations of *D.*
subobscura from Tunisia. 1967;42:109.

42. Hauschteck-Jungen E. Postmating reproductive isolation and modification of the 'sex ratio' trait
in *Drosophila subobscura* induced by the sex chromosome gene arrangement A2+3+5+7.
Genetica. 1990;83(1):31–44.

43. Verspoor RL, Smith JML, Mannion NLM, Hurst GDD, Price TAR. Strong hybrid male
incompatibilities impede the spread of a selfish chromosome between populations of a fly. Evol
Lett. 2018;2(3):169–79.

44. Fisher DN, Doff RJ, Price TAR. True polyandry and pseudopolyandry: Why does a monandrous fly
remate? BMC Evol Biol. 2013;13(1).

45. Price TAR, Bretman A, Gradilla AC, Reger J, Taylor ML, Giraldo-Perez P, et al. Does polyandry
control population sex ratio via regulation of a selfish gene? Proc R Soc B Biol Sci.
2014;281(1783).

46. Verspoor RL, Cuss M, Price TAR. Age-based mate choice in the monandrous fruit fly *Drosophila*
subobscura. Anim Behav. 2015;102:199–207.

47. Markow TA, O'Grady P. *Drosophila*: a guide to species identification and use. Elsevier. 2005;

48. Holman L, Freckleton RP, Snook RR. What use is an infertile sperm? A comparative study of
sperm-heteromorphic *Drosophila*. Evolution: International Journal or Organic Evolution; 2008. p.
374–85.

49. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. 2012;

50. Gilchrist GW, Huey RB, Serra L. Rapid evolution of wing size clines in *Drosophila subobscura*.
Genetica. 2001;112–113:273–86.

51. R Development Core Team. R Foundation for Statistical Computing. R: A language and
environment for statistical computing. R Found Stat Comput Vienna Austria URL HttpswwwR-
Proj. 2019;

52. Sirot LK, Buehner NA, Fiumera AC, Wolfner MF. Seminal fluid protein depletion and
replenishment in the fruit fly, *Drosophila melanogaster*: An ELISA-based method for tracking
individual ejaculates. Behav Ecol Sociobiol. 2009;63(10):1505–13.

53. Vahed K, Parker DJ, Gilbert JDJ. Larger testes are associated with a higher level of polyandry, but
a smaller ejaculate volume, across bushcricket species (*Tettigoniidae*). Biol Lett. 2011;7(2):261–
4.

54. Friesen CR, Powers DR, Copenhaver PE, Mason RT. Size dependence in non-sperm ejaculate
production is reflected in daily energy expenditure and resting metabolic rate. J Exp Biol.
2015;218(11):1796.

55. Kelly CD, Jennions MD. Sexual selection and sperm quantity: Meta-analyses of strategic
ejaculation. Biol Rev. 2011;86(4):863–84.

56. Hopkins BR, Sepil I, Thézénas ML, Craig JF, Miller T, Charles PD, et al. Divergent allocation of
sperm and the seminal proteome along a competition gradient in *Drosophila melanogaster*. Proc
Natl Acad Sci U S A. 2019;116(36):17925–33.

57. Hauschteck-Jungen E, Burkard W, Jungen H, Burch-Schwaller R. The loss of Y-sperm in 'sex ratio'
(SR) males of *Drosophila subobscura* is compensated. Genetica. 1987;74(1):27–30.

58. Hauschteck-Jungen E, Maurer B. Sperm dysfunction in sex ratio males of *Drosophila subobscura*.
Genetica. 1976;46(4):459–77.

59. Barreau C, Benson E, Gudmannsdottir E, Newton F, White-Cooper H. Post-meiotic transcription
in *Drosophila* testes. Development. 2008;135(11):1897–902.

60. White-Cooper H. Tissue, cell type and stage-specific ectopic gene expression and RNAi induction
in the *Drosophila* testis. Spermatogenesis. 2012;2(1):11–22.

61. Joly D, Luck N, Dejonghe B. Adaptation to long sperm in *Drosophila*: Correlated development of
the sperm roller and sperm packaging. J Exp Zoolog B Mol Dev Evol. 2008;310(2):167–78.

62. Belote JM, Zhong L. Duplicated proteasome subunit genes in *Drosophila* and their roles in
spermatogenesis. Heredity. 2009;103(1):23–31.

63. Sepil I, Hopkins BR, Dean R, Bath E, Friedman S, Swanson B, et al. Male reproductive aging arises
via multifaceted mating-dependent sperm and seminal proteome declines, but is postponable in
Drosophila. Proc Natl Acad Sci U S A. 2020;117(29):17094–103.

64. Wilkinson GS, Christianson SJ, Brand CL, Ru G, Shell W. Haldane's rule is linked to extraordinary
sex ratios and sperm length in stalk-eyed flies. Genetics. 2014;198(3):1167–81.

65. Orr HA, Irving S. Segregation distortion in hybrids between the Bogota and USA subspecies of
Drosophila pseudoobscura. Genetics. 2005;169(2):671–82.

66. Pinzone CA, Dyer KA. Association of polyandry and sex-ratio drive prevalence in natural
populations of *Drosophila neotestacea*. Proc R Soc B Biol Sci [Internet]. 2013;280(1769).
Available from: http://dx.doi.org/10.1098/rspb.2013.1397

67. Finnegan S, White NJ, Koh D, Camus MF, Fowler K, Pomiankowski A. Meiotic drive reduces egg-
to-adult viability in stalk-eyed flies. Proc R Soc B Biol Sci. 2019;286(1910).

68. Keais GL, Hanson MA, Gowen BE, Perlman SJ. X chromosome drive in a widespread Palearctic
woodland fly, *Drosophila testacea*. J Evol Biol. 2017;30(6):1185–94.

69. Dermitzakis ET, Masly JP, Waldrip HM, Clark AG. Non-mendelian segregation of sex
chromosomes in heterospecific *Drosophila* males. Genetics. 2000;154(2):687–94.

70. Esvelt KM, Gemmell NJ. Conservation demands safe gene drive. PLoS Biol. 2017;15(11):1–9.

71. Wolfner MF. Tokens of love: Functions and regulation of *Drosophila* male accessory gland
products. Insect Biochem Mol Biol. 1997;27(3):179–92.

72. Hall JC. The Mating of a Fly. Science. 1994;264(5166):1702–14.

73. Singh SR, Singh BN, Hoenigsberg HF. Female remating, sperm competition and sexual selection
in *Drosophila*. Genet Mol Res. 2002;1(3):178–215.

74. Baker RH, Denniff M, Futerman P, Fowler K, Pomiankowski A, Chapman T. Accessory gland size
influences time to sexual maturity and mating frequency in the stalk-eyed fly, *Cyrtodiopsis
dalmanni*. Behav Ecol. 2003;14(5):607–11.

75. Rogers DW, Baker RH, Chapman T, Denniff M, Pomiankowski A, Fowler K. Direct and correlated
responses to artificial selection on male mating frequency in the stalk-eyed fly *Cyrtodiopsis
dalmanni*. J Evol Biol. 2005;18(3):642–50.

592