Title: A comprehensive *in vivo* screen of yeast farnesyltransferase activity reveals

broad reactivity across a majority of CXXX sequences

Authors: June H. Kim[1], Emily R. Hildebrandt[1], Anushka Sarkar[1], Wayland Yeung[2], La

Ryel A. Waldon[1], Natarajan Kannan[1, 2], Walter K. Schmidt[1]

[1]Department of Biochemistry and Molecular Biology, University of Georgia, Athens GA

30602

[2]Institute of Bioinformatics, University of Georgia, Athens GA 30602

Corresponding author: Walter K. Schmidt, wschmidt@uga.edu (email), 706-583-8241

(phone), 706-542-1738 (fax).

Running title: Yeast FTase acts on CXXX sequences

Keywords: protein farnesylation, farnesyltransferase, yeast genetics, substrate

specificity, next-generation sequencing

Abstract (word count 208)

The current understanding of farnesyltransferase (FTase) specificity was

pioneered through investigations of reporters like Ras and Ras-related proteins that

possess a C-terminal CaaX motif that consists of 4 amino acid residues: Cysteine –

aliphatic$_1$ – aliphatic$_2$ – variable (X). These studies led to the finding that proteins with

the CaaX motif are subject to a 3-step post-translational modification pathway involving

farnesylation, proteolysis, and carboxylmethylation. Emerging evidence indicates,

however, that FTase can farnesylate sequences outside the CaaX motif and that these

sequences do not undergo the canonical 3-step pathway. In this work, we report a

comprehensive evaluation of all possible CXXX sequences as FTase targets using the

reporter Ydj1, an Hsp40 chaperone that only requires farnesylation for its activity. Our

genetic and high throughput sequencing approach reveals an unprecedented profile of

sequences that yeast FTase can recognize *in vivo*, which effectively expands the

potential target space of FTase within the yeast proteome. We also document that yeast

FTase specificity is majorly influenced by restrictive amino acids at a$_2$ and X positions

as opposed to the resemblance of CaaX motif as previously regarded. This first

complete evaluation of CXXX space expands the complexity of protein isoprenylation

and marks a key step forward in understanding the potential scope of targets for this

isoprenylation pathway.

Introduction

Isoprenylation is a post-translational modification (PTM) catalyzed by several isoprenylation enzymes (i.e., FTase, GGTase-I, GGTase-II, GGTase-III). This PTM enhances protein association with cellular membranes or strengthens protein-protein interactions (Lane and Beese 2006, Leung, Baron et al. 2007, Tate, Kalesh et al. 2015, Wang and Casey 2016). Collectively, isoprenylated proteins assist in cellular signaling, cell cycle regulation, development, aging, and various other biological processes (Greenwood, Steinman et al. 2006, Palsuledesai and Distefano 2015, Jeong, Suazo et al. 2018).

Ras signaling GTPases are often cited as classical examples of isoprenylated proteins (Willumsen, Christensen et al. 1984, Wright and Philips 2006, Zhou, Wiener et al. 2016). Ras isoprenylation is catalyzed by farnesyltransferase (FTase), which acts upon a C-terminal CaaX motif consisting of 4 amino acid residues: Cysteine – aliphatic$_1$ – aliphatic$_2$ – variable (X). FTase appends the C15 isoprenoid lipid donated by farnesyl pyrophosphate to the CaaX cysteine via a thioether linkage. After farnesylation, Ras undergoes the coupled modifications of endoproteolysis to remove the -aaX portion of the motif, followed by carboxylmethylation of the farnesylated cysteine that becomes the new C-terminal residue. These PTMs regulate Ras plasma membrane localization and

function, and have been extensively studied due to the significance of Ras GTPases in

human disease such as cancer (Tamanoi 2011, Cox, Der et al. 2015, Hobbs, Der et al.

2016). As a result, previous Ras-based investigations have led to the general model

that the three CaaX PTMs (i.e., isoprenylation, proteolysis, methylation) commonly

occur across a range of proteins that are collectively referred to as CaaX proteins. CaaX

PTMs exist in all eukaryotic species, and the impact of these PTMs on CaaX protein

biology are highly conserved across model systems (Omer, Kral et al. 1993, Cazzanelli,

Pereira et al. 2018, Ravishankar, Hildebrandt et al. 2023). This is especially evident in

the conserved similarities beween and mammalian and yeast FTase structure and

function (Kohl, Diehl et al. 1991, Gomez, Goodman et al. 1993, Omer, Kral et al. 1993).

Not all farnesylated proteins undergo all three CaaX PTMs. An example is the

Saccharomyces cerevisiae Hsp40 chaperone Ydj1 (*Sc*Ydj1) that is farnesylated on its

C-terminal CASQ sequence but not subjected to endoproteolysis or

carboxylmethylation. Farnesylation of Ydj1 is required for optimal yeast growth at

elevated temperatures (>37 °C) and subjecting Ydj1 to all three CaaX PTMs negatively

impacts this Ydj1-dependent thermotolerance growth phenotype (Caplan et al., 1992,

Hildebrandt et al., 2016). We have defined this non-canonical pathway leading to a

farnesylation-only PTM as the "shunt" farnesylation pathway (**Figure 1**). The

observation of the shunt pathway raises the question of whether the previous use of

canonically-modified CaaX protein reporters (e.g., Ras, **a**-factor), which are subject to

the additional constraints of proteolysis and carboxylmethylation, has limited the breadth

of sequences that can be identified as FTase substrates.

Many past investigations have attempted to probe FTase specificity. Initially, the

screening of FTase substrates was explored *in vitro* on a case-by-case basis using

recombinant CaaX proteins and synthetic peptides with [3H]-FPP or [3H]-GGPP, which

was inconvenient and labor intensive for large scale studies (Moores, Schaber et al.

1991, Caplin, Hettich et al. 1994). *In vitro* investigations using peptide libraries and

metabolic labeling have extended this effort but have been limited in scope and cost-

prohibitive, making it difficult to conduct systematic studies of the 8000 CXXX sequence

space (Reiss, Stradley et al. 1991, Boutin, Marande et al. 1999, Krzysiak, Scott et al.

2007, Krzysiak, Aditya et al. 2010, Wang, Dozier et al. 2014, Tate, Kalesh et al. 2015,

Suazo, Schaber et al. 2016, Storck, Morales-Sanfrutos et al. 2019). While *in silico*

prediction models based on structural analysis of mammalian FTase are available and

have potential to define isoprenylatable sequences, these models often fail at predicting

farnesylated proteins with non-canonical CaaX motifs such as *Sc*Ydj1 (CASQ),

*Hs*DNAJA2 (CAHQ), *Sc*Pex19 (CKQQ), *Hs*Lkb1 (CKQQ) and *Sc*Nap1 (CKQS) and

often lack orthologous *in vivo* reporter data to validate predictions (Collins, Reoma et al.

2000, Sapkota, Kieloch et al. 2001, Reid, Terry et al. 2004, Maurer-Stroh and

Eisenhaber 2005, Lane and Beese 2006, Maurer-Stroh, Koranda et al. 2007, London,

Lamphear et al. 2011, Berger, Yeung et al. 2022). Most of the *in vivo* reporters used to

probe CaaX space have relied on CaaX protein reporters requiring 3-step modification

and the superposition of 3 enzyme specificities for their activities (Boyartchuk, Ashby et

al. 1997, Stein, Kubala et al. 2015). Moreover, recent reports demonstrating FTase

activity on shortened and extended sequences highlight the flexibility of CaaX substrate

lengths, additionally complicating the recognized model of FTase CaaX specificity

(Ashok, Hildebrandt et al. 2020, Schey, Buttery et al. 2021).

The reliance of previous studies on canonical protein reporters and incomplete

peptide reporter sets, in conjunction with our shunt pathway observations, led us to

hypothesize that the full scope of FTase targets remains unknown. To address this gap

in knowledge, we developed *Sc*Ydj1 as a genetic reporter to elucidate the specificity of

the yeast FTase across all 8000 CXXX sequences. Our results indicate that yeast

FTase has a much broader target specificity than previously defined by the canonical

CaaX motif.

Materials and Methods

Yeast strains: Strains used in this study are listed in **Table 1**. All plasmid-

transformations into yeast were performed using a lithium acetate-based transformation

procedure unless otherwise stated (Elble 1992).

Plasmid construction for Ydj1-CKQx variants and His-Nap1: Plasmids used in this study

are listed in **Table 2**. Newly created plasmids needed for this study were constructed

using PCR-directed recombinational cloning consistent with previously reported

methods (Oldenburg, Vo et al. 1997, Hildebrandt, Cheng et al. 2016, Berger, Kim et al.

2018). For Ydj1-CKQX plasmids, pWS1132 that was linearized with *Nhe1* was co-

transformed into yWS304 along with a PCR-derived DNA fragment encoding a new

CaaX sequence. For the His-Nap1 plasmid (pWS1474), pRS316 that was linearized

with *BamHI*, *XbaI* and *XhoI* was con-transformed into BY4741 along with a PCR derived

DNA fragment encoding the *NAP1* gene. The resultant Nap1 plasmid (pWS1318) was

additionally modified to encode an octa-His tag at the 5´ end of the *NAP1* gene; the tag

was introduced by recombination using a PCR derived fragment and pWS1318 that had

been linearized with *BsaBI*. In all cases, candidate plasmids recovered from yeast

surviving appropriate genetic selection were evaluated by both restriction digest and

DNA sequence analyses (GENEWIZ / Azenta Life Sciences, South Plainfield, NJ;

Eurofins Genomics, Louisville, KY).

<u>Plasmid construction for NGS library</u>: The *YDJ1-CXXX* plasmid library (pWS1775) was

designed to represent all 8000 "XXX" amino acid combinations where each amino acid

is represented by a single unique codon by using trimer phosphoroamidites, aka "Trimer

20", incorporated into the mutagenic oligos (Integrated DNA Technologies). To create

the library, pWS1132 was first modified to introduce a silent unique *BsrGI* site directly 5´

to the SASQ encoding sequence using oligo oWS1330 (5´-

AACTATGATTCCGATGAAGAAGAACAAGGTGGCGAAGGTGTACAATCTGCATCTCA

ATGATTTTC), creating pWS1728 (*CEN URA3 YDJ1-BsrGI-SASQ*). Next, PCR-derived

DNA fragments were generated by pairing oWS1359 (5´-

TATGATTCCGATGAAGAAGAACAAGGTGGCGAAGGTGTACAATGT-iTriMix20-

iTriMix20-iTriMix20-TGATTTTCTTGATAAAAAAAGATC-3´) and oWS1308 (5´-

CAGCATATAATCCCTGCTTTA-3´) and pWS1728 template using a high-fidelity Q5

polymerase (New England Biolabs, Ipswich, MA). The PCR products were purified

(Omega E.Z.N.A Cycle Pure kit, Omega Bio-tek Inc., Norcross GA), digested with BsrGI

and AflII, re-purified, and ligated into pWS1728 (BsrGI and AflII digested) using high

concentration T4 DNA ligase (NEB M0202T). Ligations were transformed into DH5α *E.*

coli prepared using Z-Competent cell kit (G-Biosciences, St. Louis, MO) and plated to

achieve approximately 10^4 colonies per plate. In total, 606,800 colonies, each

representing a unique clone, were pooled into a single cell pellet, followed by DNA

purification (Omega E.Z.N.A Midiprep kit, Omega Bio-tek Inc., Norcross GA).

<u>Preparation of naïve yeast library containing Ydj1-CXXX variants</u>: The *E. coli*-derived

Ydj1-CXXX plasmid library was transformed into yWS304 (*ydj1Δ*) via Frozen-EZ Yeast

transformation II Kit by Zymo Research (Irvine, CA) according to manufacturer's

instructions such that over 3 million individual colonies were recovered across multiple

SC-Uracil plates on the same day. The colonies were collected by gently washing the

plates with liquid medium, and cells concentrated by centrifugation to remove excess

supernatant. The collected cells were stored at -80 °C as 100 µL aliquot stocks in 15%

glycerol at a concentration of ~2×10^9 cells per 1 mL. For all subsequent studies, freshly

thawed stock vials were used and not re-frozen for repeated use.

<u>Thermotolerance selection of Ydj1-CXXX variants *en masse* and plasmid library</u>

<u>recovery</u>: An aliquot of the naïve yeast library expressing Ydj1-CXXX variants was

thawed and used for each iteration of this assay. In brief, ~582,000 cells were diluted

into 40 mL of room temperature SC-Uracil liquid media, and this premix was divided

across 4 test tubes where sets of 2 tubes were incubated at either permissive (25 °C) or

restrictive temperature (37 °C). Each test tube containing diluted culture was rapidly

thermally equilibrated using an appropriate temperature water bath for 30 minutes

before being placed onto a rotating wheel in an incubator at the same temperature. The

cultures were incubated for approximately 24 to 48 hours until A_{600} 2.0 was reached,

cells were collected by centrifugation, and plasmids were isolated from the recovered

cell populations using a commercial kit (OMEGA Bio-Tek E.Z.N.A. Yeast Miniprep kit)

as per manufacturer's instructions. The experiment was performed twice across two

different days, with each condition performed in duplicate, for a total of 8 replicates per

temperature condition.

<u>Next-generation sequencing and data analysis</u>: The yeast plasmids recovered after

thermoselection were subject to a shortened PCR with 15 cycles with oligonucleotide

pairs oWS1408 (5´-

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGCAGCATATAATCCCTGCTTTA-

3´) and oWS1409 (5´-

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGTCCAGGGGTGGTGCAAACTATG

-3´) using a high-fidelity Q5 polymerase (New England Biolabs, Ipswich, MA) to attach

overhang sequences. The PCR products were cleaned (OMEGA Bio-Tek E.Z.N.A.

Cycle Pure Kit), quantified (Synergy H1 Hybrid Multi-Mode Microplate Reader), and

submitted to the Georgia Genomics Bioinformatics Center (GGBC, Athens, GA) for

Illumina MiSeq library assembly and sequencing (single-paired 100 bp reads starting 38

bases upstream of the CXXX region of interest).

The NGS analysis yielded over 10 million total reads, with ~96% of individual

reads at or above the quality score of Q30. All 8 replicates from 25 °C yielded high-

quality reads while only 7 replicates from 37 °C passed the quality cutoff. These 15

experimental samples, which represented about 3 million total reads, were carried

forward for downstream analysis. In parallel, NGS was used to sequence 10 replicates

of the *E. coli* plasmid library and 10 replicates of the plasmids extracted from the naïve

yeast library prior to selection.

To assess the likelihood of farnesylation of Ydj1-CXXX variants, the reads within

each replicate were first assessed to determine the number of occurrences for each

unique CXXX sequence present. The occurrence of each CXXX sequence was

summed across all replicates of the same temperature condition, then normalized to a

frequency value. The latter was calculated by dividing the occurrence value for each

unique CXXX by the total number of occurrences of all CXXX sequences (i.e.,

frequency = $\text{count}_{unique}/\text{count}_{total}$) for each of the 25 °C and 37 °C data sets. Lastly, the

frequency value of each unique CXXX sequence was used to determine a unique

enrichment factor (EF) score. This was calculated by dividing the frequency value of a

unique CXXX sequence at restrictive temperature by that of the 25 °C condition (i.e., EF

score = $\text{frequency}_{37\ °C}/\text{frequency}_{25\ °C}$). Some CXXX sequences were not recovered at

25 °C, yet present at higher temperatures (n=9; CFFM, CIFF, CILF, CIYM, CNWC,

CTFA, CVFF, CVLW, CWIA). To account for such cases, a correction factor of +1 was

applied across all frequency values at 25 °C so that 1 would be the denominator for

CXXX sequences with 0 occurrences at 25 °C. We also applied the same correction

factor of +1 to all frequency values at 37 °C for consistency.

<u>Weblogo sequence alignments:</u> The top 5% (n=400) sequences for both Ydj1-based

and Ras-based screens were evaluated by Weblogo

(http://weblogo.berkeley.edu/logo.cgi) using a custom color scheme (Crooks, Hon et al.

2004). Cys was set to blue; polar charged amino acids were set to green (Asp, Arg, Glu,

His, and Lys); polar uncharged residues were set to black (Asn, Gln, Ser, Thr, and Tyr);

branched chain amino acids were set to red (Ile, Leu and Val); all other residues were

set to purple (Ala, Gly, Met, Phe, Pro, and Trp).

Heatmap analysis: CXXX variants were clustered into 20-member groups having shared

amino acids pairs. An average EF value was determined for these groups to allow for

contextual analyses exploring the relationships between a_1 vs. a_2, a_1 vs. X, and a_2 vs. X.

The averages were then analyzed with Microsoft Excel version 16.65 using the

Conditional Formatting, Color Scales (Green - Yellow - Red) function to produce three

distinct heatmaps (HMs). The values reported within each cell of a heatmap represent

the average EF value for each respective 20-member group. The EF scores associated

with RRS data (Stein et al., 2015) were evaluated in the same manner.

To determine confidence intervals, the average HM values were averaged across

each individual row and column of the heatmap, and the averages were used to

determine a standard deviation for either the row or column set of values. A standard

deviation calculation was next used to determine a 95% confidence interval that was in

turn used to establish high and low cutoffs for identifying positive selection or negative

restriction. A pattern was deemed significant when 18 or more of each 20-member set

(i.e., >90%) were above or below the statistical cutoff.

<u>Likelihood of prenylation calculations for each CXXX variant</u>: The averaged EF values

from each of the 3 HMs were summed to produce a HM score. For example, to predict

the likelihood of prenylation for CASQ, the averages of Cys-Ala-Ser-X (n=20 varied at

the X position), Cys-Ala-X-Gln (n=20 varied at the a_2 position), and Cys-X-Ser-Gln

(n=20 varied at the a_1 position) were first calculated, then these three HM values were

summed to represent the HM score of CASQ.

<u>Ydj1-CXXX thermotolerance assay</u>: The assay was performed as previously described

(Hildebrandt, Cheng et al. 2016, Berger, Kim et al. 2018, Ashok, Hildebrandt et al.

2020). In brief, yeast cultures in SC-Uracil media were incubated at 25 °C until

saturation, then serially diluted 10-fold in YPD before being pinned onto YPD solid

medium. Plates were incubated for 2-4 days at 25 °C, 37 °C, and 39 °C prior to results

being digitally scanned face down without lids using a Cannon flat-bed scanner (300

dpi; TIFF format). Scanned images were adjusted for consistency in image rotation,

contrast, and size before being copied onto Microsoft PowerPoint version 16.65 for final

figure assembly. The experiment was performed twice in duplicate.

<u>Gel-shift assay</u>: The assay was performed as previously described (Hildebrandt, Cheng

et al. 2016, Berger, Kim et al. 2018, Schey, Buttery et al. 2021). Whole cell lysates of

mid log cells were prepared and separated by sodium dodecyl sulfate-polyacrylamide

gel electrophoresis (SDS-PAGE; 6% stacking with 9.5% resolving gel) then transferred

onto nitrocellulose. Blots were blocked with 5% milk then sequentially incubated with

rabbit anti-Ydj1 primary antibody (courtesy of A. Caplan) and HRP-conjugated goat anti-

rabbit secondary antibody (Kindle biosciences, Greenwich, CT). Fluorescence was

detected using the KwikQuant Western Blot Detection Kit (Kindle Biosciences) and a

KwikQuant Imager and as per manufacturer's instructions. Protein bands were

quantified using NIH ImageJ, and resulting values were used for calculating ratios for

prenylated and non-prenylated bands. The blots containing yeast Nap1 were treated

similarly but incubated with mouse anti-HIS primary antibody (Thermo Fisher Scientific,

Waltham, MA) and HRP-conjugated sheep anti-mouse secondary antibody (Kindle

biosciences, Greenwich, CT).

<u>Decision tree modeling:</u> Sequence motif features were generated by one-hot encoding

for each of the variable CaaX motif sites: a_1, a_2, and X. Additional binary features were

included to describe whether the variable residue was within a given set of residues. To

define these sets, we enumerated all possible amino acid combinations with a max set

size of 5. In total, 65,097 features were considered. While building the decision tree

classifier, entropy was used to evaluate the quality of potential splits. To determine

more generalizable rules, trees were allowed a maximum depth of 3 and all nodes were

required to have a minimum of 50 samples. This method was implemented using Scikit-

learn (v0.22.2) using the DecisionTreeClassifier class.

Results

<u>Many CXXX sequences sustain Ydj1-dependent growth of yeast at high temperature</u>:

Ydj1 follows an isoprenylation-only pathway in contrast to canonical reporters

(i.e., Ras GTPases, **a**-factor) that undergo additional downstream modifications

(Trueblood, Boyartchuk et al. 1997, Stein, Kubala et al. 2015) (**Figure 1**). As such, the

use of Ydj1 as a reporter to probe yeast FTase specificity reduces the risk of introducing

additional specificity bias from CaaX proteases. To evaluate the substrate scope of

FTase, we adapted methods from a previous yeast study that utilized a Ras-based

CXXX reporter (i.e., Ras Recruitment System (RRS)) along with competitive growth

enrichment and next-generation sequencing (NGS) methods (Stein, Kubala et al. 2015).

In our case, we investigated the ability of Ydj1-CXXX variants to sustain high

temperature growth in a *ydj1Δ* genetic background (**Figure S1**). A plasmid library of

Ydj1-CXXX variants was generated using single codons for each amino acid so that all

8000 CXXX sequences are represented in a relatively small library (see Material and

Methods). Compared to traditional plasmid library construction relying on fully or partly

degenerate oligonucleotides, a Trimer 20-based strategy was used with the aim of

yielding a balanced library with respect to codon redundancy (i.e., no over-

representation of amino acids with multiple codons) and no introduction of early stop

codons, which effectively reduces the number of copies needed to reach statistical

confidence for full coverage (Firth and Patrick 2008). While 110,500 independent clones

were minimally required for statistical 100% coverage, our library contains 606,800

independent clones (i.e., 6x full coverage). By contrast, constructing a library with fully

degenerate oligonucleotides to vary 3 amino acids would have required 3,200,000

independent clones for 100% coverage, leading to much higher labor and time costs for

this study.

The Ydj1-CXXX plasmid library DNA was purified from *Escherichia coli* before

being transformed into the *ydj1Δ* yeast strain to yield a naïve *ydj1Δ/YDJ1-CXXX* yeast

library. Both the *E. coli* plasmid library and the naïve yeast library were confirmed by

NGS to contain all potential Ydj1-CXXX variants (**File S1**). Ligation efficiency of vector

alone was determined to be ~0.3% relative to vector plus insert, indicating that a small

portion of the library could encode *YDJ1-SASQ* (i.e., uncut parent plasmid).

Consistently, *YDJ1-SASQ* was observed at frequencies of 0.065% and 0.071% in the *E.*

coli and naïve yeast libraries, respectively. The frequencies of all CXXX sequences from

the *E. coli* and naïve yeast libraries were also graphed, and no significant changes in

the frequency profile between the two libraries were observed (**Figure S2**). This

analysis further revealed that the libraries were not perfectly balanced such that the

extremes exhibited a ~10x range in abundance.

The naïve yeast library was propagated for approximately 8 generations at

permissive (25 °C) and selective (37 °C) temperatures in liquid media until the culture

was well saturated (A_{600} ~2.0). The selective temperature condition was expected to

enrich for farnesylated Ydj1-CXXX variants. NGS methods were then used to identify all

the CXXX sequences present in each culture, from which frequencies were determined

for each CXXX sequence in each temperature group. Similarities in experimental design

allowed for direct comparison of Ydj1 and Ras-based data sets. In the RRS study, the

likelihood of farnesylation was reported by an Enrichment Factor (RRS EF) that was

defined by the frequency of a specific CXXX sequence occurring at 37 °C divided by its

frequency at 25 °C (i.e., $\text{frequency}_{selective}/\text{frequency}_{permissive}$). A high RRS EF was

interpreted as a high possibility of farnesylation. We performed a similar calculation for

the Ydj1 data.

From our analysis, we observed that EFs for the Ydj1-based screen exhibited a

narrower range (EF: 0.036 for CWWC to 13.898 for CVFF) relative to the RRS EFs for

the Ras-based screen (RRS EF: 0.008 for CLRS to 70.667 for CYCM). We interpret

these ranges to indicate that the majority of all CXXX sequences can support growth in

the Ydj1-based screen, whereas a smaller number of sequences undergo higher

enrichment during the thermoselection process in the Ras-based screen. The

comparison also revealed that the EF profiles offered two distinct sequence landscapes,

especially for farnesylated sequences that are well characterized: non-canonical CASQ

(*Sc*Ydj1) and CKQQ (*Hs*STK11/Lkb1); canonical CVIA (*Sc* **a**-factor) and CVLS (*Hs*H-

Ras). In the Ydj1 NGS-based screen, non-canonical sequences CASQ (EF: 1.608) and

CKQQ (EF: 1.627) outperformed canonical CaaX sequences such as CVIA (EF: 0.406)

and CVLS (EF: 0.494) (**Figure 2A**). By contrast, in the Ras-based screen, CASQ (RRS

EF: 0.399) and CKQQ (RRS EF: 0.399) were significantly less enriched while CVIA

(RRS EF: 10.124) and CVLS (RRS EF: 11.804) ranked amongst the top hits (**Figure

2B**). The EFs of all CXXX sequences resulting from the Ydj1 screen are reported in **File

S2**.

Our observations suggest that the Ydj1-based screen best enriches non-

canonical sequences whereas the Ras-based screen best enriches canonical

sequences. The top 5% (n=400) of hits from the Ydj1 NGS-based assay did not exhibit

an obvious consensus sequence (**Figure 2C**); however, the same number of top hits

from the Ras-based screen were enriched with aliphatic amino acids at the a_1 and a_2

positions (**Figure 2D**). In the context of the Ras reporter, an aliphatic amino acid is

especially prominent at the a_2 position, which has been historically regarded as a

requirement for FTase specificity. The top 5% of hits from both the Ydj1-based and Ras-

based screens were also displayed as a 4D plot (**Figures 2E, F**; the size of the spot is

the 4^{th} dimension and represents relative abundance). This analysis revealed that the

top hits were more widely dispersed across CXXX sequence space in the Ydj1-based

data relative to the Ras-based data, and that the most abundant sequences differed

between the data sets. Analysis of the data as 3D plots, viewed along each of the 4D

plot axes such that Lysine (K) is the nearest amino acid, provided additional details

about the sequence space covered by each screen (**Figure S3**). In these 3D plots, the

spots representing relative abundance are stacked behind each other, and the total

number of spots occupying a particular node is not easily discerned. Although this

arrangement makes it difficult to make detailed conclusions about the depth of coverage

at each node (i.e., the number of amino acids present), it allows for clear conclusions

about restrictions (i.e., amino acids that are not tolerated independent of context). From

the perspective of a_1, both data sets lack several amino acids at a_2 (i.e., D, E, G, K, and

R) and a single amino acid at X (i.e., R) (**Figures S3A, B**). Specific to the Ras-based

data, additional amino acids were absent at a_2 (H and Y) and X (K, P, and W) or less

prevalent at a_2 (i.e., A, P, Q, S, T, and W). From the perspective of a_2, no amino acids

were absent at the a_1 position in both data sets, one amino acid was absent at the X

position in both data sets (i.e., R), and several amino acids were absent at the X

position within the Ras-based data set only (i.e., K, P, R, and W) (**Figures S3C, D**).

From the perspective of X, no amino acids were absent at the a_1 position, and the

amino acids that were absent at a_2 from the perspective of a_1 were again identified, with

one additional amino acid being absent (i.e., S) (**Figures S3E, F**). A general

interpretation of these observations is that many amino acids can be accommodated at

each position of the CXXX sequence independent of whether Ydj1 or Ras is the

reporter, except for charged residues that are strictly not tolerated at a_2 in either case.

There also appear to be additional exceptions in the context of the Ras reporter at both

the a_2 and X position that likely reflect its need for additional modification beyond initial

isoprenylation.

To further analyze our Ydj1 NGS-based data, we crosschecked CXXX

sequences that were previously identified as being farnesylated through a limited scope

genetic Ydj1-based Temperature Screen (YTS) (Berger, Kim et al. 2018). Mostly non-

canonical sequences were identified in the earlier study (n=153). Many of these

sequences exhibited high EFs in the present study (**Figure 3A**). When superposed on

the Ydj1 EF profile, most YTS identified sequences were positioned in the 4th quartile

(n=77) with fewer in the 3rd quartile (n=67) and fewest in the 2nd quartile (n=9). None of

the YTS hits were present in the 1st quartile. By contrast, superposition of the top hits

from the Ras-based screen (n=496; RRS EF > 3; the cutoff for positive hits of the Ras-

based screen) on the Ydj1 EF profile revealed positioning of sequences in the 2nd, 3rd,

and 4th quartiles of the EF profile (n=228, n=114, and n=153, respectively), with the

fewest in the 1st quartile (n=1) (**Figure 3B**). The few canonical sequences identified by

YTS (**Figure 3A**, dashed box; n=15) remained within the range displayed by RRS

sequences. Together, these results are fully consistent with our Ydj1 NGS-based

method being useful for enriching non-canonical sequences that are farnesylated in

addition to canonical sequences observed using the Ras reporter.

<u>Non-canonical CKQX sequences are targeted by FTase</u>:

Emerging evidence indicates that some CKQX sequences are farnesylated

despite the presence of non-aliphatic amino acids at a_1 and a_2 positions. CKQQ

farnesylation is well documented for human STK11/Lkb1, human Nap1L1 and yeast

Pex19, whereas CKQS derived from yeast Nap1 is farnesylated in the context of the

Ydj1 reporter (Collins, Reoma et al. 2000, Sapkota, Kieloch et al. 2001, Tate, Kalesh et

al. 2015, Storck, Morales-Sanfrutos et al. 2019, Berger, Yeung et al. 2022). We

additionally confirmed that farnesylation of CKQS occurs in the natural context of yeast

Nap1 itself using a gel-shift assay that evaluates the mobility of farnesylated species by

SDS-PAGE (**Figure S4A**). Of note, farnesylation has an opposite effect on Nap1

mobility when compared to Ydj1. Moreover, the ScanProsite tool

(https://prosite.expasy.org/scanprosite/) identifies 5179 entries across UniProtKB/Swiss-

Prot and UniProtKB/TrEMBL reference proteome sequences ending in CKQX across

eukaryotes (**File S3**). The majority of these sequences (i.e., 65%) contain CKQQ or

CKQS. Out of the 20 CKQX variants, most (n=17 positive hits) displayed a Ydj1 EF

consistent with a high likelihood of farnesylation (**Figure 4A**). The remaining sequences

(n=3) had low EFs, were well separated from the other CKQX sequences on the EF

profile, and were expected to have a low likelihood of modification (i.e., negative hits).

By comparison, none of the 20 CKQX sequences were predicted to be targets of FTase

in the Ras-based EF profile (**Figure 4B**).

The farnesylation status of Ydj1-CaaX variants were monitored by

thermotolerance and gel-shift assays (Caplan, Tsai et al. 1992, Hildebrandt, Cheng et

al. 2016). The thermotolerance assay confirmed the 17 CKQX positive hits to have

growth profiles similar to that of wildtype Ydj1, whereas the negative hits (n=3) had a

growth profile consistent with unmodified Ydj1 (**Figure 4C, Figure S4B**). In strong

agreement with the EFs and thermotolerance profiles, a prenyl-dependent gel-shift was

observed for the 17 positive Ydj1-CKQX hits and no shift observed for the three

negative hits. It should be noted that CKQR displayed a doublet gel pattern in both the

presence and absence of FTase. The reason for this doublet pattern is unknown,

however, the pattern was identical with and without FTase, so CKQR was deemed as

having 0% prenylation.

<u>Statistical analyses reveal yeast FTase specificity</u>:

Our evaluation of genetic and biochemical data suggested that FTase is not

limited to target sequences having aliphatic amino acids at a_1 and a_2. To identify factors

that could better indicate target specificity, we assessed the likelihood of prenylation for

sets of CaaX motifs that shared the same context, differing only at a single amino acid

position. Using heatmap analysis, we sought to identify potential patterns marked by

positive selection (colored in green) or negative restriction (colored in red) across a_1, a_2,

and X positions in both Ydj1-based and Ras-based results (**Figure 5**). These patterns

were identified as instances where the HM values of 18 or more of each 20-member set

(i.e., >90% of EF scores) were well above or below the statistical average for all values

(n=400) within a heatmap (**File S4**).

For the Ydj1-based data, no positive patterns were observed for the a_1 position in

combination with a_2 (**Figure 5A**), however, one restrictive pattern, D at a_1, was

observed in combination with X (**Figure 5B**). Generally, a_1 appears to tolerate many

different types of amino acids and does not seem to strongly influence reactivity with

FTase in either a positive or negative manner. When considering the a_2 position, no

positive patterns were observed, but several negative patterns were evident.

Specifically, D, K and R were restrictive in combination with either a_1 (**Figure 5A**) or X

(**Figure 5C**), with E being additionally restrictive in combination with a_1. Generally, a_2

also appears to tolerate many different types of amino acids except for charged amino

acids that negatively influence reactivity with FTase. For the X position, a positive

pattern, Q at X, was observed in combination with a_1 (**Figure 5B**). Several negative

patterns were also evident at the X position. Both P and R were restrictive in

combination with either a_1 or a_2, with K being additionally restrictive in the context of a_1.

Generally, X too seems to tolerate many different types of amino acids except for

positively charged amino acids and structurally constrained proline (**Figure 5C**). Thus,

despite repeated reports that FTase targets canonical CaaX sequences, our Ydj1-based

data strongly indicates that FTase tolerates most CXXX sequences unless amino acids

are present that restrict its specificity. Similar heatmap analysis of the Ras-based data

also revealed fewer positive than negative patterns. The two positive patterns were I at

a_2 (**Figure 5D**) and M at X (**Figure 5E**), both in combination with a_1. Most of the

negative patterns observed were primarily constrained to a_2 and X positions in all

contexts (**Figure 5C, F**). The a_1 position, on the other hand, displayed fewest negative

patterns; only two patterns were observed at E in combination with a_2 and K in

combination with X. The combined observations that both Ydj1 and Ras-based data

reveal high tolerance at the a_1 position, while a_2 and X positions have more restrictions

is consistent with previous reports on FTase specificity (Reid, Terry et al. 2004, Stein,

Kubala et al. 2015).

To fully account for the contextual information from heatmap analysis, the three

heatmap values from the Ydj1 NGS dataset were summed for each specific CaaX motif

to obtain a heatmap (HM) score. This score represents the sum of 3 averages, where

each average accounts for 20 EF data points. The advantage of the HM score is that it

normalizes statistical outliers that could skew results when using individual EF scores.

The HM scores, when a cutoff of 3 was applied, correlated well with prior evidence of

prenylation gathered from previous studies, outperforming published models such as

Prenylation Prediction Suite (PrePS), RRS, and our recently published SVM algorithm

for predicting farnesylation of non-canonical sequences (Maurer-Stroh and Eisenhaber

2005, Stein, Kubala et al. 2015, Berger, Yeung et al. 2022) (**Table 3**).

Lastly, we trained a predictive model using the results of the Ydj1-based NGS

screen to predict whether a given CXXX sequence can be modified based on a subset

of questions (decisions) and possible consequences in a decision-tree model (**Figure 6,**

Figure S5). Each question pertains to the whether certain amino acids are present in

the a_1, a_2, and X positions. The decision tree was created by systematically identifying

the best set of questions for predicting whether a CXXX sequence can be modified. This

allowed us to identify a general set of rules which govern FTase substrate specificity.

Based on our model, the biggest determinant of FTase activity is the restriction of D, E,

K, or R at the a_2 position, followed by the restriction of K, P, or R at the X position.

However, amino acids such as I, L, M and V at the a_2 position seem to neutralize the

negative effects of K, P, or R at the X position. Our findings repeatedly raise the

likelihood that restrictions, rather than tolerance, are applied to a few impermissible

amino acids at a_2 and X positions of potential substrates modified by FTase. It should

be noted that although this simple flowchart is useful in understanding general rules of

yeast FTase specificity, HM scores provide a more individualized prediction of the

prenylation status of each CXXX sequence.

Discussion

In this study, the use of Ydj1 as a reporter instead of Ras allowed for the

identification of yeast FTase targets that extend beyond the canonical CaaX consensus

sequence. Notably, this study corroborates previously published observations that non-

canonical sequences can be targets of FTase. Restrictions by a small number of amino

acids, rather than adherence to the canonical CaaX consensus, seem to guide FTase

specificity. We believe that these findings can be used to improve tools for predicting

farnesylation status that will outperform current prediction methods.

Structural analysis of mammalian FTase has previously shown that the aaX

portion of the protein substrate makes extensive Van der Waals contact with the

adjacent isoprenoid group, which is hypothesized to modulate product release (Long,

Casey et al. 2002, Reid, Terry et al. 2004). However, these interactions are not

predicted when non-aliphatic residues are at a_1 and a_2. Our finding that FTase can

indeed tolerate non-aliphatic residues at a_1 and a_2 raises the possibility that the catalytic

site of FTase may display a greater degree of structural flexibility beyond the

conformations that could be sampled from mammalian FTase in complex with canonical

substrates. Our findings also lead us to predict that binding of charged amino acids (D,

E, K, R) may be unstable in the a_2 binding pocket. Likewise, certain amino acids (K, P,

R) at the X position may be unable to coordinate with the product binding pocket of the

FTase. These situations may result in the formation of a farnesylated peptide that is

unable to be released from the enzyme, thus inhibiting FTase from turning over such

substrates efficiently. These possibilities could be resolved by future structural studies

involving non-canonical CXXX peptide substrates in complex with either yeast or

mammalian FTase, which are structurally and functionally homologous (Kohl, Diehl et

al. 1991, Gomez, Goodman et al. 1993, Omer, Kral et al. 1993).

Our study was focused on the recognition of CXXX sequences with respect to

FTase specificity. It has been observed, however, that upstream sequences can impact

FTase specificity (Cox, Graham et al. 1993). The impact of upstream sequences has

been explored by prediction algorithms, most notably by PrePS (Maurer-Stroh and

Eisenhaber 2005). It is critical to acknowledge that the Ydj1-CXXX NGS screen

recovered sequences that support high temperature growth in the context of Ydj1. Thus,

any effect of upstream sequence has not been explored in this work. Additionally, it is

possible that some Ydj1-CaaX variants affect growth in ways unrelated to prenylation

status, which could impact some of our results. Still, our YDJ1 HM scoring system

outperformed predictions made by PrePS, RRS, and SVM machine learning algorithm

(Maurer-Stroh and Eisenhaber 2005, Stein, Kubala et al. 2015, Berger, Yeung et al.

2022) (**Table 3**). Combining PrePS predictions that consider upstream sequence in

addition to the observations presented in this work is expected to provide the most

robust prediction for farnesylation of individual protein targets. As more of the

prenylation predictions are validated using additional techniques in the future, it will be

possible to determine which prediction methods are most reliable.

Many proteins undergo a process known as isoprenylation. Historically, FTase is

recognized as targeting the CaaX sequence but emerging evidence, including that

reported here, has indicated that FTase substrates are not limited to the canonical

CaaX sequence. Our methods, which primarily focused on yeast FTase, provides

insights into the broader specificity of this process which likely extends to human and

other FTase enzymes. The discovery of farnesylated non-canonical sequences such as

CKQX associated with large families of proteins like Stk11/Lkb1 and Nap1, which lack a

canonical CaaX motif, opens a new understanding of post-translational isoprenylation

and the potential for future discoveries in this field. Our results also demonstrate that the

target specificity of FTase is mainly due to restrictions at the a_2 and X positions, rather

than selection towards a canonical CaaX sequence. As Ras and other canonical

reporters rely on multiple modifications for function, previous studies using these

proteins may have reported on sequences that are suited for the combined specificities

of FTase and downstream enzymes (i.e., CaaX proteases *Sc*Ste24 or *Hs*Rce1) instead

of the specificity of FTase alone. Our study, in combination with data from previous

studies, provides new ways to advance the understanding of the specificities of each

enzymatic step associated with the post-translational isoprenylation pathway, laying out

a crucial step toward identifying its cellular targets and the extent to which they are

modified.

Data availability

Yeast strains and plasmids are available upon request. All relevant datasets for this

study are included in the supplemental files of the manuscript. **File S1** contains

frequency of CXXX sequences observed in naive libraries. **File S2** contains Enrichment

Factor results of Ydj1-based NGS. **File S3** contains CKQX hits identified on UniProt.

File S4 contains heatmap analysis. The data derived through the Ras-Recruitment

System was previously published and reanalyzed here (Stein, Kubala et al. 2015).

Acknowledgement

We thank Dr. James Hougland (Syracuse University) for early discussions about this

work, Dr. Avrom Caplan (City College of New York) for anti-Ydj1 antibody, and Somdut

Roy (Georgia Institute of Technology) for the raw Python 3 scripts for the 4D graphical

analysis. Lastly, we thank Schmidt lab members for constructive feedback and technical

assistance.

Funding and additional information

This research was supported by Public Health Service grant GM132606 from the

National Institute of General Medical Sciences (WKS, NK).

Conflict of interest

The authors have declared no competing interests exist.

References

Ashok, S., E. R. Hildebrandt, C. S. Ruiz, D. S. Hardgrove, D. W. Coreno, W. K. Schmidt and J. L. Hougland (2020). "Protein Farnesyltransferase Catalyzes Unanticipated Farnesylation and Geranylgeranylation of Shortened Target Sequences." Biochemistry **59**(11): 1149-1162.

Berger, B. M., J. H. Kim, E. R. Hildebrandt, I. C. Davis, M. C. Morgan, J. L. Hougland and W. K. Schmidt (2018). "Protein Isoprenylation in Yeast Targets COOH-Terminal Sequences Not Adhering to the CaaX Consensus." Genetics **210**(4): 1301-1316.

Berger, B. M., W. Yeung, A. Goyal, Z. Zhou, E. R. Hildebrandt, N. Kannan and W. K. Schmidt (2022). "Functional classification and validation of yeast prenylation motifs using machine learning and genetic reporters." PLoS One **17**(6): e0270128.

Boutin, J. A., W. Marande, L. Petit, A. Loynel, C. Desmet, E. Canet and J. L. Fauchere (1999). "Investigation of S-farnesyl transferase substrate specificity with combinatorial tetrapeptide libraries." Cell Signal **11**(1): 59-69.

Boyartchuk, V. L., M. N. Ashby and J. Rine (1997). "Modulation of Ras and a-factor function by carboxyl-terminal proteolysis." Science **275**(5307): 1796-1800.

Brachmann, C. B., A. Davies, G. J. Cost, E. Caputo, J. Li, P. Hieter and J. D. Boeke (1998). "Designer deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications." Yeast **14**(2): 115-132.

Caplan, A. J., J. Tsai, P. J. Casey and M. G. Douglas (1992). "Farnesylation of YDJ1p is

required for function at elevated growth temperatures in Saccharomyces

cerevisiae." J Biol Chem **267**(26): 18890-18895.

Caplin, B. E., L. A. Hettich and M. S. Marshall (1994). "Substrate characterization of the

Saccharomyces cerevisiae protein farnesyltransferase and type-I protein

geranylgeranyltransferase." Biochim Biophys Acta **1205**(1): 39-48.

Cazzanelli, G., F. Pereira, S. Alves, R. Francisco, L. Azevedo, P. Dias Carvalho, A.

Almeida, M. Corte-Real, M. J. Oliveira, C. Lucas, M. J. Sousa and A. Preto

(2018). "The Yeast Saccharomyces cerevisiae as a Model for Understanding

RAS Proteins and their Role in Human Tumorigenesis." Cells **7**(2).

Collins, S. P., J. L. Reoma, D. M. Gamm and M. D. Uhler (2000). "LKB1, a novel

serine/threonine protein kinase and potential tumour suppressor, is

phosphorylated by cAMP-dependent protein kinase (PKA) and prenylated in

vivo." Biochem J **345 Pt 3**(Pt 3): 673-680.

Cox, A. D., C. J. Der and M. R. Philips (2015). "Targeting RAS Membrane Association:

Back to the Future for Anti-RAS Drug Discovery?" Clin Cancer Res **21**(8): 1819-

1827.

Cox, A. D., S. M. Graham, P. A. Solski, J. E. Buss and C. J. Der (1993). "The carboxyl-

terminal CXXX sequence of Gi alpha, but not Rab5 or Rab11, supports Ras

processing and transforming activity." J Biol Chem **268**(16): 11548-11552.

Crooks, G. E., G. Hon, J. M. Chandonia and S. E. Brenner (2004). "WebLogo: a

sequence logo generator." Genome Res **14**(6): 1188-1190.

Elble, R. (1992). "A simple and efficient procedure for transformation of yeasts."

Biotechniques **13**(1): 18-20.

Firth, A. E. and W. M. Patrick (2008). "GLUE-IT and PEDEL-AA: new programmes for

analyzing protein diversity in randomized libraries." Nucleic Acids Res **36**(Web

Server issue): W281-285.

Giaever, G., A. M. Chu, L. Ni, C. Connelly, L. Riles, S. Veronneau, S. Dow, A. Lucau-

Danila, K. Anderson, B. Andre, A. P. Arkin, A. Astromoff, M. El-Bakkoury, R.

Bangham, R. Benito, S. Brachat, S. Campanaro, M. Curtiss, K. Davis, A.

Deutschbauer, K. D. Entian, P. Flaherty, F. Foury, D. J. Garfinkel, M. Gerstein, D.

Gotte, U. Guldener, J. H. Hegemann, S. Hempel, Z. Herman, D. F. Jaramillo, D.

E. Kelly, S. L. Kelly, P. Kotter, D. LaBonte, D. C. Lamb, N. Lan, H. Liang, H. Liao,

L. Liu, C. Luo, M. Lussier, R. Mao, P. Menard, S. L. Ooi, J. L. Revuelta, C. J.

Roberts, M. Rose, P. Ross-Macdonald, B. Scherens, G. Schimmack, B. Shafer,

D. D. Shoemaker, S. Sookhai-Mahadeo, R. K. Storms, J. N. Strathern, G. Valle,

M. Voet, G. Volckaert, C. Y. Wang, T. R. Ward, J. Wilhelmy, E. A. Winzeler, Y.

Yang, G. Yen, E. Youngman, K. Yu, H. Bussey, J. D. Boeke, M. Snyder, P.

Philippsen, R. W. Davis and M. Johnston (2002). "Functional profiling of the

Saccharomyces cerevisiae genome." Nature **418**(6896): 387-391.

Gomez, R., L. E. Goodman, S. K. Tripathy, E. O'Rourke, V. Manne and F. Tamanoi

(1993). "Purified yeast protein farnesyltransferase is structurally and functionally

similar to its mammalian counterpart." Biochem J **289 (Pt 1)**(Pt 1): 25-31.

Greenwood, J., L. Steinman and S. S. Zamvil (2006). "Statin therapy and autoimmune

disease: from protein prenylation to immunomodulation." <u>Nat Rev Immunol</u> **6**(5):

358-370.

Hildebrandt, E. R., M. Cheng, P. Zhao, J. H. Kim, L. Wells and W. K. Schmidt (2016). "A

shunt pathway limits the CaaX processing of Hsp40 Ydj1p and regulates Ydj1p-

dependent phenotypes." <u>Elife</u> **5**.

Hobbs, G. A., C. J. Der and K. L. Rossman (2016). "RAS isoforms and mutations in

cancer at a glance." <u>J Cell Sci</u> **129**(7): 1287-1292.

Jeong, A., K. F. Suazo, W. G. Wood, M. D. Distefano and L. Li (2018). "Isoprenoids and

protein prenylation: implications in the pathogenesis and therapeutic intervention

of Alzheimer's disease." <u>Crit Rev Biochem Mol Biol</u> **53**(3): 279-310.

Kohl, N. E., R. E. Diehl, M. D. Schaber, E. Rands, D. D. Soderman, B. He, S. L.

Moores, D. L. Pompliano, S. Ferro-Novick, S. Powers and et al. (1991).

"Structural homology among mammalian and Saccharomyces cerevisiae

isoprenyl-protein transferases." <u>J Biol Chem</u> **266**(28): 18884-18888.

Krzysiak, A. J., A. V. Aditya, J. L. Hougland, C. A. Fierke and R. A. Gibbs (2010).

"Synthesis and screening of a CaaL peptide library versus FTase reveals a

surprising number of substrates." <u>Bioorg Med Chem Lett</u> **20**(2): 767-770.

Krzysiak, A. J., S. A. Scott, K. A. Hicks, C. A. Fierke and R. A. Gibbs (2007). "Evaluation

of protein farnesyltransferase substrate specificity using synthetic peptide

libraries." <u>Bioorg Med Chem Lett</u> **17**(20): 5548-5551.

Lane, K. T. and L. S. Beese (2006). "Thematic review series: lipid posttranslational

modifications. Structural biology of protein farnesyltransferase and

geranylgeranyltransferase type I." J Lipid Res **47**(4): 681-699.

Leung, K. F., R. Baron, B. R. Ali, A. I. Magee and M. C. Seabra (2007). "Rab GTPases

containing a CAAX motif are processed post-geranylgeranylation by proteolysis

and methylation." J Biol Chem **282**(2): 1487-1497.

London, N., C. L. Lamphear, J. L. Hougland, C. A. Fierke and O. Schueler-Furman

(2011). "Identification of a novel class of farnesylation targets by structure-based

modeling of binding specificity." PLoS Comput Biol **7**(10): e1002170.

Long, S. B., P. J. Casey and L. S. Beese (2002). "Reaction path of protein

farnesyltransferase at atomic resolution." Nature **419**(6907): 645-650.

Maurer-Stroh, S. and F. Eisenhaber (2005). "Refinement and prediction of protein

prenylation motifs." Genome Biol **6**(6): R55.

Maurer-Stroh, S., M. Koranda, W. Benetka, G. Schneider, F. L. Sirota and F.

Eisenhaber (2007). "Towards complete sets of farnesylated and

geranylgeranylated proteins." PLoS Comput Biol **3**(4): e66.

Moores, S. L., M. D. Schaber, S. D. Mosser, E. Rands, M. B. O'Hara, V. M. Garsky, M.

S. Marshall, D. L. Pompliano and J. B. Gibbs (1991). "Sequence dependence of

protein isoprenylation." J Biol Chem **266**(22): 14603-14610.

Oldenburg, K. R., K. T. Vo, S. Michaelis and C. Paddon (1997). "Recombination-

mediated PCR-directed plasmid construction in vivo in yeast." Nucleic Acids Res

25(2): 451-452.

Omer, C. A., A. M. Kral, R. E. Diehl, G. C. Prendergast, S. Powers, C. M. Allen, J. B. Gibbs and N. E. Kohl (1993). "Characterization of recombinant human farnesyl-protein transferase: cloning, expression, farnesyl diphosphate binding, and functional homology with yeast prenyl-protein transferases." Biochemistry **32**(19): 5167-5176.

Palsuledesai, C. C. and M. D. Distefano (2015). "Protein prenylation: enzymes, therapeutics, and biotechnology applications." ACS Chem Biol **10**(1): 51-62.

Ravishankar, R., E. R. Hildebrandt, G. Greenway, N. Asad, S. Gore, T. M. Dore and W. K. Schmidt (2023). "Specific Disruption of Ras2 CAAX Proteolysis Alters Its Localization and Function." Microbiol Spectr: e0269222.

Reid, T. S., K. L. Terry, P. J. Casey and L. S. Beese (2004). "Crystallographic analysis of CaaX prenyltransferases complexed with substrates defines rules of protein substrate selectivity." J Mol Biol **343**(2): 417-433.

Reiss, Y., S. J. Stradley, L. M. Gierasch, M. S. Brown and J. L. Goldstein (1991). "Sequence requirement for peptide recognition by rat brain p21ras protein farnesyltransferase." Proc Natl Acad Sci U S A **88**(3): 732-736.

Sapkota, G. P., A. Kieloch, J. M. Lizcano, S. Lain, J. S. Arthur, M. R. Williams, N. Morrice, M. Deak and D. R. Alessi (2001). "Phosphorylation of the protein kinase mutated in Peutz-Jeghers cancer syndrome, LKB1/STK11, at Ser431 by p90(RSK) and cAMP-dependent protein kinase, but not its farnesylation at Cys(433), is essential for LKB1 to suppress cell vrowth." J Biol Chem **276**(22): 19469-19482.

Schey, G. L., P. H. Buttery, E. R. Hildebrandt, S. X. Novak, W. K. Schmidt, J. L.

Hougland and M. D. Distefano (2021). "MALDI-MS Analysis of Peptide Libraries

Expands the Scope of Substrates for Farnesyltransferase." Int J Mol Sci **22**(21).

Sikorski, R. S. and P. Hieter (1989). "A system of shuttle vectors and yeast host strains

designed for efficient manipulation of DNA in Saccharomyces cerevisiae."

Genetics **122**(1): 19-27.

Stein, V., M. H. Kubala, J. Steen, S. M. Grimmond and K. Alexandrov (2015). "Towards

the systematic mapping and engineering of the protein prenylation machinery in

Saccharomyces cerevisiae." PLoS One **10**(3): e0120716.

Storck, E. M., J. Morales-Sanfrutos, R. A. Serwa, N. Panyain, T. Lanyon-Hogg, T.

Tolmachova, L. N. Ventimiglia, J. Martin-Serrano, M. C. Seabra, B. Wojciak-

Stothard and E. W. Tate (2019). "Dual chemical probes enable quantitative

system-wide analysis of protein prenylation and prenylation dynamics." Nat

Chem **11**(6): 552-561.

Suazo, K. F., C. Schaber, C. C. Palsuledesai, A. R. Odom John and M. D. Distefano

(2016). "Global proteomic analysis of prenylated proteins in Plasmodium

falciparum using an alkyne-modified isoprenoid analogue." Sci Rep **6**: 38615.

Tamanoi, F. (2011). "Ras signaling in yeast." Genes Cancer **2**(3): 210-215.

Tate, E. W., K. A. Kalesh, T. Lanyon-Hogg, E. M. Storck and E. Thinon (2015). "Global

profiling of protein lipidation using chemical proteomic technologies." Curr Opin

Chem Biol **24**: 48-57.

Trueblood, C. E., V. L. Boyartchuk and J. Rine (1997). "Substrate specificity

determinants in the farnesyltransferase beta-subunit." Proc Natl Acad Sci U S A

94(20): 10774-10779.

Wang, M. and P. J. Casey (2016). "Protein prenylation: unique fats make their mark on

biology." Nat Rev Mol Cell Biol **17**(2): 110-122.

Wang, Y. C., J. K. Dozier, L. S. Beese and M. D. Distefano (2014). "Rapid analysis of

protein farnesyltransferase substrate specificity using peptide libraries and

isoprenoid diphosphate analogues." ACS Chem Biol **9**(8): 1726-1735.

Willumsen, B. M., A. Christensen, N. L. Hubbert, A. G. Papageorge and D. R. Lowy

(1984). "The p21 ras C-terminus is required for transformation and membrane

association." Nature **310**(5978): 583-586.

Wright, L. P. and M. R. Philips (2006). "Thematic review series: lipid posttranslational

modifications. CAAX modification and membrane targeting of Ras." J Lipid Res

47(5): 883-891.

Zhou, M., H. Wiener, W. Su, Y. Zhou, C. Liot, I. Ahearn, J. F. Hancock and M. R. Philips

(2016). "VPS35 binds farnesylated N-Ras in the cytosol to regulate N-Ras

trafficking." J Cell Biol **214**(4): 445-458.

767 **Tables**

768 **Table 1.** *Yeast strains used in this study.*

Identifier	Genotype	Reference
BY4741	*MAT**a** his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*	(Brachmann, Davies et al. 1998)
yWS304	*MAT**a** his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 ydj1Δ::KAN^R*	(Giaever, Chu et al. 2002)
yWS1632	*MAT**a** his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 ram1::KAN^R*	(Giaever, Chu et al. 2002)
yWS2542	*MAT**a** his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 ydj1Δ::NAT^R ram1Δ::KAN^R*	(Berger, Kim et al. 2018)
yWS2544	*MAT**a** his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 ydj1Δ::NAT^R*	(Berger, Kim et al. 2018)

770 **Table 2.** *Plasmids used in this study.*

Identifier	Genotype	Reference
pRS316	*CEN URA3*	(Sikorski and Hieter 1989)
pWS942	*CEN URA3 YDJ1*	(Hildebrandt, Cheng et al. 2016)
pWS1132	*CEN URA3 YDJ1-SASQ*	(Hildebrandt, Cheng et al. 2016)
pWS1286	*CEN URA3 YDJ1-CVIA*	(Hildebrandt, Cheng et al. 2016)
pWS1318	*CEN URA3 NAP1*	This study
pWS1407	*CEN URA3 YDJ1-CKQQ*	This study
pWS1411	*CEN URA3 YDJ1-CKQS*	(Berger, Yeung et al. 2022)
pWS1474	*CEN URA3 His8-NAP1*	This study
pWS1563	*CEN URA3 YDJ1-SKQQ*	This study
pWS1728	*CEN URA3 YDJ1-BsrGI-SASQ*	This study
pWS2083	*CEN URA3 YDJ1-CKQA*	This study
pWS2017	*CEN URA3 YDJ1-CKQC*	This study
pWS2018	*CEN URA3 YDJ1-CKQD*	This study
pWS2019	*CEN URA3 YDJ1-CKQE*	This study
pWS2020	*CEN URA3 YDJ1-CKQF*	This study
pWS2021	*CEN URA3 YDJ1-CKQG*	(Berger, Yeung et al. 2022)
pWS2022	*CEN URA3 YDJ1-CKQH*	(Berger, Yeung et al. 2022)
pWS2023	*CEN URA3 YDJ1-CKQI*	This study
pWS2024	*CEN URA3 YDJ1-CKQK*	This study
pWS2025	*CEN URA3 YDJ1-CKQL*	(Berger, Yeung et al. 2022)

pWS2026	*CEN URA3 YDJ1-CKQM*	This study
pWS2027	*CEN URA3 YDJ1-CKQN*	This study
pWS2028	*CEN URA3 YDJ1-CKQP*	This study
pWS2029	*CEN URA3 YDJ1-CKQR*	This study
pWS2030	*CEN URA3 YDJ1-CKQT*	This study
pWS2031	*CEN URA3 YDJ1-CKQV*	This study
pWS2033	*CEN URA3 YDJ1-CKQW*	This study

772 **Table 3**. *A comprehensive overview of in-house prediction versus published models.*

Protein[1]	Motif (n=48)	Observed[2]	%[2]	PrePS[3]	SVM[4]	RRS EF[5]	YDJ1 HM
Ras2	CIIS	+	100	1.281	1	9.011500	4.07
Hmg1	CIKS	-	41	-3.588	0	0.034717	1.93
Rho2	CIIL	+	100	0.817	1	10.313800	3.65
Ssp2	CIDL	-	0	-4.893	0	0.254070	1.82
Skt5, MiY1	CVIM	+	100	2.200	1	3.371400	3.66
Tbs1	CVKM	+	100	-2.909	0	0.066541	2.47
YDL022C-A	CSII	+	100	0.987	1	11.350000	3.34
YBR096W	CSEI	-	0	-5.759	0	0.149720	1.88
YMR265C	CSNA	+	100	-4.848	0	0.039925	3.84
Pet18	CYNA	+	100	-5.843	0	0.079849	4.95
Lih1	CSGL	-	61	-4.577	0	0.041301	2.79
Cup1	CSGK	-	59	-6.461	0	0.099811	1.75
Nap1	CKQS	+	100	-0.975	1	0.066541	4.97
Cst26	CFIF	+	100	-1.924	1	3.194000	4.81
YIL134C-A	CAPY	+	100	-2.164	1	0.199620	4.73
Atr1	CTVA	+	100	0.273	1	7.775800	4.55
Las21	CALD	+	100	-2.887	1	0.028518	4.20
YDL009C	CAVS	+	100	0.237	1	6.607500	4.69
Sua5	CIQF	+	100	-0.601	1	0.199620	5.49
NA	CAAQ	+	100	-3.280	1	0.399250	4.82
NA	CAHQ	+	100	-3.599	1	0.399250	4.74
NA	CASA	+	100	-2.693	1	0.243980	3.91
NA	CKQH	+	70	-2.442	1	0.399250	4.37
NA	CNLI	+	95	-3.804	1	0.133080	3.84
NA	CSFL	+	68	-4.425	1	0.108890	4.28
NA	CVAA	+	100	-2.789	1	0.079849	4.81
NA	CVFM	+	100	-2.939	1	2.395500	4.76
NA	CKQG	+	73	-1.876	0	0.099811	4.06

NA	CKQL	+	50	-1.680	0	0.057035	3.84
NA	CQTS	+	100	-0.922	0	0.057035	5.44
NA	CQSQ	+	100	-1.624	1	0.159700	5.27
NA	CKQA	+	95	-1.149	1	0.133080	4.57
NA	CKQC	+	100	-0.745	1	0.099811	4.12
NA	CKQD	+	67	-4.103	0	0.066541	3.51
NA	CKQE	-	19	-3.745	0	0.066541	3.36
NA	CKQF	-	28	-2.345	0	0.057035	4.62
NA	CKQH	+	73	-2.442	1	0.399250	4.37
NA	CKQI	+	70	-1.700	1	0.099811	3.90
NA	CKQK	-	0	-4.010	0	0.036295	2.22
NA	CKQM	+	100	-1.045	1	0.133080	4.50
NA	CKQN	+	88	-5.046	0	0.199620	4.48
NA	CKQP	-	13	-2.887	0	0.399250	1.97
NA	CKQQ	+	100	-0.747	1	0.399250	5.35
NA	CKQR	-	0	-4.533	0	0.399250	1.77
NA	CKQT	+	96	-1.874	1	0.079849	4.68
NA	CKQV	+	100	-1.744	1	0.133080	4.50
NA	CKQW	+	71	-5.184	0	0.199620	3.57
NA	CKQY	+	100	-3.326	0	0.399250	4.50
		# predicted correctly	30	38	17	45	
		# false positive	0	0	0	2	
		# false negative	18	10	31	1	
		% predicted correctly	62.5%	79.2%	35.4%	93.8%	
		% false positive	0.0%	0.0%	0.0%	5.1%	
		% false negative	64.3%	50.0%	75.6%	11.1%	

774   [1] Protein or gene name as reported in the Saccharomyces Genome Database

775   [2] Prenylation status and % gel-shift observed in context of Ydj1

776   [3] Values reported from (Maurer-Stroh and Eisenhaber 2005); positive cutoff > -2.

777   [4] Values reported from (Berger, Yeung et al. 2022); positive cutoff > 0.

778   [5] Values reported from (Stein, Kubala et al. 2015); positive cutoff > 3.

Figure legends

Figure 1. *The post-translational isoprenylation pathway.* Ydj1 and Lkb1/STK11 have non-canonical CaaX motifs that follow an isoprenylation-only pathway (i.e., shunted). Ras and **a**-factor have canonical CaaX motifs and are subject to the superposition of 3 modifications.

Figure S1. *Experimental scheme for genetic screening of Ydj1-CXXX variants assessed via next-generation sequencing (NGS).* A plasmid library encoding all 8000 Ydj1-CXXX variants was generated via ligation-based cloning method using PCR fragments derived using Trimer20 oligonucleotides (Integrated DNA Technologies, Newark, NJ). The library DNA was purified from *E. coli*, and plasmids were transformed into the *ydj1Δ* yeast strain. This strain cannot grow at temperatures >37 °C unless complemented by farnesylated Ydj1-CXXX variant. Populations of *ydj1Δ/YDJ1-CXXX* transformants were incubated at permissive (25 °C) and selective (37 °C) temperatures, and plasmids recovered were subject to NGS analysis. Schematic illustration created with Microsoft PowerPoint and BioRender.com.

Figure S2. *Frequencies of CXXX sequences in naïve yeast library versus E. coli plasmid library.* CXXX frequencies from the naïve yeast and *E. coli* plasmid libraries were plotted against each other. No significant change in the frequency profile of individual sequences was observed between the two libraries. Libraries were not perfectly balanced, however, across all CXXX sequences; the extremes exhibited a ~10x range in relative abundance.

Figure 2. *Comparison of results from the Ydj1-CXXX NGS screen and previously*

published Ras-CXXX NGS screen. Enrichment profiles of (**A**) Ydj1-based and (**B**) Ras-

based screens offer two distinct sequence landscapes. Annotated controls: non-

canonical *Sc*Ydj1 (CASQ) and *Hs*Lkb1/STK11 (CKQQ); canonical *Sc* **a**-factor (CVIA)

and *Hs*H-Ras (CVLS). Weblogo frequency representation of top 5% (n=400) of hits for

(**C**) Ydj1-based and (**D**) Ras-based screens. A 4D representation of the space occupied

by top 5% of hits recovered from (**E**) Ydj1-based and (**F**) Ras-based screens. The size

of the dot is the 4th dimension and corresponds to the enrichment factor of the CXXX

motif within the respective screen.

Figure S3. *3D analysis of top 5% of sequences from Ydj1-based and Ras-based data*

sets. The data described in **Figures 2E, F** was plotted in 3D from the perspective of

each axis of the 4D plot such that Lysine (K) is the nearest amino acid. The size of the

spot is the 3^{rd} dimension and represents relative abundance. The perspectives are

along the a_1 axis (**A**, **B**), the a_2 axis (**C**, **D**), and X axis (**E**, **F**) for the Ydj1-based data (**A**,

C, **E**) and the Ras-based data (**B**, **D**, **F**).

Figure 3. *Relative performance of previously identified farnesylated CXXX sequences.*

Sequences previously identified in a (**A**) Ydj1-based genetic selection or a (**B**) Ras-

based competitive growth assay were superimposed on the Ydj1 EF profile. Each

sequence is represented as a point with an associated score where the second Y-axis

refers to the scoring range used in these previous studies (YTS T-score - Ydj1-based

Temperature Screen Thermotolerance Score; RRS EF – Ras Recruitment System

Enrichment Factor). The limited number of canonical sequences that were identified by

YTS are identified by a dashed box (n=15).

Figure 4. *Evaluation of CKQX variants as FTase substrates.* Enrichment profiles of

CKQX sequences in (**A**) Ydj1-based and (**B**) Ras-based screens. **C**) Thermotolerance

and gel-shift assays of Ydj1-CKQX variants provide supporting evidence towards the

prenylation of non-canonical CKQX sequences by yeast FTase. Thermotolerance

assays were performed by culturing yeast to the same density, applying a 10x serial

dilution to the cultures, spotting dilution sets onto YPD rich media, and incubating at

indicated temperatures. Notably, plate-based selection requires a slightly higher

temperature to mimic the growth profile observed on liquid-based selection. The strains

used were transformants of yWS304 containing the indicated Ydj1-CXXX variant. Gel

shift assays were performed for the indicated Ydj1-CXXX variants in the presence and

absence of FTase activity. Total yeast extracts were analyzed by SDS-PAGE and

immunoblotting. The strains used were yWS2544 (+FTase) and yWS2542 (*ram1Δ*; -

FTase. Abbreviations: Seq = Sequence, EF = Enrichment Factor, % = percent

prenylation.

Figure S4. *Thermotolerance assay of control sequences.* **A**) Gel-shift assays were

performed for Ydj1 (CASQ) and His8-Nap1 (CKQS) in the presence and absence of

FTase activity. Total yeast extracts were analyzed by SDS-PAGE and immunoblotting.

The strains used were BY4741 (+FTase) and yWS1632 (*ram1Δ*; -FTase). **B**)

Thermotolerance assays were performed with the indicated Ydj1-CXXX variants as

described for **Figure 4**. Abbreviations: Seq = Sequence, EF = Enrichment Factor, % =

percent prenylation, N/A = not applicable.

Figure 5. *Heatmap analysis reveals contextual determinants of FTase specificity.* Ydj1-

based comparisons of **A**) a_1 vs. a_2, **B**) a_1 vs. X, and **C**) a_2 vs. X positions, and Ras-

based comparisons of **D**) a_1 vs. a_2, **E**) a_1 vs. X, and **F**) a_2 vs. X positions. Values were

generated by averaging the 20 EFs of each CXXX combinations. Color scheme: red

(low likelihood of farnesylation) to green (high likelihood of farnesylation).

Figure 6. *A simplified decision tree for predicting yeast FTase substrates.* The output of

Ydj1 EF scores was fit into a decision tree model considering each of the variable CaaX

motif sites: a_1, a_2, and X. Our results strongly suggest that a major determinant of FTase

activity depends on a select few discriminatory amino acids at a_2 (D, E, K, R) and X (K,

P, R) positions, with the potential for certain allowable amino acids at a_2 (I, L, M, V) that

can mitigate discrimination caused at the X position. For the full decision tree, see

Figure S5.

Figure S5. *Full decision tree model.* Sequence motif features were generated by one-

hot encoding each of the variable CaaX motif sites. Additional binary features were

included that describe whether the variable residue was within a given set of residues.

To define these rules, all possible residue combinations up to 5 amino acids were

considered, yielding a total of 65,097 features. While building the decision tree

classifier, entropy was used to evaluate the quality of potential splits. Trees were

allowed a maximum depth of 3 in order to determine more generalizable rules and all

nodes were required to have a minimum of 50 samples. Samples refer to the number of

CXXX variants from the previous node that fall within the category. Value refers to the

distribution of CXXX variants that fall within specified EF ranges (0-1, 1-2, 2-3, 3-4, >4).

File S1. *Frequency of CXXX sequences observed in naive libraries*

File S2. *Enrichment factor results of Ydj1-based NGS*

File S3. *CKQX hits identified on UniProt*

File S4. *Heatmap analysis*

Figure 1

Figure S1

Plasmid library prepared by PCR and ligation

E. coli transformation

E. coli library

Plasmid isolation

E. coli plasmid library

Yeast transformation

Yeast library

Yeast library aliquots stored in -80 °C

Inoculation into liquid media for thermoselection

Experimental set-up in technical duplicates

Propagation at permissive or selective temperature

Yeast DNA isolation, NGS library preparation

Illumina MiSeq 100 basepair reads

Figure S2

Figure 2

Figure S3

A Top 5% Ydj1 sequences : a₁ perspective, 'K' in front

B Top 5% Ras sequences : a₁ perspective, 'K' in front

C Top 5% Ydj1 sequences : a₂ perspective, 'K' in front

D Top 5% Ras sequences : a₂ perspective, 'K' in front

E Top 5% Ydj1 sequences : X perspective, 'K' in front

F Top 5% Ras sequences : X perspective, 'K' in front

Figure 3

Figure S4

A

B

Seq.	EF	25 °C	37 °C	39 °C
Vector	N/A			
SASQ	N/A			
SKQQ	N/A			
CASQ	1.61			
CKQQ	1.63			
CKQS	2.12			
CVIA	0.41			

Figure 4

Figure 5

Ydj1 Results

RRS Results

Figure 6

Figure S5