# Skeletal morphology of bird wings is determined by thermoregulatory demand for heat dissipation in warmer climates

Brian C. Weeks[1*], Christina Harvey[2], Joseph A. Tobias[3], Catherine Sheard[4], Zhizhuo Zhou[5,6], David F. Fouhey[5]

[1]School for Environment and Sustainability, University of Michigan; Ann Arbor, USA

[2]Department of Mechanical and Aerospace Engineering, University of California Davis; California, USA

[3]Department of Life Sciences, Imperial College London; London, UK

[4]School of Earth Sciences, University of Bristol; Bristol, UK

[5]Department of Computer Science and Engineering, University of Michigan; Michigan, USA

[6]School of Computer Science, Carnegie Mellon University; Pennsylvania, USA

*Corresponding Author: Brian C. Weeks, Email: bcweeks@umich.edu, tel: (614) 216-9266, 440 Church Street, Dana Building, Ann Arbor, MI 48109

**Author Contributions:** Conceptualization: BCW, JAT, Methodology: BCW, CH, CS, ZZ, DFF, Investigation: BCW, CH, CS, ZZ, DFF, Visualization: BCW, Writing – original draft: BCW, Writing – review & editing: CH, CS, JAT, ZZ, DFF

**Competing Interest Statement:** Authors declare that they have no competing interests.

**Classification:** BIOLOGICAL SCIENCES; Ecology

**Keywords:** Allen's Rule, heat dissipation, morphology, flight, wing evolution

**This PDF file includes:**

    Main Text

    Figures 1 & 2

**Abstract**

The tendency for animals in warmer climates to be longer-limbed (Allen's Rule) is widely attributed to the demands of thermoregulation. However, the underlying mechanism remains unclear, because variation in limb-length can typically be driven by selection for both efficient heat retention and increased heat dissipation capacity. Using comparative phylogenetic models, we find that occurrence in warmer climates is associated with longer wing bones for 1,520 species of passerine birds. The highly vascularized musculature along these bones is only uncovered during flight, when the wings function as the primary site of heat exchange, cooling the organism by dissipating excess heat generated by muscular activity. Conversely, the musculature along the wing bones is insulated by feathering when at rest, playing a negligible role in heat retention, even in colder climates. Given this asymmetry in thermoregulatory roles, we can identify the positive relationship between temperature and wing bone length as a phenotypic gradient shaped by increased demand for heat dissipation in warmer climates. Our findings provide a clear illustration of the mechanism by which global warming can drive spatial and temporal trends in appendage length, and also highlight the role of heat dissipation in reshaping even the most critical features of vertebrate anatomy.

**Significance Statement**

Animals tend to be longer-limbed in warmer climates, but it remains unclear whether this pattern is driven by selection for cold tolerance at low temperatures or efficient heat dissipation at high temperatures. We show that for 1,520 species of passerines, bird wing bones are relatively longer in warmer climates. The vascularized musculature along these bones primarily functions in heat exchange during flight, when the overwhelming thermoregulatory challenge is dissipating heat, suggesting longer wing-bone length is driven by heat dissipation demands. Our findings reveal the pervasive impacts of thermoregulatory demands on even the most important functional traits.

**Main Text**

**Introduction**

The need to balance energy production with heat dissipation is a fundamental imperative for endotherms. Animal species have overcome this thermoregulatory challenge in numerous ways, including via morphological adaptations to climatic niches. These adaptations form the basis for classic ecogeographic patterns, including the tendency to have smaller bodies [Bergmann's Rule (1)] or longer limbs [Allen's Rule (2)] in warmer climates, both of which have been reported in a wide range of animals (3–6), including humans (7). While the standard explanation for these patterns is that body size and appendage length are shaped by selection for either cold tolerance or heat dissipation, their mechanistic basis remains unclear. Unraveling these selective mechanisms is an important step towards a more complete understanding of morphological evolution (8), constraints on energy expenditure (9), and the potential utility of spatial correlations between morphology and temperature as predictors of the impacts of global warming (10).

Focusing on Allen's Rule, we ask whether demand for shedding excess heat – as opposed to conserving heat – can drive large-scale trends in appendage length, specifically in the avian wing. Much of what we know about the effects of temperature on appendage length comes from studies of the beaks and leg bones of birds (11). There is a limit to what these appendages can tell us, however, because they regulate both heat dissipation to prevent overheating (12) and to improve cold tolerance (13), complicating the identification of underlying mechanisms. A more mechanistic understanding can be achieved by focusing on the vascularized components of wings.

In birds, the high energetic demands of flight require a 10–23-fold increase in metabolism (14), and the underlying conversion of chemical energy to mechanical energy results in nearly 90% of the metabolic energy expended during flight being converted to heat (15). Consequently, during prolonged flights, less than 1% of the heat generated by a bird is retained (16). This presents a critical thermoregulatory challenge: during flight much more heat is generated than when at rest, and nearly all of it must be dissipated to the environment. The requisite increase in heat dissipation is primarily achieved through convective heat loss from the vascularized brachial areas of the wings (17). This is in stark contrast to the negligible amounts of heat lost from the wings when

individuals are not flying, and may need to retain heat, as the wings are held against the

torso with the vascularized tissue well insulated under a dense layer of plumage (18).

We characterize wing heat dissipation capacity for 1,520 species of passerine

birds (~33% of all passerines) using a neural network (19) to measure key skeletal

components of the wing – the humerus and the ulna – on digital photographs of 7,366

museum skeletal specimens originating from all continents except Antarctica (Fig. 1).

The lengths of these wing bones determine the extent of the vascularized brachial

regions of the wings, which are the main windows of heat exchange with the

environment during flight (17), and are therefore expected to determine the heat

dissipation capacity of the wings (Fig. 2 and Fig. S1). We couple these skeletal data with

estimates of primary flight feather length and body mass (20) to quantify

macroecological patterns in wing structure. Specifically, we test the hypothesis that

hotter temperatures are associated with longer wing bones, driven by demand for

increased heat dissipation capacity in warmer climates.

**Results**

After controlling for the range of geographic and aerodynamic variables that are

expected to influence wing structure, we find a significant positive relationship between

temperature and the length of the wing bones ($\beta$ = 0.05; $P$ < 0.001; Fig. 2). We are

unlikely to have missed any alternative covariate because our model has high

explanatory power, explaining nearly all variation in wing-bone length (marginal $R^2$ =

0.97; conditional $R^2$ > 0.99). We conclude that higher temperatures are strongly

associated with longer wing bones, as predicted by Allen's Rule.

The thermoregulation-associated pressures shaping wing structure play out

within a selective landscape dominated by the demands of flight. The length of the wing

bones is strongly predicted by scaling relationships, including a significant and strong

positive relationship between wing-bone length and the length of the primary flight

feathers ($\beta$ = 1.05; $P$ < 0.001; Fig. 2). The length of the primary flight feathers is highly

correlated with mass (Pearson's Correlation $(R)$ = 0.88), and when mass is included as a

predictor in the model rather than primary flight feather length, mass is similarly

positively associated with wing bone length ($\beta$ = 1.13; $P$ < 0.001; Table S4). Thus, wing

bones are longer in larger birds which have longer primary flight feathers.

Variation in the length of the wing bones relative to the length of the primary flight

feathers is expected to impact flight dynamics. To control for the impacts of the demands

of flight on wing bone length, we apply aerodynamic theory to estimate a "pitch index", a

modified index of pitch agility (21) that measures the ability of a bird to generate angular

acceleration around the pitch axis (i.e., longitudinal maneuverability; Fig. S1). We also

calculate the aspect ratio of the wings, a determinant of gliding flight efficiency (15, 22,

23). By controlling for pitch index and aspect ratio, we account for pressures associated

with maneuverability and flight efficiency on the length of the wing bones. We find that

pitch index is significantly negatively associated with wing-bone length ($\beta$ = -0.04; $P$ <

0.001). Wing aspect ratio is similarly significantly negatively associated with wing-bone

length ($\beta$ = -0.23; $P$ < 0.001).

We also control for differences in migratory strategy, as long-distance

movements are expected to exert strong selective pressure on wing shape and length

(24). We find that intermediate- and long-distance migration, both common life history

strategies in birds adapted to highly seasonal environments (25), are positively

associated with the length of the wing bones (intermediate: $\beta$ = 0.04; $P$ < 0.001; long

distance: $\beta$ = 0.05; $P$ < 0.001).

Finally, given the range of important ecological factors that vary with latitude,

such as ecological specialization (26), species richness (27), dispersal ability (24), and

geographical range size (Rapoport's Rule), we included latitude as a predictor in our

model, based on the centroid of each species' range. We find that the distance of this

centroid from the equator (absolute value of latitude) is positively associated with wing-

bone length ($\beta$ = 0.02; $P$ = 0.01). This counter-intuitive relationship may be partially

explained by migration, as our metric of migration is intended to capture major

differences in life history and flight demands rather than more nuanced differences in

migratory distance that may correlate with latitude and be associated with wing

morphology (28). Latitude is also correlated with the quotient of wingspan divided by

mass ($R$ = 0.18, $P$ < 0.001), and while mass is highly correlated with wing feather length

($R$ = 0.88), it is not perfectly correlated. Thus, latitude may be capturing some variation

in wing bone length that is driven by mass. In accordance with this idea, when mass is

included in the model rather than primary flight feather length, latitude is no longer

significantly associated with wing bone length (*SI*).

**Discussion**

The wings of flying vertebrates are generally assumed to be shaped exclusively by the demands of aerodynamics and flight efficiency (24, 29). Our results add a new dimension by revealing a significant positive relationship between temperature and wing bone length in a broad sample of passerine birds. This pattern is consistent across a range of temperature metrics, as well as robust to statistical controls for allometry (*SI*) and a suite of ecological, geographic, and aerodynamic variables expected to influence wing structure. The anatomical trend we detect across global climatic gradients supports the hypothesis that thermoregulatory demands may be a widespread and overlooked driver of bird wing evolution.

We interpret the relationship between temperature and wing bone length as an outcome of selection for increased capacity to release heat into the environment in warmer climates because wings only become meaningful sites of heat exchange during flight (17, 18). A flying bird needs to release large amounts of body-heat to avoid over-heating and to permit the sustained energy expenditure required to maintain flight. Our findings therefore align with heat dissipation limit theory (9), and suggest that this mechanism can help to explain the widespread correlation between higher temperature and longer limb-length in animals (Allen's Rule).

Variation in thermoregulatory demands may also explain widespread increases in the wing lengths of migrant and resident birds, despite ongoing reductions in body size over recent decades (30, 31). The opposing evolutionary trajectories of wing length and body size have prompted much debate, with the unexpected change in wing morphology mainly attributed to the need for increasingly efficient flight (30, 31). Our results provide new insight, suggesting that while rising temperatures appear to drive reduced body size in line with the expectations of Bergmann's Rule, similar demand for efficient heat exchange also selects for increased size of wing bones, including the carpal and "hand" bones, which contribute to the measures of increased wing length reported by previous studies (30, 31).

Nearly all aspects of bird wing size and shape, including the structure of the forelimb, are expected to be broadly important to the physics of flight (22). Accordingly, we find a strong positive relationship between wing bone length and primary flight feather length (Fig. 2), reflecting the hard constraint on total wing length necessary to

achieve flight that is imposed by body size (32). We also find significant negative

relationships between aspect ratio and wing bone length, and between pitch index and

wing bone length. This suggests that gliding flight efficiency and maneuverability are

maximized in wings with relatively short wing bones compared to the primary flight

feathers. These non-thermal selective pressures on wing structure likely interact via

trade-offs with thermal mechanisms to determine the outcome of wing evolution.

Our results reveal that bird wings are complex traits shaped not only by flight

adaptations but also by the imperative for efficient thermoregulation. The connection

between wing bone length and temperature remains strongly significant even when we

account for allometry, and a full range of other contextual factors, such as ecological

niche (primary food source), latitude, and migratory strategy (Fig. 2). We also use

aerodynamic theory to control for selection associated with flight performance (21, 22),

and further account for flight performance-associated drivers of wing structure by

constraining our data to passerines, thus avoiding potential confounding effects related

to coarse differences in flight style on wing morphology (33). For example, heat

production during flight may be significantly reduced in birds that habitually soar or glide,

but this factor is removed in passerines, all of which use flapping flight. For these

reasons, our models achieve unusually high explanatory power (conditional $R^2$ > 0.99),

suggesting that we are not overlooking any other major aerodynamic or ecological factor

driving variation in wing-bone length.

Previous studies have shown that thermoregulatory demands can affect

morphological traits functioning in food acquisition (34), with consequences for individual

fitness (35). However, these studies are inconclusive regarding the relative roles of heat

dissipation versus heat retention. Our analyses extend these previous findings by

confirming that heat dissipation is a key mechanism driving phenotypic responses to

warming temperatures, even altering the design of anatomical features fundamental to

locomotion, including flight. In the case of wing morphology, the extent to which higher

temperatures can lengthen wing bones is limited by biophysical constraints on

organismal performance. Further analyses are needed to understand how this trade-off

between thermoregulation and flight performance will impact the capacity of birds and

other volant animals to respond to global warming via morphological adaptations.

**Materials and Methods**

*Morphological and life history data*

We estimated the length of the humerus and ulna for 7,366 individual birds of 1,520 species using skeletal specimens in the University of Michigan Museum of Zoology (UMMZ). Each specimen was photographed from a consistent distance and then measurements of the ulna and humerus were generated from the photographs with a neural network-based computer vision approach (19). In brief, for this method, training data were generated by hand annotating a subset of the images of specimens using the VGG Image Annotator (36) and the Toronto Annotation Suite (37), and then precise pixel-wise segmentations for bones of interest were annotated using AI-assisted annotation tools in the Toronto Annotation Suite (37). These partial annotations (pixel segmentations were only provided for the bones of interest) were then extended to fully annotated specimens using machine learning. Specifically, a U-Net (38) was trained to determine if each pixel was a bone or not based on preliminary annotations as well as additional synthesized samples. The U-Net was used to infer labels of all pixels in the annotated images. This annotated dataset was then used to train an instance segmentation neural network [Mask-RCNN (39)] to segment the bones. From these segmentations, the length of each bone of interest was estimated as the length of the longest diagonal. The metric size of the bone was determined using its known distance from the camera on a flat surface. For complete details, and methodological validation, see (19). We photographed all of the passerine specimens at the UMMZ, but filtered our data to only include estimates for bones that the network detected with high confidence, which we define as a detection where the most likely bone class has at least a 95% probability (19). For each specimen, if multiple measurements of the humerus or ulna were available (i.e., if the model identified and measured the bones from both wings), the arithmetic mean of the measurements was used.

Estimates of the relaxed wing chord (the distance from the carpal joint to the tip of the unflattened longest primary), which we refer to as primary flight feather length, and the distance from the carpal joint to the longest secondary flight feather were recently measured for all birds (20) and are publicly available for all of our species. Migratory status data are publicly available (20), with each species classified as 1 = non-migratory, 2 = partial/intermediate migrant, or 3 = long distance migrant. Species mean mass data

and the latitude of the midpoint of the breeding range centroid were also obtained from
(20).

*Environmental data*

Temperature and precipitation data were generated by overlapping species breeding
and resident ranges obtained from Birdlife International with a 2.5-minute resolution
gridded map of the mean temperature of the warmest quarter and mean annual
precipitation averaged across 1970-2000 using WorldClim v.2.1 (40), and calculating the
mean values of these variables within each species' range. All models were also refit
using the 1970-2000 annual mean temperature and the maximum temperature of the
warmest month (40) (SI). Species were categorized as occupying 1 of 10 ecological
niches: vertivore, invertivore, scavenger, omnivore, aquatic predator, frugivore, herbivore
(aquatic), herbivore (terrestrial), nectarivore or granivore, with each species assigned to
a niche if the relevant food source accounts for at least 60% of the species' diet, based
on (20). We unified our taxonomy to the Birdtree taxonomy (41) using the crosswalk in
(20); only species with 1:1 correspondence (i.e. that were not differentially lumped or
split by the UMMZ and Birdtree taxonomy) were included.

*Flight aerodynamics*

For each species, we estimated an index of maneuverability and gliding flight efficiency.
Flight maneuverability was quantified using a 'pitch index' that is a modified version of a
pitch agility metric (21), and quantifies the capacity of a bird to generate angular
acceleration. To calculate the pitch index, we quantified wing area (*S*) and a static
margin index (*SMI*) as:

$$S = L_s(H + U) \ + \ 0.2\pi L_s \qquad \text{eqn 1}$$

$$SMI = \frac{0.25\ L_P}{M^{1/3}} \qquad \text{eqn 2}$$

Where $L_s$ is the length from the carpal joint to the tip of the first secondary feather, *H* is
the length of the humerus bone, *U* is the length of the ulna bone, $L_p$ is the length from
the carpal joint to the tip of the longest primary feather (referred to as the relaxed wing
chord) and *M* is mass. We then used these variables to calculate the Pitch Index:

$$Pitch\ Index = \frac{(SMI)M^{0.24}S}{M^{5/3}} \qquad \text{eqn 3}$$

Birds with higher pitch indices have a greater ability to generate angular

acceleration around the pitch axis (Figs. S1-S2). To estimate gliding flight efficiency, we

calculated an index of the aspect ratio of the wings as:

$$Aspect\ Ratio\ Index = \frac{(L_P+U+H)^2}{S}$$        eqn 4

We provide derivations of these equations, empirical validation, and discussion of the

necessary simplifying assumptions below.

*Statistical approach*

In our models, we used species-means for all variables. Wing-bone length (the sum of

the humerus length and length of the ulna) was modeled in a phylogenetic linear mixed

model as a function of the length of primary feather length (i.e. the relaxed wing chord),

mean temperature of the warmest quarter, the latitude at the center of the range, pitch

index, aspect ratio, migration status, and ecological niche, with species identity

incorporated as a random effect that accounts for phylogenetic relatedness. Mass, wing

area (eqn 1), and wing loading (mass/wing area) were not included as predictors, as

they were highly correlated with length of the relaxed wing chord ($R$ = 0.88, 0.95, and

0.997, respectively); none of the remaining variables were highly correlated, and

particularly importantly, temperature and latitude had a relatively low correlation in our

data ($R$ = -0.64)*.* All continuous variables were scaled to have a mean of 0 and a

standard deviation of 1 prior to model fitting; migration was included as categorical

variable, with 1 being non-migratory and 3 being long distance migrants. We also

constructed analogous models, controlling for log-transformed mass rather than relaxed

wing chord to understand the relationship between the variables and wing-bone length

relative to body size (Table S4)*.*

Models were fit using the 'MCMCglmm' function in the MCMCglmm package (42)

in R (43), with uninformative inverse Wishart priors (V = 1; nu = 0.02) for the residual

and phylogenetic variance. To account for phylogenetic relatedness and uncertainty in

those relationships, we randomly sampled 100 trees from the posterior distribution of a

comprehensive phylogeny of birds (41) constructed on the Hackett et al. (44) backbone

tree, and including species that were placed using taxonomy. Using a random tree from

the posterior, we generated starting parameter estimates by running the models for

55,000 iterations, discarding the first 5,000 iterations as burn-in and with a thinning rate

of 10. Using these starting parameter estimates, we then ran the models for 55,000

iterations, with 5,000 iterations discarded as burn-in and a thinning rate of 1,000 for each

of the 100 trees. The parameter estimates were derived by combining the posterior

distributions based on each of the 100 trees. To test that our results were robust to our

treatment of phylogenetic uncertainty, we also fit an analogous model, but used a

maximum clade credibility consensus phylogeny to capture phylogenetic relatedness

and ran the model for 500,000 iterations, with the first 50,000 discarded as burn-in and a

thinning factor of 100. We confirmed that the models had converged by examining

parameter trace plots and posterior distributions.

**Acknowledgments**

We thank Benjamin M. Winger and Brett Benz for access to skeletal specimens at the

University of Michigan Museum of Zoology. This work was supported by the David and

Lucille Packard Foundation.

**References**

1.  C. Bergmann, Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer Grösse. *Göttinger Studien* **1**, 595–708 (1847).

2.  J. A. Allen, The influence of physical conditions in the genesis of species. *Radical Review* **1**, 108–140 (1887).

3.  S. Meiri, T. Dayan, On the validity of Bergmann's rule. *J Biogeogr* **30**, 331–351 (2003).

4.  M. D. Campbell, *et al.*, Testing Bergmann's rule in marine copepods. *Ecography* **44**, 1283–1295 (2021).

5.  M. R. E. Symonds, G. J. Tattersall, Geographical variation in bill size across bird species provides evidence for Allen's Rule. *Am Nat* **176**, 188–197 (2010).

6.  A. Fröhlich, D. Kotowska, R. Martyka, M. R. E. Symonds, Allometry reveals trade-offs between Bergmann's and Allen's rules, and different avian adaptive strategies for thermoregulation. *Nature Communications* **14**, 1–13 (2023).

7.  P. T. Katzmarzyk, W. R. Leonard, Climatic influences on human body size and proportions: Ecological adaptations and secular trends. *Am J Phys Anthropol* **106**, 483–503 (1998).

8.   G. J. Tattersall, B. Arnaout, M. R. E. Symonds, The evolution of the avian
bill as a thermoregulatory organ. *Biological Reviews* **92**, 1630–1656 (2017).

9.   J. R. Speakman, E. Król, Maximal heat dissipation capacity and
hyperthermia risk: neglected key factors in the ecology of endotherms.
*Journal of Animal Ecology* **79**, 726–746 (2010).

10.  L. Tian, M. J. Benton, Predicting biotic responses to future climate warming
with classic ecogeographic rules. *Current Biology* **30**, 737–758 (2020).

11.  S. Ryding, M. Klaassen, G. J. Tattersall, J. L. Gardner, M. R. E. Symonds,
Shape-shifting: changing animal morphologies as a response to climatic
warming. *Trends Ecol Evol* **36**, 1036–1048 (2021).

12.  R. Greenberg, R. Danner, B. Olsen, D. Luther, High summer temperature
explains bill size variation in salt marsh sparrows. *Ecography* **35**, 146–152
(2012).

13.  N. R. Friedman, L. Harmáčková, E. P. Economo, V. Remeš, Smaller beaks
for colder winters: thermoregulation drives beak size evolution in
Australasian songbirds. *Evolution* **71**, 2120–2129 (2017).

14.  K. Schmidt-Nielsen, Locomotion: Energy cost of swimming, flying, and
running. *Science* **177**, 222–228 (1972).

15.  U. M. Norberg, *Vertebrate flight. Mechanics, Physiology, Morphology,*
*Ecology and Evolution.* (Springer-Verlag, 1990).

16.  A. Craig, J. Larochelle, The cooling power of pigeon wings. *Journal of*
*Experimental Biology* **155**, 193–202 (1991).

17.  S. Ward, *et al.*, Heat transfer from starlings *Sturnus vulgaris* during flight.
*Journal of Experimental Biology* **202**, 1589–1602 (1999).

18.  S. Rogalla, M. D. Shawkey, B. Vanthournout, L. D'Alba, Thermoregulation
and heat exchange in ospreys (*Pandion haliaetus*). *J Therm Biol* **96** (2021).

19.  B. C. Weeks, *et al.*, A deep neural network for high throughput
measurement of functional traits on museum skeletal specimens. *Methods*
*Ecol Evol* **14**, 347–359 (2023).

20.  J. A. Tobias, *et al.*, AVONET: morphological, ecological and geographical
data for all birds. *Ecol Lett* **25**, 581–597 (2022).

21.  C. Harvey, V. B. Baliga, J. C. M. Wong, D. L. Altshuler, D. J. Inman, Birds
can transition between stable and unstable states via wing morphing.
*Nature* **603**, 648–653 (2022).

22.  C. J. Pennycuick, *Modeling the flying bird* (Academic Press, 2008).

23.  C. Harvey, D. J. Inman, Aerodynamic efficiency of gliding birds vs
comparable UAVs: a review. *Bioinspir Biomim* **16**, 031001 (2021).

24.  C. Sheard, *et al.*, Ecological drivers of global gradients in avian dispersal
inferred from wing morphology. *Nat Commun* **11**, 1–9 (2020).

25.  B. M. Winger, G. G. Auteri, T. M. Pegan, B. C. Weeks, A long winger for
the Red Queen: rethinking the evolution of seasonal migration. *Biological*
*Reviews* **94**, 737–752 (2019).

26.  J. Belmaker, C. H. Sekercioglu, W. Jetz, Global patterns of specialization
and coexistence in bird assemblages. *J Biogeogr* **39**, 193–203 (2012).

27.  G. G. Mittelbach, *et al.*, Evolution and the latitudinal diversity gradient:
Speciation, extinction and biogeography. *Ecol Lett* **10**, 315–331 (2007).

28.  K. Marchetti, T. Price, A. Richman, Correlates of Wing Morphology with
Foraging Behaviour and Migration Distance in the Genus Phylloscopus. *J*
*Avian Biol* **26**, 177–181 (1995).

29.  A. Hedenström, L. Christoffer Johansson, Bat flight: Aerodynamics,
kinematics and flight morphology. *Journal of Experimental Biology* **218**,
653–663 (2015).

30.  V. Jirinec, *et al.*, Morphological consequences of climate change for
resident birds in intact Amazonian rainforest. *Sci Adv* **7**, eabk1743 (2021).

31.  B. C. Weeks, *et al.*, Shared morphological consequences of global warming
in North American migratory birds. *Ecol Lett* **23**, 316–325 (2020).

32.  T. N. Sullivan, M. A. Meyers, E. Arzt, Scaling of bird wings and feathers for
efficient flight. *Sci Adv* **5**, eaat4269 (2019).

33.  X. Wang, A. J. McGowan, G. J. Dyke, Avian wing proportions and flight
styles: first step towards predicting the flight modes of mesozoic birds.
*PLoS One* **6**, e28672 (2011).

34.  G. J. Tattersall, D. V. Andrade, A. S. Abe, Heat exchange from the toucan
bill reveals a controllable vascular thermal radiator. *Science* **325**, 468–470
(2009).

35.  J. A. Van Gils, *et al.*, Body shrinkage due to Arctic warming reduces red
knot fitness in tropical wintering range. *Science* **352**, 819–821 (2016).

36.  A. Dutta, A. Gupta, A. Zisserman, VGG image annotator (VIA) (2016)
http://robotics.ox.ac.uk/~vgg/software/via/.

37.  A. Kar, *et al.*, Toronto Annotation Suite (2021)
https://aidemos.cs.toronto.edu/annotation-suite/.

38.  O. Ronneberger, P. Fischer, T. Brox, U-net: Convolutional networks for
biomedical image segmentation in *International Conference on Medical*
*Image Computing and Computer-Assisted Intervention*, (Springer, Cham,
2015), pp. 234–241.

39.     K. He, G. Gkioxari, P. Dollár, R. Girshick, Mask R-CNN. *IEEE Trans*
421                 *Pattern Anal Mach Intell* (2020)
422                 https:/doi.org/10.1109/TPAMI.2018.2844175.

423     40.     S. E. Fick, R. J. Hijmans, WorldClim2: new 1km spatial resolution climate
424                 surfaces for global land areas. *International Journal of Climatology* **37**,
425                 4302–4315 (2017).

426     41.     W. Jetz, G. H. Thomas, K. Hartmann, A. O. Mooers, The global diversity of
427                 birds in space and time. *Nature* **491**, 444–448 (2012).

428     42.     J. D. Hadfield, MCMC methods for multi-response generalized linear mixed
429                 models: the MCMCglmm R package. *J Stat Softw* **33**, 1–22 (2010).

430     43.     R Core Team, R: A language and environment for statistical computing
431                 (2018) https://www.r-project.org/.

432     44.     S. J. Hackett, *et al.*, A phylogenomic study of birds reveals their
433                 evolutionary history. *Science* **320**, 1763–1768 (2008).

**Figure 1. Geographic and phylogenetic sampling of avian wing-bone morphology.**

Using measurements from 7,366 skeletal specimens, we analyze variation in wing-bone

length across 1,520 bird species, representing ~33% of passerine diversity. (**A**) Blue

bars at branch tips show the number of specimens measured per species. (**B**) Red dots

show the geographic distribution of specimen localities. Images reproduced with

permission from Cornell Lab of Ornithology.

**Figure 2. Higher temperatures are associated with increased length of avian wing-**

**bones.** (**A**) Temperature during the warmest quarter of the year is significantly positively

associated with wing-bone length after controlling for the length of the primary flight

feathers and a range of ecological and geographic variables (*Methods*). Dots show

means of the posterior distribution with bars indicating the 95% confidence interval for

each estimate; those not overlapping zero are shown in black and those overlapping

zero are shown in grey. (**B**) The vascularized section of wings is relatively large in

warmer climates**.** The proportion of the wing that is vascularized (the sum of humerus

and ulna length divided by the length of the primary flight feathers) is higher for species

in warmer areas. Dots represent species averages for each of the 1,502 species used in

our models; a linear model of the bivariate relationship between temperature and the

proportional length of the wing bones is shown.