Nuclear genetic background influences the phenotype of the *Drosophila tko^{25t}* mitochondrial protein-synthesis mutant

Howard T. Jacobs[1,2,#], Tea Tuomela[1] & Päivi Lillsunde[1]

[1]Faculty of Medicine and Health Technology, FI-33014 Tampere University, Finland; [2]Department of Environment and Genetics, La Trobe University, Melbourne, Victoria 3086, Australia

[§]Corresponding author: Howard T. Jacobs, Faculty of Medicine and Health Technology, FI-33014 Tampere University, Finland. Phone: +358503412894, E-mail: howard.jacobs@tuni.fi, howy@jacobs.net

Running title: Nuclear background effects on tko^{25t}

ABSTRACT

The *Drosophila tko²⁵ᵗ* point mutation in the gene encoding mitoribosomal protein S12 produces a complex phenotype of multiple respiratory chain deficiency, developmental delay, bang-sensitivity, impaired hearing, sugar and antibiotic sensitivity and impaired male courtship. Its phenotypic severity was previously shown to be alleviated by inbreeding, and to vary with mitochondrial genetic background. Here we show similarly profound effects conferred by nuclear genetic background. We backcrossed *tko²⁵ᵗ* into each of two standard nuclear backgrounds, Oregon R and *w¹¹¹⁸*, the latter used as recipient line in many transgenic applications requiring selection for the *white* minigene marker. In the *w¹¹¹⁸* background, *tko²⁵ᵗ* flies showed a moderate developmental delay and modest bang-sensitivity. In the Oregon R background, males showed longer developmental delay and more severe bang-sensitivity, and we were initially unable to produce homozygous *tko²⁵ᵗ* females in sufficient numbers to conduct a meaningful analysis. When maintained as a balanced stock over 2 years, *tko²⁵ᵗ* flies in the Oregon R background showed clear phenotypic improvement though were still more severely affected than in the *w¹¹¹⁸* background. Phenotypic severity did not correlate with the expression level of the *tko* gene. Analysis of tko25t hybrids between the two backgrounds indicated that phenotypic severity was conferred by autosomal, X-chromosomal and parent-of-origin dependent determinants. Although some of these effects may be *tko²⁵ᵗ*-specific, we recommend that, in order to minimize genetic drift and confounding background effects, the genetic background of non-lethal mutants should be controlled by regular backcrossing, even if stocks are usually maintained over a balancer chromosome.

INTRODUCTION

The *Drosophila* bang-sensitive mutant tko^{25t}, harboring a point mutation in the gene encoding mitoribosomal protein S12 (Royden et al. 1987; Shah et al. 1997), was originally classified as semilethal (Judd et al. 1972). It was subsequently maintained by ourselves and others, both as an inbred stock and over an FM7 balancer chromosome. The major features of the organismal phenotype, in addition to bang sensitivity, are delayed larval development, deafness (non-responsiveness to sound), impaired male courtship and hypersensitivity to antibiotics targeting the mitochondrial translation system (Toivonen et al. 2001). In an earlier study we reported that the inbred, laboratory-maintained tko^{25t} stock manifested a milder phenotype than originally described, but that a more severe phenotype could be restored by backcrossing the mutation into two different standard nuclear backgrounds (Canton S and Oregon R) to create a hybrid. Subsequently we have explored various genetic and environmental manipulations that further modified the tko^{25t} phenotype. Individual lines established from the original tko^{25t} Canton S/Oregon R hybrid, and selected at each generation for fastest development, showed varying degrees of suppression of the mutant phenotype. The three most pronounced suppressors were found to harbor segmental duplications of the region of the X chromosome encompassing the mutated *tko* gene (Kemppainen et al. 2009).

Mitochondrial genetic background was also shown to be a key determinant of tko^{25t} phenotype: in some mtDNA backgrounds, notably those previously infected with the endosymbiont *Wolbachia*, partial suppression was found (Chen et al. 2012), whilst another mtDNA haplotype isolated from the wild was synthetically lethal with tko^{25t} (Salminen et al. 2019). Here we attempted a similar analysis, but of the effect of nuclear background, keeping mtDNA haplotype and the tko^{25t} mutation

itself unchanged. Without imposing any kind of deliberate selection, we obtained divergent

outcomes, pointing to the genetic complexity of nuclear-mitochondrial interactions in a metazoan.

MATERIALS AND METHODS

Drosophila culture and crosses

Flies were cultured in standard high-sugar medium (Fernandez-Ayala et al. 2009) on a 12 h/12 h

light-dark cycle at the temperatures indicated in figures and legends (25 °C where not otherwise

stated). For long-term maintenance fly stocks were kept at 18 °C. The tko^{25t} mutation was

maintained over an FM7 balancer chromosome. tko^{25t} flies were generated for experimental analysis

by crossing tko^{25t} / FM7 females with tko^{25t} / Y males, with an initial cross of tko^{25t} / FM7 females

with FM7 / Y males to generate the latter, where needed. Eclosion timing was measured, as

previously (Kemppainen et al. 2016), using replicate sets of vials (n ≥ 3). Bang-sensitivity was

measured as recovery time after vortexing of individual flies, as previously (Toivonen et al. 2001).

Recovery time for flies that had still not recovered after 300 s was arbitrarily set to 300 s.

Single-fly genomic DNA extraction

Individual flies in a 0.5 ml tube were mashed for 5-10 s with a plastic pipette tip in 50 μl of SB (10

mM Tris-HCl, 1 mM EDTA, 25 mM NaCl, 200 μg/ml Proteinase K, pH 8.2). After incubation at

room temperature for 20-30 min the Proteinase K was inactivated by heating to 95 ºC for 1-2 min

and the mix used immediately for PCR.

Genotyping

The presence of the tko^{25t} mutation was verified by PCR amplification of a 1.49 kb fragment of the

tko gene from single-fly genomic DNA, using primer Tko-51 (5´-

GAACAAAAAACTACTGAACAAAACTCC-3´) and Tko-31 (5´-

CATTTGAACAACGTGATTAGGAAGT-3´), followed by Sanger sequencing of the agarose gel-

purified product using primers tko_TT_L1 (5´-GTGCTTTATTGATTTCGAGCGATCT-3´) and

tko_TT_R1 (5´-ACTATTGGCTCTTCTTGACGACGTG-3´).

Expression analysis by qRTPCR

Total RNA was extracted and *tko* RNA levels relative to *RpL32* were measured by quantitative

RTPCR as previously (Fernandez-Ayala et al. 2009), using primers as specified in (Kemppainen et

al. 2009).

Courtship analysis

Courtship was analyzed essentially as previously (Toivonen et al. 2001), in batches of 6 pairs of

adult male and virgin female 3 day-old flies of a given genotype. For experiments at different

temperatures, virgin flies were kept at the experiment temperature until the experiment was started.

Flies were transferred into individual mating chambers by drawing them from culture vials one by

one using a depressurized plastic bottle and releasing into the chamber. With the female and male

initially separated, mating chambers were placed in an incubator at the specified temperature. Flies

were allowed to recover from the transfer for one hour before allowed to interact with the opposite

sex and then inspected under a stereo microscope at 15-minute intervals during each hour-long

experiment to determine how many pairs copulated. Mean copulation frequency on a per batch

basis was derived from n > 5 batches of each genotype studied

Statistical analysis

Data were analyzed, where indicated, either by Student's *t* test or by one-way ANOVA using a

standard online tool (https://astatsa.com/OneWay_Anova_with_TukeyHSD/). Recovery times in the

bang-sensitivity assay were not normally distributed: therefore, raw data only are presented in the figures.

RESULTS AND DISCUSSION

Outbred tko^{25t} lines show a severely exacerbated phenotype

The tko^{25t} mutation was backcrossed over $\geq$10 generations according to the scheme outlined in Figure 1 into each of two standard nuclear backgrounds, Oregon R, used in laboratories worldwide since 1925 – see (Lindsley and Grell, 1968), and w^{1118} (Hazelrigg et al. 1984), widely used for selection of transgenic lines using the *mini-white* marker, which confers colored eyes in a white-eyed background. After back-crossing, several lines in each background were maintained over a standard X-chromosomal balancer, FM7 (Merriam, 1968). For each background, we established multiple independent back-crossed lines.

Outbred tko^{25t} lines manifest background-dependent phenotypes

All of the outbred tko^{25t} lines showed a more severe phenotype than the lab-maintained tko^{25t} stock, in terms of both developmental delay and bang-sensitivity. The tko^{25t} phenotypes in the two backgrounds were also phenotypically distinct (Fig. 2), although three independent back-crossed lines in the Oregon R background were similar to each other, as were three such lines in the w^{1118} background (Fig. 2). In initial crosses to generate tko^{25t} males from the balanced outbred stocks (see legend to Fig. 2), we observed that very few mutant males eclosed in the Oregon R background at the standard culture temperature of 25 ℃, resembling the semilethality first reported for the mutation (Fig. 2A), whereas the output of mutant males in the w^{1118} background was very close to the expected 25% (Fig. 2A). The few tko^{25t} males in the Oregon R background that eclosed were very delayed and extremely bang-sensitive, although there were too few from each line to generate

meaningful numbers. For the w^{1118} background, developmental delay at 25 ºC was around 5 days, whereas our original tko^{25t} stock showed a developmental delay of only 2-3 days (Fig. 2B). We found that more substantial numbers of tko^{25t} males in the Oregon R background eclosed at room temperature 21-22 ºC, although they were still very bang-sensitive and did not mate, making it impossible to generate tko^{25t} females in this background. We were, however, able to generate sufficient numbers of mutant males to compare developmental delay (Fig. 2C) and bang-sensitivity (Fig. 2D) in the two backgrounds.

Outbred tko^{25t} lines maintained as balanced stock show phenotypic alleviation

The tko^{25t} mutation in the outbred lines was maintained for two years over an FM7 balancer chromosome with no intentional selection. When retested, the phenotype of the lines in both backgrounds was substantially milder than in the initial tests. In the Oregon R background the mutation was no longer semilethal, and mutant males were now able to mate and give offspring. We were able to obtain sufficient numbers of mutant flies of both sexes and both backgrounds to reliably measure developmental time (Fig. 3A) and bang-sensitivity (Fig. 3B) at 25 ºC (Fig. 3). Flies outbred to the w^{1118} background still showed a milder phenotype than those in the Oregon R background, notably for bang-sensitivity (Fig. 3B), but were also less severely affected than when initially tested, immediately after the completion of the backcrosses. Developmental delay of tko^{25t} flies at 25 ºC was 3-4 d in the w^{1118} background and 4-5 d in the Oregon R background (see examples in Fig. 3), with males always more delayed than females.

tko^{25t} phenotype and tko expression are crudely correlated

In a previous study (Kemppainen et al. 2009) no strict correlation was found between the expression level of the mutant tko gene and the tko^{25t} phenotype, although duplication of the mutant gene in its natural chromosomal milieu did provide a partial phenotypic rescue (Kemppainen et al.

2009). In the present study it was clear that the phenotype of developmental delay and bang-sensitivity was more pronounced in mutant males than mutant females, in the Oregon R background compared with the w^{1118} background and in mutant flies cultured at 25 ºC than at 21-22 ºC. To re-examine the influence of *tko* expression level on phenotype we used qRTPCR to analyse *tko* RNA levels in 2-day old adults of each sex, background, genotype and culture temperature (Fig. 4). At 25 ºC the expression level of *tko* was similar in wild-type flies of each sex in the two backgrounds (Fig. 4A), being only slightly higher in Oregon R. We thus used RNA from flies of the corresponding sex and background as a control, in all subsequent assays of *tko* expression in tko^{25t} flies (Fig. 4B-4D). In all cases tested, *tko* expression was higher in tko^{25t} than control flies, and the difference was more pronounced in males, where it was greatest in the Oregon R background. These findings suggest that increased *tko* expression might be a compensatory response to (or at least a marker for) a severely deleterious phenotype. However, expression in females was almost identical in the two backgrounds, despite the phenotypic differences and was very similar at the two culture temperatures. Thus, *tko* expression level is not a simple determinant of the ko^{25t} phenotype.

Autosomal semi-dominance and parent-of-origin effects underlie strain differences in tko^{25t} phenotype

In an attempt to characterize the genetic basis of the strain background differences we crossed heterozygous tko^{25t} / FM7 females in each background with tko^{25t} males in the other, and assessed the phenotype of the first-generation progeny. Since the exact eclosion timing varies slightly between the background strains, and can also differ between experiments, we plotted the amount of developmental delay for both tko^{25t} females and males in each background, alongside the delay observed for the tko^{25t} background hybrids (Fig. 5). Female tko^{25t} hybrid progeny from tko^{25t} / FM7 mothers in the w^{1118} background were only slightly more delayed than if both parents were in the w^{1118} background (Fig. 5A). For males, the delay for hybrid progeny from these mothers was the

same as when both parents were in the w^{1118} background (Fig. 5B). Hybrid tko^{25t} progeny of both sexes showed an intermediate delay when the female parent was from the Oregon R background (Fig. 5A, 5B). These findings suggest a mixed contribution from autosomal and maternal determinants. The latter could be mitochondrial, although the mtDNA of the two lines was identical at the start of the back-cross. It may also indicate a form of parental imprinting (Lloyd, 2000; Anaka et al. 2009), although the existence of imprinting in *Drosophila* has been questioned (Coolon et al. 2012). Alternatively it could be a classic maternal effect, e.g. involving non-coding RNAs (Soleimani et al. 2020), localized (Hoch and Jäckle 1993) or pioneer transcription factors (Gaskill et al. 2021) or their mRNAs, or other regulatory proteins (Zhang et al. 2018). A simple role for the X-chromosome in mutant eclosion timing seems to be excluded, since the w^{1118} background appears to be dominant only when inherited from the mother, and in both sexes. Although clear and mostly statistically significant, the differences in delay between the crosses and sexes are too small to enable a meaningful next-generation cross to be conducted, in order to try to tease out the contributions of autosomal, mitochondrial and possible X-chromosomal inheritance.

We next examined bang-sensitivity in the progeny of these crosses (Fig. 6). Although the recovery times are not normally distributed, making standard statistical tests unreliable, it is clear from the cumulative plots for each sex that a rather similar pattern of dominance and parent-of-origin effects is seen as for developmental delay. In the hybrid crosses, the phenotype of tko^{25t} progeny from mothers of the w^{1118} background but Oregon R fathers (green data plots) is very similar to that of tko^{25t} flies of the pure w^{1118} background (red data plots). tko^{25t} females with Oregon R mothers but w^{1118} fathers (blue data plots, Fig 6A) are intermediate between the two pure lines (red *versus* black data plots), whereas tko^{25t} males with Oregon R mothers but w^{1118} fathers (blue data plots, Fig. 6B) are almost identical with tko^{25t} males of the pure OR background (blue *versus* black data plots, Fig.

6B). Thus, the data indicate an overall 'dominance' of the milder w^{1118} cytoplasm or epigenome, as well as a negative influence on recovery from mechanical shock of the Oregon R X-chromosome.

tko^{25t} courtship analysis is compromised by background effects

We attempted to analyse a third phenotypic parameter of tko^{25t} flies, namely defective male courtship (Toivonen et al. 2001). However, this did not prove meaningful for the following reasons. Immediately after the back-cross, males in the Oregon R background were extremely weak and unable to mate. Furthermore, previous investigators have found that mutation of the *white* locus itself impairs male courtship behavior, although flies do eventually mate in sufficient numbers to maintain *white*⁻ lines. Working on the assumption that the w^{1118} mutation itself was the cause of the line's courtship defect, as previous authors have inferred (Xiao et al. 2017),we created a line of Oregon R flies back-crossed into the w^{1118} background, but retaining red eyes (this was also true of the tko^{25t} back-cross into w^{1118}, since *tko* and *w* map very close together on the X-chromosome). Red-eyed tko^{25t} males in the w^{1118} background were courtship-impaired (Fig. 7) in a similar manner as inbred tko^{25t} males studied previously (Toivonen et al. 2001), but red-eyed wild-type males in the w^{1118} background were also courtship-impaired (Fig. 7), rendering the main experiment meaningless. Although seemingly at odds with the previous findings on the male-courtship effects of *white* (Xiao et al. 2017), it should be noted that the males derived from the back-cross undertaken by the previous authors gave only 6 successful copulations out of 16 trials (Xiao et al. 2017). Furthermore, these authors showed that the absence of eye color *per se* was not the cause of the w^{1118} courtship defect (Xiao et al. 2017), and that the genotype of females in the assay also influenced the copulation success of w^{1118} males. In conclusion, although tko^{25t} males in the Oregon R background were unable to mate, the less severe courtship defect in the w^{1118} background was impossible to quantify reliably. Note that, despite the male courtship defect of w^{1118} males, they are not bang-sensitive, as confirmed here (Fig. 2D).

Concluding comments

After its initial isolation in 1972, the tko^{25t} mutant has been maintained in various backgrounds in different laboratories during 50 years, and its exact history is practically impossible to reconstruct. During this time, as a result of inbreeding and inadvertent selection in many different laboratories, the phenotype has become milder (Fig. 2A). A restoration of the original semilethal phenotype was achieved by backcrossing into standard reference strain Oregon R. However, despite being maintained subsequently over a balancer chromosome with no deliberate selection, its phenotype has again become partially alleviated, during 2 years of standard laboratory maintenance. For any mutant strain with a nonlethal phenotype, especially one like tko^{25t}, where reproductive competence and developmental timing are affected, we would recommend that the strain be periodically backcrossed into the original strain background and the phenotype rechecked, before the strain is used in any experimental context.

Given that the phenotype recorded after back-crossing into two commonly used and broadly wild-type nuclear backgrounds (apart from the w mutation in w^{1118}) was so different (semilethal versus fully viable, 5-10-fold difference in median recovery times from mechanical shock, ~30% increase in developmental delay, mild male courtship defect versus inability to mate) it would be advisable to conduct such a back-cross additionally into several other reference strains. Although partial phenotypic suppression of tko^{25t} by functional absence (or heterozygosity) of $white$ cannot be excluded, it seems more likely that other background alleles not knowingly under selection are responsible for the phenotypic differences. This problem of cryptic mutations in inbred reference strains is a widespread phenomenon. The rapid generation time and use of different culture conditions in different laboratories, not to mention possible human error in stock passaging, mean that different isolates of the same reference strain are unlikely to be or to remain genetically

identical, and thus may confer different traits upon a given mutation such as tko^{25t}. The *Drosophila melanogaster* Genetic Reference Panel DmGRP (Mackay et al. 2012), a collection of outbred/inbred fly lines, is an extremely valuable tool in this regard, with a vast array of applications, both potential and realized (e.g. see; Spierer et a 2021; Havula et al.2022). It could be more widely used by the community, to identify background polymorphisms that influence the phenotypic expression of mutations such as tko^{25t}, especially given the increasing recognition of the importance and flexibility of *Drosophila* in modeling human diseases (Verheyen, 2022). The same issue plagues mouse genetics, for example, where disease-modeling mutations can have no phenotype in one background, yet are lethal in another. The creation of a reference collection of outbred/inbred mouse strains (the CC resource, equivalent to DmGRP) to model complex traits in a genetically diverse population has yielded hundreds of such lines (Churchill et al. 2004; Abu Toamih Atamni et al. 2018).

Ideally, unexpected phenotypes should be verified by back-crossing to a different isolate of the same background strain. Given our previous observations of profound effects of different mtDNA backgrounds on tko^{25t} (Chen et al. 2012; Salminen et al. 2019), we would recommend that attention should also be paid to mitochondrial genotype, in such back-crossing schemes, regardless of whether the expected phenotype is obviously 'mitochondrial'. Note that the phenotypic differences observed in our back-crossed tko^{25t} lines is not the result of genetic heterogeneity within the wild-type lines, even though this may be considerable, as inferred previously for Oregon R (Lints and Bourgois 1987). Back-crossed tko^{25t} lines within each background had relatively uniform phenotypes.

The *tko* gene product is part of a complex structure, the mitoribosome, containing dozens of other polypeptides and RNA moieties. Furthermore, the mitoribosome directly influences the functional

expression of the 13 mtDNA-encoded polypeptides and the dozens of nuclear-coded polypeptides with which they interact, both transiently and within stable complexes, to maintain essential biological processes. As such, it is hardly surprising that mutations in *tko* impose all kinds of selective pressure that can be compensated by diverse changes in genotype or gene expression. But in these respects *tko* is far from unusual. Thus, the possible influence of genetic background, of selection operating for a supposedly recessive mutation even in the heterozygous condition, and of parent-of-origin effects, should not be discounted for any mutation under study in *Drosophila* or in any model organism.

DATA AVAILABILITY

All data in this paper is freely available on request. Source data files for each numerical figure are available as spreadsheet tables.

ACKNOWLEDGMENTS

We thank Lotta Kulmala and Eveliina Teeri-Kahelin for technical assistance

FUNDING

HTJ was supported in this work by grants from the Academy of Finland (283157, 307431 and 324730).

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest regarding the contents of this article.

REFERENCES

Abu Toamih Atamni H, Nashef A, Iraqi FA. 2018. The Collaborative Cross mouse model for dissecting genetic susceptibility to infectious diseases. *Mamm Genome*. 29:471–487. doi: 10.1007/s00335-018-9768-1.

Anaka M, Lynn A, McGinn P, Lloyd VK. 2009. Genomic imprinting in *Drosophila* has properties of both mammalian and insect imprinting. *Dev Genes Evol*. 219:59–66. doi: 10.1007/s00427-008-0267-3.

Chen S, Oliveira MT, Sanz A, Kemppainen E, Fukuoh A, et al. 2012. A cytoplasmic suppressor of a nuclear mutation affecting mitochondrial functions in *Drosophila*. *Genetics*. 192:483–493. doi: 10.1534/genetics.112.143719.

Churchill GA, Airey DC, Allayee H, Angel JM, Attie AD, et al. 2004. The Collaborative Cross, a community resource for the genetic analysis of complex traits. *Nat Genet*. 36:1133–1137. doi: 10.1038/ng1104-1133.

Coolon JD, Stevenson KR, McManus CJ, Graveley BR, Wittkopp PJ. 2012. Genomic imprinting absent in *Drosophila melanogaster* adult females. *Cell Rep*. 2:69–75. doi: 10.1016/j.celrep.2012.06.013.

Fernandez-Ayala DJ, Sanz A, Vartiainen S, Kemppainen KK, Babusiak M, et al. 2009 Expression of the *Ciona intestinalis* alternative oxidase (AOX) in *Drosophila* complements defects in mitochondrial oxidative phosphorylation. *Cell Metab*. 9:449–460. doi: 10.1016/j.cmet.2009.03.004.

Gaskill MM, Gibson TJ, Larson ED, Harrison MM. 2021.GAF is essential for zygotic genome activation and chromatin accessibility in the early *Drosophila* embryo. *Elife*. 10:e66668. doi: 10.7554/eLife.66668.

George J, Tuomela T, Kemppainen E, Nurminen A, Braun S, et al. 2019. Mitochondrial dysfunction generates a growth-restraining signal linked to pyruvate in *Drosophila* larvae. *Fly (Austin)*. 13:12–28. doi: 10.1080/19336934.2019.1662266

Havula E, Ghazanfar S, Lamichane N, Francis D, Hasygar K, et al. 2022 Genetic variation of macronutrient tolerance in *Drosophila melanogaste*r. *Nat Commun.* 13:1637. doi: 10.1038/s41467-022-29183-x.

Hazelrigg TI, Lewis R, Rubin GM. 1984. Transformation of *white* locus DNA in *Drosophila*: dosage compensation, zeste interaction and position effects. *Cell.* 36:469–481.

Hoch M, Jäckle H. 1993. Transcriptional regulation and spatial patterning in *Drosophila*. *Curr Opin Genet Dev*. 3:566–573. doi: 10.1016/0959-437x(93)90092-4.

Judd BH, Shen MW, Kaufman TC. 1972. The anatomy and function of a segment of the X chromosome of *Drosophila melanogaster*. *Genetics*. 71:139–156. doi: 10.1093/genetics/71.1.139.

Kemppainen E, Fernández-Ayala DJ, Galbraith LC, O'Dell KM, Jacobs HT. 2009. Phenotypic suppression of the *Drosophila* mitochondrial disease-like mutant *tko*(25t) by duplication of the mutant gene in its natural chromosomal context. *Mitochondrion*. 9:353–363. doi: 10.1016/j.mito.2009.07.002.

Kemppainen E, George J, Garipler G, Tuomela T, Kiviranta E, et al. 2016. Mitochondrial dysfunction plus high-sugar diet provokes a metabolic crisis that inhibits growth. *PLoS One*. 11:e0145836. doi: 10.1371/journal.pone.0145836.

Lindsley DL, Grell EH. 1968. Genetic variations of *Drosophila melanogaster*. *Publs Carnegie Instn* 627: 469pp.

Lints F, Bourgois M. 1987. Phenotypic and genotypic differentiation in cage populations of *Drosophila melanogaster*. I. Duration of development, thorax size and weight. *Genet Sel Evol*. (1983). 19:155–170. doi: 10.1186/1297-9686-19-2-155.

Lloyd V. 2000. Parental imprinting in *Drosophila*. *Genetica*. 109:35–44. doi: 10.1023/a:1026592318341.

Mackay TF, Richards S, Stone EA, Barbadilla A, Ayroles J, et al. 2012. The *Drosophila melanogaster* Genetic Reference Panel. *Nature*. 482:173–178. doi: 10.1038/nature10811.

Merriam J. 1968. FM7: first multiple seven. *Drosophila Information Service*. 43:64.

Royden CS, Pirrotta V, Jan LY. 1987. The *tko* locus, site of a behavioral mutation in *D. melanogaster*, codes for a protein homologous to prokaryotic ribosomal protein S12. *Cell*. 51:165–173. doi: 10.1016/0092-8674(87)90144-9.

Salminen TS, Cannino G, Oliveira MT, Lillsunde P, Jacobs HT, et al. 2019. Lethal interaction of nuclear and mitochondrial genotypes in *Drosophila melanogaster*. *G3 (Bethesda)*. 9:2225–2234. doi: 10.1534/g3.119.400315.

Salminen TS, Oliveira MT, Cannino G, Lillsunde P, Jacobs HT, et al. 2017. Mitochondrial genotype modulates mtDNA copy number and organismal phenotype in *Drosophila*. *Mitochondrion*. 34:75–83. doi: 10.1016/j.mito.2017.02.001.

Shah ZH, O'Dell KM, Miller SC, An X, Jacobs HT. 1997. Metazoan nuclear genes for mitoribosomal protein S12. *Gene*. 204:55–62. doi: 10.1016/s0378-1119(97)00521-0.

Soleimani S, Valizadeh Arshad Z, Moradi S, Ahmadi A, et al. 2020. Small regulatory noncoding RNAs in *Drosophila melanogaster*: biogenesis and biological functions. *Brief Funct Genomics*. 19:309–323. doi: 10.1093/bfgp/elaa005.

Spierer AN, Mossman JA, Smith SP, Crawford L, Ramachandran S, et al. 2021. Natural variation in the regulation of neurodevelopmental genes modifies flight performance in *Drosophila*. *PLoS Genet*. 17:e1008887. doi: 10.1371/journal.pgen.1008887.

Toivonen JM, O'Dell KM, Petit N, Irvine SC, Knight GK, et al. 2001. *Technical knockout*, a *Drosophila* model of mitochondrial deafness. *Genetics*. 159:241–254. doi: 10.1093/genetics/159.1.241.

Verheyen EM. 2022. The power of *Drosophila* in modeling human disease mechanisms. *Dis Model Mech*. 15:dmm049549. doi: 10.1242/dmm.049549.

Xiao C, Qiu S, Robertson RM. 2017. The *white* gene controls copulation success in *Drosophila melanogaster*. *Sci Rep*. 7:7712. doi: 10.1038/s41598-017-08155-y.

Zhang Z, Krauchunas AR, Huang S, Wolfner MF. 2018. Maternal proteins that are phosphoregulated upon egg activation include crucial factors for oogenesis, egg activation and embryogenesis in *Drosophila melanogaster*. *G3 (Bethesda)*. 8:3005–3018. doi: 10.1534/g3.118.200578.

LEGENDS TO FIGURES

Figure 1

Backcrossing scheme for *tko^{25t}*

At each backcross generation, heterozygous (*tko^{25t}* / +) females were identified by single-organism genomic DNA extraction, PCR and sequencing, after pre-mating. Progeny from wild-type females was discarded. Several parallel lines were established in each backcross, both to the *w^{1118}* and to the Oregon R backgrounds. The FM7 -balanced, 'autosomally backcrossed' lines were established in parallel.

Figure 2

Phenotype of *tko^{25t}* in different nuclear backgrounds

(A) Proportion of *tko^{25t}* males amongst eclosing progeny from crosses of the type *tko^{25t}* / FM7 x FM7 / Y, conducted at 25 ºC, after back-crossing of *tko^{25t}* into the Oregon R background (OR bkd), to create lines OR1, OR3 and OR5 as indicated, or into the *w^{1118}* background, to create lines w3, w4 and w5. The dotted line indicates the expected proportion of *tko^{25t}* / Y progeny (25%). (B) Eclosion times (means $\pm$ SD) for *tko^{25t}* flies generated in multiple crosses using the original *tko^{25t}* stock. In the three experiments shown (orig 1, orig 2, orig 3), mothers were *tko^{25t}* / FM7 and carrying different autosomal balancers (respectively, CyO, TM3-Sb and TM3-Ser) with the scored *tko^{25t}* progeny free of all balancer markers, some of which are known to interact with *tko^{25t}* (see Fig. S2 of George et al. 2019, which presents some of the same data). All controls are *tko^{25t}* / FM7 heterozygotes in the same background, which, in both back-cross backgrounds, gave indistinguishable eclosion times as wild-type males in the same background (Fig. S1A). (C) Eclosion times (means $\pm$ SD) for similar crosses as in (A), but carried out at room temperature (21-22 ºC). (D) Bang-sensitivity assay (recovery times) for progeny flies of the indicated genotypes from the crosses in (C), assayed at 22

ºC. Note that not all data points are visible, due to overlaps and coincident values. Box plots show 25th and 75th percentiles of each distribution and medians (bold lines).

Figure 3

Phenotype after 2 years in culture of *tko²⁵ᵗ* backcrossed into different nuclear backgrounds

(A) Eclosion times (means $\pm$ SD) and (B) bang-sensitivity recovery times (box-plot nomenclature as in Fig. 2) for *tko²⁵ᵗ* flies and controls of the indicated back-crossed lines, after 2 years in culture as balanced stocks. Crosses of the type *tko²⁵ᵗ* / FM7 x *tko²⁵ᵗ* / Y were conducted at 25 ºC, with controls being *tko²⁵ᵗ* / FM7 heterozygotes. In (B) the same data for *tko²⁵ᵗ* flies are shown at two different scales (note that, for clarity, the individual data points for Oregon R background are omitted in the right-hand panel, since they extend beyond the scale used). Note again that not all data points are visible, due to overlaps and coincident values. Control flies (see Fig. 2D) were not bang-sensitive.

Figure 4

Expression level of *tko* in *tko²⁵ᵗ* flies backcrossed into two nuclear backgrounds

tko RNA levels by qRTPCR (means $\pm$ SD), for flies of the indicated sex, background and culture temperature, normalized (A) to values for one isolate of Oregon R wild-type females and (B) to mean values for control flies of the same sex and background (Oregon R wild-type or *w¹¹¹⁸*, as appropriate). *, **, *** denote significantly different data classes (pairwise comparisons between genotypes, same sex and culture temperature, by Student's *t* test, $p < 0.05$, 0.01 or 0.001, respectively). For (A), despite experiment-to-experiment variation, *tko* expression was generally lower in the *w¹¹¹⁸* background in both sexes and at all temperatures tested, as in the example shown.

Figure 5

Eclosion timing in *tko²⁵ᵗ* strain-background crosses

Developmental delay (days, d, means $\pm$ SD) at 25 ºC of (A) female and (B) male *tko²⁵ᵗ* progeny from the indicated crosses between *tko²⁵ᵗ* / FM7 females and *tko²⁵ᵗ* / Y males in the backgrounds as shown (OR – Oregon R, *w* – *w¹¹¹⁸*). Dotted lines indicate values for the non-hybrid crosses, for ease of comparison. Data from (A) and (B) are from the same crosses, but with the sexes shown separately, for clarity. Control flies were *tko²⁵ᵗ* / FM7 heterozygous females, which show an identical eclosion timing as wild-type males (see Fig. S1A), with only the SD shown (means were all set to zero, so as to exclude minor, strain-specific differences in eclosion timing from the analysis).

Figure 6

Bang-sensitivity in *tko²⁵ᵗ* strain-background crosses

Recovery times (plotted cumulatively, over time) at 25 ºC for (A) female and (B) male *tko²⁵ᵗ* progeny from the indicated crosses between *tko²⁵ᵗ* / FM7 females and *tko²⁵ᵗ* / Y males of the indicated parental strain backgrounds (OR – Oregon R background, *w* – *w¹¹¹⁸* background). Control progeny from all crosses were not bang-sensitive. $n \geq 60$ females or 53 males of each genotype.

Figure 7

Male courtship effects of *tko²⁵ᵗ* and *w¹¹¹⁸*

Percentage of successful copulation (mean $\pm$ SEM) of multiple samplings of 6 pairs of flies, at the indicated temperatures. In all assays, male and female flies were of the same genotype and background, and all had red eye color. n – number of batches analyzed for each genotype as indicated. Note that, to exclude the possibility of contamination, the *tko* locus from the red-eyed control flies back-crossed into the *w¹¹¹⁸* background was sequenced and confirmed to be *tko⁺*.

LEGENDS TO SUPPLEMENTARY FIGURE

Figure S1

Supplementary data on eclosion timing of backcrossed strains

(A) Eclosion times (means $\pm$ SD) for control flies from the original tko^{25t} stock and w^{1118} and

Oregon R backgrounds, as indicated. Dashed vertical lines demarcate separate experiments. Despite

minor variations from experiment to experiment, the eclosion times for wild-type males were the

same as for tko^{25t} / FM7 heterozygous females of the given background. (B) Eclosion times (means

$\pm$ SD) for flies of the indicated genotypes, cultured at 18 °C, from the Oregon R (OR) background.

Experimental cross: tko^{25t} / FM7 x FM7 / Y. Note that the FM7 balancer itself confers a modest

developmental delay (FM7 female homozygotes not shown, for clarity). Control cross conducted in

parallel. % figures above the bars indicate the % of progeny of that phenotype class (expected

frequency = 25%). (C) Eclosion times (means $\pm$ SD) for male flies of the indicated genotypes,

cultured at 18 °C, from the two backgrounds, Oregon R (OR) and w^{1118}. Data for OR reproduced

from (B).

Original stock

tko^{25t} / FM7   x   + / Y   (or w^{1118} / Y)

Individual pre-mated female genotyped

tko^{25t} / +   x   + / Y

8 further repeats

tko^{25t} / +   x   + / Y

Balanced line 'autosomally' backcrossed in parallel

tko^{25t} / +   x   FM7 / Y

same

Creation of balanced, backcrossed stock

tko^{25t} / FM7   x   FM7 / Y

tko^{25t} / tko^{25t} tko^{25t} / Y

Backcrossed tko^{25t} flies for experimental analysis

Jacobs et al, Figure 1

Jacobs et al, Figure 2

Jacobs et al, Figure 3

Jacobs et al, Figure 4

A

B

parental strain backgrounds

♀   ♂

- OR x OR
- OR x *w*
- *w* x OR
- *w* x *w*

Jacobs et al, Figure 6

genotype:

Oregon R   *tko*[25t] *w*[1118] bkd   *w*[+] *w*[1118] bkd

% copulation

temperature (°C)   25 25 29   25 29   25
n =   5   5   9   10 10   9

Jacobs et al, Figure 7

A

25 °C

FM7 / *tko²⁵ᵗ* ♀ (white bars)
+ / Y ♂ (black bars)

B

18 °C

- □ *tko²⁵ᵗ* / FM7 (OR)
- ■ FM7 / Y
- 🟥 *tko²⁵ᵗ* / Y (OR)
- 🟧 *tko²⁵ᵗ* / Y (*w¹¹¹⁸*)
- 🟨 + / Y (OR)
- ⬜ + / Y (*w¹¹¹⁸*)

C

18 °C

from panel B

Jacobs et al, Figure S1