Inhaled CpG increases survival and synergizes with checkpoint inhibition in lymphangioleiomyomatosis

Mayowa,M, Amosu[1], Jacob, C, McCright[1], Bennett, E, Yang[1], Juan, Grano de Oro Fernandez[1], Kaitlyn, A, Moore[1], Havish, S, Gadde[1], Michele, L, Kaluzienski[1], Katharina, Maisel[1]*

[1]Fischell Department of Bioengineering, University of Maryland, College Park, MD 20742

*Correspondence to: Dr. Katharina Maisel, Fischell Department of Bioengineering, University of Maryland, College Park, MD 20742, United States, E-mail: maiselka@umd.edu

Running title: CpG as novel LAM immunotherapy

Contributions: MA led all major experiments and data analysis; JM, BY, JF, KAM, HG, and MK assisted in experiments and experimental analysis. KM supervised experimental planning, data analysis and manuscript drafting. MA, BY, JF, and KM drafted, edited, and finalized the manuscript.

Support: Funding was received from the Fischell Department of Bioengineering (KM), T32 GM080201 (MA), Vogel Endowed Summer Fellowship (MA), LAM Foundation (KM), and American Lung Association (KM). We would also like to acknowledge the BioWorkshop and the University of Maryland School of Medicine's Pathology Histology Core for aid in sample preparation, data collection and experimental analysis.

Descriptor: 8.19 LAM: Fundamental Mechanisms

This article has an online data supplement, which is accessible from this issue's table of content online at **www.atsjournals.org**

Manuscript Word Count: 3500

ABSTRACT

Rationale: Lymphangioleiomyomatosis (LAM) is a devastating disease primarily found in women of reproductive age that leads to cancer-like cystic destruction of the lungs. Recent work has shown that LAM causes immunosuppression and that checkpoint inhibitors can be used as LAM treatment. IN lung cancer, TLR agonist, in particular TLR9 agonist CpG has been shown to be effective.

Objectives: Here we investigate the use of TLR9 agonist CpG as LAM immunotherapy in combination with checkpoint inhibitor, anti-PD1 and assess induced changes in anti-LAM immunity.

Methods: We used a murine model of metastatic LAM to determine survival after intranasal treatment with TLR9 agonist CpG at two doses and in combination the checkpoint inhibitor immunotherapy, anti-PD-1. We used histology and flow cytometry to assess overall inflammation as well as changes in the immune response upon treatment.

Measurements and Main Results: We found that local administration of CpG enhances survival in a murine model of LAM and that a lower dose more effectively balanced the inflammation induced by CpG with the anti-LAM therapeutic benefits. We also found that CpG reduces regulatory T cell infiltration in LAM lungs and that CD4 helper T cells are skewed toward pro-inflammatory phenotypes. We also found that CpG treatment is effective in both early stage and progressive disease and that CpG is synergistic with previously tested anti-PD1 therapy.

Conclusions: We have found that TLR9 agonist CpG can be used as LAM immunotherapy and effectively synergizes with anti-PD1 therapy in LAM.

Abstract Word Count: 241

Keywords: TLR agonist, regulatory T cells, metastatic murine LAM, combination immunotherapy

INTRODUCTION

Lymphangioleiomyomatosis (LAM) is a progressive rare lung disease, primarily affecting

women of child-bearing age (1). LAM patients can experience a range of debilitating

symptoms from shortness of breath, lung collapse, and eventual respiratory failure. LAM

is characterized by the persistent abnormal growth of smooth muscle-like cells (LAM

cells) and the formation of thin-walled cysts predominantly in the lungs, with

involvement of the surrounding lymphatics and kidneys. LAM is caused by inactivating

mutations in tumor suppressor genes tuberous sclerosis complex 1 and 2 (TSC1 /

TSC2) that negatively regulate cell growth through inhibition of a signaling protein

complex called mammalian target of rapamycin (mTOR). Currently, sirolimus

(rapamycin) is the only FDA approved drug for LAM (2), which inhibits the constitutively

activated mTOR complex thus slowing disease progression (3,4). However, rapamycin

does not promote disease regression or lung regeneration (5,6). Furthermore, patient

responses to sirolimus are incomplete and not always well tolerated for life-long use (5).

Therefore, there is an urgent need for true curative treatments that can both eliminate

LAM nodules and restore healthy lung function in patients suffering from LAM.

LAM is designated as a neoplastic growth, and recent research debate suggests it has

similarities to cancer (7). Immunosuppression is a common feature of many cancers,

including lung cancers(8)(9), which over time conditions immune cells to ignore the

tumor cells thus promoting cancer progression (10–12). In human LAM tissues, a lack of

T-cell infiltration suggests immunosuppression due to unresponsiveness of lymphocytes

to LAM antigens (13). LAM lungs typically have elevated levels of secreted factors like

monocyte chemoattractant protein 1 (MCP-1) (14) which may contribute to recruitment

of immunosuppressive cells (15,16). Prior work by us and others has also shown that LAM causes overexpression of checkpoint molecules PD-1 and CTLA-4 on T cells and LAM nodules have elevated expression of PD-1's binding partner, PD-L1 (17,18). Anti-PD1/PD-L1 and anti-CTLA-4 therapies were shown to enhance survival in murine models of LAM (17,18), suggesting that countering immunosuppression in LAM could be an effective strategy for immunotherapy.

In addition to the previously investigated checkpoint inhibitor treatments, reducing immunosuppression in LAM could be accomplished by reactivating antigen presenting cells (APCs) in the local tumor environment. Recent studies investigating targeted activation of APC's in cancer show enhanced anti-tumor immunity through improved T cell responses (19), and increased responsiveness of immunosuppressed myeloid cells to immunotherapy (20). Local tumor treatments inducing APC activation have also been shown to stimulate systemic anti-tumor immunity and regression of distant tumors (21). APCs are particularly responsive to molecules that mimic microbial pathogens to stimulate toll-like receptors (TLRs). When engaged through TLR activation, APCs like dendritic cells and macrophages secrete inflammatory cytokines and promote the differentiation of cytotoxic T cells (22,23). Several TLR agonists have received FDA approval for treatment of various cancers. For example, BCG is a TLR4 agonist that is used as immunotherapy for bladder cancer. Currently, there are 5 clinical trials investigating the use of TLR3, and TLR7/8 agonists as cancer vaccine adjuvants (NCT04364230), monotherapy (NCT04819373), or in combination with immunotherapy (NCT04278144, NCT03486301, NCT04840394). Three active clinical trials are also investigating targeted delivery of TLR9 agonists for treatment

of various cancers (NCT02668770, NCT05607953, NCT04935229). TLR9 agonists, like CpG, have been approved for use in humans as adjuvants in vaccination and CpG in particular has been explored as therapy in multiple cancers including non-small cell lung cancer (24–26). TLR9 activation has also been shown to be synergistic with immune checkpoint inhibition, making resistant tumors more responsive to therapy (27–31). Given the current efforts at leveraging TLR9 agonist use in cancer therapy, we propose to investigate its potential for therapy in LAM. Here, we investigate the efficacy of TLR9 agonist, CpG with or without simultaneous checkpoint inhibition as immune activating treatments for LAM and assess how CpG modulates anti-LAM immunity.

METHODS

Mice

7 to 9-week old female C57BL6/J mice (Jackson Laboratory) were used for all studies. Study protocols were approved by the UMD Institutional Animal Care and Use Committee (IACUC).

LAM induction

TSC2-null kidney-derived epithelial tumor (TTJ) cells, were acquired from the Krymskaya Lab at the University of Pennsylvania (17). TTJ cells (5.0 x 10^5) were administered in 100 μL of PBS via tail vein injection to induce LAM disease in 8-9 week old female mice. Lung lesions typically develop within 3-7 days.

Survival Studies

Mice received CpG class B (5 μg/10 μg) 2x/week (Invivogen, tlrl-1826-5), in 50 μL of PBS, intranasally starting 4 days post LAM induction. Mice receiving combination

therapy, received intranasal CpG and 300 µg anti-PD-1 antibody (BioXCell, BE0146-A0) in 200 µL of PBS via intraperitoneal (IP) injection 2x/week. Control groups received 50 µL of PBS intranasally and 300 µg of rat IgG isotype control (BioXCell, BE0089) by IP injection. Mice were monitored for symptoms of disease and body weight changes and removed from studies based on experimental timing or humane endpoints. At endpoint, mice were euthanized and lung tissues collected for histological analysis.

Histology

Tissues were fixed overnight in 4% PFA at 4ºC, embedded in paraffin wax, sectioned at 5 µm, and stained with hematoxylin and eosin (H&E). LAM nodule counts, average nodule diameters, and general tissue inflammation were quantified using QuPath Imaging software.

Flow Cytometry

After LAM induction, mice received a single intranasal treatment dose of CpG (5 µg or 10 µg) in 50 µL PBS or 50 µL of PBS as a vehicle control. At 1 and 3 days post treatment, lung tissues were perfused and collected for flow cytometric analysis. Left lung lobes were dissociated by enzymatic digestion with 1 mg/mL collagenase D (Roche, 11088882001), and 1 mg/mL collagenase IV (Worthington Biochemical, LS004188) in DMEM with 5% FBS for 1 hour. Single cell suspensions were stained to identify eosinophils (CD64-, Siglec F+, Ly6G-), neutrophils (CD64-, Siglec F-, Ly6G+) dendritic cells (CD11c+ MHC-II^{high}), alveolar macrophages (CD64+, SiglecF^{high}, CD11b-), interstitial macrophages (CD64+, SiglecF^{low}, CD11b+, MHC-II+), B Cells (CD19+ or B220+), T Cells (CD3+, CD4+ or CD8+), regulatory T Cells (CD3+, CD4+, FoxP3+), Th1 T cells (CD3+, CD4+, FoxP3-, Tbet+), Th2 T cells (CD3+, CD4+, FoxP3-, Gata3hi),

and Th17 T Cells (CD3+, CD4+, FoxP3-, ROR-γt+). Cells were analyzed by flow cytometry using a BD FACSCelesta (BD Biosciences) and FlowJo (FlowJo LLC). Gating strategies can be found in the data supplement (**Fig E1**).

ELISA

For cytokine analysis, lungs (smallest right lobe) were snap-frozen and stored at -80°C. For tissue homogenates, snap-frozen lungs were homogenized at 4°C at 20% w/v in protein extraction reagent (T-PER, Thermo Fisher Scientific) containing Halt™ Protease Inhibitor Cocktail (Thermo Fisher Scientific). Lung tissue homogenates were centrifuged at 15,000 g for 10 min at 4°C. Supernatants were aliquoted and stored at -80°C. Supernatants were diluted 1:10 with 1% BSA ELISA assay diluent and ELISA was performed according to manufacturer's instruction (ELISA MAX™ Standard Set, Biolegend).

Statistical Analysis

Data were analyzed using Graphpad Prism (GraphPad, La Jolla, CA).

Survival studies were assessed using the log-rank (Mantel-Cox) test. Unpaired Welch t tests were used to compare pairs of samples or groups. Mann–Whitney test was used to compare the groups of data that failed to satisfy parametric assumptions. Brown-Forsythe and Welch One-way ANOVA were used for comparisons between ≥3 groups. Differences were considered statistically significant when $p < 0.05$. *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

RESULTS

Treatment with Intranasal CpG Increases Median Survival in Murine LAM and is dose dependent

To determine the therapeutic benefit of TLR9-agonist CpG in LAM disease, we performed survival studies comparing the effect of administering routine doses of intranasal CpG during the course of LAM disease in our metastatic mouse model. We found that intranasal CpG increases survival in murine LAM from 32.5 to 45 days (**Fig 1A**) and that this effect is dose dependent, with lower dose CpG (5μg) resulting in higher median survival (60 days) than higher dose CpG (10μg, 45 days). We conducted a cross sectional study after 3 weeks (Day 22) and 6 total treatment administrations to investigate the effect of treatment with CpG in LAM lungs at a point where the treatment groups begin to diverge in survival studies. Histological analysis of these tissues showed that treatment with CpG slows the development of LAM-like nodules in mice as indicated by the reduced tumor burden and nodule sizes in treated vs untreated lungs (**Fig 1B**). We found that higher dose CpG (10ug), significantly decreased the nodule count, more that the lower dose (5ug), which was surprising given the improvement in overall survival with lower dose CpG (**Fig 1C**). However, when we considered that at both doses, the combined measure of tumor burden and inflammation are similar (**Fig 1C**), we found that higher dose CpG causes higher levels of inflammation and may actually contribute to increased morbidity in mice with LAM. Consistent with this assessment, the histological images in the higher dose CpG show considerable immune cell infiltration and thickened alveolar walls compared to lower dose CpG (**Fig 1B**).

Intranasal CpG modulates Immune cell infiltration in LAM Lungs

We next sought to evaluate the effects of CpG treatment on immune cell infiltration in LAM. We conducted flow cytometry analysis of immune cells at two cross-sectional timepoints, day 5 and day 22, of our initial survival study. We found that after the first dose of CpG, given early in the course of disease on day 5, populations of eosinophils, alveolar macrophages, dendritic cells, and B and T cells were significantly decreased at both low and higher dose CpG. In contrast, both neutrophils and interstitial macrophages, and monocytes were increased in response to CpG treatment (**Fig 2A, Supplementary Figure 2**). By day 22, CpG treatment in LAM mice resulted in decreased cell counts in all identified immune populations except B cells and interstitial macrophages when compared to counts in untreated LAM controls (**Fig 2B**). Comparing the expansion of immune cells within treatment groups between Day 5 and Day 22, we found up to 5 fold increases in T and B cell counts in CpG treated mice compared to up to 10 fold increases in disease controls (**Figures E3-4**). Between Day 5 and Day 22, untreated mice had greatest expansion of interstitial macrophages and cDCs, while CpG treated mice had greatest expansion in cDCs and T Cells (**Supplementary Figures 3-4**). B cells in untreated mice were the only cell population with little expansion from Day 5 to 22. In CpG treated lungs, there was no expansion in eosinophil, alveolar macrophage and monocyte populations. This suggests that CpG stimulates inflammation in the tissue, and recruits a different assortment of cells while restraining the inflammation seen in untreated LAM.

Intranasal CpG shifts T Cells toward inflammatory phenotypes and reduces regulatory T cells

Although overall CD4+ and CD8+ T cell numbers were not different between treated and untreated LAM lungs (**Figure E2**), CpG treated LAM lungs show signs of reduced immunosuppression. Flow cytometry analysis reveals that CD8+ T cells have increased expression of activation marker ICOS (**Fig 3A**) and the proportion of effector memory CD8+ T cells is significantly increased in our CpG treatment groups (**Fig 3B**). This suggests that CpG treatment activates cytotoxic CD8+ T cells. Cytotoxic T cells are the main producers of IFN-γ, so we measured IFN-γ in lung tissue homogenates by ELISA and found that CpG-treated lungs have increased levels of IFN-γ (**Fig 3A**), suggesting increased cell-mediated cytotoxicity.

Among CD4+ helper T cells subsets, we found significantly decreased regulatory T cell numbers in CpG treated mice (**Fig 3C**). We did not see decreases in populations of Th1 or Th17 cells (**Fig 3C**), therefore we can attribute the overall decrease in CD4 T cells in the CpG treatment groups to a decline in regulatory T cells. This is a positive indicator of decreased immunosuppression. Furthermore, we found increased proportions of Th1 and Th17 cells (**Fig 3C**) suggesting that T cells in CpG treated LAM lungs are polarized towards more proinflammatory phenotypes.

Late intervention CpG treatment in LAM mice

Given the encouraging results in administering CpG early in LAM disease, and since LAM can be diagnosed at a variety of disease stages, we next evaluated CpG's therapeutic potential in more advanced disease. We induced LAM in mice and allowed disease to develop for 14 days, before beginning intranasal treatment with CpG. We

found that lower dose CpG (5 µg) had no significant effect on survival (**Fig 4A**), but a higher dose of CpG (20 µg) showed some therapeutic benefit (**Fig 4A**). We also found that the CpG treatment did not affect tissue inflammation, nodule count, or nodule size in more advanced disease (**Fig 4B**). However, we did observe that nodules in CpG-treated mice showed histological signs of necrosis in LAM nodules, unlike untreated LAM controls (**Fig 4A**)

We then compared the immune response in CpG treated LAM mice with advanced disease to untreated LAM controls. We evaluated the effects of CpG treatment on the immune response 1 and 3 days following the first intranasal treatment of CpG given 14 days post disease induction. Unlike the striking responses seen during early intervention, CpG given later in disease did not significantly affect immune cell infiltration in LAM lungs. We found only a decrease in eosinophils and cDCs at day 15; all other populations of identified immune cells remained unchanged (**Fig 5**).

Combination CpG and anti-PD1 therapy work synergistically to increase Survival in Murine LAM

We have previously shown that treatment with anti-PD-1 antibody was effective at prolonging survival in murine LAM (17). In addition, CpG has been shown to enhance anti-tumor immunity in pre-clinical cancer models when given in combination with chemotherapy or checkpoint inhibition(32–34). Therefore, we conducted a survival study to evaluate combination therapy of CpG with checkpoint inhibition to determine potential synergistic effects in LAM treatment. We found that combination CpG and anti-PD-1 therapy increased survival in mice (**Fig 6A**) compared to CpG or anti-PD1 treatment alone. The median survival in mice receiving combination therapy was 65

days compared to 45 days in the CpG treatment group and 49 days in the anti-PD1 treatment group. This suggests that that TLR9 activation can make LAM more responsive to checkpoint inhibition. Histological analysis of tissues collected at symptomatic endpoints showed that combination CpG and anti-PD-1 therapy resulted in reduced LAM nodule counts but had no effect on overall inflammation as a result of increased nodule sizes (**Fig 6B-C**). We observed signs of necrosis (areas with less nuclear hematoxylin staining) in these larger nodules (**Fig 6B**) that were not present in nodules of untreated LAM controls. Tumor necrosis is often associated with enhanced cytotoxic T-cell responses as a result of immune checkpoint inhibition.

DISCUSSION

Our research offers preclinical evidence to support the benefit of adjuvant-mediated immunotherapy in the treatment of LAM. We found that intranasal treatment with CpG at early stage disease increased survival in mice with LAM. This treatment slowed the progression of disease in the lungs, and decreased the LAM nodule burden in treated mice while increasing the population of cytotoxic and effector CD8+ T cells and decreasing the population of regulatory T cells in the LAM microenvironment. We have previously shown that anti-PD-1 therapy increases survival in mice with LAM and others have found that in cancer immunotherapy, checkpoint inhibitors can synergize with immune adjuvants. Here, we observed synergistic effects of systemic checkpoint inhibition when combined with local CpG adjuvant therapy. This finding suggests that immune activating therapies, particularly combination therapies, could be an effective strategy to treat LAM.

Striking the balance between good/bad inflammation

A sophisticated balance of pro- and anti-inflammatory signals helps regulate immune responses in the body. Immunotherapies influence these pathways, tipping the scale towards inflammation to produce impressive anti-tumor responses. However, the inflammation induced can also lead to immune mediated adverse effects and inflammatory toxicities. These side effects have hindered the development of immunotherapies as alternative cancer treatments. Choosing the right dose and treatment schedule to maintain equilibrium between treatment efficacy and runaway inflammation could be key to avoiding undesirable tissue damage and severe symptoms. Our finding that lower dose CpG therapy produces less inflammation in the lungs and increases survival in LAM mice despite the increase in nodule count (compared to higher dose CpG), suggests that not all inflammation is helpful for the anti-LAM immune response. In fact, too much inflammation can cause damage in an already delicate tissue environment and be detrimental to treatment outcomes. A recent clinical trial looking at inhaled TLR9 agonist for the treatment of lung cancer reported several adverse events in patients all related to inflammatory immune responses (NCT03326752) (35). For CpG, lower doses may be effective to reign in inflammation while stimulating enough of an inflammatory immune response to drive anti-LAM immunity. Nanoparticle drug delivery techniques can also be employed to target even smaller doses of CpG to specific cell types, which could reduce tissue-wide inflammation while enhancing therapeutic immune responses to inhibit LAM nodule proliferation. Because LAM nodules are composed of many cell types including fibroblasts that may contribute to LAM pathology (36–38), targeting treatment to

immune cells could minimize stromal cell exposure to CpG stimulation and curb fibrosis and collagen deposition (tissue remodeling) (39).

The timing of drug administration is another variable in optimizing treatment efficacy. In cancer treatment, immunotherapies like immune checkpoint inhibitors are commonly administered to patients at a high dose every 2-3 weeks, while researchers typically administer these drugs to mice twice weekly. This striking difference suggests that further pharmacokinetic analyses are required prior to translating LAM immunotherapies from mice to humans. In clinical trials assessing safety and efficacy of TLR9 agonists for cancer, the most common dosing schedule was 1-15 mg CpG once per week (NCT00070629, NCT00299728, NCT00292045, NCT00031278), while murine studies including ours can vary from giving a single dose of CpG to up to two doses per week. Our study involved repeated stimulation of TLR9 and it is possible that the frequency of CpG dosing in mice may induce a tolerizing immune response known as TLR or endotoxin tolerance. TLR tolerance is characterized by reduced pro-inflammatory cytokine production and transient unresponsiveness of immune cells upon re-exposure to TLR agonists within 5 days of a prior stimulation (40–45). This could hinder the ability of our therapeutic strategy to maintain an active inflammatory response in the lungs. Tolerance could mean an eventual restoration of the LAM microenvironment to its initial immunosuppressive state, which could explain why CpG, while beneficial, is not curative. Researchers found that delaying a secondary stimulation until after the refractory period (5 days) avoided the induction of TLR tolerance (46). Therefore, adjusting the timing of our treatment regimen in future studies to once weekly or every 2 weeks in mice could prolong the therapeutic effect.

Our results in context of current LAM therapy and potential for clinical translation

Combination therapy with existing LAM treatments: Because Rapamycin is the best treatment currently available to LAM patients, assessing its compatibility with CpG will be critical for consideration in clinical applications. Rapamycin is an immunosuppressive drug and could affect the immune response induced by CpG treatment. More specifically, the mTOR complex 1 is also involved in TLR9 activation through the myD88 signaling pathway, so inhibition of mTOR by Rapamycin treatment could potentially attenuate the inflammatory response induced by CpG treatment in antigen presenting cells. LAM is treated using low dose Rapamycin and how this low dose will interact with immune activating treatments in LAM is an unstudied area. Given that 40% of LAM patients have partial or no response to treatment with Rapamycin (47), TLR9 activation could offer these individuals a therapeutic strategy with similar benefits to standard of care.

Translating cancer immunotherapies to LAM: Several immunotherapies are currently approved for treatment of lung cancer and may be considered for LAM. These include monoclonal antibodies against EGFR and VEGF/VEGFR pathways and immune checkpoint inhibitors targeting PD-1/PD-L1 or CTLA-4. Clinical trials for adoptive T cell therapy and lung cancer vaccines are ongoing (48) and several immunotherapies are approved for advanced lung cancers (49). Similarly, clinical trials for LAM therapies have explored the use of FDA-approved anti-angiogenic and anti-fibrotic therapies targeting pathways involved in the progression of lung cancer and idiopathic pulmonary fibrosis (Bevacizumab, Nintedanib, Saracatinib), but so far with limited success. Immune checkpoint inhibitors have not yet been tested in LAM patients, but there is

promising research in mouse models to support this treatment strategy (17,18).

Furthermore, our work showing TLR9 agonist treatments can enhance survival in

murine LAM could be translated into the clinic in future, particularly since TLR9 agonists

have been approved for several cancers and are explored in clinical trials for lung

cancer. Finally, results from lung cancer studies suggest that combination of immune

checkpoint inhibitors and TLR9 agonists lead to successful anti-lung cancer immune re-

activation. Our data suggests that this strategy could be successful in LAM as well and

thus there is potential for designing LAM combination immunotherapies in the future.

Conclusion

Our work demonstrates that TLR9 agonist CpG can enhance survival in a mouse model

of LAM and synergizes with anti-PD1checkpoint inhibitor immunotherapy. Our work

highlights a new potential area to develop as LAM treatments, combination

immunotherapies, and lays the foundation for clinical translation of LAM vaccines.

REFERENCES

1. Lymphangioleiomyomatosis (LAM. Breathe (Sheff. 2017;13(1):64–71.

2. Singla A, Gupta N, Apewokin S, McCormack FX. Sirolimus for the treatment of lymphangioleiomyomatosis. Expert Opinion on Orphan Drugs. 2017;5(11):907–21.

3. Elia D, Cssandro R, Torre O, Harari S. Long Term Sirolimus therapy for LAM: a real life experience. European Respiratory Journal. 2017;50(suppl 61).

4. Yao J, Taveira-DaSilva AM, Jones AM, Julien-Williams P, Stylianou M, Moss J. Sustained effects of sirolimus on lung function and cystic lung lesions in lymphangioleiomyomatosis. Am J Respir Crit Care Med. 2014;190(11):1273–82.

5. McCormack FX, Inoue Y, Moss J, Singer LG, Strange C, Nakata K, et al. Efficacy and Safety of Sirolimus in Lymphangioleiomyomatosis. New England Journal of Medicine. 2011 Apr 28;364(17):1595–606.

6. Bissler JJ, McCormack FX, Young LR, Elwing JM, Chuck G, Leonard JM, et al. Sirolimus for Angiomyolipoma in Tuberous Sclerosis Complex or Lymphangioleiomyomatosis. New England Journal of Medicine. 2008 Jan 10;358(2):140–51.

7. Prizant H, Hammes SR. Minireview: Lymphangioleiomyomatosis (LAM): The "Other" Steroid-Sensitive Cancer. Endocrinology. 2016 Sep;157(9):3374–83.

8. Tormoen GW, Crittenden MR, Gough MJ. Role of the immunosuppressive microenvironment in immunotherapy. Adv Radiat Oncol. 2018;3(4):520–6.

9. Wu AA, Drake V, Huang HS, Chiu S, Zheng L. Reprogramming the tumor microenvironment: tumor-induced immunosuppressive factors paralyze T cells. Oncoimmunology. 2015;4(7).

10. Gonzalez H, Hagerling C, Werb Z. Roles of the immune system in cancer: from tumor initiation to metastatic progression. Genes Dev. 2018;32(19–20):1267–84.

11. Blomberg OS, Spagnuolo L, Visser KE. Immune regulation of metastasis: mechanistic insights and therapeutic opportunities. Dis Model Mech. 2018;11(10).

12. Janssen LME, Ramsay EE, Logsdon CD, Overwijk WW. The immune system in cancer metastasis: friend or foe? Journal for ImmunoTherapy of Cancer. 2017;5(1).

13. Han F, Dellacecca ER, Barse LW, Cosgrove C, Henning SW, Ankney CM, et al. Adoptive T-Cell Transfer to Treat Lymphangioleiomyomatosis. Am J Respir Cell Mol Biol. 2020 Jun;62(6):793–804.

14. Pacheco-Rodriguez G, Kumaki F, Steagall WK, Zhang Y, Ikeda Y, Lin JP, et al. Chemokine-Enhanced Chemotaxis of Lymphangioleiomyomatosis Cells with Mutations in the Tumor Suppressor TSC2 Gene1. The Journal of Immunology. 2009 Feb 1;182(3):1270–7.

15. Mittal P, Wang L, Akimova T, Leach CA, Clemente JC, Sender MR, et al. The CCR2/MCP-1 Chemokine Pathway and Lung Adenocarcinoma. Cancers. 2020 Dec;12(12):3723.

16. Gok Yavuz B, Gunaydin G, Gedik ME, Kosemehmetoglu K, Karakoc D, Ozgur F, et al. Cancer associated fibroblasts sculpt tumour microenvironment by recruiting monocytes and inducing immunosuppressive PD-1+ TAMs. Sci Rep. 2019 Feb 28;9(1):3172.

17. Maisel K, Merrilees MJ, Atochina-Vasserman EN, Lian L, Obraztsova K, Rue R, et al. Immune Checkpoint Ligand PD-L1 Is Upregulated in Pulmonary Lymphangioleiomyomatosis. Am J Respir Cell Mol Biol. 2018 Dec;59(6):723–32.

18. Liu HJ, Lizotte PH, Du H, Speranza MC, Lam HC, Vaughan S, et al. TSC2-deficient tumors have evidence of T cell exhaustion and respond to anti–PD-1/anti–CTLA-4 immunotherapy. JCI Insight. 2018 Apr 19;3(8):e98674.

19. Chang H, Zou Z, Wang Q, Li J, Jin H, Yin Q, et al. Targeting and Specific Activation of Antigen-Presenting Cells by Endogenous Antigen-Loaded Nanoparticles Elicits Tumor-Specific Immunity. Adv Sci. 2020 Jan;7(1):1900069.

20. De Henau O, Rausch M, Winkler D, Campesato LF, Liu C, Cymerman DH, et al. Overcoming resistance to checkpoint blockade therapy by targeting PI3Kγ in myeloid cells. Nature. 2016 Nov;539(7629):443–7.

21. Suek N, Campesato LF, Merghoub T, Khalil DN. Targeted APC Activation in Cancer Immunotherapy to Enhance the Abscopal Effect. Front Immunol. 2019 Apr 2;10:604.

22. Barr TA, Brown S, Ryan G, Zhao J, Gray D. TLR-mediated stimulation of APC: Distinct cytokine responses of B cells and dendritic cells. Eur J Immunol. 2007 Nov;37(11):3040–53.

23. Grassin-Delyle S, Abrial C, Salvator H, Brollo M, Naline E, Devillier P. The Role of Toll-Like Receptors in the Production of Cytokines by Human Lung Macrophages. J Innate Immun. 2020;12(1):63–73.

24. Brignole C, Marimpietri D, Paolo D. Therapeutic targeting of TLR9 inhibits cell growth and induces apoptosis in neuroblastoma. Cancer Res. 2010;70(23):9816–26.

25. Jahrsdorfer B, Weiner GJ. CpG oligodeoxynucleotides as immunotherapy in cancer. Update Cancer Ther. 2008;3(1):27–32.

26. CPG 7909 Injection in Non-Small Cell Lung Cancer [Internet]. Available from: https://ClinicalTrials.gov/sh

27. Cheng Y, Lemke-Miltner CD, Wongpattaraworakul W, Wang Z, Chan CHF, Salem AK, et al. In situ immunization of a TLR9 agonist virus-like particle enhances anti-PD1 therapy. J Immunother Cancer. 2020 Oct 1;8(2):e000940.

28. Gallotta M, Assi H, Degagné É, Kannan SK, Coffman RL, Guiducci C. Inhaled TLR9 Agonist Renders Lung Tumors Permissive to PD-1 Blockade by Promoting Optimal CD4+ and CD8+ T-cell Interplay. Cancer Res. 2018 Sep 1;78(17):4943–56.

29. Reilley MJ, Morrow B, Ager CR, Liu A, Hong DS, Curran MA. TLR9 activation cooperates with T cell checkpoint blockade to regress poorly immunogenic melanoma. Journal for ImmunoTherapy of Cancer. 2019 Nov 26;7(1):323.

30. Sato-Kaneko F, Yao S, Ahmadi A, Zhang SS, Hosoya T, Kaneda MM, et al. Combination immunotherapy with TLR agonists and checkpoint inhibitors suppresses head and neck cancer. JCI Insight. 2017 Sep 21;2(18):e93397.

31. Mangsbo SM, Sandin LC, Anger K, Korman AJ, Loskog A, Tötterman TH. Enhanced tumor eradication by combining CTLA-4 or PD-1 blockade with CpG therapy. J Immunother. 2010 Apr;33(3):225–35.

32. Cheng Y, Wang C, Wang H, Zhang Z, Yang X, Dong Y, et al. Combination of an autophagy inhibitor with immunoadjuvants and an anti-PD-L1 antibody in multifunctional nanoparticles for enhanced breast cancer immunotherapy. BMC Med. 2022 Oct 28;20(1):411.

33. Gu F, Hu C, Cao W, Li C, Xia Q, Gao Y, et al. Tumor Microenvironment Multiple Responsive Nanoparticles for Targeted Delivery of Doxorubicin and CpG Against Triple-Negative Breast Cancer. Int J Nanomedicine. 2022;17:4401–17.

34. Zhang M, Fang Z, Zhang H, Cui M, Wang M, Liu K. Reversing tumor immunosuppressive microenvironment via targeting codelivery of CpG ODNs/PD-L1 peptide antagonists to enhance the immune checkpoint blockade-based anti-tumor effect. European Journal of Pharmaceutical Sciences. 2022 Jan 1;168:106044.

35. Garon E, Spira AI, Johnson M, Bazhenova L, Leach J, Candia A, et al. Abstract CT224: Phase Ib/II, open label, multicenter study of inhaled DV281, a Toll-like receptor 9 agonist, in combination with nivolumab in patients with advanced or metastatic non small cell lung cancer (NSCLC). Cancer Research. 2019 Jul 1;79(13_Supplement):CT224–CT224.

36. Clements D, Dongre A, Krymskaya VP, Johnson SR. Wild Type Mesenchymal Cells Contribute to the Lung Pathology of Lymphangioleiomyomatosis. PLOS ONE. 2015 May 15;10(5):e0126025.

37. Clements D, Miller S, Babaei-Jadidi R, Adam M, Potter SS, Johnson SR. Cross talk between LAM cells and fibroblasts may influence alveolar epithelial cell behavior in lymphangioleiomyomatosis. American Journal of Physiology-Lung Cellular and Molecular Physiology. 2022 Feb;322(2):L283–93.

38. Babaei-Jadidi R, Dongre A, Miller S, Castellanos Uribe M, Stewart ID, Thompson ZM, et al. Mast-Cell Tryptase Release Contributes to Disease Progression in Lymphangioleiomyomatosis. Am J Respir Crit Care Med. 2021 Aug 15;204(4):431–44.

39. Trujillo G, Meneghin A, Flaherty KR, Sholl LM, Myers JL, Kazerooni EA, et al. TLR9 Differentiates Rapid from Slowly Progressive Forms of Idiopathic Pulmonary Fibrosis. Sci Transl Med. 2010 Nov 10;2(57):57ra82.

40. Biswas SK, Lopez-Collazo E. Endotoxin tolerance: new mechanisms, molecules and clinical significance. Trends in Immunology. 2009 Oct 1;30(10):475–87.

41. Yeo SJ, Yoon JG, Hong SC, Yi AK. CpG DNA Induces Self and Cross-Hyporesponsiveness of RAW264.7 Cells in Response to CpG DNA and Lipopolysaccharide: Alterations in IL-1 Receptor-Associated Kinase Expression. The Journal of Immunology. 2003 Jan 15;170(2):1052–61.

42. Tsukada K, Kitazawa T, Fukushima A, Okugawa S, Yanagimoto S, Tatsuno K, et al. Macrophage tolerance induced by stimulation with Toll-like receptor 7/8 ligands. Immunology Letters. 2007 Jul 31;111(1):51–6.

43. Naqvi I, Giroux N, Olson L, Morrison SA, Llanga T, Akinade TO, et al. DAMPs/PAMPs induce monocytic TLR activation and tolerance in COVID-19 patients; nucleic acid binding scavengers can counteract such TLR agonists. Biomaterials. 2022 Jan 28;121393.

44. Butcher SK, O'Carroll CE, Wells CA, Carmody RJ. Toll-Like Receptors Drive Specific Patterns of Tolerance and Training on Restimulation of Macrophages. Front Immunol. 2018;9:933.

45. Kwok SK, Lee JY, Park SH, Cho ML, Min SY, Park SH, et al. Dysfunctional interferon-alpha production by peripheral plasmacytoid dendritic cells upon Toll-like receptor-9 stimulation in patients with systemic lupus erythematosus. Arthritis Res Ther. 2008;10(2):R29.

46. Bourquin C, Hotz C, Noerenberg D, Voelkl A, Heidegger S, Roetzer LC, et al. Systemic Cancer Therapy with a Small Molecule Agonist of Toll-like Receptor 7 Can Be Improved by Circumventing TLR Tolerance. Cancer Research. 2011 Jul 28;71(15):5123–33.

47. Revilla-López E, Berastegui C, Méndez A, Sáez-Giménez B, Ruiz de Miguel V, López-Meseguer M, et al. Long-term results of sirolimus treatment in lymphangioleiomyomatosis: a single referral centre experience. Sci Rep. 2021 May 13;11(1):10171.

48. Zhong S, Cui Y, Liu Q, Chen S. CAR-T cell therapy for lung cancer: a promising but challenging future. J Thorac Dis. 2020 Aug;12(8):4516–21.

49. Non-Small Cell Lung Cancer Treatment (PDQ®)–Health Professional Version - NCI

[Internet]. 2022 [cited 2023 Feb 2]. Available from:

https://www.cancer.gov/types/lung/hp/non-small-cell-lung-treatment-pdq

FIGURE LEGENDS

Figure 1. *Intranasal CpG increases survival in murine model of metastatic LAM, decreases tissue-wide inflammation and overall LAM nodule burden.* **A)** Aggregate data from three survival experiments LAM n=34, CpG (10ug) n=22, CpG (5ug) n=15. **B)** Representative H&E stained histological images of mouse lungs from treatment groups: untreated LAM control, CpG 5 µg, and CpG 10 µg. Scale bar = 500 µm. **C)** Quantification of LAM nodules and Inflamed tissue regions between treatment groups 22 days post LAM induction and after 6 total treatment administrations. 3 sections analyzed per mouse. Statistical analysis was performed using log-rank test (survival studies) and one-way ANOVA (histological analysis). * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001.

Figure 2. *CpG treatment modulates immune cell infiltration in LAM.* Data shows total cell numbers in the left lung lobe of granulocytes and antigen presenting cells at day 5 or day 22 after LAM induction and 1 or 18 days respectively after CpG treatment has started. Data is representative of n= 2+ experiments and n>8 mice. Statistical analysis was performed using one-way ANOVA. * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001.

Figure 3. *Helper T cells in CpG treated LAM lungs shift away from immunosuppressive phenotypes and CD8 T cells express activation markers.* Data shows T cell populations in the left lung lobe of 22 days after LAM induction and 18 days after beginning CpG treatment. **A)** ICOS expression CD8+T Cells and cytokine levels in

LAM lungs. **B)** Effector memory, central memory, and naïve CD8+ T cell subsets. **C)** CD4 helper T cell subsets: total cell numbers and % of CD4+ cells. Data is representative of n= 2+ experiments and n>5 mice. Statistical analysis was performed using one-way ANOVA. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure 4. *Late intervention intranasal CpG increases survival in murine model of metastatic LAM at higher dose.* **A)** Aggregate data from three survival experiments LAM n=34, CpG (10ug) n=14, CpG (5ug) n=16. **B)** Representative H&E stained histological images of mouse lungs from treatment groups: untreated LAM control, CpG 5 µg, and CpG 20 µg. Scale bar = 500 µm. **C)** Quantification of LAM nodules and Inflamed tissue regions between treatment groups of mice at symptomatic endpoints from survival studies in (A). 3 sections analyzed per mouse. Statistical analysis was performed using log-rank test (survival studies) and one-way ANOVA (histological analysis). * $p<0.05$.

Figure 5. *CpG treatment modulates immune cell populations in response to late intervention (starting at day 14) intranasal CpG in LAM lungs (Day 1 & Day 3 after CpG treatment).* Total cell numbers of granulocytes and antigen presenting cells infiltrating the lungs 1 and 3 days after CpG treatment obtained via flow cytometry. Data is representative of n= 3 experiments and n = 8+ mice. Statistical analysis was performed using one-way ANOVA. * $p<0.05$, ** $p<0.01$, *** $p<0.001$.

Figure 6. *CpG treatment is synergistic with ICI Therapy.* **A)** Aggregate data from four

survival experiments LAM n=34, CpG (10 µg) n=22, α-PD-1 n=16, CpG (10 µg) + α-

PD-1 n=20. **B)** Representative H&E stained histological images of mouse lungs from

treatment groups: untreated LAM control, CpG (10 µg), α-PD-1, CpG (10 µg) + α-

PD-1 . Scale bar = 500 µm. **C)** Quantification of LAM nodules and Inflamed tissue

regions between treatment groups of mice at symptomatic endpoints from survival

studies in (A). 3 sections analyzed per mouse. Statistical analysis was performed

using log-rank test (survival studies) and one-way ANOVA (histological analysis). *

$p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure E1: Flow Cytometry Gating Strategy

Figure E2: T cells after CpG treatment on day 22. Data shows CD4+ and CD8+ T cell

numbers, ICOS expression, and cell population distribution between effector

memory, central memory and naïve subsets. Data is representative of n= 3

experiments and n>10 mice. Statistical analysis was performed using one-way

ANOVA. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure E3: Total cell counts of granulocytes and antigen presenting cells on day 5 and

22 after CpG treatment. Statistical analysis was performed using two-way ANOVA. *

$p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure E4: Total cell counts of T and B cells on day 5 and 22 after CpG treatment.

Statistical analysis was performed using two-way ANOVA. * p<0.05, ** p<0.01, ***

p<0.001, **** p<0.0001.

FIGURES

Figure 1

Intranasal CpG increases survival in murine model of metastatic LAM, decreases tissue-wide inflammation and overall LAM nodule burden

Figure 2

CpG treatment modulates immune cell infiltration in LAM

Figure 3

Helper T cells in CpG treated LAM lungs shift away from immunosuppressive phenotypes and CD8 T cells express activation markers

Figure 4

Late intervention intranasal CpG increases survival in murine model of metastatic LAM at higher dose

Figure 5

CpG treatment modulates immune cell populations in response to late intervention

Figure 6

CpG treatment is synergistic with ICI Therapy.

Figure E1

Flow Cytometry Gating Strategy

Figure E2

T cells after CpG treatment on day 22

Figure E3

Total cell counts of granulocytes and antigen presenting cells on day 5 and 22 after CpG treatment

Figure E4

Total cell counts of T and B cells on day 5 and 22 after CpG treatment