Nucleus Accumbens Local Circuit for Cue-Dependent Aversive Learning

Andrew Belilos[1], Cortez Gray[2], Christie Sanders[1], Destiny Black[2], Elizabeth Mays[2], Christopher

Richie[1], Ayesha Sengupta[1], Holly Hake[1], T. Chase Francis[2]*

Affiliations:

1. Intramural Research Program, National Institute on Drug Abuse, National Institutes of

Health, Baltimore, Maryland 21224, USA

2. Department of Drug Discovery and Biomedical Sciences, College of Pharmacy,

University of South Carolina, Columbia, South Carolina 29208, USA

*Correspondence: francit@mailbox.sc.edu

Keywords: Substance P, Tachykinin, Medium Spiny Neuron, Long-term Potentiation, AMPA

receptors, Acetylcholine, Cholinergic Interneuron, Fear, Aversion, Learning

Summary

Response to threatening environmental stimuli requires detection and encoding of

important environmental features that dictate threat. Aversive events are highly salient which

promotes associative learning about stimuli that signal this threat. The nucleus accumbens is

uniquely positioned to process this salient, aversive information and promote motivated output,

through plasticity on the major projection neurons in the brain area. We uncovered a nucleus

accumbens core local circuit whereby excitatory plasticity facilitates learning and recall of

discrete aversive cues. We demonstrate that putative nucleus accumbens substance P release

and long-term excitatory plasticity on dopamine 2 receptor expressing projection neurons is

required for cue-dependent fear learning. Additionally, we found fear learning and recall were

dependent on distinct projection-neuron subtypes. Our work demonstrates a critical role for

Nucleus Accumbens substance P in cue-dependent aversive learning.

Introduction

Aversive stimuli are inherently salient, as animals need to recognize these threats to

survive. Consequently, aversive stimuli elicit rapid learning about and response to cues that

predict aversive outcomes. Canonically, the major projection neurons of the Nucleus

Accumbens (NAc) are studied in terms of motivational or incentive salience, a cognitive process

that facilitates approach or avoidance in response to a stimulus. These responses, however,

require recognition of a salient stimulus to promote these motivated outcomes. The NAc

responds to salient cues in humans [1], is required for aversive learning [2,3], which is cue-

dependent in the NAc core [3–5], suggesting the NAc core may facilitate Pavlovian learning

independently of the motivated response. If true, it is likely consolidation and subsequent recall

of these salient events requires long-term plasticity to facilitate persistent memories. These

NAc-dependent synaptic mechanisms to enable recognition of threatening, aversive stimuli

have yet to be determined.

Within the NAc, medium spiny projection neuron (MSN) subtypes are thought to promote

opposing outcomes, with dopamine 1 (D1) receptor MSN activity driving rewarding and positive

outcomes, and dopamine 2 (D2) receptor MSN activity driving aversive outcomes, suppressed

reward, and punishment [6–15]. This understanding has been updated and it is now known MSN

subtype activity participates in more complex and pivotal roles in guiding learned behavior. In

addition, NAc core D2-MSNs have been shown to be required for discrimination [16] and reversal

learning [17,18], and reward [19–21]. These additional known roles of D2-MSNs may hinge on the

ability of D2-MSNs to alter their response to salient stimuli. Thereby, stimulus-dependent activity

(e.g., through potentiation of excitatory transmission on D2-MSNs), during learning, may provide

a means of facilitating learning about aversion associated stimuli.

We recently uncovered a mechanism in which glutamatergic long-term excitatory

potentiation (LTP) is elicited on D2-MSNs exclusively by activity of D1-MSNs [22] and required the

activation of cholinergic interneurons by release of the neuropeptide substance P. Both

acetylcholine and substance P are released by salient stimuli including drugs of abuse and

aversive stimuli [23,24] and these neuromodulators contribute to many of the same behaviors as

activation of D2-MSNs. Additionally, phasic activity changes of cholinergic neurons in response

to cue-dependent associative outcomes have been observed in primates [25,26] and rodents [27].

Like D2-MSN neuronal activity, striatal acetylcholine is necessary for behavioral flexibility [28],

discrimination [29], reversal learning [30], and aids in the expression of cue-motivated behavior [31].

These similarities, and the fact that this circuit promotes plasticity position this circuit as a likely

candidate for signaling aversion-related outcomes. Therefore, we hypothesized this NAc core

substance P is necessary for supporting aversive fear learning through plasticity on D2-MSNs.

In this paper, we aimed to determine synaptic and circuit mechanisms which underpin

NAc core dependent aversive learning. Using pharmacology and gene knockout technology, we

found substance P receptors in cholinergic neurons were required for cue-dependent aversive

learning, in a fear conditioning context. These findings were corroborated through enhanced D1-

MSN activity and acetylcholine release in response to aversive foot shocks and D2-MSN activity

in response to learned cues. Disrupting this circuit at any synaptic and cellular node diminished

learning about cues. Our work provides a new perspective on how MSN subtypes interact to

facilitate learning about aversive cues associated with foot shock.

Results

Neurokinin 1 receptor signaling is required for cue-dependent aversive learning

Since substance P is released by strongly salient stimuli [23,24] and is involved in many of

the same aversive outcomes as D2-MSNs activation, we first tested whether putative substance

P release and binding to its primary receptor, the neurokinin 1 (NK1) receptor, was necessary

for aversive learning. A Pavlovian fear conditioning model was used as a model of aversive

learning, where freezing was the main measure of learning (**Figure 1A**). To determine if

substance P was required for associative aversive learning, mice were injected (i.p.) with the

brain penetrant neurokinin 1 receptor antagonist L-733,060 (10 mg/kg) prior to fear conditioning.

NK1 receptor antagonism suppressed cue-dependent, but not context dependent freezing

(**Figure 1B**), without affecting locomotion (**Figure S1**). Intraperitoneal injection of L-733,060

prior to cue recall had no effect on cue recall (**Figure 1C**). These results suggest substance P

signaling during aversive cue conditioning is required for the full expression of fear learning.

The NK1 receptor is found throughout the brain, including in regions known to be

associated with fear conditioning (e.g., amygdala and periaqueductal grey). In the NAc, the NK1

receptor (encoded by the *Tacr1* gene) is found in all choline acetyltransferase (ChAT)

expressing cholinergic cells. Therefore, we used a clustered regularly interspaces short

palindromic repeats (CRISPR) Cas9 system, generated an AAV expressing guide RNAs for the

Tacr1 gene (*Tacr1*-gRNAs), and injected it in the NAc core of ChAT-Cre/Cas9 mice, which

express Cas9 exclusively ChAT cells. RNA scope *in situ* hybridization was used to assess the

efficacy of the knockout. Since we were unable to separate ChAT cells expressing or not

expressing the *Tacr1* guide RNAs (see Methods), all *Chat* expressing cells in the NAc core

where GFP expression was observed were quantified for *Tacr1* expression. A significant

reduction in the *Tacr1* transcript in *Chat* positive cells was observed and no difference was

observed in *Tacr1* expression in *Chat* negative cells (**Figure 1D**). Additionally, a functional and

near complete inhibition of substance P-induced increase in ChAT firing was observed in mice

expressing Cas9 and the *Tacr1*-gRNAs (**Figure 1E**). Next, we tested the functional knockout in

fear conditioning behavior. NK1 receptor knockout suppressed cue-dependent, but not context-

dependent freezing (**Figure 1F**), suggesting the cue-dependent reduction in freezing due to

NK1 receptor antagonism was indeed an effect selective for the NAc core. In agreement with

this finding, local NAc core infusion of L-733,060 during conditioning was capable of

suppressing cue-dependent but not context-dependent recall compared to saline infused mice

(**Figure 1G**). Overall, these experiments indicate substance P signaling is required for the full

expression of cue-dependent recall of aversion-associated cues.

D1-MSNs are required for cue-dependent fear conditioning

The primary source of substance P in the striatum is D1-MSNs which exclusively express and release the peptide and we have demonstrated activation of D1-MSNs releases substance P [22]. Since substance P signaling is required for aversive learning and substance P is released in the striatum in response to salient aversive stimuli [23], we predicted D1-MSN activity would increase in response to salient, aversive stimuli. To selectively assay activity of D1-MSNs, Dyn-Cre mice were injected with AAV-DIO-GCaMP7f and implanted with fibers in the NAc core for assessment of *in vivo* calcium activity with fiber photometry as a means of assessing correlated neural activity (**Figure 2A**). We first acquired signals in response to random interval, unpredictable foot shocks. D1-MSN calcium activity scaled with the magnitude of foot shock (**Figure 2B-C**) but did not significantly vary across shock trials (**Figure 2D**). This scaling was not affected by injection of L-733,060 (**Figure S2A**). Next, we assayed activity during conditioning and cue recall. During conditioning, foot shock but not the novel cue increased D1-MSN calcium activity (**Figure 2E**). D1-MSN calcium activity was not increased by the cue during cue recall (**Figure 2F**), despite an increase in freezing (**Figure S2B**). Additionally, individual freeze events were not associated with changes in D1-MSN calcium activity (**Figure S2C**). We then scaled foot shock intensity to determine if the magnitude of the foot shock controlled both the D1-MSN calcium response and subsequent learning. As expected, freezing to the associated cue increased with foot shock intensity (**Figure 2G**). Interestingly, the area under the curve (AUC) of the shock-induced calcium signal during conditioning correlated significantly with the percent freezing (**Figure 2H**). Taken together, these results indicate the strength of D1-MSN activity during conditioning may be necessary for scaling cue-dependent aversive learning and recall.

To test the necessity of D1-MSN activity during cue-dependent aversive learning, we

used a temporally selective optogenetic strategy to inhibit D1-MSNs during cue-dependent

behavior. Dyn-Cre mice were injected with AAV-DIO-eNpHR3.0 and implanted with optogenetic

fibers in the NAc core (**Figure 2I**). Optogenetic inhibition of D1-MSNs did not affect locomotion

(**Figure S2D-E**). D1-MSNs were inhibited selectively for 10 sec during the cue and shock during

conditioning. D1-MSN inhibition during the cue and shock in conditioning diminished cue recall

but not context recall (**Figure 2J**). In another group of mice, we selectively inhibited D1-MSNs

during every other 10 sec cue to determine if D1-MSN inhibition reversibly inhibited cue-

dependent freezing. No change in cue recall was observed from D1-MSN inhibition (**Figure 2K-**

L). Altogether, these results indicate D1-MSN activity contributes to cue-dependent learning, but

not recall.

Substance P signaling is required for shock-driven acetylcholine release

Substance P is released from D1-MSNs in response to salient stimuli [23,32] and has a role

in aversion [33,34]. Additionally, substance P release promotes increased activity of ChAT neurons

via the NK1 receptor [22,35]. In addition, previous studies have indicated ChAT neuron exhibit

phasic activity patterns in response to aversive stimuli *in vivo* [26] . Therefore, it was predicted

substance P would cause ChAT activity and acetylcholine release *in vivo*. An adeno-associated

virus (AAV) containing the GPCR-activation-based sensor for acetylcholine (GRAB-ACh3.0) [36]

was injected into the NAc core of mice and expressed in neurons (**Figure 3A**). *Ex vivo* slice

studies demonstrated bath application of low (1 µM) and high (100 µM) concentrations of

acetylcholine significantly increase bulk fluorescence intensity in a concentration dependent

manner (**Figure S3A**). To evaluate acetylcholine binding *in vivo*, the virus was injected, a fiber

was implanted in the NAc core, and fluorescent transients were observed with fiber photometry.

In an open field chamber, mice were injected with the broad muscarinic antagonist scopolamine,

which has been observed to block acetylcholine binding to the GRAB-ACh3.0 sensor [36].

Scopolamine significantly diminished the average magnitude of the GRAB-ACh signal and

suppressed the amplitude of all transients (**Figure S3B**). These results verify that the GRAB-

ACh3.0 sensor causes fluorescence through acetylcholine release and binding to the GRAB-

Ach3.0 receptor sensor.

Next, to determine if acetylcholine is released by aversive stimuli in a substance P-

dependent manner, foot shocks (10 total) were given at random, unpredictable intervals in

behavior naïve mice while recording GRAB-ACh3.0 fluorescent activity in the NAc core. Shock

alone produced a decrease in fluorescence ("pause") followed by an increase ("rebound") in

acetylcholine-mediated fluorescence (**Figure 3B**), suggesting acetylcholine is released in a

multi-phasic pattern. This phasic pause and rebound has been observed in rewarding stimuli in

other studies [46]. Similar to D1-MSN calcium activity, increasing the intensity of the shock

increased the amount of rebound acetylcholine fluorescence following the shock without

significantly changing the AUC of the pause, despite habituation to repeated shock (**Figure 3C-**

D). Unlike D1-MSNs, this acetylcholine release habituated to the shock over repeated

presentations (**Figure 3C-D**). Next to determine if the NK1 receptor was responsible for the

rebound increase in acetylcholine fluorescence, L-733,060 was injected (*i.p.*) and mice were

exposed to random foot shocks at 0.6 mA. Antagonism of the NK1 receptor attenuated the

increase in acetylcholine-mediated fluorescence AUC during the rebound only (**Figure 3E-G**)

and did not diminish habituation to the shock in the rebound condition (**Figure 3G**). These

results suggest aversive stimuli increase acetylcholine release via substance P after foot shock

and acetylcholine release scales with the intensity of an aversive stimulus.

To determine how aversive conditioned stimuli affect this signal, mice injected with

GRAB-ACh3.0 in the NAc core were run through fear conditioning. Three prominent phasic

events were observed in acetylcholine-mediated fluorescence: a response to the novel cue, a

pause, and a rebound (**Figure 3H**). Like in the condition of the shock alone, the rebound

response was attenuating (**Figure S3C**). The rebound fluorescence, but not the other features,

was significantly diminished in mice treated with L-733,060 compared with saline injected mice

(**Figure 3I-J**). During cue recall, acetylcholine fluorescence was only increased at the

termination of the cue in saline treated mice but was increased at the start of the cue and the

termination of the cue in L-733,060 treated mice (**Figure S3C-D**). This increase in acetylcholine

fluorescence observed in L-733,060 treated mice is attenuated after the first presentation of the

cue during cue recall (**Figure S3D**). This may suggest that the cue itself still appears to be novel

to the mice with NK1 receptor antagonism. Interestingly, the rebound AUC significantly

correlated with the time spent freezing to the cue during cue recall (**Figure 3K**). Overall, these

results indicate that the increase in fluorescence following the shock is important for learning

about cues predicting aversive foot shock.

Shock-induced acetylcholine is important for learning about a conditioned aversive cue

To determine the necessity of NAc core ChAT cells are influencing learning about cues

predicting foot shocks, we injected ChAT-Cre mice with an AAV-DIO-eNpHR3.0 or AAV-DIO-

eYFP construct and inhibited ChAT neurons at two different time points during fear conditioning

to align inhibition to time windows of GRAB-ACh fluorescence: during the conditioning cue or

immediately after the foot shock (**Figure 4A**). Inhibition of ChAT cells had no observed effects

on locomotion (**Figure S4**). Mice that received inhibition during the cue showed no change in

context or cue recall compared to eYFP controls (**Figure 4B**). However, 5 sec of inhibition of

ChAT cells following the shock, during the observed rebound, significantly diminished freezing

to cues predicting shocks (**Figure 4C**), suggesting acetylcholine from ChAT neurons during the

rebound condition is necessary for the full expression of aversive cue recall.

Substance P causes long-term potentiation (LTP) of excitatory transmission on D2-

MSNs through enhanced acetylcholine release and activation of the muscarinic 1 receptor

signaling pathway [22]. This mechanism would provide a cell-type specific mechanism for

enhancing activity of D2-MSNs to either promote aversive outcomes or to facilitate learning

about aversive stimuli. To first determine if aversive conditioning and subsequent acetylcholine

release acts through muscarinic 1 receptors on D2-MSNs, mice were implanted with cannulas in

the NAc core and infused with the muscarinic 1 receptor-like antagonist pirenzepine (4 ng/µL,

bilateral) prior to conditioning or prior to cue retrieval. Compared to saline infused controls, cue

recall was suppressed when pirenzepine was infused prior to conditioning (**Figure 4D**), but not

prior to cue recall (**Figure 4E**).

Substance P promotes plasticity on D2-MSNs during cue-dependent aversive learning

D2-MSNs are likely to receive this excitatory muscarinic 1 receptor signaling more

strongly as D1-MSNs also express the inhibitory muscarinic 4 receptor which could suppress

this signaling. To determine if muscarinic 1 receptor plasticity occurred on D2-MSNs in a

substance P-dependent manner, we conditioned mice to a cue after injection with saline or L-

733,060 and conducted whole-cell electrophysiology recordings 1 day after cue recall.

Substance P-dependent LTP on D2-MSNs is primarily induced through insertion of GluR2-

lacking AMPA receptors [22]. These GluR2-lacking AMPA receptors are sensitive to intracellular

peptides at positive membrane potentials and excitatory synaptic currents at positive potentials

show inward rectification when these receptors have been inserted in the synaptic membrane.

Therefore, we assessed inward rectification of electrically evoked excitatory post-synaptic

currents (EPSCs), which would be suggestive of increased membrane insertion of GluR2-

lacking AMPA receptors. Mice that received conditioning (Cue + Shock) displayed significant

rectification at +40 mW as compared to non-conditioned mice (Cue only) (**Figure 5A**).

Additionally, mice injected with L-733,060 displayed no difference in rectification compared to

non-conditioned mice, suggesting that suppressing substance P signaling was responsible for

LTP observed on D2-MSNs (**Figure 5A**). As a secondary measure, we compared sensitivity of

electrically evoked currents on D2-MSNs from conditioned and non-conditioned mice to the

selective GluR2-lacking AMPA receptor antagonist 1-naphthyl acetyl spermine trihydrochloride

(NASPM) (100 µM). NASPM significantly diminished currents in conditioned mice relative to

non-conditioned mice, verifying the presence of more GluR2-lacking AMPA receptors (**Figure**

5B). Furthermore, amplitude, but not frequency of spontaneous excitatory post-synaptic

currents (sEPSCs) was increased on D2-MSNs (**Figure 5C**). Shock only mice or context only

mice displayed no changes in inward rectification on D2-MSNs (**Figure S4**). D1-MSNs did not

display inward rectification from any groups assayed (**Figure 5D and S4**). Results indicate

signatures of substance P-dependent LTP are expressed selectively in D2-MSNs following cued

aversive learning.

D2-MSNs activity is necessary for recall of aversive-conditioned cues

If LTP is selectively expressed on D2-MSNs and is dependent on aversive cue

associations, then it is likely D2-MSNs are active in response to aversively conditioned cues. To

verify this activity in real time and *in vivo*, A2a-cre mice were injected with AAV-DIO-GCaMP7f

in the NAc core and implanted with fibers for fiber photometry, A2a-Cre mice were injected with

AAV-DIO-GCaMP7f and implanted with fibers for photometry (**Figure 6A**) and subjected to 10

unpredictable foot shocks at various foot shock intensities. D2-MSNs responded similarly to all

foot shock intensities (**Figure 6B**) and L-733,060 did not alter D2-MSN calcium response to foot

shock (**Figure S6A**). In contrast to D1-MSNs, D2-MSN calcium activity did not increase with

increasing foot shock intensity (**Figure 6C**) and responses were variable across shock events

(**Figure 6D**). Like D1-MSNs, D2-MSNs responded to foot shock during conditioning, but not the

cue (**Figure 6E**). Unlike D1-MSNs, D2-MSNs were activated during cue recall (**Figure 6F**) in

response to the associated aversive cue and this signal did not habituate over repeated plays of

the cue (**Figure S6B**). Furthermore, D2-MSN calcium activity did not increase in response to

individual freezing events, suggesting freezing alone did not explain D2-MSN calcium signals

(**Figure S6C**). To determine if NK1 receptors were required during conditioning to cause this

increase to the cue, L-733,060 was injected in mice prior to conditioning. NK1 receptor

antagonism did not affect D2-MSN calcium response to the shock during conditioning (**Figure**

6G). Next, mice conditioned in the presence of L-733,060 were exposed to conditioned cues.

Photometry mice injected with L-733,060 showed significantly reduced freezing to the

conditioned cue compared to saline injected mice (**Figure S6D**). D2-MSN calcium activity did

not increase in response to conditioned cues in L-733,060 treated mice (**Figure 6H-I**). In

addition, AUC during the cue correlated with freezing (**Figure 6J**). Taken together, this suggests

NK1 receptors are required for D2-MSN activity in response to conditioned aversive cues.

Next, to determine if D2-MSN activity was required for recall of associated cues, AAV-

DIO-eNpHR3.0 was injected, and fibers were implanted into the NAc of A2a-cre mice for

selective optogenetic inhibition of D2-MSNs (**Figure 6K**). To compare the necessity of D2-

MSNs to D1-MSNs in conditioning, mice received optogenetic inhibition of D2-MSNs during both

cues during conditioning (**Figure 6K**). Surprisingly, inhibition of D2-MSNs during the cue in

conditioning significantly diminished freezing in the context (**Figure 6L**). As expected, D2-MSN

inhibition during conditioning had no effect on freezing to cues during cue recall when D2-MSNs

were not inhibited (**Figure 6M**). Despite this, inhibition of D2-MSNs during cue recall diminished

freezing (**Figure 6M**), suggesting D2-MSNs are required for the full expression of freezing to

aversively conditioned cues. To test this outcome without the potential confounds of inhibition

during cue conditioning, D2-MSNs were inhibited during the 10 second cue on every other cue

(**Figure 6N**). D2-MSN inhibition reversibly inhibited freezing to the cue (**Figure 6O**) without

affecting freezing to the context (**Figure 6N**). This suppression in freezing was not dependent

on locomotor effects (**Figure S6E-F**). Taken together, associative aversive stimuli promote

activity of D2-MSNs in response to associative cues and this response is required expression of

cue-dependent aversive learning.

Discussion

Our work has demonstrated a di-synaptic, local circuit within the NAc core that is critical

for processing aversive cues, with necessary roles for NAc core D1-MSN and D2-MSN activity

at different stages of learning and recall (**Figure 7**). While the activity does not appear to be

required for the full expression of the fear behavior, activity of these subtypes significantly

modulates the overall expression of the cue-dependent behavior, suggesting it does indeed play

a critical role in learning. We have shown that D1-MSN activity and substance P is required for

learning during cue-dependent aversive conditioning and D2-MSNs are required for recall of

conditioned cues. This D2-MSN activity change is likely due to excitatory LTP [22] produced

through muscarinic receptor activation, which would enhance the likelihood of D2-MSN

activation during excitatory input that encodes salient stimuli. Overall, our results provide

evidence for the involvement of the Nucleus Accumbens core in Pavlovian learning that might

support later motivated output.

This work is consistent with previous studies showing MSNs respond to different aspects

of associative reward conditions, where D1-MSNs respond to outcomes and D2-MSNs respond

to associative cues [21]. We found similar outcomes in response to an aversive stimulus, where

D2-MSNs were active in response to a learned cue but not to novel cues and D1-MSNs were

active in response to the shock alone. Together these findings indicate MSN subtypes are

responding to salient events, including rewarding events, further supporting the hypothesis that

the NAc core supports learning about salient cues [1]. Furthermore, the duration of MSN subtype

responses is temporally different in both rewarding [21] and aversive-associated stimuli. D1-MSNs

rapidly respond to the non-associative, salient stimuli and D2-MSNs tend to display extended

activity following associative cue presentation. We speculate D2-MSNs may be actively

engaged in either retaining expression of the learned association or suppressing additional

associations. More work will be needed to test the temporal nature of D1-MSN and D2-MSN

activity in response to associative conditioning and learning.

We provide a role for D1-MSNs and the neuropeptide substance P in mediating aversive

learning. D1-MSNs are thought to facilitate reward; however, the data presented here and data

from others assessing activity in the NAc core [38,39] and NAc shell [40,41] suggest aversive stimuli

activate D1-MSNs. Our data indicates this activity participates in associative learning through

the release of peptides, activation of cholinergic neurons, and release of acetylcholine.

Interestingly, scaling up D1-MSN activity appears to scale acetylcholine release, which is

consistent with increasing substance P levels [22]. While acetylcholine may act as an attentional

signal facilitating the salience of associative stimuli [42,43], we have discovered substance P works

upstream to promote ChAT activation, acetylcholine release, and plasticity on D2-MSNs, which

may be dependent on the concentration of neuromodulator release in response to scaled stress.

This plasticity facilitates information transfer from excitatory brain regions to the NAc core. As

the effect of substance P and acetylcholine plasticity is cue-dependent but not context-

dependent, it is likely that extensive excitatory projections arising from limbic regions that are

involved in Pavlovian conditioning including the basolateral amygdala [44,45] or other regions

involved in aversion such as the paraventricular thalamus [47] may facilitate this D2-MSN activity

during cue recall. In contrast, strong projections to D1-MSNs from the ventral hippocampus [40,49]

may facilitate D1-MSN activity for substance P release. Future experiments will need to assess

what inputs undergo substance P, acetylcholine-dependent potentiation.

Our work shows acetylcholine release plays a significant role in aversive learning and its

release is likely driven by substance P/NK1 receptor signaling. This could result from varying

the upstream activity of D1-MSNs, subsequent substance P release, and the activation of ChAT

neurons. Our previous work has shown that the activity of ChAT cells can be graded by varying

concentration of substance P [22]. We demonstrated D1-MSN activity increases with foot shock

intensity, likely leading to increasing putative substance P release, and this might be a method

of grading acetylcholine release and learning or response to an aversive stimulus. Other factors

are likely contributing to this effect as well. We cannot rule out the role of dopamine input via D2

receptors [46], midbrain GABA input [48], thalamic input [50], intrinsic inhibition [51] or other mechanisms

in generation of these cue-dependent outcomes. In fact, substance P and ChAT acetylcholine is

known to promote dopamine release [52–55]. Dopamine release could accompany the increase in

acetylcholine observed in response to novelty [56] or initial phasic transients of foot shock.

Therefore, selective plasticity could occur due to coordinated acetylcholine and dopamine

release during learning.

Despite a non-habituating D1-MSN calcium signal following repeated foot shocks, the

observed acetylcholine signal habituated to repeated foot shocks. After binding of substance P

to the NK1 receptor and activation, rapid beta-arrestin-mediated endocytosis occurs [57,58] and

attenuates additional responses to substance P for 5-15 min [22,35,57]. At this point, NK1 receptor

activation of ChAT cells would be less sensitive to substance P, thereby decreasing the size of

the signal. Therefore, since we observed the acetylcholine release to be NK1 receptor

dependent, we predict the habituation of the signal from shock 1 to shock 10 is due to NK1

receptor internalization. This, along with the NK1 receptor antagonist attenuation of the

acetylcholine signal, indicates the increase in acetylcholine release is most likely due to release

of substance P and activation of the NK1 receptor on ChAT neurons. Such a mechanism may

suppress the likelihood of additional stimuli promoting plasticity that underlies associative

learning. In addition, the D2-MSN foot shock signal is also rapidly attenuated and much smaller

after repeated shocks. Mice only required two cue shock pairings for associating, perhaps

indicating the plasticity on D2-MSNs occurs quickly, within the first few associations. This may

further suggest that particularly salient stimuli can produce this plasticity rapidly.

One curious aspect of the findings is the activity observed in D1-MSNs, D2-MSNs, and

acetylcholine is induced by foot shock alone. It is likely the release of substance P and

subsequently acetylcholine interacts with other input on D2-MSNs to promote plasticity. This

may include activity from glutamatergic inputs and ventral tegmental area dopamine. In fact,

some evidence suggests the pause in cholinergic neurons is partially mediated by dopamine

input to the striatum [46] and NAc D2-MSN spine dynamics in response to learned cues is

dependent on dopamine [16,51]. Furthermore, coordinated pre-synaptic and post-synaptic activity

via glutamatergic input and muscarinic 1 receptor signaling on D2-MSNs may provide a window

for spike timing dependent plasticity. These secondary sources may also explain the difference

in the role of D2-MSNs in contextual and cue recall, as inhibition of D2-MSNs during

conditioning suppressed context recall as well. These other sources will need to be investigated

in future studies to fully understand the mechanisms that drive these outcomes.

Overall, we demonstrate substance P signaling and plasticity in the NAc core is critical

for learning about salient, aversive cues. We uncovered a new role of D1-MSN local

communication to facilitate acetylcholine plasticity on D2-MSNs that is relevant to cue-

dependent aversive learning. Enhanced processing of salient stimuli, through this substance P,

D2-MSN mechanism could lead to exacerbated response to aversive stimuli and could enhance

susceptibility to aversive stimuli that cause stress or anxiety disorders. Therefore, targeting

these neurobiological mechanisms may be useful in therapies aiming to treat anxiety and mood

disorders.

Acknowledgements

We would like to thank Dr. Geoffrey Schoenbaum for his assistance in conceptualization

of the fear conditioning and latent inhibition experiments. This work was supported by that

National Institutes of Mental Health R00MH123673, University of South Carolina College of

Pharmacy, and the National Institute on Drug Abuse Intramural Research Program.

Construction and packaging of the AAV vector expressing mTACR1 gRNAs was performed by

the National Institute on Drug Abuse Genetic Engineering and Viral Vector Core,

RRID:SCR_022969.

Author contributions

Conceptualization, T.C.F.; Methodology and Investigation, A.B., C.G., C.S., C.R., D.B., E.M.,

A.S., H.H., and T.C.F.; Writing, T.C.F.; Review and Editing, A.B., C.G., A.S., C.R., and T.C.F.;

Supervision, T.C.F.

Declaration of Interests

The authors declare no competing interests.

Main Text Figure Legends

Figure 1. NAc substance P signaling is required for cue-dependent aversive learning. (A)

Fear conditioning paradigm for aversive learning. **(B)** Injection of the NK1 receptor antagonist

L-733,060 (10 mg/kg) or saline was given prior to conditioning. NK1 receptor antagonist

suppressed freezing in response to the cue during cue recall ($p < 0.001$) and no difference in

freezing was observed in the context ($p > 0.05$) (n = 20, 22 mice). **(C)** L-733,060 prior to cue

recall does not change the percent of time freezing to conditioned cues ($p > 0.05$) (n = 7, 8). **(D)**

Fluorescent image of EGFP-KASH expression in the NAc core. Scale bar 100 µm. Insert:

Expression of EGFP-KASH (green), *Chat* RNA (red), *Tacr1* RNA (grey/white), and DAPI (blue).

CRISPR-Cas9 mediated knockout of *Tacr1* decreased measured *Tacr1* RNA puncta in *Chat*

RNA positive cells ($p < 0.001$) but not *Chat* RNA negative cells ($p > 0.05$) (n = 4 mice per

group). **(E)** Graph and representative traces showing Cas9 and *Tacr1*-gRNA was required to

suppress substance P(SP)-dependent activation of ChAT cells (Interaction $p<0.0001$; post-hoc

Baseline vs. SP ChAT-Cre/Cas9^{Tacr1} $p > 0.05$, Mice/cells(N/n) = 4/10; Baseline vs. SP ChAT-

Cre^{Tacr1}, ChAT-Cre/Cas9, or ChAT-Cre $p < 0.0001$) (N/n = 4/9, 4/8, 3/7). **(F)** *Tacr1* knockout

decreased freezing to the cue ($p < 0.001$), but not the conditioning context ($p > 0.05$) (n = 14, 15

mice). **(G)** Local NAc infusion of L-733,060 during conditioning suppressed freezing to

conditioned cues ($p < 0.01$) but not the conditioned context ($p > 0.05$) (n = 7, 8). Data are

represented as mean ± SEM. For detailed statistics see **Table S1**. See also **Figure S1**.

Figure 2. D1-MSN activity is required for cue-dependent conditioning. (A) Representation of a Dyn-Cre mouse injected with the calcium indicator GCaMP7f and a representative image of its expression in the NAc core with a fiber track. Scale bar 200 µm. **(B)** Heat maps and time plots representing average z-scores of D1-MSN calcium-driven fluorescence over increasing foot shock intensity (0.1-1.0 mA). **(C)** AUC from GCaMP7f fluorescent signals increased with shock intensity (Interaction $p < 0.0001$, n = 8-10 mice per shock intensity). **(D)** No difference was observed across individual shock trials at 0.6 mA ($p > 0.05$, n = 10 mice). **(E)** Averaged z-score D1-MSN calcium activity during fear conditioning. **(F)** Average z-score D1-MSN calcium activity in response to the condition cue during cue recall. **(G)** Conditioning foot shock intensity increasingly enhances freezing to the cue during recall ($p < 0.0001$). **(H)** AUC of the D1-MSN GCaMP7f signal significantly correlates with freezing to the cue during recall ($p < 0.01$) (n = 5-13 mice per group) **(I)** Expression of eNpHR3.0-eYFP, fiber track for optogenetic inhibition of the NAc core, and cartoon displaying bilateral implant and expression of the optogenetic construct in mice. **(J)** Inhibition of D1-MSNs during the cue in conditioning suppressed later cue recall ($p < 0.01$) but did not affect context recall ($p > 0.05$) (n = 11, 10 mice). **(K)** Halorhodopsin inhibition of D1-MSNs during cue recall had no effect on subsequent context recall. **(L)** Inhibition of D1-MSNs during cue recall had no effect on freezing (Laser off vs. on, $p > 0.05$) (n = 9, 10 mice). Data are represented as mean ± SEM. For detailed statistics see **Table S1**. See also **Figure S2**.

Figure 3. Acetylcholine release during foot shock is substance P-dependent. (A) Representative image of GRAB-ACh3.0 expression in the NAc core and cartoon of fiber placement. **(B)** Heatmaps and time courses of fluorescent GRAB-ACh3.0 signals from 10 random foot shocks at intensities of 0.1-1.0 mA amplitudes. Time courses are taken from average z-scores from shock 1 and shock 10. **(C and D)** AUC measurement of acetylcholine

release at varying shock intensities for shock 1 and shock 10. **(C)** No differences were observed

during the AUC (inverse to make positive) during pauses in acetylcholine-mediated GRAB-

ACh3.0 fluorescence across shock intensities or repeated shocks ($p > 0.05$). **(D)**. Acetylcholine

release varied across both AUC and shock intensity during the post shock rebound (Cue 1 vs.

Cue 10, $p < 0.05$) (n = 9 mice). **(E)** Heat map and time course of GRAB-ACh3.0 fluorescent

response after L-733,060 administration. **(F)** L-733,060 had no effect on the size of the pause

across shocks ($p > 0.05$). **(G)** NK1 receptor antagonism suppressed AUC of the acetylcholine

signal varied across both drug treatment (AUC Saline vs. L-733,060 during Cue 1, $p < 0.05$) and

shock presentation (Cue 1 vs. Cue 10, $p < 0.05$) (n = 5, 6 mice). **(H-I)** Phasic changes in GRAB-

ACh3.0 fluorescence during aversive cue conditioning in saline and L-733,060 treated mice. **(J)**

L-733,060 significantly decreases the rebound AUC GRAB-ACh3.0 fluorescence during cue

conditioning ($p < 0.05$, n = 8, 12). **(K)** The AUC during the rebound in conditioning significantly

correlated with percent freezing to the cue during recall ($p < 0.01$, n = 12 mice). Data are

represented as mean ± SEM. For detailed statistics see **Table S1**. See also **Figure S3**.

Figure 4. ChAT-mediated cholinergic signaling is required for cue-dependent aversive

learning. (A) Expression of halorhodopsin in ChAT cells in the NAc core. Schematic showing

inhibition paradigms: during the cue in conditioning or after the shock and cue during

conditioning. **(B)** ChAT inhibition during the cue had no effect on context or cue recall ($p > 0.05$,

n = 6 mice per group). **(C)** Inhibition of ChAT cells following the cue and shock during

conditioning significantly decreased freezing to the cue during recall ($p < 0.001$), but did not

affect context recall ($p > 0.05$) (n = 5, 9). **(D)** Selective infusion of the muscarinic 1 receptor

antagonist pirenzepine (bilateral) prior to conditioning suppressed cue-dependent ($p < 0.01$) but

not conditioning context dependent freezing ($p > 0.05$) (n = 10, 9). **(E)** Infusion of pirenzepine

prior to cue recall had no effect on freezing ($p > 0.05$) in mice showing similar context recall (n =

6, 7). Data are represented as mean ± SEM. For detailed statistics see **Table S1**. See also

Figure S4.

Figure 5. Excitatory plasticity on D2-MSNs is caused by cue-dependent aversive learning

(A) Cue/shock conditioning enhanced inward rectification of the I-V curve generated from

electrically evoke excitatory currents on D2-MSNs (Interaction $p < 0.001$) from the cue/shock

group compared to cue only (post-hoc $p < 0.05$) and cue/shock with L-733,060 (post-hoc $p <$

0.05). Rectification index indicated more rectification for D2-MSNs from cue/shock mice

(Interaction, $p < 0.01$) compared to cue only (post-hoc, $p < 0.01$) and cue/shock with L-733,060

(post-hoc, $p < 0.05$) (N/n (Mice/cells) = 5/10, 5/10, 4/7). **(B)** NASPM suppressed currents more

from D2-MSNs from cue/shock mice compared to cue only mice (25-30 min average, $p < 0.05$)

(N/n = 4/6, 4/5). **(C)** Spontaneous excitatory post-synaptic current (sEPSC) amplitude ($p <$

0.05), but not frequency ($p > 0.05$) is increased in D2-MSNs following fear conditioning (N/n =

6/14, 5/9, 4/7). **(D)** D1-MSNs do not show inward rectification across conditions ($p > 0.05$) (N/n

= 6/8, 4/9, 4/5). Data are represented as mean ± SEM. For detailed statistics see **Table S1**. See

also **Figure S5**.

Figure 6. D2-MSNs are required for cue-dependent aversive learning. (A) A cartoon of an

A2a-Cre mouse injected with the calcium indicator GCaMP7f and a representative image of its

expression in the NAc core with a fiber track. Scale bar 200 µm. **(B)** Heat maps and time plots

representing average z-scores of D2-MSN calcium-driven fluorescence over increasing foot

shock intensity (0.1-1.0 mA). **(C)** AUC of an average of all 10 foot shocks indicated was no

change in response across foot shock intensity ($p > 0.05$). **(D)** AUC fluorescence from D2-MSN

GCaMP7f does not change across all foot shock trials at 0.6 mA intensity ($p > 0.05$,) (n = 10

mice). **(E)** D2-MSNs responded to the first conditioning shock, but not the cue. **(F)** Following

conditioning, calcium signals from D2-MSNs were responsive to the conditioned cue. **(G)** D2-

MSN calcium signals responded to the foot shock during conditioning after being injected with L-

733,060. **(H)** Cue-driven calcium transients from D2-MSNs were not observed in mice injected

with L-733,060. **(I)**   AUC fluorescent transients during cue recall from mice expressing

GCaMP7f in D2-MSNs were larger than in mice expressing GCaMP7f in D1-MSNs ($p < 0.01$)

and in mice injected with L-733,060 expressing GCaMP7f in D2-MSNs ($p < 0.05$) (n = 11, 8,

8).**(J)** AUC during the cue significantly correlated with freezing to the cue ($p < 0.05$, n = 8, 8).

(K) eNpHR3.0-eYFP expression in D2-MSNs within the NAc core and cartoon depicting bilateral

inhibition of the NAc core. **(L)** Inhibition of D2-MSNs during the cue in conditioning decreased

freezing during context recall ($p < 0.05$). **(M)**   Inhibition of D2-MSNs reversibly suppressed

freezing to recalled cues (Over cues $p < 0.01$; Laser off vs on $p < 0.01$; post-hoc laser on eYFP

vs. eNpHR3.0 $p < 0.01$) (n = 8, 8). **(N-O)** Inhibition of D2-MSNs reversibly suppressed freezing

to recalled cues (Over cues $p < 0.01$; Laser off vs on $p < 0.0001$; post-hoc laser on eYFP vs.

eNpHR3.0 $p < 0.05$) in mice that showed no difference in context recall ($p < 0.05$) (n = 9, 6

mice). Data are represented as mean ± SEM. For detailed statistics see **Table S1**. See also

Figure S6.

Figure 7. Model for the role of the NAc core in cue-dependent fear learning. NAc circuitry

involved in aversive cue learning (left). Shock activates D1-MSNs releasing substance P and

activating ChAT neurons. ChAT neurons release acetylcholine selectively during cue/shock

pairings and promotes glutamatergic long-term potentiation (LTP) on D2-MSNs rendering them

more responsive to the cue. Timing of putative activity and release of neurotransmitters from

NAc neuronal subtypes during cue conditioning and recall (right).

STAR Methods

RESOURCE AVAILABILITY

Lead contact

Further information and requests or resources and reagents should be directed to and will be

fulfilled by the lead contact, T. Chase Francis (francit@mailbox.sc.edu).

Materials availability

Plasmids generated in this study have been deposited to Addgene (pOTTC1553, Cat #:

131682; pOTTC1642, Cat #: 195018) and are publicly available as of the date of publication. All

data reported in this paper will be shared by the lead contact upon request and can be found on

the Open Science Framework platform (DOI 10.17605/OSF.IO/4FE3M).

Data and code availability

Python code generated to extract calcium and acetylcholine events is available through

Github.com (https://github.com/FrancisLab-Repository/Photometry-Event-Extract.git). In some

experiments, ezTrack was used to quantify freezing behavior or locomotion [61].

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Animals

Female and male mice 10-15 weeks old were used in all studies. For GRAB-ACh3.0

virus, cannula infusion experiments, and systemic drug injections without imaging or

electrophysiology, C57Bl6/J mice (JAX: 000664) were used. For optogenetic inhibition

experiments either B6;129S-Pdyntm1.1(cre)Mjkr/LowlJ (Dyn-Cre; JAX: 027958) or Tg(Adora2a-

cre)KG139Gsat/Mmucd (A2a-Cre; MMRRC: 031168-UCD). For CRISPR-Cas9 experiments,

B6;129-Gt(ROSA)26Sortm1(CAG-cas9*,-EGFP)Fezh/J (LSL-Cas9; JAX: 024857) mice were

crossed with B6;129S6-Chattm2(cre)Lowl/J (ChAT-Cre) mice for conditional expression of Cas9

in ChAT-Cre cells. All transgenic animals were bred in house. Mice were fed *ad libitum* and

were housed in a reverse light/dark (12/12 hr). Experimental procedures were approved and

conducted in accordance with NIDA-IRP ACUC, University of South Carolina IACUC, and NIH

guidelines.

METHOD DETAILS

Surgical methods

Mice used in fluorescence imaging, optogenetic, or CRISPR knockdown experiments were anesthetized with ketamine/xylazine for intracranial AAV injection, hair was removed, scalps were cleaned with ethanol and betadine, and lidocaine was used prior to scalp incision. Body temperature was maintained with an electric heating pad. Mice received bilateral stereotactic injections (0.3-0.5 µL) of either AAV9-hsyn-GRAB-ACh3.0 (Addgene; plasmid# 121922; packaged by Vigene Biosciences)[36], AAV9-syn-FLEX-jGCaMP7f (Addgene; 104492-AAV9), AAV5-EF1a-DIO-eNpHR3.0-eYFP (Addgene; 26966-AAV5), or AAV1-mU6-Tac1r-gRNA1::EF1a-NucEnv-EGFP::hU6-Tacr1-gRNA2 in the NAc core (created and packaged by NIDA GEVVC Core) (A/P: 1.6, Lat: 1.6, D/V: -4.4, 10° angle). For photometry or optogenetics, mice were then implanted with either a 400 µm photometry fiber (Doric) or 200 µm core optogenetic fiber fixed with head screws and dental cement. Following surgery, mice were injected with warm saline to replenish fluids and given carprofen (5 mg/kg) for pain relief during recovery.

In vivo **fiber photometry,** *ex vivo* **imaging, and** *in vivo* **optogenetics**

In vivo fiber photometry was used to assess cell-type specific calcium activity (GCaMP7f) or acetylcholine release (GRAB-ACh3.0) using a Tucker-Davis Technologies LUX RZ10 fiber photometry system. Signals were tested and mice were habituated for 3-5 days prior behavioral recordings. For all recordings, sensors were excited at 470 nm to acquire the fluorescent emission signal and the isosbestic point for GFP at 405 nm as a normalization control. Excitation signals were generated by two light-emitting diodes and emission signals were collimated through a Doric fluorescent mini cube before being processed and digitized using the Synapse software suite. Analysis was performed offline using a custom Python script

which extracted fluorescent signals, smoothed the signal with a linear fit to the 405 nm signal

and applying the regression to the 470 nm signal, calculated $\Delta F/F$, aligned signals based on a

timestamp, extracted fluorescent signals around events, and normalized signals (z-score) based

on the 10 seconds prior to an event. Area Under the Curve (AUC) was assessed to find mean

changes in calcium or acetylcholine activity. The following intervals were used for AUC

calculation in the described experiment: for foot shock during fear conditioning or during shock

alone, 0-5 sec from the initiation of the shock; for cue recall, 0-10 sec from the start of the cue;

for shock-related decrease in ACh-mediated fluorescence, 0.5-2 sec after from the initiation of

the shock; for shock-related rebound in ACh-mediated fluorescence, 2.5 to 7 sec from the

initiation of the shock.

For *ex vivo* imaging of GRAB-ACh3.0, slices were prepared as discussed in the

electrophysiology methods section. Acetylcholine (1 or 100 µM) was bath applied after a 2 min

baseline. Videos were acquired with a 40x objective at a 10 Hz frame rate using QCapturePro7

and analyzed offline with ImageJ. A subregion 100x100 µm devoid of fiber tracts in the NAc

core was chosen in each image and analyzed for fluorescence intensity over time. Fluorescence

traces were flattened using a linear regression fit and normalized (z-score) to the 1 min prior to

acetylcholine bath application.

For optogenetic inhibition experiments, mice were habituated to patch cables 5 days

prior to behavior. Percent transmission was calculated during creation of optogenetic fibers and

594 nm light was delivered at 3-5 mW from a DPSS laser (OptoEngine LLC). Optogenetic

inhibition was delivered during the onset of the cue and one second after the offset of the cue

for all conditioning cues and on every other cue during cue recall.

Behavior

Pavlovian fear conditioning was conducted over a 3 day period (**Figure 1A**) in Med

Associates Fear conditioning chambers contained in sound attenuating cabinets. Prior to each

phase of fear conditioning, mice were habituated in a separate room for at least 1 hour in the

lighting condition of the test. Conditioning was conducted in white light with vanilla scent (10%

vanilla extract in ethanol) chamber with a grid floor. Following a 2 min baseline, 2 white-noise

(80 dB) cues co-terminating in a 0.6 mA foot shock separated by a 30 sec interval. The next day

mice were exposed to the conditioning context for 5 min. On the third day, mice were placed in

a new context where floors were covered with white, opaque plexiglass, a curved white

plexiglass insert, no light (red light in behavior room), and vinegar scent (1% acetic acid in

water). After a 2 min baseline, 5-6 cues at 30 sec intervals were played back. In all tests,

freezing was measured using VideoFreeze (Med Associates), which has been thoroughly

validated in relation to hand scoring [59]. Freezing thresholds were set at least 30, 0.033 sec

frames of frozen motion and motion indices below 6 for mice without fiber or optogenetic tethers

and below 500 for mice with optical tethers. For mice exposed to foot shock only, after a 2 min

baseline each mouse was exposed to an unpredictable foot shock set to random intervals at

least 30 sec between each shock. Mice received one intensity of foot shock per day, and each

subsequent day were subjected to increasing foot shock intensities or decreasing foot shock

intensities to control for previous foot shock intensities from interfering with the signal for any

subsequent foot shock.

Electrophysiology

Whole-cell electrophysiological recordings were done one day following cue recall. Mice

were anesthetized with Euthasol, brains were extracted, and cut (300 µM) in ice cold N-Methyl

D-Glucamine (NMDG) cutting artificial cerebrospinal fluid (ACSF) solution (in mM): 92 NMDG,

20 HEPES, 25 glucose, 30 sodium bicarbonate, 1.2 sodium phosphate, 2.5 potassium chloride,

5 sodium ascorbate, 3 sodium pyruvate, 2 thiourea, 10 magnesium, 0.5 calcium chloride,

osmolarity 305-310 mOsm, pH 7.34. Immediately after slicing, slices were placed in warm

(32°C) NMDG solution for 2-5 min, then transferred to HEPES holding solution at room

temperature which was the same as NMDG solution except NMDG was replaced with 92 mM

sodium chloride and contained 1 mM magnesium chloride and 2 mM calcium chloride.

Recording began 1 hr after brain slicing was complete and was performed at 33°C. Slices were

perfused at a rate of 2 mL/min with ACSF (in mM): 125 sodium chloride, 26 sodium bicarbonate,

and 11 glucose, 2.5 potassium chloride, 1.25 sodium phosphate, 2.4 calcium chloride, 1

magnesium chloride, osmolarity 305-310 mOsm, pH 7.34. To block inhibitory currents, synaptic

recordings were done holding cells at -70 mV in the presence of picrotoxin (100 µM) and with a

cesium methanesulfonate internal solution (in mM): 117 cesium methanesulfanate, 20 HEPES,

2.8 sodium chloride, 4 magnesium ATP, 0.3 sodium GTP, 0.4 EGTA, pH 7.30, osmolarity 280-

287 mOsm. For IV curves, spermine (100 mM) was added to the pipette solution and currents

were recorded in the presence of APV to isolate AMPA receptor currents. NASPM (100 µM)

was used to isolate currents from inserted GluR2-lacking AMPA receptors. For cell-attached

ChAT recordings, ACSF was used as the internal solution.

Whole-cell currents or cell attached currents were obtained with borosilicate glass

pipettes (whole cell: 1.7-3.0 MΩ; cell attached: 3.0-4.0 MΩ) and signals were amplified and

digitized using a Multiclamp 700B amplifier (4kHz low-pass filter for synaptic, 1kHz for cell-

attached or spontaneous EPSCs) and Digidata 1322 digitizer (10 kHz), respectively (Molecular

Devices). Synaptic currents were evoked with a concentric bipolar stimulating electrode (FHC)

and evoked currents were stabilized for 5-10 min prior to data acquisition. Series resistance

drifts of more than 20% led to exclusion of the cell from analysis. D1- and D2-MSNs were

identified under the visual guidance of a BX61 Olympus microscope and Tdtomato

fluorescence. Cholinergic neurons were identified by cell size (40-60 µm diameters) and high

baseline firing rates.

CRISPR-Cas9 for *Tacr1* knockout

The SpCas9 guide RNA seed sequences within the first exon of mouse *TACR1* were identified and scored using the Guide Design Tool at CRISPR.mit.edu. Two gRNAs (gRNA 1 A55: CCGTATAGGCGGCTGCCCAA, and gRNA 2 B108: TTCCGTGGTGGGCAACGTAG) were chosen based on their likelihood for functioning as a "nickase-compatible" pair. These sequences were inserted into plasmids already containing RNA pol III promoters and SpCas9 gRNA scaffold sequences. (gifts from Charles Gersbach; Addgene plasmid # 53187; http://n2t.net/addgene:53187; RRID:Addgene_53187, Addgene plasmid # 53188; http://n2t.net/addgene:53188; RRID:Addgene_53188). Then each intact gRNA expression cassette was amplified by PCR and both were inserted into an AAV expression vector that contained a EGFP-KASH reporter (pOTTC1553, Addgene #131682), resulting in plasmid pOTTC1642 (Addgene #195018). All cloning reactions used PCR amplification and ligation-independent cloning (In-Fusion, Takara Inc.). Reactions involving non-AAV plasmids were transformed into Stellar competent cells (Takara Inc.) and those involving AAV plasmids were transformed into NEB Stable competent cells (New England Biolabs).

The pOTTC1642 viral vector was packaged by triple transfection along with pHelper and rep/cap plasmids (serotype 1) and purified by affinity chromatography on GE AVB columns [60]. Purified vectors were titered by droplet digital PCR with an assay that recognizes the EF1a reporter. Titered vectors were tested for GFP expression by transduction into rat primary cortical neurons prior to *in vivo* use.

Immunohistochemistry and *in situ* hybridization

Expression of GRAB-ACh3.0 or eNpHR3.0-eYFP was identified by histological staining for GFP. Briefly, brains slices (40 μm) from perfused (4% paraformaldehyde) brains were washed with 1X PBS, blocked with 1X PBS buffer containing 0.01% Triton X, 10% Normal

Donkey Serum and incubated with a primary anti-body for GFP made in chicken (Aves lab, cat#:

GFP-1020). Slices were washed and incubated with a secondary anti-chicken antibody made in

donkey conjugated to Alexa Fluor 488 (Jackson immuno, cat#: 703-545-155), stained with DAPI

and mounted on slides. All images were acquired on a Fluoview 1000 BX51WI confocal

microscope (Olympus).

In situ hybridization was performed on fresh frozen 16 μm slices according to the

ACDBio specifications as described previously [22]. EGFP-KASH maintained its fluorescence

through the protocol. The probes used were as follows: *Chat* (Mm-Chat-C1 (nucleotide target

region 1090-1952), GenBank #: NM_009891.2); *Tacr1* (Mm-Tacr1-C3 (nucleotide target region

845-1775), GenBank #: NM_009313.5). All images were taken within a week of the protocol and

image intensities were maintained across slices and slides. Quantification was done in the NAc

core, where EGFP-KASH expression was found. Due to the cre-independent expression of

EGFP-KASH and expression of GFP in ChAT cells, we were unable to specifically identify cells

with or without expression and, therefore, all *Chat* positive cells in the expression region were

analyzed. RNA was quantified using ImageJ by selecting 10 individual pixel values for every

image and averaging to find the intensity values of a single RNA puncta. Intensity values of

Tacr1 in ChAT positive cells were assessed by selecting perinuclear RNA, quantifying intensity,

and dividing the total intensity by the intensity of a single puncta value to calculate total RNA

puncta. A total of 4 slices from each animal was taken and averaged. Slice averages are

represented as individual points.

QUANTIFICATION AND STATISTICAL ANALYSIS

Detailed statistics can be found in **Table S1**. Data was analyzed and statistics were

calculated using Graphpad Prism. Tests were considered significant if $p < 0.05$. For two-sample

data, two-sided, un-paired or paired *t*-tests were used. For more than 2 variables over repeated

time points one-way or two-way repeated measure (RM) ANOVAs were used followed by post-

697 hoc tests using Sidak's multiple comparison test. In most cases with an interaction, main effects

698 were reported. For electrophysiology experiments, all are represented as N/n (Mice/cells).

699 Animal exclusions for behavioral experiments were made only when viral expression was

700 absent unilaterally or bilaterally following histological staining. All data is represented as Mean ±

701 standard error. Cartoons were made using BioRender during the subscription period.

703 **References**

704 1. Zaehle, T., Bauch, E.M., Hinrichs, H., Schmitt, F.C., Heinze, H.-J., Voges, J., Heinze, H.-
705   J., and Bunzeck, N. (2013). Nucleus Accumbens Activity Dissociates Different Forms of
706   Salience: Evidence from Human Intracranial Recordings. The Journal of Neuroscience
707   *33*, 8764 LP – 8771. 10.1523/JNEUROSCI.5276-12.2013.

708 2. Schoenbaum, G., and Setlow, B. (2003). Lesions of Nucleus Accumbens Disrupt
709   Learning about Aversive Outcomes. Journal of Neuroscience *23*, 9833–9841.

710 3. Dutta, S., Beaver, J., Halcomb, C.J., and Jasnow, A.M. (2021). Dissociable roles of the
711   nucleus accumbens core and shell subregions in the expression and extinction of
712   conditioned fear. Neurobiol Stress *15*, 100365.
713   https://doi.org/10.1016/j.ynstr.2021.100365.

714 4. Wendler, E., Gaspar, J.C.C., Ferreira, T.L., Barbiero, J.K., Andreatini, R., Vital, M.A.B.F.,
715   Blaha, C.D., Winn, P., and Da, C. (2014). Neurobiology of Learning and Memory The
716   roles of the nucleus accumbens core , dorsomedial striatum , and dorsolateral striatum in
717   learning□: Performance and extinction of Pavlovian fear-conditioned responses and
718   instrumental avoidance responses. Neurobiol Learn Mem *109*, 27–36.
719   10.1016/j.nlm.2013.11.009.

720 5. Ray, M.H., Russ, A.N., Walker, R.A., and McDannald, M.A. (2020). The Nucleus
721   Accumbens Core is Necessary to Scale Fear to Degree of Threat. The Journal of
722   Neuroscience *40*, 4750 LP – 4760. 10.1523/JNEUROSCI.0299-20.2020.

723 6. Francis, T.C., and Lobo, M.K. (2017). Emerging Role for Nucleus Accumbens Medium
724   Spiny Neuron Subtypes in Depression. Biol Psychiatry *81*, 645–653.
725   10.1016/j.biopsych.2016.09.007.

726 7. Lee, H.J., Weitz, A.J., Bernal-Casas, D., Duffy, B.A., Choy, M.K., Kravitz, A. v., Kreitzer,
727   A.C., and Lee, J.H. (2016). Activation of Direct and Indirect Pathway Medium Spiny
728   Neurons Drives Distinct Brain-wide Responses. Neuron *91*, 412–424.
729   10.1016/j.neuron.2016.06.010.

8.   Kravitz, A. v, Tye, L.D., and Kreitzer, A.C. (2012). Distinct roles for direct and indirect
pathway striatal neurons in reinforcement. Nat Neurosci *15*, 816–818. 10.1038/nn.3100;
10.1038/nn.3100.

9.   Francis, T.C., Chandra, R., Friend, D.M., Finkel, E., Dayrit, G., Miranda, J., Brooks, J.M.,
Iniguez, S.D., O'Donnell, P., Kravitz, A., et al. (2015). Nucleus accumbens medium spiny
neuron subtypes mediate depression-related outcomes to social defeat stress. Biol
Psychiatry *77*, 212–222. 10.1016/j.biopsych.2014.07.021.

10.   Smith, R.J., Lobo, M.K., Spencer, S., and Kalivas, P.W. (2013). Cocaine-induced
adaptations in D1 and D2 accumbens projection neurons (a dichotomy not necessarily
synonymous with direct and indirect pathways). Curr Opin Neurobiol *23*, 546–552.
10.1016/j.conb.2013.01.026.

11.   Lobo, M.K., Covington 3rd, H.E., Chaudhury, D., Friedman, A.K., Sun, H., Damez-Werno,
D., Dietz, D.M., Zaman, S., Koo, J.W., Kennedy, P.J., et al. (2010). Cell type-specific loss
of BDNF signaling mimics optogenetic control of cocaine reward. Science *330*, 385–390.
10.1126/science.1188472.

12.   Ferguson, S.M., Phillips, P.E., Roth, B.L., Wess, J., and Neumaier, J.F. (2013). Direct-
pathway striatal neurons regulate the retention of decision-making strategies. J Neurosci
33, 11668–11676. 10.1523/JNEUROSCI.4783-12.2013; 10.1523/JNEUROSCI.4783-
12.2013.

13.   Tai, L.H., Lee, A.M., Benavidez, N., Bonci, A., and Wilbrecht, L. (2012). Transient
stimulation of distinct subpopulations of striatal neurons mimics changes in action value.
Nat Neurosci *15*, 1281–1289. 10.1038/nn.3188; 10.1038/nn.3188.

14.   Bock, R., Shin, J.H., Kaplan, A.R., Dobi, A., Markey, E., Kramer, P.F., Gremel, C.M.,
Christensen, C.H., Adrover, M.F., and Alvarez, V.A. (2013). Strengthening the accumbal
indirect pathway promotes resilience to compulsive cocaine use. Nat Neurosci *16*, 632–
638. 10.1038/nn.3369; 10.1038/nn.3369.

15.   Calipari, E.S., Bagot, R.C., Purushothaman, I., Davidson, T.J., Yorgason, J.T., Pena,
C.J., Walker, D.M., Pirpinias, S.T., Guise, K.G., Ramakrishnan, C., et al. (2016). In vivo
imaging identifies temporal signature of D1 and D2 medium spiny neurons in cocaine
reward. Proc Natl Acad Sci U S A *113*, 2726–2731. 10.1073/pnas.1521238113 [doi].

16.   Iino, Y., Sawada, T., Yamaguchi, K., Tajiri, M., Ishii, S., Kasai, H., and Yagishita, S.
(2020). Dopamine D2 receptors in discrimination learning and spine enlargement. Nature
579, 555–560. 10.1038/s41586-020-2115-1.

17.   MacPherson, T., Morita, M., Wang, Y., Sasaoka, T., Sawa, A., and Hikida, T. (2016).
Nucleus accumbens dopamine D2-receptor expressing neurons control behavioral
flexibility in a place discrimination task in the IntelliCage. Learning and Memory *23*, 359–
364. 10.1101/lm.042507.116.

18.   Macpherson, T., Kim, J.Y., and Hikida, T. (2022). Nucleus Accumbens Core Dopamine
D2 Receptor-Expressing Neurons Control Reversal Learning but Not Set-Shifting in
Behavioral Flexibility in Male Mice. Front Neurosci *16*, 885380.
10.3389/fnins.2022.885380.

19.   Soares-Cunha, C., Coimbra, B., Domingues, A.V., Vasconcelos, N., Sousa, N., and
Rodrigues, A.J. (2018). Nucleus Accumbens Microcircuit Underlying D2-MSN-Driven
Increase in Motivation. eNeuro *5*, ENEURO.0386-18.2018. 10.1523/ENEURO.0386-
18.2018.

20.   Soares-Cunha, C., Coimbra, B., David-Pereira, A., Borges, S., Pinto, L., Costa, P.,
Sousa, N., and Rodrigues, A.J. (2016). Activation of D2 dopamine receptor-expressing
neurons in the nucleus accumbens increases motivation. Nat Commun *7*, 11829.
10.1038/ncomms11829 [doi].

21.   Owesson-White, C., Belle, A.M., Herr, N.R., Peele, J.L., Gowrishankar, P., Carelli, R.M.,
and Wightman, R.M. (2016). Cue-Evoked Dopamine Release Rapidly Modulates D2
Neurons in the Nucleus Accumbens During Motivated Behavior. J Neurosci *36*, 6011–
6021. 10.1523/JNEUROSCI.0393-16.2016.

22.   Francis, T.C., Yano, H., Demarest, T.G., Shen, H., and Bonci, A. (2019). High-Frequency
Activation of Nucleus Accumbens D1-MSNs Drives Excitatory Potentiation on D2-MSNs.
Neuron *103*. 10.1016/j.neuron.2019.05.031.

23.   Commons, K.G. (2010). Neuronal pathways linking substance P to drug addiction and
stress. Brain Res *1314*, 175–182. 10.1016/j.brainres.2009.11.014.

24.   Nakamura, Y., Izumi, H., Shimizu, T., Hisaoka-Nakashima, K., Morioka, N., and Nakata,
Y. (2013). Volume Transmission of Substance P in Striatum Induced by Intraplantar
Formalin Injection Attenuates Nociceptive Responses via Activation of the Neurokinin 1
Receptor. J Pharmacol Sci *121*, 257–271. 10.1254/jphs.12218FP.

25.   Aosaki, T., Tsubokawa, H., Ishida, A., Watanabe, K., Graybiel, A.M., and Kimura, M.
(1994). Responses of tonically active neurons in the primate's striatum undergo
systematic changes during behavioral sensorimotor conditioning. J Neurosci *14*, 3969–
3984. 10.1523/JNEUROSCI.14-06-03969.1994.

26.   Ravel, S., Legallet, E., and Apicella, P. (2003). Responses of Tonically Active Neurons in
the Monkey Striatum Discriminate between Motivationally Opposing Stimuli. The Journal
of Neuroscience *23*, 8489 LP – 8497. 10.1523/JNEUROSCI.23-24-08489.2003.

27.   Schulz, J.M., Pitcher, T.L., Savanthrapadian, S., Wickens, J.R., Oswald, M.J., and
Reynolds, J.N.J. (2011). Enhanced high-frequency membrane potential fluctuations
control spike output in striatal fast-spiking interneurones in vivo. J Physiol *589*, 4365–
4381. 10.1113/jphysiol.2011.212944.

28. Aoki, S., Liu, A.W., Zucca, A., Zucca, S., and Wickens, J.R. (2015). Role of Striatal
Cholinergic Interneurons in Set-Shifting in the Rat. The Journal of Neuroscience *35*, 9424
LP – 9431. 10.1523/JNEUROSCI.0490-15.2015.

29. Okada, K., Nishizawa, K., Fukabori, R., Kai, N., Shiota, A., Ueda, M., Tsutsui, Y., Sakata,
S., Matsushita, N., and Kobayashi, K. (2014). Enhanced flexibility of place discrimination
learning by targeting striatal cholinergic interneurons. Nat Commun *5*, 3778.
10.1038/ncomms4778.

30. Ragozzino, M.E., Mohler, E.G., Prior, M., Palencia, C.A., and Rozman, S. (2009).
Acetylcholine activity in selective striatal regions supports behavioral flexibility. Neurobiol
Learn Mem *91*, 13–22. 10.1016/j.nlm.2008.09.008.

31. Collins, A.L., Aitken, T.J., Huang, I.-W., Shieh, C., Greenfield, V.Y., Monbouquette, H.G.,
Ostlund, S.B., and Wassum, K.M. (2019). Nucleus Accumbens Cholinergic Interneurons
Oppose Cue-Motivated Behavior. Biol Psychiatry *86*, 388–396.
10.1016/j.biopsych.2019.02.014.

32. Sandweiss, A.J., and Vanderah, T.W. (2015). The pharmacology of neurokinin receptors
in addiction: prospects for therapy. Subst Abuse Rehabil *6*, 93–102.
10.2147/SAR.S70350.

33. Tomaz, C., and Nogueira, P.J.C. (1997). Facilitation of memory by peripheral
administration of substance P. Behavioural Brain Research *83*, 143–145.
https://doi.org/10.1016/S0166-4328(97)86058-5.

34. Ebner, K., and Singewald, N. (2006). The role of substance P in stress and anxiety
responses. Amino Acids *31*, 251–272. 10.1007/s00726-006-0335-9.

35. Aosaki, T., and Kawaguchi, Y. (1996). Actions of substance P on rat neostriatal neurons
in vitro. The Journal of neuroscience *16*, 5141–5153.

36. Jing, M., Li, Y., Zeng, J., Huang, P., Skirzewski, M., Kljakic, O., Peng, W., Qian, T., Tan,
K., Zou, J., et al. (2020). An optimized acetylcholine sensor for monitoring in vivo
cholinergic activity. Nat Methods *17*, 1139–1146. 10.1038/s41592-020-0953-2.

37. Lubow, R.E. (1973). Latent inhibition. Psychol Bull *79*, 398.

38. Soares-Cunha, C., de Vasconcelos, N.A.P., Coimbra, B., Domingues, A.V., Silva, J.M.,
Loureiro-Campos, E., Gaspar, R., Sotiropoulos, I., Sousa, N., and Rodrigues, A.J. (2020).
Nucleus accumbens medium spiny neurons subtypes signal both reward and aversion.
Mol Psychiatry *25*, 3241–3255. 10.1038/s41380-019-0484-3.

39. Francis, T.C., Chandra, R., Gaynor, A., Konkalmatt, P., Metzbower, S.R., Evans, B.,
Engeln, M., Blanpied, T.A., and Lobo, M.K. (2017). Molecular basis of dendritic atrophy
and activity in stress susceptibility. Mol Psychiatry *22*. 10.1038/mp.2017.178.

40. Pignatelli, M., Tejeda, H.A., Barker, D.J., Bontempi, L., Wu, J., Lopez, A., Palma Ribeiro,
S., Lucantonio, F., Parise, E.M., Torres-Berrio, A., et al. (2021). Cooperative synaptic and
intrinsic plasticity in a disynaptic limbic circuit drive stress-induced anhedonia and passive
coping in mice. Mol Psychiatry *26*, 1860–1879. 10.1038/s41380-020-0686-8.

41. He, Z.-X., Xi, K., Liu, K.-J., Yue, M.-H., Wang, Y., Yin, Y.-Y., Liu, L., He, X.-X., Yu, H.-L.,
Xing, Z.-K., et al. (2023). A Nucleus Accumbens Tac1 Neural Circuit Regulates
Avoidance Responses to Aversive Stimuli. Int J Mol Sci *24*. 10.3390/ijms24054346.

42. Picciotto, M.R., Higley, M.J., and Mineur, Y.S. (2012). Acetylcholine as a neuromodulator:
cholinergic signaling shapes nervous system function and behavior. Neuron *76*, 116–129.
10.1016/j.neuron.2012.08.036.

43. Oldenburg, I.A., and Ding, J.B. (2011). Cholinergic modulation of synaptic integration and
dendritic excitability in the striatum. Curr Opin Neurobiol *21*, 425–432.
https://doi.org/10.1016/j.conb.2011.04.004.

44. Britt, J.P., Benalioua, F., McDevitt, R.A., Stuber, G.D., Wise, R.A., and Bonci, A. (2012).
Synaptic and behavioral profile of multiple glutamatergic inputs to the nucleus
accumbens. Neuron *76*, 790–803. 10.1016/j.neuron.2012.09.040;
10.1016/j.neuron.2012.09.040.

45. Gale, G.D., Anagnostaras, S.G., Godsil, B.P., Mitchell, S., Nozawa, T., Sage, J.R.,
Wiltgen, B., and Fanselow, M.S. (2004). Role of the Basolateral Amygdala in the Storage
of Fear Memories across the Adult Lifetime of Rats. The Journal of Neuroscience *24*,
3810 LP – 3815. 10.1523/JNEUROSCI.4100-03.2004.

46. Gallo, E.F., Greenwald, J., Yeisley, J., Teboul, E., Martyniuk, K.M., Villarin, J.M., Li, Y.,
Javitch, J.A., Balsam, P.D., and Kellendonk, C. (2022). Dopamine D2 receptors modulate
the cholinergic pause and inhibitory learning. Mol Psychiatry *27*, 1502–1514.
10.1038/s41380-021-01364-y.

47. Zhu, Y., Wienecke, C.F.R., Nachtrab, G., and Chen, X. (2016). A thalamic input to the
nucleus accumbens mediates opiate dependence. Nature *530*, 219–222.
10.1038/nature16954.

48. Al-Hasani, R., Gowrishankar, R., Schmitz, G.P., Pedersen, C.E., Marcus, D.J., Shirley,
S.E., Hobbs, T.E., Elerding, A.J., Renaud, S.J., Jing, M., et al. (2021). Ventral tegmental
area GABAergic inhibition of cholinergic interneurons in the ventral nucleus accumbens
shell promotes reward reinforcement. Nat Neurosci *24*, 1414–1428. 10.1038/s41593-021-
00898-2.

49. MacAskill, A.F., Cassel, J.M., and Carter, A.G. (2014). Cocaine exposure reorganizes cell
type- and input-specific connectivity in the nucleus accumbens. Nat Neurosci *17*, 1198–
1207. 10.1038/nn.3783.

50.   Matsumoto, N., Minamimoto, T., Graybiel, A.M., and Kimura, M. (2001). Neurons in the
875   thalamic CM-Pf complex supply striatal neurons with information about behaviorally
876   significant sensory events. J Neurophysiol *85*, 960–976. 10.1152/jn.2001.85.2.960.

877   51.   Zhang, Y.-F., Reynolds, J.N.J., and Cragg, S.J. (2018). Pauses in Cholinergic
878   Interneuron Activity Are Driven by Excitatory Input and Delayed Rectification, with
879   Dopamine Modulation. Neuron *98*, 918-925.e3. 10.1016/j.neuron.2018.04.027.

880   52.   Guzman, R.G., and Kendrick, K.M. (1993). Effect of substance P on acetylcholine and
881   dopamine release in the rat striatum: a microdialysis study. Brain Res *622*, 147–154.

882   53.   Brimblecombe, K.R., and Cragg, S.J. (2015). Substance P Weights Striatal Dopamine
883   Transmission Differently within the Striosome-Matrix Axis. Journal of Neuroscience *35*,
884   9017–9023. 10.1523/JNEUROSCI.0870-15.2015.

885   54.   Cachope, R., Mateo, Y., Mathur, B.N., Irving, J., Wang, H.L., Morales, M., Lovinger, D.M.,
886   and Cheer, J.F. (2012). Selective activation of cholinergic interneurons enhances
887   accumbal phasic dopamine release: setting the tone for reward processing. Cell Rep *2*,
888   33–41. 10.1016/j.celrep.2012.05.011 [doi].

889   55.   Threlfell, S., Lalic, T., Platt, N.J., Jennings, K.A., Deisseroth, K., and Cragg, S.J. (2012).
890   Striatal dopamine release is triggered by synchronized activity in cholinergic interneurons.
891   Neuron *75*, 58–64. 10.1016/j.neuron.2012.04.038 [doi].

892   56.   Kutlu, M.G., Zachry, J.E., Melugin, P.R., Tat, J., Cajigas, S., Isiktas, A.U., Patel, D.D.,
893   Siciliano, C.A., Schoenbaum, G., Sharpe, M.J., et al. (2022). Dopamine signaling in the
894   nucleus accumbens core mediates latent inhibition. Nat Neurosci *25*, 1071–1081.
895   10.1038/s41593-022-01126-1.

896   57.   McConalogue, K., Corvera, C.U., Gamp, P.D., Grady, E.F., and Bunnett, N.W. (1998).
897   Desensitization of the neurokinin-1 receptor (NK1-R) in neurons: effects of substance P
898   on the distribution of NK1-R, Galphaq/11, G-protein receptor kinase-2/3, and beta-
899   arrestin-1/2. Mol Biol Cell *9*, 2305–2324. 10.1091/mbc.9.8.2305.

900   58.   Bowden, J.J., Garland, A.M., Baluk, P., Lefevre, P., Grady, E.F., Vigna, S.R., Bunnett,
901   N.W., and McDonald, D.M. (1994). Direct observation of substance P-induced
902   internalization of neurokinin 1 (NK1) receptors at sites of inflammation. Proceedings of
903   the National Academy of Sciences *91*, 8964–8968. 10.1073/pnas.91.19.8964.

904   59.   Anagnostaras, S., Wood, S., Shuman, T., Cai, D., LeDuc, A., Zurn, K., Zurn, J.B., Sage,
905   J., and Herrera, G. (2010). Automated Assessment of Pavlovian Conditioned Freezing
906   and Shock Reactivity in Mice Using the Video Freeze System. Frontiers in Behavioral
907   Neuroscience *4*.

908   60.   Howard, D.B., and Harvey, B.K. (2017). Assaying the Stability and Inactivation of AAV
909   Serotype 1 Vectors. Hum Gene Ther Methods *28*, 39–48. 10.1089/hgtb.2016.180.

910   61.   Pennington, Z.T., Dong, Z., Feng, Y., Vetere, L.M., Page-Harley, L., Shuman, T., and Cai,
911   D.J. (2019). ezTrack: An open-source video analysis pipeline for the investigation of
912   animal behavior. Sci Rep *9*, 19979. doi: 10.1038/s41598-019-56408-9

A

	Day 1	Day 2	Day 3
	Conditioning	Context Recall	Cue Recall
🔊	⌐‾⌐ x2 10 sec, 80 dB		⌐‾⌐ x5 10 sec, 80 dB
⚡	⌐‾⌐ 2 sec, 0.6 mA		
💡	White	White	Red
	Vanilla	Vanilla	Vinegar

B ● Saline ■ L-733,060

Freezing (%)
Injection Before Conditioning
Context   Cue ***

C

Freezing (%)
Injection Before Cue Recall
Context n.s.   Cue n.s.

D eGFP-KASH *Chat* Tacr1 DAPI

AAV.mU6-Tac1ngRNA1;
EF1a-eGFP-KASH;
hU6-Tacr1-gRNA2

LSL-Cas9 X ChAT-Cre

AC
NAcC

AC

DAPI   eGFP-KASH

Chat   Tacr1

△ ChAT-Cre
▼ ChAT-Cre/Cas9
▲ ChAT-Cre^Tacr1-gRNAs
▲ ChAT-Cre/Cas9^Tacr1-gRNAs

Tacr1 RNA molecules per cell
Chat+ Cells *   *Chat*- Cells n.s.

E

ChAT
Cell Attached

Spontaneous Firing Frequency (Hz)
Baseline SP Wash Out ****

Baseline   Substance P
ChAT-Cre^Tacr1-gRNAs

ChAT-Cre/Cas9^Tacr1-gRNAs
5 sec

F ▲ ChAT-Cre^Tacr1-gRNAs ▲ ChAT-Cre/Cas9^Tacr1-gRNAs

Freezing (%)
Context   Cue ***

G During conditioning
● Saline ● L-733,060

Freezing (%)
Context   Cue **

A ⚡ 2 sec, 0.6 mA, X10
random interval

Dyn-Cre

AAV-DIO-GCaMP7f

fiber

NAc
core

αGFP

Average Z-score (ΔF/F)
-4 -2 0 2 4 6 8

B

0.1mA

0.3mA

0.6mA

1.0mA

Shock Trial

Time (sec)

0.1 mA

0.3 mA

0.6 mA

1.0 mA

Z-score (ΔF/F)

Time (sec)

C

AUC

Shock Intensity (mA)

D

AUC

Shock Trial

E ⚡ 10 sec x2

2 sec, 0.6 mA

Cue
Shock

Z-score

Conditioning

Time (sec)

F 🔊 10 sec X5
30 sec

Cue

Average Z-score

Cue Recall

Time (sec)

G

Freezing to Cue (%)

Foot Shock Intensity (mA)

H

AUC during conditioning

r^2=0.31
p<0.01**

Freezing (%)

0.1
0.3
0.6
1.0

I

Dyn-Cre

AAV-DIO-eNpHR3.0

fiber

DAPI
αGFP

NAc
core

J Inhibition during conditioning

● eYFP
● eNpHR3.0

Laser ▭ 10 sec

▭ 10 sec x2
2 sec, 0.6 mA

Freezing (%)

Context

Cue

**

K Inhibition during cue recall

Laser ▭ 10 sec

🔊 ▭ 10 sec x3
30 sec

Freezing (%)

Context

L ⚠ 532 nm

Freezing (%)

Novel Context

Cue
1 2 3 4 5 6

Off On
Laser

A
2 sec, 0.6 mA, X10
random interval

Average z-score
-2   0   2

AAV-GRAB-ACh3.0

B

0.1 mA

Shock 1
0.1mA

Shock 10
0.1mA

0.3 mA

0.3mA

0.3mA

0.6 mA

0.6mA

0.6mA

1.0 mA

1.0mA

1.0mA

Shock Trial

Z-score

Time (sec)

Time (sec)

C
● Shock 1   ▲ Shock 10

Pause

Inverse AUC

Shock Intensity (mA)

D
Rebound

AUC

Shock Intensity (mA)

E
L-733,060   0.6 mA

Shock Trial

Z-score

Shock 1

Shock 10

Time (sec)

F
● Saline
■ L-733,060

Pause

Absolute AUC

Shock Trial
0.6mA

G
Rebound

AUC After Shock

Shock Trial
0.6mA

H
Cue
Shock

Z-score

Saline
Conditioning

I
Cue
Shock

Z-score

L-733,060
Conditioning

Time (sec)

J

AUC

Pause   Rebound

K

AUC Rebound

$r^2 = 0.53$
$p < 0.01$

Freezing (%)

A AAV-DIO-eNpHR3.0

ChAT-Cre

During Cue
Laser ▬▬▬ 10 sec

🔊 ⎍⎍ 10 sec x2 OR

⚡ 2 sec, 0.6 mA

After Shock
▬▬ 5 sec

⎍⎍ 10 sec x2

2 sec, 0.6 mA

B ● eYFP ● eNpHR3.0

Freezing (%)

Context | Cue

Inhibition During Cue

C

Freezing (%)

Context | Cue   ***

Inhibition After Shock

D ● Saline ● Pirenzepine

Freezing (%)

Context | Cue   **

Infusion Before Conditioning

E

Freezing (%)

Context | Cue

Infusion Before Cue Recall

A ● Cue ● +Shock ● +Shock+L-733,060

D2-MSNs

Vm (mV)

Norm. I (%)

100pA
25msec

Rectification Index

D2-MSNs

n.s.
**
*

B

NASPM

EPSC amplitude (% baseline)

Time (min)

NASPM
Baseline

NASPM
Baseline

C

10pA
1sec

sEPSC Amp (pA)

sEPSC Freq (Hz)

D2-MSNs

**

D

D1-MSNs

Vm (mV)

Norm. I (%)

Rectification Index

D1-MSNs