Using pose estimation and 3D rendered models to study leg-mediated self-righting by lanternflies

Theodore Bien[1][§], Benjamin H. Alexander[1][§], Chengpei Li[1], Natalie Goeler-Slough[1], S. Tonia Hsieh[2], Suzanne Amador Kane[1][*]

[1] Physics and Astronomy Department, Haverford College, 370 Lancaster Ave, Haverford, Pennsylvania, 19041 USA

[2] Department of Biology, BioLife Building, 1900 North 12th Street, Temple University, Philadelphia, 19122 USA

[§] Contributed equally

[*] Corresponding author

Running title: Modeling leg-mediated self-righting

Abstract

The ability to upright quickly and efficiently when overturned on the ground (terrestrial self-righting) is crucial for living organisms and robots. Previous studies have mapped the diverse behaviors used by various animals to self-right on different substrates, and proposed physical models to explain how body morphology can favor specific self-righting methods. However, to our knowledge no studies have quantified and modeled all of an animal's limb motions during these complicated behaviors. Here, we studied terrestrial self-righting for nymphs of the invasive spotted lanternfly (*Lycorma delicatula*), an insect that must frequently recover from being overturned after jumping and falling in its native habitat. These nymphs self-righted successfully in 92-100% of trials on three substrates with different friction and roughness, with no significant

difference in the time or number of attempts required. They accomplished this using three

stereotypic sequences of movements. To understand these motions, we combined 3D poses

tracked on multi-view high-speed video with articulated 3D models created using

photogrammetry and Blender rendering software. The results were used to calculate the

mechanical properties (e.g., potential and kinetic energy, angular speed, stability margin,

torque, force, etc.) of these insects during righting trials. We used an inverted physical

pendulum model (a "template") to estimate the kinetic energy available in comparison to the

increase in potential energy required to flip over. While these insects began righting using

primarily quasistatic motions, they also used dynamic leg motions to achieve final tip-over.

However, this template did not describe important features of the insect's center of mass

trajectory and rotational dynamics, necessitating the use of an "anchor" model comprising the

3D rendered body model and six articulated two-segment legs to model the body's internal

degrees of freedom and capture the role of the legs' contribution to inertial reorientation. This

anchor elucidated the sequence of highly coordinated leg movements these insects used for

propulsion, adhesion, and inertial reorientation during righting, and how they frequently pivot

about a body contact point on the ground to flip upright. In the most frequently used method,

diagonal rotation, these motions allowed nymphs to spin their bodies to upright with lower force

with a greater stability margin compared to the other less frequently-used methods. We provide

a concise overview of necessary background on 3D orientation and rotational dynamics, and the

resources required to apply these low-cost modeling methods to other problems in

biomechanics.

Introduction

Overturned insects benefit from being able to reassume an upright orientation quickly and with

low metabolic energy investment, thereby mitigating the risk of predation, hunger, and

desiccation. This behavior, which is called terrestrial self-righting when performed while on the

ground, has been studied in insects (Full et al. 1995; Faisal and Matheson 2001; Li et al. 2019;

Pace and Harris 2021; Zhang et al. 2021), spiders (O'Donnell 2018), frogs (Malashichev 2006),

turtles (Domokos and Várkonyi 2007) and rats (Pellis et al. 1991). Animals use a rich, diverse

range of body and appendage motions to rotate from supine to upright on the ground (Li et al.

2019). For example, (Frantsevich 2004) identified 20 different modes of self-righting in a study

of 116 species of beetles. Ongoing research aims at understanding how the methods used by a

specific organism to self-right depend on its morphology and locomotor abilities, as well as the

substrate's properties (Davis et al. 2011). Animal righting behaviors also have inspired designs

for robots that can recover easily from upsets—a particular concern for operating in complex

terrains (Saranli et al. 2004; Wang et al. 2022)—using coordinated motions of artificial legs

(Saranli et al. 2004), wings (Li et al. 2016), and tails (Casarez and Fearing 2017).

Here we present the first study to our knowledge that measures and models the detailed

motions of an animal's body and all of its legs during terrestrial self-righting. We studied this

behavior for nymphs of the spotted lanternfly (*Lycorma delicatula*), an invasive insect pest

rapidly expanding its range in the U.S. (Urban and Leach 2022). Spotted lanternflies frequently

jump and fall in their native habitats to disperse, evade predators, and respond to environmental

disturbances (Kim et al. 2011). They therefore need to be able to self-right quickly and

effectively under a variety of circumstances. In an earlier study, we found that falling spotted

lanternfly nymphs are highly successful at self-righting both in the air and during landing (Kane

et al. 2021). Here we turn our attention to how spotted lanternfly nymphs reorient to upright

when starting from rest on their backs.

High-speed multi-angle video was used to record these insects as they attempted to self-right

on several different substrates. These results were used to construct an ethogram of observed

righting behaviors for this species and to determine how they depended on substrate and life

stage. We also performed 3D tracking to quantify the detailed body and leg motions during

righting (pose estimation). To interpret these data, we built on previous studies of terrestrial self-

righting that considered how the animal's gravitational potential energy and kinetic energy vary

during righting (e.g., (Domokos and Várkonyi 2007; Li et al. 2019)). If an organism performs the

motions required to self-right slowly enough that it remains approximately statically stable at

every moment, it can be said to reorient quasistatically. Alternatively, it can use dynamic

motions (inertial reorientation), e.g., by impulsively swinging an appendage (e.g., arm, leg or

tail) so as to generate a reaction counter-torque on the body (Yeadon and King 2017). For a

body in contact with the ground, such maneuvers generate ground reaction torques that can

both enhance the ability to right and stabilize the final orientation after overturning (Frohlich

1979; Davis et al. 2011). Living organisms and robots can take advantage of combinations of

quasistatic and dynamic righting strategies, so we considered both possibilities in our analysis.

Earlier studies have proposed physical models with varying levels of detail ("templates and

anchors" (Full and Koditschek 1999)) to determine whether animals meet the mechanical

requirements for righting either quasi-statically or dynamically (Kessens et al. 2014; Li et al.

2016; Kessens and Dotterweich 2017; Othayoth and Li 2021). To this purpose, we created a

detailed 3D rendered model of spotted lanternflies' bodies using photogrammetry and Blender

computer graphics software. We then used this model to calculate the mechanical properties

required to test these predictions of righting strategies based on the tracked 3D kinematics of

the insects' body and legs measured from high-speed video.

Fig. 1 A) Fourth instar spotted lanternfly nymphs showing the definition of body length, L_{body}, and the anatomical landmarks tracked on video (blue circles) B) Arena and video camera used to film terrestrial self-righting. C) Video frame showing the four camera views recorded of spotted lanternfly nymphs as they attempted to self-right. D) Substrates used in experiments. (Scale bar = 10 mm). E) Illustration of the geometry used to define the projection of the dorsal-ventral axis on the spatial Z direction used to quantify overturned vs righted orientation, $-1 \leq Z_{dv}$

≤ +1 and the tarsus (foot) z-position, z_{tarsus}, in the body frame used to quantify foot motions.

This figure shows a scenario when the body and spatial frames do not coincide. F) Example of a

3D rendered model of a spotted lanternfly nymph used to estimate mechanical properties for

modeling. G) Illustration of a 3D model of the spotted lanternfly nymph's body in the orientation

used to define the body (xyz) and spatial (XYZ) frames in the reference orientation, and the

relationship between coordinate axes, roll, pitch and yaw rotations and Euler angles. (G) (H)

Frame from the animation showing tracked landmarks (Fig. 1A) with the 3D body model and six

jointed rod legs. (I-J) Animation frames showing examples of the overturned, flipping, and

righted resting poses defined in the text.

Materials and methods
112

Data analysis and statistical methods
113

114 All image processing and data analysis were performed using MATLAB vR2023b (Mathworks,

115 Natick MA, USA); specific MATLAB functions are given in italics. Results are reported as mean

116 ± error bars based on instrumental uncertainties, unless noted otherwise.

Insect collection, handling & morphometrics
117

Healthy, intact spotted lanternfly nymphs (Fig. 1A) were collected from tree of heaven (*Ailanthus*

altissima) and wild grape vines (*Vitis vinifera*) in southeastern Pennsylvania (40°00' 30.2" N;

75°18' 22.0" W) in June 22-July 9, 2021. All specimens were collected mid-morning through

mid-afternoon when they are most active (Kim et al. 2011). Experiments were performed in the

laboratory at ambient conditions (23.2 [22.5, 24.1] °C, and relative humidity 62 [42, 70] %

(mean, full range) within 5 minutes of collection to avoid specimen fatigue, hunger, or

desiccation. To determine if self-righting behavior depended on life stage, we studied the two

largest nymphal stages (3rd and 4th instars); earlier stages were too small for tracking on video.

A combination of coloration and body morphometrics were used to determine the life stage of

each specimen studied here, as previously described in (Bien et al. 2023). The body (head-

caudal) length, L_{body}, was measured to ± 0.05 mm using calipers and whole insect mass, M, was

measured to ± 0.4 mg using an analytical balance (Fig. 1A), giving the following values: 3rd

instar: L_{body} = 9.2 ± 0.7 mm, M = 31 ± 11 mg; 4th instar: L_{body} = 11.7 ± 0.8 mm, M = 59 ± 13 mg

(mean ± SD).

Self-righting experiments

Spotted lanternfly nymphs were recorded while attempting to self-right on flat, horizontal (± 0.05

deg) substrates made of posterboard, sandpaper and polished glass. (Fig. 1 B-D) Experiments

were performed for 5 trials for each of 15 unique individuals for each of four different conditions

(combination of instar and substrate considered); the limited number of trials per individual was

chosen to minimize the effects of either learning or fatigue. This gave 299 total trials overall (75

trials/condition for all but the 4th instars on sandpaper, for which one video was corrupted, so

only 74 trials were analyzed).

At the start of each trial, we gently held a spotted lanternfly nymph for a few seconds to induce

tonic immobility (Faisal and Matheson 2001). The nymph was then placed on its back on the

substrate and allowed to attempt to self-right for ≤ 30 s. If the specimen failed to self-right in this

time, the trial was terminated and recorded as a failure. An "attempt" at self-righting was

defined as the time from when a specimen on its back began to lift its dorsum off the ground

until it either turned upright successfully or else fell on its back after failing to right. Successful

self-righting was defined as taking place at the moment when a specimen first came to rest

stably on all of its feet with its dorsum uppermost. The mean attempt time was also measured

for each specimen as the ratio of the time spent attempting to right divided by the number of

attempts, including any time during which it was temporarily quiescent.

Video analysis

High-speed videos of each trial were recorded using a Chronos 1.4 video camera (color, 1280 x

1024 pixels, 1000 fps; shutter speed 500 microsec, Krontech, Burnaby, BC, Canada) with a 12

mm focal length lens (f/# = 7, model M1214-MP2, Computar, Torrance, CA, USA). A total of four

views (main camera and three mirror views) were filmed to allow 3D tracking of all body parts

and legs with minimal occlusion. Data for 3D camera calibrations were obtained using wand

calibration with easyWand software (wand score 0.64, reconstruction error 2.7-4.5 pixels, 7

pixel/mm) (Theriault et al. 2014) and the vertical direction was determined by tracking a falling

ball. To approximate a natural light environment, the arena was illuminated with an LED light

source (SL-200W, Godox, Shenzhen, China) covered by a diffusing softbox so as to provide

7,000 lux at the specimen, similar to the intensity measured in the understory of trees where the

nymphs were collected.

Repeated viewings of all recorded videos were used to determine the stereotypic sequences of

poses assumed while attempting to self-right, along with their frequency, success rate and

timescales, the number of self-righting attempts, how the legs, feet, and body moved during

different behaviors, and the behaviors associated with failed righting attempts. Note that we use

"pose" to describe the body's position and orientation in space combined with its posture (the

detailed configuration of its leg joint angles and body parts.)

The insect's pose was determined quantitatively using the program Direct Linear Transformation

data viewer (DLTdv) (Hedrick 2008) to measure the image plane coordinates of multiple body

anatomical landmarks. (Fig. 1A) These results were then used to compute the 3D XYZ

coordinates of each landmark in the laboratory-fixed spatial frame. Manual tracking was

necessary in most cases to measure the coordinates at sufficiently high resolution and accuracy

to support 3D reconstruction of 16-18 body landmarks in four camera views for hundreds of

frames in each video. Attempts to use automated tracking using DeepLabCut (Mathis et al.

2018) for this study were complicated by frequent occlusions, the small size of the features

tracked, and the similar appearance of the legs and head; this resulted in points often failing to

be tracked or to be tracked too inaccurately for 3D reconstruction, even after repeated rounds of

supervised training. Custom MATLAB code was used to analyze the coordinates for velocity,

angular velocity, kinetic energy, and potential energy as a function of time and body pose.

Tracked coordinates and body orientation axes were smoothed by local regression using robust

quadratic polynomial regression over a moving window of 25 ms (for slower-moving body parts)

and 5 ms (for faster-moving legs) (MATLAB's *smooth* with the "rloess" option, which ignores

data that lie over 6 mean absolute deviations from the regression). Numerical derivatives were

computed using quadratic polynomial regression using the same 25 ms time window (Li et al.

2023). We classified the rotational motions during self-righting using the orientation of the

rotational axis in spatial and body fixed coordinates (defined in "Measuring and representing 3D

orientation" below), the location of the pivot point about which rotations occur, and the distance,

d, of the insect's center of mass (COM) from the axis of rotation. The location of the XY ground

plane on video was measured to 0.25 mm SD along Z by fitting a plane to a cloud of foot points

visually determined to be in contact with the ground at a wide variety of locations throughout a

trial.

Substrate characterization

To test how surface properties affect terrestrial self-righting, we characterized the roughness of

the three substrate materials used: posterboard for 3rd and 4th instars; and 80 grit sandpaper

painted white and polished glass for 4th instars only. (Fig. 1D) The feet of spotted lanternflies

adhere to surfaces using a combination of their tarsal claws and adhesive pads (arolia)

(Frantsevich et al. 2008). To characterize potential grasping by tarsal claws, we compared the

feature sizes for each substrate with the claw dimensions of 3rd and 4th instar spotted lanternfly

nymphs (> 1 µm radius tips and tip-tip spacing 660-849 µm) reported in (Avanesyan et al.

2019). The sizes of surface features of the sandpaper and posterboard were measured from

micrographs using ImageJ (Schneider et al. 2012)**,** and found to be large enough to allow

nymphs to grasp the fibers of posterboard and sandpaper asperities using their tarsal claws.

(Table 1) Polished glass, however, is reported to have sub-nm roughness values too small to

afford a grip by these specimens (Yates and Duffy 2008). Combined with the fact that insect

arolia adhere best to smooth substrates (Al Bitar et al. 2010), this suggests that spotted

lanternfly nymphs should adhere to polished glass using their arolia, but not their claws.

In addition to needing to hold onto and exert forces on the substrate, insects can rotate upright

about a pivot point formed by a body part resting on the ground if frictional forces are sufficient

at this contact region to prevent sliding. To characterize the friction between the exoskeleton

and the substrate, freshly euthanized specimens were placed on their dorsum on an inclined

plane tilted to find the minimum angle, ψ, at which sliding first occurs. The coefficient of static

friction between the exoskeleton cuticle and substrate was then computed as: $\mu_s = \tan \psi$

(Halliday et al. 2013). While we were able to measure multiple instances of sliding on the

inclined plane for glass and posterboard, the frictional force between the exoskeleton and

sandpaper was so great that rolling occurred before sliding in only 2 of 83 inclined plane trials,

so we were only able to determine a lower bound on the corresponding frictional coefficient.

(Table 1)

Based on the combined effects of foot and back interactions, we expected that spotted lanternfly

nymphs should right more effectively on sandpaper (which had the greatest frictional coefficient

tested and substrate feature size amenable to tarsal claw grasping) than on lower-friction

posterboard or glass.

Table 1 Substrate feature size and coefficients of static friction between the insect's dorsum and

substrate. [§] Roughness of polished glass from (Yates and Duffy 2008). (The frictional coefficient

was so high during the sandpaper trials that only 2 trials out of 83 total resulted in sliding instead

of rolling.)

Substrate	Feature size	Coefficient of static friction relative to insect exoskeleton cuticle (mean ± SD)	N specimens (3rd, 4th instars); n trials
Posterboard	40 ± 8 μm diameter fibers (median ± SD)	0.36 ± 0.09	10 (5, 5); 63
Glass	< 1 nm[§]	0.55 ± 0.13	9 (5, 4); 73
Sandpaper	570 ± 100 μm diameter pores (mean ± SD)	≥1.2 ± 0.4	2 (1, 1); 2

3D rendered model and calculations of mechanical properties

We created 3D rendered models of spotted lanternfly nymphs to use in analyzing the tracked

3D coordinates. Photogrammetry and microscopy were used to determine the 3D geometry of

the entire body and legs. Six 4[th] instar nymphs also were dissected for use in separately

measuring the dimensions and mass of the body and each leg. These data were used with

Blender rendering software (www.blender.org, January 16, 2023) to create realistic 3D models

of the spotted lanternfly nymph body; for details see (Li et al. 2023). (Fig. 1F) Three different 3D

models were created to match the slightly different morphologies of the specimens analyzed in

detail below to provide measures of uncertainty for the mechanical properties. (See

Supplementary Dataset S1 for Blender files.) The 3D body models were used to create .stl files

that contained the triangular mesh describing its surface geometry; these were used with the

MATLAB package RigidBodyParams (Semechko 2023) to compute the body's COM

coordinates.

Mechanical properties during self-righting were computed using an "anchor" model in which the

body (head, abdomen and thorax) was treated as a single rigid object represented by the 3D

body model, and each leg was treated as an articulated assembly of two rigid rod-like segments

(one segment each for the combined tibia and tarsus, and one for the combined femur and

shorter coxa) based on the measured leg segment masses and lengths. The pose of the

anchor's body and legs at each timestep was determined using the results from 3D video

tracking. (Fig. 1H) Because an earlier growth allometry study has shown that the body

morphology of spotted lanternfly nymphs scales isometrically with body length (Bien et al.

2023), we analyzed the mechanical properties of tracked data for different specimens using 3D

models scaled to a standardized body length and mass as follows. First, we multiplied the

tracked coordinates for each trial analyzed by a scaling factor so the specimen's body length in

rescaled coordinates corresponded to a standardized value of L_{body} = 8.9 mm. The rescaled

coordinates were then analyzed in combination with a 3D model with this body length assuming

a whole insect mass of 28.4 mg typical of this body length.

Measuring & representing 3D orientation

We now consider how to represent orientation in 3D (attitude) during self-righting. In this study,

attitude was defined as the 3D orientation of *xyz* axes defined by the insect's anatomical axes

(cranial-caudal, lateral, and dorsal-ventral, respectively), in a body frame of reference with

respect to the *XYZ* axes of the laboratory-fixed spatial frame. (Fig. 1G) Attitude coordinates

describe an ordered sequence of rotations that transform the body frame from a reference

orientation to the orientation of interest with respect to the spatial frame. We use Euler angle

attitude coordinates ψ, θ, φ (roll, pitch, yaw), which describe rotations about each of the *xyz*

body axes, respectively. In the reference orientation used here, (ψ, θ, φ) = (0,0,0) corresponded

to the body frame *xyz* axes aligned with the spatial frame *XYZ* axes. (Fig. 1G) The rotations

used to relate the Euler angles to attitude were performed in as the order yaw, pitch, roll (Tait-

Bryan convention, *zy'x''* order), where, e.g., y' refers to the new orientation of the pitch (lateral)

axis after the first yaw rotation by φ about z, and x'' refers to the direction of the cranial-caudal

axis for roll rotations after rotation by θ about the y'-axis. (Fig. S1A) Note that if the rotations are

performed in a different order, this results in a different 3D orientation (Schaub and Junkins

2003). (Fig. S1B) This also means that, in general, the Euler angles (ψ, θ, φ) do not correspond

to the angles between the final *x''' y''' z'''* body axis orientation and the *XYZ* spatial axes. (Fig.

S1C) We therefore reported the *Z*-component of the body dorsal-ventral axis ($-1 \leq Z_{dv} \leq +1$) in

the spatial frame (Fig. 1E) as a measure of the body's orientation during righting. When the

insect is overturned and flat on its back, then Z_{dv} = +1, and when it is upright with its dorsal-

ventral axis pointed downward along the vertical, Z_{dv} = -1. (Fig. 1I, K) We refer to the pose when

Z_{dv} = 0 as the flipping point. (Fig. 1J)

To avoid mathematical ambiguities that can arise when using Euler angles for calculating

rotations (e.g., gimbal lock), we first performed calculations on orientations computed from the

tracked coordinates using quaternion-based functions in MATLAB, and then converted the

result to Euler angles (Wang et al. 2023). Given the measured 3D orientations of the body and

appendages in the spatial frame at times *t* and *t* + Δt , we used MATLAB functions *absor* (Matt

2024) and *angvel* to compute the corresponding angular velocity vector, $\vec{\omega}(t)$, of each object.

The rotation axis, $\hat{r}$, rotation angle, $\Delta\theta$, and angular speed, ω, in rad s^{-1} then were computed

using $\vec{\omega}(t) = \frac{\Delta\theta}{\Delta t}\hat{r} = \omega\,\hat{r}$, and the angular acceleration, α, was calculated as $\alpha = d\vec{\omega}(t)/dt$.

Stability metrics

The stability margin of poses during locomotion can be computed as a test for static mechanical

stability (McGhee and Frank 1968; Ting et al. 1994). To this purpose, we used custom MATLAB

code and the tracked insect body coordinates to compute the support polygon (the convex hull

formed by all body parts in contact with the ground). (Fig. 2A) A tracked leg segment or point on

the body was defined as contacting the ground if it lay within a distance comparable to the

tracking uncertainty of the ground plane. The stability margin (*SM*) then was computed as the

shortest distance from the boundary of the support polygon to the projection of the insect's COM

on the ground. If only pushing forces are exerted by the organism on the ground, then *SM* > *0*

for stable equilibrium, *SM* = 0 for metastable equilibrium, and *SM* < 0 for unstable poses. The

value of *SM* is at its maximum (the ideal stability margin, *ISM*) when the projected COM agrees

with the centroid of the support polygon. The ratio *SM:ISM* ≤ 100% then measures how close

the insect's pose is to optimal static stability. However, for insects that can exert pulling forces

on the ground, either through adhesion with their arolia or by grasping the ground with tarsal

claws, static stability can be achieved for *any* value of *SM* given a sufficiently high pulling force.

Analyzing energy during righting

To determine to what extent an organism's motions during righting contribute to inertial

reorientation, we next consider how to compute its potential and kinetic energy. Many animals,

including spotted lanternflies, must increase their gravitational potential energy (hereafter,

potential energy or *PE*) between the initial overturned and final righted poses, as well as

possibly overcoming one or more intervening *PE* maxima or "barriers" (*PE$_{barrier}$*). (Fig. 2B)

Following (Domokos and Várkonyi 2007; Othayoth et al. 2021), we computed $PE = MgZ$ for

each pose observed during self-righting, where $g = 9.81$ m s^{-2} is the acceleration of gravity and

Z is the insect's COM height relative to the ground. (Fig. 2B) The potential energy landscape for

terrestrial self-righting is then the value of PE calculated for all possible poses consistent with

the organism's morphology and range of motion about each of its locomotor degrees of freedom

that can result in a transition from overturned to upright (Othayoth et al. 2021). By analogy with

free energy landscape models in physical chemistry, it has been proposed that self-righting by

animals and robots can be modeled as a transition between the initial and final configurations,

with the optimal trajectory in the PE landscape defined by following local PE basins so as to

minimize any intervening energy barriers (Kessens et al. 2012; Li et al. 2019). However, in this

study, the spotted lanternfly nymphs were observed to sometimes lift their bodies off the

substrate and use all their legs during righting, so it was not feasible to compute PE for the full

multidimensional configuration space during self-righting. Instead, we computed a simplified PE

landscape for the 3D body model with one point of the body in contact with the ground for all

possible values of pitch and roll to allow a comparison with measured values of body pitch and

roll and with earlier studies of cockroach self-righting (Li et al. 2019).

Purely dynamic self-righting requires the organism to accumulate enough kinetic energy $KE(t)$,

to enable it to rotate upright while increasing its potential energy from the current value, $PE(t)$, to

the maximum during righting, PE_{max} (Robins et al. 1998; Kessens and Dotterweich 2017; Xuan

and Li 2020). This requirement can be stated as $KE(t) \geq \Delta PE(t) = PE_{max} - PE(t)$. This time

dependence indicates that the timing of a dynamic motion (e.g., a push that increases the

body's rotational KE or a rapid leg swing) determines the effectiveness of its contribution to

inertial reorientation. For example, if an inverted pendulum is near the metastable apex of its

trajectory, at which $\Delta PE \approx 0$, then *any* nonzero KE for rotations with the correct direction result

in tip-over; farther from the apex, the inverted pendulum requires a greater kinetic energy to

overcome the *ΔPE* required to reach the apex and tip it over. In addition, not all contributions to

an object's kinetic energy are available for conversion to work against gravity so as to promote

righting. For example, the kinetic energy associated with rotations about a vertical axis or with

horizontal translational motions is irrelevant for righting. For this reason, we define an available

kinetic energy, $KE_{avail}(t)$, that can contribute to righting rotations and discuss below how to

calculate it after developing the physical models used to interpret our tracked data.

To describe the contribution of inertial reorientation to righting at time *t*, we define a

dimensionless righting number, $RN(t) = KE_{avail}(t)/\Delta PE(t)$. If $RN(t) \geq 1$, then the object has

enough available kinetic energy to right at time *t* without exerting any additional torques (purely

dynamic righting). At the other limit, $RN \ll 1$, righting is purely quasistatic. If $RN < 1$ but not $\ll$

1, RN quantifies how much the available kinetic energy contributes to inertial reorientation by

reducing the ground reaction torques required to right.

Fig. 2 A) Top view of a 3D model of a spotted lanternfly nymph (red) showing the definition of

the stability margin. The support polygon (gray) is the convex hull formed by all body parts in

353 contact with the ground support. The stability margin (dashed line) is the shortest distance from

354 the boundary of the support polygon to the projection of the COM on the ground, while the ideal

355 stability margin (gray line) is the shortest distance from the support polygon's in-plane centroid

356 (×) to the boundary. (B) Schematic gravitational potential energy vs pose diagram showing the

357 definition of the increase, $\Delta PE(t)$, in potential energy required to surmount the potential energy

358 barrier starting at time t. Insets show sample images of a spotted lanternfly nymph in the

359 overturned and righted poses. (Color online)

360 ## Rotational kinematics and dynamics

361 Using tracked coordinates and the 3D anchor model, we also computed a variety of measures

362 for interpreting the rotational kinematics and dynamics during righting. We first briefly review

363 some necessary background; for a full treatment of the relevant topics in advanced classical

364 mechanics see (Goldstein 1980). An object's angular acceleration, α, depends on net torque

365 (moment of force), τ_{net}, as (Fig. 3A):

366   $\tau_{net} = r_{\perp}\,F = I_{rot}\,\alpha,$         (1)

367 where F is force, $r_{\perp}$ is the magnitude of the moment arm (the vector from the point at which the

368 force acts to the axis of rotation) perpendicular to the force, and the moment of inertia, I_{rot},

369 defines resistance to changes in angular velocity (Goldstein 1980). An object's moment of

370 inertia, I_{rot}, depends on both the axis of rotation and its relation to the distribution of its mass,

371 M, via (Fig. 3B):

372   $I_{rot} = \int_{M} r^2 dm$         (2)

373 Note that I_{rot} is computed in the body frame of the object, independent of its orientation in the

374 spatial frame. For a rod with mass m and length L rotating about an axis passing through its

375 COM, Equation (2) gives $I_{rod} = \frac{1}{12}mL^2$. (Fig. 3C) We used this relationship to compute the

376 moment of inertia of each of the two segments used to model each leg in the 3D model. We

also needed to account for the distance between the COM of each segment and the whole

insect COM using the parallel axis theorem. This states that, given the moment of inertia, I_{COM},

for rotations about an axis that passes through an object's COM, the value for rotations about

the same axis displaced a distance *d* from the COM, I_d, is (Fig. 3E, F):

$$I_d = I_{COM} + Md^2 \qquad\qquad (3)$$

The tracked body coordinates were used with Equation 3 to determine the relevant geometry for

computing each leg segment's moment of inertia at a given time.

Insect bodies have complex shapes that influence their 3D rotational dynamics. In the most

general case, the moment of inertia for rotations about the COM of a 3D object is a 3×3 matrix,

the inertia tensor $\overleftrightarrow{I_{COM}}$. There always exists a body coordinate frame in which the inertia tensor

is a diagonal matrix, $\overleftrightarrow{I_{COM}} = \begin{bmatrix} I_x & 0 & 0 \\ 0 & I_y & 0 \\ 0 & 0 & I_z \end{bmatrix}$, in which the *xyz* axes are called the principal axes and

the object can undergo stable rotations for zero net torque about the *x, y,* or *z* principal axis with

moment of inertia, $I_x, I_y,$ or I_z, respectively. For example, Fig. 3G shows the principal axes for

the 3D model of the spotted lanternfly body, which happen to agree with the body frame's roll,

pitch, and yaw axes. In the most general case, the moment of inertia can be computed for

rotation about any axis, $\hat{r}$, that passes through the object's COM (Fig. 3H) as:

$$I_{\hat{r}} = \hat{r}^T \cdot \overleftrightarrow{I_{COM}} \cdot \hat{r} \qquad\qquad (4)$$

Values of the moment of inertia and principal axes for the spotted lanternfly's body were

computed using the MATLAB package RigidBodyParams (Semechko 2023) with the 3D body

model. We then used Equations (3) and (4) with the body axes, calculated rotation axes, and

pivot point to determine the moment of inertia of the body, which was added to the contributions

of each leg segment to give the whole insect moment of inertia at each time step.

Fig. 3 A) Geometry used in defining angular velocity, ω, force, F, moment arm, r, and the moment of inertia, I_{rot}, for a point mass, m. Geometry for defining moment of inertia for the general case of (B) a distributed mass and C) a rod rotating about its center of mass (COM). The moment of inertia depends on the axis of rotation about the COM (D) and the distance, d, from the COM to the axis of rotation (E), as in the case of an inverted physical pendulum (F). The geometry for defining spin rotations about the COM and orbital motion of the COM is also shown in F. (G) Moments of inertia about each principal axis for the case where they

correspond to the body's roll, pitch and yaw axes. (H) Calculation of the moment of inertia about

an arbitrary axis of rotation, $\hat{r} = \frac{\vec{\omega}}{\omega}$. (Color online)

Following (Li et al. 2016), we also created a template that modeled the motion of the insect

during righting as that of an inverted physical pendulum rotating about a fixed pivot point on the

ground. (Fig. 3F) Nymphs indeed were observed to always maintained several points of no-slip

contact with the ground during each successful righting attempt. This template also assumes

that the relative motion of body and legs was slow enough to allow them to be approximated as

a rigid object over times comparable to a few video frames. Using this model, we estimated the

$KE_{avail}(t)$ for rotating the insect upright as the kinetic energy due to translation of its COM

(orbital motion) and rotation about the COM (spin), assuming the same angular velocity for the

orbital, $\omega_{orbital}$, and body roll and pitch spin, ω_{spin}, motions. (Fig. 3F) For the inverted physical

pendulum, this value of KE_{avail} comprises the kinetic energy of all motions that could contribute

to increasing the height of the COM, and hence increasing *PE*, while excluding the kinetic

energy of motions that leave COM height unchanged (e.g., purely horizontal translations,

rotations about a vertical axis, counter-rotations of two or more body components about the

same axis with angular momenta that sum to zero, etc.) We computed for each time step the

whole insect's COM, its distance *d* from the pivot point, its COM speed, $v_{orbital}$, and the spin

moment of inertia of the whole insect about its COM (Fig. 3D) This allowed us to compute the

orbital angular velocity, $\omega_{orbital} = v_{orbital}/d$, and total moment of inertia about a pivot point on

the ground a distance d away from the COM, $I_{pendulum}$ using Equation 3 (Halliday et al. 2013).

The pendulum's kinetic energy was then computed as $KE_{avail} = \frac{1}{2}I_{pendulum}\omega_{orbital}^{2}$.

The insect's moment of inertia is necessary for computing its kinetic energy during righting, but

it has other potential influences on righting success based on fundamental principles of

rotational dynamics (Halliday et al. 2013). To see why, we note that an inverted physical

pendulum with mass M has natural period, $T_{pendulum} = 2\pi\sqrt{\dfrac{I_{pendulum}}{Mgd}}$ (Fig. 3F) This period sets

the timescale for failure if the animal falls due to losing traction or to being unable to sustain

sufficient force to stay raised, and the timescale for rotating upright from a given starting

position. For example, gymnasts hold their arms outstretched when walking along a beam to

increase the time allowed for making corrective motions to avoid falling. Furthermore, the inertial

torque exerted on the body by corrective relative rotations of their arms is proportional to the

arms' moment of inertia, so increasing arm extension, and hence the moment of inertia, allows

the gymnast to compensate for body tilting using smaller angular acceleration. In righting, the

opposite is true: for an overturned animal, lower values of I_{pitch} and I_{roll} facilitate rotating

upright more rapidly, while a larger value of I_{yaw} allows for corrective motions to prevent yaw

rotations, which are unhelpful in the overturned pose. Second, the time, t_{rot}, required to rotate

any object with moment of inertia I_{rot} acted on by a constant torque, τ, through an angle θ is $\theta =$

½ α t_{rot}^2 = ½ τ / I_{rot} t_{rot}^2, so $t_{rot} = \sqrt{2\pi\,{}^{I_{rot}}/_{\tau}}$. To compensate for an increase in moment of

inertia, the animal therefore either must exert a greater torque to turn over in the same time, or

exert the same torque for a longer duration. It can exert a greater torque by either exerting a

greater force or by using a larger moment arm. However, the magnitude and duration of force

production are limited because righting is a strenuous activity for insects (Full et al. 1995), and

the maximum moment arm is limited by the length of the legs and their limited ranges of motion.

Using the template, we also estimated the net ground reaction torque acting on the insect with

respect to the pivot point, $\tau_{net} = I_{pendulum}\alpha_{orbital}$, where $\alpha_{orbital} = d\omega_{orbital}/dt$ is orbital

angular acceleration. During self-righting, the forces acting on the insect include gravity, which

acts only on the insect's COM, and ground reaction forces at each contact between the body

and ground. The net ground reaction torque for rotations about the pivot point is therefore:

$$\tau_{reaction} = \tau_{net} - \tau_{grav} = I_{pendulum}\alpha_{orbital} - M\,g\,d = r_{\perp}F_{reaction} \tag{5}$$

where $F_{reaction}$ is the net ground reaction force, and r_{COM} and $r_{\perp}$ are the moment arms for the

force of gravity and the ground reaction force, respectively. We estimated r_{COM} as the distance

between the pivot point and the projection of the insect's COM on the ground, and $r_{\perp}$ using the

mean distance from all tarsi in contact with the ground and the effective pivot point. We also

computed the mean reaction force exerted by each leg (Full et al. 1995) by dividing the net force

by the number of tarsi in contact with the ground (n_{tarsi}) at each timestep: $F_{leg} = F_{reaction} / n_{tarsi} = $

$\tau_{reaction} / (n_{tarsi}\,r_{\perp})$.

Leg coordination

We also analyzed how leg motions might influence inertial reorientation. Because the relevant

leg motions require changing the height of the tarsus above the body dorsal plane, we

computed the value of z_{tarsus} for each leg vs time for comparison with other measures during and

after righting (Fig. 1E). Following, (Othayoth and Li 2021) the coordination between the z-

positions of the jth and kth tarsi, z_j and z_k, was analyzed using the discrete-time normalized

cross-correlation, $z_j * z_k(t_{lag})$, a type of inner product between two functions, one of which is

offset in time from the other by t_{lag} (the lag time):

$$z_j * z_k(t_{lag}) = \frac{1}{n|\,z_j(0)||z_k(0)|}\sum_{i=1}^{n} z_j(t_i)\,z_k(t_i + t_{lag}) \tag{6}$$

where the sum is performed over all n time samples (Press 2007). The magnitude (similarity) of

the normalized cross-correlation of two signals is a measure of how correlated one signal

remains with another when shifted by a positive or negative time offset called the lag time,

where +1 is the maximum similarity possible for in-phase coordination, and -1 the maximum for

out-of-phase motion. For example, the cross-correlation of any function with itself (its

autocorrelation) is equal to 1 at zero lag time, while the autocorrelation of a random signal with

zero mean decays rapidly to zero as lag time increases. (Fig. S6) The autocorrelation of a

periodic signal has periodic peaks at nonzero lag times at integer multiples of the signal period,

T_j, with peak similarities that indicate the timescale over which the signal maintains correlation

with itself. Fig. S6 illustrates this behavior using several examples: a sine wave has peaks with

similarity = 1 at all values of T_j, while a sine wave with added noise in its period, phase or

amplitude has periodic peaks with similarities that decrease with increasing absolute lag time. If

two sinusoidal signals have the same period but one is shifted by a constant phase offset, then

their cross-correlation function has its first peak at a lag time equal to that corresponding to the

phase offset.

We performed correlation analysis from the first frame analyzed to when the trajectory reached

its apex using *xcorr* in MATLAB, after standardizing z_{tarsus} to zero mean to remove trivial

correlations between constant values. First, the normalized autocorrelation of z_{tarsus} for each

tarsus was computed as a function of lag time to measure the timescale over which each leg's

motion remained correlated, as well as its maximum peak similarity at nonzero lag time.

Second, to determine the phase offset and maximum similarity of each pair's motions, the

cross-correlation function was measured between each pair of z_{tarsus} values for the forelegs and

midlegs (the hindlegs were not considered because they were primarily used for relatively static

stabilizing, propelling or pulling motions). (Fig. S7-9)

Results

Terrestrial self-righting behaviors

Spotted lanternfly nymphs were found to self-right using a sequence of stereotypic motions.

(Fig. 4A, Movie 1) Initially, the nymph lay flat on its back (overturned). Upon recovering from

tonic immobility, the nymph moved its legs in probing motions in all possible directions

(searching) in apparent attempts to locate the surface, similar to those described for beetles and

spiders prior to righting (Frantsevich 2004; O'Donnell 2018). Once a foot made contact with the

ground dorsally, the other legs were quickly elevated and extended so as to grasp the ground

(planting). Next, the nymph primarily used its forelegs, with some assistance from the midlegs,

to push against the ground so as to pitch and raise its cranial end off the ground (lifting). It then

attempted to flip upright (righting) using a combination of propelling and swinging leg motions

(Fig. 4A, Movie 1); we discuss more specifics of these movements below. During the post-tip-

over recovery phase in successful trials, the nymph used its legs to achieve the pose typical of

resting and walking (righted), in which its cranial-caudal axis was pitched upward from the

horizontal. If self-righting was unsuccessful, the nymph toppled onto its side or dorsum (failed),

after which it either rested or resumed searching.

Spotted lanternfly nymphs self-righted successfully (92-100% all trials) on all three substrates

for both life stages studied. (Fig. 4B; Table S1) We performed linear regression to test whether

success rate, time/attempt and number of attempts were correlated with trial number to test the

hypotheses that: 1) fatigue might result in poorer righting performance upon repeated attempts,

or, conversely, that 2) learning from repeated attempts could improve righting outcomes. (Fig.

S2A-C, Table S2) Only the number of attempts for the 3rd instar nymphs on posterboard

increased significantly with trial number. Given this result and the balanced study design, we

consequently pooled data across trials and individuals before performing further tests. The

percent of self-righting trials resulting in success did not depend significantly on the combination

of instar and substrate (Fig. 4B; Table S3). All self-righting trials were successful for all 3rd instar

specimens for all three substrates and for all 4th instar nymphs on sandpaper. All 4th instars

succeeded in the first trial and in at least 1 additional trial on posterboard and glass. The only

trials in which a specimen failed to eventually self-right involved five failures by two 4th instars on

posterboard and five failures by five 4th instars on posterboard or glass. Significance testing

using Kruskal-Wallis revealed that the total time required to self-right, the time/attempt, and the

number of attempts did not depend significantly on the combination of substrate and life stage.

(Fig. S2D-F; Table S4)

Fig. 4 A) Ethogram showing stereotypic poses assumed by spotted lanternfly nymphs during

terrestrial self-righting. B) Percent successful trials for each combination of substrate and life

stage (instar) studied. (75 total trials per condition: N = 15, 5 trials/specimen). Sequences of

closeup still images for (C) diagonal rotating about a pivot point on the caudal end (4th instar on

sandpaper), (D) lifted rotating with the legs but not the body contacting the substrate (4th instar

on sandpaper); and (E) pitching about a pivot point on the caudal end (3rd instar on

posterboard). (C-E) Times relative to the first image shown at top left in each frame. (Color

online)

Fig. 4C-E and Movie 1 show examples of the rotational body and leg motions used by spotted

lanternfly nymphs for self-righting. In 96.7% (289:299) of all trials, this involved the nymph

rotating its body upright about a rotational axis intermediate between pitching and rotating while

pivoting about a contact point between its body and the ground ("diagonal rotating"). (Fig. 4C)

During diagonal rotating, the hindlegs remained planted while the fore- and midlegs swung from

the dorsal to ventral sides, consistent with swinging motions expected to promote inertial

reorientation. Once the forelegs contacted the ground, the specimen was close to the resting

pose for this righting mode because its cranial-caudal axis was already elevated relative to the

ground.

In 3.0% of trials (9:299; four 4th instars on sandpaper and 1 on posterboard, three 3rd instars on

posterboard), the nymph rotated upright with only its legs, but not its body, contacting the

ground. (Fig. 4D, Movie 1) This lifted rotating behavior required the nymph to position itself on

all three of its legs on the side nearer the substrate while swinging its raised contralateral legs

dorsal-ventrally so as to flip rapidly upright in a single motion.

Only one specimen in one trial succeeded in righting by pitching about a pivot point formed by

its caudal end resting on the ground (0.3% = 1:299; one 3rd instar on posterboard), although

several other specimens unsuccessfully attempted to right using this mode. (Fig. 4E, Movie 1)

During pitching, the nymph first used its hindlegs to pull on the ground so as to rotate and lift its

body onto a support tripod formed by its hindlegs and caudal end. It then swung its fore- and

midlegs from the dorsal to ventral side consistent with attempting to grasp the ground and to

provide inertial torques for righting. Once the forelegs touched the ground, the nymph used all

its legs to lever its body forward about the caudal end as a pivot point to complete righting.

Pooling data across all trials, life stages and substrates, we found no significant difference

among the time to self-right, time/attempt or number of attempts for diagonal rotating and lifted

rotating. (Table 2, Table S5, Fig. S2G-I) While the mean time/attempt for the single pitching trial

agreed with the mean of the other two righting methods, the total time required to self-right by

pitching and the number of attempts before success lay near their outer ranges.

Table 2 Summary statistics for different methods used to self-right by spotted lanternfly nymphs;

also see Table S5 for results of Kruskal-Wallis tests.

	Diagonal rotating (289 trials, 60 individuals)	Lifted rotating (9 trials, 7 individuals)	Pitching 1 trial
Time to self-right (s) mean ± SEM (SD)	0.98 ± 0.10 (1.73)	1.72 ± 1.10 (3.28)	7.65
Time/attempt (s) mean ± SEM (SD)	0.31 ± 0.01 (0.17)	0.31 ± 0.05 (0.14)	0.32
Number of attempts mean ± SEM (SD)	3.3 ± 0.2 (3.7)	4.0 ± 1.8 (5.4)	24

Failure to self-right was associated with several factors. Some specimens became fatigued and

stopped trying to right after repeated attempts, consistent with earlier reports that self-righting is

a strenuous activity (Full et al. 1995). (Fig. S3, Movie 1) Slipping at the contact point with either

glass or posterboard sometimes compromised the nymph's ability to maintain a secure foothold

or stationary pivot point. (Fig. S3A) In other cases, the specimen appeared unable to exert

sufficient torque to flip upright; e.g., for some failed attempts on posterboard, the nymph lifted

and rested its body on its caudal end, then tried to rotate upright by a combination of swinging

its legs from the dorsal to ventral side and pulling on the ground while pivoting on its caudal end,

but finally fell back onto its dorsum instead of righting. (Fig. S3B)

Mechanical properties during terrestrial self-righting

We tracked body and leg landmarks for the one successful pitching trial and for two trials each

for diagonal rotating and lifted rotating (5 total trials, each for a different individual). Movie S1

illustrates the kinematic uniformity among trials that used each righting strategy. Results were

scaled to standard body mass and dimensions to allow direct comparisons between data for

different specimens. The results of the analyses are plotted in Fig. 5 and S4. We refer to the

approximate interval of time in which the dorsal-ventral axis changes orientation most rapidly as

"active overturning" (see gray shaded regions in Fig. 5).

The tracking results were used to test the following predictions of the inverted physical

pendulum template. First, the angular speeds for body spin, ω_{spin}, and orbital motion, $\omega_{orbital}$,

should be equal, so the body should flip its dorsal-ventral axis orientation and reach its

maximum potential energy at the same time. (Fig. 5A-I) While ω_{spin} and $\omega_{orbital}$ were similar

initially during all three methods, during active overturning the body's angular speed was

significantly greater than that for orbital motion, resulting in the body rotating to upright before

the insect's COM reached its apex. (Table 3, Fig. 5J-L) Second, the template predicts that

during overturning the insect's COM trajectory should be a circular arc curving downward and

with radius comparable to the maximum height. (Fig. 3F) The measured trajectories were

approximately circular during active overturning until the apex for all but one diagonal rotation

trial, although only those for lifted rotation moved in an approximately vertical plane as

predicted. (Fig. S5) These discrepancies indicate that the rigid object approximation used in the

template did not capture fully all of the rotational dynamics during righting, highlighting the

importance of measuring the dynamic leg motions in detail and using the more realistic anchor

model.

We now consider how energy varied throughout the righting process. (Fig. 5D-I) All specimens

that successfully righted started out overturned with legs raised and ended in an upright resting

pose. During planting, because the body rested on its dorsum and the legs extended laterally

and dorsally to the ground, potential energy decreased to a global minimum from its initial value

when overturned, PE_i. The total PE then increased monotonically when the nymph lifted its body

in preparation for attempting to right and during active overturning. The increase in total PE

between its initial and maximum value was the same within measurement uncertainty for

diagonal rotating and pitching, and somewhat lower for lifted rotating. (Table 3)

The comparison of potential and kinetic energy during righting revealed that spotted lanternfly

nymphs relied primarily on quasistatic righting until shortly before flipping. The available kinetic

energy estimated using the template was negligible compared to $\Delta PE(t)$ until the specimen

began active overturning: specifically, $RN(t) = KE_{avail}(t)/\Delta PE(t) < 25\%$ until 36, 31 and 26ms

before the apex for diagonal rotating, lifted rotating and pitching, respectively. During active

overturning, inertial reorientation contributed substantially ($RN > 50\%$) to righting within

approximately 23-28 ms of the apex. (Fig. 5G-I) However, as mentioned above, the template's

estimate for KE_{avail} did not include the body's faster spin rotation that resulted in flipping before

the apex, which indicated dynamical righting played a greater role than computed using the

template.

Fig. 6 shows the measured self-righting trajectories in 2D body pitch-roll configuration space.

The total *PE* computed for the legs and the body at its actual height relative to the substrate is

indicated by the colored trajectory markers. The heatmap shows *PE* computed for the 3D body

model, with no contributions from legs or the body's vertical displacement from the ground. The

measured trajectories and PE landscapes for diagonal rotating, lifted rotating, and pitching by

spotted lanternfly nymphs were similar to those shown in (Li et al. 2019) for diagonal rotating,

rolling on the ground (as opposed to lifted rotating), and pitching for a simple geometrical model

of self-righting by cockroaches. For the PE landscape model used in the latter study, the

predicted frequency of righting methods (rolling, diagonal rotating, and pitching, from highest to

lowest frequency) based on PE barrier height agreed with the empirical data for Madagascar

hissing and American cockroaches. However, the frequency of righting methods used by

spotted lanternflies did not agree with this order, presumably due to the importance of their legs

and body height from the ground in determining total PE, and the important contribution of their

internal motions to inertial reorientation.

The stability margin had very different values during the three righting methods. During diagonal

rotating, the nymph always maintained a positive stability margin near its ideal value by keeping

one body point and 3-6 leg parts in contact with the ground. (Fig. S6D-F) By contrast, the

stability margin was always negative during lifted rotating and pitching until the nymph was

upright because the support polygon formed by the legs (for both methods) and caudal end (for

pitching only) in contact with the ground did not enclose the COM's projection on the ground

until after flipping. (Fig. S6A-C) During pitching the nymph sometimes remained stationary when

the stability margin was negative, indicating that it used its claws or arolia to adhere to the

substrate.

While the mean ground reaction torques were similar within measurement uncertainty for all

three righting methods, the mean and maximum force per leg, F_{leg}, were lower during diagonal

rotating than lifted rotating and pitching. This was because the most commonly-used method

had more ground contacts and a somewhat greater moment arm, $r_\perp$. (Table 3, Fig. S6) ground

reaction torque and F_{leg}, were positive until flipping, consistent with the legs providing an

overturning torque during righting for all three methods. (Fig. S6G-I) During recovery, the

direction of F_{leg} varied as expected as the legs executed corrective motions required to reduce

the body's spin so that it came to rest in the righted pose.

Moment of inertia calculations were consistent with the frequencies of observed righting

strategies. (See Table 3 for summary statistics, and Fig. S7 for plots of time dependence.) First,

spotted lanternfly nymphs could significantly change their moment of inertia by flexing or

extending their legs, which accounted for 65-72% of their total spin moment of inertia; this

enhanced their ability to generate inertial torques using leg swings. Second, the body's mean

spin moment of inertia during diagonal rotating and lifted rotating, was 64% and 74%,

respectively, of that for pitching. This means that either ground reaction or inertial torques

performed for a given time resulted in a lower increase in ω_{spin} during pitching than the other

two methods; this agreed with the measured values of ω_{spin} shown in Fig. 5J-L. Third, yaw

rotations of the 3D body model had a greater moment of inertia than for roll or pitch (I_{roll}, I_{pitch},

I_{yaw} = 0.46 ± 0.05, 1.25 ± 0.01, 1.50 ± 0.04 x 10^{-10} kg m^2, respectively), which indicates that a

greater torque is required to achieve the a given yaw angular acceleration than for pitch or roll.

Pure yaw rotations not only do not contribute to righting when the nymph is on its back, they

sometimes even lead to righting failure (Movie 1); therefore, a greater resistance to increases in

yaw rate is helpful for righting.

The analysis of the tarsal motion vs time was consistent with the nymphs swinging their forelegs

and midlegs to facilitate inertial reorientation. For example, in Fig. 4C and 5M, we see that

during diagonal rotating, the uppermost foreleg and midleg were successively swung from the

dorsal to ventral sides just before flipping. These legs also accelerated more rapidly during the

dorsal-to-ventral stroke than when retracted in the reverse direction, as required for generating

a reaction torque to promote righting (Frohlich 1979). The cross-correlation analysis revealed

strong correlation between the forelegs and midlegs for diagonal rotating and lifted rotating

(maximum similarity mean, full range 0.65 [0.46, 0.94] and 0.64 [0.34, 0.95], respectively), but

lower correlation between the motions of these legs for pitching (maximum similarity 0.45 [0.37,

0.55]). During pitching, the forelegs were swung periodically, consistent with either providing

inertial torque or attempting to grasp the substrate. (See Fig. S9-13 and Table S6 for full

results.)

Table 3 Summary statistics from analysis of the kinematics and dynamics during self-righting. Here, apex refers to the point at which

the insect's potential energy (*PE*) was at a maximum. Data are quoted as mean ± error bars unless otherwise stated. Error bars for

PE were based on tracking errors and those for *KE* based on uncertainty in computing moment of inertia and angular speed. For

other quantities, error bars were computed from 95% bootstrapped confidence intervals (MATLAB *bootci*, 1000 samples) added in

quadrature with the variation between trials, when relevant.

	Diagonal rotating	Lifted rotating	Pitching
Number trials analyzed (specimens)	2 (2)	2 (2)	1 (1)
Active overturning duration (s)	0.078 ± 0.006	0.052± 0.019	0.51± 0.01
$PE_{max} - PE_i$ (J)	$7.7 \pm 1.1 \times 10^{-7}$	$5.8 \pm 0.5 \times 10^{-7}$	$7.0 \pm 1.0 \times 10^{-7}$
Time from Z_{dv} = 0 to apex (ms)	20 ± 5	8 ± 5	102 ± 5
$\omega_{spin} : \omega_{orbital}$			
before active overturning	121 ± 30%	97 ± 14%	75 ± 1%
active overturning to apex	242 ± 31%	118 ± 9%	131 ± 1%
$I_{spin\ legs} : I_{total\ spin}$ (all data)	68 ± 2%	72% ± 2%	65 ± 1%
I_{body} (kg m^2) active overturning to apex	$6.0 \pm 0.2 \times 10^{-118}$	$6.8 \pm 1.2 \times 10^{-11}$	$9.2 \pm 0.2 \times 10^{-11}$
Total torque (Nm) start to apex	$7.3 \pm 0.9 \times 10^{-7}$	$9.1 \pm 1.0 \times 10^{-7}$	$6.6 \pm 0.2 \times 10^{-7}$
Force/leg (BW) start to apex	0.19 ± 0.04	0.92 ± 0.55	0.33 ± 0.02
Moment arm (mm) start to apex	3.7 ± 0.5	2.8 ± 0.8	3.2 ± 0.1

SM/ISM start to apex	76 ± 4%	-58 ± 10%	-361 ± 19%
number leg ground contacts (mean ± SD) start to apex	4.8 ± 0.1	3.5 ± 0.5	3.4 ± 0.1

Fig. 5 Results of analysis of the tracked coordinates and 3D model plotted for diagonal rotating, lifted rotating, and pitching. Note that different time intervals were plotted for each righting mode to display data for approximately equal relative time intervals before and during active overturning (gray shaded region) Fig. S4A,B show plots for the other diagonal rotating and lifted rotating trials, and Fig

S4C shows full results for the pitching trial. (A-C) show the z-component of the dorsal-ventral axis, which transitions from ≈ +1 when

overturned to 0 at the flipping point to a final value ≥ -1 with the body pitched upward in the resting pose. (D-F) show the time

behavior of gravitational potential energy of the whole insect and its body, and available kinetic energy, K_{avail}, computed from the

inverted physical pendulum model (see text for details). These plots also illustrate the increase in potential energy, $\Delta PE(t)$, required

to right at any given time. The plots in (G-I) compare the time dependence of $\Delta PE(t)$, K_{avail} and the righting number, RN, used to

characterize the contribution of inertial reorientation to righting. Note especially that the energy scale for pitching in (I) is

approximately an order of magnitude lower than that for the other two methods. The plots in (J-L) compare the time behavior of the

angular velocity of the body's spin motion along its pitch and roll axes with the orbital angular velocity computed from the inverted

physical pendulum model; see Fig. 3F for the definitions of $\omega_{orbital}$ and ω_{spin}; note that ω_{spin} only includes contributions from roll and

pitch rotations. (M-O) show the motion of the tarsi (feet) along in the z-axis in the body frame, as defined in Fig. 1E. The light gray

vertical line denotes the flipping point (Z_{dv} = 0) when the specimen transitions from overturned, and the dark gray vertical line the

apex of its trajectory when total PE is at its maximum. (Fig. 1I-K)

Fig. 6 The measured pitch and roll angles vs time (black lines) for righting by A, B) diagonal rotating, C,D) lifted rotating, C) pitching

shown superimposed on contour plots and heatmaps showing the gravitational potential energy (PE) landscape for the 3D rendered

model of the spotted lanternfly body computed for each combination of elevation and bank angles with the lowest point on the body

resting on the substrate. The actual measured values of the trajectory are also plotted using markers with colors corresponding to the

computed whole insect potential energy at each time, using the colormap shown. Note that the heatmaps in A-D) were computed

using a 4th instar 3D model and E) for a 3rd instar 3D model.

Discussion

This study showed that determining the motions of all legs as well as the body's orientation

during righting yielded new insights into how insects perform terrestrial self-righting. Specifically,

the complex, agile leg motions observed during diagonal rotating and lifted rotating by spotted

lanternfly nymphs were different from those reported for other insects and proposed for legged

robots (a topic we return to below). The nymphs constantly moved their legs to facilitate

overturning by first raising and tilting their bodies, then pushing and pulling on the substrate

(quasistatic righting), and finally swinging their legs in a coordinated fashion to generate inertial

torques (dynamic self-righting). Each of the three observed righting methods involved a

stereotypic sequence of motions involving all legs, with no one pair of legs playing a

predominant role. These dynamic motions precluded assuming the low-dimensional potential

energy landscape used effectively in earlier studies (Domokos and Várkonyi 2007; Li et al.

2019) because the insect's COM height, and hence the potential energy, depends significantly

on the leg joint angles as well as body orientation and height from the ground (Kessens et al.

2012). While a geometrical model comprising an ellipsoidal body and two legs was sufficient to

describe terrestrial self-righting by discoid cockroaches (Othayoth and Li 2021), a 3D rendered

body model and six articulated legs was necessary to represent the antero-posteriorly

asymmetrical body shape of spotted lanternfly nymphs and the important contributions of their

long, massive legs. Using this anchor model revealed that, for these insects, the legs constitute

such a large fraction of the total mass that their configuration is important for understanding their

dynamic mechanical properties. The relative geometry of the body and legs (i.e., body width to

leg length ratio and the legs' point of insertion relative to the body midline) also impacts an

insect's ability to self-right. For example, a wide, flattened body and relatively short legs can limit

an overturned insect's ability to grasp and push dorsally against the substrate (Faisal and

Matheson 2001). Spotted lanternflies instead have legs long compared to their body width and

the coxae of the fore- and midlegs are located closer to the body outer edges than the midline.

Taken together, this allows them to reach their legs both laterally and dorsally, allowing them to

use their legs effectively, even when overturned, as long levers to facilitate lifting and rotating.

Terrestrial self-righting methods used by spotted lanternfly nymphs had no significant

dependence among success rate (92-100%), time required, number of attempts, substrate, and

life stage. By contrast, the righting strategies used by beetles and stink bugs (Sasaki and

Nonaka 2016; Pace and Harris 2021; Zhang et al. 2021) and the times required for righting

(Zhang et al. 2021) depended significantly on substrate properties. The forces spotted lanternfly

nymphs exerted on the ground by were consistent with those measured for terrestrial self-

righting by discoid cockroaches (≤ 5.1 body weight/leg) (Full et al. 1995), wedging and pulling by

beetles (Evans 1977; Evans and Forsythe 1984), and adhesion on glass by adult spotted

lanternflies (1-2 body weight/leg) (Frantsevich et al. 2008). Spotted lanternflies are known to

cling tenaciously to a wide variety of natural surfaces (Kim et al. 2011; Kane et al. 2021) using a

combination of their tarsal claws and arolia (Avanesyan et al. 2019). Their use of multiple legs

for grasping and adhesion likely provides them with adaptations for righting on the varied

surfaces encountered in their native habitats, which range from smooth, waxy leaves to rugged

ground (Sasaki and Nonaka 2016; Kane et al. 2021).

Spotted lanternfly nymphs favored self-righting by diagonal rotating, in which they used their

legs to pivot about a body contact point, similar to motions used during righting by some

species of beetles and cockroaches (Frantsevich 2004; Li et al. 2019). This method was found

to ensure a greater stability margin and to require a lower force per leg during righting than the

other two methods observed. All three methods were found to be primarily quasistatic until

shortly before the flipping point, when rapid body spin and dynamic swinging leg motions

became important. Spotted lanternfly nymphs also sometimes righted by using their legs to

entirely lift their bodies off the substrate while overturning (lifted rotating), but rarely by pitching--

a method commonly used by some species of beetles, cockroaches and stinkbugs (Frantsevich

2004; Li et al. 2019; Pace and Harris 2021). Pitching was found to correspond to a large,

negative stability margin requiring stabilization by adhesion forces applied by a small number of

legs until immediately before overturning; this means that even a momentary loss of contact

with the ground would destabilize this method before tip-over, as observed during failed pitching

attempts. (Movie 1)

It is also illuminating to compare the righting motions observed for spotted lanternfly nymphs to

those proposed for righting by legged robots (Peng et al. 2017; Wang et al. 2022). The planting

and lifting motions used by these insects correspond to steps required for robots to assume a

starting posture suitable for using the legs to support, balance, and rotate the body. Because

the nymphs cannot passively rotate upright given their flattened body geometry, they must

generate the necessary overturning torque either by using pushing or pulling motions with the

supporting feet or by employing dynamic swinging motions by the legs (Saranli et al. 2004).

Interestingly, the motions used during diagonal rotating resemble those used in "pivoting"

methods for object manipulation by robots, similar to how humans move a heavy object by lifting

and balancing it on an edge (Yoshida et al. 2006). During the recovery phase, the legs

supported and lowered the nymph's body, avoiding jarring and tumbling on impact. After righting

was completed, the legs were repositioned to serve as supports and to assume the typical

resting pose.

We present tracking results for a small number of trials that is consistent with the number

analyzed in earlier studies (Burrows et al. 2015; Othayoth and Li 2021), due to the difficulty of

manually tracking dozens of points over thousands of video frames. Ongoing improvements in

unsupervised tracking methods based on deep learning promise to facilitate gathering larger

tracking datasets for inter- and intraspecies comparisons. This would also allow expanding the

study of terrestrial self-righting to consider a wider range of insect species with diverse body and

foot morphologies (e.g., tarsal claws vs adhesive pads). In future work, it would be interesting to

see if these results for self-righting generalize to a wider range of substrates, including granular,

compliant, and inhomogeneous materials. The analysis methods used in this study can be

applied to a wide range of animals for which legs, tails and long necks are employed during

righting, and therefore promise a new source of inspiration for robotic models of righting. On a

practical note, this study demonstrates the feasibility of creating 3D models of external

morphology using free software packages and low-cost photogrammetry for specimens too

small for 3D scanning, and for studies that do not require the resolution and full 3D imaging

capabilities of microCT. Given that this combination of 3D tracking and modeling has proved

productive for understanding rotational dynamics during insect jumping (Li et al. 2023) and

terrestrial self-righting (this study), these methods could also be extended to explore a wider

range of motions (e.g., modes of locomotion) in combination with 3D prints based on realistic 3D

models (Behm et al. 2018).

825   Funding

826   This work was supported by Haverford College and a National Science Foundation CAREER

827   award to STH (IOS-1453106). Support for participation in this symposium was provided by all

828   divisions of the Society for Integrative and Comparative Biology and the National Science

829   Foundation (IOS-2326876). Funding to pay the Open Access publication charges for this article

830   was provided by Haverford College.

Acknowledgements

832 The authors thank Robert Beyer and Kent Watson for instrumental design and fabrication

833 assistance.

Competing interests

835 No competing interests declared.

Data availability

837 The data underlying this article are available in the article and in its online supplementary

838 material.

References

Al Bitar L, Voigt D, Zebitz CPW, Gorb SN. 2010. Attachment ability of the codling moth Cydia
pomonella L. to rough substrates. Journal of Insect Physiology 56:1966–72.
Avanesyan A, Maugel TK, Lamp WO. 2019. External morphology and developmental changes
of tarsal tips and mouthparts of the invasive spotted lanternfly, Lycorma delicatula
(Hemiptera: Fulgoridae). PLoS One 14.
Behm JE, Waite BR, Hsieh ST, Helmus MR. 2018. Benefits and limitations of three-dimensional
printing technology for ecological research. BMC Ecol 18:32.
Bien T, Alexander BH, White E, Hsieh ST, Kane SA. 2023. Sizing up spotted lanternfly nymphs
for instar determination and growth allometry. PLOS ONE 18:e0265707.
Burrows M, Cullen DA, Dorosenko M, Sutton GP. 2015. Mantises Exchange Angular
Momentum between Three Rotating Body Parts to Jump Precisely to Targets. Current
Biology 25:786–89.
Casarez CS, Fearing RS. 2017. Dynamic terrestrial self-righting with a minimal tail. In: 2017
IEEE/RSJ International Conference on Intelligent Robots and Systems (IROS)
Presented at the 2017 IEEE/RSJ International Conference on Intelligent Robots and
Systems (IROS). p. 314–21.
Davis M, Gouinaud C, Fauroux J-C, Vaslin P. 2011. A review of self-righting techniques for
terrestrial animals. In: International Workshop for Bio-inspired Robots.
Domokos G, Várkonyi PL. 2007. Geometry and self-righting of turtles. Proceedings of the Royal
Society B: Biological Sciences 275:11–17.

Evans MEG. 1977. Locomotion in the Coleoptera Adephaga, especially Carabidae. Journal of Zoology 181:189–226.

Evans MEG, Forsythe TG. 1984. A comparison of adaptations to running, pushing and burrowing in some adult Coleoptera: especially Carabidae. Journal of Zoology 202:513–34.

Faisal AA, Matheson T. 2001. Coordinated righting behaviour in locusts. Journal of Experimental Biology 204:637–48.

Frantsevich L. 2004. Righting kinematics in beetles (Insecta: Coleoptera). Arthropod structure & development 33:221–35.

Frantsevich L, Ji A, Dai Z, Wang J, Frantsevich L, Gorb SN. 2008. Adhesive properties of the arolium of a lantern-fly, Lycorma delicatula (Auchenorrhyncha, Fulgoridae). Journal of Insect Physiology 54:818–27.

Frohlich C. 1979. Do springboard divers violate angular momentum conservation? American Journal of Physics 47:583–92.

Full R, Yamauchi A, Jindrich D. 1995. Maximum single leg force production: cockroaches righting on photoelastic gelatin. Journal of Experimental Biology 198:2441–52.

Full RJ, Koditschek DE. 1999. Templates and anchors: neuromechanical hypotheses of legged locomotion on land. Journal of Experimental Biology 202:3325–32.

Goldstein H. 1980. Classical mechanics Addison-Wesley.

Halliday D, Resnick R, Walker J. 2013. Fundamentals of physics John Wiley & Sons.

Hedrick TL. 2008. Software techniques for two- and three-dimensional kinematic measurements of biological and biomimetic systems. Bioinspir Biomim 3:034001.

Kane SA, Bien T, Contreras-Orendain L, Ochs MF, Tonia Hsieh S. 2021. Many ways to land upright: novel righting strategies allow spotted lanternfly nymphs to land on diverse substrates. Journal of The Royal Society Interface 18:20210367.

Kessens CC, Dotterweich J. 2017. Ground-based self-righting using inertial appendage methods. In: Unmanned Systems Technology XIX Presented at the Unmanned Systems Technology XIX. SPIE. p. 38–50.

Kessens CC, Lennon CT, Collins J. 2014. A metric for self-rightability and understanding its relationship to simple morphologies. In: 2014 IEEE/RSJ International Conference on Intelligent Robots and Systems Presented at the 2014 IEEE/RSJ International Conference on Intelligent Robots and Systems. p. 3699–3704.

Kessens CC, Smith DC, Osteen PR. 2012. A framework for autonomous self-righting of a generic robot on sloped planar surfaces. In: 2012 IEEE International Conference on Robotics and Automation Presented at the 2012 IEEE International Conference on Robotics and Automation. p. 4724–29.

Kim JG, Lee E-H, Seo Y-M, Kim N-Y. 2011. Cyclic Behavior of Lycorma delicatula (Insecta: Hemiptera: Fulgoridae) on Host Plants. J Insect Behav 24:423.

Li C, Kessens CC, Young A, Fearing RS, Full RJ. 2016. Cockroach-inspired winged robot reveals principles of ground-based dynamic self-righting. In: 2016 IEEE/RSJ International Conference on Intelligent Robots and Systems (IROS) Presented at the 2016 IEEE/RSJ International Conference on Intelligent Robots and Systems (IROS). p. 2128–34.

Li C, Wöhrl T, Lam HK, Full RJ. 2019. Cockroaches use diverse strategies to self-right on the ground. Journal of Experimental Biology 222:jeb186080.

Li C, Xu AJ, Beery E, Hsieh ST, Kane SA. 2023. Putting a new spin on insect jumping performance using 3D modeling and computer simulations of spotted lanternfly nymphs. Journal of Experimental Biology 226:jeb246340.

Malashichev YB. 2006. One-sided limb preference is linked to alternating-limb locomotion in anuran amphibians. Journal of Comparative Psychology 120:401–10.

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018. DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. Nature Neuroscience 21:1281–89.

Matt J. 2024. Absolute Orientation - Horn's method. (https://www.mathworks.com/matlabcentral/fileexchange/26186-absolute-orientation-horn-s-method).

McGhee RB, Frank AA. 1968. On the stability properties of quadruped creeping gaits. Mathematical Biosciences 3:331–51.

O'Donnell D. 2018. Mechanics of righting behavior in the tarantula (Theraphosidae) Wake Forest University.

Othayoth R, Li C. 2021. Propelling and perturbing appendages together facilitate strenuous ground self-righting. eLife 10:e60233.

Othayoth R, Xuan Q, Wang Y, Li C. 2021. Locomotor transitions in the potential energy landscape-dominated regime. Proceedings of the Royal Society B: Biological Sciences 288.

Pace CM, Harris M. 2021. Flat on its back: the impact of substrate on righting methods of the brown marmorated stink bug, Halyomorpha halys. J Comp Physiol A.

Pellis VC, Pellis SM, Teitelbaum P. 1991. A descriptive analysis of the postnatal development of contact-righting in rats (Rattus norvegicus). Developmental Psychobiology 24:237–63.

Peng S, Ding X, Yang F, Xu K. 2017. Motion planning and implementation for the self-recovery of an overturned multi-legged robot. Robotica 35:1107–20.

Press WH. 2007. Numerical recipes 3rd edition: The art of scientific computing Cambridge university press.

Robins A, Lippolis G, Bisazza A, Vallortigara G, Rogers LJ. 1998. Lateralized agonistic responses and hindlimb use in toads. Animal Behaviour 56:875–81.

Saranli U, Rizzi AA, Koditschek DE. 2004. Model-Based Dynamic Self-Righting Maneuvers for a Hexapedal Robot. The International Journal of Robotics Research 23:903–18.

Sasaki M, Nonaka T. 2016. The Reciprocity of Environment and Action in Self-Righting Beetles: The Textures of the Ground and an Object, and the Claws. Ecological Psychology 28:78–107.

Schaub H, Junkins JL. 2003. Analytical mechanics of space systems Aiaa.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. Nature Methods 9:671–75.

Semechko A. 2023. Rigid body parameters of closed surface meshesGitHub. (https://github.com/AntonSemechko/Rigid-Body-Parameters).

Theriault DH, Fuller NW, Jackson BE, Bluhm E, Evangelista D, Wu Z, Betke M, Hedrick TL. 2014. A protocol and calibration method for accurate multi-camera field videography. Journal of Experimental Biology 217:1843–48.

Ting LH, Blickhan R, Full RJ. 1994. Dynamic and static stability in hexapedal runners. Journal of Experimental Biology 197:251–69.

Urban JM, Leach H. 2022. Biology and Management of the Spotted Lanternfly, Lycorma delicatula (Hemiptera: Fulgoridae), in the United States. Annual Review of Entomology 68.

Wang K, Xin G, Xin S, Mistry M, Vijayakumar S, Kormushev P. 2023. A unified model with inertia shaping for highly dynamic jumps of legged robots. Mechatronics 95:103040.

Wang S, Wang K, Zhang C, Dai JS. 2022. Kinetostatic backflip strategy for self-recovery of quadruped robots with the selected rotation axis. Robotica 40:1713–31.

Xuan Q, Li C. 2020. Coordinated appendages accumulate more energy to self-right on the ground. IEEE Robotics and Automation Letters 5:6137–44.

Yates BW, Duffy AM. 2008. Statistical analysis of the metrological properties of float glass. In:
Advances in X-Ray/EUV Optics and Components III Presented at the Advances in X-
Ray/EUV Optics and Components III. SPIE. p. 105–15.
Yeadon MR, King MA. 2017. Computer simulation modelling in sport. In: Biomechanical
Evaluation of Movement in Sport and Exercise. 2nd ed Routledge.
Yoshida E, Blazevic P, Hugel V, Yokoi K, Harada K. 2006. Pivoting a large object: whole-body
manipulation by a humanoid robot. Applied Bionics and Biomechanics 3:227–35.
Zhang J, Li J, Li C, Wu Z, Liang H, Wu J. 2021. Self-righting physiology of the ladybird beetle
Coccinella septempunctata on surfaces with variable roughness. Journal of Insect
Physiology 130:104202.
969
970
971