Title: Library preparation method and DNA source influence endogenous DNA recovery from

100-year-old avian museum specimens

Running title: WGS of historical avian museum specimens

Amie E. Settlecowski [†1]

Ben D. Marks[1]

Joseph D. Manthey [†2]

[†]corresponding author

[1]Bird Collection, Field Museum of Natural History, Chicago, Illinois, USA

[2]Department of Biological Sciences, Texas Tech University, Lubbock, Texas, USA

Abstract

Museum specimens collected prior to cryogenic tissue storage are increasingly being used as

genetic resources, and though high throughput sequencing is becoming more cost-efficient,

whole genome sequencing (WGS) of historical DNA (hDNA) remains inefficient and costly due

to its short fragment sizes and high loads of exogenous DNA, among other factors. It is also

unclear how sequencing efficiency is influenced by DNA source. We aimed to identify the most

efficient method and DNA source for collecting WGS data from avian museum specimens. We

analyzed low coverage WGS from 60 DNA libraries prepared from four American Robin (*Turdus*

migratorius) and four Abyssinian Thrush (*Turdus abyssinicus*) specimens collected in the 1920s.

We compared DNA source (toepad versus incision-line skin clip) and three library preparation

methods: 1) double-stranded, single tube (KAPA); 2) single-stranded, multi-tube (IDT); and 3)

single-stranded, single-tube (Claret Bioscience). We found that the multi-tube ssDNA method

resulted in significantly greater endogenous DNA content, average read length, and sequencing

efficiency than the other tested methods. We also tested whether a predigestion step reduced

exogenous DNA in libraries from one specimen per species and found promising results that

warrant further study. The ~10% increase in average sequencing efficiency of the best

performing method over a commonly implemented dsDNA library preparation method has the

potential to significantly increase WGS coverage of hDNA from bird specimens. Future work

should evaluate the threshold for specimen age at which these results hold and how the

combination of library preparation method and DNA source influence WGS in other taxa.

Introduction

Museum specimens collected prior to cryogenic tissue storage have long been used as genetic resources to address questions in ecology, evolutionary biology, and conservation (Habel, Husemann, Finger, Danley, & Zachos, 2014; Wandeler, Hoeck, & Keller, 2007). Genetic studies using these specimens have increased with the advent of high throughput sequencing methods, which in comparison to prior Sanger sequencing methods, drastically increased the data returned from each destructive sampling (Burrell, Disotell, & Bergey, 2015). Now museum specimens commonly facilitate genomic studies via reduced representation (Bi et al., 2013; Linck, Hanna, Sellas, & Dumbacher, 2017; McCormack, Tsai, & Faircloth, 2016) and even whole genome sequencing (e.g. van der Valk, Díez-del-Molino, Marques-Bonet, Guschanski, & Dalén, 2019; Wu et al., 2022). Despite its increasing prevalence and dropping cost, sequencing whole genomes of museum specimens remains expensive because of the degraded nature of the historical DNA (hDNA).

Historical DNA tends to consist of short fragment lengths (McDonough, Parker, McInerney, Campana, & Maldonado, 2018; Straube et al., 2021; Tsai, Schedl, Maley, & McCormack, 2020) that are smaller than the recommended library sizes for the most cost-efficient sequencing setups (i.e., Illumina NovaSeq 6000 S4 flowcell, 200 or 300 cycles). As a result, many sequencing cycles are directly wasted by a lack of base pairs to sequence or indirectly wasted on adapter read through (Straube et al., 2021). Historical DNA libraries also tend to consist of low proportions of DNA from the focal specimen (hereafter endogenous DNA). The rest of the library may consist of exogenous DNA from (1) microbes that have colonized the museum specimen or (2) other environmental microbes, (3) contaminating DNA from researchers or (4) other museum specimens, and (5) more recent DNA samples (Fulton & Shapiro, 2019).

Altogether, the degraded nature of hDNA results in the recovery of lower proportions of

endogenous DNA sequence data (Burrell et al., 2015) and necessitates increased sequencing

effort per specimen to recover similar WGS coverage to modern, high quality DNA libraries.

This inefficiency limits the use of historical DNA from museum specimens to address population

genomic questions that require larger sample sizes in addition to sufficient coverage to address

questions about selection and demography (Lou, Jacobs, Wilder, & Therkildsen, 2021).

Ancient DNA researchers have identified that single-stranded (Gansauge & Meyer, 2013)

and single-tube library preparation methods (Carøe et al., 2018), and those that ligate adapters to

unmodified DNA molecule ends (Kapp, Green, & Shapiro, 2021), increase the amount of

degraded ancient DNA molecules that are converted into genomic libraries. However, the most

implemented ancient library preparation methods are non-proprietary (Gansauge et al., 2017;

Henneberger, Barlow, & Paijmans, 2019), thus requiring a high level of startup effort. Early

ssDNA methods were also more expensive to implement than double-stranded DNA (dsDNA)

libraries, and their improvement in sequencing efficiency did not warrant the additional effort and

cost to implement for all but the most degraded ancient DNA samples (Wales et al., 2015). That

is perhaps why only two studies to date have evaluated the influence of ssDNA versus dsDNA

library preparation on shotgun sequencing of historical specimens. Sproul and Maddison (2017)

found that ssDNA libraries–in comparison to dsDNA libraries–prepared from 16 whole beetle

specimens resulted in more retained reads following quality filtering, but no difference in

endogenous DNA content. Similarly, Hahn et al. (2022) recently found no difference in

endogenous DNA content or insert length between ssDNA and dsDNA libraries prepared from

twelve taxonomically diverse wet collection vertebrate specimens. Additional studies of the

influence of library preparation on WGS of museum specimens that control for taxonomy,

locality, and collection age of specimens will be valuable moving forward. Early ssDNA methods

have been modified to reduce costs and ease implementation (Gansauge et al., 2017) and ssDNA

methods are now commercially available as kits facilitating further study of their impact on

hDNA sequencing efficiency.

Thus far, the majority of research investigating how to maximize the recovery of genetic

data from non-ancient museum specimens has focused on the influence of DNA source or

extraction method on DNA yield (Hahn et al., 2022; Hawkins, Flores, McGowen, & Hinckley,

2022; McDonough et al., 2018; Pacheco et al., 2022; Straube et al., 2021; Tsai et al., 2020; Zacho

et al., 2021). However, DNA yield does not necessarily predict sequencing success or efficiency

(McDonough et al., 2018; Straube et al., 2021; Zacho et al., 2021) because it is not possible to

estimate the proportions of endogenous versus exogenous extracted DNA. For example, a recent

study of hundreds of historical genomic DNA libraries built from samples of museum bird

specimens found that those built from specimens of smaller species (which generally produce

smaller samples) unintuitively had a higher proportion of endogenous sequence data (Irestedt et

al., 2022). Moreover, a few studies have shotgun sequenced DNA from multiple sources on the

same specimen and found differences in endogenous DNA content across sampling sites in fluid-

preserved garter snake specimens (Zacho et al., 2021), prepared mammal skins (McDonough et

al., 2018), and formalin-fixed specimens of a dozen vertebrate taxa (Hahn et al., 2022). Despite

research indicating that differences in hDNA sourced from toepads versus incision-line clips in

bird specimens could influence high-throughput sequencing results, no studies have evaluated

their difference in endogenous DNA content and sequencing efficiency.

Bird study skin specimens have been an especially prolific source of hDNA research

(Billerman & Walsh, 2019) in part due to preservation methods (i.e., skin drying) that are not

catastrophic to DNA preservation relative to methods such as formalin-fixation. Bird study skins

have been the foci of some of the earliest studies of hDNA (Mundy, Unitt, & Woodruff, 1997),

the source for some of the first implementations of reduced representation, high throughput

sequencing methods using museum specimens (Linck et al., 2017; McCormack et al., 2016), and

some of the largest studies implementing WGS of hDNA to date (Irestedt et al., 2022). In this

study we aim to maximize the potential of hDNA from bird study skins by identifying whether

DNA source and library preparation method influence the endogenous DNA content and

sequencing efficiency of hDNA libraries, and by introducing a pre-digestion step prior to DNA

extraction to reduce exogenous DNA.

In this study we test three library preparation methods that vary in 1) the number of

cleanups and tube transfers that occur before the library amplification step (one vs two) and 2)

whether they convert single or double-stranded DNA into library molecules. Each cleanup and

tube transfer is an opportunity to lose DNA molecules of the target length (greater than number

of sequencing cycles) due to the inherent imprecision of SPRI bead cleanups in addition to

pipette error. Methods that are optimized with one, in comparison to two, tube transfers should

transform more DNA molecules of target length into library molecules, thus maximizing library

complexity and sequencing efficiency. Double-stranded DNA library preparation methods cannot

convert ssDNA into library molecules, though as described above, hDNA is expected to consist

in some proportion of single strand molecules due to degradation. Thus, we expect that dsDNA

libraries prepared from hDNA will have reduced sequencing efficiency and possibly endogenous

DNA content compared to that of ssDNA libraries.

We also test the influence of DNA source—toepads versus incision-line skin clips—on

endogenous DNA content and sequencing efficiency. A previous study indicated that toepads

consist of longer DNA fragments than skin clips (Tsai et al., 2020), another possible source of

hDNA from birds (Töpfer, Gamauf, & Haring, 2011). Libraries prepared from samples consisting

of longer DNA fragments should maximize the sequencing capacity, resulting in longer read

lengths on average and greater sequencing efficiency. To test these expectations, we prepared

shotgun DNA libraries from a toepad and skin clip from eight approximately 100-year-old bird

specimens via three methods: 1) double-stranded, single tube (KAPA HyperPrep Kit); 2) single-

stranded, multi-tube (IDT xGen™ ssDNA & Low-Input DNA Library Prep Kit); and 3) single-

stranded, single-tube (Claret Bioscience SRSLY® NanoPlus Kit). We sought to reduce

exogenous DNA by implementing a predigestion step prior to DNA extraction, to our knowledge,

for the first time on bird specimens. To qualitatively evaluate the influence of the predigestion we

also prepared libraries from replicate toepad and skin clip DNA extractions not subjected to

predigestion from two of the eight specimens (Figure 1).

METHODS

2.1. Sampling

We sampled eight bird specimens: four Abyssinian Thrush (*Turdus abyssinicus*; hereafter

thrushes) and four American Robin (*Turdus migratorius*; hereafter robins; Table 1). We chose the

thrush specimens based on their inclusion in another ongoing project and chose to bolster our

sample size for this study with the robin specimens because they are a closely related, similar

species that is well-represented in North American natural history collections. Moreover, the

thrushes were collected in the tropics and the robins were collected in a temperate region which

could influence the drying time of the study skins and in turn, possible degradation due to rot or

the microbial load within dried skin (Irestedt et al., 2022). We chose specimens collected within

one year of each other to control for DNA degradation due to time since specimen preparation.

We collected two samples from each specimen—a toepad and a "skin punch" from the

incision-line through the pectoral apterium (following Tsai et al., 2020)—to evaluate whether

tissue source differed in proportion of endogenous DNA. We also took replicate samples from

one specimen of each species (Table 1) to qualitatively evaluate the effect of sample predigestion

prior to DNA extraction on the exogenous DNA load. The experimental design is summarized in

Figure 1.

We followed stringent sampling precautions to limit the introduction of contaminant

DNA: We (1) wore surgical masks and gloves throughout sampling, (2) took samples in the

collections away from any specimen preparation laboratory, (3) did not enter any modern

molecular DNA or specimen preparation laboratory prior to sampling, (4) prepared the work

surface and all other supplies (e.g., forceps, optivisor, writing utensil) by cleaning with freshly

prepared 10% bleach followed by 70% isopropanol or ethanol, and (5) immediately placed

samples in sterile microcentrifuge tubes that were unpackaged in a sterile lab and not opened

prior to beginning molecular lab work. We used a fresh pair of gloves and sterile scalpel blade for

each sample to minimize contamination between samples.

2.2. Molecular laboratory work

We followed stringent ancient DNA clean lab protocols to minimize contamination

during molecular laboratory work (Fulton & Shapiro, 2019). We completed all pre-PCR steps in

an ancient DNA facility in the Department of Human Genetics at the University of Chicago in a

non-human specific room. We performed each step prior to PCR in a maximum batch size of 12

samples, introduced a negative control in each batch of extractions and library preparations, and

then carried these controls through the remaining steps of lab work.

We extracted DNA via a phenol-chloroform protocol followed by ethanol precipitation

with minor modifications to the protocol presented in Tsai et al. (2020). We performed an NEB

PreCR DNA repair treatment following the sequential reaction protocol on each DNA extraction.

This treatment repairs DNA damage from hydrolysis and oxidative stress, among other

mechanisms, that results in deaminated cytosines, nicks, and other DNA damage incurred with

age. A previous study found that a different NEB repair kit optimized for formalin-fixed

specimens increased the yields of libraries prepared from historical beetle specimens by

approximately 30% (Sproul & Maddison, 2017). Following the damage repair treatment, we

performed a Qiagen MinElute column cleanup and resuspended the DNA in 50 µL of PCR-grade

water. Next, we measured the DNA yield and distribution of DNA fragment sizes using the Qubit

High Sensitivity dsDNA (Thermo Fisher Scientific) and Agilent Bioanalyzer High-Sensitivity

DNA Kit assays, respectively, following the DNA extraction and again following DNA repair

and cleanup. We performed the same assays for each extraction negative control to monitor for

contamination.

We prepared three shotgun sequencing libraries from each DNA extraction and negative

control. Each of the three libraries were prepared via a different method: 1) double-stranded,

single tube (KAPA HyperPrep Kit); 2) single-stranded, multi-tube (IDT xGen™ ssDNA & Low-

Input DNA Library Prep Kit); and 3) single-stranded, single-tube (Claret Bioscience SRSLY®

NanoPlus Kit). We largely followed manufacturer protocols with the following modifications:

during the KAPA adapter ligation step we ligated 25 µM iTru Stubs (Glenn et al., 2019) to each

library molecule. For all cleanups we used a homebrew SPRI bead-solution (Rohland & Reich,

2012) and for each cleanup step in the KAPA and IDT protocols we performed 1.2x SPRI

concentration cleanups. We indexed each library via amplification with 2.5 µM of a unique pair

of iTru5 and iTru7 indexed primers (Glenn et al., 2019) and KAPA HiFi HotStart Uracil+

ReadyMix. For amplification we split each adapter-ligated library into two replicates of 25 µL

each and ran a nine- to twelve-cycle PCR, depending on input DNA amount and method, on the

first replicate; then we estimated the yield of the first replicate via a Qubit High Sensitivity

dsDNA assay and ran a 10- to 12-cycle PCR on the second adapter-ligated library replicate. We

combined amplified replicates for each library and performed a final SPRI cleanup. Finally, we

measured the average molecule size and calculated the concentration of adapter ligated molecules

for each sample library via an Agilent Bioanalyzer High-Sensitivity DNA Kit assay and qPCR

with the KAPA Library Quantification Kit. We submitted a final library pool to Texas Tech

University Center for Biotechnology & Genomics for sequencing. They first checked that

libraries were sequencable with an Illumina MiSeq nano run followed by 100 base pair (bp)

paired-end sequencing on one Illumina NovaSeq SP flowcell.

2.3. Bioinformatics

We received demultiplexed sequence data as raw fastq files from the sequencing facility

and ran FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) to assess quality

and adapter contamination by library preparation method. We trimmed 10 bp from the beginning

of every IDT library read 2 via Seqtk *trimfq* (https://github.com/lh3/seqtk) to remove a low-

complexity polynucleotide tail that facilitates adapter ligation in this method. Duplicate reads

resulting from PCR were identified and removed via Super Deduper

(https://github.com/s4hts/HTStream). Then we used SeqPrep

(https://github.com/jstjohn/SeqPrep) to simultaneously identify adapter contamination and

overlapping paired reads, and then trim adapters and merge reads as necessary. We trimmed

bases from both read ends via four bp sliding window to a minimum quality of 15 and then

removed reads that were less than 30 bp long via Trimmomatic v2.X (Bolger, Lohse, & Usadel,

2014). Finally, we removed any remaining reads that were comprised of more than 50% of one

nucleotide via remove_low_complex.py (distributed as part of the NF-Polish sequence polishing

pipeline described in Irestedt et al. (2022)). We aligned cleaned sequencing reads to the Rufous-

bellied Thrush (*T. rufiventris*) reference genome (ASM1318643v1) via BWA 0.7.17 *mem* (Li,

2013) and indexed mapped reads with Samtools 1.9 *index* (Danecek et al., 2021; Li et al., 2009).

Following sequence cleaning and alignment we evaluated adapter contamination and sequence

quality via FastQC. We used MapDamage 2.0 (Ginolhac, Rasmussen, Gilbert, Willerslev, &

Orlando, 2011; Jónsson, Ginolhac, Schubert, Johnson, & Orlando, 2013) to estimate the

frequency of C to T and G to A misincorporations that result from a transition to uracil during

DNA degradation over time through hydrolysis. For each library we output sequencing metrics

via Samtools 1.9 *stats*.

2.4. Analyses

We tested whether there were differences in DNA yield and mean DNA fragment size

between different sources (toepad vs. skin clip) via paired-t tests. We evaluated whether DNA

source, library preparation method, or an interaction between them resulted in differences in 1)

endogenous DNA content, 2) sequencing efficiency, and 3) mean read length via two-way,

repeated-measures ANOVAs. For any two-way ANOVA that resulted in a significant interaction,

we performed a one-way, repeated measures ANOVA for each method to evaluate whether there

were significant differences by DNA source. For any two-way ANOVA that resulted in a

significant effect of either independent variable, we performed subsequent pairwise, paired-t tests

between all library preparation methods. We accounted for multiple-testing in all post-hoc one-

way and paired-t tests by adjusting p-values via the BH method (Benjamini & Hochberg, 1995).

We expected that samples with larger mean DNA fragment sizes would also have longer

mean read lengths and as a result, greater endogenous DNA content and sequencing efficiency.

To test this hypothesis while controlling for any effect of DNA source and library preparation

method we defined two linear models for each of the following response variables: endogenous

DNA content, sequencing efficiency, and read length. Each null model included the library

preparation method and DNA source as fixed effects and sample as a random effect. The

alternative model also included mean DNA fragment length as a fixed effect. To test whether

mean DNA fragment length had a significant influence on each response variable we performed a

likelihood ratio test of the null and alternative model.

Finally, we sought to qualitatively evaluate the effect of predigestion on replicate

samples. To do so, we plotted the difference in each metric of interest between the replicate

samples. All statistical analyses were completed in R v4.1.0 (R Core Team, 2021). ANOVAs

and t-tests were conducted with the package *rstatix v0.7.0* (Kassambara, 2021), linear mixed

models were built in *lme4* v1.1-27.1 (Bates, Mächler, Bolker, & Walker, 2015), and we used the

suite of functions in *tidyverse* v1.3.1 (Wickham et al., 2019) for data parsing, manipulation, and

visualization.

RESULTS

3.1. DNA Yield and Size

All 60 DNA extractions were successful in terms of producing measurable amounts of

DNA with an average of 589.1 nanograms (ng) per sample and a minimum of 48.6 ng in the skin

clip control replicate from robin specimen 162188 (Table 1). All samples retained enough DNA

through the DNA repair and cleanup to progress to library preparation by each of the three

methods. In general, the DNA repair resulted in an upshift in the distribution of DNA fragment

lengths (Figures S1A and S2A). There was no statistical difference in DNA yield and size

between toepad and skin clips immediately following extraction or after the DNA repair and

cleanup (Figure 2). However, the lack of statistical difference in DNA size is driven by the large

variance in skin clips (Table 2). The mean size of DNA extracted from the toepad sample is

greater than that of the skin clip sample for all but two specimens: robin specimen 83114 and

thrush specimen 66823 (Table 1). For example, these two specimens bias the distribution of the

post-repair skin clip mean size (post-repair M = 315.63, Mdn = 65.49, SD = 651.92) upwards, but

not the toepad mean size (post-repair M = 83.94, Mdn = 83.94, SD = 7.13).

3.2. Endogenous DNA content and sequencing efficiency

Sequencing returned a total of approximately 1.45 billion raw reads and, per library, an

average of 11.23 million raw reads (SD = 1.84) and 9.88 million mapped reads (SD = 9.78) per

library excluding non-predigested replicates. Detailed sequencing metrics for each library are

reported in the supplementary material (Table S1).

There was a significant difference across library-preparation methods, but not DNA

source in endogenous DNA content and sequencing efficiency with IDT outperforming SRSLY

and KAPA in both metrics (Table 4). Similar to the results described above for DNA size, the

toepad samples outperform the corresponding skin clip samples except for specimens 83114 and

66823 (Figure 3A, 3B) so we summarize the results by method and source (Table 2). The average

endogenous DNA content of IDT toepad and skin clip libraries is 88.7% (SD = 0.014) and 81.2%

(SD = 0.140) respectively, 0.06% and 1.4% greater than that of KAPA, and 2.4% and 1% greater

than that of SRSLY. The average sequencing efficiency of IDT toepad and skin clip libraries is

35.9% (SD = 0.045) and 29.8% (SD = 0.245), respectively. In comparison to IDT, KAPA toepad

and skin clip libraries are 11% and 1.8% less efficient, respectively, and SRSLY toepad and skin

clip libraries are 7.8% and 1.8% less efficient. There was also a significant difference among

methods in mean read length with IDT producing longer reads than KAPA and SRSLY with a

significant interaction between method and DNA source (Table 2, Table 4, Figure 3C). IDT

toepad libraries produced significantly longer reads than IDT skin clip libraries (Table 4, Figure

3C).

The tests of the influence of input DNA fragment size on sequencing outcomes produced

mixed results. Including DNA size significantly improved the fit of the linear models for

endogenous DNA content ($\chi^2_1 = 7.558$, p = 0.006), sequencing efficiency ($\chi^2_1 = 45.771$, p <

0.001), and read length ($\chi^2_1 = 25.308$, p < 0.001). However, in the linear models including DNA

size, it was only a significant predictor of endogenous DNA content and read length based on

confidence intervals of the coefficient estimate. In some cases, the trend for the relationship

between DNA size and the dependent variable are contrary to our expectations (Figure S3).

There was evidence of contamination in two of the 20 libraries based on the proportion of

base pair differences between the mapped reads and the reference genome and endogenous DNA

content (Figure S4, Table 1). The skin clip library of thrush specimen 83109 exhibited

approximately 1% more differences from the reference genome than all other samples (Table 1,

Figure S4) in addition to a comparatively low proportion of endogenous DNA content across

preparation methods (IDT = 48.0%, KAPA = 47.7%, SRSLY = 51.9%). The replicate skin clip

library of robin specimen 162188 that was not subjected to predigestion also had a low

proportion of endogenous DNA content across preparation methods (IDT = 17.4%, KAPA =

33.4%, SRSLY = 41.5%), but it was similar to all other libraries in the proportion of mapped

bases that differed from the reference (Figure S4). There was no indication of contamination in

the high performing skin clips samples from specimens 83114 and 66823 (Table 1, Figure S4)

that biased the skin clip averages of most metrics upward.

The influence of sample predigestion was unclear (Table 1, Figure S5). The differences in

DNA yield, mean DNA size, and endogenous DNA content were inconsistent between the

control and predigested replicates from robin 162188 and thrush 83114. The difference between

the replicates in sequencing efficiency and mean read length were marginal.

4. Discussion

4.1. WGS of hDNA from 100-year-old bird study skins

We have demonstrated via shallow sequencing of 60 hDNA libraries that ssDNA library

preparation methods outperform dsDNA methods in sequencing efficiency and, to a lesser extent,

in returning endogenous DNA content from WGS ~100-year-old bird specimens. In contrast to

our predictions, the IDT multi-tube, ssDNA method outperformed the Claret Biosciences single-

tube, ssDNA method and we discuss possible explanations below. We also confirm previous

research suggesting that toepads provide consistently larger DNA fragments and demonstrate that

hDNA from toepads rather than skin clips is a better source for WGS. We show that, though skin

clips may sometimes outperform toepads for a given specimen, toepads have less variance and

therefore less unexpected sequencing outcomes. Altogether, we've shown that toepads are a

better source of DNA and ssDNA library preparations are a better method for collecting WGS

from 100-year old bird specimens. Below we elaborate further on the nuances of our findings and

conclude with broader implications for WGS of historical DNA from museum specimens in

natural history collections.

4.2. Library preparation method and DNA source influenced sequencing

Library preparation method influenced all key metrics; but contrary to our predictions, the

SRSLY ssDNA, single-tube method did not outperform the other two methods. Instead, the IDT

ssDNA, multi-tube method resulted in greater endogenous DNA content, sequencing efficiency,

and read lengths than either of the other methods (Figure 3, Table 4). The margin of difference

between IDT and the other methods was much greater for sequencing efficiency and average read

length than endogenous DNA content (Table 2), suggesting that IDT produced more complex

libraries. The only other study to compare ssDNA and dsDNA methods for shotgun sequencing

of historical specimens prepared libraries from beetle specimens of various ages using the same

ssDNA, multi-tube method that we implemented and a different dsDNA, single-tube method.

Those results showed no difference in endogenous DNA content between methods, and that

ssDNA libraries maintained more sequencing reads through quality filtering and trimming

(Sproul & Maddison, 2017). We find this consistent with our results and suspect that controlling

for taxonomy and specimen age in our experimental design facilitated detecting the small

difference that library preparation method made in endogenous DNA content. That IDT

outperformed SRSLY makes some sense given that SRSLY was originally developed for cell-

free DNA which averages 30 bp long (Troll et al., 2019); though, the commercial kit provides

several versions of the protocol optimized for different purposes, and we implemented the

version for moderate length DNA inserts, less than 200 bp. That protocol for moderate length

DNA inserts includes two-sided SPRI cleanups following adapter ligation and indexing PCR, as

compared to the IDT ssDNA method which uses single-sided cleanups. We suspect that the two-

sided cleanups in SRSLY limited conversion of DNA molecules on the larger end of the DNA

size distribution into library molecules (Figures S1, S2). This is consistent with SRSLY

producing shorter average read lengths than IDT. Still, SRSLY outperformed the KAPA dsDNA

method overall in sequencing efficiency and outperformed IDT in sequencing efficiency for skin

clips in a few specimens (Figure 3A). It is possible that SRSLY may be the better method for

samples that are more degraded as a result of age or DNA source. More recently another ssDNA

single-tube method that builds upon SRSLY was developed specifically for ancient DNA

samples (Kapp et al., 2021); though the nonproprietary status limits ease of implementation. It

may be worthwhile to optimize the SRSLY clean up steps to maximize conversion of the largest

DNA fragments as the IDT method costs $1.89\times$ more than SRSLY per reaction.

Most statistical tests of the effect of DNA source on our metrics of interest were not

significant, though toepads seem to perform better for WGS when considering the influence of

the large variance in the skin clip metrics. Toepads had much smaller variance in DNA yield,

DNA size, endogenous DNA content, sequencing efficiency, and average read length than skin

clips. For six of the eight specimens, toepads clearly provided greater endogenous DNA content,

sequencing efficiency, and average read length than the corresponding skin clip. In general, this

result is reflected by the mean, and more so the median values, of these metrics for toepads as

compared to skin clips (Table 2). That phenol chloroform extraction of toepads does not yield

more DNA but does result in longer DNA fragments than that of skin clips supports previous

research (Tsai et al., 2020). Our results are also in line with previous studies based on five fluid-

preserved garter snake specimens (Zacho et al., 2021) and three dried mammal skins from

different species (McDonough et al., 2018) that showed that different DNA sources have

differences in endogenous DNA content. It is possible that hDNA sampled from bird specimen

toepads produces better WGS data than that of skin clips because of their different structural

makeups. The keratinized, scaly skin of bird feet may provide a better environmental barrier to

water—which promotes DNA degradation via hydrolysis and also overall tissue degradation—

and to invading microbes that would increase exogenous DNA. Toepad cells may also have lower

innate water content due to desiccation during keratinization (Bengtsson et al., 2012). The role of

the keratin structure in maintaining better DNA for WGS is supported by the work of

McDonough and colleagues (2018) which showed that of bone, skin, and claw samples from

dried mammal specimens, claw samples had the highest or near highest proportion of endogenous

DNA and in a qPCR analysis, the highest copy number of nuclear genomic markers.

4.3. Predigestion, high performing samples, and potential contamination

Interpretation of the effect of predigestion on reducing exogenous DNA content and

increasing sequencing efficiency is limited by the small sample size for which we sequenced

replicate predigested and control samples. The lack of clear signal of predigestion effect in DNA

yield and size is unsurprising given the variation across all samples (Figure 2). Moreover, the

lack of any potential signal is unsurprising given that of the eight samples included as replicates

to evaluate predigestion, three were exceptions to the general trends identified by the larger

study. Both skin clips from thrush 83114 were among the few skin clip samples that performed

uncharacteristically better than all other samples and the control skin clip replicate from robin

from 162188 resulted in the low endogenous DNA content of all libraries (Table 1) indicating

high levels of exogenous DNA or contamination. Notably, the predigested skin clip replicate

sample from robin 162188 did not show signs of contamination suggesting that predigestion may

have reduced exogenous DNA in only two minutes of predigestion time. In contrast, the skin clip

from thrush specimen 83109 was one of four samples with the shortest, one-minute predigestion

and also showed a clearer signal of contamination based on low endogenous DNA content and a

larger genetic distance from the reference than other samples (Table 1, Figure S4). Finally, it is

likely that predigestion reduced total DNA yield, though enough DNA remained for all 20

samples to prepare three successfully sequenced libraries. Altogether we suggest this preliminary

investigation is a promising avenue to maximize endogenous hDNA from museum bird

specimens for WGS and warrants further research.

The primary source of the larger variance in most metrics for skin clips were two

specimens for which the skin clips not only outperformed the corresponding toepad from the

same specimen but all other samples. Robin 66823 and thrush 83114 returned the first and second

largest DNA yield, DNA size, endogenous DNA content, average read length, and highest

sequencing efficiency. Notably, thrush 83114 was one of the two specimens included in the

predigestion study, and both the predigested and non-predigested skin clips were high performing

samples and had consistent values across metrics. There are multiple explanations for why the

samples outperformed all others, the first being contamination. However, neither of these samples

show clear signs of contamination like the lower endogenous DNA content or large genetic

distance from the reference genome mentioned for the skin clip from thrush 83109. The only

explanation of contamination we consider plausible is contamination by modern DNA from the

same or a closely related species, perhaps from a more recently collected specimen in the same

drawer. This possibility highlights the need for stringent sampling procedures during sample

collection from museum specimens for hDNA purposes. Importantly, a population genomics

study involving deeper sequencing of these samples would allow assembly of mitochondrial

genomes that would enable identification of multiple individuals within one sequencing library.

Another possible explanation for the higher performance of these skin clips is that these

specimens received different treatments at the time of collection than the other specimens in the

study. At the time of collection it was common practice to treat bird (and mammal) skins with

arsenic-containing solutions for tanning as well as protection from pests (Marte, Péquignot, &

Von Endt, 2006), and arsenic has been demonstrated as a DNA polymerase inhibitor (Töpfer et

al., 2011). It is possible that the specimens with high performing skin clips were accidentally

skipped in some treatment that ultimately promoted DNA damage in the other skins. The last

and, in our opinion, most likely possibility is that this variation represents real variation in DNA

quality between specimens. Such large variations are not uncommon when working with hDNA

(e.g. McDonough et al., 2018) and ancient DNA (e.g. Wales et al., 2015) and highlights the value

of identifying methods and DNA sources that can consistently return an expected amount of

WGS data like we have found for toepads and the IDT ssDNA, single-tube method.

5. Conclusion

We have shown that for 100-year old museum bird study skin specimens, of those

combinations we tested, the combination of toepads and ssDNA library preparation, in this case

the IDT xGen™ ssDNA & Low-Input DNA Library Prep Kit, provide the best WGS data. Our

results regarding toepads, in combination with previous studies of endogenous DNA content in

other taxa, can be reasonably extended beyond birds to suggest that keratinous sources of hDNA

may be the best source for WGS and should motivate additional investigations of this hypothesis.

We also have shown that when comparing WGS from ssDNA and dsDNA methods, ssDNA

methods provide a larger increase in sequencing efficiency than endogenous DNA content,

suggesting that they successfully convert more hDNA molecules into sequenceable library

molecules and likely lead to more complex libraries better suited for WGS at the depth required

for population genomic studies. Further study of the impact of library preparation method on

sequencing efficiency that controls for variation among specimens and also evaluates the role of

age of the specimen is necessary to identify the threshold at which an ssDNA method is or is not

warranted. Also, it may be worthwhile to attempt to further optimize SRSLY cleanups to

minimize bias against converting larger fragment hDNA molecules into library molecules as

SRSLY is currently ~89% less expensive per reaction than the IDT method. Finally, our

inclusion of a predigestion step to reduce external exogenous DNA did not yield straightforward

results, though it did provide some evidence that it limited contamination in one of four samples

for which we made a direct comparison. Importantly we showed that a brief (less than 15

minute), gentle (37C° incubation) predigestion step does not preclude successful library

construction, and thus we will cautiously include this step in our own protocols moving forward.

Altogether this study identifies how to maximize WGS data collected from 100-year old bird

specimens and provides some general insights on how to increase the quality and quantity of

WGS data recovered from hDNA of museum specimens overall.

Acknowledgments

This work was supported by NSF Grant No. 1953796 to JDM and BDM. We would like to thank

Dr. Manaasa Raghavan for providing access to the ancient DNA facility at the University of

Chicago and the Raghavan Genoscape Lab, especially Dr. Constanza de la Fuentes, for

facilitating our pre-PCR lab work. Mohamed Fokar at the TTU Center for Biotechnology &

Genomics provided sequencing support. The TTU Center for Biotechnology & Genomics

acquisition of the NovaSeq6000 was supported by NIH grant 1S10OD025115-01. We

acknowledge the High Performance Computing Center (HPCC) at Texas Tech University

(http://www.hpcc.ttu.edu) and the Pritzker DNA Lab at Field Museum of Natural History for

providing computational and molecular resources, respectively, that have contributed to the

research results reported within this paper.

Conflict of interest statement

IDT has been a corporate sponsor of the Field Museum of Natural History as recently as 2020,

but no funds from IDT were used to directly support the research that is the subject of the

publication.

References

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models
using lme4. *Journal of Statistical Software, 67*(1), 1-48. doi:10.18637/jss.v067.i01
Bengtsson, C. F., Olsen, M. E., Brandt, L. Ø., Bertelsen, M. F., Willerslev, E., Tobin, D. J., . . .
Gilbert, M. T. P. (2012). DNA from keratinous tissue. Part I: Hair and nail. *Annals of*
Anatomy, 194(1), 17-25. doi:10.1016/j.aanat.2011.03.013
Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*
(Methodological), 57(1), 289-300. doi:10.1111/j.2517-6161.1995.tb02031.x
Bi, K., Linderoth, T., Vanderpool, D., Good, J. M., Nielsen, R., & Moritz, C. (2013). Unlocking
the vault: next-generation museum population genomics. *Molecular Ecology, 22*(24),
6018-6032. doi:10.1111/mec.12516
Billerman, S. M., & Walsh, J. (2019). Historical DNA as a tool to address key questions in avian
biology and evolution: a review of methods, challenges, applications, and future
directions. *Molecular Ecology Resources, 19*(5), 1115-1130. doi:10.1111/1755-
0998.13066
Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics, 30*(15), 2114-2120. doi:doi:10.1093/bioinformatics/btu170
Burrell, A. S., Disotell, T. R., & Bergey, C. M. (2015). The use of museum specimens with high-
throughput DNA sequencers. *Journal of Human Evolution, 79*, 35-44.
doi:10.1016/j.jhevol.2014.10.015
Carøe, C., Gopalakrishnan, S., Vinner, L., Mak, S. S. T., Sinding, M. H. S., Samaniego, J. A., . . .
Gilbert, M. T. P. (2018). Single-tube library preparation for degraded DNA. *Methods in*
Ecology and Evolution, 9(2), 410-419. doi:10.1111/2041-210X.12871

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., . . . Li, H. (2021). Twelve years of SAMtools and BCFtools. *GigaScience, 10*(2). doi:10.1093/gigascience/giab008

Fulton, T. L., & Shapiro, B. (2019). Setting up an Ancient DNA Laboratory. In B. Shapiro, A. Barlow, P. D. Heintzman, M. Hofreiter, J. L. A. Paijmans, & A. E. R. Soares (Eds.), Ancient DNA: Methods and Protocols. Methods in Molecular Biology (Vol. 1963, pp. 1-15). doi:10.1007/978-1-4939-9176-1_1

Gansauge, M.-T., Gerber, T., Glocke, I., Korlević, P., Lippik, L., Nagel, S., . . . Meyer, M. (2017). Single-stranded DNA library preparation from highly degraded DNA using T4 DNA ligase. *Nucleic Acids Research, 45*(10), e79. doi:10.1093/nar/gkx033

Gansauge, M.-T., & Meyer, M. (2013). Single-stranded DNA library preparation for the sequencing of ancient or damaged DNA. *Nature Protocols, 8*(4), 737-748. doi:10.1038/nprot.2013.038

Ginolhac, A., Rasmussen, M., Gilbert, M. T. P., Willerslev, E., & Orlando, L. (2011). mapDamage: testing for damage patterns in ancient DNA sequences. *Bioinformatics, 27*(15), 2153-2155. doi:10.1093/bioinformatics/btr347

Glenn, T. C., Nilsen, R. A., Kieran, T. J., Sanders, J. G., Bayona-Vásquez, N. J., Finger, J. W., . . . Faircloth, B. C. (2019). Adapterama I: universal stubs and primers for 384 unique dual-indexed or 147,456 combinatorially-indexed Illumina libraries (iTru & iNext). *PeerJ, 7*, e7755. doi:10.7717/peerj.7755

Habel, J. C., Husemann, M., Finger, A., Danley, P. D., & Zachos, F. E. (2014). The relevance of time series in molecular ecology and conservation biology. *Biological Reviews, 89*(2), 484-492. doi:10.1111/brv.12068

Hahn, E. E., Alexander, M. R., Grealy, A., Stiller, J., Gardiner, D. M., & Holleley, C. E. (2022). Unlocking inaccessible historical genomes preserved in formalin. *Molecular Ecology Resources, 22*(6), 2130-2147. doi:10.1111/1755-0998.13505

Hawkins, M. T. R., Flores, M. F. C., McGowen, M., & Hinckley, A. (2022). A comparative analysis of extraction protocol performance on degraded mammalian museum specimens. *Frontiers in Ecology and Evolution, 10*, 984056. doi:10.3389/fevo.2022.984056

Henneberger, K., Barlow, A., & Paijmans, J. L. A. (2019). Double-Stranded Library Preparation for Ancient and Other Degraded Samples. In B. Shapiro, A. Barlow, P. D. Heintzman, M. Hofreiter, J. L. A. Paijmans, & A. E. R. Soares (Eds.), Ancient DNA: Methods and Protocols. Methods in Molecular Biology (Vol. 1963, pp. 65-73). New York, NY: Springer New York. doi:10.1007/978-1-4939-9176-1_8

Irestedt, M., Thörn, F., Müller, I. A., Jønsson, K. A., Ericson, P. G. P., & Blom, M. P. K. (2022). A guide to avian museomics: Insights gained from resequencing hundreds of avian study skins. *Molecular Ecology Resources*. doi:10.1111/1755-0998.13660

Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P. L. F., & Orlando, L. (2013). mapDamage2.0: fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics, 29*(13), 1682-1684. doi:10.1093/bioinformatics/btt193

Kapp, J. D., Green, R. E., & Shapiro, B. (2021). A fast and efficient single-stranded genomic library preparation method optimized for ancient DNA. *Journal of Heredity, 112*(3), 241-249. doi:10.1093/jhered/esab012

Kassambara, A. (2021). rstatix: Pipe-Friendly Framework for Basic Statistical Tests. In: R package version 0.7.0. https://CRAN.R-project.org/package=rstatix.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv. Quantitative Biology*, 1303.3997. doi:arXiv:1303.3997

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., . . . Subgroup, G. P. D. P. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics, 25*(16), 2078-2079. doi:10.1093/bioinformatics/btp352

Linck, E. B., Hanna, Z. R., Sellas, A., & Dumbacher, J. P. (2017). Evaluating hybridization capture with RAD probes as a tool for museum genomics with historical bird specimens. *Ecology and Evolution, 7*(13), 4755-4767. doi:10.1002/ece3.3065

Lou, R. N., Jacobs, A., Wilder, A. P., & Therkildsen, N. O. (2021). A beginner's guide to low-coverage whole genome sequencing for population genomics. *Molecular Ecology, 30*(23), 5966-5993. doi:10.1111/mec.16077

Marte, F., Péquignot, A., & Von Endt, D. W. (2006). *Arsenic in taxidermy collections: history, detection, and management.* Paper presented at the Collection Forum.

McCormack, J. E., Tsai, W. L., & Faircloth, B. C. (2016). Sequence capture of ultraconserved elements from bird museum specimens. *Molecular Ecology Resources, 16*(5), 1189-1203.

McDonough, M. M., Parker, L. D., McInerney, N. R., Campana, M. G., & Maldonado, J. E. (2018). Performance of commonly requested destructive museum samples for mammalian genomic studies. *Journal of Mammalogy, 99*(4), 789-802. doi:10.1093/jmammal/gyy080

Mundy, N. I., Unitt, P., & Woodruff, D. S. (1997). Skin from feet of museum specimens as a non-destructive source of DNA for avian genotyping. *The Auk, 114*(1), 126-129.

Pacheco, C., Lobo, D., Silva, P., Álvares, F., García, E. J., Castro, D., . . . Godinho, R. (2022). Assessing the performance of historical skins and bones for museomics using wolf specimens as a case study. *Frontiers in Ecology and Evolution, 10*, 970249. doi:10.3389/fevo.2022.970249

Rohland, N., & Reich, D. (2012). Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture. *Genome Research, 22*(5), 939-946. doi:10.1101/gr.128124.111

Sproul, J. S., & Maddison, D. R. (2017). Sequencing historical specimens: successful preparation of small specimens with low amounts of degraded DNA. *Molecular Ecology Resources, 17*(6), 1183-1201. doi:10.1111/1755-0998.12660

Straube, N., Lyra, M. L., Paijmans, J. L. A., Preick, M., Basler, N., Penner, J., . . . Hofreiter, M. (2021). Successful application of ancient DNA extraction and library construction protocols to museum wet collection specimens. *Molecular Ecology Resources, 21*(7), 2299-2315. doi:10.1111/1755-0998.13433

Team, R. C. (2021). R: A language and environment for statistical computing. In. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Töpfer, T., Gamauf, A., & Haring, E. (2011). Utility of arsenic-treated bird skins for DNA extraction. *BMC research notes, 4*(1), 197. doi:10.1186/1756-0500-4-197

Troll, C. J., Kapp, J., Rao, V., Harkins, K. M., Cole, C., Naughton, C., . . . Green, R. E. (2019). A ligation-based single-stranded library preparation method to analyze cell-free DNA and synthetic oligos. *Bmc Genomics, 20*(1). doi:10.1186/s12864-019-6355-0

Tsai, W. L. E., Schedl, M. E., Maley, J. M., & McCormack, J. E. (2020). More than skin and bones: Comparing extraction methods and alternative sources of DNA from avian museum specimens. *Molecular Ecology Resources, 20*(5), 1220-1227. doi:10.1111/1755-0998.13077

van der Valk, T., Díez-del-Molino, D., Marques-Bonet, T., Guschanski, K., & Dalén, L. (2019).
Historical Genomes Reveal the Genomic Consequences of Recent Population Decline in
Eastern Gorillas. *Current Biology, 29*(1), 165-170.e166.
doi:https://doi.org/10.1016/j.cub.2018.11.055
Wales, N., Carøe, C., Sandoval-Velasco, M., Gamba, C., Barnett, R., Samaniego, J. A., . . .
Gilbert, M. T. P. (2015). New insights on single-stranded versus double-stranded DNA
library preparation for ancient DNA. *BioTechniques, 59*(6), 368-371.
doi:10.2144/000114364
Wandeler, P., Hoeck, P. E., & Keller, L. F. (2007). Back to the future: museum specimens in
population genetics. *Trends in Ecology & Evolution, 22*(12), 634-642.
doi:10.1016/j.tree.2007.08.017
Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D. A., François, R., . . . Yutani,
H. (2019). Welcome to the tidyverse. *Journal of open source software, 4*(43), 01686.
doi:10.21105/joss.01686
Wu, M. Y., Lau, C. J., Ng, E. Y. X., Baveja, P., Gwee, C. Y., Sadanandan, K., . . . Rheindt, F. E.
(2022). Genomes from historic DNA unveil massive hidden extinction and terminal
endangerment in a tropical Asian songbird radiation. *Molecular Biology and Evolution*.
doi:10.1093/molbev/msac189
Zacho, C. M., Bager, M. A., Margaryan, A., Gravlund, P., Galatius, A., Rasmussen, A. R., &
Allentoft, M. E. (2021). Uncovering the genomic and metagenomic research potential in
old ethanol-preserved snakes. *PLOS ONE, 16*(8), e0256353.
doi:10.1371/journal.pone.0256353

Data Accessibility and Benefit-Sharing

Raw sequence reads will be deposited in NCBI Sequence Read Archive (SRA) (BioProject

XXX). Individual bam files and code used in data analysis for this project are available on Dryad

(XXXX). Code used in data analysis is also available at: github.com/amiesett/WGS-hDNA-birds.

Benefit sharing: We are sharing all research benefits of this project by providing raw data,

bioinformatic scripts, and results in public databases. No monetary benefits are expected from

these research results.

Author Contributions

622   All authors contributed to project design and manuscript completion. JDM and BDM acquired

623   funding. AES performed lab work and data analysis, with support from JDM, and wrote the first

624   draft of the manuscript.

Table 1. Summary of extracted DNA and sequencing results for each sample.

Species	Sample	Post-extraction Yield (ng/mg)	Post-extraction Mean DNA size (bp)	Post-repair Yield (ng/mg)	Post-repair Mean DNA size (bp)	Method	Type	Raw reads (millions)	Clean reads (millions)	Mapped reads (millions)	Mean read length	Dist.to ref.	Endog. DNA content	Seq. eff.
T. abyssinicus	83107					IDT	skin	10.54	3.38	2.96	71.99	0.018	0.851	0.193
T. abyssinicus	83107	400.03	57.36	136.10	68.02	KAPA	skin	60.29	15.17	12.45	59.75	0.014	0.820	0.119
T. abyssinicus	83107					SRSLY	skin	12.85	5.23	4.64	75.14	0.015	0.822	0.267
T. abyssinicus	83107					IDT	toepad	12.62	5.51	5.12	97.84	0.017	0.900	0.382
T. abyssinicus	83107	698.97	65.75	332.89	83.90	KAPA	toepad	85.06	28.05	25.05	78.48	0.012	0.889	0.224
T. abyssinicus	83107					SRSLY	toepad	8.81	3.36	2.89	67.02	0.017	0.853	0.213
T. abyssinicus	83109					IDT	skin	11.26	3.29	1.69	62.71	0.035	0.480	0.086
T. abyssinicus	83109	376.78	57.93	73.28	61.74	KAPA	skin	50.69	12.09	5.77	53.49	0.027	0.477	0.057
T. abyssinicus	83109					SRSLY	skin	10.08	3.74	2.13	65.56	0.030	0.519	0.119
T. abyssinicus	83109					IDT	toepad	10.95	4.56	4.18	88.02	0.018	0.896	0.323
T. abyssinicus	83109	858.85	66.68	338.13	85.08	KAPA	toepad	63.05	21.06	18.86	79.73	0.013	0.890	0.232
T. abyssinicus	83109					SRSLY	toepad	9.29	3.46	2.97	66.34	0.017	0.853	0.206
T. abyssinicus	83114					IDT	skin	46.55	32.78	30.90	100.17	0.009	0.916	0.640
T. abyssinicus	83114	1350.36	78.71	576.64	218.79	KAPA	skin	32.03	15.21	14.27	109.38	0.013	0.919	0.472
T. abyssinicus	83114					SRSLY	skin	9.56	5.06	4.69	104.24	0.017	0.903	0.503
T. abyssinicus	83114					IDT	toepad	24.48	10.02	8.80	89.44	0.016	0.856	0.309
T. abyssinicus	83114	800.00	53.96	55.17	98.74	KAPA	toepad	45.38	15.44	13.93	83.94	0.015	0.889	0.250
T. abyssinicus	83114					SRSLY	toepad	7.62	3.34	3.01	85.32	0.018	0.874	0.332
T. abyssinicus	83114[†]					IDT	skin	42.34	28.30	26.73	101.24	0.009	0.917	0.615
T. abyssinicus	83114[†]	4254.55	96.90	1872.73	211.98	KAPA	skin	34.26	15.74	14.75	107.05	0.013	0.919	0.447
T. abyssinicus	83114[†]					SRSLY	skin	9.35	4.80	4.44	101.67	0.017	0.902	0.474
T. abyssinicus	83114[†]					IDT	toepad	28.09	11.70	10.77	99.56	0.015	0.891	0.368
T. abyssinicus	83114[†]	1901.96	56.07	202.94	107.53	KAPA	toepad	62.51	22.23	20.22	84.52	0.013	0.899	0.266
T. abyssinicus	83114[†]					SRSLY	toepad	14.23	6.31	5.75	85.21	0.016	0.876	0.344

Species	Sample	Post-extraction		Post-repair		Method	Type	Raw reads (millions)	Clean reads (millions)	Mapped reads (millions)	Mean read length	Dist.to ref.	Endog. DNA content	Seq. eff.
		Yield (ng/mg)	Mean DNA size (bp)	Yield (ng/mg)	Mean DNA size (bp)									
T. abyssinicus	83115					IDT	skin	14.52	4.28	3.80	78.74	0.017	0.858	0.197
T. abyssinicus	83115	942.98	56.71	526.94	69.01	KAPA	skin	80.02	19.29	16.20	59.36	0.014	0.843	0.117
T. abyssinicus	83115					SRSLY	skin	11.55	3.30	2.65	54.80	0.015	0.803	0.120
T. abyssinicus	83115					IDT	toepad	43.02	17.60	16.22	94.73	0.013	0.893	0.345
T. abyssinicus	83115	564.00	59.43	217.47	83.13	KAPA	toepad	14.25	4.92	4.37	78.08	0.018	0.881	0.232
T. abyssinicus	83115					SRSLY	toepad	12.70	5.55	5.02	81.83	0.015	0.862	0.324
T. migratorius	162188					IDT	skin	15.77	4.77	3.97	60.97	0.018	0.822	0.146
T. migratorius	162188	375.19	52.58	597.41	58.71	KAPA	skin	2.14	0.74	0.60	60.52	0.018	0.803	0.162
T. migratorius	162188					SRSLY	skin	9.77	3.52	2.94	61.09	0.018	0.823	0.176
T. migratorius	162188					IDT	toepad	21.99	9.50	8.58	95.19	0.016	0.879	0.354
T. migratorius	162188	501.80	57.60	436.09	76.70	KAPA	toepad	10.85	4.53	3.96	78.38	0.019	0.866	0.275
T. migratorius	162188					SRSLY	toepad	8.36	3.60	3.15	76.19	0.019	0.868	0.279
T. migratorius	162188[†]					IDT	skin	6.56	4.11	0.86	80.05	0.020	0.174	0.100
T. migratorius	162188[†]	173.57	54.16	47.37	62.90	KAPA	skin	14.98	6.38	2.90	59.05	0.018	0.334	0.110
T. migratorius	162188[†]					SRSLY	skin	8.85	3.79	2.07	63.26	0.017	0.415	0.135
T. migratorius	162188[†]					IDT	toepad	9.14	3.95	3.55	89.65	0.019	0.875	0.333
T. migratorius	162188[†]	231.03	59.13	98.59	77.89	KAPA	toepad	31.17	12.31	10.86	78.33	0.017	0.873	0.263
T. migratorius	162188[†]					SRSLY	toepad	10.20	4.42	3.92	78.11	0.019	0.870	0.293
T. migratorius	175413					IDT	skin	8.18	2.72	2.31	78.30	0.019	0.807	0.208
T. migratorius	175413	197.25	50.97	59.33	62.37	KAPA	skin	15.67	4.99	3.86	60.91	0.018	0.770	0.144
T. migratorius	175413					SRSLY	skin	11.30	4.57	3.84	70.22	0.017	0.792	0.227
T. migratorius	175413					IDT	toepad	12.34	6.03	5.66	100.66	0.017	0.894	0.442
T. migratorius	175413	501.09	56.22	219.23	86.23	KAPA	toepad	32.87	13.07	11.71	81.68	0.016	0.882	0.280
T. migratorius	175413					SRSLY	toepad	7.51	3.53	3.24	86.20	0.017	0.868	0.372

Species	Sample	Post-extraction		Post-repair		Method	Type	Raw reads (millions)	Clean reads (millions)	Mapped reads (millions)	Mean read length	Dist.to ref.	Endog. DNA content	Seq. eff.
		Yield (ng/mg)	Mean DNA size (bp)	Yield (ng/mg)	Mean DNA size (bp)									
T. migratorius	175421					IDT	skin	6.68	2.31	1.97	64.10	0.019	0.840	0.180
T. migratorius	175421	463.63	52.32	111.02	62.95	KAPA	skin	56.18	17.57	14.70	63.39	0.015	0.833	0.159
T. migratorius	175421					SRSLY	skin	14.82	5.83	5.05	67.82	0.017	0.838	0.222
T. migratorius	175421					IDT	toepad	21.44	9.64	8.98	98.88	0.016	0.891	0.398
T. migratorius	175421	838.10	60.18	234.00	82.44	KAPA	toepad	51.02	19.65	17.51	78.30	0.015	0.879	0.259
T. migratorius	175421					SRSLY	toepad	7.29	3.33	3.03	83.03	0.018	0.868	0.343
T. migratorius	66823					IDT	skin	60.98	51.78	49.16	95.45	0.007	0.920	0.736
T. migratorius	66823	1150.86	387.24	950.45	1923.43	KAPA	skin	35.56	24.12	22.91	106.55	0.010	0.921	0.659
T. migratorius	66823					SRSLY	skin	9.88	6.23	5.87	105.58	0.017	0.917	0.605
T. migratorius	66823					IDT	toepad	5.13	2.24	2.03	85.15	0.020	0.885	0.321
T. migratorius	66823	223.70	53.26	47.36	75.28	KAPA	toepad	50.77	19.75	17.26	73.40	0.015	0.871	0.241
T. migratorius	66823					SRSLY	toepad	11.86	4.31	3.63	61.16	0.018	0.855	0.182

[†]Extraction replicate not subjected to predigestion

628 **Table 2.** Summary (median, mean, standard deviation) of DNA yield and mean DNA size for predigested toepads and skin clips.

	Yield (ng/mg)				Mean size (bp)			
	Post-extraction		Post-repair		Post-extraction		Post-repair	
Source	Mdn	M (SD)	Mdn	M (SD)	Mdn	M (SD)	Mdn	M (SD)
toepad (n=8)	631.49	623.314 (217.606)	226.62	235.042 (135.626)	58.52	59.136 (4.985)	83.51	83.937 (7.132)
skin (n=8)	431.83	657.135 (427.479)	331.52	378.896 (329.864)	57.04	99.227 (116.707)	65.49	315.628 (651.918)

Table 3. Summary (median, mean, standard deviation) of sequencing metrics for predigested toepads and skin clips.

	Mean read length (bp)						Endogenous DNA content						Sequencing efficiency					
	IDT		KAPA		SRSLY		IDT		KAPA		SRSLY		IDT		KAPA		SRSLY	
Source	Mdn	M (SD)	Mdn	M (SD)	Mdn	M (SD)	Mdn	M (SD)	Mdn	M (SD)	Mdn	M (SD)	Mdn	M (SD)	Mdn	M (SD)	Mdn	M (SD)
toepad (n=8)	94.959	93.74 (5.598)	78.428	78.998 (3.059)	79.005	75.883 (9.771)	0.892	0.887 (0.014)	0.882	0.881 (0.009)	0.865	0.863 (0.008)	0.35	0.359 (0.045)	0.245	0.249 (0.021)	0.302	0.281 (0.072)
skin (n=8)	75.145	76.554 (14.799)	60.717	71.669 (22.587)	69.018	75.557 (19.097)	0.846	0.812 (0.14)	0.827	0.798 (0.14)	0.823	0.802 (0.123)	0.195	0.298 (0.245)	0.151	0.236 (0.212)	0.225	0.28 (0.179)

Table 4. Summary of the statistical analyses of endogenous DNA content, sequencing efficiency, and mean read length, and their results. All the p-values presented for t-tests are corrected for multiple testing via the Benjamini and Hochberg (1995) method.

Response variable	Test	DF	Test-statistic	p-value
Endogenous DNA content	ANOVA - source	1,7	0.066	0.805
	ANOVA - method	2,14	6.241	0.012
	ANOVA - source:method	2,14	0.739	0.496
	T-test - IDT vs. KAPA	15	6.631	< 0.001
	T-test - IDT vs. SRSLY	15	2.476	0.026
	T-test - KAPA vs. SRSLY	15	-2.570	0.026
Sequencing efficiency	ANOVA - source	1,7	0.066	0.805
	ANOVA - method	2,14	6.241	0.012
	ANOVA - source:method	2,14	0.739	0.496
	T-test - IDT vs. KAPA	15	2.492	0.037
	T-test - IDT vs. SRSLY	15	2.765	0.037
	T-test - KAPA vs. SRSLY	15	1.283	0.219
Mean read length (bp)	ANOVA - source	1,7	1.333	0.286
	ANOVA - method	2,14	14.181	< 0.001
	ANOVA - source:method	2,14	6.778	0.009
	T-test - IDT vs. KAPA	15	3.879	0.004
	T-test - IDT vs. SRSLY	15	3.001	0.013
	T-test - KAPA vs. SRSLY	15	-0.184	0.856
Mean read length (bp) - IDT	ANOVA - source	1,7	7.179	0.096
Mean read length (bp) - KAPA	ANOVA - source	1,7	0.818	1.0000
Mean read length (bp) - SRSLY	ANOVA - source	1,7	0.002	1.0000

Figure 1. Study sampling design. The illustration on the left shows where incision-line skin clips and toepads are sampled from the specimens. The flowchart on the right is an overview of the experimental design and the laboratory procedures. Toepads and skin clips are depicted as triangular and trapezoidal shapes, respectively. Photographs by JDM and illustrations by L. Nassef.

Figure 2. Source tissue impacts on DNA quantity and quality. For each sample, DNA yield (ng/mg) measured A) following DNA extraction and B) following DNA repair and cleanup is plotted by DNA source (toepad or skin clip). For each sample, mean DNA fragment size (bp) C) following DNA extraction and D) following DNA repair are plotted by DNA source. Samples from robin specimen 66823 are not plotted in C and D because the mean DNA size of its skin clip (Table 1) limits visualization of the variation in the size of the other samples. Gray lines connect toepad and skin clip data points from the same specimen. Summary statistics are also plotted by source: the mean and median are represented by dashed and solid lines, respectively, the upper and lower limits of the boxes represent the 75% and 25% quantiles, and the error bars represent the standard deviation.

Figure 3. Association of sequencing library type and sequence data characteristics. For each library, A) sequencing efficiency, or the proportion of raw bases that uniquely mapped to the reference genome, B), endogenous DNA content, or the proportion of cleaned bases that uniquely mapped to the reference genome, and C) the mean read length (bp) is plotted by DNA source (toepad or skin clip). Gray lines connect toepad and skin clip data points from the same specimen. Summary statistics are also plotted by method and source: the mean and median are represented by dashed and solid lines, respectively, the upper and lower limits of the boxes represent the 75% and 25% quantiles, and the error bars represent the standard deviation.