Horizontal gene transfer and CRISPR targeting drive phage-bacterial host interactions and co-

evolution in pink berry marine microbial aggregates

Running title: Phage-host co-evolution in marine microbial aggregates

James C. Kosmopoulos*[1,2,3], Danielle E. Campbell*[#3,4,5], Rachel J. Whitaker[3,7,8], Elizabeth G.

Wilbanks[#3,6]

[1] Department of Bacteriology, University of Wisconsin-Madison, Madison, Wisconsin, USA

[2] Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, Wisconsin,

USA

[3] Microbial Diversity 2020, University of Chicago Marine Biological Laboratory, Woods Hole, MA,

USA

[4] Department of Medicine, Division of Infectious Diseases, Washington University School of

Medicine, St. Louis, MO, USA

[5] Edison Family Center for Genome Sciences & Systems Biology, Washington University School of

Medicine, St. Louis, MO, USA

[6] Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara,

Santa Barbara, California, USA

[7] Department of Microbiology, University of Illinois, Urbana, Illinois, USA

[8] Carl R. Woese Institute for Genomic Biology, University of Illinois, Urbana, Illinois, USA

*These authors contributed equally to this work.

[#] Address correspondence to ewilbanks@ucsb.edu or cdanielle@wustl.edu

ABSTRACT

Bacteriophages (phages), viruses that infect bacteria, are the most abundant components of microbial communities and play roles in community dynamics and host evolution. The study of phage-host interactions, however, is made difficult by a paucity of model systems from natural environments and known and cultivable phage-host pairs. Here, we investigate phage-host interactions in the "pink berry" consortia, naturally-occurring, low-diversity, macroscopic aggregates of bacteria found in the Sippewissett Salt Marsh (Falmouth, MA, USA). We leverage metagenomic sequence data and a comparative genomics approach to identify eight compete phage genomes, infer their bacterial hosts from host-encoded clustered regularly interspaced short palindromic repeats (CRISPR), and observe the potential evolutionary consequences of these interactions. Seven of the eight phages identified infect the known pink berry symbionts *Desulfofustis* sp. PB-SRB1, *Thiohalocapsa* sp. PB-PSB1, and *Rhodobacteraceae* sp. A2, and belong to entirely novel viral taxa, except for one genome which represents the second member of the *Knuthellervirus* genus. We further observed increased nucleotide variation over a region of a conserved phage capsid gene that is commonly targeted by host CRISPR systems, suggesting that CRISPRs may drive phage evolution in pink berries. Finally, we identified a predicted phage lysin gene that was horizontally transferred to its bacterial host, potentially via a transposon intermediary, emphasizing the role of phages in bacterial evolution in pink berries. Taken together, our results demonstrate that pink berry consortia contain diverse and variable phages, and provide evidence for phage-host co-evolution via multiple mechanisms in a natural microbial system.

IMPORTANCE

Phages (viruses that infect bacteria) are important components of all microbial systems, where they drive the turnover of organic matter by lysing host cells, facilitate horizontal gene transfer (HGT), and co-evolve with their bacterial hosts. Bacteria resist phage infection, which is often costly or lethal,

through a diversity of mechanisms. One of these mechanisms are CRISPR systems, which encode

arrays of phage-derived sequences from past infections to block subsequent infection with related

phages. Here, we investigate bacteria and phage populations from a simple marine microbial

community known as "pink berries" found in salt marshes of Falmouth, Massachusetts, as a model of

phage-host co-evolution. We identify eight novel phages, and characterize a case of putative CRISPR-

driven phage evolution and an instance of HGT between phage and host, together suggesting that

phages have large evolutionary impacts in a naturally-occuring microbial community.

INTRODUCTION

Phages, viruses that infect bacteria, occur in all microbial ecosystems, often outnumbering

bacteria by 10 to 1, and play pivotal roles in altering community structure (Andersson & Banfield,

2008; Bergh *et al.*, 1989; Breitbart *et al.*, 2018), mediating horizontal gene transfer (HGT) (Breitbart *et*

al., 2018; Hall *et al.*, 2017; Schneider, 2021), and driving bacterial evolution (Campbell *et al.*, 2020;

Koskella & Brockhurst, 2014; Martiny *et al.*, 2014). Though some phage-host interactions can be

beneficial, phage infection canonically ends with the lysis and death of the host to release progeny

phage particles for transmission to new host cells. Thus, there is strong selection for bacteria to evolve

mechanisms to resist infection. Likewise, phages must evolve to overcome those resistances to

survive. This phage-bacterial host coevolution is often described as an "arms race" (Hampton *et al.*,

2020).

Bacteria have evolved a wide range of phage defense systems, such as clustered regularly

interspaced short palindromic repeat (CRISPR) loci, which act as microbial adaptive immune systems.

During a new phage infection, CRISPR systems incorporate short segments of phage-derived

sequence, known as "protospacers," into CRISPR arrays as "spacers" (Barrangou *et al.*, 2007; Jansen

et al., 2002; Jinek *et al.*, 2012). CRISPR systems further encode mechanisms to degrade invading

phage DNA that matches an existing spacer, allowing the host to resist infection. CRISPR arrays thus

serve as a genetic record of phages a host has encountered and can be leveraged to identify phage

hosts from sequence data (Childs *et al.*, 2014; England *et al.*, 2018).

In-depth analysis of CRISPR targeting offers insights into phage-host interactions. CRISPR

systems often target conserved phage sequences, thus conferring protection from groups of related

phages (Barrangou *et al.*, 2007; Deveau *et al.*, 2008; Mojica *et al.*, 2009). Phages can accumulate

mutations within protospacers which allow the phage to escape CRISPR defenses (Andersson &

Banfield, 2008; Deveau *et al.*, 2008; Sun *et al.*, 2013). Together, these patterns in CRISPR spacer and

protospacer nucleotide variation can shed light on phage-host co-evolution.

Here, we investigate phage-bacterial interactions in microbial consortia known as "pink

berries." Pink berries are macroscopic microbial aggregates found in the Sippewissett Salt Marsh of

Falmouth, MA (Seitz *et al*., 1993; Wilbanks *et al.*, 2014). These aggregates are primarily composed of

phototrophic, sulfide-oxidizing and sulfate-reducing bacteria that together form a syntrophic sulfur

cycle (Wilbanks *et al.*, 2014). Though phage sequences have been found in pink berries previously

(Wilbanks *et al.*, 2022), the interactions of phage and their host remain uncharacterized. We show that

pink berries host diverse, novel phages that drive bacterial evolution through HGT and putative

CRISPR-driven arms race dynamics.

RESULTS

Pink berries contain novel phages that are variable between individual aggregates

Pink berries were strategically sampled and pooled during sequencing library preparation to

observe both the breadth of diversity across individual pink berries, and to deeply sample the total

diversity of pink berries. Thus, we sequenced two metagenomes from single pink berries, one with low

sequencing depth (LS06-2018-s01) and one with high sequencing depth (LS06-2018-s02), and one

metagenome from three pink berries homogenized together and sequenced at very high depth (LS06-

2018-s03) (Table 1). Co-assembly of the three pink berry metagenomes yielded 184 contigs totaling

4.35 Mb in length (Table 1). Two phage sequence prediction tools, VIBRANT (Kieft *et al.*, 2020) and

ViralVerify (Antipov *et al.*, 2020), identified nine full-length, circular phage genomes which were the

targets of our downstream analyses (Suppl. Fig. 1). The phages were named according to their hosts

predicted by CRISPR spacer-protospacer matches (described in *Pink berry phages are targeted by*

bacterial CRISPR systems).

Functional annotations were obtained for 236 of 632 putative protein-coding genes predicted

across the nine complete viral genomes (Suppl. Data 1). One virus was found to have large amounts of

homology to eukaryote-infecting Circular Rep-encoding Single Stranded (CRESS) DNA viruses and

may infect a nematode host, which are frequently observed grazing on bacteria in pink berries. This

virus was excluded from further analysis to focus on the primary pink berry bacterial components and

their phages ("Pink berry virus MD00"; Suppl. Data 1). Functional annotations for the remaining 8

phage genomes predicted nucleotide metabolism proteins, head and packaging proteins, integration

and excision proteins, transcriptional regulators, and various lytic proteins (Fig. 1, Suppl. Data 1).

Importantly, all phages of interest here are predicted to exhibit strictly lytic lifecycles, as they lack the

genes for a temperate lifecycle, such as an integrase. There were also predictions for several auxiliary

metabolic genes (Breitbart *et al.*, 2007), which we inferred based on functional predictions outside the

core functions required for the phage lifecycle. One such AMG was a *darB*-like antirestriction gene

encoded on the Thiohalocapsa phage MD04 genome (Suppl. Data 1). Interestingly, *darB* has been

shown to methylate phage DNA to resist host restriction modification (RM) systems (Iida *et al.*, 1987;

Iyer *et al.*, 2017). Prior work found pink berry bacteria employ numerous, diverse RM systems, and

revealed that putative pink berry phages have been shown to contain similar methylation profiles as

their hosts (Wilbanks *et al.,* 2022). The presence of AMGs such as *darB* suggest that pink berry

phages have adapted to increase their fitness in *Thiohalocapsa* hosts, consistent with an armsrace-like

process of coevolution.

Phage taxonomy is based on genome similarity (Turner *et al.*, 2021), and forms the basis of

inferring the diversity of phages within a community. We hypothesized that pink berry phages would

be at least as diverse as their pool of potential bacterial hosts, which span multiple phyla. vConTACT

(Bin Jang, *et al.*, 2019), a genome-wide protein similarity-based approach, was used to infer phage

taxonomy for the eight phage genomes of interest. Although nominal protein similarity was detected

between all phages of interest and the Viral RefSeq protein database during gene annotation (Suppl.

Data 1), only Desulfofustis phage MD02, Thiohalocapsa phage MD04, and Rhodobacteraceae phage

MD07 are connected to a known phage in the vConTACT network (Fig. 2, Suppl. Data 2). Further,

only one phage genome, Desulfofustis phage MD02, had sufficient protein similarity to cluster with a

cultured phage reference genome, Pseudomonas phage PMBT14, which is currently the only species

of the genus *Knuthellervirus* (Suppl. Data 2). None of the phage genomes of interest clustered with

each other. The remaining five genomes of interest lacked sufficient protein similarity for a connection

in the vConTACT network, indicating that these phages represent novel and undescribed diversity.

Relative bacterial abundances are similar across pink berries (Wilbanks *et al.*, 2014), which is

likely due to the constraints of the syntrophic metabolic interactions between constituent bacteria. To

assess the distribution of pink berry-associated bacteria and phages, genome-wide read coverages were

analyzed for individual pink berry metagenomes (Fig. 3). In agreement with previous observations, we

found that the relative abundance of pink berry bacteria is relatively homogenous across samples (Fig.

3A). In contrast, phage presence and abundance are highly variable between different pink berry

communities (Fig. 3B). Only two phages, Rhodobacteraceae phage MD05 and Pink berry phage

MD08, are similarly abundant in each pink berry metagenome, while read mapping to the remaining

six phages suggests they are distributed unevenly between individual aggregates (Fig. 3). Additionally,

to observe whether the phage genomes of interest are present in pink berry metagenomes from

previous years, we mapped reads from a metagenome of 10 pink berries sampled in 2011 (Wilbanks *et*

al., 2014). This revealed near-complete coverage of Desulfofustis phage MD01 and Rhodobacteraceae

phage MD06 (Suppl. Fig. 2), indicating that these phages have persisted over seven years. In contrast,

read mapping to Desulfofustis phage MD02 and Thiohalocapsa phage MD04 genome largely occurs at

highly conserved regions, and is likely the consequence of non-specific cross-mapping (Suppl. Fig. 2).

Likewise, Thiohalocapsa phage MD03, Rhodobacteraceae phages MD05 and MD07, and Pink berry

phage MD00 had <1% or no genome coverage. Since neither this study nor the study from 2011

(Wilbanks *et al.,* 2014) are exhaustive surveys of pink berry diversity, it is difficult to determine the

mechanisms behind the emergence of these six phages. Taken together, these results suggest that

although pink berries have relatively simple and conserved bacterial community structures, their

phages are highly variable over both space and time.

Pink berry phages are targeted by bacterial CRISPR systems

Bacterial CRISPR arrays serve as a record of past phage infection and can be used to infer

hosts for phage genomes (Childs *et al.*, 2014; England *et al.*, 2018). Two independent CRISPR spacer

prediction tools identified a total of 48 unique repeat sequences from four reference genomes for

known pink berry-associated bacteria: *Desulfofustis* sp. PB-SRB1 (GenBank: JAEQMT010000010.1),

Flavobacteriales bacterium (GenBank: DNTB01000031.1), *Rhodobacteraceae* sp. A2 (GenBank:

JAERIM010000001.1), and *Thiohalocapsa* sp. PB-PSB1 (GenBank: CP050890.1) (Wilbanks *et al.*,

2014). Parsing the remaining available pink berry genome, *Oceanicaulis alexandrii* sp. A1 (GenBank:

JAERIO010000015.1), did not yield any CRISPRs. Because CRISPR repeat sequences are conserved

within bacterial species (Lange *et al.*, 2013; Mojica *et al.*, 2000), they can be used to identify adjacent

spacer sequences in unassembled metagenomic short reads (England *et al.*, 2018; Skennerton *et al.*,

2013). Using the CRISPR repeat sequences from reference genomes, NARBL (England *et al.*, 2018)

identified 2,802 unique CRISPR spacer sequences from the set of merged metagenomic reads from

LS06-2018-s01, LS06-2018-s02, and LS06-2018-s03. Of these, 798 spacers were adjacent to repeats

associated with *Desulfofustis* sp. PB-SRB1, 71 were adjacent to *Flavobacteriales* repeats, 349 were

adjacent to *Rhodobacteraceae* sp. A2 repeats, and 1,584 unique spacers were adjacent to repeats

associated with *Thiohalocapsa* sp. PB-PSB1.

To look for evidence of previous phage-bacteria interactions, metagenomic CRISPR spacer

sequences were aligned to the eight complete phage genome assemblies (Fig. 1 & 4A). Of the 2,802

unique spacer sequences extracted from the merged set of metagenomic reads, 163 unique spacers

aligned to seven phage contigs of interest with at least 80% identity over the entire spacer length (Fig.

1 & 4A, Suppl. Data 3). Spacers from three out of the four potential host taxa aligned to phage

genomes, while no spacers from *Flavobacteriales* aligned to any phage genome of interest.

Thiohalocapsa sp. PB-PSB1 was predicted to be the host of Thiohalocapsa phages MD03 and MD04,

Rhodobacteraceae sp. A2 was predicted to be the host of Rhodobacteraceae phages MD05, MD06,

and MD07, and *Desulfofustis* sp. PB-SRB1 was predicted to be the host of Desulfofustis phages MD01

and MD02 (Fig. 4A, Suppl. Data 3). For the two most prevalent and abundant phages identified,

Desulfofustis phage MD02 (Suppl. Fig. 3A) and Thiohalocapsa phage MD04 (Suppl. Fig. 3B), phage

genome and targeting CRISPR spacer coverages were positively correlated. Moreover, although most

host spacers matched to a single virus, two spacers from the *Rhodobacteraceae* host aligned to

Thiohalocapsa phage MD04 and Desulfofustis phage MD02 (Fig. 4A). We do not predict these phages

to have infected the *Rhodobacteraceae* bacterium, since there was only one alignment each to

Thiohalocapsa phage MD04 and Desulfofustis phage MD02 compared to 126 and three alignments

from the two phage genomes to *Thiohalocapsa* and *Desulfofustis* spacers, respectively (Suppl. Data 3).

Additionally, the alignments from the two *Rhodobacteraceae* spacers to Desulfofustis phage MD02

were weaker than the alignments from the *Desulfofustis* spacers (Fig. 4A, Suppl. Data 3). Alignments to multiple host taxa may be due to CRISPR systems targeting motifs that are present in several phage lineages.

The vast majority of CRISPR spacer-protospacer matches occurred only once in the dataset. However, four spacers from *Thiohalocapsa* aligned imperfectly to two distinct protospacers on the genome of Thiohalocapsa phage MD04 (*Thiohalocapsa* sp. PB-PSB1 spacers 09-2, 21-2, 21-11, and 21-164 in Suppl. Data 3). An additional two spacers from *Desulfofustis* sp. PB-SRB1 are reverse complements of each other and target the same protospacer sequence on both Desulfofustis phage MD01 and MD02 (*Desulfofustis* sp. PB-SRB1 spacers 01-103 and 02-11) (Figs. 1, 4A, 4B, Suppl. Data 3). The shared CRISPR targeting of Desulfofustis phage MD01 and MD02 occurred in a conserved gene predicted to encode a phage capsid protein (Fig. 1, Suppl. Data 1) (ORFs DPMD01_45 and DPMD02_11, respectively), and corresponding to genomic regions with high read coverage relative to the remainder of the phage genomes (Fig. 1, Suppl. Fig. 2). A third spacer (01-64) targets a distinct protospacer within this same capsid gene on Desulfofustis phage MD02. A third capsid gene from an incomplete viral contig was found to be homologous to these two variants from Desulfofustis phage MD01 and MD02 and is 92% identical and of similar length. The capsid gene from this incomplete phage contig aligns with the same three spacers targeting Desulfofustis phage MD01 and MD02 (Fig. 4B, Suppl. Data 4). Nucleotide variation at these protospacers inferred by read mapping shows that other variants of these protospacers likely exist in related pink berry phages not assembled here (Fig. 4B). We obtained metagenome-wide allelic variants spanning the entire capsid gene of Desulfofustis phage MD01, Desulfofustis phage MD02, and Incomplete phage contig 1 and observed a near three-fold increase in the number of variants over CRISPR-targeted regions (Suppl. Fig. 4). Taken together, these results suggest that this conserved phage capsid gene is an active site of diversification.

Horizontal gene transfer between pink berry phages and their hosts

Desulfofustis phage MD02 and Thiohalocapsa phage MD04 were found to contain discrete

regions of high read coverage compared to the rest of the genome (Fig. 1, Suppl. Fig. 2). We

hypothesized that these regions may be the result of read mapping from homologous regions of

bacterial chromosomes or other phage genomes.

The high coverage region on Desulfofustis phage MD02 (genome coordinates 170-1760) did

not align to any region of the *Desulfofustis* sp. PB-SRB1 host reference genome (GenBank:

JAEQMT000000000.1) or to any other bacterial genomes in RefSeq. This region also did not

successfully align to any other contigs in the co-assembly, indicating that the coverage at this region is

not the result of conservation of this sequence among other members of the metagenome. Because of

the circularity of these genomes, it is possible that the high read coverage at these regions is attributed

to terminal redundancy from circular permutation (Garneau *et al.*, 2017; Grossi *et al.*, 1983).

The high coverage region of Thiohalocapsa phage MD04 (genome coordinates 21,035-22,077)

encodes a glucosaminidase domain-containing protein (ORF TPMD04_36) predicted to function as the

phage lysin. TPMD04_36 is homologous to two predicted ORFs (NCBI N838_07070 and

N838_07065) in the *Thiohalocapsa* sp. PB-PSB1 genome (GenBank CP050890.1), which are adjacent

to a predicted transposase (Fig. 5). This observation prompted an investigation into a possible

transposon-mediated HGT event between Thiohalocapsa phage MD04 and its host, *Thiohalocapsa* sp.

PB-PSB1.

Alignment of the Thiohalocapsa phage MD04 and *Thiohalocapsa* PB-PB1 genomes revealed

that the phage lysin gene and the host pseudogene are in frame with each other, except the host ORF

N838_07070 contains a single-nucleotide insertion at position 1,668,396 that results in a premature

stop codon (Fig. 5). After closer observation of the region surrounding the pseudogene on the

Thiohalocapsa sp. PB-PSB1 genome, we observed a transposon of the IS4 family with terminal

inverted repeats and numerous direct repeats indicative of past transposase activity (Suppl. Fig. 5). The

Thiohalocapsa phage MD04 genome contains an imperfect copy of a 17-bp direct repeat, differing by

only one nucleotide, at the N-terminal of the TPMD04_36 ORF. Though MD04 was frequently

targeted by host CRISPRs (including at protospacers directly adjacent to the lysin gene), no spacer-

protospacer alignments were observed within the lysin gene, likely the result of selection against

CRISPR self-targeting. Together, this finding suggests a past transposon-mediated HGT event may

have resulted in the transfer of the phage lysin gene from Thiohalocapsa phage MD04, or a related

ancestral phage, to its host.

DISCUSSION

Marine phages are the most numerous biological components of the global ocean,

outnumbering their bacterial hosts by tenfold (Breitbart *et al.*, 2018), and play vital ecosystem roles as

predators that turn over organic matter through bacterial lysis (Heldal & Bratbak, 1991; Maranger &

Bird, 1995; Proctor *et al.*, 1988; Steward *et al.*, 1996) and as agents of HGT impacting bacterial

community structure and function (Anantharaman *et al.*, 2014; Breitbart *et al.*, 2018; Kieft *et al.*, 2021;

Tuttle & Buchan, 2020). Pink berries are marine microbial aggregates with a microscale sulfur cycle,

and have been used as a model system to study cryptic biogeochemical cycling (Wilbanks *et al.*,

2014). Though phages have been identified within the pink berry metagenomes (Wilbanks *et al.*,

2022), the full diversity of phages and their impacts on pink berry communities remain largely

unexplored. Here, we investigated these simple, naturally-occurring microbial communities as a model

for phage-host co-evolution.

We co-assembled three pink berry samples, recovering eight complete phage genomes

spanning a total of 350 Kb and infecting three different bacterial species within the consortia,

Desulfofustis sp. PB-SBR1, Rhodobacteraceae sp. A2, and *Thiohalocapsa* sp. PSB1. We found that

pink berry-associated phages are highly diverse and largely novel, as seven of the eight complete

phage genomes analyzed fail to cluster with any known phage sequences. One pink berry phage,

Desulfofustis phage MD02, is only the second member of the genus *Knuthellervirus*. The other

member of the *Knuthellervirus*, Pseudomonas phage PMBT14, infects *Pseudomonas fluorescens*,

another marine organism, suggesting this phage genus infects diverse hosts.

Although the composition of the pink berry bacterial community is similar across individual

aggregates, we found that phage presence and abundance is highly heterogeneous across samples. Pink

berries are free-living microbial aggregates that exist at the sediment-water interface of intertidal

ponds, with no obvious physical barrier to phage entry into the system. This raises ecological

questions about the mechanisms underlying phage ingress into a pink berry aggregate and their

persistence within, loss, or exclusion from the community.

CRISPRs are a common phage-resistance system employed by bacteria and archaea

(Barrangou *et al.*, 2007; Horvath & Barrangou, 2010; Jansen *et al.*, 2002). We identified an astounding

2,731 unique CRISPR spacer sequences from pink berry-associated *Desulfofustis*, *Rhodobacteraceae*,

and *Thiohalocapsa* hosts, 163 of which (~6%) target a complete phage genome we assembled. This

discrepancy suggests that the true diversity of phages pink berry-associated bacteria encounter is far

greater than what we report here. Seven of the eight phages investigated were targeted by described

pink berry bacterial CRISPR systems, and the eighth, Pink berry phage MD08, may infect other pink

berry-associated taxa that do not encode CRISPR defenses or for which we do not yet have a high-

quality reference genome. Finally, we were able to observe diversification of a CRISPR-targeted

conserved capsid gene, which is inconsistent with diversification across the rest of the phage genomes.

Although we cannot establish a causative relationship between CRISPR targeting and phage variation,

these observations are consistent with a model of CRISPR-driven evolution causing positive selection

in a phage structural protein. We further observed a positive correlation between CRISPR spacer

abundance and target phage abundance for two of the most prevalent and abundant phages identified.

This suggests that these CRISPR spacers are positively selected for within individual pink berry

consortia, and is consistent with observations from diverse microbial systems (Somerville *et al.*, 2022;

Meaden *et al.*, 2021).

Phages and other mobile genetic elements are powerful mediators of HGT (Breitbart *et al.*,

2018; Hall *et al.*, 2017; Schneider, 2021). HGT between bacterial species within the pink berry

consortia has been previously reported (Wilbanks *et al.*, 2022), yet the role of phages in HGT and

bacterial genome evolution in this system remains to be explored. We identified a predicted phage

lysin gene that was horizontally transferred from Thiohalocapsa phage MD04 to its *Thiohalocapsa*

host likely via a transposon intermediary. It is unclear if the lysin gene encoded on the *Thiohalocapsa*

sp. PB-PSB1 genome is functional; a nonsense mutation in this ORF suggests it is a pseudogene, and

was perhaps selected for to avoid deleterious effects of expressing this potentially lethal protein.

Future work should aim to experimentally determine how these phages impact the evolution of both

individual bacterial hosts and entire pink berry aggregates through HGT.

Taken together, our results demonstrate that pink berry communities contain diverse and

variable phage consortia, which are highly targeted by host-encoded CRISPR systems. We leveraged

metagenomic sequence data to better understand phage-host co-evolution occurring through CRISPR

evasion and HGT. Pink berries offer a simple yet relatively unexplored, naturally-occurring model of

phage invasion into and exclusion from microbial communities. The potential roles of phages in pink

berry syntrophy and community-wide metabolic exchanges remain to be explored, but it is now clear

that phages are notable members of these microbial consortia.

METHODS

Sampling

Pink berries, their surrounding sediment, and seawater were collected from pond LS06

(41.57587, -70.63781) in the Little Sippewissett Salt Marsh in Falmouth, MA, on July 17, 2018, using

sterile 50-mL conical tubes. Samples LS06-2018-s01 and LS06-2018-s02 each contained one pink

berry aggregate, and sample LS06-2018-s03 contained three pink berry aggregates. These samples

were transported to the lab and immediately processed for DNA extraction.

DNA isolation and sequencing

Pink berry samples were each mechanically homogenized in 1 mL of TE buffer and

centrifuged at 1,000 xg for 1 min to pellet particulate matter. The supernatant was removed and

subjected to a Wizard Genomic DNA Purification Kit (Promega Catalog No. A1120) according to the

manufacturer's instructions. Purified DNA was fragmented, and sequencing adapters and barcodes

were ligated with the Nextera DNA Flex Library Prep Kit (Illumina Catalog No. 20018705) using

Nextera DNA CD indexes (Illumina Catalog No. 20018708). DNA yield was measured with a Qubit

High Sensitivity dsDNA assay kit (ThermoFisher Catalog No. Q32851), and DNA from LS06-2018-

s01, LS06-2018-s02, and LS06-2018-s03 were pooled at a ratio of 1:1:10. After pooling, DNA was

purified with Ampure XP beads (Beckman Coulter Catalog No. A63881) according to the

manufacturer's instructions. DNA sequencing was performed on an Illumina HiSeq 2500 using a

2x250nt protocol at the University of Illinois at Urbana-Champaign Roy J. Carver Biotechnology

Center.

Metagenome co-assembly & read mapping

For each sample, reads were quality checked with FASTQC v0.11.9 (Andrews, 2010), trimmed

using Trimmomatic v0.39 (Bolger *et al.*, 2014), and adapter sequences were removed. Trimmomatic

filtered reads using a sliding window of 4 base pairs, a minimum average base quality score of 15, a

minimum quality score for retention on the leading and trailing ends of 2, and a minimum read length

of 100 bases. The resulting trimmed and filtered reads were merged into a single set of reads and co-

assembled using Metaviral SPAdes v3.15.2 (Antipov *et al.*, 2020) with default parameters to maximize

the recovery of complete, circular phage genomes. To estimate phage and bacterial abundances,

Bowtie2 v2.4.5 (Langmead & Salzberg, 2012) was used to map reads from the set of merged reads or

from each pink berry sample to assembled phage contigs and to representative host genomes from

NCBI BioProject PRJNA684324: *Desulfofustis* sp. PB-SRB1 (GenBank: JAEQMT010000010.1),

Rhodobacteraceae sp. A2 (GenBank: JAERIM010000001.1), and *Thiohalocapsa* sp. PB-PSB1

(GenBank: CP050890.1) (Wilbanks *et al.*, 2022). Read mapping statistics were obtained from Bowtie2

alignments using samtools v1.15.1 (Danecek *et al.*, 2021).

Phage sequence identification, binning, and annotation

ViralVerify v1.1 (Antipov *et al.*, 2020) was used with default settings to categorize the

metagenome contigs as putatively bacterial or viral. ViralComplete v1.1 (Antipov *et al.*, 2020) was

used with default settings to identify viral contigs that represent complete phage genomes. To verify

these predictions, VIBRANT v1.2.1 (Kieft *et al.*, 2020) was used with default settings on the same

metagenome contigs. All resulting putative viral contigs that were estimated to be complete viral

genomes by both prediction tools were targeted for downstream analyses and annotation. vConTACT

v2.0 (Bin Jang *et al.*, 2019) was used with the RefSeq v211 Viral database (Brister *et al.*, 2015) to

cluster the viral contigs of interest with existing phage genomes and to approximate phage taxonomy.

The viral contigs of interest were passed through Pharokka v1.0.1

(github.com/gbouras13/pharokka), using PHANOTATE v1.5.0 (McNair *et al.*, 2019) to predict genes

and PHROGs v3 (Terzian *et al.*, 2021) to provide initial protein annotations. Protein functions were

also predicted using Phyre2 (Kelley *et al.*, 2015), BLASTp v2.11.0 (Altschul *et al.*, 1990) with the

NCBI RefSeq v211 virus amino acid database and the non-redundant amino acid database (Brister *et*

al., 2015; O'Leary *et al.*, 2016), and HMMER v3.2.1 (Eddy, 2011) with Pfam-a v35.0 (Mistry *et al.*,

2021) and TIGRFAMs v15.0 (Li *et al.*, 2021). The resulting predictions from each method were

manually reviewed for each protein, and a consensus annotation was inferred (Suppl. Data 1). Clinker

v0.0.23 (Gilchrist & Chooi, 2021) was used to identify conserved genes among phage genomes, which

were then aligned with tBLASTx v2.11.0 (Camacho *et al.*, 2009) against all the contigs in the

metagenome co-assembly. Metagenomic reads from pink berries sampled in 2011 (SRA:

SRR13297012) were mapped to the viral contigs of interest with Bowtie2 using the same methods

described above.

CRISPR spacer-protospacer analysis and host prediction

CRISPRclassify v1.1.0 (Nethery *et al.*, 2021) and MinCEd v0.4.2 (Bland *et al.*, 2007) were

used to identify CRISPR repeat sequences from representative genomes of pink berry taxa from NCBI

BioProject PRJNA684324: *Desulfofustis* sp. PB-SRB1 (GenBank: JAEQMT010000010.1),

Oceanicaulis alexandrii (GenBank: JAERIO000000000.1), *Rhodobacteraceae* sp. A2 (GenBank:

JAERIM010000001.1), *Thiohalocapsa* sp. PB-PSB1 (GenBank: CP050890.1) (Sayers *et al.*, 2020;

Wilbanks *et al.*, 2022), and *Flavobacteriales* bacterium (GenBank: DNTB01000031.1). The identified

repeats from each tool were combined and dereplicated to obtain a list of repeats found in the genomes

of pink berry taxa. Since CRISPR arrays are often misassembled with short-read data, putative spacer

sequences from the trimmed and filtered reads of each LS06-2018 metagenome were identified with

NARBL (England *et al.*, 2018) using the dereplicated set of repeats identified from the reference

genomes, an approximate repeat size of 36, and a minimum coverage of supporting neighbor spacers

of 2. Since repeat sequences are highly conserved between bacterial species (Kunin *et al.*, 2007), any

spacer identified by NARBL was inferred to belong to the same species as the reference genome from

which its associated repeat came. The resulting spacers were aligned to the viral contigs of interest

using Spacerblast v0.7.7 (Collins & Whitaker, 2022). Viral contigs that aligned to spacers with at least

80% identity over the full length of the spacer were considered to have a host match with the

bacterium whose genome contained the spacer. The merged set of filtered and trimmed reads were

aligned to spacers identified by NARBL with Bowtie2 and read coverage statistics were obtained with

samtools as described above.

Metagenome reads were mapped to the regions containing conserved phage genes identified

with Clinker or tBLASTx, above, with spacer-protospacer alignments with Bowtie2. Mapped reads

were converted to multiple sequence alignments using SAM4WebLogo in JVarkit v2021.10.13

(Lindenbaum, 2015) and sequence logos were visualized with WebLogo (Crooks *et al.*, 2004). Allele

variants for conserved phage genes were called by using Snippy v3.2 (github.com/tseemann/snippy)

with default settings on metagenome reads. Variants that resulted before the "--mincov" and "--

minfrac" filters were applied were used downstream to maximize the number of possible variants

recovered. Variant statistics were obtained and visualized using vcfR v1.13.0 (Knaus & Grünwald,

2017) with default settings, except 100-bp size windows were used instead of 1000-bp.

Horizontal gene transfer analysis

Uneven sequence coverage patterns on phage genomes are sometimes attributed to HGT

between the phage and its host genome, especially if the region aligns to the host genome and/or

contains genes that facilitate HGT (Kleiner *et al.*, 2020). Viral genomes of interest and their coverages

from the merged set of metagenome reads were visualized in IGV v2.11.4 (Thorvaldsdóttir *et al.*,

2013). For any discrete regions of the viral genomes of interest that had much higher read coverage

(>3x) than the surrounding region, those regions were aligned to their predicted host genomes, if

available, using BLASTn v2.11.0 (Camacho *et al.*, 2009). Predicted regions of HGT between phages

and their hosts were aligned with Mauve v2015-02-13 (Darling *et al.*, 2004). Conserved repeats

flanking a putative transposon were initially identified by reciprocal BLASTn using "blastn-short".

The transposon region with its inverted repeats in the *Thiohalocapsa* sp. PB-PSB1 genome was

identified with ISEScan v1.7.2.3 (Xie & Tang, 2017), and other repeats were identified and annotated

manually in Geneious Prime v2022.1.1 (www.geneious.com/prime).

Data availability

ACKNOWLEDGMENTS

This research was conducted as a part of the 2020 Microbial Diversity course at the Marine

Biological Laboratory (MBL), which was funded by NSF Award ID 1822263, DOE Award ID DE-

SC0016127, and the Simons Foundation. We gratefully acknowledge the insights and support

provided by all participants and instructors, especially Whitney England, Laura Suttenfield, and

George O'Toole. D.E.C. was supported by NIH T32 DK077653-29 and Crohn's & Colitis Foundation

Research Fellowship Award #935619. E.G.W. gratefully acknowledges support from the Whitman

Fellowship at the MBL. This research was sponsored by the U.S. Army Research Office and

accomplished under cooperative agreement W911NF-19-2-0026 for the Institute for Collaborative

Biotechnologies.

Conceptualization, J.C.K., D.E.C., E.G.W., and R.J.W.; Data Curation, J.C.K., D.E.C. and

R.J.W.; Analysis, J.C.K. and D.E.C.; Funding Acquisition, R.J.W.; Investigation, J.C.K. and D.E.C.;

Methodology, J.C.K., D.E.C., E.G.W., and R.J.W.; Project Administration, J.C.K., D.E.C., E.G.W.,

and R.J.W.; Resources, R.J.W.; Software, J.C.K. and D.E.C.; Supervision, D.E.C., E.G.W., R.J.W.;

Validation, J.C.K. and D.E.C.; Visualization, J.C.K. and D.E.C.; Writing—Original Draft, J.C.K. and

D.E.C.; Writing—Review and Editing, J.C.K., D.E.C., E.G.W., and R.J.W.

REFERENCES

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment

search tool. J*ournal of molecular biology, 215*(3), 403–410. https://doi.org/10.1016/S0022-

2836(05)80360-2

Anantharaman, K., Duhaime, M. B., Breier, J. A., Wendt, K. A., Toner, B. M., & Dick, G. J. (2014).

Sulfur oxidation genes in diverse deep-sea viruses. *Science*, *344*(6185), 757-760.

Andersson, A. F., & Banfield, J. F. (2008). Virus population dynamics and acquired virus resistance in

natural microbial communities. *Science*, 320(5879), 1047-1050.

Andrews, S. (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data [Online].

Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Antipov, D., Raiko, M., Lapidus, A., & Pevzner, P. A. (2020). Metaviral SPAdes: assembly of viruses

from metagenomic data. *Bioinformatics (Oxford, England)*, *36*(14), 4126–4129.

https://doi.org/10.1093/bioinformatics/btaa490.

Barrangou, R., Fremaux, C., Deveau, H., Richards, M., Boyaval, P., Moineau, S., ... & Horvath, P.

(2007). CRISPR provides acquired resistance against viruses in prokaryotes. *Science*,

315(5819), 1709-1712.

Bergh, Ø., Børsheim, K. Y., Bratbak, G., & Heldal, M. (1989). High abundance of viruses found in

aquatic environments. *Nature, 340*(6233), 467-468.

Bin Jang, H., Bolduc, B., Zablocki, O., Kuhn, J. H., Roux, S., Adriaenssens, E. M., Brister, J. R.,

Kropinski, A. M., Krupovic, M., Lavigne, R., Turner, D., & Sullivan, M. B. (2019).

Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing

networks. *Nature biotechnology, 37*(6), 632–639. https://doi.org/10.1038/s41587-019-0100-8

Bland, C., Ramsey, T. L., Sabree, F., Lowe, M., Brown, K., Kyrpides, N. C., & Hugenholtz, P. (2007).

CRISPR recognition Tool (CRT): A tool for automatic detection of Clustered Regularly

Interspaced Palindromic repeats. *BMC Bioinformatics, 8*(1). https://doi.org/10.1186/1471-

2105-8-209

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer FOR Illumina

sequence data. *Bioinformatics, 30*(15), 2114–2120.

https://doi.org/10.1093/bioinformatics/btu170

Breitbart, M., Bonnain, C., Malki, K., & Sawaya, N. A. (2018). Phage puppet masters of the marine

microbial realm. *Nature Microbiology* 3, 754–766. https://doi.org/10.1038/s41564-018-0166-y

Breitbart, M., Thompson, L. R., Suttle, C. A., & Sullivan, M. B. (2007). Exploring the vast diversity of

marine viruses. *Oceanography (Wash DC)* 20, 135–139.

Brister, J. R., Ako-Adjei, D., Bao, Y., & Blinkova, O. (2015). NCBI viral genomes resource. *Nucleic*

Acids Research, 43. https://doi.org/10.1093/nar/gku1207

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics, 10*(421), https://doi.org/10.1186/1471-2105-10-421

Campbell, D. E., Ly, L. K., Ridlon, J. M., Hsiao, A., Whitaker, R. J., & Degnan, P. H. (2020). Infection with Bacteroides phage BV01 alters the host transcriptome and bile acid metabolism in a common human gut microbe. *Cell reports, 32*(11), 108142.

Childs, L. M., England, W. E., Young, M. J., Weitz, J. S., & Whitaker, R. J. (2014). CRISPR-Induced Distributed Immunity in Microbial Populations. *PLoS ONE, 9*(7). https://doi.org/10.1371/journal.pone.0101710

Collins, A. J., & Whitaker, R. J. (2022). CRISPR Comparison Toolkit (CCTK): Rapid Identification, Visualization, and Analysis of CRISPR Array Diversity. *BioRxiv* 2022.07.31.502198. Advance online publication. https://doi.org/10.1101/2022.07.31.502198

Crooks, G. E., Hon, G., Chandonia, J. M., & Brenner, S. E. (2004). WebLogo: a sequence logo generator. Genome research, 14(6), 1188–1190. https://doi.org/10.1101/gr.849004

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of samtools and BCFtools. GigaScience, 10(2). https://doi.org/10.1093/gigascience/giab008

Darling, A. C., Mau, B., Blattner, F. R., & Perna, N. T. (2004). Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome research, 14*(7), 1394–1403. https://doi.org/10.1101/gr.2289704

Deveau, H., Barrangou, R., Garneau, J. E., Labonté, J., Fremaux, C., Boyaval, P., Romero, D. A., Horvath, P., & Moineau, S. (2008). Phage response to CRISPR-encoded resistance in Streptococcus thermophilus. *Journal of bacteriology*, 190(4), 1390-1400.

Eddy, S. R. (2011). Accelerated profile HMM searches. *PLoS Computational Biology*, *7*(10).

https://doi.org/10.1371/journal.pcbi.1002195

England, W. E., Kim, T., & Whitaker, R. J. (2018). Metapopulation Structure of CRISPR-Cas

Immunity in Pseudomonas aeruginosa and Its Viruses. *MSystems*, *3*(5).

https://doi.org/10.1128/msystems.00075-18

Garneau, J. R., Depardieu, F., Fortier, L. C., Bikard, D., & Monot, M. (2017). PhageTerm: a tool for

fast and accurate determination of phage termini and packaging mechanism using next-

generation sequencing data. *Scientific reports*, *7*(1), 8292. https://doi.org/10.1038/s41598-017-

07910-5

Gilchrist, C. L., & Chooi, Y. H. (2021). Clinker & clustermap. js: Automatic generation of gene

cluster comparison figures. *Bioinformatics, 37*(16), 2473-2475.

Grossi, G. F., Macchiato, M. F., & Gialanella, G. (1983). Circular permutation analysis of phage T4

DNA by electron microscopy. *Zeitschrift fur Naturforschung. Section C, Biosciences*, *38*(3-4),

294–296. https://doi.org/10.1515/znc-1983-3-422

Hall, J. P., Brockhurst, M. A., & Harrison, E. (2017). Sampling the mobile gene pool: Innovation via

horizontal gene transfer in bacteria. Philosophical Transactions of the Royal Society B:

Biological Sciences, 372(1735), 20160424. https://doi.org/10.1098/rstb.2016.0424

Hampton, H. G., Watson, B. N., & Fineran, P. C. (2020). The arms race between bacteria and their

phage foes. *Nature*, *577*(7790), 327–336. https://doi.org/10.1038/s41586-019-1894-8

Heldal, M., & Bratbak, G. (1991). Production and decay of viruses in aquatic environments. *Mar.

Ecol. Prog. Ser*, *72*(3), 205-212.

Horvath, P., & Barrangou, R. (2010). CRISPR/Cas, the immune system of bacteria and archaea.

Science, 327(5962), 167-170.

Iida, S., Streiff, M. B., Bickle, T. A., & Arber, W. (1987). Two DNA antirestriction systems of bacteriophage P1, darA, and darB: characterization of darA□ phages. *Virology*, *157*(1), 156-166.

Iyer, L. M., Burroughs, A. M., Anand, S., de Souza, R. F., & Aravind, L. (2017). Polyvalent proteins, a pervasive theme in the intergenomic biological conflicts of bacteriophages and conjugative elements. *Journal of bacteriology*, *199*(15), e00245-17.

Jansen, R., Embden, J. D. V., Gaastra, W., & Schouls, L. M. (2002). Identification of genes that are associated with DNA repeats in prokaryotes. *Molecular microbiology*, *43*(6), 1565-1575.

Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J. A., & Charpentier, E. (2012). A programmable dual-RNA–guided DNA endonuclease in adaptive bacterial immunity. *science*, *337*(6096), 816-821.

Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N., & Sternberg, M. J. E. (2015). The Phyre2 web portal for protein modeling, prediction and analysis. Nature Protocols 10, 845-858.

Kieft, K., Zhou, Z., & Anantharaman, K. (2020). VIBRANT: automated recovery, annotation and curation of microbial viruses, and evaluation of viral community function from genomic sequences. *Microbiome*, *8*(1), 1-23.

Kieft, K., Zhou, Z., Anderson, R. E., Buchan, A., Campbell, B. J., Hallam, S. J., Hess, M., Sullivan, M. B., Walsh, D. A., Roux, S., & Anantharaman, K. (2021). Ecology of inorganic sulfur auxiliary metabolism in widespread bacteriophages. *Nature Communications, 12*(1). https://doi.org/10.1038/s41467-021-23698-5

Kleiner, M., Bushnell, B., Sanderson, K. E., Hooper, L. V., & Duerkop, B. A. (2020). Transductomics: sequencing-based detection and analysis of transduced DNA in pure cultures and microbial communities. *Microbiome*, *8*(1), 1-17.

Koskella, B., & Brockhurst, M. A. (2014). Bacteria–phage coevolution as a driver of ecological and

evolutionary processes in microbial communities. *FEMS Microbiology Reviews*, *38*(5), 916–

931. https://doi.org/10.1111/1574-6976.12072

Kunin, V., Sorek, R., & Hugenholtz, P. (2007). Evolutionary conservation of sequence and secondary

structures in CRISPR repeats. *Genome biology*, *8*(4), 1-7.

Lange, S. J., Alkhnbashi, O. S., Rose, D., Will, S., & Backofen, R. (2013). CRISPRmap: an automated

classification of repeat conservation in prokaryotic adaptive immune systems. *Nucleic acids*

research, *41*(17), 8034-8044.

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods,*

9(4), 357–359. https://doi.org/10.1038/nmeth.1923

Li, W., O'Neill, K. R., Haft, D. H., DiCuccio, M., Chetvernin, V., Badretdin, A., Coulouris, G.,

Chitsaz, F., Derbyshire, M. K., Durkin, A. S., Gonzales, N. R., Gwadz, M., Lanczycki, C. J.,

Song, J. S., Thanki, N., Wang, J., Yamashita, R. A., Yang, M., Zheng, C., … Thibaud-Nissen,

F. (2021). RefSeq: expanding the Prokaryotic Genome Annotation Pipeline reach with protein

family model curation. *Nucleic acids research*, *49*(D1), D1020–D1028.

https://doi.org/10.1093/nar/gkaa1105.

Lindenbaum, P. (2015). JVarkit: java-based utilities for Bioinformatics. *figshare*. Journal contribution.

https://doi.org/10.6084/m9.figshare.1425030.v1

Maranger, R., & Bird, D. F. (1995). Viral abundance in aquatic systems: a comparison between marine

and fresh waters. *Marine Ecology Progress Series*, *121*, 217-226.

Martiny, J. B., Riemann, L., Marston, M. F., & Middelboe, M. (2014). Antagonistic coevolution of

marine planktonic viruses and their hosts. *Annual review of marine science*, *6*, 393-414.

McNair, K., Zhou, C., Dinsdale, E. A., Souza, B., & Edwards, R. A. (2019). PHANOTATE: A novel

approach to gene identification in phage genomes. *Bioinformatics*, *35*(22), 4537–4542.

https://doi.org/10.1093/bioinformatics/btz265

Meaden, S., Biswas, A., Arkhipova, K., Morales, S., Dutilh, B., Westra, E., Fineran, P. (2021). High

viral abundance and low diversity are associated with increased CRISPR-Cas prevalence

across   microbial   ecosystems.   *Current   bioloigy,   32*(1),   P220-227.E5.

https://doi.org/10.1016/j.cub.2021.10.038

Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E., Tosatto, S.,

Paladin, L., Raj, S., Richardson, L. J., Finn, R. D., & Bateman, A. (2021). Pfam: The protein

families   database   in   2021.   *Nucleic   acids   research*,   *49*(D1),   D412–D419.

https://doi.org/10.1093/nar/gkaa913

Mojica, F. J., Díez-Villaseñor, C., Soria, E., & Juez, G. (2000). Biological significance of a family of

regularly spaced repeats in the genomes of Archaea, Bacteria and mitochondria. *Molecular

microbiology*, *36*(1), 244-246.

Mojica, F. J., Díez-Villaseñor, C., García-Martínez, J., & Almendros, C. (2009). Short motif

sequences determine the targets of the prokaryotic CRISPR defence system. *Microbiology*,

155(3), 733-740.

Knaus, B. J., & Grünwald, N. J. (2017). vcfr: a package to manipulate and visualize variant call format

data in R. *Molecular ecology resources*, *17*(1), 44-53.

Nayfach, S., Camargo, A. P., Schulz, F., Eloe-Fadrosh, E., Roux, S., & Kyrpides, N. C. (2020).

CheckV assesses the quality and completeness of metagenome-assembled viral genomes.

Nature Biotechnology, 39(5), 578–585. https://doi.org/10.1038/s41587-020-00774-7

590 Nethery, M. A., Korvink, M., Makarova, K. S., Wolf, Y. I., Koonin, E. V., & Barrangou, R. (2021).

591   CRISPRclassify: Repeat-Based Classification of CRISPR Loci. *The CRISPR journal*, *4*(4),

592   558–574. https://doi.org/10.1089/crispr.2021.0021

593 O'Leary, N. A., Wright, M. W., Brister, J. R., Ciufo, S., Haddad, D., McVeigh, R., Rajput, B.,

594   Robbertse, B., Smith-White, B., Ako-Adjei, D., Astashyn, A., Badretdin, A., Bao, Y.,

595   Blinkova, O., Brover, V., Chetvernin, V., Choi, J., Cox, E., Ermolaeva, O., … Pruitt, K. D.

596   (2016). Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion,

597   and functional annotation. *Nucleic acids research*, *44*(D1), D733–D745.

598   https://doi.org/10.1093/nar/gkv1189

599 Proctor, L. M., Fuhrman, J. A., & Ledbetter, M. C. (1988). Marine bacteriophages and bacterial

600   mortality. *Eos*, *69*, 1111-1112.

601 Sayers, E. W., Cavanaugh, M., Clark, K., Pruitt, K. D., Sherry, S. T., Yankie, L., & Karsch-Mizrachi,

602   I. (2020). GenBank. *Nucleic Acids Research*.

603 Schneider, C. L. (2021). Bacteriophage-mediated horizontal gene transfer: transduction.

604   *Bacteriophages: Biology, Technology, Therapy*, 151-192.

605 Schneider, T. D., & Stephens, R. M. (1990). Sequence logos: a new way to display consensus

606   sequences. *Nucleic acids research*, *18*(20), 6097-6100. https://doi.org/10.1093/nar/18.20.6097

607 Seitz, A. P., Nielsen, T. H., & Overmann, J. (1993). Physiology of purple sulfur bacteria forming

608   macroscopic aggregates in Great Sippewissett Salt Marsh, Massachusetts. *FEMS microbiology*

609   *ecology*, *12*(4), 225-235.

610 Skennerton, C. T., Imelfort, M., & Tyson, G. W. (2013). Crass: identification and reconstruction of

611   CRISPR from unassembled metagenomic data. *Nucleic acids research, 41*(10), e105-e105.

Somerville, V., Schowing, T., Chabas, H., Schmidt, R., Ah, U., Bruggmann, R., Engel, P. (2022).

Extensive diversity and rapid turnover of phage defense repertoires in cheese-associated

bacterial communities. *Microbiome, 10*(1), 137. https://doi.org/10.1186/s40168-022-01328-6

Steward, G. F., Smith, D. C., & Azam, F. (1996). Abundance and production of bacteria and viruses in

the Bering and Chukchi Seas. *Marine Ecology Progress Series*, *131*, 287-300.

Su, G., Morris, J. H., Demchak, B., & Bader, G. D. (2014). Biological network exploration with

Cytoscape 3. *Current protocols in bioinformatics*, *47*(1), 8-13.

Sun, C. L., Barrangou, R., Thomas, B. C., Horvath, P., Fremaux, C., & Banfield, J. F. (2013). Phage

mutations in response to CRISPR diversification in a bacterial population. *Environmental*

microbiology, 15(2), 463-470.

Terzian, P., Olo Ndela, E., Galiez, C., Lossouarn, J., Pérez Bucio, R. E., Mom, R., Toussaint, A., Petit,

M. A., & Enault, F. (2021). PHROG: families of prokaryotic virus proteins clustered using

remote homology. *NAR Genomics and Bioinformatics*, *3*(3).

Thorvaldsdóttir, H., Robinson, J. T., & Mesirov, J. P. (2013). Integrative Genomics Viewer (IGV):

high-performance genomics data visualization and exploration. *Briefings in bioinformatics*,

14(2), 178-192.

Turner, D., Kropinski, A. M., & Adriaenssens, E. M. (2021). A Roadmap for Genome-Based Phage

Taxonomy. *Viruses*, *13*(3), 506. https://doi.org/10.3390/v13030506

Tuttle, M. J., & Buchan, A. (2020). Lysogeny in the oceans: lessons from cultivated model systems

and a reanalysis of its prevalence. *Environmental microbiology*, *22*(12), 4919-4933.

Wilbanks, E. G., Doré, H., Ashby, M. H., Heiner, C., Roberts, R. J., & Eisen, J. A. (2022).

Metagenomic methylation patterns resolve bacterial genomes of unusual size and structural

complexity. *The ISME Journal*, 1-11.

635 Wilbanks, E. G., Jaekel, U., Salman, V., Humphrey, P. T., Eisen, J. A., Facciotti, M. T., Buckley, D.

636   H., Zinder, S. H., Druschel, G. K., Fike, D. A., & Orphan, V. J. (2014). Microscale sulfur

637   cycling in the phototrophic pink berry consortia of the Sippewissett Salt Marsh. *Environmental*

638   *Microbiology, 16*(11), 3398–3415. https://doi.org/10.1111/1462-2920.12388

639 Xie, Z., & Tang, H. (2017). ISEScan: automated identification of insertion sequence elements in

640   prokaryotic genomes. *Bioinformatics*, *33*(21), 3340-3347.

641   **Table 1. Summary of pink berry metagenome sequencing and co-assembly.**

	LS06-2018-s01	LS06-2018-s02	LS06-2018-s03
Total read length (Gb)	0.41	1.07	7.19
Quality trimmed total read length (Gb)	0.35	0.91	6.94
Perc. of total read length after quality trimming	87%	85%	97%
	Co-assembly		
Contigs	184		
N50 (bp)	50,490		
L50	23		
Avg. read coverage (±SD)	12.89 ± 39.63		

Read sample
— LS06-2018-s01
— LS06-2018-s02
— LS06-2018-s03

Protein annotation category
- ▶ Lysis
- ▶ Auxiliary metabolic gene
- ▶ Head-tail connector
- ▶ Head and packaging
- ▶ DNA, RNA, and nucleotide metabolism
- ▷ Integration and excision
- ▶ Tail protein
- ▶ Transcription regulation
- ▷ Other/hypothetical protein

Spacer target host
- ▲ *Desulfofustis*
- ▲ *Thiohalocapsa*
- ▲ *Rhodobacteraceae*

Region highlights
- HGT event
- Conserved gene

Figure 1. Complete phage genomes vary in abundance across samples and are targeted by

bacterial CRISPR spacers. Normalized read coverage by position for each sample are given.

Coverage values were normalized to the total number of trimmed and filtered reads for each sample.

Horizontal arrows indicate ORFs predicted by PHANOTATE (McNair *et al.*, 2019), and their colors

correspond to predicted functional categories. Triangles indicate genome positions of protospacers,

colored by host taxonomy of the corresponding spacer. Regions highlighted with a blue background

indicate a conserved gene between phage genomes inferred by Clinker (Gilchrist & Chooi, 2021).

Region highlighted with a red background was found to be an HGT event between the phage and host.

Figure 2. Pink berry consortia host largely novel phages. Each node represents a phage genome and edges are genome relatedness inferred by vConTACT (Bin Jang *et al.*, 2019). Reference genomes are from the RefSeq Prokaryotic virus database v211 (Brister *et al.*, 2015). The group circled with a solid line shows the phage genus of interest, *Knuthellervirus*. Groups circled in dashed lines show related phages infecting the same host. The network was visualized in Cytoscape v3.9.0 (Su *et al.*, 2014). Only phages which are first-neighbors to pink berry phage genomes are shown.

Figure 3. Phage presence and abundance are highly variable between pink berry communities. Average read coverages for host (A) and phage (B) genomes were converted to reads per kilobase million (RPKM) using the total number of filtered and trimmed reads per sample.

Figure 4. CRISPR spacer-phage genome alignments reveal hosts and a conserved protospacer.
(A) Spacerblast (Collins & Whitaker, 2022) alignment results with at least 80% nucleotide identity over the entire spacer length were visualized in Cytoscape v3.9.0 (Su *et al.*, 2014). Circular nodes represent unique spacers from bacterial contigs and are colored by taxonomy. Triangular nodes are phage contigs. Solid edges represent percent nucleotide identity over the entire spacer length. The dashed edge shows the connection between Desulfofustis phage MD01 and *Desulfofustis* sp. PB-SRB1

repeat 01 spacer 64, which is below 80% identity and included in part (C). The nucleotide sequences

of phage protospacers within conserved capsid genes that were identified on phage contigs were

aligned to (B) *Desulfofustis* sp. PB-SRB1 repeat 01 spacer 103 and (C) *Desulfofustis* sp. PB-SRB1

repeat 01 spacer 64. Spacer 02-11 is the reverse complement of 01-103 and is not shown. The resulting

metagenome-wide variation in protospacer sequences from mapping reads to protospacers are shown

as sequence logos (Crooks *et al.*, 2004; Schneider & Stephens, 1990).

Figure 5. Region of homology between Thiohalocapsa phage MD04 and its host. Locally colinear

blocks aligned with Mauve (Darling *et al.*, 2004) are shown in red, with traces inside representing

nucleotide similarity. Genome tracks show genome coordinates and ORFs. Conserved 17-bp direct

repeat sequences (DR) are shown in blue. A single-nucleotide insertion (green) in the *Thiohalocapsa*

ORF N838_07070 results in a premature stop codon and a pseudogene annotation.