Phantom touch illusion

Unexpected phenomenological effects of tactile gating in the absence of tactile stimulation

Artur Pilacinski[1,2,*], Marita Metzler[1], Christian Klaes[1]

[1] Chair for Neurotechnology, Ruhr-University Bochum, Germany.

[2] CINEICC, University of Coimbra, Portugal

* Corresponding author: art.pilacinski@gmail.com

Abstract

We report the presence of a tingling sensation perceived during self touch without physical stimulation. We used immersive virtual reality scenarios in which subjects touched their body using a virtual object. This touch resulted in a tingling sensation corresponding to the location touched on the virtual body. We called it "phantom touch illusion" (PTI). Interestingly the illusion was also present when subjects touched invisible (inferred) parts of their limb. We reason that this PTI results from tactile gating process during self-touch. The presence of PTI when touching invisible body arms indicates that tactile gating is not exclusively based on vision, but rather on multi-sensory, top-down input involving body schema. This finding shows that representations of own body are defined top-down, beyond the available sensory information.

Introduction

You can't tickle yourself. If you try sliding a finger along your forearm, the tickle sensation will be much weaker than if there was an insect crawling down your skin. This is because the nervous

system cancels out the predicted sensory input caused by your own movements (Blakemore et al., 2000). This mechanism is called tactile gating. Tactile gating has been described on neuronal

level as downregulation of neuronal firing in response to tactile receptors activation (Angel and Malenka, 1982; Blakemore et al., 1998; Bays et al., 2006). It affects several levels of processing

of tactile information, including spinal cord, motor, premotor and somatosensory cortex (Seki and Fetz, 2012). The involvement of these different levels of processing might reflect the complex

nature of tactile gating, linked to multisensory representation of touch based on sensory information, body schema, predictive encoding or active touch (Fuehrer et al., 2022; Chapman

and Beauchamp, 2006).

While some authors suggest that tactile gating is sensory-driven (Williams and Chapman, 2002),

recent evidence suggests that it indeed results from predictions encoding self touch. Fuehrer et al. (2022) showed that the size of tactile gating depends on whether the touch pattern can be

predicted or not. While according to the authors this effect suggests that tactile gating relies on prediction, they also show that it occurs for non-predicted stimuli, suggesting that non-specific

attenuation may also be involved (Fuehrer et al., 2022). Regardless of specificity, the presence of predictive processing in tactile gating makes one ask what if these predictions are confronted

with lack of actual tactile input to attenuate. But what happens with tactile gating if there is no afferent tactile signal? Here we tested this using an immersive virtual reality (VR) scenario in

which we asked subjects to self-touch using a virtual stick. We were inspired by the phenomenon of "phantom" touch sensation that has been anecdotally reported by people using virtual reality

environments (Alexdottir et al. 2022). This sensation is only broadly described as a vague feeling of touch when - for example - the user gets in contact with another user's avatar in a VR

environment. We predict that, if this sensation results from predicted touch, it should likewise reflect the phenomenological properties of the neural activity meant to attenuate predicted

touch on one's own skin such as moving together with the visual object.

Methods

62 *Participants*

Twenty-one participants (nine males) took part in the study. The participants were 21-42 years

64 old. The participants had none to little prior experience with virtual reality, no expert knowledge

in touch perception, and were completely naive to the experimental hypothesis. The experiment

66 was conducted by three experimenters according to a standard protocol. All procedures were

approved by the local ethics committee of the Ruhr-University Bochum and were performed in

68 accordance with the declaration of Helsinki.

70 *Experimental setup and virtual reality scenes*

We used Unity (Unity Technology, San Francisco, USA) version 2021.3.8f1 and Oculus Quest 2

72 (Meta, California, USA) head-mounted display (HMD) in hand tracking mode. The scenes took

place in a simulated virtual environment where several realistically scaled daily objects were

74 placed on a table in front of the participant (Figure 1 B). Among them there were two sticks,

approximately 30 cm long and 2 cm wide. The subjects could only see their virtual hands but not

76 their forearms or other body parts.

78 Figure 1. A) Schematic depiction of stimulation with the virtual stick in prone and supine positions. B) A

screenshot of the VR stimulation scene in prone and supine positions. C) Stimulation sites: fingertips,

80 fingers (phalanges), palm and forearm. The forearms were invisible to subjects (see panel B).

84   *Procedure*

At the beginning, the HMD was put on the subjects and we ensured they could see the scene and

86   their virtual hands clearly. Then, subjects familiarized themselves with the scene such as freely

moving and touching the objects on the table. After they appeared acclimatized to the VR

88   environment, the experimental phase began. We asked the subjects to grab one of the virtual

sticks by the end with their right hand, and repeatedly stroke their left virtual hand with the

90   stick's end. We did not further instruct the subjects about any detailed way of stroking the hand

such as the exact location to start touching the hand or rhythm. We said that the stick can

92   immerse a bit in the virtual skin, but told them explicitly to be careful to not put the stick through

the touched hand and to keep the hands apart (at stick length) and avoid touching their own

94   body at all times. After approximately 30 to 60 seconds we asked the subjects "if they could feel

anything". If they responded negatively, we let them touch their hand with the stick longer and

96   repeated the question after a while.We did not record the exact time of asking these questions

nor their number. Importantly, we did not cue the subjects as to what the sensation in the

98   "receiving" hand could be.

If the subjects responded positively, we asked them to describe the feeling and noted down their

100   response. We then asked if the sensation moved consistently with the position of the stick on the

skin. Then, we asked the subjects to stroke the middle of the inner side of their palm and rate

102   the intensity of the sensation on a scale from 0 to 10 (0 meant no sensation; we did not further

specify what these levels exactly meant). Afterwards we asked them to touch the middle of their

104   fingers and rate the sensation there (on the same scale as above). Lastly, we asked them to rate

the sensation's intensity in the fingertips. Next, we asked subjects to touch the top of their hand

106   and repeat stroking all three parts mentioned earlier. After obtaining ratings, we asked the

subject to stroke their (not visible) forearm. We then repeated the procedure for the right hand

108   (with the left hand holding the stick). If subjects spontaneously stroke their hand in a different

position than the above order, we did not instruct them to change it, instead we continued the

110   procedure as above until probing all described locations.

Results

All but one of the subjects reported a tingling sensation when touching parts of their hand with a stick. The sensation was usually described as "tingling/static/prickle/electric" or "like wind passing through the hand" (See Figure 2 B and Supplementary Table 1). We call this sensation "phantom touch illusion" (PTI). All subjects that reported this sensation likewise reported that it was consistent with the position of the stick on the avatarized hand. Interestingly, some subjects spontaneously reported they "can feel something" before we asked them for the first time. All subjects described feeling surprised by the sensation caused by PTI. One subject even asked if we were using some device for remote tactile stimulation.

Most importantly, twelve of the subjects (Figure 2, Supplementary Figure 1; Supplementary Table 1) also reported the sensation to be present when touching their non-visible forearm. Two subjects were visibly startled by this feeling.

Figure 2. A) Left-hand results of phantom touch intensity ratings across different locations on the hand and the forearm. Box colors represent the relevant parts of the hand as depicted on the right; gray dots

represent individual data points; horizontal line represents the median; whiskers represent 1.5 times the interquartile range. Compare Supplementary Figure 1 for the whole range of results across all positions in

both hands. B) Frequencies of verbal descriptions of the PTI sensation.

The reported intensity of the stimulation differed across subjects (see Figure 2) but did not significantly differ across stimulation sites and position, as shown by repeated-measures non-

parametric (Friedman) ANOVA for the left hand ($\chi^2[7]=12.2$, $p=0.09$). Additional post hoc tests showed that only the differences between different hand parts and the forearm were at

significance levels below $p<0.1$ though not crossing alpha level of 0.05. This was due to a larger proportion of 'no sensation' answers for the forearm condition. As we did not control exact

stimulation spots, we did not run more detailed tests regarding distinct intensities of the phantom sensation at different spots of the hand surface within each area of interest.

Lastly, several subjects reported PTI when actively touching objects available on the scene (see Supplementary Table 1). Unfortunately, we added this condition only in the course of the

experiment and did not consistently measure the strength of the sensation and whether it corresponded to the touching hand parts. We also did not provide a consistent instruction on

whether subjects had to just touch or grasp the object which might have further influenced the results (i.e., the number of subjects who reported PTI).

Discussion

148 In this study we used immersive virtual reality to simulate scenes of self-touch with a tool without accompanying tactile stimulation. Our subjects reported a tingling sensation at the spots where

150 the virtual stick touched their virtual hand. This happened despite the lack of any physical stimulation. Interestingly, the majority of the subjects reported feeling this phantom sensation

152 when touching their invisible forearm too. We called this sensation "phantom touch illusion" to relate to an apparently similar phenomenon reported in the VR community for social touch (see

154 e.g. Alexdottir et al., 2022).

We ruled out that the sensation was caused by heat, airflow or other physical sensation caused

156 by the moving hand by: 1) ensuring separation of both hands by about 30 cm, that is twice the

distance at which temperature sensation is typically perceived on the skin in normal conditions

(Gheogiu et al., 2001); 2) testing the procedure with a plastic sheet separating both hands which did not affect the sensation.

While our results suggest that touch gating may be predominantly guided by vision because it was induced by a virtual stick and required subjects to look at it, the presence of PTI when

subjects touched their invisible forearm is puzzling. It may indicate that tactile gating uses input from other senses than vision, for example proprioception, or top-down input from the body

schema. The influence of other modalities and input sources (such as body schema) on tactile gating needs to be investigated.

The very overlapping descriptions of the PTI quality across some of our subjects for both passive (stick) and active (object) touch suggest that PTI does not represent predictions about specific

features of object texture or that it does not differ between passive (self) and active (object) touch. Although recent work from Kilteni and Ehrsson (2022) suggests that tactile attenuation

and gating may be distinct processes aimed at modulating intensity and precision of touch, respectively. It is, however, worth mentioning that although we did not investigate touch quality

or precision in detail, we believe there may be subtle differences in tactile gating depending on predicted touch qualities such as suggested by Fueherer et al. (2022). On the other hand, the

same study showed that tactile gating is not limited to predictable stimuli. This lack of prediction-specific gating speaks in favor of the fact that our reported PTI findings likewise represent

stimulus- unspecific cortical events.

An important question remains about the source of neural activity causing PTI.

178 Electrophysiological data shows multi-level processing of tactile gating: cancellation signals are present in the spinal cord and cortical activity in somatosensory, motor and premotor areas

180 (Blakemore et al., 1998; Seki and Fetz, 2012). This multi-level processing likely reflects top-down, predictive control of tactile gating processes before movement is initiated (Seki and Fetz, 2012;

182 Colino et al., 2015). The presence of conscious experience of PTI suggests that visual cues for self-generated touch modulate activity in the somatosensory cortex which is the site for tactile

184 sensations (Gallace and Spence, 2008). However, the fact that some subjects felt PTI when stimulating their invisible forearm, suggests that its nature may be originating more from top-

186 down modulation of somatosensory cortex through the body schema rather than being purely

sensory-driven. Posterior parietal cortex has been often suggested to be a critical site for both

188   the body schema and action predictions (e.g., Frederiks, 1969; Ehrsson et al;, 2005; Desmurget and Sirigu, 2009) and a state comparator across different sensori-motor modalities (Medendorp

190   and Heed, 2019). As such, it may provide top-down input to the somatosensory cortex in the absence of direct vision of the stimulated skin surfaces.

Difference to embodiment illusions

194   While our finding involves feeling touch that is not physical, it does not seem to relate to widely-described embodiment illusions using virtual reality, such as rubber hand illusion (Botvinick and

196   Cohen, 1998). These illusions use simultaneous visual stimulation delivered to an artificial body part (which for example looks like a hand) and tactile stimulation to the user's own corresponding

198   body part which results in an induced feeling that stimulating the artificial hand is like that of the real one. The main difference of phantom touch to these illusions is that they did not involve any

200   sort of tactile stimulation that would then be "transferred" to the virtual hand. We can not completely rule out, however, that the phenomenological effects reported in these illusions

202   involve some sort of predictive tactile gating like we believe our PTI results from. This needs more research.

Open questions

One subject did not have PTI despite a prolonged "induction" phase. We told them specifically to not give a false positive report if they don't feel anything. On a side note, this subject was tested

in a significantly colder room than the others and was visibly distracted by movable objects in the scene. We had no chance to repeat testing with this subject to specifically check whether room

temperature had any effect on their PTI. This subject's lack of phantom sensation during self-touch suggests PTI shows context specificity that may be prone to attentional or other cognitive

demands. Such context influences may be similar to effects reported before, where context seemingly affected object grasping, reflecting qualitative features (temperature) of objects that

users grasp in VR (Blaga et al., 2020). Interestingly, this subject still reported feeling PTI when touching a cup, which indicates there may be differences between active and passive touch as

suggested by previous research (Kilteni and Ehrsson, 2022).

It was recently suggested (e.g., Lush et al., 2020; Fuehrer et al. 2022), embodiment illusions such as the rubber hand may be prone to the influence of demand characteristics of the experimental conditions and phenomenological traits such as suggestibility or phenomenological control. In the case of our experiment, we minimized this confound as we did not provide subjects with any cues about what they should expect and all subjects who had PTI expressed surprise when they felt it. Likewise, the high level of cross-subject congruence in verbal descriptions of the sensation further suggests PTI does not arise from demand characteristics. However, any possible modulatory effects of cognitive demands on PTI, such as mentioned above for temperature perception, seem worth investigating.

Conclusions

We demonstrated an illusory phantom touch sensation emerging during self touch stimulation in the absence of actual tactile stimulus. This PTI was present also when subjects stimulated an invisible, inferred body part. PTI reflects cortical processing of tactile gating resulting in conscious tactile sensation. Moreover, its presence in the absence of the vision of the body parts suggests that tactile gating is not just sensory-driven but additionally influenced by the body schema.

Acknowledgments

This research was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project ID 122679504 – SFB 874. The authors would like to thank Gabriella Andrietta for help with data collection and Jon Walbrin for discussions.

Literature

Angel, R. W., & Malenka, R. C. (1982). Velocity-dependent suppression of cutaneous sensitivity during movement. Experimental neurology, 77(2), 266-274.

Alexdottir, S., & Yang, X. (2022, October). Phantom Touch phenomenon as a manifestation of the Visual-Auditory-Tactile Synaesthesia and its impact on the users in virtual reality. In *2022 IEEE International Symposium on Mixed and Augmented Reality Adjunct (ISMAR-Adjunct)* (pp. 727-732). IEEE.

246 Bays, P. M., Flanagan, J. R., & Wolpert, D. M. (2006). Attenuation of self-generated tactile sensations is predictive, not postdictive. PLoS biology, 4(2), e28.

248 Blaga, A. D., Frutos-Pascual, M., Creed, C., & Williams, I. (2020). Too hot to handle: An evaluation of the effect of thermal visual representation on user grasping interaction in virtual reality. In

250 *Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems* (pp. 1-16).

Blakemore, S. J., Wolpert, D. M., & Frith, C. D. (1998). Central cancellation of self-produced tickle

252 sensation. *Nature neuroscience*, *1*(7), 635–640. https://doi.org/10.1038/2870

Blakemore, S. J., Wolpert, D., & Frith, C. (2000). Why can't you tickle yourself?. *Neuroreport*,

254 *11*(11), R11–R16. https://doi.org/10.1097/00001756-200008030-00002

Botvinick, M., & Cohen, J. (1998). Rubber hands 'feel touch that eyes see. *Nature*, *391*(6669), 756-

256 756.

Chapman, C. E., Beauchamp, E. (2006) Differential controls over tactile detection in humans by

258 motor commands and peripheral reafference. *J. Neurophysiol.* **96**, 1664–1675.

Critchley, M. (1953). *The parietal lobes.* Williams and Wilkins.

260 Desmurget, M., & Sirigu, A. (2009). A parietal-premotor network for movement intention and motor awareness. *Trends in cognitive sciences*, *13*(10), 411-419.

262 Fuehrer, E., Voudouris, D., Lezkan, A., Drewing, K., & Fiehler, K. (2022). Tactile suppression stems from specific sensorimotor predictions. *Proceedings of the National Academy of Sciences*,

264 *119*(20), e2118445119.

Colino, F. L., Buckingham, G., Cheng, D. T., van Donkelaar, P., & Binsted, G. (2014). Tactile gating

266 in a reaching and grasping task. *Physiological reports*, *2*(3), e00267. https://doi.org/10.1002/phy2.26

268 Gallace, A., & Spence, C. (2008). The cognitive and neural correlates of "tactile consciousness": A multisensory perspective. *Consciousness and cognition*, *17*(1), 370-407.

270 Gheoghiu, V. A., Polczyk, R., & Kappeller, C. (2003). The warmth suggestibility scale: A procedure for measuring the influence of suggestion on warm

272 sensations. Personality and Individual Differences, 34(2), 2 19-23 1

Kilteni, K., & Ehrsson, H. H. (2022). Predictive attenuation of touch and tactile gating are distinct

274 perceptual phenomena. Iscience, 25(4), 104077.

276 Lush, P., Botan, V., Scott, R. B., Seth, A. K., Ward, J., & Dienes, Z. (2020). Trait phenomenological control predicts experience of mirror synaesthesia and the rubber hand illusion. *Nature communications*, *11*(1), 1-10.

278 Medendorp, W. P., & Heed, T. (2019). State estimation in posterior parietal cortex: Distinct poles of environmental and bodily states. *Progress in neurobiology*, *183*, 101691.
280 https://doi.org/10.1016/j.pneurobio.2019.101691

Seki, K., & Fetz, E. E. (2012). Gating of sensory input at spinal and cortical levels during
282 preparation and execution of voluntary movement. *Journal of Neuroscience*, *32*(3), 890-902.

Williams, S. R., & Chapman, C. E. (2002). Time course and magnitude of movement-related gating
284 of tactile detection in humans. III. Effect of motor tasks. *Journal of neurophysiology*, *88*(4), 1968-1979.

Supplementary materials

Supplementary figure 1

308   Left- and right-hand results of phantom touch intensity ratings across different locations on the hand

and the forearm. Box colors represent the relevant parts of the hand as depicted on the right; horizontal

310   line is the median; gray dots represent individual data points; whiskers represent 1.5 times the

interquartile range.

Supplementary Table 1

Individual subjects' ratings of PTI intensity, subject data, specific verbal descriptions and their

classifications.

Subjects	Left								Right								Object touch	Moving	Gender	Subject statements	Classification
	Supine				Prone				Supine				Prone								
	Palm	Fingers	Fingertips	Forearm	Palm	Fingers	Fingertips	Forearm	Palm	Fingers	Fingertips	Forearm	Palm	Fingers	Fingertips	Forearm					
1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	1	1	m	Like touching soap bubbles/static	Tingling
2	3	2	3	5	0	0	0		4	2	3	6	3	3	3		1	1	w	Tingling / tickling	Tingling
3	1	1	1	1	1	1	1	1										1	m	Thought I am using stimulator	Tingling
4	1	1	1	1														1	w	Tingling (formigamento)	Tingling
5	2	2	2	4														0	w	Warm feeling, like wood texture	Warm
6	4	2	2		3	3			3	2	2	0	4,5	4,5		0		1	w	Tingling (formigamento)	Tingling
7	3	5	3	0	6	7	6	0	6	6	6	0						1	w	Tingling (formigamento)	Tingling
8	2	1	1	1	1,5	0,5	0,5		2	1	1		0,5	0,5	0,5	0,5		1	w	Tickle	Tickle
9	3	5	5	0	0	5			0	0	0	0	0	0	0			1	m	Like wind going through the hand	Airflow
10	1	1	1	0	1	1	1	0	1	1	1	0	1	1	1	0		1	m		Tingling
11	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	w		
12	5	4	7	2	0	3	5		5	6	7	1	1	5	7		1	1	m	Like bubble of soap	Foam
13	4	6	6	5	8	8	8		5	6	6	0	4	7	7		0	1	w	Like air coming through the hand	Airflow
14	3	3	4	2	3	3	3		2	2	2	1	2	2	2		0	1	w	Like air coming through the hand	Airflow
15	4	6	5	1	7	7	7	1	5	7	7	5	5	7	8	3	1	1	w	It is like a little wind	Airflow
16	4	6	4	2	6	3	1	0	2	2	0	2	3	2	0	4	1	1	w	Like the stick touch, it is soft like velvet	Velvet
17	3	2	2	2	0	1	1	2	3	1	1	1	0	1	1	1	1	1	w	Slihtly cold feeling, as if the hairs on palm and forearm raises	Cold / Tingling
18	3	2	1	4	4	3	1	3	3	2	1	3	3	3	1	3		1	m	mostly prickling, on pinky fingers and forearm warm feeling	Tingling
19	1	1	1	1	2	1	0	0	3	1	1	1	0	1	0	0		0	m	on hand: tickle; on forearm: weak pressure	Tingling
20	6	5	7	3	3	4	3	1	7	7	5	3	5	7	6	3		0	m	on hand: like striking the hair; fingertips had warm feeling	Tingling
21	2	2	4	2	1	4	1	2	4	5	5	0	2	2	3	0		0	m	sensation: tickle	Tingling