IQGAP2 regulates blood-brain barrier immune dynamics

Ketaki A. Katdare[1], Andrew Kjar[2], Natasha M. O'Brown[3], Emma H. Neal[4], Alexander G. Sorets[2], Alena

Shostak[5], Wilber Romero-Fernandez[5], Alexander J. Kwiatkowski[4], Kate Mlouk[4], Hyosung Kim[4], Rebecca

P. Cowell[4], Katrina R. Schwensen[4], Kensley B. Horner[4], John T. Wilson[2,4], Matthew S. Schrag[1,5,6], Sean G.

Megason[3], Ethan S. Lippmann[1,2,4,5,6*]

[1]Vanderbilt Brain Institute, Vanderbilt University, Nashville, TN, USA

[2]Department of Biomedical Engineering, Vanderbilt University, Nashville, TN, USA

[3]Department of Systems Biology, Harvard Medical School, Boston, MA, USA

[4]Department of Chemical and Biomolecular Engineering, Vanderbilt University, Nashville, TN, USA

[5]Department of Neurology, Vanderbilt University Medical Center, Nashville, TN, USA

[6]Vanderbilt Memory and Alzheimer's Center, Vanderbilt University Medical Center, Nashville, TN, USA

*Correspondence to Ethan S. Lippmann (ethan.s.lippmann@vanderbilt.edu)

Abstract

Brain endothelial cells (BECs) play an important role in maintaining central nervous system (CNS)

homeostasis through blood-brain barrier (BBB) functions. BECs express low baseline levels of adhesion

receptors, which limits entry of leukocytes. However, the molecular mediators governing this phenotype

remain mostly unclear. Here, we explored how infiltration of immune cells across the BBB is influenced

by the scaffold protein IQ motif containing GTPase activating protein 2 (IQGAP2). In mice and zebrafish,

we demonstrate that loss of Iqgap2 increases infiltration of peripheral leukocytes into the CNS under

homeostatic and inflammatory conditions. Using single-cell RNA sequencing and immunohistology, we

further show that BECs from mice lacking Iqgap2 exhibit a profound inflammatory signature, including

extensive upregulation of adhesion receptors and antigen-processing machinery. Human tissue analyses

also reveal that Alzheimer's disease is associated with reduced hippocampal IQGAP2. Overall, our results

implicate IQGAP2 as an essential regulator of BBB immune privilege and immune cell entry into the CNS.

Introduction

Maintenance of central nervous system (CNS) homeostasis is crucial for ensuring normal functions in neurons and glial cells, which are sensitive to exogenous molecules in circulation (*1*). The brain is insulated from these factors with the help of a specialized partition known as the blood-brain barrier (BBB). The BBB is composed of barrier-forming brain endothelial cells (BECs) with unique cellular machinery that regulate the entry of macromolecules and solutes into the brain (*2*), including transporters that control the bidirectional exchange of nutrients and waste and tight junctions that prevent passive leakage of blood components into the brain (*3*).

In addition to modulating molecular transport, BECs also act as a selective interface between the peripheral immune system and the brain (*4*). Until recently, the CNS was considered to be completely isolated from the peripheral immune system and therefore an immune-privileged organ (*5*, *6*). However, recent evidence shows that the CNS is under constant immune surveillance to identify and resolve mediators of injury (*7*, *8*). For example, microglia are tissue resident innate immune cells that continually inspect the CNS parenchyma (*9*, *10*). Further, interstitial fluid and cerebrospinal fluid provide drainage pathways for prospective antigens to reach the periphery and stimulate immune cells (*11–14*). As such, initiation of an inflammatory response in the CNS can lead to recruitment of leukocytes across the BBB. BECs facilitate this process through the expression of various receptors, such as leukocyte adhesion molecules (LAMs), that allow interactions with and extravasation of leukocytes into tissue beds (*15*, *16*). In addition, BEC chemokine signaling and antigen presentation (*17–19*) also play a key role in orchestrating leukocyte extravasation. While the underlying molecular mechanisms of leukocyte extravasation are similar across all organs, BECs have been shown to express very low levels of LAMs under homeostatic conditions, making them refractory to mild inflammatory cues (*1*, *20*). However, despite several exquisite single-cell RNA sequencing (scRNA-seq) studies identifying differentially expressed genes between BECs and peripheral endothelial cells (*21–26*), as well as scRNA-seq profiles of different cell types in the

neurovascular unit (*24, 25*), the mechanisms underlying suppression of LAM expression in BECs and

general limitation of immune cell extravasation across the BBB remain poorly understood.

IQ motif containing GTPase activating protein 1 (IQGAP1), a ubiquitously expressed scaffolding protein,

has been recently implicated in facilitating leukocyte trafficking across peripheral endothelium (*27, 28*).

Historically, IQGAP1 was known as a regulator of cellular signaling due to its role as a scaffolding protein

(*29*). It also acts as an oncogene driving hepatocellular carcinogenesis (*30, 31*). Interestingly, IQ motif

containing GTPase activating protein 2 (IQGAP2), a related member of the same scaffolding protein

family, acts as a tumor suppressor to counteract the oncogenic effects of IQGAP1 (*32, 33*). Since IQGAP2

is believed to suppress IQGAP1 function, and IQGAP2 expression is predicted in multiple neurovascular

support cells such as astrocytes and microglia (*21, 34–36*), we hypothesized that IQGAP2 may influence

inflammatory responses and leukocyte extravasation at the BBB. Herein, using multiple *in vivo* models, we

show that loss of Iqgap2 increases leukocyte infiltration into the CNS under various conditions. scRNA-

seq of BECs from wildtype and Iqgap2[-/-] mice further reveals an upregulation of multiple inflammatory

genes and signaling pathways involved in BBB-immune cell interactions. Further, using postmortem human

brain tissue, we determined that IQGAP2 was reduced in the hippocampus of patients with AD. Overall,

our results benchmark IQGAP2 as a key molecular player involved in BBB-immune crosstalk and

leukocyte entry into the CNS.

Results

Global loss of Iqgap2 increases peripheral immune cell infiltration into the brain in a mouse model of acute neuroinflammation

To initially assess whether murine Iqgap2 influences leukocyte infiltration into the brain, we delivered interleukin 1-beta (IL1β) into the lateral ventricles of wildtype and Iqgap2$^{-/-}$ mice (129S background) to induce acute neuroinflammation (*37, 38*). After 24 hours, infiltration of leukocytes was assessed by immunohistochemical labeling of CD45+ cells in cortical brain sections. We measured a significant increase in the number of CD45+ cells in the cortex of Iqgap2$^{-/-}$ mice as compared to their wildtype littermates (Figure 1). We also confirmed that delivery of saline to the lateral ventricles in a similar fashion did not stimulate CD45+ cell infiltration into the brains of both wildtype and Iqgap2$^{-/-}$ mice (Figure 1B). These data suggest that Iqgap2 constrains leukocyte infiltration into the mouse CNS following a central inflammatory challenge.

Global loss of Iqgap2 increases immune cell infiltration in experimental autoimmune encephalomyelitis

As dysregulated immune cell infiltration is a hallmark of several neurodegenerative conditions (*39–45*), we sought to next evaluate whether loss of Iqgap2 affects immune cell access to the CNS in the presence of an inflammatory neurodegenerative condition. As such, to monitor the effects of Iqgap2 loss under a chronic inflammatory insult (*46*), we induced experimental autoimmune encephalomyelitis (EAE) in wildtype and Iqgap2$^{-/-}$ 129S mice (Figure 2A). To validate our experimental strategy, we concurrently induced EAE in C57BL/6 mice and observed robust development of disease (Supplementary Figure 1). We measured a significant increase in CD45+ cells in the Iqgap2$^{-/-}$ lumbar spinal cord compared to wildtype 129S mice 30 days after EAE induction (Figure 2B). Interestingly, this increase in infiltrating leukocytes did not produce significant differences in disease severity, probability of survival, or demyelination (Figure 2C-E),

suggesting the infiltration events may be decoupled from pathology. Overall, these data further indicate that

loss of Iqgap2 contributes to increased leukocyte extravasation into the CNS under extended

neuroinflammation.

Loss of Iqgap2 increases infiltration of peripheral immune cells into the brain in zebrafish in the

absence of inflammation

Our data in multiple inflammatory mouse models suggest that Iqgap2 plays an important role in BBB

immune privilege. To corroborate these findings in an additional species, we generated zebrafish crispants

by direct injection of Cas9 protein with multiple sgRNAs to target genes of interest in single-cell embryos

with both endothelial cells (kdrl:mCherry) and immune cells (mpeg1:EGFP) transgenically labeled. We

specifically assessed the presence of mpeg+ macrophage lineage cells in the brains of 5 days post

fertilization (dpf) zebrafish, which have a functional BBB (*47*), that were either uninjected or targeting *tyr*

or *iqgap2.* While uninjected fish are expected to retain normal function of all genes, *tyr* crispants should

have mosaic knockout of tyrosinase*,* a protein involved in pigment production that is not expected to affect

leukocyte infiltration; this serves as an additional CRISPR injection control. Both controls displayed

similarly low numbers of mpeg+ cells in the brain, while *iqgap2* crispants displayed a significant increase

in the number of mpeg+ cells (Figure 3B). Since mpeg also labels brain resident microglia, we used an

established method to label microglia with Neutral Red dye (*48*) and distinguish these cells from infiltrating

leukocytes. Uninjected controls and *iqgap2* crispants were therefore treated with Neutral Red and all

mpeg+/Neutral Red+ double-positive microglia and mpeg+/Neutral Red- infiltrating leukocytes were

quantified throughout the entire zebrafish brain. We measured a significant increase in the total number of

mpeg+ cells, similar to previous experiments, but not double-positive microglia in the *iqgap2* crispants,

and the number of mpeg+/Neutral Red- infiltrating leukocytes was significantly increased (Figure 3C).

These data suggest that *iqgap2* is essential for restricting the infiltration of leukocytes into the CNS under

homeostatic conditions in zebrafish, as its mosaic depletion enhanced the entry of peripheral leukocytes

into the brain.

Global loss of Iqgap2 yields a profound inflammatory transcriptomic profile in mouse BECs

Our data indicate that Iqgap2 plays an important role in restricting peripheral immune access to the CNS

and in modulating responses to inflammatory insults. Iqgap2 is a large scaffolding protein known to

orchestrate many different cellular functions such as regulating cytoskeletal organization, cytokinesis, and

carcinogenesis (*29*), suggesting it could govern many different signaling axes that would influence BBB

function and cellular crosstalk within the neurovascular unit. After confirming general expression of Iqgap2

in mouse brain and enriched vessel fractions (Supplementary Figure 2), we performed scRNA-seq on BECs

isolated from wildtype and Iqgap2$^{-/-}$ mice to better understand how loss of Iqgap2 affects BBB function.

Here, to generate an endothelial cell-enriched population for sequencing, we isolated antibody-labeled

CD31+ cells from dissociated mouse brains using fluorescence-activated cell sorting (Figure 4A). After

implementing quality control metrics, cells were first analyzed using dimension-reduction by uniform

manifold approximation and projection (UMAP) and unsupervised clustering to obtain 12 unique identities:

endothelial cells (EC), PLVAP-expressing endothelial cells (EC_plvap), hemoglobin-expressing

endothelial cells (EC_hb), endothelial/stromal cells or pericyte-like cells (EC/PC), endothelial/stromal cells

or astrocyte-like cells (EC/AC), astrocytes (AC), B cells, T cells, monocytes (MNC), microglia (MG),

oligodendrocytes (OLG), and fibroblasts (Supplementary Figure 3). Endothelial cells were the largest

represented cell type, followed by immune cells such as monocytes and T cells that are also predicted to

express CD31 (Figure 4B and Supplementary Figure 3); other cell types likely represent some small

contamination in the sorting process. Each cell type was annotated using previously established marker

genes (*49*) and all non-endothelial clusters were filtered out of the dataset for these initial analyses.

Wildtype (WT) and Iqgap2$^{-/-}$ (KO) genotypes were equally represented in the EC cluster (Figure 4C), and

bulk gene expression comparisons between WT and KO BECs show 928 differentially regulated genes

(DEGs) (Figure 4D). Since leukocyte extravasation occurs predominantly at post-capillary venules (*50–*

52), we examined these DEGs along the neurovascular tree. We used unsupervised clustering to subcluster

BECs into 6 sub-populations, and based on marker gene expression, these subclusters were further

classified into arterial (A; genes *Hey1, Bmx, and Sema3g*), capillary (C; genes *Slc16a2, Car4, and Mfsd2a*),

and venous (V; genes *Icam1, Slc38a5, and Vwf*) zonal identities (Figure 4E and Supplementary Figure 3)

(*21*). Further analysis suggests that the strongest DEGs are shared among zonal identities and include genes

involved in antigen presentation, interleukin receptor subunits, and adhesion molecules. Unique DEGs

identified in the A and V zonal identities did not show any significant functional enrichment whereas a

similar analysis of unique DEGs identified in the C zonal identity suggest subtle changes in BEC function

(Figure 4F and Supplementary Table 1).

Across these zones, we were able to confirm that loss of Iqgap2 does not significantly affect expression of

most canonical BBB genes, including junction proteins and nutrient transporters such as *Cdh5, Cldn5, Ocln,*

Tjp, and *Slc2a1* (Supplementary Figure 4A). At the protein level, total vessel density was unchanged, and

we found no obvious deficits in expression of claudin-5, occludin, ZO-1, and Glut1 in the Iqgap2$^{-/-}$ mice

(Supplementary Figure 5). We did observe significant differences in gene expression for certain transporters

that facilitate exchange of amino acids and metabolites across the BBB, such as *Abcb1a*, *Slc7a1*, *Slc7a5*,

and *Slc16a1* (Supplementary Figure 4B). In addition, major regulator genes involved in BBB functional

development and maintenance like *Mfsd2a* (*53*) and *Ctnnb1* (*54–56*) were significantly downregulated in

the Iqgap2$^{-/-}$ BECs, suggesting possible connections to BBB dysfunction (Supplementary Figure 4C).

Interestingly, we also saw significant upregulation of several LAMs and chemokine receptors. Although

interaction with leukocytes and response to inflammation is primarily facilitated by venous ECs, we found

that loss of Iqgap2 significantly upregulates expression of leukocyte receptors and signaling molecules like

Vcam1, Icam1, and *Ackr1* across multiple vascular zones (Figure 5A). Using immunohistochemistry, we

confirmed upregulation of Vcam1 in cortical vasculature of Iqgap2$^{-/-}$ mice (Figure 5B). The widespread

expression of vascular Vcam1 in Iqgap2$^{-/-}$ mice was particularly striking, given that Vcam1 expression on

BECs has been implicated in brain inflammation and cognitive decline in mice (*57*). We have also recently

shown that vascular VCAM-1 expression is significantly increased in Alzheimer's disease cortex relative

to asymptomatic age-matched controls, further highlighting its links to human neurodegeneration (*58*).

To extend our understanding of potential pathways in BECs affected by global loss of Iqgap2, we performed

gene set enrichment analysis (GSEA) for KEGG signaling pathways. GSEA indicated that pathways

involved in response to infections like Kaposi sarcoma-associated herpes virus infection (KSHV), human

T-cell leukemia virus 1 infection (HTLV), human immunodeficiency virus 1 infection (HIV), and human

papillomavirus infection (HPV) were upregulated. In addition, other pathways facilitating immune

interactions like cell adhesion (*Vcam1, Icam1*), antigen processing and presentation (*Psme2, Hspa5, Canx,*

Calr), and TNF signaling (*Cxcl1, Csf1, Ptgs2*) were also significantly upregulated in the Iqgap2$^{-/-}$ BECs

(Figure 5C and Supplementary Figure 6). These results indicate that *Iqgap2* loss shifts both the

transcriptional profile and protein expression of BECs towards an activated, inflammatory state.

Due to the significant upregulation of LAMs in the Iqgap2$^{-/-}$ BECs, we further analyzed cell-cell interactions

between BECs and other cell types identified in the scRNA-seq dataset using CellChat (*59*). We found that

BEC-immune cell interactions were overrepresented in Iqgap2$^{-/-}$ mice. Quantification of these results

suggested an increase in cell-cell interactions between BECs and microglia as well as BECs and peripheral

immune cells like monocytes, T cells, and B cells. We then assessed the predicted directionality of these

interactions by analyzing known receptor-ligand pairs. BECs were predicted to be the "senders" whereas

immune populations, especially monocytes, were the primary "receivers" (Supplementary Figure 7). To

understand whether these changes were due to Iqgap2 loss in a specific cell type, we performed DEG

analyses in all clusters annotated as immune populations. Monocytes had the highest number of significant

DEGs, followed by microglia and T cells (Supplementary Figure 8A). KEGG pathway analysis of DEGs

in the monocytes indicated upregulation of pathways such as leukocyte transendothelial migration and

regulation of actin cytoskeleton (Supplementary Figure 8B). These data indicate that Iqgap2 may also play

an important role in modulating BEC-monocyte communication, which could additionally contribute to the

overall inflammatory phenotype observed in the BECs in Iqgap2$^{-/-}$ mice.

IQGAP2 in postmortem human brain tissue

To putatively assess connections between IQGAP2 and human disease states, we evaluated IQGAP2

protein distribution patterns in postmortem human hippocampal tissue from patients with Alzheimer's

disease (AD) and in cases without AD. Using a custom polyclonal antibody raised against a peptide with

selective homology to human IQGAP2, we immunostained and quantified vascular-associated and

parenchymal IQGAP2 signal in human hippocampal sections. IQGAP2 staining was strongly detected

along blood vessels (identified by collagen expression), with more diffuse and punctate signal observed in

the parenchyma (Figure 6). A significant decrease in IQGAP2 levels was found in parenchymal tissue (non-

vascular regions) in AD patients (Figure 6). Connections between immune cell entry into the brain and

neurodegeneration are becoming increasingly scrutinized (*60*, *61*), highlighting the potential importance of

this finding.

Discussion

Peripheral endothelial cells display high levels of LAMs and can respond swiftly to local and systemic inflammatory cues (*62*). This is followed by rapid infiltration of leukocytes into surrounding tissue beds. In comparison, BECs are comparably immune quiescent under homeostatic conditions and generally express low levels of LAMs. This allows the BBB to more selectively control the activation of downstream inflammatory pathways and extravasation of leukocytes into the CNS (*63*, *64*). Although peripheral immune responses are essential for the resolution of CNS injury, it is well established that age-related neurological deficits and chronic neurodegeneration may be exacerbated in part by unwarranted entry of leukocytes into the brain (*39*, *41*, *65–67*). Further, BECs upregulate transcriptional signatures of inflammatory responses during aging and neurological disease (*24*, *25*, *68*). As such, identifying mechanisms that regulate BEC inflammatory responses is critical for understanding the pathological progression of these diseases and developing strategies to decrease leukocyte extravasation. Our study provides evidence that IQGAP2 plays an important role in BBB immune dynamics and the propensity of leukocytes to enter the CNS after immune stimulation.

To our knowledge, the role of IQGAP proteins in BBB integrity has not been studied. *IQGAP2* belongs to the *IQGAP* family of scaffolding proteins involved in orchestrating a wide array of intracellular signaling and cytoskeleton dynamics (*29*). The multidomain structure of these proteins acts as a framework for complex formation of signaling proteins, thus influencing many downstream cellular processes. *IQGAPs* were historically considered to be modulators of cytoskeletal architecture. However, it has become apparent that their role extends into other physiological processes like glomerular filtration in the kidney, cardiomyocyte function in the heart, smooth muscle cell contraction in lung airways, and metabolism in the liver (*29*, *69*, *70*). *IQGAP2* has also been studied in the context of its tumor-suppressive characteristics (*71*). Previous studies identified *IQGAP2* as a novel tumor suppressor gene specifically linked to the development of hepatocellular carcinoma. More recently, *IQGAP2* inactivation has been linked to other malignancies like gastric cancer (*72*), prostate cancer (*73*), and bladder cancer (*74*). Moreover, reduced

expression of *IQGAP2* is associated with worsened cancer pathology and poor clinical outcomes (*73, 75,*

76). Our study provides new context for the role of IQGAP2 in physiological processes related to immune

cell trafficking.

One limitation of our study is that we cannot exclusively ascribe the observed CNS leukocyte infiltration

to loss of Iqgap2 expression in a particular cell type. *IQGAP2* has generally high expression in many

immune cell subtypes (*77*), but this has not previously been associated with specific phenotypes like tissue

extravasation and responsiveness to inflammatory cytokines. Our current data may suggest that cell-to-cell

communication between BECs and immune cells plays a role in BBB inflammation when Iqgap2 is lost.

However, more extensive scRNA-seq profiling of the neurovascular unit in Iqgap2$^{-/-}$ mice, as well as

lineage-specific Iqgap2 knockout models, will be necessary in future studies to fully clarify cell-intrinsic

and cell-extrinsic effects of IQGAP2 on BBB immune privilege.

An additional unanswered question is whether changes in IQGAP2 expression contribute to human

neurodegenerative disease progression through modulation of immune cell recruitment to the CNS. In the

acute and chronic inflammatory animal models used in this study, we did not observe significant differences

in pathology. In the EAE model, it is possible that the 30-day timepoint does not reflect increased damage

caused by infiltrating leukocytes due to other compensatory mechanisms (e.g. significant damage could

occur at an earlier time point followed by regeneration), especially when considering the biphasic disease

severity in the Iqgap2$^{-/-}$ mice and that deaths generally occurred during periods where disease severity

worsened. In human hippocampal tissue, we observed a decrease in IQGAP2 in patients with AD. Single-

cell datasets indicate parenchymal IQGAP2 expression is restricted to microglia, while vascular IQGAP2

expression could potentially be localized to perivascular macrophages or fibroblasts (*21, 24, 36*).

Interestingly, microglia have been recently connected to T cell infiltration in mouse models of tauopathy

(*78*) and general aging (*79*), and T cell infiltration and resultant activation of microglia has been shown to

exacerbate neurodegeneration in engineered human cell-based models of AD (*80*). In the context of our

255 study, these findings motivate future exploration into whether loss of Iqgap2 influences pathology and

256 immune cell infiltration in mouse and human models of AD.

257 Overall, our work reveals a novel role for IQGAP2 in regulating BBB immune dynamics. While the cell-

258 specific effects of IQGAP2 are currently unclear, our collective data suggest that this protein plays an

259 important conserved and previously unrecognized role in suppressing BEC inflammatory responses and

260 regulating immune cell trafficking to the CNS through non-cell autonomous mechanisms. Future work will

261 determine these mechanisms of action and their relevance to brain disorders.

Methods

Methods

<u>Mouse maintenance and procedures</u>

<u>Colony maintenance</u>: All mouse protocols were approved by the Institutional Animal Care and Use

Committee at Vanderbilt University. Iqgap2$^{-/-}$ mice (129S background) were obtained from Jackson

Laboratory (strain 025452). Male and female Iqgap2$^{-/-}$ mice and wildtype littermate controls were used for

all experiments. Mice were at least 8 weeks of age at the time of use and were housed continuously in an

environmentally controlled facility in a 12-hour light/dark cycle with *ad libitum* access to food and water.

<u>Genotyping</u>: Mice were ear-tagged and tail snips were collected at approximately 2 weeks of age. Genomic

DNA was extracted using an Extracta DNA Prep kit for tissue (Quantabio) per manufacturer instructions.

DNA was extracted in the Extraction buffer at 95°C for 30 minutes, cooled to room temperature and mixed

with Stabilization buffer before being stored at -20°C. The reactions were performed using the Apex Hot

Start Taq BLUE Master Mix (Apex Bioresearch) on a ProFlex PCR system (Applied Biosystems). A

touchdown cycling protocol was used with an initial annealing temperature of 65°C gradually lowered to

60°C over the course of 10 cycles. Genotyping primers used were as follows: Mutant Reverse–

ATTTGTCACGTCCTGCACGACG, Wildtype Reverse–TGGCCTTCCTCCCTTAAAGT, and Common

Forward–TGACTCAGAGGGCACATGGT. PCR products were run on a 2% agarose-TAE buffered gel

supplemented with SYBR Safe DNA Gel Stain (Invitrogen) and imaged using a LI-COR Odyssey Fc gel

imager (Supplementary Figure 9).

<u>Tissue collection</u>: Mice were deeply anesthetized using high-dose isoflurane and euthanized by transcardial

perfusion of 1X DPBS (Gibco), followed by 4% paraformaldehyde (PFA, Thermo Fisher Scientific). Brains

and lumbar segment of spinal cords were extracted and postfixed in 4% PFA overnight followed by

cryopreservation in sucrose gradient solutions (15% and 30%, respectively). The tissue was then embedded

in OCT medium (Tissue-Tek) and 15 µm (brain) and 25 µm (spinal cord) thick sections were cut and stored

at -80⁰C.

<u>Microvessel isolation</u>: Mice were deeply anesthetized using high-dose isoflurane and euthanized by

decapitation. Brains were extracted and collected in ice-cold PBS. Microvessels from the cortex were

isolated as previously described (81). In brief, cortices were dissected from remaining brain tissue,

homogenized in PBS using a tissue homogenizer (Wheaton), and collected by centrifugation. Homogenized

tissue was resuspended in a 15% dextran solution (~70,000 kDa, Sigma) and centrifuged at 10,000xg to

separate the vessel fraction from the remaining tissue. The vessel fraction was washed with PBS and filtered

using a 40 µm cell strainer (Corning). For protein extraction, the vessel fraction was incubated in RIPA

buffer (Sigma) supplemented with 1% v/v protease and phosphatase inhibitor cocktails (Sigma) for 30-60

minutes on ice. Cell debris was separated by centrifugation (12,000xg for 15 minutes at 4⁰C) and the

supernatant was stored at -20⁰C. Protein concentration was quantified using a Pierce BCA Protein Assay

(Thermo Fisher Scientific) according to the manufacturer instructions. For immunohistochemical analysis,

microvessel suspension was placed on glass slides and allowed to dry at room temperature. Dried

microvessels were then fixed with 4% parafolmaldehyde solution and labelled with Lectin DyLight 488 for

30 minutes at room temperature before mounting in Prolong Gold Antifade Mountant (Invitrogen).

<u>Intracerebroventricular injection of IL1β</u>: Male and female Iqgap2$^{-/-}$ mice and wildtype littermates were

used as experimental animals. All animals were at least 12 weeks old at the time of the surgery. Under

isoflurane anesthesia, mice were unilaterally injected into the lateral ventricle using a stereotactic apparatus

at coordinates of AP = -0.3 mm, ML = -1 mm, and DV = -3 mm. After injections, mice were returned to

prewarmed home cages for recovery. Each mouse received 20 ng/µL of IL1β solution (20 ng in 1 µL PBS)

or an equivalent volume of sterile PBS. 24 hours after surgery, mice were transcardially perfused with PBS

followed by 4% paraformaldehyde and brains were extracted for immunohistological analysis. For

quantification, CD45+ cells were manually counted in each section under blinded conditions. Vasculature

was labelled using a fluorescence-conjugated GLUT1 antibody to solely quantify CD45+ cells that had

extravasated out of the vessels into the brain parenchyma.

Experimental autoimmune encephalomyelitis (EAE): Female Iqgap2$^{-/-}$ mice and wildtype littermates were

used as experimental animals. All animals were between 9 and 13 weeks of age at the time of induction.

EAE kits (Hooke Laboratories) targeting MOG$_{35-55}$ antigen were used. 100 µL of MOG$_{35-55}$/Complete

Freund's Adjuvant emulsion was injected subcutaneously at the scruff of the neck and near the base of the

tail resulting in a total injection volume of 200 µL into each mouse. At 2 and 24 hours post injection of

emulsion, 100 µL of pertussis toxin (4 µg/mL) was injected intraperitoneally. Clinical scores were evaluated

starting 7 days post induction as follows: score 1, flaccid tail; score 2, weak hind limbs; score 3, hind limb

paralysis; score 4, quadriplegia. Clinical scores were recorded every day for the first week after

development of symptoms followed by every other day thereafter. Premature deaths were recorded. 30 days

following induction, mice were transcardially perfused with PBS followed by 4% paraformaldehyde. Brains

and spinal cords were extracted for immunohistological analysis. For quantification, CD45+ cells were

manually counted in each section under blinded conditions. Vasculature was labelled using a fluorescence-

conjugated GLUT1 antibody to ensure that CD45+ cells had extravasated out of the vessels. For quantifying

EAE pathology in spinal cord, total area and demyelination area were calculated using the "Measure" tool

in ImageJ by manually outlining regions of interest as indicated by Luxol Fast Blue stain.

Zebrafish maintenance and procedures

Zebrafish were maintained at 28.5°C following standard protocols (82). All zebrafish work was approved

by the Harvard Medical Area Standing Committee on Animals under protocol number IS00001263-3. Adult

fish were maintained on a standard light-dark cycle from 8 am to 10 pm. Adult fish, aged 3 months to 1.5

years, were crossed to produce embryos and larvae. For imaging live larvae, 0.003% phenylthiourea (PTU)

was used beginning at 1 dpf to inhibit melanin production. These studies used the AB wildtype strains and

the transgenic reporter strains (Tg(kdrl:HRAS-mCherry)[s896] (*83*), abbreviated as Tg(kdrl:mCherry), and

Tg(mpeg1:EGFP)[gl22] (*84*), abbreviated as Tg(mpeg1:EGFP). Mosaic *iqgap2* crispants were generated by

injection of 7 μM Cas9 protein complexed with four sgRNAs (5'-AGTAGCCTCGATTTACAGG-3', 5'-

GCACTTTGTCAGTCACGGAA-3', 5'-CAGGACAGCGCGAGCACTG-3', and 5'-AAAGTCCGCGCG

CAGTTTA-3') to target multiple sites in the *iqgap2* transcript, and *tyr* control crispants were similarly

targeted with four sgRNAs (5'-GCCGCACACAGAGCCGTCGC-3', 5'-GGATGCATTATTACGTGTCC

-3', 5'- GACTCTACATCGGCGGATGT-3', and 5'-GTATCCGTCGTTGTGTCCGA-3'). To distinguish

between microglia and macrophages, 4 dpf larvae were exposed to 2.5 μg/ml of Neutral Red (Millipore

Sigma: N7005) in embryo water for 3 hours at 28.5°C. Larvae were washed at least three times to remove

the residual dye and then microglia were assessed the next day as previously described (*48*). Zebrafish

larvae were immobilized by tricaine exposure and live imaged on a Leica SP8 line scanning confocal

microscope. Quantification of mpeg+ and Neutral Red+ cells was manually performed on blinded z-stack

images that spanned the entire larval head using ImageJ.

<u>Human brain tissue and preparation</u>

Human brain tissue was obtained at autopsy and prepared as previously described (*85*). De-identified brain

tissue was obtained from the Vanderbilt Brain and Biospecimen Bank at Vanderbilt University Medical

Center. Written informed consent for brain donation was obtained from patients or their surrogate decision

makers. All brain tissue collection was authorized by the Institutional Review Board at Vanderbilt

University Medical Center. Demographics and neuropathological information for each donor are listed in

Supplementary Table 2.

Human brain tissue was obtained at autopsy and immersion fixed in 10% formalin (Thermo Fisher

Scientific) at 4°C for 1-3 days. The fixative solution was then removed and the tissued rinsed with 1x TBS

(Corning) three times for 5 minutes each. The tissue was placed in sterile 10% sucrose (Millipore Sigma)

/1x TBS/0.02% sodium azide (NaN$_3$, Millipore Sigma) until tissue sank and then 30% sucrose/1x

TBS/0.02% NaN$_3$ for overnight at 4°C or until the tissue sank. The tissue block was affixed to the stage of

vibratome using cyanoacrylate cement and fully submerged in 1x TBS. Hippocampal sections were

prepared at 50 μm thickness. Floating tissue sections were transferred to 15 mL Falcon tubes with antigen

retrieval buffer (10 mM citric acid pH 6.0, Millipore Sigma) containing 0.05% Tween-20 (Millipore Sigma)

and heated to 95°C for 20 minutes in a block heater. Hippocampal sections were then washed with 100 mM

glycine (Millipore Sigma)/1x TBS/0.1% Triton X-100 (Millipore Sigma) buffer for 30 minutes followed

by permeabilization with 0.3% Triton X-100/1x TBS buffer for 30 minutes and two washes for 5 minutes

each with 1x TBS at room temperature.

<u>Development of a custom antibody against human IQGAP2 protein</u>

A peptide corresponding to amino acid residues 1460-1474 of human IQGAP2 (RSIKLDGKGEPKGAK)

was synthesized with an amino-terminal cysteine and conjugated to maleimide-activated Keyhole Limpet

Haemocyanin (KLH), maleimide-activated bovine serum albumin (BSA), and SulfoLink resin using

manufacturer protocols (Thermo Fisher). The peptide-KLH conjugate was used to immunize rabbits

(Cocalico Biologicals). Rabbit antisera were tested for the presence of antibodies recognizing the IQGAP2

peptide by dot blot analysis using the peptide-BSA conjugate. The rabbit antibodies were affinity-purified

from the antisera using the peptide-SulfoLink resin, where 5 mL of rabbit sera was diluted 1:1 with PBS

and passed over a 2 mL peptide-SulfoLink column. After extensive washing with PBS, bound antibodies

were eluted with 8.5 mL 0.1 M Glycine (pH 2.2) and collected in a tube containing 1.5 mL of 1 M Tris (pH

8). Antibody solution was stored at -80°C.

<u>Immunofluorescent staining</u>

<u>Mouse tissue</u>: Tissue slices were retrieved from the -80ºC freezer and allowed to thaw at room temperature

for 10-15 minutes. Sections were washed with 1X PBS with 0.03% Triton X-100 to remove the OCT

medium. Sections were then blocked using a goat serum blocking buffer and incubated in primary antibody

solution overnight at 4°C. After incubation, primary antibody solution was thoroughly washed off and

sections were incubated in secondary antibody solution for 2 hours at room temperature. All antibodies and

corresponding dilutions used for immunohistochemistry are listed in Supplementary Tables 3-4. Following

final washes, tissue was mounted in Prolong Gold Antifade Mountant with DAPI (Invitrogen) and slides

were allowed to dry overnight before imaging on a Leica DMi8 epifluorescence microscope. All acquired

images were processed and quantified using ImageJ software. For quantification of vascular Vcam1

expression, mean Vcam1 intensity was calculated within CD31+ vessels using ImageJ.

<u>Human tissue</u>: Immunohistochemical labeling in hippocampal tissue slices was performed as previously

described (*85*, *86*) with minor modifications. Tissue slices were incubated with primary antibodies for 48

hours at 4ºC followed by secondary antibodies for 2 hours at room temperature. All antibodies and

corresponding dilutions used for immunohistochemistry are listed in Supplementary Tables 3-4. Neuritic

plaques, neurofibrillary tangles and related AD pathological structures were additionally stained using 1

µM 4,4'-[(2-methoxy-1,4-phenylene) di-(1E)-2,1-ethenediyl] bisphenol (MX-04) (Tocris) for 15 minutes

at room temperature. Confocal images were acquired using the Zeiss LSM 710 confocal laser-scanning

microscope (Carl Zeiss AG) with a 20× air/dry or 63× oil objective and 10 µm z-stack scanning projections

with a step interval of 1 µm or one scanning projection, with a minimum resolution of 1500 x 1500 pixels.

Vascular IQGAP2 expression was quantified using mean IQGAP2 intensity within Collagen+ area and

parenchymal IQGAP2 expression was quantified by gating Collagen- area using ImageJ.

<u>Luxol Fast Blue staining</u>

Tissue slices were retrieved from the -80ºC freezer and allowed to thaw at room temperature for 10-15

minutes. Sections were first allowed to dry overnight at room temperature and then immersed in a 70%

ethanol solution overnight to facilitate defatting. Luxol Fast Blue stain (Abcam) was applied to the sections

and incubated for 5 to 6 hours at 60ºC. Excess stain was washed off by consecutive dipping in fresh absolute

ethanol. Slides were differentiated briefly using lithium carbonate solution and washed with distilled water

and alcohol solution. Lastly, sections were counterstained with Cresyl Etch Violet, washed with distilled

water, dehydrated with absolute alcohol and mounted in DPX mounting medium (Sigma-Aldrich).

Mounting medium was cured overnight before imaging on a Leica DMi8 inverted microscope.

<u>Western blotting</u>

Protein samples from mouse tissue were prepared by diluting 20-40 µg of protein with 1X Laemmli buffer

(Biorad) supplemented with beta-mercaptoethanol (Sigma) and Ultrapure water (Gibco) to a final volume

of 20-30 µL. Samples were then boiled at 95ºC for 5 minutes, cooled on ice, and centrifuged briefly.

Samples were then loaded into 4-20% Criterion TGX Midi protein gels (Biorad) along with Precision Plus

Dual Color Protein ladder (Biorad) and run at 80-120V. Protein gels were then transferred onto iBlot2

Nitrocellulose membranes (Thermo Fisher Scientific) using an iBlot2 transfer device (Thermo Fisher

Scientific). Membranes were cut to size and blocked for at least 30 minutes at room temperature in Intercept

TBS Blocking buffer (LI-COR Biosciences) on a shaker. Membranes were submerged in primary

antibodies diluted in blocking buffer with 0.05% Tween20 (Sigma) and incubated at 4ºC overnight.

Following primary antibody incubation, membranes were washed in wash buffer (1X tris buffered saline

with 0.05% Tween-20). Membranes were incubated in secondary antibodies diluted in the wash buffer at

room temperature for 2 hours. All primary and secondary antibody information is listed in Supplementary

Tables 3-4. Blots were imaged using a LI-COR Odyssey Clx or Fc Imager and bands were quantified using

Image Studio Lite software.

Single-cell RNA sequencing

Fresh cortical tissue was homogenized and delipidated using an Adult Brain Dissociation Kit (Miltenyi
Biotec) according to the manufacturer's instructions. The resultant single-cell suspension was incubated
with TruStain FcX (Biolegend) for 10 minutes at 4°C to prevent non-specific antibody binding of Fc
receptors, labeled with a secondary conjugated CD31 antibody (1:2000, eBioscience) for 30 minutes, and
counterstained with DAPI. Live CD31+ cells were flow sorted using a 4-laser FACS Aria III sorter (BD
Biosciences) at the Vanderbilt Flow Cytometry Shared Resource. Live cells were resuspended in DMEM
(Gibco) supplemented with 2% FBS to obtain a concentration of 700-1200 cells/µL. RNA extraction, 10X
Genomics Chromium 5' scRNAseq Library Prep, and sequencing on an Illumina NovaSeq6000 sequencer
was performed at the VANTAGE core facility.

Single-cell RNA sequencing analysis

All gene by counts data were read into Seurat (v.4). The initial data were filtered to retain only cells with
RNA counts between 1,200 and 20,000 (with less than 10% being mitochondrial). Within each biological
experiment, data were log normalized. The top 2,000 variable features (as identified by variance) were
selected. Samples were combined using CCA (Seurat v.4) and then standardized. UMAP based on the first
50 principal components was used to reduce dimensionality for visualization and clustering (*87*). Predicted
doublets were filtered out using DoubletFinder (*88*). Unsupervised clustering was achieved using the KNN-
graph approach in Seurat (*89*), and clusters were annotated with SingleR (*90*) based on previously published
datasets (*49*). Predominantly endothelial clusters (0,2,4,5) were rescaled and clustered based on the first 15
principal components. Clusters were annotated as arterial, capillary, or venous based on the expression of
established marker gene sets (*21*). Approximately 2,000 ECs were analyzed per condition. Differentially
expressed genes (DEGs) were computed in Seurat using the FindMarkers() function with built-in

Bonferroni correction. KEGG or Reactome gene set enrichment analysis (GSEA) was computed on

significant DEGs using WebGestalt with a false discovery rate cutoff of 0.05 and weighted set cover

redundancy reduction (*91*). Communication between cell types was predicted using the CellChat R

packages (*59*). For visualization, *EnhancedVolcano* (*92*), *UpSetR* shiny app (*93*), *iGraph* (*94*), and *ggvenn*

were employed.

<u>Statistical analysis</u>

Statistical analysis for single-cell RNA sequencing data was conducted in R. All other analysis was

performed in GraphPad Prism 9.0.0.

Data availability

Raw sequencing data files have been deposited in the ArrayExpress collection under accession code E-

MTAB-12687. All code used for the single-cell RNA sequencing analyses is publicly available at:

https://github.com/LippmannLab/IQGAP2_WT_KO_BEC_scRNAseq.

Author contributions

KAK, ESL, and EHN conceived the study. KAK and ESL designed the majority of experiments with input

from the other authors. KAK, NMO, EHN, AGS, AS, AJK, HK, and RPC conducted all experiments. AK

performed all bioinformatics analyses. KM assisted with data quantification. WRF performed human tissue

histology. KRS and KBH provided support on animal husbandry and takedowns for experiments. MSS

provided human tissue samples and contributed to interpretation of histological images. JTW and SGM

provided input on experimental planning and data interpretation.

Acknowledgments

Funding was provided by a Chan Zuckerberg Initiative Ben Barres Early Career Acceleration Award (grant

2019-191850 to ESL) and NIH grant R21 NS106510 (to ESL). AK and EHN were supported by the

National Science Foundation Graduate Research Fellowship Program. NMO was supported by a Damon

Runyon Postdoctoral Fellowship. Support for RNA sequencing was provided by Vanderbilt Technologies

for Advanced Genomics core facility, which is supported by the CTSA Grant (NIH grant 5UL1 RR024975),

the Vanderbilt Ingram Cancer Center (NIH grant P30 CA68485), the Vanderbilt Vision Center (NIH grant

P30 EY08126), and NIH/NCRR grant G20 RR030956. Support for histology was provided by Vanderbilt

Translational Pathology Shared Resource core facility, which is supported by P30 CA068485. Flow

cytometry experiments were performed in the Vanderbilt Flow Cytometry Shared Resource, which is

supported by P30 CA068485 and the Vanderbilt Digestive Disease Research Center (P30 DK058404). The

custom IQGAP2 antibody in this study was produced by the Vanderbilt Antibody and Protein Resource,

which was supported by P30 CA068485. Image acquisition was performed in part through the use of the

Vanderbilt Cell Imaging Shared Resource, which is supported by NIH grants P30 CA68485, P30 DK20593,

P30 EY08126, and S10RR027396. The authors thank Dr. Jose Gomez for early assistance with the Iqgap2^{-}

$^{/-}$ mice and Dr. Eric Shusta for sharing microarray data that initially indicated the potential relevance of

IQGAP2. Some figures in this manuscript were created in part using BioRender.

Competing interests

493 The authors declare no competing interests.

    Figures

Figure 1: Global loss of Iqgap2 increases infiltration of peripheral leukocytes in a mouse model of

acute neuroinflammation.

A) Schematic representation of experimental design for assessing leukocyte numbers in mouse brains 24

hours after intracerebroventricular saline or IL1β delivery.

B) Representative images and quantification of CD45+ immune cells (magenta) and vasculature (green)

in wildtype (WT) and Iqgap2$^{-/-}$ (KO) mouse brain cortex following treatment with saline or IL1β. Data

are represented as mean ± SD, and each data point represents an individual mouse, where at least 5

images were quantified per mouse. For saline treatment, N=3 WT mice and N=3 KO mice, and for

506 IL1β treatment, N=4 WT mice and N=6 KO mice. Statistical significance was calculated using one-

507 way ANOVA with Tukey's multiple comparison's test (*, $p<0.05$, **, $p<0.01$, ****$p<0.0001$).

Figure 2: Global loss of Iqgap2 increases infiltration of peripheral leukocytes in EAE.

A) Schematic representation of experimental design for assessing response to EAE 30 days following

induction.

B) Representative images and quantification of CD45+ immune cells (magenta) and vasculature (green) in wildtype (WT) and Iqgap2$^{-/-}$ (KO) lumbar spinal cord at 30 days following EAE induction. Data are represented as mean ± SD, and each data point represents an individual mouse. At least 8 images were quantified per mouse. N=8 WT mice, N=6 KO mice. Statistical significance was calculated using the unpaired student's t-test (**, $p<0.01$).

C) EAE score curve for wildtype (WT) and Iqgap2$^{-/-}$ (KO) mice following EAE induction. Data are presented as mean ± SEM. N=8 WT mice, N=6 KO mice. Statistical significance was calculated using the unpaired student's t-test on area under the curve.

D) Probability of survival in WT versus KO mice following EAE induction. N=10 WT and KO mice. Statistical significance was calculated using Log-rank (Mantel-Cox) test.

E) Representative images and quantification of demyelinating lesions in WT and KO lumbar spinal cord section stained with Luxol Fast Blue at 30 days following EAE induction. Data are represented as mean ± SD, and each data point represents an individual mouse, where 1 image was quantified per mouse. N=8 WT mice, N=6 KO mice. Statistical significance was calculated using the unpaired student's t-test.

Figure 3: Mosaic loss of *iqgap2* expression increases infiltration of peripheral immune cells into the zebrafish brain.

A) Schematic representation of experimental design for assessing leukocyte numbers in the larval zebrafish brain. Double transgenic (kdrl:mCherry; mpeg:EGFP) single-cell embryos were injected with Cas9 protein and sgRNAs to target genes of interest. These mosaic crispants were then allowed to develop normally, and mpeg+ leukocytes were quantified in the brain at 5 dpf.

B) Representative 100 µm thick maximum intensity projection images and quantification of macrophage

lineage cells (mpeg:EGFP) in the brains of uninjected (UN) controls, *tyr* crispant controls, and *iqgap2*

crispants. Vasculature is marked with the kdrl:mCherry transgene (magenta). N=30 fish (UN), 15 fish

(*tyr*), and 52 fish (*iqgap2*). Data are represented as mean ± SD, and each data point represents an

individual fish. Statistical significance was calculated using a one-way ANOVA with Tukey's multiple

comparison's test (****, $p<0.0001$).

C) Representative images and quantification of infiltrating leukocytes versus tissue-resident microglia in

the brains of *iqgap2* crispants versus uninjected (UN) controls. Representative 30 µm thick maximum

intensity projection images of the dorsal (top) or ventral (bottom) brain regions and quantification of

mpeg+ (green) macrophage lineage cells and mpeg+/Neutral Red+ (magenta) microglia. Yellow

arrowheads indicate individual mpeg+/Neutral Red- infiltrating leukocytes in the ventral brain of

iqgap2 crispants. N=16 fish (UN) and 20 fish (*iqgap2*). Data represented as mean ± SD, and each data

point represents an individual fish. Statistical significance was calculated using an unpaired Student's

t-test (****, $p<0.0001$).

Figure 4: Global loss of Iqgap2 yields extensive transcriptional changes in BECs.

A) Schematic representation of experimental design. Whole brain cortices were isolated from wildtype (WT) and Iqgap2$^{-/-}$ (KO) mice, dissociated to a single cell suspension using enzyme-based dissociation techniques, and sorted to enrich for CD31+ cells before sequencing.

B) UMAP cell annotations based on unsupervised clustering. EC = endothelial cells, EC/PC = EC/stromal cells (pericytes), EC/AC = EC/stromal cells (astrocytes), MNC = monocytes, MG = microglia.

C) UMAP comparison between WT (grey) and KO (pink) cells.

D) Volcano plot highlighting differentially expressed genes in the EC cluster. 928 genes were significantly altered, with 584 upregulated and 344 downregulated by loss of Iqgap2. Red dots indicate genes with $p < 0.05$ and >2-fold change in expression.

E) UMAP of endothelial zonal identity based on unsupervised clustering. A = arterial, C = capillary, V = venous.

F) Venn diagram showing number of DEGs shared between zonal identities.

Figure 5: Global loss of Iqgap2 produces a widespread inflammatory phenotype in BECs.

A) Split violin plots indicating differential gene expression of select inflammatory markers across vascular zones between wildtype (WT) and Iqgap2^-/- (KO) BECs. Statistical significance was calculated using Wilcox rank-order tests with Bonferroni correction (***, p<0.001).

B) Representative images and quantification of vascular Vcam1 expression in WT versus KO mouse cortex. Quantification was performed across N=3 biological replicates. Data are represented as mean ± SD (black bars). Biological replicates are represented as squares and measurements from individual images are represented as circles color coded to each replicate. Statistical significance was calculated using the student's unpaired t-test (*, p<0.05).

C) GSEA analysis for signaling pathways upregulated in KO versus WT BECs. Each node represents an enriched gene set belonging to the labeled canonical pathway. Nodes are colored based on p-value, and thickness of the connecting lines indicates similarity of overlapping genes represented in connected gene sets. KSHV = Kaposi sarcoma-associated herpes virus infection, HTLV = Human T-cell leukemia

580   virus 1 infection, HIV = Human immunodeficiency virus 1 infection, PPER = Protein processing in

581   endoplasmic reticulum, HPV = Human papillomavirus infection.

Figure 6: IQGAP2 distribution in human hippocampus.

A) Representative confocal microscopy images of IQGAP2 (green) in human hippocampal tissue from AD and non-AD donors. Images were produced from 10 μm z-stack scanning projections with a step interval of 1 μm. Vasculature was stained with Collagen (red) and β-amyloid and neuritic plaques, neurofibrillary tangles and other tau aggregates were stained with Methoxy-X04 (blue).

B) Mean IQGAP2 intensity in vascular regions was quantified solely one scanning projection in collagen+ area, while IQGAP2 intensity in parenchyma was quantified by gating samples to

591   exclude collagen+ area. Data are presented as mean $\pm$ SD, where each data point represents an

592   individual donor with at least five images quantified per donor. N=7 non-AD donors and N=7 AD

593   donors. Statistical significance was calculated using unpaired student's t-test (**, $p<0.01$).

References

1. C. P. Profaci, R. N. Munji, R. S. Pulido, R. Daneman, The blood–brain barrier in health and disease:
Important unanswered questions. *J. Exp. Med.* **217**, e20190062 (2020).

2. R. Daneman, A. Prat, The Blood–Brain Barrier. *Cold Spring Harb. Perspect. Biol.* **7**, a020412 (2015).

3. B. Obermeier, R. Daneman, R. M. Ransohoff, Development, maintenance and disruption of the blood-
brain barrier. *Nat. Med.* **19**, 1584–1596 (2013).

4. B. Engelhardt, C. Coisne, Fluids and barriers of the CNS establish immune privilege by confining
immune surveillance to a two-walled castle moat surrounding the CNS castle. *Fluids Barriers CNS*
8, 4 (2011).

5. A. Louveau, T. H. Harris, J. Kipnis, Revisiting the Mechanisms of CNS Immune Privilege. *Trends*
Immunol. **36**, 569–577 (2015).

6. B. Engelhardt, P. Vajkoczy, R. O. Weller, The movers and shapers in immune privilege of the CNS.
Nat. Immunol. **18**, 123–131 (2017).

7. L. Marchetti, B. Engelhardt, Immune cell trafficking across the blood-brain barrier in the absence and
presence of neuroinflammation. *Vasc. Biol.* **2**, H1–H18 (2020).

8. J. A. Mapunda, H. Tibar, W. Regragui, B. Engelhardt, How Does the Immune System Enter the Brain?
Front. Immunol. **13**, 805657 (2022).

9. G. Raivich, Like cops on the beat: the active role of resting microglia. *Trends Neurosci.* **28**, 571–573
(2005).

10. S. S. Ousman, P. Kubes, Immune surveillance in the central nervous system. *Nat. Neurosci.* **15**, 1096–
1101 (2012).

11. J. D. Laman, R. O. Weller, Drainage of Cells and Soluble Antigen from the CNS to Regional Lymph
Nodes. *J. Neuroimmune Pharmacol.* **8**, 840–856 (2013).

12. A. Louveau, B. A. Plog, S. Antila, K. Alitalo, M. Nedergaard, J. Kipnis, Understanding the functions
and relationships of the glymphatic system and meningeal lymphatics. *J. Clin. Invest.* **127**, 3210–
3219 (2017).

13. B. Engelhardt, R. O. Carare, I. Bechmann, A. Flügel, J. D. Laman, R. O. Weller, Vascular, glial, and
lymphatic immune gateways of the central nervous system. *Acta Neuropathol. (Berl.)* **132**, 317–338
(2016).

14. S. P. das Neves, N. Delivanoglou, S. Da Mesquita, CNS-Draining Meningeal Lymphatic Vasculature:
Roles, Conundrums and Future Challenges. *Front. Pharmacol.* **12**, 655052 (2021).

15. B. Engelhardt, R. M. Ransohoff, The ins and outs of T-lymphocyte trafficking to the CNS: anatomical
sites and molecular mechanisms. *Trends Immunol.* **26**, 485–495 (2005).

16. B. Engelhardt, R. M. Ransohoff, Capture, crawl, cross: the T cell code to breach the blood–brain
barriers. *Trends Immunol.* **33**, 579–589 (2012).

17. S. Aydin, J. Pareja, V. M. Schallenberg, A. Klopstein, T. Gruber, N. Page, E. Bouillet, N. Blanchard, R. Liblau, J. Körbelin, M. Schwaninger, A. J. Johnson, M. Schenk, U. Deutsch, D. Merkler, B. Engelhardt, Antigen recognition detains CD8+ T cells at the blood-brain barrier and contributes to its breakdown. *Nat. Commun.* **14**, 3106 (2023).

18. C. E. Fain, J. Zheng, F. Jin, K. Ayasoufi, Y. Wu, M. T. Lilley, A. R. Dropik, D. M. Wolf, R. C. Rodriguez, A. Abudumijiti, Z. P. Tritz, S. M. Bouchal, L. L. Pewe, S. L. Urban, Y. Chen, S.-Y. Chang, M. J. Hansen, J. M. Kachergus, J. Shi, E. A. Thompson, H. E. Jensen, J. T. Harty, I. F. Parney, J. Sun, L.-J. Wu, A. J. Johnson, Discrete class I molecules on brain endothelium differentially regulate neuropathology in experimental cerebral malaria. *Brain*, awad319 (2023).

19. M. A. Lopes Pinheiro, A. Kamermans, J. J. Garcia-Vallejo, B. van het Hof, L. Wierts, T. O'Toole, D. Boeve, M. Verstege, S. M. van der Pol, Y. van Kooyk, H. E. de Vries, W. W. Unger, Internalization and presentation of myelin antigens by the brain endothelium guides antigen-specific T cell migration. *eLife* **5**, e13149 (2016).

20. I. Galea, The blood–brain barrier in systemic infection and inflammation. *Cell. Mol. Immunol.* **18**, 2489–2501 (2021).

21. M. Vanlandewijck, L. He, M. A. Mäe, J. Andrae, K. Ando, F. Del Gaudio, K. Nahar, T. Lebouvier, B. Laviña, L. Gouveia, Y. Sun, E. Raschperger, M. Räsänen, Y. Zarb, N. Mochizuki, A. Keller, U. Lendahl, C. Betsholtz, A molecular atlas of cell types and zonation in the brain vasculature. *Nature* **554**, 475–480 (2018).

22. M. F. Sabbagh, J. S. Heng, C. Luo, R. G. Castanon, J. R. Nery, A. Rattner, L. A. Goff, J. R. Ecker, J. Nathans, Transcriptional and epigenomic landscapes of CNS and non-CNS vascular endothelial cells. *eLife* **7**, e36187 (2018).

23. J. Kalucka, L. P. M. H. de Rooij, J. Goveia, K. Rohlenova, S. J. Dumas, E. Meta, N. V. Conchinha, F. Taverna, L.-A. Teuwen, K. Veys, M. García-Caballero, S. Khan, V. Geldhof, L. Sokol, R. Chen, L. Treps, M. Borri, P. de Zeeuw, C. Dubois, T. K. Karakach, K. D. Falkenberg, M. Parys, X. Yin, S. Vinckier, Y. Du, R. A. Fenton, L. Schoonjans, M. Dewerchin, G. Eelen, B. Thienpont, L. Lin, L. Bolund, X. Li, Y. Luo, P. Carmeliet, Single-Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell* **180**, 764-779.e20 (2020).

24. A. C. Yang, R. T. Vest, F. Kern, D. P. Lee, M. Agam, C. A. Maat, P. M. Losada, M. B. Chen, N. Schaum, N. Khoury, A. Toland, K. Calcuttawala, H. Shin, R. Pálovics, A. Shin, E. Y. Wang, J. Luo, D. Gate, W. J. Schulz-Schaeffer, P. Chu, J. A. Siegenthaler, M. W. McNerney, A. Keller, T. Wyss-Coray, A human brain vascular atlas reveals diverse mediators of Alzheimer's risk. *Nature* **603**, 885–892 (2022).

25. F. J. Garcia, N. Sun, H. Lee, B. Godlewski, H. Mathys, K. Galani, B. Zhou, X. Jiang, A. P. Ng, J. Mantero, L.-H. Tsai, D. A. Bennett, M. Sahin, M. Kellis, M. Heiman, Single-cell dissection of the human brain vasculature. *Nature* **603**, 893–899 (2022).

26. E. A. Winkler, C. N. Kim, J. M. Ross, J. H. Garcia, E. Gil, I. Oh, L. Q. Chen, D. Wu, J. S. Catapano, K. Raygor, K. Narsinh, H. Kim, S. Weinsheimer, D. L. Cooke, B. P. Walcott, M. T. Lawton, N. Gupta, B. V. Zlokovic, E. F. Chang, A. A. Abla, D. A. Lim, T. J. Nowakowski, A single-cell atlas of the normal and malformed human brain vasculature. *Science* **375**, eabi7377 (2022).

27. D. P. Sullivan, P. J. Dalal, F. Jaulin, D. B. Sacks, G. Kreitzer, W. A. Muller, Endothelial IQGAP1
regulates leukocyte transmigration by directing the LBRC to the site of diapedesis. *J. Exp. Med.* **216**,
2582–2601 (2019).

28. D. P. Sullivan, P. J. Dalal, D. B. Sacks, W. A. Muller, Calcium Signaling Regulates Leukocyte
Transendothelial Migration through the Action of Endothelial Cell IQGAP1, Calmodulin, and
CaMKIIδ. *FASEB J.* **34**, 1–1 (2020).

29. J. M. Smith, A. C. Hedman, D. B. Sacks, IQGAPs choreograph cellular signaling from the membrane
to the nucleus. *Trends Cell Biol.* **25**, 171–184 (2015).

30. Q. Dai, Q. Ain, M. Rooney, F. Song, A. Zipprich, Role of IQ Motif-Containing GTPase-Activating
Proteins in Hepatocellular Carcinoma. *Front. Oncol.* **12**, 920652 (2022).

31. X. Jin, Y. Liu, J. Liu, W. Lu, Z. Liang, D. Zhang, G. Liu, H. Zhu, N. Xu, S. Liang, The Overexpression
of IQGAP1 and β-Catenin Is Associated with Tumor Progression in Hepatocellular Carcinoma In
Vitro and In Vivo. *PLOS ONE* **10**, e0133770 (2015).

32. C. D. White, H. Khurana, D. V. Gnatenko, Z. Li, R. D. Odze, D. B. Sacks, V. A. Schmidt, IQGAP1
and IQGAP2 are Reciprocally Altered in Hepatocellular Carcinoma. *BMC Gastroenterol.* **10**, 125
(2010).

33. V. A. Schmidt, C. S. Chiariello, E. Capilla, F. Miller, W. F. Bahou, Development of Hepatocellular
Carcinoma in *Iqgap2* -Deficient Mice Is IQGAP1 Dependent. *Mol. Cell. Biol.* **28**, 1489–1502 (2008).

34. Y. Zhang, K. Chen, S. A. Sloan, M. L. Bennett, A. R. Scholze, S. O'Keeffe, H. P. Phatnani, P.
Guarnieri, C. Caneda, N. Ruderisch, S. Deng, S. A. Liddelow, C. Zhang, R. Daneman, T. Maniatis,
B. A. Barres, J. Q. Wu, An RNA-Sequencing Transcriptome and Splicing Database of Glia, Neurons,
and Vascular Cells of the Cerebral Cortex. *J. Neurosci.* **34**, 11929–11947 (2014).

35. Y. Zhang, S. A. Sloan, L. E. Clarke, C. Caneda, C. A. Plaza, P. D. Blumenthal, H. Vogel, G. K.
Steinberg, M. S. B. Edwards, G. Li, J. A. Duncan, S. H. Cheshier, L. M. Shuer, E. F. Chang, G. A.
Grant, M. G. H. Gephart, B. A. Barres, Purification and Characterization of Progenitor and Mature
Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse. *Neuron* **89**, 37–
53 (2016).

36. L. He, M. Vanlandewijck, M. A. Mäe, J. Andrae, K. Ando, F. Del Gaudio, K. Nahar, T. Lebouvier,
B. Laviña, L. Gouveia, Y. Sun, E. Raschperger, Å. Segerstolpe, J. Liu, S. Gustafsson, M. Räsänen,
Y. Zarb, N. Mochizuki, A. Keller, U. Lendahl, C. Betsholtz, Single-cell RNA sequencing of mouse
brain and lung vascular and vessel-associated cell types. *Sci. Data* **5**, 180160 (2018).

37. M. G. Proescholdt, S. Chakravarty, J. A. Foster, S. B. Foti, E. M. Briley, M. Herkenham,
Intracerebroventricular but not intravenous interleukin-1β induces widespread vascular-mediated
leukocyte infiltration and immune signal mRNA expression followed by brain-wide glial activation.
Neuroscience **112**, 731–749 (2002).

38. X. Liu, D. P. Nemeth, D. B. McKim, L. Zhu, D. J. DiSabato, O. Berdysz, G. Gorantla, B. Oliver, K.
G. Witcher, Y. Wang, C. E. Negray, R. S. Vegesna, J. F. Sheridan, J. P. Godbout, M. J. Robson, R.
D. Blakely, P. G. Popovich, S. D. Bilbo, N. Quan, Cell-Type-Specific Interleukin 1 Receptor 1
Signaling in the Brain Regulates Distinct Neuroimmune Activities. *Immunity* **50**, 764–766 (2019).

39.   S. Amor, F. Puentes, D. Baker, P. van der Valk, Inflammation in neurodegenerative diseases.
Immunology **129**, 154–169 (2010).

40.   D. Gate, E. Tapp, O. Leventhal, M. Shahid, T. J. Nonninger, A. C. Yang, K. Strempfl, M. S. Unger,
T. Fehlmann, H. Oh, D. Channappa, V. W. Henderson, A. Keller, L. Aigner, D. R. Galasko, M. M.
Davis, K. L. Poston, T. Wyss-Coray, CD4[+] T cells contribute to neurodegeneration in Lewy body
dementia. *Science* **374**, 868–874 (2021).

41.   D. Gate, N. Saligrama, O. Leventhal, A. C. Yang, M. S. Unger, J. Middeldorp, K. Chen, B. Lehallier,
D. Channappa, M. B. De Los Santos, A. McBride, J. Pluvinage, F. Elahi, G. K.-Y. Tam, Y. Kim, M.
Greicius, A. D. Wagner, L. Aigner, D. R. Galasko, M. M. Davis, T. Wyss-Coray, Clonally expanded
CD8 T cells patrol the cerebrospinal fluid in Alzheimer's disease. *Nature* **577**, 399–404 (2020).

42.   M. S. Unger, E. Li, L. Scharnagl, R. Poupardin, B. Altendorfer, H. Mrowetz, B. Hutter-Paier, T. M.
Weiger, M. T. Heneka, J. Attems, L. Aigner, CD8+ T-cells infiltrate Alzheimer's disease brains and
regulate neuronal- and synapse-related gene expression in APP-PS1 transgenic mice. *Brain. Behav.*
Immun. **89**, 67–86 (2020).

43.   R. H. Earls, J.-K. Lee, The role of natural killer cells in Parkinson's disease. *Exp. Mol. Med.* **52**, 1517–
1525 (2020).

44.   B. Hemmer, M. Kerschensteiner, T. Korn, Role of the innate and adaptive immune responses in the
course of multiple sclerosis. *Lancet Neurol.* **14**, 406–419 (2015).

45.   J. Stephenson, E. Nutma, P. Van Der Valk, S. Amor, Inflammation in CNS neurodegenerative
diseases. *Immunology* **154**, 204–219 (2018).

46.   C. S. Constantinescu, N. Farooqi, K. O'Brien, B. Gran, Experimental autoimmune encephalomyelitis
(EAE) as a model for multiple sclerosis (MS). *Br. J. Pharmacol.* **164**, 1079–1106 (2011).

47.   N. M. O'Brown, S. G. Megason, C. Gu, Suppression of transcytosis regulates zebrafish blood-brain
barrier function. *eLife* **8**, e47326 (2019).

48.   C. E. Shiau, K. R. Monk, W. Joo, W. S. Talbot, An Anti-inflammatory NOD-like Receptor Is
Required for Microglia Development. *Cell Rep.* **5**, 1342–1352 (2013).

49.   L. Zhao, Z. Li, J. S. L. Vong, X. Chen, H.-M. Lai, L. Y. C. Yan, J. Huang, S. K. H. Sy, X. Tian, Y.
Huang, H. Y. E. Chan, H.-C. So, W.-L. Ng, Y. Tang, W.-J. Lin, V. C. T. Mok, H. Ko,
Pharmacologically reversible zonation-dependent endothelial cell transcriptomic changes with
neurodegenerative disease associations in the aged brain. *Nat. Commun.* **11**, 4413 (2020).

50.   B. Rossi, S. Angiari, E. Zenaro, S. L. Budui, G. Constantin, Vascular inflammation in central nervous
system diseases: adhesion receptors controlling leukocyte-endothelial interactions. *J. Leukoc. Biol.*
89, 539–556 (2011).

51.   W. A. Muller, Getting Leukocytes to the Site of Inflammation. *Vet. Pathol.* **50**, 7–22 (2013).

52.   S. Nourshargh, R. Alon, Leukocyte Migration into Inflamed Tissues. *Immunity* **41**, 694–707 (2014).

53.   A. Ben-Zvi, B. Lacoste, E. Kur, B. J. Andreone, Y. Mayshar, H. Yan, C. Gu, Mfsd2a is critical for
the formation and function of the blood–brain barrier. *Nature* **509**, 507–511 (2014).

54.   R. Daneman, D. Agalliu, L. Zhou, F. Kuhnert, C. J. Kuo, B. A. Barres, Wnt/β-catenin signaling is
required for CNS, but not non-CNS, angiogenesis. *Proc. Natl. Acad. Sci.* **106**, 641–646 (2009).

55.   J. M. Stenman, J. Rajagopal, T. J. Carroll, M. Ishibashi, J. McMahon, A. P. McMahon, Canonical
Wnt Signaling Regulates Organ-Specific Assembly and Differentiation of CNS Vasculature. *Science*
322, 1247–1250 (2008).

56.   S. Liebner, M. Corada, T. Bangsow, J. Babbage, A. Taddei, C. J. Czupalla, M. Reis, A. Felici, H.
Wolburg, M. Fruttiger, M. M. Taketo, H. von Melchner, K. H. Plate, H. Gerhardt, E. Dejana, Wnt/β-
catenin signaling controls development of the blood–brain barrier. *J. Cell Biol.* **183**, 409–417 (2008).

57.   H. Yousef, C. J. Czupalla, D. Lee, M. B. Chen, A. N. Burke, K. A. Zera, J. Zandstra, E. Berber, B.
Lehallier, V. Mathur, R. V. Nair, L. N. Bonanno, A. C. Yang, T. Peterson, H. Hadeiba, T. Merkel, J.
Körbelin, M. Schwaninger, M. S. Buckwalter, S. R. Quake, E. C. Butcher, T. Wyss-Coray, Aged
blood impairs hippocampal neural precursor activity and activates microglia via brain endothelial cell
VCAM1. *Nat. Med.* **25**, 988–1000 (2019).

58.   H. Kim, K. Leng, J. Park, A. G. Sorets, S. Kim, A. Shostak, R. J. Embalabala, K. Mlouk, K. A.
Katdare, I. V. L. Rose, S. M. Sturgeon, E. H. Neal, Y. Ao, S. Wang, M. V. Sofroniew, J. M. Brunger,
D. G. McMahon, M. S. Schrag, M. Kampmann, E. S. Lippmann, Reactive astrocytes transduce
inflammation in a blood-brain barrier model through a TNF-STAT3 signaling axis and secretion of
alpha 1-antichymotrypsin. *Nat. Commun.* **13**, 6581 (2022).

59.   S. Jin, C. F. Guerrero-Juarez, L. Zhang, I. Chang, R. Ramos, C.-H. Kuan, P. Myung, M. V. Plikus, Q.
Nie, Inference and analysis of cell-cell communication using CellChat. *Nat. Commun.* **12**, 1088
(2021).

60.   H. D. Mason, D. B. McGavern, How the immune system shapes neurodegenerative diseases. *Trends*
Neurosci. **45**, 733–748 (2022).

61.   T. R. Hammond, S. E. Marsh, B. Stevens, Immune Signaling in Neurodegeneration. *Immunity* **50**,
955–974 (2019).

62.   M. P. Bevilacqua, Endothelial-Leukocyte Adhesion Molecules. *Annu. Rev. Immunol.* **11**, 767–804
(1993).

63.   W. F. Hickey, B. L. Hsu, H. Kimura, T-lymphocyte entry into the central nervous system. *J. Neurosci.*
Res. **28**, 254–260 (1991).

64.   K. Rössler, C. Neuchrist, K. Kitz, O. Scheiner, D. Kraft, H. Lassmann, Expression of leucocyte
adhesion molecules at the human blood-brain barrier (BBB). *J. Neurosci. Res.* **31**, 365–374 (1992).

65.   G. Yilmaz, D. N. Granger, Cell adhesion molecules and ischemic stroke. *Neurol. Res.* **30**, 783–793
(2008).

66.   T. K. Roberts, Leukocyte transmigration across the blood-brain barrier: perspectives on neuroAIDS.
Front. Biosci. **15**, 478 (2010).

67.   B. W. Dulken, M. T. Buckley, P. Navarro Negredo, N. Saligrama, R. Cayrol, D. S. Leeman, B. M.
George, S. C. Boutet, K. Hebestreit, J. V. Pluvinage, T. Wyss-Coray, I. L. Weissman, H. Vogel, M.

M. Davis, A. Brunet, Single-cell analysis reveals T cell infiltration in old neurogenic niches. *Nature* **571**, 205–210 (2019).

68. M. B. Chen, A. C. Yang, H. Yousef, D. Lee, W. Chen, N. Schaum, B. Lehallier, S. R. Quake, T. Wyss-Coray, Brain Endothelial Cells Are Exquisite Sensors of Age-Related Circulatory Cues. *Cell Rep.* **30**, 4418-4432.e4 (2020).

69. A. C. Hedman, J. M. Smith, D. B. Sacks, The biology of IQGAP proteins: beyond the cytoskeleton. *EMBO Rep.* **16**, 427–446 (2015).

70. A. Sen, S. Youssef, K. Wendt, S. Anakk, Depletion of IQ motif-containing GTPase activating protein 2 (IQGAP2) reduces hepatic glycogen and impairs insulin signaling. *J. Biol. Chem.* **299**, 105322 (2023).

71. Yanyun Xie, Anil Kapoor, Hao Peng, Jean-Claude Cutz, Lijian Tao, Damu Tang, IQGAP2 Displays Tumor Suppression Functions. *J. Anal. Oncol.* **4** (2015).

72. L. Xu, Y. Shao, L. Ren, X. Liu, Y. Li, J. Xu, Y. Ye, IQGAP2 Inhibits Migration and Invasion of Gastric Cancer Cells via Elevating SHIP2 Phosphatase Activity. *Int. J. Mol. Sci.* **21**, 1968 (2020).

73. Y. Xie, L. Zheng, L. Tao, Downregulation of IQGAP2 Correlates with Prostate Cancer Recurrence and Metastasis. *Transl. Oncol.* **12**, 236–244 (2019).

74. F. Song, R. Kotolloshi, M. Gajda, M. Hölzer, M.-O. Grimm, D. Steinbach, Reduced IQGAP2 Promotes Bladder Cancer through Regulation of MAPK/ERK Pathway and Cytokines. *Int. J. Mol. Sci.* **23**, 13508 (2022).

75. Q. Dai, F. Song, X. Li, F. Huang, H. Zhao, Comprehensive analysis of the expression and prognosis for IQ motif-containing GTPase-activating proteins in hepatocellular carcinoma. *BMC Cancer* **22**, 1121 (2022).

76. Z. Deng, L. Wang, H. Hou, J. Zhou, X. Li, Epigenetic regulation of IQGAP2 promotes ovarian cancer progression via activating Wnt/β-catenin signaling. *Int. J. Oncol.* **48**, 153–160 (2016).

77. M. Uhlén, L. Fagerberg, B. M. Hallström, C. Lindskog, P. Oksvold, A. Mardinoglu, Å. Sivertsson, C. Kampf, E. Sjöstedt, A. Asplund, I. Olsson, K. Edlund, E. Lundberg, S. Navani, C. A.-K. Szigyarto, J. Odeberg, D. Djureinovic, J. O. Takanen, S. Hober, T. Alm, P.-H. Edqvist, H. Berling, H. Tegel, J. Mulder, J. Rockberg, P. Nilsson, J. M. Schwenk, M. Hamsten, K. von Feilitzen, M. Forsberg, L. Persson, F. Johansson, M. Zwahlen, G. von Heijne, J. Nielsen, F. Pontén, Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015).

78. X. Chen, M. Firulyova, M. Manis, J. Herz, I. Smirnov, E. Aladyeva, C. Wang, X. Bao, M. B. Finn, H. Hu, I. Shchukina, M. W. Kim, C. M. Yuede, J. Kipnis, M. N. Artyomov, J. D. Ulrich, D. M. Holtzman, Microglia-mediated T cell infiltration drives neurodegeneration in tauopathy. *Nature* **615**, 668–677 (2023).

79. X. Zhang, R. Wang, H. Chen, C. Jin, Z. Jin, J. Lu, L. Xu, Y. Lu, J. Zhang, L. Shi, Aged microglia promote peripheral T cell infiltration by reprogramming the microenvironment of neurogenic niches. *Immun. Ageing* **19**, 34 (2022).

80. M. Jorfi, J. Park, C. K. Hall, C.-C. J. Lin, M. Chen, D. Von Maydell, J. M. Kruskop, B. Kang, Y. Choi, D. Prokopenko, D. Irimia, D. Y. Kim, R. E. Tanzi, Infiltrating CD8+ T cells exacerbate Alzheimer's disease pathology in a 3D human neuroimmune axis model. *Nat. Neurosci.* **26**, 1489–1504 (2023).

81. Y.-K. Lee, H. Uchida, H. Smith, A. Ito, T. Sanchez, The isolation and molecular characterization of cerebral microvessels. *Nat. Protoc.* **14**, 3059–3081 (2019).

82. M. Westerfield, *The Zebrafish Book: A Guide for the Laboratory Use of Zebrafish (Brachydanio Rerio)* (1993).

83. N. C. Chi, R. M. Shaw, S. De Val, G. Kang, L. Y. Jan, B. L. Black, D. Y. R. Stainier, Foxn4 directly regulates tbx2b expression and atrioventricular canal formation. *Genes Dev.* **22**, 734–739 (2008).

84. F. Ellett, L. Pase, J. W. Hayman, A. Andrianopoulos, G. J. Lieschke, mpeg1 promoter transgenes direct macrophage-lineage expression in zebrafish. *Blood* **117**, e49-56 (2011).

85. W. Romero-Fernandez, C. Carvajal-Tapia, A. Prusky, K. A. Katdare, E. Wang, A. Shostak, L. Ventura-Antunes, H. J. Harmsen, E. S. Lippmann, K. Fuxe, J. A. MacGurn, D. O. Borroto-Escuela, M. S. Schrag, Detection, visualization and quantification of protein complexes in human Alzheimer's disease brains using proximity ligation assay. *Sci. Rep.* **13**, 11948 (2023).

86. A. G. Nackenoff, T. J. Hohman, S. M. Neuner, C. S. Akers, N. C. Weitzel, A. Shostak, S. M. Ferguson, B. Mobley, D. A. Bennett, J. A. Schneider, A. L. Jefferson, C. C. Kaczorowski, M. S. Schrag, PLD3 is a neuronal lysosomal phospholipase D associated with β-amyloid plaques and cognitive function in Alzheimer's disease. *PLOS Genet.* **17**, e1009406 (2021).

87. E. Becht, L. McInnes, J. Healy, C.-A. Dutertre, I. W. H. Kwok, L. G. Ng, F. Ginhoux, E. W. Newell, Dimensionality reduction for visualizing single-cell data using UMAP. *Nat. Biotechnol.* **37**, 38–44 (2019).

88. C. S. McGinnis, L. M. Murrow, Z. J. Gartner, DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst.* **8**, 329-337.e4 (2019).

89. T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W. M. Mauck, Y. Hao, M. Stoeckius, P. Smibert, R. Satija, Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21 (2019).

90. D. Aran, A. P. Looney, L. Liu, E. Wu, V. Fong, A. Hsu, S. Chak, R. P. Naikawadi, P. J. Wolters, A. R. Abate, A. J. Butte, M. Bhattacharya, Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nat. Immunol.* **20**, 163–172 (2019).

91. Y. Liao, J. Wang, E. J. Jaehnig, Z. Shi, B. Zhang, WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs. *Nucleic Acids Res.* **47**, W199–W205 (2019).

92. K. Blighe, S. Rana, M. Lewis, EnhancedVolcano: publication-ready volcano plots with enhanced colouring and labeling, (2018); https://bioconductor.org/packages/release/bioc/vignettes/EnhancedVolcano/inst/doc/EnhancedVolcano.html.

857   93.   A. Lex, N. Gehlenborg, H. Strobelt, R. Vuillemot, H. Pfister, UpSet: Visualization of Intersecting
858   Sets. *IEEE Trans. Vis. Comput. Graph.* **20**, 1983–1992 (2014).

859   94.   G. Csardi, T. Nepusz, The igraph software package for complex network research. *InterJournal*
860   **Complex Systems**, 1695 (2006).

861