1 **Temporal Dynamics of Genetically Heterogeneous Extended-Spectrum Cephalosporin**

2 **Resistant *Escherichia coli* Bloodstream Infections**

3 **Running Title:** Molecular Epidemiology of ESC-R *E. coli* BSIs in Hematological Patients

4 William C Shropshire[a], Benjamin Strope[b], Selvalakshmi Selvaraj Anand[b], Jordan Bremer[a],

5 Patrick McDaneld[c*], Micah M Bhatti[d], Anthony R Flores[e], Awdhesh Kalia[b], Samuel A

6 Shelburne[a,f#]

7 [a]Department of Infectious Diseases and Infection Control, The University of Texas MD

8 Anderson Cancer Center, Houston, TX, USA

9 [b]Program in Diagnostic Genetics and Genomics, MD Anderson Cancer Center School of Health

10 Professions, Houston, TX, USA

11 [c]Division of Pharmacy, The University of Texas MD Anderson Cancer Center, Houston, TX,

12 USA

13 [d]Department of Laboratory Medicine, The University of Texas MD Anderson Cancer Center,

14 Houston, TX, USA

15 [e]Division of Infectious Diseases, Department of Pediatrics, McGovern Medical School,

16 University of Texas Health Science Center, Houston, TX, USA

17 [f]Department of Genomic Medicine, The University of Texas MD Anderson Cancer Center,

18 Houston, TX, USA

19 *Patrick McDaneld is currently an employee of bioMérieux Diagnostics but was not an employee

20 at the time of his contributions to the study.

21   #Address for correspondence: Samuel A Shelburne, MD, PhD, 1901 East Rd., South Campus

22   Research Building, 4SCR5.1042, Houston, TX 77054; phone: (713) 792-3629, e-mail:

23   sshelburne@mdanderson.org

ABSTRACT

Extended-spectrum cephalosporin resistant *Escherichia coli* (ESC-R-*Ec*) is an urgent public

health threat with sequence type clonal complex 131 (STc131), phylogroup B2 strains being

particularly concerning as the dominant cause of ESC-R-*Ec* infections. To address the paucity

of recent ESC-R-*Ec* molecular epidemiology data in the United States, we used whole genome

sequencing (WGS) to fully characterize a large cohort of invasive ESC-R-*Ec* at a tertiary care

cancer center in Houston, Texas collected from 2016-2020. During the study timeframe, there

were 1154 index *E. coli* bloodstream infections (BSIs) of which 389 (33.7%) were ESC-R-*Ec*.

Using time series analyses, we identified a temporal dynamic of ESC-R-*Ec* distinct from ESC-

susceptible *E. coli* (ESC-S-*Ec*), with cases peaking in the last six months of the calendar year.

WGS of 297 ESC-R-*Ec* strains revealed that while STc131 strains accounted for ~45% of total

BSIs, the proportion of STc131 strains remained stable across the study time frame with

infection peaks driven by genetically heterogeneous ESC-R-*Ec* clonal complexes. Bla_{CTX-M}

variants accounted for most β-lactamases conferring the ESC-R phenotype (89%; 220/248

index ESC-R-*Ec*), and amplification of bla_{CTX-M} genes was widely detected in ESC-R-*Ec* strains,

particularly in carbapenem non-susceptible, recurrent BSI strains. $Bla_{CTX-M-55}$ was significantly

enriched within phylogroup A strains, and we identified $bla_{CTX-M-55}$ plasmid-to-chromosome

transmission occurring across non-B2 strains. Our data provide important information regarding

the current molecular epidemiology of invasive ESC-R-*Ec* infections at a large tertiary care

cancer center and provide novel insights into the genetic basis of observed temporal variability

for these clinically important pathogens.

IMPORTANCE

Given that *E. coli* is the leading cause of worldwide ESC-R *Enterobacterales* infections, we

sought to assess the current molecular epidemiology of ESC-R-*Ec* using a WGS analysis of

many BSIs over a five-year period. We identified fluctuating temporal dynamics of ESC-R-*Ec*

infections, which has also recently been identified in other geographical regions such as Israel.

Our WGS data allowed us to visualize the stable nature of STc131 over the study period and

demonstrate a limited, but genetically diverse group of ESC-R-*Ec* clonal complexes are

detected during infection peaks. Additionally, we provide a widespread assessment of β-

lactamase gene copy number in ESC-R-*Ec* infections and delineate mechanisms by which such

amplifications are achieved in a diverse array of ESC-R-*Ec* strains. These data suggest that

serious ESC-R-*Ec* infections are driven by a diverse array of strains in our cohort and impacted

by environmental factors suggesting that community-based monitoring could inform novel

preventative measures.

INTRODUCTION

Extended-spectrum β-lactamase (ESBL) producing *Enterobacterales* (ESBL-E) were ranked as a serious antimicrobial resistance (AMR) threat by the Centers for Disease Control and Prevention in 2019 (1). A recent study of multi-drug resistant (MDR) infections in hospitalized patients indicated a 53.3% increased incidence rate in estimated ESBL-E cases from 2012 to 2017 in the United States (US) (2). Other sources of US data have indicated similar trends, such as a 138% increase in resistance to 3rd generation cephalosporins amongst invasive *E. coli* strains from 2009 to 2016 (3, 4). Furthermore, multiple population-based studies indicate that ESBL-E bloodstream infections may have a higher likelihood of adverse outcomes compared to non-ESBL-E bloodstream infections (5-7).

The key impact of ESBL enzymes is 3rd generation cephalosporin resistance (*i.e.,* extended-spectrum cephalosporin resistance; ESC-R). Although ESBL enzymes account for the vast majority of the associated ESC-R phenotype, there are other mechanisms in *Enterobacterales*, such as plasmid-borne AmpC enzymes (8). Therefore, broadening our scope of potential ESC-R mechanisms provides a more holistic approach to understanding transmission dynamics of these ESC-R genes (9). Amongst ESC-R *Enterobacterales*, some 50% or greater of infections are due to *Escherichia coli* (10, 11). ESC-R *E. coli* (ESC-R-*Ec*) in the 1980s and 1990s primarily harbored ESBL variants of TEM and SHV enzymes (12); however, this changed in the early 2000s with the increasing spread of CTX-M enzymes (13). The rise of CTX-M enzyme producing *E. coli* coincided with a marked increase in sequence type (ST) 131 strains (14). In particular, fluoroquinolone resistant sub-populations of ST131 (*i.e.,* ST131-*H*30 subclade C) strongly associated with *bla*$_{CTX-M-15}$ and *bla*$_{CTX-M-27}$ carriage, have been shown to be the predominant cause of ESC-R-*Ec* infections in the US (15). However, whether the increasing number of ESC-R-*Ec* infections in the US is driven by further expansion of ST131 strains is not well understood.

Application of WGS to large cohorts of temporally diverse *E. coli*, including ESBL producing

isolates, from local and nationwide surveys has revealed common as well as unique molecular

epidemiology features (16-23). Consistently, *E. coli* strains causing human infections are pauci-

clonal with domination of a small number of STs relative to the overall population (16-18, 24).

Many analyses indicate ST131 rose to become the leading *E. coli* ST isolated from humans by

2010 (19, 25). However, studies in England and Calgary have found that ST131 has remained

at a relatively steady proportion of overall *E. coli* infections over the past five to ten years (16,

19, 22) whereas investigations in France, Spain, and Norway have identified a continued

increase in emergent ST131 subclades (17, 20, 21). Geographical variation in relative ST131

proportion changes may reflect differential ST131 cladal adaptation within ecological niches,

with advantageous traits failing to fixate in populations due to negative frequency-dependent

selection, reflected by the eventual stability of ST131 within the broader *E. coli* population (26).

A comprehensive US study found that ST131 accounted for 50% of all ESC-R-*Ec* strains in

2012 with an estimated increase of 50% from 2007 to 2011 in the percentage of ST131 *E. coli*

(27). Given the lack of recent longitudinal data analyzing ESC-R-*Ec* in the US, we sought to

gain insight into the molecular epidemiology of ESC-R-*Ec* by analyzing bloodstream infection

causing strains at our institution from 2016-2020.

RESULTS

Observed peaks in ESC-R-*Ec* BSIs occurred temporally within our sampling frame.

We identified total *Escherichia coli* bloodstream infections (*Ec*-BSI) at our institution that occurred from March 1st, 2016, to December 31st, 2020. There were 1370 total *Ec*-BSIs with 84.2% (1154/1370) being index *Ec*-BSI cases and 15.8% (216/1370) being recurrent (see methods for definitions). The overall ESC-R-*Ec* proportion was 33.7% (389/1154) among index *Ec*-BSI cases. The proportion of ESC-R-*Ec* infections that were recurrent (121/510, 23.7%) was significantly higher compared to recurrent extended-spectrum cephalosporin susceptible *E. coli* (ESC-S-*Ec*) cases (95/860, 11%, χ^2 *P*-value < 0.001).

The non-normalized temporal trends of ESC-R-*Ec* within this 4+ year timeframe is presented in Table S1. Given that changes in absolute yearly ESC-R-*Ec* might be reflective of variance in the total population seen at our institution, we utilized monthly admissions as a surrogate of the hospital population to calculate prevalence rates. We found no statistically significant linear increase in ESC-R-*Ec* rates over the course of the study (linear regression one-sample t-test *P*-value=0.20). Notably, when calculating *Ec*-BSI prevalence rates stratified by ESC status, we identified an oscillating pattern of total *Ec*-BSI prevalence rates corresponding to peaks and valleys (dotted grey loess curve; **Figure 1**) that primarily correlated with ESC-R-*Ec* prevalence rates (solid red loess curve; **Figure 1**) rather than ESC-S-*Ec* prevalence rates (solid blue curve, **Figure 1**). Stratifying the sampling frame based on these observed increased infection peaks (**Figure 1; Table 1**), ESC-R-*Ec* prevalence rates were greatest during window 2 (1.89 BSI cases/1000 admissions), window 4 (2.09 BSI cases/1000 admissions), and window 6 (2.24BSI cases/1000 admissions). The mean peak ESC-R-*Ec* prevalence rate (2.07 BSI cases/1000 admissions) was 60% greater than the average of 1.29 BSI cases/1000 admissions during non-peak periods (P < 0.001 for monthly ESC-R-*Ec* rates during peak vs. non-peak windows by Student's t-test).

We performed time series analyses and found that while both ESC-R-*Ec* and ESC-S-*Ec* had

non-significant monotonic trends, there was evidence of a non-monotonic trend for ESC-R-*Ec*

that occurred up to the start of the COVID-19 pandemic in March 2020 (WAVK test-statistic

=2.7; *P*-value = 0.016), which was not evident in ESC-S-*Ec* (WAVK test-statistic =0.2; *P*-value =

0.877). To determine whether ESC-R-*Ec* and ESC-S-*Ec* prevalence rates were distinct, we

compared pairwise month-to-month prevalence and found no statistically significant correlation

(R=0.14, Pearson *P*-value = 0.31; Fig. S1A) suggesting independent temporal trends. Given the

periodic trends noted in **Figure 1,** we assessed whether variation in *Ec*-BSIs corresponded to

the time of year by dividing the index infections into the 1st or 2nd half of the year. We found a

significantly higher average of ESC-R-*Ec* (7.9 $\pm$ 0.5 vs. 5.5 $\pm$ 0.6 BSIs/month; *P*-value = 0.003)

but not ESC-S-*Ec* (13.8 $\pm$ 0.5 vs. 12.5 $\pm$ 0.6; *P*-value =0.09) occurring in the last 6 months

relative to the initial 6 months of each year (Fig. S1B) respectively. Taken together, these data

trends suggest temporal dynamics in ESC-R-*Ec* independent of ESC-S-*Ec* strains with a

potential seasonal component.

Peaks in ESC-R *Ec*-BSIs are composed of genetically heterogeneous strains from

multiple sequence type complexes.

From the 389 index and 121 recurrent ESC-R-*Ec* strains identified during the study period, we

generated assemblies for 297 ESC-R-*Ec* isolates including 64% (248/389) of index and 41%

(49/121) of recurrent ESC-R-*Ec* strains. The quality of assemblies is shown in Table S2. In total

there were 7 phylogroups detected across the full sampling frame (**Table 2**). The majority of

index ESC-R-*Ec* belonged to phylogroup B2 (50.8%; 126/248), followed by A (19.8%; 49/248),

D (12.1%; 30/248), F (7.7%; 19/248), B1 (6.5%; 16/248), C (2.8%; 7/248), and G (0.4%; 1/248).

We identified a similar distribution for recurrent ESC-R-*Ec* strains with B2 (38.8%; 19/49) as the

most common phylogroup followed by A (34.7%; 17/49), D (12.2%; 6/49), B1 (8.2%; 4/49), C

(4.1%; 2/49), and F (2.0%; 1/49). In terms of sequence types (STs) (**Table 2**), ST131 was the

most common for both index (n = 102, 41%) and recurrent strains (n = 17, 35%). A further

breakdown of each of the most common STs detected by phylogroup can be found in **Table 2**.

To investigate the molecular epidemiology underlying the temporal dynamics of our ESC-R-*Ec*

population, we analyzed isolates characterized by phylogroup across infection windows (**Figure**

2). **Figure 2A-B** provides an illustration of the phylogroup distribution throughout the infection

windows which remained remarkably stable over time with no statistically significant trends

(Fisher's exact test simulated *P*-value= 0.95). There were sporadic increases in the absolute

numbers of various phylogroups during the peak infection periods such as phylogroup A during

windows 2 and 6, as well as phylogroup B1 and phylogroup F during window 4. We collapsed

the "non-peak" (*i.e.,* windows 1, 3, 5, and 7) and "peak" (*i.e.,* windows 2, 4, and 6) periods

together and again found no statistically significant differences in phylogroup proportions

between peak and non-peak periods (Fisher's exact test *P*-value= 0.42). The dominant B2

phylogroup accounted for 58% of infections during the non-peak windows and 47% during the

peak windows (Fisher's exact test *P*-value= 0.15).

We grouped index strains by sequence type clonal complexes (*i.e.,* single locus ST variants

within a 7-housekeeping gene schema) to further analyze temporal trends. The leading

sequence type complexes (STc) were STc131 (n = 109, 44%), STc10 (n = 35, 14%), STc405 (n

= 16, 7%), STc14 (n = 14, 6%), STc648 (n = 14 (6%), and STc38 (n = 10, 4%). For testing

purposes, we collapsed STcs with less than 5 strains into an "uncommon" category (n = 32,

13%). Similar to our phylogroup analysis, we did not identify any significant trends for a

particular STc over the study period (Fisher's exact test simulated *P*-value= 0.91). However,

there were temporary absolute increases in specific STc infections within peak infection

windows such as "uncommon" STcs within window 2, STc14 and STc648 within window 4, and

STc10 within window 6 (**Figure 2C-D**). Importantly, STc131 remained stable in prevalence

across each of the infection windows (χ^2 P-value = 0.9) with STc131 strains accounting for 50%

of infections during non-peak windows compared to 40% of infections during peak infection

windows (Fisher's exact test P-value= 0.15). Temporal information regarding isolates

sequenced per window including the proportion of isolates that were STc131 is provided in

Table S3. There were no peak infection periods in which STc131 strains accounted for a higher

percentage of infections relative to non-peak periods (Table S3). Similarly, there were no

statistically significant proportion differences across windows in ST1193, the other most

commonly prevalent B2 isolates (Fisher's exact test P-value= 0.48) nor in the second most

common, STc10 (Fisher's exact test P-value= 0.74). Thus, these data indicate that peaks of

ESC-R-*Ec* observed temporally over our study period were driven by a diverse combination of

ESC-R-*Ec* isolates rather than expansion of STc131 strains.

Bla_{CTX-M} variants are dominant encoding genes conferring inferred ESC-R phenotype.

We searched the WGS data for genes potentially conferring the ESC-R phenotype, which we

defined as those encoding ESBLs, *i.e.*, TEM, SHV, CTX-M encoding genes with a 2be Jacoby-

Bush classification (12), as well as CMY and carbapenemase encoding genes. Of the 248 index

isolates, 236 (95%) had at least one gene expected to confer the ESC-R phenotype with CTX-M

variants being present in 220 strains (89%) and CMY variants found in 21 strains (8.4%) (*N.b.,*

strains could contain multiple ESC-R encoding genes). A single ST131 strain carried an ESBL

bla_{TEM-19}, two strains carried ESBL bla_{SHV-12}, one ST361 strain carried $bla_{OXA-232}$, and three

STc10 strains harbored bla_{NDM-5}. The most common CTX-M encoding gene variants detected

within index ESC-R-*Ec* isolates were $bla_{CTX-M-15}$ (n=134), $bla_{CTX-M-27}$ (n=29), $bla_{CTX-M-55}$ (n=26),

$bla_{CTX-M-14}$ (n=19), and multiple bla_{CTX-M} variant carriage (n=7). A similar distribution was

observed for recurrent ESC-R-*Ec* isolates: $bla_{CTX-M-15}$ (n=26), $bla_{CTX-M-27}$ (n=7), $bla_{CTX-M-55}$ (n=8),

and $bla_{CTX-M-14}$ (n=3). Of the 12 index strains that did not carry a known ESC-R phenotype

determinant, six contained bla_{OXA-1}, five encoded non-ESBL TEM enzymes, and only one isolate

lacked a known exogenous β-lactamase encoding gene. The preponderance of CTX-M with a

smaller role of CMY in conferring the ESC-R phenotype in *E. coli* is consistent with previously

published results in the US (9, 10).

Temporal whole genome analysis indicates low level of clonality amongst ESC-R-*Ec.*

To gain further insights into the molecular epidemiology driving the temporal dynamics of ESC-

R-*Ec*, we used WGS data to analyze the population structure for all 297 isolates. The core gene

inferred maximum likelihood phylogeny based on 2,695 core genes (*i.e.,* genes present in ≥99%

of the 297 ESC-R *Ec*-BSI Isolates) is shown in **Figure 3**. Consistent with previous WGS studies

of ESC-R *E. coli* population structure (9, 18), we observed significant variation in the degree of

genetic similarity amongst strains of distinct STcs. For example, there was only an average of

28 core gene SNPs separating a given STc14 strain from its nearest neighbor compared to

1518 for STc10 strains. Similarly, we identified several previously described subclades within

the broader STc131 population (15, 20, 28). By overlying infection window with our WGS

phylogeny, we sought to determine whether the increased resolution would facilitate detection of

highly related strains driving infection peaks. However, in line with our phylogroup and STc-

based analyses, we did not observe large numbers of phylogenetically clustered index strains

occurring during infection windows (**Figure 3**). Consistent with the notion of similar genetic

diversity being present during peak and non-peak infection windows, there was no difference in

pairwise core gene SNP distance between one isolate and its most closely related strain during

the various windows (Kruskal-Wallis test *P*-Value=0.81; Fig S2).

To search for clusters of closely genetically related strains both within peak infection windows

and throughout the study period, we measured the pairwise SNP distances of index isolates

using recombination free core genome alignments from the 12 most commonly observed STs in

our cohort, which make up 79.8% of the total population (198/248). We identified seven groups

of two or more isolates (*i.e.*, genetic clusters) from unique patients with <10 core genome

pairwise SNPs (Fig S3). The maximum number of closely related isolates within a peak infection

window was two strains whereas the maximum cluster size was three strains, with duration

between isolates from the same cluster ranging from 5 days to 777 days (Fig. S3). These data

indicate that even at the WGS level, peaks of ESC-R-*Ec* are driven by genetically heterogenous

isolates with few signals indicating clonal outbreaks across each infection window.

ESC-R encoding genes are highly enriched within *Ec*-BSI sub-populations.

We combined the β-lactamase encoding gene data with our WGS based phylogeny to assess

the relationship between *E. coli* genetic content and ESC-R mechanisms (**Figure 3**). There was

a strong association of phylogroup B2 isolates harboring $bla_{CTX-M-15}$ genes as 60.4% of (81/134)

$bla_{CTX-M-15}$ harboring strains were present in this phylogroup (row-wise Fisher's exact test

adjusted *P*-value = 0.01). Specifically, this association was driven by the presence of $bla_{CTX-M-15}$

in 72 STc131 isolates (adj. *P*-value = 0.009), the vast majority of which are part of the previously

identified $bla_{CTX-M-15}$ carrying *H*30Rx lineage (15). Similarly, there was also enrichment of the

second most commonly detected CTX-M encoding gene variant, $bla_{CTX-M-27}$, in B2 index isolates

(26/29 $bla_{CTX-M-27}$ harboring strains; adj. *P*-value < 0.0001) again driven by STc131 isolate

carriage (23/29; Adj. *P*-value = 0.0005). Specifically, there were 19/23 $bla_{CTX-M-27}$ positive ESC-

R-*Ec* belonging to the ST131-C1-M27 clade, which has a strong association with $bla_{CTX-M-27}$

carriage (28) as seen in the lower right quadrant of the core gene phylogeny (**Figure 3**). The

third most prevalent CTX-M encoding gene variant, $bla_{CTX-M-55}$, was enriched within phylogroup

A (12 phylogroup A strains out of 26 $bla_{CTX-M-55}$ containing index isolates; adj. *P*-value = 0.004)

whereas it was only present in one B2 isolate (1/26, Adj. *P*-value < 0.0001). The final major

*bla*CTX-M variant in our cohort, *bla*CTX-M-14 was carried by a diverse array of genetically

heterogenous strains (**Figure 3**). In contrast to the *bla*CTX-M encoding genes in which there were

several large clusters of genetically adjacent strains encoding the same CTX-M variant, carriage

of CMY encoding genes was sporadically observed across phylogroups. Similar to the

phylogroup trend and phylogeny analyses, we did not observe any statistically significant

association between ESC-R mediating enzyme variants and infection windows (χ^2 adj. *P*-value

> 0.05).

Accessory genomes are distinct between B2 and non-B2 phylogroups.

We further explored genomic differences within ESC-R-*Ec* by comparing accessory genome

content (*i.e.,* defined as genes carried in >5% and <95% of the total ESC-R-*Ec* population)

across the full cohort. There was a total of 25,631 genes in the pan genome with 6,060

accessory genes. Through a cluster analysis using the Jaccard distance of the accessory

genome between samples, we observed a strong association between phylogroup, clonal

complex, and the accessory genome (**Figure 4**)**.** A neighbor joining (NJ) tree constructed from a

Jaccard distance matrix shows a clear delineation of B2 and non-B2 isolates that split at the root

of the tree (**Figure 4**). Using a network graph analysis of the accessory genome (see methods)

(Fig. S4), there was separation of B2 and non-B2 isolates into ten clusters highly correlating to

STc (**Figure 4**). Indeed, each of the clonal complexes with ≥5 isolates had at least one

statistically significant association (Adj. *P*-value <0.05) with an accessory genome Markov

Cluster (MCL) group (Fig. S5). Furthermore, the MCL clustering algorithm delineated the sub-

clade structure of STc131 where each of the ST131-*H*30Rx subclades C1 (Cluster 2) and C2

(Cluster 3) were separated by accessory genome (**Figure 4**). ST131-*H*41-O16 subclade A

(Cluster 9) was found exclusively in one MCL group (**Figure 4**) demonstrating unique accessory

genome between ST131 C1/C2 and A subclades. A complimentary principal component

analysis demonstrated clustering of accessory content by phylogroup (Fig. S6A), STc (Fig.

S6B), MCL group (Fig. S6C), and ESBL gene (Fig. S6D) further highlighting the separation of

B2 and non-B2 groups by accessory genome content. The emergent ST1193 lineage, which

made up all STc14 (n=15) strains, made up a separate MCL cluster (Cluster 8) within B2

isolates, however, there were no clear associations with a particular bla_{CTX-M} variant. These

analyses indicate potential differential pathways to ESC-R development dependent on

phylogroup background.

Significant copy number amplifications of enriched CTX-M encoding genes occur within

phylogroups

Given the growing appreciation of β-lactamase gene amplification in contributing to AMR

phenotypes (29-31), we sought to characterize gene copy number variation of ESC-R genes to

determine whether ESC-R gene amplification correlated with our previously identified gene

enrichment associations. **Table 3** provides copy number estimates of bla_{CTX-M} variants stratified

by the three most common phylogroups. Tables S4 and S5 provide copy numbers of other β-

lactamase encoding genes and overall estimated copy numbers of AMR genes respectively. For

the 268 bla_{CTX-M} genes, the median copy number variant (CNV) was 1.2✗ (Interquartile Range

[IQR] = 1.3) with a range of 0.4 to 12.9. The median CNV across CTX-M variants was similar

with $bla_{CTX-M-15}$ (n=161, CNV=1.1✗; IQR =1.2), $bla_{CTX-M-27}$ (n=37, CNV=1.4✗; IQR =0.8), $bla_{CTX-M-55}$

(n=37, CNV=1.5✗; IQR =1.4), and $bla_{CTX-M-14}$ (n=37, CNV=1.4✗; IQR =2.2) having statistically

significant greater than 1 copy (one-sample Wilcoxon signed rank adj. *P*-value <0.01).

We and others have previously identified that copy number increases in narrow-spectrum and

extended-spectrum β-lactamase encoding genes contribute to the development of carbapenem

resistance (32, 33). Therefore, we stratified our ESC-R-*Ec* population by carbapenem

susceptibility (*i.e.,* carbapenem non-susceptible vs. susceptible) to determine if there were

statistically significant amplifications in carbapenem-susceptible ESC-R-*Ec* isolates. The

carbapenem non-susceptible ESC-R-*Ec* had greater estimated copy numbers of bla_{CTX-M}

(median CNV = 2.7✗; IQR = 4.2) compared to carbapenem susceptible ESC-R-*Ec* (median CNV

= 1.1✗; IQR = 1.0; two-sample Wilcoxon rank sum test *P*-value = 3e-7). Thus, we further

characterized bla_{CTX-M} gene amplification across phylogroup solely within ESC-R-*Ec* that are

carbapenem susceptible (**Figure 5A).** We found statistically significant amplifications for $bla_{CTX-M-15}$

and $bla_{CTX-M-27}$ in phylogroup B2 isolates and $bla_{CTX-M-55}$ in phylogroup A isolates, which

aligned with the enrichment analysis. These statistically significant trends persisted specifically

within STc131 for $bla_{CTX-M-15}$/$bla_{CTX-M-27}$ and STc10 for $bla_{CTX-M-55}$ (Fig. S7). To assess whether

bla_{CTX-M} amplifications occurred as an adaptive response, we sought to identify isolates from the

same patient that were the same clonal complex and had the same bla_{CTX-M} variants, *i.e.,* were

likely clonal, recurrent infections (**Figure 5B**). We identified 16 pairs from 14 patients in which

56% (9/16) second ESC-R *Ec*-BSI isolates were carbapenem non-susceptible following their

antecedent carbapenem susceptible isolate. We found that there was a statistically significant

increase (Wilcoxon Signed Rank Test *P*-value=0.03) in bla_{CTX-M} copy numbers from the first

isolate (median CNV=1.02; IQR=0.86) to the second isolate (median CNV=1.80; IQR=3.32).

Notably, these bla_{CTX-M} amplifications occurring within sequential isolates were observed for

$bla_{CTX-M-15}$/$bla_{CTX-M-27}$ and $bla_{CTX-M-55}$ in STc131 and STc10 respectively (**Figure 5B**). To gain

insights into how β-lactamase gene amplification affected AMR phenotypes, we characterized

the minimum inhibitory concentration (MIC) distribution for recurrent vs. non-recurrent (*i.e.,* first

isolate) populations for piperacillin-tazobactam (TZP), ceftazidime (CAZ), cefepime (FEP),

ertapenem (ETP), and meropenem (MEM) antimicrobial susceptibility testing (Fig. S8A-D). For

all tested β-lactams, the MIC90 was higher for the recurrent relative to the non-recurrent isolates

(Fig. S8). Thus, there was a general shift in MIC distributions which correlated with recurrent

isolates that had increased bla_{CTX-M} gene copy numbers.

F-type plasmids are primary vectors of $bla_{CTX-M-55}$ in non-STc131 strains.

Previous studies have focused largely on ST131 subclade associations with $bla_{CTX-M-15}$ and

$bla_{CTX-M-27}$ (15, 28) including our characterization of $bla_{CTX-M-15}$ and bla_{OXA-1} co-carriage on similar,

multireplicon F-type plasmids contributing to carbapenem resistance (32, 33). Given the rarity of

$bla_{CTX-M-55}$ molecular epidemiology data from North America (34), we focused on $bla_{CTX-M-55}$

inasmuch as it was highly prevalent throughout our sampling frame. We used a long-read

approach to complete assemblies of nine $bla_{CTX-M-55}$ positive strains and found that most (n=7)

carried $bla_{CTX-M-55}$ in a highly conserved IS*Ecp1* context on ~100kb IncFIB/IncFIC plasmids

(**Figure 6A**). Whereas the cargo gene region was relatively variable, the F-type plasmid regions

responsible for transfer, replication, and stability were highly conserved (blastn ID/coverage

≥99%). For the full cohort of $bla_{CTX-M-55}$ isolates (n=38), 31 isolates had IncFIB/IncFIC amplicon

hits from PlasmidFinder (>80% ID;>80% COV). Short-read pileup analysis to pMB8960_1 (*i.e.,*

the most conserved IncFIB/IncFIC $bla_{CTX-M-55}$ positive isolate), showed that 30/31 isolates had

>80% breadth of coverage and >100× coverage depth for pMB8960_1.

Our long read analyses allowed for identification of $bla_{CTX-M-55}$ amplification arising from plasmid

copy number variation (MB8960 and MB9029) as well as plasmid-to-chromosome transfer (*i.e.,*

segmental duplication) from two isolates (MB6206 and MB7355) as evidenced by tracking target

site duplications (TSDs) and plasmid content from IS*Ecp1-*$bla_{CTX-M-55}$ (**Figure 6B**). This included

an isolate in which IS*Ecp1-*$bla_{CTX-M-55}$ disrupts the *envZ* gene, which is part of a two-component

regulatory system (*i.e.,* EnvZ-OmpR) that regulates expression of outer membrane porin genes

(35). A carbapenem susceptible ESC-R-*Ec* isolate (MB7355) was identified with two

independent insertions within the chromosome likely originating from the IncFIB/IncFIC plasmid

as evidenced through TSD tracking (**Figure 6B;** one insertion into *fhu*A gene demonstrated).

Another vector of $bla_{CTX-M-55}$ was identified in two ST167 isolates in a substantially differing

context with $bla_{CTX-M-55}$ upstream of IS*26*-v1 on smaller IncX1 plasmids (**Figure 6B**). The IS*26*-v1

upstream of $bla_{CTX-M-55}$ had truncated IS*Ecp1* (**Figure 6B;** purple striped CDS) and were within

unique antimicrobial resistance gene transposons. These analyses together show that $bla_{CTX-M-55}$

is a major contributor to the ESC-R-*Ec* phenotype in our cohort with predominantly

IncFIB/IncFIC carriage.

DISCUSSION

The past twenty years in the US have seen a consistent rise in ESC-R-*Ec* causing infection,

primarily due to ESBL enzymes (2-4). While the high prevalence of ST131 observed in the US is

concerning, it is not clear if these pathogens are responsible for the increased frequency of

ESC-R-*Ec* (2). We sought to leverage our longitudinal collection of bloodstream isolates to

address the recent temporal dynamics of *E. coli* at our tertiary care cancer center. Although we

observed that STc131 strains accounted for nearly 50% of infecting isolates, the STc131

prevalence rates remained fairly stable over time with multiple MDR clones accounting for

peaks in infection rates observed during our study period.

We chose to focus on BSIs because of the clear distinction between colonization and infection

relative to other potential sample sites. Our large number of fully sequenced isolates and

extended study timeframe provide critical information regarding contemporary molecular

epidemiology of invasive ESC-R-*Ec* infections in a major US city. Although our finding of ~35%

Ec-BSIs being ESC-R is high relative to other studies from the US (4), it is consistent with the

~40% proportion of ESC-R in *Ec*-BSI occurring in solid-organ transplant recipients (36).

Whereas recent longitudinal ESC-R-*Ec* epidemiological studies from the US are lacking, we can

compare our findings to those recently published from Europe (16-21). A similar theme has

emerged from these investigations where for several years following introduction into the

population, STc131 strains dramatically increase before becoming a stable proportion of overall

E. coli infections (16-21). Given that STc131 had been identified as the dominant antimicrobial

resistant *E. coli* STc in the US for at least 10 years prior to the start of our sampling (27, 37), our

finding of a relatively stable contribution of STc131 is consistent with other studies (16-21) and

suggest that, at least for our population, STc131 may have reached a stable level in the broader

ESC-R-*Ec* population.

A key strength of our study was our ability to investigate the molecular epidemiology underlying

the temporal variability of ESC-R-*Ec* prevalence rates. We found that ESC-R-*Ec* rates generally

peaked within the last six months of the calendar year and were independent of ESC-S-*Ec*

rates. Seasonality of both ESC-R-*Ec* infection and colonization have been previously reported

with generally higher rates identified during warmer months (38-40). For example, a recent

pooled observational cohort analysis of ESBL positive *E. coli* and *K. pneumoniae* (ESBL-E/K)

carriage in the Netherlands found that the highest prevalence occurred in August (OR 1.88,

95% CI 1.02-3.49) and September (OR 2.25, 95% CI 1.30-3.89) compared to January (39). In

Houston, the highest temperatures of the year are generally observed between June and

September (https://www.weather.gov/hgx/climate_iah_normals_summary), which might facilitate

proliferation of *E. coli* strains leading to colonization and subsequent infection at a delayed

interval. Consistent with environmental factors being a major driver of the temporal variability,

we observed low genetic relatedness amongst *E. coli* strains during the peak windows of

infection rather than closely related clones. On a few occasions, we identified strains from

distinct patients that were closely related (*i.e.,* < 10 SNPs) consistent with hospital acquisition,

but most strains were genetically diverse suggesting a community source. Given the recent

success of wastewater surveillance in identifying SARS-CoV-2 and influenza transmission

dynamics (41, 42), we envision that a similar approach could be used to improve understanding

of how community prevalence of ESC-R pathogens impacts subsequent infections.

In previous works we have identified that non-carbapenemase producing, carbapenem resistant

Enterobacterales (non-CP-CRE) have increased copy numbers of genes encoding ESC-R

mediating enzymes such as bla_{CTX-M} variants (32, 33). This study expands on our previous copy-

number assessment of β-lactamase encoding genes mediating the ESC-R phenotype in

carbapenem-susceptible *E. coli* strains. This allowed us to find that carbapenem-susceptible

ESC-R-*Ec* strains have lower bla_{CTX-M} copy numbers relative to carbapenem-resistant ESC-R-*Ec*

strains consistent with an important role of bla_{CTX-M} amplification in the transition from ESC-R to

the non-CP-CRE phenotype. Moreover, via an analysis of paired isolates from the same

patients, we found significant bla_{CTX-M} amplifications suggesting that such amplifications are

likely an adaptive mechanism to antimicrobial exposure in the host as has been observed by our

group and others in the laboratory environment (32, 33, 43). Together with recently published

data from ESBL-positive *E. coli* strains isolated in St. Louis (34), our long-read analyses

demonstrate the diverse mechanisms by which *E. coli* can increase bla_{CTX-M} encoding gene copy

numbers including augmented plasmid copy numbers, multiple chromosomal copies in distinct

locations, and *in situ* amplifications mediated by insertion sequences such as IS*Ecp1* and IS*26*.

Although gene amplifications have long been thought to confer a major fitness cost and thus be

transient in response to antimicrobials, incorporation of bla_{CTX-M} encoding genes into particular

chromosomal locations has been found to have a low fitness cost, allowing for stable

amplifications across many generations in the absence of antimicrobial pressure (44, 45). We

hypothesize that the high rates of recurrence observed amongst ESC-R-*Ec* relative to ESC-S-

Ec strains might, in part, be due to bla_{CTX-M} amplifications occurring amongst colonizing strains in

the gastrointestinal tract which would allow for their persistence during carbapenem treatment

and subsequent re-infection. Such an amplification of $bla_{CTX-M-14}$ was recently described in an *E.*

coli strain recurrently isolated from a patient's urine over many months (34).

Furthermore, we were able to identify IncFIB/IncFIC plasmids as the primary vector of IS*Ecp1*-

$bla_{CTX-M-55}$ carriage in our cohort where amplification was occurring through both increase in

plasmid copy numbers as well as plasmid-to-chromosome horizontal gene transfer. Although we

did not identify a 'clonal expansion' of chromosomal IS*Ecp1*-$bla_{CTX-M-55}$, the independent transfer

of these vectors into two different phylogroup chromosomal backgrounds suggest the potential

for stable chromosomal propagation. A recent study in China found a temporal trend with

increasing chromosomal $bla_{CTX-M-55}$ albeit without clarity to whether this was occurring within

clonal populations (46). While little is known regarding $bla_{CTX-M-55}$ prevalence in North America

(34), the enrichment of this bla_{CTX-M} variant in the second most commonly detected

phylogroup/STc in our cohort suggest these IncFIB/IncFIC::IS*Ecp1*-$bla_{CTX-M-55}$ plasmids are a

significant contributor to ESC-R-*Ec* in our region.

Although our study has many strengths, there are important limitations. First, this was a single-

center investigation at a hospital serving an immunocompromised population and thus external

generalizability is unclear. This weakness is somewhat mitigated by our center serving a

geographically dispersed group of patients and because the impact of ESC-R-*Ec* is particularly

severe amongst immunocompromised persons. Moreover, our finding of temporal variability

amongst *E. coli* infections has been observed amongst broader populations such as a recent

nationwide report from Israel (47). Second, the diverse nature of observed clonal complexes

and subclones meant that there were often small numbers of non-STc131 clonal complexes

during specific infection windows. Larger sampling from a more geographically dispersed area is

needed to address these issues. A final weakness is that the beginning of the COVID epidemic

likely led to a major disruption in infection epidemiology at our institution given the nearly 50%

decrease in admissions observed in the spring and summer of 2020. Whereas both the absolute

numbers of and proportion of ESC-R-*Ec* BSIs had been steadily increasing from 2017 through

2019, both significantly decreased during 2020. The reasons for these findings are not clear, but

they stand in contrast to preliminary data released by the CDC indicated an increase in ESBL

Enterobacterales in 2020 nationwide (48).

In summary, we present a comprehensive whole genome-based analyses of ESC-R-*Ec* from

the US analyzing recently collected strains. These data show that STc131 strains have reached

a stable proportion of our ESC-R-*Ec* population. Additionally, we provide molecular insights into

the temporal variability of ESC-R-*Ec* prevalence which is being increasingly appreciated but is

not well understood mechanistically (16-21). The genetically diverse nature of ESC-R-*Ec* strains

identified in our study indicate that efforts to mitigate serious ESC-R-*Ec* infections likely will

need to address broad environmental reservoirs such as through vaccination and highlight the

need to better understand ESC-R-*Ec* transmission and temporal dynamics outside of the

hospital setting.

METHODS

Study design and sample collection

Escherichia coli bloodstream infections (*Ec*-BSI) occurred from March 1st, 2016, to December 31st, 2020, at MD Anderson Cancer Center (MDACC) in Houston, TX with March 1st, 2016, chosen given this was the date of implementation of the Epic Electronic Health Record system. We defined an index *Ec*-BSI isolate as first occurrence of a positive blood culture and a recurrent *Ec*-BSI isolate as a positive blood culture that occurred at least 14 days from a previous *Ec*-BSI. Antibiotic susceptibility testing of all *Ec*-BSI isolates was performed by the MDACC microbiology laboratory using the Accelerate PhenoTest™ BC (Accelerate Diagnostics), ETEST® (bioMérieux), and VITEK® 2 (bioMérieux). *Ec*-BSI Isolates were considered as extended-spectrum cephalosporin resistant (ESC-R) if they had a ceftriaxone (CRO) minimum inhibitory concentration (MIC) $\geq$ 4 μg/mL and/or a predicted ESBL phenotype.

Genomic sequencing

ESC-R-*Ec* isolates were sequenced by Illumina NovaSeq 6000 or Illumina MiSeq as previously described (33). There were 297 ESC-R isolates that had paired-end 150 base pair reads that passed quality control assessed using fastqc-v0.11.9. We performed long-read sequencing on an nine isolates using the Oxford Nanopore Technologies GridION platform as described previously (33).

Genome analysis and molecular typing

Quality assessed, paired-end 150 bp reads were assembled via SPAdes-v3.15.2 (49) with default parameters and the inclusion of the isolate option. Quality assessment of output SPAdes assembly contigs was performed via QUAST-v5.2.0 (50) to determine N50 and contiguity. Assemblies with (a) genomes < 4 Mb or > 6 Mb; or (b) Contigs >500 were excluded from

analysis. The EzClermont *in silico* phylotyping tool was utilized with the 2013 Clermont PCR

typing method (Waters, N EzClermont Github: https://github.com/nickp60/EzClermont). *E. coli*

multilocus sequence typing (MLST) using the Achtman definition was performed with the mlst-

v2.19.0 *in silico* contig scanning tool (Seemann T mlst GitHub:

https://github.com/tseemann/mlst), which features the PUBMLST database (51). The COpy

Number Variant quantifICation Tool (convict-v1.0) was used to identify antimicrobial resistance

(AMR) genes and estimate gene copy numbers as described previously (Shropshire, W convict

GitHub: https://github.com/wshropshire/convict) (33). Briefly, short-read data is used as input to

KmerResistance-v2.2.0 (52, 53) to identify potential AMR genes from the resfinder database

(Accessed 2022-10-01). Through a coverage depth ratio of AMR gene to housekeeping gene

that accounts for variance in coverage depths, convict estimates gene copy numbers. The high-

performance computing (HPC) cluster, Seadragon, that is hosted through MDACC was used to

perform genomic analyses.

Pan genome analysis

Draft and complete genomes were annotated using the Prokka-v1.14.6 (54) using default

parameters. Annotated gff files from prokka were used as input for pangenome analysis using

Roary-v3.13.0 (55) using a protein family ID threshold of 95%. A core gene alignment was

completed using Mafft-v7.508 with the Roary pipeline using a core gene threshold of 99%. This

core gene alignment file was used as input to IQTree2-v2.2.0-beta to create a core gene

inferred maximum likelihood phylogeny. ModelFinder was utilized to determine most appropriate

DNA model with the best fit model: UNREST+FO+I+G4 chosen according to BIC. A final

bootstrap consensus tree was created using 1000 SH-like aLRT replicates in addition to 1000

Ultrafast Bootstrap support values. A core genome alignment was performed on the top 12

sequence types observed in our cohort using Snippy-v4.6.0 using default variant calling

parameters. The following were the chosen references for alignment: (ST131: CP049085.2;

ST405: CP103633.1; ST1193: MB11100; ST648: CP103657.1; ST744: MB7355; ST10:

CP049081.1; ST167: CP103540.1; ST38: CP103559.1; ST617: CP103645.1; ST410:

CP103533.1; ST224: CP103609.1; ST361: CP103704.1). A core genome alignment was used

as input for Gubbins (56) to create a recombination-free, maximum likelihood tree using

IQTree20v.2.2.0 using the aforementioned parameters. The recombination free fasta output was

used to extract pairwise SNP sites (snp-sites-v2.5.1) (57) and SNP distances (snp-dists-v0.8.2)

respectively. A pairwise SNP distance cutoff of 10 for window independent analyses of

transmission was based on the recombination-free, core-genome SNP differences of eight

pairwise comparisons of ST131 recurrent isolates where the median pairwise SNP difference

was 5.5 SNPs and the maximum observed was nine SNPs.

The accessory genome was assessed using the gene_presence_absence.csv file output from

Roary. A Jaccard distance matrix was created using this binary accessory genome file after

removing genes present in <5% and >95% of the total population. This distance matrix was

used as input to create a neighbor joining tree using the BIONJ algorithm (58). We performed a

principal component analysis of this distance matrix using base R functions. The graph network

analysis was accomplished using the same gene_presence_absence.csv input using graphia

(59, 60). Edge transformations using a k-nearest neighbors algorithm (n=8) were employed to

reduce the number of edges from 44k to 1724. Clustering was performed using the Markov

Cluster algorithm (MCL) with an inflation value=1.5 (61). Visualization of data was accomplished

using ggplot2-v3.3.5 (62) and ggtree-v.3.3.1 (63).

Long read analysis

A short- and long-read assembly pipeline (Shropshire, W flye_hybrid_assembly_pipeline

GitHub: https://github.com/wshropshire/flye_hybrid_assembly_pipeline) was used to close

complete genomes of ONT sequenced data as described previously (33). Incomplete

assemblies were re-assembled using Unicycler-v0.5.0 and manually curated for errors using

short- and long-read pileups and visualizing with the integrated genome browser (IGV-v2.14.1).

Comparison of plasmid vectors with BLAST ring image generator (BLAST) was performed using

the Proksee server while querying AMR genes with the Comprehensive Antimicrobial

Resistance Database (CARD) (64).

Statistical analysis

Prevalence rates were plotted across time using ggplot2 with a smoothing curve (*i.e.,* a loess

function) to visualize prevalence trends. Time series data was assessed for monotonic trends

using the Mann-Kendall Trend test and non-monotonic using the WAVK statistic. Correlations

between ESC-R-*Ec* and ESC-S-*Ec* were measured with a Pearson's product moment

correlation test. We used χ^2 test (Fisher's exact when observations <5) to determine statistical

differences in proportions across phylogroup/clonal complexes. A Kruskal-Wallis global test to

determine differences in gene copy numbers across groups. Pairwise and one-way Wilcoxon

Rank sum tests were to determine group level and single mean differences respectively with a

Bonferroni correction for multiple comparisons. All statistics was computed on R (4.0.4) and

α=0.05 parameter was used across all tests.

Data Availability

WGS data sequenced during this study period was submitted to NCBI and can be accessed

within BioProject PRJNA924946. WGS data from previous studies can be accessed from

BioProject PRJNA603908 and PRJNA836696 (33, 65). All R Scripts can be made available

through request of the corresponding author.

Acknowledgements

We would like to thank the MDACC clinical microbiology lab for all their hard work in identifying,

handling, and transferring these pathogenic strains of interest to us for our research projects.

Core grant CA016672(ATGC) and NIH 1S10OD024977-01 grant provide funding for the

Advanced Technology Genomics Core (ATGC) sequencing facility at MD Anderson Cancer

Center. WCS is supported through the National Institute of Allergy and Infectious Diseases

(NIAID) T32 AI141349 Training Program in Antimicrobial Resistance. Support for this study

was also provided by the NIAID R21AI151536 and P01AI152999 for S.A.S. The research in the

A.K. laboratory is supported by NIGMS 1R01GM133904-01 and the Welch Foundation

Research Grant AU-1998-20190330. The authors acknowledge the support of the High-

Performance Computing for research facility at the University of Texas MD Anderson Cancer

Center for providing computational resources that have contributed to the research results

reported in this paper.

REFERENCES

1. CDC. 2019. Antibiotic Resistance Threats in the United States, 2019. U.S. Department of Health and Human Services, CDC, Atlanta, GA.
2. Jernigan JA, Hatfield KM, Wolford H, Nelson RE, Olubajo B, Reddy SC, McCarthy N, Paul P, McDonald LC, Kallen A, Fiore A, Craig M, Baggs J. 2020. Multidrug-Resistant Bacterial Infections in U.S. Hospitalized Patients, 2012–2017. New England Journal of Medicine 382:1309-1319.
3. Kaye KS, Gupta V, Mulgirigama A, Joshi AV, Scangarella-Oman NE, Yu K, Ye G, Mitrani-Gold FS. 2021. Antimicrobial Resistance Trends in Urine Escherichia coli Isolates From Adult and Adolescent Females in the United States From 2011 to 2019: Rising ESBL Strains and Impact on Patient Management. Clin Infect Dis 73:1992-1999.
4. Begier E, Rosenthal NA, Gurtman A, Kartashov A, Donald RGK, Lockhart SP. 2021. Epidemiology of Invasive Escherichia coli Infection and Antibiotic Resistance Status Among Patients Treated in US Hospitals: 2009-2016. Clin Infect Dis 73:565-574.
5. Mackinnon MC, McEwen SA, Pearl DL, Lyytikäinen O, Jacobsson G, Collignon P, Gregson DB, Valiquette L, Laupland KB. 2021. Mortality in Escherichia coli bloodstream infections: a multinational population-based cohort study. BMC Infectious Diseases 21:606.
6. Mackinnon MC, McEwen SA, Pearl DL, Parfitt EC, Pasquill K, Steele L, Laupland KB. 2021. Escherichia coli bloodstream infections in the western interior of British Columbia, Canada: a population-based cohort study. Epidemiology and Infection 149:1-28.
7. Trecarichi EM, Giuliano G, Cattaneo C, Ballanti S, Criscuolo M, Candoni A, Marchesi F, Laurino M, Dargenio M, Fanci R, Cefalo M, Delia M, Spolzino A, Maracci L, Nadali G, Busca A, Del Principe MI, Daffini R, Simonetti E, Dragonetti G, Zannier ME, Pagano L, Tumbarello M. 2019. Bloodstream infections caused by Escherichia coli in onco-haematological patients: Risk factors and mortality in an Italian prospective survey. PLOS ONE 14:e0224465.
8. Philippon A, Arlet G, Jacoby GA. 2002. Plasmid-Determined AmpC-Type β-Lactamases. Antimicrobial Agents and Chemotherapy 46:1-11.
9. Zamudio R, Boerlin P, Beyrouthy R, Madec J-Y, Schwarz S, Mulvey MR, Zhanel GG, Cormier A, Chalmers G, Bonnet R, Haenni M, Eichhorn I, Kaspar H, Garcia-Fierro R, Wood JLN, Mather AE. 2022. Dynamics of extended-spectrum cephalosporin resistance genes in Escherichia coli from Europe and North America. Nature Communications 13:7490.
10. Tamma PD, Sharara SL, Pana ZD, Amoah J, Fisher SL, Tekle T, Doi Y, Simner PJ. 2019. Molecular Epidemiology of Ceftriaxone-Nonsusceptible Enterobacterales Isolates in an Academic Medical Center in the United States. Open Forum Infectious Diseases 6:ofz353.
11. Castanheira M, Deshpande LM, Mendes RE, Canton R, Sader HS, Jones RN. 2019. Variations in the Occurrence of Resistance Phenotypes and Carbapenemase Genes Among Enterobacteriaceae Isolates in 20 Years of the SENTRY Antimicrobial Surveillance Program. Open Forum Infect Dis 6:S23-S33.
12. Bush K, Jacoby GA. 2010. Updated Functional Classification of β-Lactamases. Antimicrobial Agents and Chemotherapy 54:969-976.
13. Perez F, Endimiani A, Hujer K, Bonomo R. 2007. The continuing challenge of ESBLs. Current Opinion in Pharmacology 7:459-469.
14. Stoesser N, Sheppard AE, Pankhurst L, De Maio N, Moore CE, Sebra R, Turner P, Anson LW, Kasarskis A, Batty EM, Kos V, Wilson DJ, Phetsouvanh R, Wyllie D, Sokurenko E, Manges AR, Johnson TJ, Price LB, Peto TE, Johnson JR, Didelot X, Walker AS, Crook DW, Modernizing Medical Microbiology Informatics G. 2016.

Evolutionary History of the Global Emergence of the Escherichia coli Epidemic Clone
ST131. MBio 7:e02162.

15.   Price LB, Johnson JR, Aziz M, Clabots C, Johnston B, Tchesnokova V, Nordstrom L,
Billig M, Chattopadhyay S, Stegger M, Andersen PS, Pearson T, Riddell K, Rogers P,
Scholes D, Kahl B, Keim P, Sokurenko EV. 2013. The Epidemic of Extended-Spectrum-
β-Lactamase-Producing Escherichia coli ST131 Is Driven by a Single Highly Pathogenic
Subclone, H30-Rx. mBio 4:e00377-13-e00377.

16.   Lipworth S, Vihta K-D, Chau K, Barker L, George S, Kavanagh J, Davies T, Vaughan A,
Andersson M, Jeffery K, Oakley S, Morgan M, Hopkins S, Peto TEA, Crook DW, Walker
AS, Stoesser N. 2021. Ten-year longitudinal molecular epidemiology study of
Escherichia coli and Klebsiella species bloodstream infections in Oxfordshire, UK.
Genome Medicine 13:1-13.

17.   Royer G, Darty MM, Clermont O, Condamine B, Laouenan C, Decousser J-W, Vallenet
D, Lefort A, De Lastours V, Denamur E. 2021. Phylogroup stability contrasts with high
within sequence type complex dynamics of Escherichia coli bloodstream infection
isolates over a 12-year period. Genome Medicine 13:77.

18.   Verschuuren TD, van Hout D, Arredondo-Alonso S, Fluit AC, Reuland EA, Top J,
Schurch AC, Bosch T, Bonten MJM, Kluytmans J, Willems RJL. 2021. Comparative
genomics of ESBL-producing Escherichia coli (ESBL-Ec) reveals a similar distribution of
the 10 most prevalent ESBL-Ec clones and ESBL genes among human community
faecal and extra-intestinal infection isolates in the Netherlands (2014-17). J Antimicrob
Chemother 76:901-908.

19.   Kallonen T, Brodrick HJ, Harris SR, Corander J, Brown NM, Martin V, Peacock SJ,
Parkhill J. 2017. Systematic longitudinal survey of invasive Escherichia coli in England
demonstrates a stable population structure only transiently disturbed by the emergence
of ST131. Genome Research 27:1437-1449.

20.   Gladstone RA, McNally A, Pöntinen AK, Tonkin-Hill G, Lees JA, Skytén K, Cléon F,
Christensen MOK, Haldorsen BC, Bye KK, Gammelsrud KW, Hjetland R, Kümmel A,
Larsen HE, Lindemann PC, Löhr IH, Marvik Å, Nilsen E, Noer MT, Simonsen GS,
Steinbakk M, Tofteland S, Vattøy M, Bentley SD, Croucher NJ, Parkhill J, Johnsen PJ,
Samuelsen Ø, Corander J. 2021. Emergence and dissemination of antimicrobial
resistance in Escherichia coli causing bloodstream infections in Norway in 2002–17: a
nationwide, longitudinal, microbial population genomic study. The Lancet Microbe
2:e331-e341.

21.   Rodríguez I, Figueiredo AS, Sousa M, Aracil-Gisbert S, Fernández-de-Bobadilla MD,
Lanza VF, Rodríguez C, Zamora J, Loza E, Mingo P. 2021. A 21-year survey of
Escherichia coli from bloodstream infections (BSI) in a tertiary hospital reveals how
community-hospital dynamics of B2 phylogroup clones influence local BSI rates.
Msphere 6:e00868-21.

22.   Peirano G, Lynch T, Matsumara Y, Nobrega D, Finn TJ, Devinney R, Pitout JDD. 2020.
Trends in Population Dynamics of Escherichia coli Sequence Type 131, Calgary,
Alberta, Canada, 2006–2016. Emerging Infectious Diseases 26:2907-2915.

23.   Mills EG, Martin MJ, Luo TL, Ong AC, Maybank R, Corey BW, Harless C, Preston LN,
Rosado-Mendez JA, Preston SB, Kwak YI, Backlund MG, Bennett JW, Mc Gann PT,
Lebreton F. 2022. A one-year genomic investigation of Escherichia coli epidemiology
and nosocomial spread at a large US healthcare network. Genome Medicine 14.

24.   Manges AR, Geum HM, Guo A, Edens TJ, Fibke CD, Pitout JDD. 2019. Global
Extraintestinal Pathogenic Escherichia coli (ExPEC) Lineages. Clin Microbiol Rev
32:e00135-18.

25.   Banerjee R, Johnson JR. 2014. A new clone sweeps clean: the enigmatic emergence of
Escherichia coli sequence type 131. Antimicrob Agents Chemother 58:4997-5004.

26.   McNally A, Kallonen T, Connor C, Abudahab K, Aanensen DM, Horner C, Peacock SJ,
662   Parkhill J, Croucher NJ, Corander J. 2019. Diversification of Colonization Factors in a
663   Multidrug-Resistant Escherichia coli Lineage Evolving under Negative Frequency-
664   Dependent Selection. mBio 10:e00644-19.
665   27.   Johnson JR, Porter S, Thuras P, Castanheira M. 2017. The Pandemic H30 Subclone of
666   Sequence Type 131 (ST131) as the Leading Cause of Multidrug-Resistant Escherichia
667   coli Infections in the United States (2011-2012). Open Forum Infect Dis 4:ofx089.
668   28.   Matsumura Y, Pitout JD, Gomi R, Matsuda T, Noguchi T, Yamamoto M, Peirano G,
669   DeVinney R, Bradford PA, Motyl MR, Tanaka M, Nagao M, Takakura S, Ichiyama S.
670   2016. Global Escherichia coli Sequence Type 131 Clade with blaCTX-M-27 Gene.
671   Emerg Infect Dis 22:1900-1907.
672   29.   Hansen KH, Andreasen MR, Pedersen MS, Westh H, Jelsbak L, Schonning K. 2019.
673   Resistance to piperacillin/tazobactam in Escherichia coli resulting from extensive IS26-
674   associated gene amplification of blaTEM-1. J Antimicrob Chemother 74:3179-3183.
675   30.   Rodriguez-Villodres A, Gil-Marques ML, Alvarez-Marin R, Bonnin RA, Pachon-Ibanez
676   ME, Aguilar-Guisado M, Naas T, Aznar J, Pachon J, Lepe JA, Smani Y. 2019. Extended-
677   spectrum resistance to beta-lactams/beta-lactamase inhibitors (ESRI) evolved from low-
678   level resistant Escherichia coli. J Antimicrob Chemother 75:77-85.
679   31.   Nicoloff H, Hjort K, Levin BR, Andersson DI. 2019. The high prevalence of antibiotic
680   heteroresistance in pathogenic bacteria is mainly caused by gene amplification. Nat
681   Microbiol 4:504-514.
682   32.   Shropshire WC, Aitken SL, Pifer R, Kim J, Bhatti MM, Li X, Kalia A, Galloway-Peña J,
683   Sahasrabhojane P, Arias CA, Greenberg DE, Hanson BM, Shelburne SA. 2021. IS26-
684   mediated amplification of blaOXA-1 and blaCTX-M-15 with concurrent outer membrane
685   porin disruption associated with de novo carbapenem resistance in a recurrent
686   bacteraemia cohort. Journal of Antimicrobial Chemotherapy 76:385-395.
687   33.   Shropshire WC, Konovalova A, McDaneld P, Gohel M, Strope B, Sahasrabhojane P,
688   Tran CN, Greenberg D, Kim J, Zhan X, Aitken S, Bhatti M, Savidge TC, Treangen TJ,
689   Hanson BM, Arias CA, Shelburne SA. 2022. Systematic Analysis of Mobile Genetic
690   Elements Mediating Beta-Lactamase Gene Amplification in Noncarbapenemase-
691   Producing Carbapenem-Resistant Enterobacterales Bloodstream Infections. Msystems
692   7:e00476-22.
693   34.   Mahmud B, Wallace MA, Reske KA, Alvarado K, Muenks CE, Rasmussen DA, Burnham
694   C-AD, Lanzas C, Dubberke ER, Dantas G. 2022. Epidemiology of Plasmid Lineages
695   Mediating the Spread of Extended-Spectrum Beta-Lactamases among Clinical
696   Escherichia coli. Msystems 7:e00519-22.
697   35.   Cai SJ, Inouye M. 2002. EnvZ-OmpR Interaction and Osmoregulation in Escherichia
698   coli. Journal of Biological Chemistry 277:24155-24161.
699   36.   Anesi JA, Lautenbach E, Tamma PD, Thom KA, Blumberg EA, Alby K, Bilker WB,
700   Werzen A, Tolomeo P, Omorogbe J, Pineles L, Han JH. 2021. Risk Factors for
701   Extended-Spectrum beta-lactamase-Producing Enterobacterales Bloodstream Infection
702   Among Solid-Organ Transplant Recipients. Clin Infect Dis 72:953-960.
703   37.   Johnson JR, Urban C, Weissman SJ, Jorgensen JH, Lewis JS, Hansen G, Edelstein PH,
704   Robicsek A, Cleary T, Adachi J, Paterson D, Quinn J, Hanson ND, Johnston BD,
705   Clabots C, Kuskowski MA. 2012. Molecular Epidemiological Analysis of Escherichia coli
706   Sequence Type ST131 (O25:H4) and blaCTX-M-15 among Extended-Spectrum-β-
707   Lactamase-Producing E. coli from the United States, 2000 to 2009. Antimicrobial Agents
708   and Chemotherapy 56:2364-2370.
709   38.   Gradel KO, Nielsen SL, Pedersen C, Knudsen JD, Østergaard C, Arpi M, Jensen TG,
710   Kolmos HJ, Søgaard M, Lassen AT, Schønheyder HC. 2016. Seasonal Variation of
711   Escherichia coli, Staphylococcus aureus, and Streptococcus pneumoniae Bacteremia

According to Acquisition and Patient Characteristics: A Population-Based Study.
Infection Control & Hospital Epidemiology 37:946-953.

39. Wielders CCH, Van Duijkeren E, Van Den Bunt G, Meijs AP, Dierikx CM, Bonten MJM,
Van Pelt W, Franz E, De Greeff SC. 2020. Seasonality in carriage of extended-spectrum
β-lactamase-producing Escherichia coli and Klebsiella pneumoniae in the general
population: a pooled analysis of nationwide cross-sectional studies. Epidemiology and
Infection 148:1-17.

40. Deeny SR, Van Kleef E, Bou-Antoun S, Hope RJ, Robotham JV. 2015. Seasonal
changes in the incidence of Escherichia coli bloodstream infection: variation with region
and place of onset. Clinical Microbiology and Infection 21:924-929.

41. Maryam S, Ul Haq I, Yahya G, Ul Haq M, Algammal AM, Saber S, Cavalu S. 2022.
COVID-19 surveillance in wastewater: An epidemiological tool for the monitoring of
SARS-CoV-2. Front Cell Infect Microbiol 12:978643.

42. Wolken M, Sun T, McCall C, Schneider R, Caton K, Hundley C, Hopkins L, Ensor K,
Domakonda K, Kalvapalle P, Persse D, Williams S, Stadler LB. 2023. Wastewater
surveillance of SARS-CoV-2 and influenza in preK-12 schools shows school,
community, and citywide infections. Water Res 231:119648.

43. Johnson TJ, Danzeisen JL, Youmans B, Case K, Llop K, Munoz-Aguayo J, Flores-
Figueroa C, Aziz M, Stoesser N, Sokurenko E, Price LB, Johnson JR. 2016. Separate F-
Type Plasmids Have Shaped the Evolution of the H 30 Subclone of Escherichia coli
Sequence Type 131. mSphere 1:e00121-16.

44. Yoon E-J, Gwon B, Liu C, Kim D, Won D, Park SG, Choi JR, Jeong SH. 2020. Beneficial
Chromosomal Integration of the Genes for CTX-M Extended-Spectrum β-Lactamase in
Klebsiella pneumoniae for Stable Propagation. Msystems 5:e00459-20.

45. Yoon E-J, Choi YJ, Kim D, Won D, Choi JR, Jeong SH. 2022. Amplification of the
Chromosomal bla CTX-M-14 Gene in Escherichia coli Expanding the Spectrum of
Resistance under Antimicrobial Pressure. Microbiology Spectrum:e00319-22.

46. Zeng S, Luo J, Li X, Zhuo C, Wu A, Chen X, Huang L. 2021. Molecular Epidemiology
and Characteristics of CTX-M-55 Extended-Spectrum beta-Lactamase-Producing
Escherichia coli From Guangzhou, China. Front Microbiol 12:730012.

47. Feldman SF, Temkin E, Wulffhart L, Nutman A, Schechner V, Shitrit P, Shvartz R,
Schwaber MJ, Carmeli Y. 2022. Effect of temperature on Escherichia coli bloodstream
infection in a nationwide population-based study of incidence and resistance.
Antimicrobial Resistance & Infection Control 11:1-9.

48. CDC. 2022. COVID-19: US impact on antimicrobial resistance, special report 2022. U.S.
Department of Health and Human Services, CDC, Atlanta, GA.

49. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes
De Novo Assembler. Current Protocols in Bioinformatics 70:e102.

50. Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. 2018. Versatile genome
assembly evaluation with QUAST-LG. Bioinformatics 34:i142-i150.

51. Jolley KA, & Maiden, M. C. 2010. BIGSdb: scalable analysis of bacterial genome
variation at the population level. BMC bioinformatics 11:595.

52. Clausen PTLC, Zankari E, Aarestrup FM, Lund O. 2016. Benchmarking of methods for
identification of antimicrobial resistance genes in bacterial whole genome data. Journal
of Antimicrobial Chemotherapy 71:2484-2488.

53. Clausen PTLC, Aarestrup FM, Lund O. 2018. Rapid and precise alignment of raw reads
against redundant databases with KMA. BMC Bioinformatics 19.

54. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics
30:2068-9.

55.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush D,
762   Keane JA, Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome analysis.
763   Bioinformatics 31:3691-3.
764   56.   Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J,
765   Harris SR. 2015. Rapid phylogenetic analysis of large samples of recombinant bacterial
766   whole genome sequences using Gubbins. Nucleic Acids Res 43:e15.
767   57.   Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, Harris SR. 2016.
768   SNP-sites: rapid efficient extraction of SNPs from multi-FASTA alignments. Microbial
769   Genomics 2:e000056.
770   58.   Gascuel O. 1997. BIONJ: an improved version of the NJ algorithm based on a simple
771   model of sequence data. Molecular Biology and Evolution 14:685-695.
772   59.   Freeman TC, Horsewell S, Patir A, Harling-Lee J, Regan T, Shih BB, Prendergast J,
773   Hume DA, Angus T. 2022. Graphia: A platform for the graph-based visualisation and
774   analysis of high dimensional data. PLOS Computational Biology 18:e1010310.
775   60.   Harling-Lee JD, Gorzynski J, Yebra G, Angus T, Fitzgerald JR, Freeman TC. 2022. A
776   graph-based approach for the visualisation and analysis of bacterial pangenomes. BMC
777   Bioinformatics 23:1-15.
778   61.   Van Dongen S. 2008. Graph clustering via a discrete uncoupling process. SIAM Journal
779   on Matrix Analysis and Applications 30:121-141.
780   62.   Wickham H. 2016. ggplot2: elegant graphics for data analysis Springer-Verlag New York
781   63.   Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. 2017. ggtree: package for visualization and
782   annotation of phylogenetic trees with their covariates and other associated data.
783   Methods in Ecology and Evolution 8:28-36.
784   64.   Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, Lago BA, Dave BM,
785   Pereira S, Sharma AN, Doshi S, Courtot M, Lo R, Williams LE, Frye JG, Elsayegh T,
786   Sardar D, Westman EL, Pawlowski AC, Johnson TA, Brinkman FS, Wright GD, McArthur
787   AG. 2017. CARD 2017: expansion and model-centric curation of the comprehensive
788   antibiotic resistance database. Nucleic Acids Res 45:D566-D573.
789   65.   Shelburne SA, Kim J, Munita JM, Sahasrabhojane P, Shields RK, Press EG, Li X, Arias
790   CA, Cantarel B, Jiang Y, Kim MS, Aitken SL, Greenberg DE. 2017. Whole-Genome
791   Sequencing Accurately Identifies Resistance to Extended-Spectrum beta-Lactams for
792   Major Gram-Negative Bacterial Pathogens. Clin Infect Dis 65:738-745.

794 **TABLES**

Table 1: Windows of *E. coli* Bloodstream Infection Rates per 1000 Hospital Admissions from 2016-03-01 to 2020-12-31							
Window	1	2 (ESC-R Peak 1)	3	4 (ESC-R Peak 2)	5	6 (ESC-R Peak 3)	7
Date Range (Days)	16/03/01 – 16/06/30 (121)	16/07/01 – 17/04/30 (394)	17/05/01 – 18/06/30 (425)	18/07/01 – 18/12/31 (183)	19/01/01 – 19/04/30 (119)	19/05/01 – 19/12/31 (244)	20/01/01 – 20/12/31 (365)
Total *E. coli* BSI per 1000 admissions	4.44	4.77	4.31	4.96	3.86	5.60	5.11
ESC-R-*Ec* per 1000 admissions	0.92	1.89	1.16	2.09	1.07	2.24	1.63
ESC-S-*Ec* per 1000 admissions	3.51	2.88	3.15	2.87	2.80	3.36	3.48

Table 2: Overview of Sequence Type Distribution Across Each Phylogroup from Sequenced Index and Recurrent ESC-R *E. coli* Isolates

Typing Group[a,b]	Index (n=248)	Recurrent (n=49)
Phylogroup B2 (n=145)	**126 (50.8%)**	**19 (38.8%)**
ST131 (n=129)	102 (80.9%)	17 (89.5%)
ST1193 (n=15)	14 (11.1%)	1 (5.3%)
Other ST (n=11)	10 (7.9%)	1 (5.3%)
Phylogroup A (n=66)	**49 (19.8%)**	**17 (34.7%)**
ST10 (n=13)	8 (16.3%)	5 (29.4%)
ST744 (n=11)	9 (18.4%)	2 (11.8%)
ST167 (n=11)	8 (16.3%)	3 (17.6%)
ST44 (n=7)	2 (4.1%)	5 (29.4%)
Other ST (n=24)	22 (44.9%)	2 (11.8%)
Phylogroup D (n=36)	**30 (12.1%)**	**6 (12.2%)**
ST405 (n=19)	16 (53.3%)	3 (50.0%)
ST38 (n=9)	7 (23.3%)	2 (33.3%)
Other ST (n=8)	7 (23.3%)	1 (16.7%)
Phylogroup F (n=20)	**19 (7.7%)**	**1 (2.0%)**
ST648 (n=15)	14 (73.7%)	1 (100.0%)
Other ST (n=5)	5 (26.3%)	0 (0.0%)
Phylogroup B1 (n=20)	**16 (6.5%)**	**4 (8.2%)**
ST224 (n=6)	5 (31.3%)	1(25.0%)
Other ST (n=13)	11 (57.9%)	3 (75.0%)
Phylogroup C (n=9)	**7 (2.8%)**	**2 (4.1%)**
ST410 (n=7)	5 (71.4%)	1 (50.0%)

| Other ST (n=2) | 2 (28.6%) | 1 (50.0%) |

a'Other ST' consists of sequence types detected at ≤5 total counts within each phylogroup.

bPhylogroup G consists of one observation (ST501) thus was not included in table.

Table 3: Copy Number Variants of β-lactamase Encoding Genes

	CTX-M-15	CTX-M-27	CTX-M-55	CTX-M-14	Other CTX-M
Total					
Count	161	37	37	23	10
Mean (SD)	1.62 (1.29)	1.87 (1.89)	2.50 (2.65)	2.61 (2.57)	3.55 (3.35)
Median (IQR)	1.11 (1.19)	1.40 (0.80)	1.50 (1.43)	1.42 (2.21)	1.31 (5.36)
Range	0.37-8.72	0.42-9.10	0.62-12.9	0.64-10.8	0.60-9.45
Phylogroup B2					
Count	92	33	1	6	3
Mean (SD)	1.41 (0.96)	1.12 (0.36)	3.41	2.53 (2.06)	4.80 (3.66)
Median (IQR)	1.03 (0.54)	1.20 (0.354)	NA	1.75 (1.83)	6.54 (3.33)
Range	0.44-5.84	0.42-9.10	NA	0.88-6.33	0.60-7.26
Phylogroup A					
Count	31	3	19	5	2
Mean (SD)	1.66 (1.72)	1.12 (0.36)	2.39 (2.09)	1.65 (1.42)	1.08 (0.13)
Median (IQR)	1.07 (0.97)	1.20 (0.35)	1.68 (1.82)	1.01 (0.31)	1.08 (0.09)
Range	0.50 - 8.72	0.72 - 1.44	0.62 - 7.20	0.78-4.16	0.98-1.17
Phylogroup D					
Count	21	1	5	4	3
Mean (SD)	2.02 (1.48)	1.8	1.25 (0.49)	1.80 (0.98)	3.90 (4.81)
Median (IQR)	2.00 (1.62)	NA	1.30 (0.53)	1.43 (0.54)	1.31 (4.25)
Range	0.37-5.14	NA	1.00-1.96	1.10-3.26	0.94-9.45

SD = Standard Deviation; IQR = Interquartile Range; Copy Number Variants are estimated based on CONVICT results

FIGURES

Figure 1. Number of Bacteremia Events Observed per Month from March 2016 to

December 2020 at MD Anderson Cancer Center. Number of index *E. coli* bacteremia events

per 1000 hospital admissions for both ESC-R (red) and ESC-S (blue) isolates. A local

regression curve for each respective ESC category is provided with 95% confidence intervals

(gray area). The dotted gray line represents the prevalence trend for total *E. coli* bacteremia

events. Windows of peak infection estimates were observed in windows 2, 4, and 6 as

demarcated above by vertical dotted black lines with estimated peak date indicated within peak

regions (vertical dotted red lines).

809 **Figure 2. Temporal Distribution of ESC-R *Ec*-BSI Isolates.** The proportional distribution and

810 absolute frequency counts by phylogroup **(A-B)** and clonal complex **(C-D)** respectively across

811 each infection window. **(A-C)** Group totals are listed above each of the windows. **(B-D)** Data

812 shown are line graphs connecting month-to-month frequency counts colored by proportions of

813 phylogroup **(B)** and clonal complex **(D)** as indicated in the legend.

Figure 3. Population Structure of ESC-R *Ec*-BSI isolates. A core gene alignment inferred

maximum likelihood phylogenetic tree with phylogroups highlighted within each monophyletic

group. Branch tip colors indicate clonal complex number as shown in the figure inset with clonal

complexes observed <5 counts grouped as "Uncommon". Rings corresponding to infection

window (1), recurrence (2), and β-lactamases are labelled respectively. Internal nodes labelled

with black circles have SH-aLRT ≥ 80% and UFboot ≥ 95%.

Figure 4. Accessory Genome Clustering by ST131 and non-ST131 ESC-R Isolates.

Jaccard distance matrix inferred NJ tree of accessory genome content with metadata indicated

in legend. Phylogroups and chosen STs/clades are highlighted within tree and labelled

accordingly. MCL = Markov Clustering group; STc = Sequence Type Clonal Complex

Figure 5. *Bla*$_{CTX-M}$ Amplification Occurring within ESC-R-*Ec* Isolates. (A) Copy number estimates of *bla*$_{CTX-M}$ variants stratified by

phylogroup indicated in facetted plots that are ESC-R and carbapenem susceptible. One-sample Wilcoxon rank sum test for

amplification (*i.e.*, copy number > 1) performed across each *bla*$_{CTX-M}$ variant with * indicating *P*-value<0.05 **(B)** Estimated copy

number of *bla*$_{CTX-M}$ variants of paired ESC-R serial isolates (selection criteria as outlined in the text). Shapes and colors indicate

clonal complex and CTX-M variant as noted in the legend. Dotted lines connect paired isolates. Paired Wilcoxon-signed rank test

was statistically significant with * indicating *P*-value<0.05.

Figure 6. Characterization of mobile genetic elements associated with transmission and amplification of *bla*CTX-M-55 within

non-B2 isolates. (A) BRIGs output of similar IncFIB/IncFIC plasmids with conserved carriage of IS*Ecp1-bla*CTX-M-55 carriage

(Underlined) in 4 different non-B2 phylogroups. Inner ring indicates GC content/skew with coding DNA sequences in reverse/forward

orientation. Next seven rings indicate blastn identities to pMB6206_1 as indicated in legend. Outer ring includes plasmid, mobile

genetic element, and antimicrobial resistance gene content as identified by the Comprehensive Antimicrobial Resistance Database

(CARD). **(B)** Blast nucleotide and gene amplification comparisons of differential *bla*CTX-M-55 vectors responsible for plasmid

transmission as well as potential vertical transmission via chromosomal insertion. Normalized coverage depth of conserved ~10kb

region that harbors IS*Ecp1-bla*CTX-M-55 (~10kb) as demarcated by dotted black lines in **(A).** Inferred plasmid copy number vs. plasmid-

to-chromosome events as indicated by tracking target site duplications of IS*Ecp1-bla*CTX-M-55 on complete assemblies as well as mean

coverage depths of full-length plasmid. Coding sequences within blastn comparisons are identified as followed: IS*Ecp1* (purple),

*bla*CTX-M-55 (green), AMR genes (red), IS*26* (white), IS*26*-v1 (beige), other MGE (black), other CDS (gray). Black horizontal line

demarcates IncFIB/IncFIC positive isolates from high copy number IncX1 plasmids harboring *bla*CTX-M-55 from ST167 isolates.

Parentheticals following plasmid label indicate plasmid length and copy number. CDS with striped, purple label indicates IS*Ecp1*

truncated by IS*26*-v1 upstream of *bla*CTX-M-55.