Experience with mating receptivity cues affects sexual behaviour of male guppies, but not their strength of preference towards receptive females

Versara Goberdhan[1], Iulia Darolti[1], Judith E. Mank[1], Alberto Corral-Lopez[1*]

1 Department of Zoology and Biodiversity Research Centre, University of British Columbia

*corresponding author: corral@zoology.ubc.ca

Abstract

Females are traditionally presented as the choosier sex, selecting males based on the quality of their traits. Yet, there is increasing evidence that male mate choice is also important, even in species without male parental care. Social environment and learning are key factors in determining mate preference, and animals are able to use the information they gather from previous experience to potentially increase their odds of obtaining a high-quality mate. We examined how the social environment affects male mate choice in the guppy (*Poecilia reticulata*). We evaluated whether male guppies with previous social experience of female receptivity cues learn to prefer and adapt their behavioural repertoire towards females with higher receptiveness levels, as this represents an optimal use of time and energy and is more likely to result in insemination. For this, we measured sexual preference and behaviour for receptive females in no-choice and dichotomous choice tests using guppy males experienced or naïve to female receptivity cues. Experience with receptivity cues did not change the strength of preference towards receptive females. However, male guppies that had previous experience with female receptivity cues adapted their mating tactic compared to naïve males. The change in mating tactics but lack of preference towards receptive females shows that the influence of social learning is present but might be weaker than predicted in this species. Furthermore, these results provide further support to studies of female mate choice suggesting mating status is not a key factor driving the strength of sexual preferences in natural populations.

Keywords: mating status; alternative mating tactics; social learning; sexual selection; male choice

Introduction

Mate choice is a fundamental evolutionary process, as it influences which individuals successfully pass their genes to the next generation (Darwin, 1871; Andersson, 1996). While females historically have been studied as the choosy sex, there is compelling evidence of male mate choice, suggesting that males can be choosy even in species without male parental care (Edward & Chapman, 2011; Herdman et al., 2004; Rosenqvist, 1990; Werner & Lotem, 2003; Wong & Jennions, 2003).

Previous experience can play an important role in forming mating preferences. The social environment can create opportunities for animals to collect information about their surroundings. Animals regularly use this information to adapt their behaviour and efficiently obtain and compete for mates, as well of for other resources that lead them to increase their fitness (Bailey & Moore, 2012; Danchin et al., 2004; Fowler-Finn & Rodríguez, 2012; Valone & Templeton, 2002). In systems in which males perform sexual displays to solicit copulation consent, previous experience can help males to better assess female quality or to avoid soliciting females that signal unwillingness to mate (Akinyemi & Kirk 2019, Dukas 2005, Rather et al., 2022). Similarly, males can adapt their behavioural repertoire or shift to alternative mating tactics based on previous encounters with females (Bailey et al, 2010, Řežucha & Reichard, 2014). Considering the effect of social environment is therefore paramount when evaluating male mate choice patterns.

In fish, male mate choice has been documented in many species (Schlupp, 2018), with males selecting for female traits associated with higher fecundity. For instance, males exhibit preference for larger females with higher fecundity potential in species such as eastern mosquitofish, *Gambusia holbrooki* (Bisazza et al., 1989; Head et al., 2015, Hoysak & Godin, 2007), sailfin mollies, *Poecilia latipinna* (Gumm & Gabor, 2005), and Atlantic mollies, *Poecilia mexicana* (Plath et al., 2006). In fish species with internal fertilization, preference for females showing higher receptiveness levels is theoretically expected, as it is more likely to result in insemination and therefore higher reproductive fitness per unit effort (Bondurianski, 2001, Jordan et al., 2014). However, more research effort is needed to understand pre-copulatory male mate choice in relation to female mating status.

Guppies, fish native to streams of north-eastern South America and the Caribbean, are a traditional

model for sexual selection studies. The species is sexually dimorphic, with smaller and colorful

males, and have a non-resource based promiscuous mating system (Houde, 1997). Studies of male

mate choice in the guppy have revealed that males prefer larger, more fecund females (Corral-

Lopez et al., 2018, Dosen and Montgomerie, 2004, Herdman et al., 2004., Jeswiet et al., 2012).

Additionally, guppy males adapt their mating tactics and sexual effort in relation to female mating

status (Guevara-Fiore et al., 2009; 2010). Studies in guppies have likewise been crucial for our

understanding of the role of social environment in sexual selection, and have shown that females

shift their preferences for male color traits depending on early social experiences (Rosenqvist &

Houde, 1997, Macario et al., 2017). Similarly, rearing conditions and previous success in mating

affect subsequent male mating behavioural tactics (Guevara-Fiore, 2012; Guevara-Fiore & Endler,

2018).

While the role the social environment in sexual selection and male mate choice for specific female

traits have been studied in guppies, this system provides the additional opportunity to understand

how prior social experiences can affect male preference for female mating status. Here we study

whether previous social experience with female receptivity cues affect behaviour and strength of

preference of male guppies in relation to female mating status. To empirically test this, we

experimentally manipulated the social environment of male guppies and quantified their

preference and behavioral repertoire in the presence of receptive and non-receptive females. To

avoid potential biases introduced by choice of experimental design paradigm (Dougherty &

Shuker, 2015), we assessed the role of previous experience in guppy male choice for receptive

females using a combination of dichotomous choice and no choice tests. Given theoretical

expectations of fitness maximization, we predicted that male guppies would use social learning to

shift their preference levels to favour receptive females. Similarly, we predicted that the social

environment would affect how males adapt their behavioural repertoire depending on whether they

encounter receptive or non-receptive females.

Methods

Study system

All guppies used in this experiment originated from a laboratory-adapted stock population,
originally collected from the high predation region of the Quaré River (Trinidad & Tobago).
Aquaria contained gravel, water filters, and aquatic plants, and all experiments were approved by
institutional animal ethics protocols. Fish were raised at a water temperature of 25 °C with a 12:12
light:dark schedule, and fed a daily diet of flake food (Hikari Fancy Food) and live *Artemia* brine
shrimp.
71
We collected newborn guppies from a stock aquarium and held them in nursery aquaria until they
could be accurately sexed by the development of a gonopodium, a modified anal fin (Houde, 1997;
Liley, 1966), at which point we removed males and held them in male-specific aquaria in groups
of seven individuals. During this time, males were not allowed to visualize any females. Once
males reached sexual maturity, as evidenced by the development of male colouration, we randomly
allocated them into two experimental treatments, experienced and naïve. To ensure all males had
similar age and social experience when tested in behavioural experiments and due to logistic
reasons, we performed the experiment in two batches that account for half of the individuals each.
80
To study the preference and behaviour of naïve and experienced males towards females with
different mating status, we exposed them to receptive and non-receptive females in dichotomous
and no-choice tests (see details below). Sexual receptiveness towards males strongly correlates
with the female guppy reproductive cycle. Levels of female receptiveness are highest following
parturition of live offspring and for a period of approximately three days in which new ova are
commonly fertilized. Receptiveness levels decrease linearly for the following days until they reach
minimum levels approximately ten days post-parturition and are maintained in minimum levels
until parturition of a new clutch of offspring (approximately 28 days; Liley, 1966, Houde, 1997).
Receptiveness in virgin females presents a similar pattern during first reproductive cycle (Houde,
1997). Following theoretical expectations of receptiveness levels towards males and methods in
Guevara-Fiore et al. (2010), we housed small groups of virgin females with males in a 1:1 ratio
and used them in behavioural tests the following day (receptive females) or 14 days after (non-
receptive females).
94
95 *Dichotomous choice preference tests*
96

To assess potential differences in preference for receptive females between naïve and experienced
males, we measured time associating with receptive and non-receptive females in dichotomous
choice tests. We performed two dichotomous choice tests, with an intervening 45 days treatment
exposure to females for experienced but not for naïve males. This testing protocol allowed us to
determine whether experienced males acquire information about receptivity during their extended
exposure to females.

First, to measure baseline preference for receptive females (pre-treatment test), we performed an
initial dichotomous choice test on 62 reproductively mature males of similar age (approximately
four months old). We photographed each male after behavioural testing using a Canon EOS Rebel
T7i camera in a small glass aquarium (5 x 5 x 5 cm) with white walls and a scale for sizing. Camera
colour calibration was performed daily with a Calibrite ColorChecker (X-Rite Inc.). Next, we
transferred males to treatment 10L aquaria for a 45-day period. Specifically, we transferred each
male tested in the pre-treatment test to a separate tank, with half of the males placed with two other
virgin males, two virgin females and two non-virgin females of similar age (experienced
condition), and the other half placed with two other virgin males of similar age (naïve condition).
Naïve males had restricted visual access to tanks with females.

We used photographs to identify males from the pre-treatment tests following the 45 days of
experimental treatment, and transferred them in a 3L aquarium three days prior to a second
dichotomous choice test (post-treatment test) to allow for sperm replenishment. This avoids biases
in preference measurements due to lack in motivation (Pilastro et al., 2002). While biases due to
motivation were only expected for experienced males, naïve males were likewise transferred to 3L
aquarium three days prior to their second behavioral test. Males were then presented for
dichotomous choice post-treatment tests.

We performed all behavioral tests in a circular arena (diameter = 47 cm) sheltered to prevent
disruption. We filmed the arena for 15-minute periods using a OBSBOT webcam (1080P at 30
fps) after a five-minute acclimatization period. For accurate identification of fish with tracking
software, we placed them in the experimental arena in 20 second intervals. We placed females first
in the arena, randomizing the order of placing receptive and nonreceptive females. Additionally,

to account for any olfactory cues, we changed water in the arena between tests. To minimize stress,

each fish was netted, placed in a glass bowl and transferred to the testing apparatus. For

consistency, the tests were always conducted in the morning for a period of 4-6 hours.

We used idTracker to track the position of males and females in video recordings (Perez-Escudero

et al., 2014) and to quantify the distance between the male and female for each video frame. To

calculate preference for receptive females, we defined the time that a male associated with each

female as the number of frames in the video recording that male was < 4 cm (less than two female

guppy average body lengths) to each female. Preference ratio was calculated as:

138
$$\frac{(time\ spent\ with\ receptive - time\ spent\ with\ nonreceptive)}{total\ amount\ of\ time\ spent\ with\ both\ females}$$

To evaluate differences in preferences for receptive females between experienced and naïve males,

we used a Linear Mixed Model with preference ratio as the dependent variable, and the time of

testing, experience to mating receptivity cues and the interaction of these as fixed factors. We

included the experimental batch as a random factor in the model. Given singularity issues caused

by low variance in batch effects, we performed an analogous linear model including batch as a

fixed effect. Significance tests were computed using a Wald t-distribution with Kenward-Roger

approximation using the parameters package (Lüdecke et al., 2020). All analyses were performed

in R (v. 4.1.3; R Core Team, 2022).

To assess potential differences in morphology or colouration patterns between fish used across

treatments, we extracted measurements from male photographs taken right after pre-treatment

tests. We quantified the number of pixels with carotenoid colouration, black melanic colouration,

body size (fish standard length) and tail size in the photographs using ImageJ (Schneider et al.,

2012). We used a linear model with each measurement as the dependent variable and social

treatments as fixed effect in R (v. 4.1.3; R Core Team, 2022).

155
156   *No choice preference tests*
157

To assess potential differences in preference for receptive females, as well as differences in the sexual behaviour repertoire of naïve and experienced males toward receptive females, we performed no-choice tests with receptive and non-receptive females on 122 males from the 31 experienced and 31 naïve aquaria, described above. The males not used in the dichotomous choice tests were presented to either a receptive or nonreceptive female, with 62 experienced males (n=31 with nonreceptive females and n= 31 with receptive females) and 60 naïve males (n=30 with nonreceptive females and n= 30 with receptive females). We excluded two naïve males due to uncertainty with fish labelling. All males were removed from treatment aquaria three days prior to experimentation for sperm replenishment, and to avoid biases in preference measurements due to lack in motivation (Pilastro et al., 2002).

Tests were conducted in a similar fashion to the dichotomous choice tests except only one female was present for each test, alongside a male, with the female placed first in the testing apparatus. Additionally, to account for any olfactory cues, we changed the water between each test. In order to minimize stress, each fish was netted, placed in a glass bowl and transferred to the testing apparatus. For consistency the tests were always conducted in the morning for 4 – 8 hours.

A single observer scored male sexual behaviour in video recordings in a random order and quantified the following behaviours as defined in Liley (1966): i) number of sigmoid displays, every time a male positioned himself in front of the female with an S-shaped posture soliciting copulation; ii) number of sneak attempts, unsolicited attempts of inseminating a female from behind by thrusting his gonopodium at the female's urogenital pore. We also calculated latency to initial sexual behaviour. Following the procedure described for dichotomous choice tests, we used idTracker software (Perez-Escudero et al., 2014), to calculate the distance between the male and female for each frame in the video recordings and extracted the time spent following a female for each trial (number of frames < 4 cm).

To compare the number of sigmoid displays in experienced and naïve males, we fit a statistical model using a Poisson distribution and a logit link function for the conditional mean in the package glmmTMB (Brooks et al., 2017). We used the mating status of the female, experience to mating receptivity cues and the interaction of these as fixed factors. We included the number of tank and

experimental batch as random factors in the model. For sneak attempts and latency to first sexual
behaviour, we used analogous models including a zero inflation linear predictor. For time
following the female, we used an analogous structure in a Linear Mixed Model fitted with lmer
package (Bates et al., 2007). We evaluated the adequacy of our fitted models using scaled-residuals
quantile-quantile plots, residual versus predicted values plots and a zero-inflation test in the
DHARMa package (Hartig, 2018). We processed the parameters of our statistical models using
Wald tests obtained via the parameters package (Lüdecke et al., 2020). We obtained post-hoc
comparisons of the male response between female receptivity levels at pre-treatment and post-
treatment time of testing in the previous models using the emmeans package with the tukey-
adjustment method for multiple comparisons (Lenth et al., 2019). All analyses were performed in
R (v. 4.1.3; R Core Team, 2022).
200

Results
201
202
Male colour and morphology analyses
Average proportion of orange or black coloration did not differ between males used for
experienced or naïve treatments (mean ± SE; orange coloration: naïve males 7.33 ± 3.47,
experienced males 7.14 ± 2.47, $F_{df=1}$ = 121, p = 0.729; black coloration: naïve males 2.27 ± 1.02,
experienced males 2.20 ± 0.86, $F_{df=1}$ = 0.154, p = 0.695; Supplementary Figure 1). Similarly, there
was no significant overall difference between naïve and experienced males in morphological traits
(body size: naïve males 1.55 ± 0.42, experienced males 1.57 ± 0.42, $F_{df=1}$ = 0.00, p = 0.99; tail
size: naïve males 0.51 ± 0.08, experienced males 0.48 ± 0.09, $F_{df=1}$ = 0.75, p = 0.38; Supplementary
Figure 1).
212

Dichotomous Choice Preference Tests
Neither experienced nor naïve males changed their strength of preference towards receptive
females after 45 days of treatment (time of testing: $estimate_{pre-treatment}$: -0.01 ± 0.04, t = -0.37, p =
0.72; Figure 1; Table1). In addition, we found no overall differences in the strength of preference
for receptive females between experienced and naïve males, or in the rate of change in preference
between experienced and naïve males following the 45 days of treatment (male social treatment:

estimate$_{experienced}$: 0.01 ± 0.04, t $= 0.45$, p $= 0.65$; male social treatment x time of testing : estimate$_{pre\text{-}}$
$_{treatment\ x\ experienced}$: 0.04 ± 0.06, t $= 0.72$, p $= 0.48$; Figure 1; Table 1).

No Choice Preference Tests

There was no significant difference between naïve and experienced males in their levels of display

behaviour or the number of displays that were performed towards receptive versus non-receptive

females (Figure 2a; Table 2). However, we found that, unlike naïve males, experienced males

significantly increased the number of displays towards receptive females compared to non-

receptive females (female status x social treatment: estimate$_{receptive\ x\ experienced}$: 0.26 ± 0.08, t $= 3.01$,

p $= 0.002$; Figure 2a; Table 2).

Experienced males exhibited significantly more sneak attempts than naïve males (male social

treatment: estimate$_{experienced}$: 0.49 ± 0.22, t $= 2.19$, p $= 0.028$; Figure 2b; Table 2), and this difference

was due to greater frequency of sneak attempts by experienced males toward non-receptive

females. We observed no difference in sneak frequency between experienced and naïve males in

tests with receptive females (log$_{mean \pm SE}$; non-receptive females: naïve males 1.16 ± 0.18,

experienced males 1.65 ± 0.14, t-ratio$_{df=119}$ $= -2.19$, p $= 0.030$; log$_{mean \pm SE}$; receptive females: naïve

males 1.14 ± 0.16, experienced males 1.16 ± 0.15, t-ratio$_{df=119}$ $= -0.07$, p $= 0.94$; Fig. 2b; Table 2).

Additionally, naïve and experienced males showed no significant difference in the time spent

following females in the trials, or in overall time spent following receptive and non-receptive

females (Figure 2c; Table 2).

Experienced males showed decreased latency to perform sexual behavior compared to naïve males

in tests with non-receptive and receptive females (male social treatment: estimate$_{experienced}$: $-1.51 \pm$

0.36, z $= -4.17$, p < 0.001; Figure 2d; Table 2) However, post-hoc tests indicate that this decrease

was significant just in tests with non-receptive females (log$_{mean \pm SE}$; non-receptive females: naïve

males 4.28 ± 0.26, experienced males 2.76 ± 0.25, t-ratio$_{df=115}$ $= 4.17$, p < 0.001; mean $\pm$ SE;

receptive females: naïve males 3.75 ± 0.26, experienced males 3.23 ± 0.25, t-ratio$_{df=115}$ $= 1.44$, p $=$

0.15; Fig. 2d; Table 2).

248
249   **Discussion**
250

We used dichotomous choice and no-choice tests to investigate how previous experience with
female receptivity cues alter guppy male sexual behaviour and strength of preference for female
mating status. Our results showed that males with previous access to female receptivity cues
exhibited significantly greater frequency of coercive sexual behaviours and lower latency to first
sexual behaviour to non-receptive females than naïve males. In addition, only experienced males
significantly increased their number of displays towards receptive females compared to the number
of displays performed with non-receptive females. However, previous experience with receptivity
cues did not affect the strength of guppy male preference for female mating status.
259
Previous studies evaluating how female mating status affect male mating behavior showed higher
levels of coercive copulation attempts towards non-receptive females and higher levels of sigmoid
displays towards receptive females (Guevara-Fiore et al., 2010a;2010b). Our results match these
patterns only in males with previous experiences with female receptivity cues, suggesting a key
role of social learning driving preferences for high quality females in this species. Our
experimental design does not allow to disentangle which mechanism leads to changes in behaviour
between experienced and naïve males. Prior experience with females might lead to males better
recognizing which mating tactics provide higher success, as previously observed in species such
as *Drosphila melanogaster* (Dukas, 2005, Saleem et al., 2014, Balaban-Feld & Valone., 2017), or
eastern mosquitofish (Bisazza et al., 1996 but see Iglesias-Carrasco et al., 2019). Alternatively,
changes in encounter rates of females and in mating success artificially created by the two social
environments used in our experimental setup are known to affect male mating tactics (Cattelan et
al., 2016; Devigili et al., 2015; Jordan & Brooks, 2012, Guevara-Fiore & Endler, 2018).
Ultimately, the behavioural patterns observed in males with access to female receptivity cues
correspond to theoretical predictions of fitness maximization, once accounting for the lower
energetic requirements of sneak attempts of sperm insemination in relation to more costly sigmoid
displays aiming to engage female with sexual consent (Devigli et al., 2013, Head et al., 2010).
277
Contrary to our prediction, males with social learning experience of female receptivity cues did
not became choosier or increase their preference towards receptive females in dichotomous choice
tests. It may be that that our experimental treatment might have changed the perception of naïve
and experienced males in future reproductive opportunities, potentially biasing the investment of

naïve males in sexual behaviors that we observed in the first sexual encounter of their pre-treatment

test (Fischer et al., 2008, Aich et al., 2021). Furthermore, determining the costs of sexual

behaviours are challenging in benign lab environments where food is not a limiting factor. Future

work incorporating resource limitation will be helpful to determine ecologically-relevant effects

of social learning in male mating preferences. Yet, it is important to note that our results are

concordant with a recent meta-analysis showing no evidence that mating status is an important

factor for preference. Specifically, across species, virgin females are as choosy as mated females

across reproductive isolation, inbreeding avoidance, and sexually transmitted disease scenarios

(Richardson & Zuk, 2022). Our study here presents a similar finding, as males with no previous

mating experience (naïve males) presented similar preferences to males with mating experience

(experienced males). Hence, our study focusing on male mate choice on preference for higher

quality females is in broad agreement to previous observations on the role of mating status for

female preference across species.

Using both dichotomous and no-choice approaches allowed for a broader picture of male

preference in the context of social learning. It is possible that mating preferences may be stronger

in choice tests compared to no-choice design, as males can select the female that is more likely to

result in insemination (Dougherty, 2020). However, there is arguably an increased risk of being

rejected by the only potential mate in a no-choice tests, and this could make males more careful in

tuning their mate strategy to female receptivity cues (Dougherty & Shuker, 2015). Our tests using

these two complementary experimental paradigms are concordant in that we did not observe

changes in overall preference for receptive females or differences in overall sexual behaviour

levels based on female receptivity status. However, our observations of changes in rates of sexual

display and coercive copulations depending on female mating status suggest that future mate

choice studies should incorporate both methodologies.

The changes we observe in male guppies in behavioural repertoire and latency towards females

based on previous experience with female receptivity cues add to the evidence suggesting that

social environment and learning from previous experience can affect male sexual behaviour.

Overall, our results are consistent with the idea that male guppies use social learning to efficiently

tune their mating tactics, soliciting copulation in higher rates to receptive females and performing

higher coercive copulation attempts towards non-receptive females.

Acknowledgements

We thank Wouter van der Bijl for help with photography and fish care, as well as Lydia Fong,

Yuying Lin and Jacelyn Shu for help with fish care. **Funding**: We gratefully acknowledge funding

from a Canada 150 Research Chair and NSERC (J.E.M). **Author contributions**: J.E.M. and AC-

L. conceptualized the study. V.G. and I.D. performed research. A.C-L and V.G. performed formal

analyses and visualization. J.E.M. contributed to funding acquisition of the project. A.C-L., J.E.M.,

and V.G. wrote the original draft. All authors contributed to the final version of the manuscript.

References

Aich U, Bonnet T, Head ML, Jennions MD. 2021. Disentangling the effects of male age and
mating history: Contrasting effects of mating history on precopulatory mating behavior and
paternity success. *Evolution*, *75*(11), 2867-2880.

Akinyemi AO, Kirk WD. 2019. Experienced males recognise and avoid mating with non-virgin
females in the western flower thrips. *PLoS One*, *14*(10), e0224115.

Andersson M, Iwasa Y. 1996. Sexual selection. Trends *Ecology Evoloution*.11(2):53-8.

Bailey NW, Moore AJ. 2012. Runaway sexual selection without genetic correlations: social
environments and flexible mate choice initiate and enhance the Fisher process. *Evolution;*
International Journal of Organic Evolution, *66*(9), 2674–2684.

Bailey NW, Gray B, Zuk M. 2010. Acoustic experience shapes alternative mating tactics and
reproductive investment in male field crickets. *Current Biology*, *20*(9), 845-849.

Balaban-Feld J, Valone TJ. 2017. Prior information and social experience influence male
reproductive decisions. *Behavioral Ecology*, *28*(5), 1376-1383.

Bates D, Sarkar D, Bates MD, Matrix L. 2007. The lme4 Package. *R package version* **2**, 74.

Bisazza A, Marconato A, Marin G. 1989. Male mate preferences in the mosquitofish *Gambusia*
holbrooki. *Ethology,* 1989;83:335–343.

Bisazza A, Pilastro A, Palazzi R, Marin G. 1996. Sexual behaviour of immature male eastern
mosquitofish: a way to measure intensity of intra-sexual selection? *Journal of Fish*
Biology, *48*(4), 726-737.

Bonduriansky R. 2001. The evolution of male mate choice in insects: a synthesis of ideas and
evidence. *Biological Reviews*, *76*(3), 305-339.

Brooks ME, Kristensen K, Van Benthem KJ, Magnusson A, Berg CW, Nielsen A, ... & Bolker
BM. 2017. glmmTMB balances speed and flexibility among packages for zero-inflated
generalized linear mixed modeling. *The R journal*, *9*(2), 378-400.

Cattelan S, Evans JP, Pilastro A, Gasparini C. 2016. The effect of sperm production and mate
availability on patterns of alternative mating tactics in the guppy. *Animal behaviour*, *112*, 105-
110.

Corral-López A, Kotrschal A, Kolm N. 2018. Selection for relative brain size affects context-
dependent male preference for, but not discrimination of, female body size in guppies. *Journal of
Experimental Biology*, *221*(12), jeb175240.

Danchin E, Giraldeau LA, Valone TJ, Wagner RH. 2004. Public information: from nosy
neighbors to cultural evolution. *Science*, 305: 487–491.

Darwin CR. 1871. The Descent of Man, and Selection in Relation to Sex. John Murray, London.

Devigili A, Doldan-Martelli V, Pilastro A. 2015. Exploring simultaneous allocation to mating
effort, sperm production, and body growth in male guppies. *Behavioral Ecology*, 26(4),
1203e1211.

Devigili A, Kelley JL, Pilastro A, Evans JP. 2013. Expression of pre- and postcopulatory traits
under different dietary conditions in guppies. *Behavioral Ecology*, 24(3), 740e749.

Dosen LD, Montgomerie R. 2004. Female size influences mate preferences of male
guppies. *Ethology*, *110*(3), 245-255.

Dougherty LR. 2020. Designing mate choice experiments. *Biological Reviews*, *95*(3), 759–781.

Dougherty LR, Shuker DM. 2015. The effect of experimental design on the measurement of
mate choice: a meta-analysis. *Behavioral Ecology*, *26*(2), 311-319.

Dukas R. 2005. Experience improves courtship in male fruit flies. *Animal Behaviour*, *69*(5),
1203-1209.

Edward DA, Chapman T. 2011. The evolution and significance of male mate choice. *Trends in
Ecology & Evolution*, *26*(12), 647–654.

Fischer K, Perlick J, Galetz T. 2008. Residual reproductive value and male mating success: older
males do better. *Proceedings of the Royal Society B: Biological Sciences*, *275*(1642), 1517-1524.

Fowler-Finn KD, Rodríguez RL. 2012. The evolution of experience-mediated plasticity in mate
preferences. *Journal of Evolutionary Biology*, *25*(9), 1855–1863.

Guevara-Fiore P. 2012. Early social experience significantly affects sexual behaviour in male
guppies. *Animal Behaviour*, *84*(1), 191–195.

Guevara-Fiore P, Endler JA. 2018. Female receptivity affects subsequent mating effort and mate
choice in male guppies. *Animal Behaviour*, *140*, 73–79.

Guevara-Fiore P, Skinner A, Watt PJ. 2009. Do male guppies distinguish virgin females from
recently mated ones? *Animal Behaviour*, 77, 425-431.

Guevara-Fiore P, Stapley J, Watt PJ. 2010. Mating effort and female receptivity: How do male
guppies decide when to invest in sex? *Behavioral Ecology and Sociobiology*, 64(10), 1665e1672.

Gumm JM, Gabor CR. 2005. Asexuals looking for sex: conflict between species and mate-
quality recognition in sailfin mollies *Poecilia latipinna*. *Behavioral Ecology and*
Sociobiology. 2005;58:558–565.

Hartig F. 2018. DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression
models. *R package version 0.3*.

Head ML, Jacomb F, Vega-Trejo R, Jennions MD. 2015. Male mate choice and insemination
success under simultaneous versus sequential choice conditions. *Animal Behaviour* 103: 99–105.

Head ML, Wong BB, Brooks R. 2010. Sexual display and mate choice in an energetically costly
environment. *PLoS One*, *5*(12), e15279.

Herdman EJE, Kelly CD, Godin JGJ. 2004. Male Mate Choice in the Guppy (Poecilia reticulata):
Do Males Prefer Larger Females as Mates? *Ethology*, *110*(2), 97–111.

Houde AE, 1997. Sex, color, and mate choice in guppies. Princeton, NJ: Princeton University
Press.

Hoysak DJ, Godin JGJ. 2007. Repeatability of male mate choice in the mosquitofish, *Gambusia*
holbrooki. *Ethology.* 2007;113:1007–1018.

Iglesias-Carrasco M, Fox RJ, Vincent A, Head ML, Jennions MD. 2019. No evidence that male
sexual experience increases mating success in a coercive mating system. *Animal Behaviour*, *150*,
201-208.

Jeswiet SB, Lee-Jenkins SS, Godin JGJ. 2012. Concurrent effects of sperm competition and
female quality on male mate choice in the Trinidadian guppy (Poecilia reticulata). *Behavioral*
Ecology, *23*(1), 195-200.

Jordan LA, Brooks RC. 2012. Recent social history alters male courtship preferences. *Evolution*,
66(1), 280e287.

Jordan LA, Kokko H, Kasumovic M. 2014. Reproductive foragers: male spiders choose mates by
selecting among competitive environments. *The American Naturalist*, *183*(5), 638-649.

Lenth R, Singmann H, Love J, Buerkner P, Herve M. 2019. Package 'emmeans'. R. Package
Version, 1

Liley NR. 1966. Ethological Isolating Mechanisms in Four Sympatric Species of Poeciliid
Fishes. *Behaviour. Supplement*, *14*, III–197.
413
414   Lüdecke D, Ben-Shachar MS, Patil I, Makowski D. 2020. Extracting, computing and exploring
415   the parameters of statistical models using R. *Journal of Open Source Software*, *5*(53), 2445.
416
Macario A, Croft DP, Endler JA, Darden SK. 2017. Early social experience shapes female mate
choice in guppies. *Behavioral Ecology*, *28*(3), 833-843.

Pérez-Escudero A, Vicente-Page J, Hinz R *et al.* 2014. idTracker: tracking individuals in a
group by automatic identification of unmarked animals. *Nat Methods* **11**, 743–748.

Pilastro A, Evans JP, Sartorelli S, Bisazza A. 2002. Male phenotype predicts insemination
success in guppies. *Proceedings of the Royal Society B: Biological Sciences*, *269*(1498), 1325.

Plath M, Seggel U, Burmeister H, Heubel KU, Schlupp I. 2006. Choosy males from the
underground: male mating preferences in surface- and cave-dwelling Atlantic mollies *Poecilia*
mexicana. *Naturwissenschaften.* 2006;93:103–109.

R Core Team, 2022. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Rather PA, Herzog AE, Ernst DA, Westerman EL. 2022. Effect of experience on mating
behaviour in male Heliconius melpomene butterflies. *Animal Behaviour*, *183*, 139-149.

Řežucha R, Reichard M. 2014. The effect of social environment on alternative mating tactics in
male Endler's guppy, Poecilia wingei. *Animal Behaviour*, *88*, 195-202.

Richardson J, Zuk M. 2022. Unlike a virgin: a meta-analytical review of female mating status in
studies of female mate choice. *Behavioral Ecology*, 1–18.

Rosenqvist G. 1990. Male mate choice and female-female competition for mates in the pipefish
Nerophis ophidion. *Animal Behaviour*, *39*(6), 1110–1115.

Rosenqvist G, Houde A. 1997. Prior exposure to male phenotypes influences mate choice in the
guppy, Poecilia reticulata. *Behavioral Ecology*, *8*(2), 194-198.

Saleem S, Ruggles PH, Abbott WK, Carney GE. 2014. Sexual experience enhances Drosophila
melanogaster male mating behavior and success. *PloS one*, *9*(5), e96639.

Schlupp I. 2018. Male mate choice in livebearing fishes: an overview. *Current Zoology*, *64*(3),
393–403.

Schneider C, Rasband W, Eliceiri K. 2012. NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* **9**, 671–675.

Valone TJ, Templeton JJ. 2002. Public information for the assessment of quality: a widespread
social phenomenon. *Philosophical Transactions of the Royal Society B: Biological Sciences.* 357:
1549–1557.

Werner NY, Lotem A. 2003. Choosy males in a haplochromine cichlid: first experimental
evidence for male mate choice in a lekking species. *Animal Behaviour*, *66*(2), 293–298.

Wong BBM, Jennions MD. 2003. Costs influence male mate choice in a freshwater fish.
Proceedings. Biological Sciences, *270 Suppl 1*(Suppl 1).

453
Figure 1. The effect of previous experience with female receptivity cues in guppy male
preference for receptive females. Preference ratios were calculated as total time spent associating
with receptive females by the total time associating with receptive and non-receptive females in
dichotomous choice tests. Tests were performed to males naïve and experienced with female
receptivity cues before (pre-treatment, black circles) and after (post-treatment, gray squares) a 45-
day treatment in their respective experimental condition. Larger circles and squares indicate
average preference ratio for each treatment and time of testing with 95% CI bars. We found no
differences in the amount spent with receptive or nonreceptive females between naïve and
experienced males in pre-treatment or post-treatment tests (see Table 1).
463
464
465

466
Figure 2. The effect of previous experience with female receptivity cues in male sexual
behaviour. Quantification of (a) the number of sigmoid displays, (b) number of sneak attempts,
(c) total time spent following females, and (d) latency to first sexual behaviour performed towards
non-receptive and receptive females in guppy males naïve (black circles) and experienced (gray
squares) with female receptivity cues. Larger circles and squares indicate average values of each
behaviours with 95% CI bars. For number of displays, experienced males significantly increased
the number of displays towards receptive females compared to non-receptive females (female
status x social treatment: p = 0.002; see Table 2). Stars indicate significance in post-hoc
comparisons of the male response between female receptivity levels at pre-treatment and post-
treatment time of testing (p < 0.001 ***, p < 0.05 *).

Table 1. Results for a Linear Mixed Model comparing preference for receptive females performed to male guppies naïve and experienced with female receptivity cues before and after a 45-day treatment in their respective experimental condition.

Parameter	Estimate	SE	t (df:119)	P – value
Intercept	-0.02	0.03	-0.88	0.60
Male social treatment (experienced)	0.01	0.04	0.45	0.65
Time of testing (pre-treatment	-0.01	0.04	-0.37	0.72
Batch (2)	-0.02	0.03	-0.62	0.53
Male social treatment:time of testing	0.04	0.06	0.72	0.48

521 **Table 2.** Statistical tests for models comparing potential differences in behaviour in no choice tests
522 with non-receptive and receptive females performed to male guppies naïve and experienced with
523 female receptivity cues. Parameters with significant differences in bold.
524

Behaviour	Parameter	Estimate	SE	Statistic$_{df}$	P - value
Display	Intercept	2.57	0.20	z : 12.70	**< 0.001**
	Female status (receptive)	-0.05	0.06	z : -0.85	0.39
	Male social treatment (experienced)	-0.05	0.17	z : -0.28	0.77
	Female status:male social treatment	0.26	0.08	z : 3.01	**0.002**
Sneak	Intercept	1.15	0.17	z : 6.48	**< 0.001**
	Female status (receptive)	-0.01	0.20	z : -0.07	0.94
	Male social treatment (experienced)	0.49	0.22	z : 2.19	**0.028**
	Female status:male social treatment	-0.47	0.26	z : -1.79	0.07
	Zero Inflation: Intercept	0.49	0.10	z : -3.37	**<0.001**
Latency	Intercept	4.27	0.26	z : 16.42	**< 0.001**
	Female status (receptive)	-0.52	0.36	z : -1.42	0.15
	Male social treatment (experienced)	-1.51	0.36	z : -4.17	**< 0.001**
	Female status:male social treatment	0.99	0.51	z: 1.93	0.053
	Zero Inflation: Intercept	0.00	0.00	z : -0.00	0.996
Time spent following	Intercept	351.27	44.13	t$_{3.7}$: 7.95	**0.002**
	Female status (receptive)	24.51	42.85	t$_{61.5}$: 0.57	0.56
	Male social treatment (experienced)	13.70	51.52	t$_{105.5}$: 0.26	0.79
	Female status:male social treatment	-61.06	59.30	t$_{-61.8}$: -1.03	0.31

Supplementary Materials for:

Male guppies (*Poecilia reticulata*) exposed to mating receptivity cues change their sexual behaviour, but not their preference, towards receptive females

Versara Goberdhan, Iulia Darolti, Judith E. Mank, Alberto Corral-Lopez*

*corresponding author: corral@zoology.ubc.ca

Table S1. Results of linear model comparing colouration and morphological traits in male guppies used for social experience experimental treatments

Trait	Sum Sq	Mean Sq	F value (df = 1)	P value
Orange Colouration	1.1	1.09	0.12	0.72
Black Colouration	0.14	0.13	0.15	0.69
Standard Length (cm)	0.00	0.00	0.00	0.93
Tail Length (cm)	0.012	0.01	3.08	0.08

Figure. S1. **Body morphology and colouration in experienced and naïve males**. No significant differences were found between randomly assigned males to naïve (n = 31) and experienced (n = 31) treatments for (a) tail length, (b) body length, (c) proportion of black, or (d) proportion of orange. For all boxplots, horizontal lines indicate medians, boxes indicate the interquartile range, and whiskers indicate all points within 1.5 times the interquartile range.