Structural stability of the oyster gut microbiome contributes to the host's physiological tolerance to temperature fluctuations on tropical rocky shores

Bovern Suchart Arromrak[1]; Adrian Tsz Chun Wong[1]; Tin Yan Hui[1]; Kin Sum Leung[1]; Gray A. Williams[1]; Monthon Ganmanee[2]; Thierry Durand[3]; Jetty Chung Yung Lee[4]; & Juan D. Gaitan-Espitia[1*]

[1] The Swire Institute of Marine Science and School of Biological Sciences, The University of Hong Kong, Hong Kong, SAR, China.

[2] Department of Animal Production Technology and Fisheries, Faculty of Agricultural Technology, King Mongkut's Institute of Technology Ladkrabang, Bangkok 10520, Thailand

[3] Institut des Biomolécules Max Mousseron (IBMM), Pôle Chimie Balard Recherche, UMR 5247, Université de Montpellier, CNRS, ENSCM, F-34293 Montpellier Cedex 05, France

[4] School of Biological Sciences, The University of Hong Kong, Hong Kong, SAR, China

* Correspondence: juadiegaitan@gmail.com

Abstract

Extreme thermal conditions on rocky shores are challenging to the survival of intertidal ectotherms, especially during emersion periods. Yet, many species are highly successful in these environments in part due to their ability to regulate intrinsic mechanisms associated with physiological stress and their metabolic demands. More recently, there has been a growing awareness that other extrinsic mechanisms, such as animal-associated microbial communities, can also influence the tolerance and survival of ectotherms under stressful conditions. The extent to which intrinsic and extrinsic mechanisms are functionally linked as part of the overall adaptive response of intertidal animals to temperature change and stress is, however, poorly understood. Here, we examined the dynamics and potential interactions of intrinsic and extrinsic mechanisms in the ecology of the tropical high shore rock oyster, *Isognomon nucleus*. We found that oysters modulate their internal biochemistry (PUFA-oxidised metabolites including $5\text{-}F_{2t}\text{-}IsoP$, $10\text{-}F_{4t}\text{-}NeuroP$, $13\text{-}F_{4t}\text{-}NeuroP$, and $16\text{-}F_{1t}\text{-}PhytoP$) as part of their adaptive regulation to cope with physiological stress during periods of extreme temperatures when emersed. While we detected extrinsic microbiome changes in alpha diversity, the overall taxonomic and functional structure of the microbiome showed temporal stability with no association with the host biochemical profiles. Our finding here suggests that the microbiome taxonomic and functional structure is maintained by a stable host control (not associated to the host biochemistry) and/or that the microbiome (independent of the host) is resilient to the temperature fluctuations and extremes. This microbiome stability is likely to contribute to the oyster, *I. nucleus* thermal tolerance, in addition to the intrinsic biochemical adjustment, to survive in the thermally challenging intertidal environment.

Introduction

The tropical rocky shore is one of the most extreme environments on Earth. This system is characterised by a marked heterogeneous thermal landscape influenced by tidal dynamics (Helmuth and Hofmann, 2001; Raffaelli and Hawkins, 1999). Across this vertical gradient defined by the tides, environmental temperatures shape the physiology, behaviour, and spatial distribution of tropical intertidal ectotherms (Chow, 2004; Garrity, 1984; Giomi et al., 2016; Ng et al., 2017; Samakraman et al., 2010; Williams and Morritt, 1995). For ectotherms occupying the high shore habitat, thermal and oxidative stress (i.e., excessive production of reactive oxygen species - ROS) during desiccation periods of extended emersion, are the two major physiological challenges faced (Birben et al., 2012; Freire et al., 2011; Sokolova and

Pörtner, 2001). To survive under these challenging conditions, high shore intertidal ectotherms

modulate intrinsic mechanisms associated with their internal physiology (e.g., metabolic rate)

and biochemical activities. This modulation can be observed through, for instance, the

regulation of ROS level via polyunsaturated fatty acids (PUFA; Letendre et al., 2009; Sokolova

and Pörtner, 2001; Wang et al., 2018; Yin et al., 2017), and/or the control of the gene expression

associated with heat shock response (HSR) and oxidative stress response (OSR; Wang et al.,

2022). In addition to that, some of these organisms are able to enter into a hypometabolic state

(i.e., metabolic depression; Hui et al., 2020; Marshall and McQuaid, 2011) to avoid excessive

energy consumption when exposed to high temperatures during low tides. Together, these

functional characteristics are regarded as important, adaptive traits for intertidal ectotherms to

cope and ameliorate the extreme stress experienced on tropical rocky shores.

Survival under stressful environmental conditions is also hypothesized to be influenced

by extrinsic mechanisms such as host-associated microbial communities (i.e., microbiome;

Ahmed et al., 2019; Bang et al., 2018; Fan et al., 2013). The structure and functions of these

microbiomes have been suggested to play important roles in shaping tolerances and

biochemical regulation of animals to elevated temperatures (Herrera et al., 2020; Moghadam et

al., 2018; Tignat-Perrier et al., 2022). Under such conditions, functional microbiomes that are

resilient/responsive to environmental changes are fundamental to the host's health and survival

(Klepac et al., 2015; Kremer et al., 2018; Muñoz et al., 2019; Soen, 2014). For instance, Epstein

et al. (2019) demonstrated that the resilience of the coral, *Pocillopora acuta,* to a stressful

thermal event may be attributed to the stability of the microbiome composition and structure.

In contrast, a dysbiotic host state is commonly associated with a breakdown in microbiome

stability (Fan et al., 2013; McDevitt-Irwin et al., 2019). This is typically reflected by a marked

variation among host-associated microbiomes, a signal indicating the host losing control of the

microbial community and/or differential sensitivities of microbiomes to extreme disturbances

(Carey and Duddleston, 2014; Lokmer and Mathias Wegner, 2015; Zhou et al., 2014). This

pattern is in-line with the "Anna Karenina Principle", which states, "all happy families are all

alike; each unhappy family is unhappy in its own way" (Díaz-almeyda et al., 2022; Pita et al.,

2018; Zaneveld et al., 2017). In other words, stressed organisms may host more stochastic

microbiomes than healthy organisms. Although microbiome stability seems important for the

correct functioning of the host, some studies have suggested that flexibility in the microbiome

can also provide adaptive capacities to the host, allowing it to cope with rapidly changing

environments (Maher et al., 2020; Voolstra and Ziegler, 2020). Ziegler et al. (2017), for

example, showed that the coral *Acropora hyacinthus* exhibits changes in its microbiome

composition and structure that are aligned with the local thermal environmental conditions.
Similarly, several studies have also shown seasonal variation in the host-associated microbiome
of many marine organisms such as bivalves and echinoderms, as a potential plastic/flexible
response to environmental change (Bäckhed et al., 2015; Feng et al., 2021; Pierce and Evan,
2019). Microbiome stability or flexibility may, therefore, be a sign of host adaptive responses
to environmental changes. However, the influence of these contrasting patterns is context-
dependent and thus, their contribution to the tolerances and survival of animal hosts may differ
in extreme environments such as tropical rocky shores.
Previous studies have shown that the capacity for plastic adaptive responses of animal
hosts to environmental stress may be modulated by the interactions between intrinsic
(physiology and biochemical changes of the host) and extrinsic (host's microbiome structure
and function) mechanisms (Fontaine and Kohl, 2022; Marangon et al., 2021; Muñoz et al., 2019;
Walke et al., 2021). It is unclear, however, to what extent these mechanisms are functionally
linked as part of the whole organism's adaptive response to survive in an extreme environment
such as the tropical intertidal rocky shore. In this study, we investigated the intrinsic and
extrinsic mechanisms underpinning tolerances and survival of a tropical intertidal ectotherm
(oyster *Isognomon nucleus*) by assessing the influence of natural variation in environmental
temperature on i) host physiology (i.e., biochemical dynamics) ii) the diversity, structure, and
function of the host-associated microbiome and iii) the interplay between these mechanisms.
The integration of intrinsic and extrinsic mechanisms offers a holobiont perspective for
understanding the stress tolerance regulation of ectotherms in extreme and fluctuating
environments (Alberdi et al., 2016; Apprill, 2017; Bang et al., 2018).
121

Material and Methods

Site and species descriptions

Host-microbiome interactions were investigated in the high shore intertidal oyster,
Isognomon nucleus (Bivalvia: Isognomonidae). Animals were collected from Ko Sichang,
Thailand (latitude: 13°08'52''N; longitude: 100°48'11''E, (Samakraman et al., 2010). In this
area, *I. nucleus* is abundant and forms a distinctive band at the supra-littoral region (~2.5 - 3 m
above the mean sea level; Samakraman et al., 2010), where rock temperatures vary daily
(emersion period) from ~30°C at dawn to ~50°C at noon (hot season: March to June; Hui et al.,
2020). The oyster exhibits strong variations in physiological and metabolic adjustments with
elevated body temperatures, providing an ideal system to investigate the functional role of the

microbiome to facilitate the host's survival in this thermally extreme and variable environment

(Giomi et al., 2016; Hui et al., 2020).

Sample collection and processing

136   A total of 80 individuals of *I. nucleus* were randomly selected at four different

timepoints within an emersion period (Fig. 1A; 20 animals at each time) with contrasting

thermal conditions (09:15 hr, 12:30 hr, 16:12 hr, and 17:40 hr; hereafter, morning, noon, early-

afternoon, and late-afternoon, respectively; Table S1). This cycle represents major changes in

the environmental temperature as a result of the tidal dynamics and air/sun exposure (Hui et al.,

2020). Body temperatures were measured from 10 randomly selected individuals at each

timepoint using a digital thermometer ($\pm$ 0.1 $^{\circ}$C, TM-947SD, Lutron, Taiwan) where a fine-

tipped type-K thermocouple was inserted into the bodies through the valve opening. At each

time point, the whole tissue of five collected oysters was quickly removed at the site with a

sterile dissecting kit and snap-frozen in liquid nitrogen on shore for oxidative stress

quantification (as a proxy for host physiological and biochemical dynamics). The guts of

another five individuals were also dissected at each timepoint on-site, snap frozen in liquid

nitrogen for gut microbiome assessment. All collected samples (20 for oxidative stress and 20

for microbiome analysis across four time points) were stored in a portable liquid nitrogen tank

and transported to The University of Hong Kong within three days, where they were held in a

-80°C freezer until processing and analyses.

Rock surface temperature and tidal dynamics

154   To quantify environmental conditions experienced by *Isognomon nucleus* throughout

the tidal cycle, rock surface temperatures and tidal levels were recorded. To measure and track

surface rock temperature, i-Button loggers (DS1922L, Maxim, US) were firstly embedded in

waterproofing resin (Scotchcast, 3M, US), placed in plastic Falcon tube caps and affixed, using

marine epoxy (A-788, Z-Spar, US; Ng et al., 2021), onto the shore at the height where *I. nucleus*

were collected. Rock surface temperatures were logged by these i-Buttons every 5 minutes at a

resolution of 0.5 °C (n = 5). Additionally, a time-lapse camera (TLC200Pro, Brinno, Taiwan)

was set adjacent to the study site to monitor the immersion state of the *I. nucleus* band during

the day ($\sim$ 05:00 hr to 18:00 hr) to assess the tidal states. Images were captured every 10 minutes

and the tidal level was scored according to whether the band was underwater (immersed),

exposed to air (emersed), or being splashed by waves during transitions from
immersion/emersion (awash).
166
167 *Determination of polyunsaturated fatty acids-oxidised metabolites*
168
The levels of oxidative stress (i.e., associated with ROS production) of the oysters were
determined based on the concentration of non-enzymatic polyunsaturated fatty acids (PUFA)-
oxidised metabolites (hereafter, PUFA-oxidised metabolites; Basu, 2010; Durand et al., 2009;
Leung et al., 2015). A modified Folch extraction method was used to extract the PUFA-oxidised
metabolites from the oysters sampled (n=5) at each timepoint (t=4) (Dupuy et al., 2016; Folch
et al., 1957). In brief, 0.05g of oyster flesh was suspended in a 5 ml ice-cold Folch solution
(chloroform: methanol, 2:1, v/v) with 0.01% BHT (w/v) and homogenized at 24,000 rpm in 20
secs bursts on ice using a polytron benchtop homogenizer (T25, Ultra-Turrax, IKA, Germany).
Homogenization was performed twice, and the two parts of the Folch solution extracts were
pooled. The pooled solutions were added with 2 ml of 0.9% NaCl to introduce phase separation.
The mixture was then shaken on ice for 30 min and centrifuged at $2000 \times g$ for 10 min at room
temperature. The lower chloroform phase was transferred into a 30 ml glass vial. The extracted
samples were then added with potassium hydroxide (1M in methanol) and 1X PBS (pH7.4).
Samples were then shaken overnight in the dark at room temperature for alkaline hydrolysis.
The reaction was terminated by adding 1M of hydrochloric acid, methanol, 20 mM of formic
acid, and 40 mM of formic acid. Deuterated internal standards (5ng prepared in methanol) were
also added after the hydrolysis. Purification of the oxidised PUFA metabolites was performed
with the mixed anion solid phase extraction (MAX SPE, Waters, Milford, MA USA). Before
the start of the solid phase extraction, the SPE column was preconditioned with methanol and
20 mM formic acid. One sample was then loaded into a column and cleaned with 2%
ammonium hydroxide and 20 mM formic acid. The final eluent was collected with
hexane/ethanol/acetic acid (70/29.4/0.6, v/v/v). All the samples were then placed on a heating
block and heated at 37°C under a stream of nitrogen gas until they were completely dried. The
dried samples were then re-suspended in 100 µL methanol and cleaned through a 0.45 µm PTFE
filter to remove insoluble impurities, and then immediately analysed by liquid chromatography-
tandem mass spectrometry (LC-MS/MS).
In our study, we used the Sciex X500R QTOF LC-MS/MS system (Sciex Applied
Biosystems, MA, USA) which consisted of an Exion LC liquid chromatography with a C18
column (150×2.1 mm, 2.6 µm particle size, Phenomenex, Torrance, CA, USA) which was

maintained at 40°C for LC-MS/MS analysis. A mobile phase consisting of 0.1% aqueous acetic

acid (solvent A) and 0.1% acetic acid/methanol (solvent B) was used. The flow rate was set to

300 µl/min and the injection volume to 10 µl. For each sample analysis, the gradient condition

was maintained at 20% of solvent B for 2 min and between 20% and 98% for 8 min and then

held for 5 min. Then, the percentage of solvent B was reduced to 20% in 1 min and held for an

additional 5 min to equilibrate to the initial conditions. The X500R QTOF system was operated

in negative electrospray ionization (ESI) mode. The spray voltage was set to -4500V and

nitrogen was used as the curtain gas. The ionization chamber temperature was set at 350°C, and

the pressure of the ion source gases 1 and 2 were 35 and 45 psi, respectively. The declustering

potential (DP) was set to -80V and the collision energy (CE) was -10V for the QTOF MS. The

scan mode was set to multiple reaction monitoring (MRM). All data collected by the X500R

QTOF system was analyzed by the Sciex operating system (version 1.2.0.4122). The

quantification of each analyte was determined by correlating the peak area to its corresponding

deuterated internal standard peak. For analytes without corresponding deuterated internal

standards, the following deuterated internal standards, 5(S)-HETE-d_8, 15-F_{2t}-IsoP-d_4, and 4-F_{4t}-

NeuroP-d_4 were used for quantification. We identified fifteen PUFA-oxidised metabolites that

are known to be the products of non-enzymatic lipid peroxidation, based on previous work by

Dupuy et al.(2016), Galano et al.(2017) and Lee et al.(2018). These metabolites were derived

from the oxidation of the following PUFA: (1) Arachidonic acid (ARA) (2) Adrenic acid (AdA)

(3) Docosahexaenoic acid (DHA) (4) α-Linolenic acid (ALA) (5) Eicosapentaenoic acid (EPA).

DNA extraction and 16S rRNA gene sequencing

Total genomic DNA was extracted from the oysters' guts (~0.03g) using the commercial

DNA extraction kit, Dneasy Power Soil Kit (Qiagen GmbH, Hilden, Germany), following the

manufacturers protocol with some modifications (i.e., the solution mixture was incubated at 55°C

in a water bath for 30 minutes after the bead-beating step). The extracted DNA (100µl) was

then collected in a microcentrifuge tube and preserved in a -80°C freezer. All the DNA samples

were then sent to Ramaciotti Centre for Genomics at the University of New South Wales,

Sydney Australia for 16S rRNA amplicon paired-end sequencing (2X 300bp) on an Illumina

Miseq Platform. Before the amplicon sequencing, the V3-V4 variable region of the 16S rRNA

gene from each DNA sample was amplified using the following primers: (1) 341F (Forward

Primer);5'- CCTACGGGNGGCWGCAG-3' (2) 805R (Reverse Primer); 5'-

GACTACHVGGGTATCTAATCC-3'(Herlemann et al., 2011; Klindworth et al., 2013). Raw

reads received were the demultiplexed paired-end sequences in a FASTQ format and they were

deposited in NCBI Sequence Read Archive (SRA) under Bioproject number PRJNA931123.

The data will be made available once published.

16S rRNA gene sequence processing

Raw demultiplexed paired-end sequences were processed following the Quantitative

Insights into Microbial Ecology (QIIME2 ver. 2021.4) pipeline (Bolyen et al., 2019). Briefly,

the raw data was first visualized with the demux plug-in function for quality-checking and

identifying poor-quality base regions to filter. The sequences were then trimmed, filtered,

denoised, chimeras-removed, and merged using the DADA2 plug-in function in QIIME2

(Callahan et al., 2016). The sequences were identified at a single nucleotide threshold

(Amplicon Sequence Variants; ASV; Callahan et al., 2017). Additional steps of quality control

were performed to reduce non-target sequences by aligning (vsearch alignment method) the

ASV sequences to the 99% representative sequences from the greengenes database and only

ASV with a similarity threshold of 0.65% and coverage threshold of 0.5% were retained

following https://github.com/biocore/deblur/issues/139#issuecomment-282143373. Singletons

and sequences with less than 349bp were also removed with filter-features and filter-sets

function in QIIME2, respectively. The quality-filtered sequences were then taxonomically

assigned with a naïve-Bayes classifier trained on the V3-V4 region of the 16S gene in the Silva

v138 database (Quast et al., 2013), using the classify-sklearn function in the feature-classifier

plugin (Bolyen et al., 2019). The output from this generated a taxonomy table with taxonomic

naming for each ASV found within our samples. ASV sequences that were identified as

mitochondria, chloroplast, archaea, and eukaryotes were filtered out. The quality-filtered ASV

sequences were then ready to be used for downstream analysis.

Profiling microbiome functions

The functional microbiome was predicted using the software Phylogenetic Investigation

Communities by Reconstruction of Unobserved States version 2 (PICRUSt2; Douglas et al.,

2020). The predicted functions associated with the microbiome (using the quality-filtered

sequences) were inferred based on the MetaCyc pathway abundance, using the PICRUSt plug-

in within QIIME2.

Data analysis

Unless otherwise specified, the R statistical programming language (ver. 4.2.0, R Core Team) and R Studio was used to generate graphics and statistical analyses. The oysters' body temperature data violated the statistical assumption of homogeneity of variances (Levene's test, p-value < 0.05). Hence, the statistical analysis for this trait across different timepoints was assessed using a generalized linear model (GLM) with a gamma distribution. The likelihood-ratio chi-square test was used to determine the statistical significance and is indicated if the p-value was < 0.05 (R package "Car"; Fox and Weisberg, 2019). Post-hoc testing for pairwise comparison was made using the estimated marginal means with Tukey's tests from the R package emmeans (Lenth, 2023)

Individual PUFA-oxidised metabolites were first evaluated for normality (Shapiro-Wilk's Test) and homogeneity of variance (Levene's test) assumptions. Metabolites that fulfilled both assumptions were tested with one way Analysis of variance (ANOVA) to investigate the variation in metabolites' concentrations across timepoints, followed by *post-hoc* Tukey's Honest significant difference (Tukey's HSD). For metabolites that violated either one of the assumptions, they were evaluated with GLM (gamma distribution) following a similar procedure as with the body temperature datasets.

Assessment of PUFA-oxidised metabolites' beta diversity (with all the 15 metabolites or only the 4 significant metabolites) was performed by first generating a Euclidean distance matrix between samples. A principle Coordinate Analysis (PCoA) plot was then created to visualize the PUFA-oxidised metabolites clustering across different timepoints, using the ggplot2 R package (Wickham, 2008). Permutational multivariate analysis of variance (PERMANOVA) was performed with 999 permutations to test for the differences in PUFA-oxidised metabolites profile across timepoints (Oksanen et al., 2022). We also performed a permutational multivariate dispersion test (PERMDISP) to assess differences in the dispersion of metabolite profiles between timepoints with 999 permutations followed by pairwise comparisons. All multivariate metabolites analyses were performed using the package vegan in R (Oksanen et al., 2022).

Oysters' gut microbiome analysis was conducted by first combining the ASV count data, taxonomy table, and metadata into a simplified data matrix for ease of downstream processing, using the phyloseq() function from the phyloseq R package (McMurdie and Holmes, 2013). Rarefaction curves (total ASVs captured against sampling depth) were generated for each sample with the ggrare() function from the ranacapa R package (Kandlikar et al., 2018). All

samples approached the asymptote level, indicating that the total ASV richness for each sample has been captured (Fig. S1). Each of the samples was then rarefied to a sampling depth of 21,282 reads (a cut-off that balances the maximum number of sequences retained per sample and minimizing the number of samples removed) for downstream analysis (i.e., alpha and beta diversity; McKnight et al., 2019; Weiss et al., 2017). Using the rarefied data, the microbiome alpha diversity was determined based on two indices – ASV richness and Shannon-Wiener (H') diversity. ASV richness was calculated with the Base-R function, while Shannon-Wiener (H') diversity index was generated using the diversity() function from the Vegan package (Oksanen et al., 2022). Differences in the microbiome alpha diversity across the timepoints were assessed with ANOVA and followed by a *post-hoc* Tukey HSD test. The ANOVA statistical assumptions of normality (Shapiro-Wilk test) and homogeneity of variance (Levene's test) for all datasets were tested and fulfilled (p-value > 0.05). Linear regression analysis was also conducted to determine the relationship between the oysters' microbiome alpha diversity (i.e., ASV richness and Shannon-Wiener (H') index) and body temperature (i.e., average of the 10 random independent body temperature replicates collected at each timepoint). The statistical assumptions for the linear regression model were fulfilled based on the normality test (Shapiro-Wilk test) and residuals plot (variance homogeneity).

Beta diversity of the oysters' gut microbiome taxonomic and functional profiles was evaluated by generating the Bray-Curtis dissimilarity distance matrix between samples, and a PCoA plot was then created to visualize microbiome taxonomic and functional clusters across different timepoints, using the ggplot2 R package (Wickham, 2008). PERMANOVA was performed with 999 permutations to test for the differences in the microbiome composition and structure between timepoints (Oksanen et al., 2022). PERMDISP with 999 permutations was performed to test for differences in the taxonomic and functional microbiome profile dispersion followed by pairwise comparison. All multivariate analyses were conducted using the package vegan in R.

For the microbial taxa profiling, the unrarefied quality-filtered ASV sequences were binned to the family taxonomic level for each sample. Differential abundance analysis of microbiome was performed with family taxonomic level data and analysed with Analysis of Composition Microbiome with Bias Correction (ANCOM-BC) from the ANCOMBC R package (Lin and Peddada, 2020; Nearing et al., 2022).

Mantel test (Spearman correlation method) was employed to determine the correlation between the *I. nucleus* microbiome taxonomic structure and the PUFA-oxidised metabolites using their dissimilarity matrices across timepoints using the mantel() function in vegan

package (Oksanen et al., 2022), which tested whether higher dissimilarity in microbiome was associated with higher dissimilarity in metabolite profiles. Since the oysters' microbiome and metabolites were sampled from independent individuals, we averaged each microbial taxon and each metabolite concentration across replicates at each timepoints before generating a dissimilarity distance matrix between samples – Bray-Curtis (microbiome) and Euclidean (metabolites).

The GraphPad Prism software (San Diego, California, USA) was used to generate the figure for oyster's body temperatures, individual PUFA-oxidised metabolites and the 10 dominant microbe's relative abundance across the timepoints.

Results

Tidal dynamics and temporal changes in environmental and body temperatures

Oysters were regularly exposed to the air as a result of the tidal regime (Fig. 1A) and this temporal change was aligned to changes in the rock surface temperature across the day, with the highest temperature recorded at noon (45-50°C, Fig. 1A). The body temperature of the oysters varied temporally (Fig. 1B; GLM $\mathcal{X}^2_{(3)} = 1073.9$, p-value < 0.05), increasing from 32.11 ± 0.12 °C (mean ± SD) in the morning to 39.99 ± 0.95 °C (mean ± SD) at noon, after which body temperatures drop steadily to ~ 35 °C in the afternoon. Besides early-afternoon (35.4°C ± 0.46) and late-afternoon (35.18 °C ± 0.46), all pairwise comparisons showed significant differences in body temperatures between timepoints. Such temporal variation in body temperature encompasses the thermal window (noon's temperature) where oysters are expected to enter a state of metabolic depression (body temperatures falling within the metabolic depression range, i.e., cBP1 – cBP3; Fig. 1C), as well as windows where they remain active (body temperatures before cBP1, Fig. 1C).

Temporal assessment of PUFA-oxidised metabolites

There was no significant difference in the metabolites profile (contributed by fifteen metabolites) between timepoints (Table 2), but PERMDISP indicated a significant difference in their dispersion (Table 3; Fig. 3A). Individual analyses revealed that only four PUFA-oxidised metabolites (i.e., $5\text{-}F_{2t}\text{-IsoP}$, $10\text{-}F_{4t}\text{-NeuroP}$, $13\text{-}F_{4t}\text{-NeuroP}$, and $16\text{-}F_{1t}\text{-PhytoP}$) significantly varied between timepoints (Table 1, Fig. 2). An additional PERMANOVA analysis for these four metabolites showed a significant difference in the clusters between the morning metabolites profile and all the other timepoints during the emersion period (Table 2;

Fig. 3B). Multivariate dispersion of the four PUFA-oxidised metabolites also differed between

timepoints (Table 3; Fig. 3B). With the exception of two pairwise comparisons (i.e., noon vs.

early-afternoon, and early-afternoon vs. late-afternoon), all the other timepoints showed

significant differences.

Temporal characterization of oyster-microbiomes

The 16S rRNA amplicon sequencing generated a total of 2,032,567 reads from 30

samples which were reduced to 1,126,884 after denoising and quality control. These reads were

then used to characterize the microbial communities of *I. nucleus* across the different time

points. Due to the low sequence counts, samples BRM3 (5612 reads) and DRM4 (7519 reads)

were removed after each sample was rarefied to a sampling depth of 21,282 reads.

The ASV richness and Shannon-Wiener index of *I. nucleus* were significantly different

across the emersion period (Fig. 4; ASV richness: $_{3,14)}$ = 3.67; p-value < 0.05; Shannon-Wiener

index: $F_{(3,14)}$ = 3.76; p-value < 0.05). However, no specific temporal trends were detected for

all pairwise comparisons in ASV richness. Significant differences were, however, found for the

Shannon-Wiener index between morning and noon timepoints. Oyster body temperature

significantly influenced ASV richness and Shannon-Wiener indexes (Fig. S2). Overall,

morning timepoints showed the highest alpha diversity. This diversity then drastically declined

at noon, with partial recovery at the last timepoint – late-afternoon.

The top 10 most abundant microbial taxa (at the family level) associated with *I. nucleus*

were   *Mycoplasmataceae,*   *Helicobacteraceae,*   *Spirochaetaceae,*   *Xenococcaceae,*

Rhodobacteceae,   *Rickettsiales*   unknown*,*   *Arcobactericeae,*   *Caldilineaceae,*

Alphaproteobacteria unknown and *Saprospiraceae* (Fig. 5 and 6). These top 10 families made

up on average 57.09%, 80.88%, 81.42%, and 68.20% of the overall relative abundance in

morning, noon, early-afternoon, and late-afternoon, respectively. Among them, only two

families (*Helicobacteraceae* and *Arcobacteraceae*) showed significant differences in the

relative abundance between timepoints based on the ANCOM-BC analysis (Fig. 6;

Helicobacteraceae: morning vs early-afternoon; *Arcobacteraceae*: morning vs noon, and

morning vs early-afternoon).

No significant difference between timepoints was detected in the taxonomic and

functional microbiome structures (Fig. 7, Table 2), but significant differences were detected in

their dispersion (Fig. 7, Table 3). Major differences in the microbiome dispersion based on

taxonomy were found between morning and early-afternoon, which corresponds to the pre-and

post-noon period, respectively. For the functional microbiome, the pair early-afternoon-late-afternoon was the cause of the difference in dispersion. Despite the changes found in some bacterial taxa and microbiome dispersions, no Mantel correlation was observed between the microbiome taxonomic profile and the PUFA-oxidised metabolite profile, either with the 15 (all) or 4 metabolite groups (Table 4).

Fig. 1. The body temperature of *Isognomon nucleus* varies across sampling timepoints within a tidal cycle. (A) Thermal (black line) and tidal conditions (blue line) as experienced by *I. nucleus* at the study site. Thermal condition is represented by the mean rock surface temperature (averaged over 3 – 5 i-Button temperature loggers), and tidal condition during the day is categorized into three states: "E" (exposed), "A" (awash), and "I" (immersed). Black arrows indicate the sampling time points of *I. nucleus* to assess its physiological responses (i.e., biochemical dynamics) and microbiome taxonomic/functional profile (B) Boxplots showing *I. nucleus* body temperatures (n=10) across different timepoints (Centre line represent the median; box limits indicate the first and third quartiles). Statistical significance (p-value < 0.05) is indicated if the letters above each bar are different. (C) Schematic diagram of *I. nucleus* cardiac thermal performance curves (TPC) over a gradient of increasing temperature (adapted from Hui et al. (2020)). Coloured text boxes indicated the sampling timepoints (body temperatures) of the *I. nucleus* in the TPC. The abbreviated labels (with temperatures) and arrows pointing to the yellow circles refer to the different critical body temperature for cardiac thermal performances (see Hui et al. (2020); cBP–- critical breakpoints, FBP–- Final breakpoint, FLT Flatline temperature). Metabolic depression breadth is defined as between cBP1 and cBP3.

Fig. 2. Four PUFA-oxidised metabolites exhibit variation in concentration across the emersion period. Bar graphs showing the metabolites derived from Arachidonic acid: (A) 5-F_{2t}-IsoP (B) 15-F_{2t}-IsoP; from Adrenic acid: (C) 7-F_{2t}-Dihomo-IsoP (D) 17- F_{2t}-Dihomo-IsoP; from Docosahexaenoic acid: (E) 4-F_{4t}-NeuroP (F) 10-F_{4t}-NeuroP (G) 13-F_{4t}-NeuroP (H) 20-F_{4t}-NeuroP; from α-Linolenic acid: (I) 9-D_{1t}-PhytoP (J) 9-F_{1t}-PhytoP (K) 16-F_{1t}-PhytoP (L) 16-B_{1t}-PhytoP; from Eicosapentaenoic acid: (M) 5-F_{3t}-IsoP (N) 15-F_{3t}-IsoP (O) 8-F_{3t}-IsoP, across different sampling timepoints in *I. nucleus*. IsoP: isoprostane; NeuroP: neuroprostane; PhytoP: phytoprostanes. Data are presented as mean concentration and + 1 standard deviation (error bar) for each metabolite (n=5) at the given timepoints. Statistical significance (p-value < 0.05) is indicated if the letters above each bar are different.

**Fig. 3. Four PUFA-oxidised metabolites drive variation in the sample clusters across the
emersion period.** PCoA based on Euclidean distance measured between samples from different
timepoints based on (A) all PUFA-oxidised metabolites (B) four PUFA-oxidised metabolites:
5-F_{2t}-IsoP, 10-F_{4t}-NeuroP, 13-F_{4t}-NeuroP, and 16-F_{1t}-PhytoP. The colours of the datapoints and
ellipses (95% confidence interval of the true multivariate means) indicate their corresponding
timepoints.

**Fig. 4. *I. nucleus*-associated microbiome ASV richness and Shannon-Wiener diversity
across the emersion period.** Boxplot showing the alpha diversity index of (A) ASV richness,
and (B) Shannon-Wiener index (H') of the oyster-associated microbiome at different timepoints.
(Centre line represents the median; box limits indicate the first and third quartiles; the lower
and upper whiskers indicate the smallest and largest values that within 1.5 times of the IQR
(Inter-quartile range) from the first and third quartiles, respectively. The data points shown in
each timepoint is replicate values. Statistical significance ($p < 0.05$) is indicated if the letters
above each boxplot are different between any two timepoints.

Fig. 5. Relative abundance of top 30 most dominant microbial taxa associated with the *I. nucleus's* gut microbiome at each sampling timepoints. ASV counts were aggregated into the family taxonomic level. Each bar represents a single individual replicate of oyster gut microbiome.

Fig. 6. *Helicobacteraceae* and *Arcobacteraceae* mean relative abundance vary across the emersion period. Bar graphs showing the top 10 dominant family level taxa in *I. nucleus*'s gut microbiome at each timepoint: (A) *Mycoplasmataceae* (B) *Helicobacteraceae* (C) *Spirochataceae* (D) *Xenococcaceae* (E) *Rhodobacteraceae* (F) *Rickettsiales* unknown (G) *Arcobacteraceae* (H) *Caldilineaceae* (I) *Alphaproteobacteria* unknown (J) *Saprospiraceae*. For F and I, the taxa were assigned with the lowest taxonomic level name identifiable. Data are

461 presented as mean relative abundance and + 1 standard deviation (error bar) for each taxon at
462 the given timepoints. Statistical significance (p-value < 0.05) is indicated if the letters above
463 each bar are different.
464

Fig. 7. *I. nucleus*-**associated microbiome taxonomic and functional structures exhibit**
overall stability across the emersion period. PCoA based on the Bray-Curtis distance metric
measured between samples from different timepoints, based on (A) taxonomy (ASV level) and
(B) functions (MetaCyc pathways). The colours of the datapoints and ellipses (95% confidence
interval of the true multivariate means) indicate their corresponding timepoints.

Table 1. Statistical analysis result (ANOVA or GLM with gamma distribution) testing the
effect of timepoints on the concentration of each of the PUFA-oxidised metabolites.

PUFA	Metabolite	Test Statistic	p-value
Arachidonic acid	$5\text{-}F_{2t}\text{-}IsoP$	$F_{(3,14)} = 5.08$	**< 0.05**
	$15\text{-}F_{2t}\text{-}IsoP$	$F_{(3,14)} = 1.72$	> 0.05
Adrenic acid	$7\text{-}F_{2t}\text{-}Dihomo\text{-}IsoP$	$F_{(3,14)} = 3.10$	> 0.05
	$17\text{-}F_{2t}\text{-}Dihomo\text{-}IsoP$	GLM $\mathcal{X}^2_{(3)} = 0.66$	> 0.05
Docosahexaenoic acid	$4\text{-}F_{4t}\text{-}NeuroP$	$F_{(3,14)} = 3.13$	> 0.05
	$10\text{-}F_{4t}\text{-}NeuroP$	$F_{(3,14)} = 8.43$	**< 0.05**
	$13\text{-}F_{4t}\text{-}NeuroP$	GLM $\mathcal{X}^2_{(3)} = 27.46$	**< 0.05**
	$20\text{-}F_{4t}\text{-}NeuroP$	$F_{(3,14)} = 1.23$	> 0.05
α-Linolenic acid	$9\text{-}D_{1t}\text{-}PhytoP$	$F_{(3,14)} = 0.58$	> 0.05
	$9\text{-}F_{1t}\text{-}PhytoP$	GLM $\mathcal{X}^2_{(3)} = 7.92$	> 0.05
	$16\text{-}F_{1t}\text{-}PhytoP$	$F_{(3,14)} = 3.72$	**< 0.05**
	$16\text{-}B_{1t}\text{-}PhytoP$	GLM $\mathcal{X}^2_{(3)} = 6.66$	> 0.05
Eicosapentaenoic acid	$5\text{-}F_{3t}\text{-}IsoP$	$F_{(3,14)} = 2.49$	> 0.05
	$15\text{-}F_{3t}\text{-}IsoP$	GLM $\mathcal{X}^2_{(3)} = 0.32$	> 0.05
	$8\text{-}F_{3t}\text{-}IsoP$	$F_{(3,14)} = 0.81$	> 0.05

Table 2. Permutation multivariate analysis of variance (PERMANOVA) to investigate differences in the PUFA-oxidised metabolites or microbiome structure in taxonomy/functions across the emersion period. The number of permutations performed was 999 times. The numerator and denominator degree of freedom were 3 and 14, respectively.

Category	Distance metric	$pseudo\text{-}F$	R^2	p-value
PUFA-oxidised metabolites (all)	Euclidean	2.14	0.31	> 0.05
PUFA-oxidised metabolites (four)*	Euclidean	4.02	0.46	**< 0.05**
Microbiome (Taxonomy)	Bray-Curtis	1.18	0.20	> 0.05
Microbiome (Function)	Bray-Curtis	1.49	0.24	> 0.05

*The four metabolites were: $5\text{-}F_{2t}\text{-}IsoP$, $10\text{-}F_{4t}\text{-}NeuroP$, $13\text{-}F_{4t}\text{-}NeuroP$, and $16\text{-}F_{1t}\text{-}PhytoP$

Table 3. Permutational analysis of multivariate dispersions (PERMDISP) analysis to investigate differences in the PUFA-oxidised metabolites or microbiome structure in taxonomy/functions across the emersion period. The number of permutations performed was 999 times.

Category	Distance metric	F-value	p-value
PUFA-oxidised metabolites (all)	Euclidean	6.88	**< 0.05**
PUFA-oxidised metabolites (four)*	Euclidean	3.51	**< 0.05**
Microbiome (Taxonomy)	Bray-Curtis	3.55	**< 0.05**
Microbiome (Functions)	Bray-Curtis	3.74	**< 0.05**

*The four metabolites were: $5\text{-}F_{2t}\text{-}IsoP$, $10\text{-}F_{4t}\text{-}NeuroP$, $13\text{-}F_{4t}\text{-}NeuroP$, and $16\text{-}F_{1t}\text{-}PhytoP$

488 Table 4. Mantel's spearman's rank correlation (r) and p-value to investigate the association
489 between the *I. nucleus* microbiome taxonomic structure (mean abundance of each taxon across
490 all timepoints) and PUFA-oxidised metabolite profile (mean concentration across timepoints).
491 The number of permutations performed was 23(maximum possible).
492

Association with microbiome taxonomic structure	Mantel's r	p-value
PUFA-oxidised metabolites (all)	0.7134	> 0.05
PUFA-oxidised metabolites (Four)*	0.1343	> 0.05

493 *The four metabolites were: $5\text{-}F_{2t}\text{-}IsoP$, $10\text{-}F_{4t}\text{-}NeuroP$, $13\text{-}F_{4t}\text{-}NeuroP$, and $16\text{-}F_{1t}\text{-}PhytoP$

Discussion

Organisms living on the intertidal rocky shore regularly experience drastic changes in
environmental temperature (Harley, 2008; Somero, 2002; Stillman, 2002). For many sessile
ectotherms, this represents a physiological challenge due to their inability to escape from heat
stress (e.g. seeking thermal refuge) and so, they rely on various intrinsic (metabolic and
physiological plasticity) and extrinsic (host-associated microbiome) mechanisms to cope with
temperature fluctuations (Bang et al., 2018; Giomi et al., 2016; Huang et al., 2020; Lathlean et
al., 2014). As with other ectotherms, the body temperature of the high shore oyster, *Isognomon*
nucleus was highly influenced by changes in ambient temperatures throughout the day. Such
changes are known to induce physiological stress and metabolic depression (Hui et al., 2020),
with the associated oxidative stress (Hermes-Lima et al., 2015). Here, linked to these
physiological conditions, we found adjustments in four major non-enzymatic PUFA-oxidised
metabolites (i.e., $5\text{-}F_{2t}\text{-}IsoP$, $10\text{-}F_{4t}\text{-}NeuroP$, $13\text{-}F_{4t}\text{-}NeuroP$, and $16\text{-}F_{1t}\text{-}PhytoP$) that reflect
temporal variation in the oxidative stress experienced by the oysters. Contrary, no clear
temporal extrinsic adjustments in the structure and/or function of the gut microbiome were
detected despite the temporal changes in environmental conditions. This suggests that the oyster
gut microbiome is temporally stable and insensitive to extreme thermal conditions (> 40ºC in
body temperature) that are physiologically stressful to the oyster host. The differences in the
host and microbiome sensitivity to external changes was further supported in the lack of
temporal correlation between the intrinsic (biochemical profile) and extrinsic (microbiome
taxonomic structure) mechanisms. In summary, the microbiome stability may be important for
the overall homeostasis of the oysters, contributing to their thermal tolerance and survival in
the challenging tropical intertidal rocky environment.

The intrinsic mechanism underpinning oysters' adaptive response to temporal environmental changes

For high-shore intertidal ectotherms, metabolic depression is an adaptive response that allows them to cope with extreme and stressful thermal conditions (Hui et al., 2020; Marshall et al., 2011; Sokolova and Pörtner, 2001). This physiological adjustment is a common signal that the animal is under thermal stress and acts as a strategy to conserve energy required for critical functions during normal metabolic resumption (i.e., less stressful thermal conditions; Hand and Hardewig, 1996; Marshall and McQuaid, 2020). Metabolic depression is also known to be an important mechanism to reduce/avoid the over-production of harmful ROS that results from the rapid increase in metabolic demands under elevated temperatures (Abele et al., 2002; Storey and Storey, 2004). Hui et al. (2020) demonstrated that the oyster *I. nucleus* consistently enters into metabolic depression when body temperature approaches ~ 37°C (a value that is regularly surpassed on daily basis at around noon on tropical rocky shores). This suggests that these animals should enter a state of metabolic depression when exposed by the tide during noon time and resume their metabolism when body temperatures lower in late afternoon/evening, or when they are re-immersed. During the low tide, oysters are exposed to anaerobic conditions because of the closure of the shell, which prevents oxygen exchange with the environment (Chapman et al., 2011). However, as with other intertidal bivalves, oysters can display intermittent gaping behaviour during periods of aerial exposure as a way to enhance oxygen intake and evaporative cooling, particularly when experiencing thermal stress (Lent, 1969; Mcmahon, 1988; Nicastro et al., 2010). In our study, all individuals shut their valves and no intermittent gaping behaviour was observed either during the high thermal stress or the recovering period (Hui, T.Y., unpublished data). Therefore, oxygen depletion is a plausible stress in the oysters particularly under elevated body temperatures as experienced during noon time, and anaerobic respiration is likely to contribute to the oysters' metabolic demand during such thermally stressful conditions.

As part of the metabolic depression, intertidal ectotherms modulate biochemical molecules such as polyunsaturated fatty acids (PUFA) to cope with the elevated thermal and oxidative stress conditions in their environment (Dubousquet et al., 2016; Pernet et al., 2008). The higher metabolic demand driven by higher temperatures generate ROS during and after thermal stress and/or under hypoxia (Byrne et al., 1990; Freire et al., 2011; Rivera-Ingraham et al., 2013; Welker et al., 2013). PUFAs such as ARA, DHA, and EPA are known to be effective molecules to mitigate the effect of ROS (Lund and Steenfeldt, 2011; Yin et al., 2017). This is

because these PUFA contain unsaturated double bonds making them susceptible to ROS reaction (particularly to peroxide, HO•), and thus could be an agent for ROS removal (Fadhlaoui and Lavoie, 2021; Leung et al., 2015; Sargent, 1976). The non-enzymatic peroxidation of these PUFAs, therefore, generates by-products that can be used as biomarkers of the ROS level (Galano et al., 2017; Miller et al., 2014; Yonny et al., 2016). Using this as a measure of the oyster's biochemical response to thermal environmental changes, we found that there were four metabolites (i.e., $5\text{-}F_{2t}\text{-}IsoP$ (Isoprostane), $10\text{-}F_{4t}\text{-}NeuroP$ (Neuroprostane), $13\text{-}F_{4t}\text{-}NeuroP$ (Neuroprostane), and $16\text{-}F_{1t}\text{-}PhytoP$ (Phytoprostane)) that showed visible temporal patterns. Such patterns may be linked to the sensitivity of the metabolites to ROS changes in the oyster. Elevation of $5\text{-}F_{2t}\text{-}IsoP$, $10\text{-}F_{4t}\text{-}NeuroP$, and $13\text{-}F_{4t}\text{-}NeuroP$ during noon, and $16\text{-}F_{1t}\text{-}PhytoP$ at post-noon timepoints, clearly indicates that oysters experienced a rise in the ROS level, a signal of oxidative stress (Galano et al., 2017; Hermes-Lima and Zenteno-Savín, 2002; Storey and Storey, 2004; Sussarellu et al., 2012; Welker et al., 2013). This could be caused by an increase in metabolic demand and/or hypoxia (when individuals shut their valves and show no gaping behaviour). ALA and its non-enzymatic oxidised by-products (phytoprostanes) are known to be the major molecules to cope with oxidative stress in plants (Durand et al., 2009; Imbusch and Mueller, 2000; Jahn et al., 2008; Loeffler et al., 2005; Thoma et al., 2003). For filter-feeding bivalves, ALA can be obtained from phytoplankton, which is a major part of their diet (Sargent, 1976; Zhukova, 2019). In our study, the high levels of $16\text{-}F_{1t}\text{-}PhytoP$ (Phytoprostane) metabolites as compared to other three PUFA-oxidised metabolites in the *I. nucleus,* imply that the PUFA ALA could also be the predominant fatty acid in alleviating oxidative stress in this oyster. Thus, for *I. nucleus,* phytoplankton, through their fatty acid composition, are likely to provide key benefits for their physiological regulation and survival in the thermally and oxidatively stressful conditions at the high shore environment.

The structural profile of the ROS-sensitive metabolites exhibits clear temporal variation and inter-individual variation in the *I. nucleus*. The large inter-individual variability in the PUFA-metabolites profile at noon/afternoon relative to morning timepoint might be related to the thermal and oxidative stress conditions experienced by the oysters during those periods of elevated temperatures. This variability suggests inter-individual differences in physiological plasticity, regulation, and sensitivities to environmental stress. The result here also indicates that the polyunsaturated fatty acids (ARA, DHA, and ALA) from which these four PUFA-oxidised metabolites were derived, may play a key role acting as 'resolving' lipid mediators needed by the oysters to return to homeostasis (Lee et al., 2020; Roy et al., 2017).

The extrinsic mechanism underpinning the oysters' adaptive response to temporal environmental changes

Host-associated microbiome diversity, structure, and functions are more frequently considered as important components of animal adaptation and acclimation to environmental change (Bang et al., 2018; Foster et al., 2017; Voolstra and Ziegler, 2020). Alterations in microbiome diversity can be associated with the direct influence of the host's physiology and/or with host-independent environmental effects (Li et al., 2018; Li et al., 2022; Muñoz et al., 2019; Shiu et al., 2017). In our study, the composition and diversity (ASV richness and Shannon diversity) of the oyster gut-associated microbiome varied with time which coincided with variation in the oyster's body temperature, suggesting that the thermal condition experienced by the host may modulate the microbial alpha diversity via either direct/indirect physio-biochemical influences (Apprill, 2017; Pita et al., 2018; Sorek et al., 2014). In the morning, the oyster's gut had a significantly more diverse microbiome diversity, which may be linked to the less thermally stressed host (Hui et al., 2020), and/or to the filter-feeding activities before being exposed by the receding tide (Parris et al., 2019). However, at noon, a major decline in diversity was observed, which may be associated with the thermal physiological stress experienced by the host (body temperature is within the metabolic depression range; Hui et al., 2020) and oxidative stress (indicated by the rise in the PUFA-oxidised metabolites). This decline in diversity is also a common pattern seen in many marine organisms under physiological stress (Green and Barnes, 2010; Hartman et al., 2020; Li et al., 2018). For example, Lokmer & Wegner (2015) showed that thermal stress and/or infection caused a marked decline in the alpha diversity of the microbiome in the oyster *Crassostrea gigas*. Interestingly, in our study, during the late afternoon timepoint, we observed an increase in the microbiome diversity, driven by an increase in the extremely low abundant taxa, most of them undetectable at noon. This trend suggests that the stress condition experienced during the day did not cause an irreversible breakdown in the host-microbiome relationship, in which diversity seems resilient probably as a consequence of the predictability and magnitude of the environmental change (Huus and Ley, 2021; Longford et al., 2019; Narrowe et al., 2015; Webster et al., 2008). It is unclear, however, whether this trend would be maintained when stressful conditions are extreme and unpredictable, such as in the case of marine heatwaves.

Whilst the temporal changes exhibited by the *I. nucleus* microbiome and their association with changes in the oysters' body temperatures suggest that the microbiome composition may be actively regulated by the host, differential sensitivities of certain bacterial

taxa to external environmental changes may also be an important mechanism driving the
temporal variation in the microbiome composition. *Helicobacteraceae* and *Arcobacteraceae*
are the only two dominant family-level microbial taxa associated with the temporal
thermal/oxidative stress variation of the host. In particular, there was an elevated relative
abundance of these two bacteria groups at noon and/or at the afternoon timepoints, which may
suggest that their abundances during these periods were promoted by specific
host/environmental conditions, facilitating their growth probably by reducing competition
(Garren et al., 2016; Sharp and Foster, 2022; Shaver et al., 2017). *Arcobacteraceae* is a group
of common opportunistic bacteria found in marine bivalves and is known to be associated with
disease-causing bacteria such as *Vibrio* (de Lorgeril et al., 2018; Lokmer and Mathias Wegner,
2015). In many studies, the elevation of the *Arcobacteraceae* has been associated with the
thermal stress condition experienced by the host (Lokmer and Mathias Wegner, 2015; Neu et
al., 2021; Shiu et al., 2017). Members of *Arcobacteraceae* are also known to be microaerophilic
as they tend to grow better under a low-oxygen environment (Vandamme and De Ley, 1991).
This may explain the dominant presence of *Arcobacteraceae* within the oysters' microbiome
due to the prevailing hypoxic and thermally stressful conditions as experienced by the oyster
hosts during emersion. Similarly, the presence and abundance *Helicobacteraceae* can be
associated to oxygen levels in the marine environment. For instance, certain members of this
family are capable of oxidising sulfide, and thus, they might contribute with a detoxification
function to the oyster host (Kodama and Watanabe, 2004; Nakagawa et al., 2017; Neu et al.,
2021). The accumulation of sulfide can be detrimental or lethal to marine bivalves (Le Moullac
et al., 2008), and it is typically associated with hypoxic stress (Le Moullac et al., 2008; Pal et
al., 2018). Such stress is hypothesised to occur in *I. nucleus* during aerial exposure, as no
intermittent gaping behaviour was detected. When oxygen is low, sulfide production is
promoted by the activity of hydrogen sulfide-producing bacteria such as *Desulfobacterota*
members (present in the *I. nucleus*; Dordević et al., 2021; Flannigan et al., 2011; Pimentel et
al., 2021). It is unclear, however, the extent to which the dynamics in the abundance of sulfide-
producing bacteria and *Helicobacteraceae* members (with potential detoxification role)*,* are
functionally interconnected. While this seems likely, we are unaware of any study exploring
this functional link, and it would be important to investigate such species' interaction and its
effects on the physiology and fitness of the host.
Host tolerance and resilience to environmental stress are known to be conditioned by
the stability/flexibility of its associated microbiome. Despite the changes in the alpha diversity
and relative abundance of some of the most abundant bacteria (family level) in *I. nucleus*, the

structural and functional microbiome profiles showed an overall stability throughout the day (emersion period). Such stability could be potentially explained by the ability of the host to control/regulate the microbiome during a disturbance event (Pita et al., 2018; Rädecker et al., 2022; Rocca et al., 2019; Sharp et al., 2017). Alternatively, the stability of the microbiome may reflect its own resilience (or lower sensitivity), rather than the host control/regulation, to changing and stressful environments (Soen, 2014). This characteristic could be essential for host tolerance and acclimation to environmental challenges such as thermal stress. For example, Santoro et al. (2021) demonstrated that the stability of inoculated "beneficial microorganisms" in the coral *Mussismilia hispida* provided a thermal protection effect, reducing heat stress and preventing the associated mortality (40% increase in the survival rate) of the hosts. Contrary, disruptions in the microbiome structure and function can lead to a weakened thermal tolerance and performance of the host, and a decrease in survival rate under prolonged heat stress condition, particularly in early ontogenetic stages such as in the tadpoles of the green frog *Lithobates clamitans* (Fontaine et al., 2022). Overall, regardless of the mechanisms, the temporal stability in the microbiome structure and function found in this study may imply a fundamental ecological role for *I. nucleus* survival in the thermally extreme tropical intertidal environment (Bang et al., 2018).

No correlation between microbiome structure and PUFA-oxidised metabolite profile

Over the decades, many studies have documented the role and action of PUFA (e.g., as probiotics) in modulating dynamics in the gut microbiome of different animals (Fu et al., 2021; Huyben et al., 2020; Lau et al., 2022; Ma et al., 2022). Hence, it is expected that stress-driven changes in these biochemicals and their by-products would influence the oyster gut microbiome. However, we found no correlation between the PUFA-oxidised metabolites and microbiome taxonomic structure in *I. nucleus*, despite the temporal and inter-individual variation in the PUFA-oxidised metabolite profile across the emersion period. The lack of association between the microbial and PUFA-oxidised metabolites profiles of *I. nucleus*, suggests two possibilities that are not mutually exclusive. Firstly, as mentioned previously, physiological/biochemical and microbiome profiles may have different sensitivities to changes in environmental conditions. Secondly, such microbiome profiles and biochemical responses may not show strong variation and/or alignment because of local adaptation. Here, the thermal window and oxygen levels experienced by the oysters, although extreme for many ectotherms, can be

considered within the tolerance range of the species. Several studies have shown that the
microbiome-metabolites correlation is detectable only under severe post-disturbance events and
unhealthy host conditions (Mallott et al., 2022; Nguyen et al., 2021; Paix et al., 2019; Pimentel
et al., 2021; Shankar et al., 2015; Walke et al., 2021), which go beyond tolerance levels. Thus,
in this study, oysters may be in a condition where stress is within the tolerance range, and thus
no correlation can be observed between these two components (i.e., host biochemical profile
and microbiome structure). Future studies aiming to explore the functional regulation and
interplay between intrinsic and extrinsic mechanisms in ectotherms living in extreme
environments require the assessment of both components when animals are closer to or surpass
their physiological tolerance limits at sublethal levels.

695   Conclusion

Intrinsic (e.g., non-enzymatic PUFA-oxidised metabolites) and extrinsic (e.g.,
microbiome profile) mechanisms linked to the ecophysiology and tolerance/sensitivity of the
oyster *Isognomon nucleus*, showed temporal variation indicative of thermal and oxidative stress.
Such trend was tightly aligned to changes in environmental and body temperatures during the
tidal cycle where oysters were fully emersed. Although the microbial alpha diversity in the
oyster gut varied along this temporal and environmental cycle, no major changes were detected
in the overall structure and function of the associated microbial communities. This microbiome
stability along periods of extreme temperature and oxygen deprivation, could be explained by
the potential regulatory influence of the host (consistent with the Anna Karenina Principle)
and/or the insensitivity of these microorganisms to the environmental changes experienced in
the tropical rocky shores. When assessing both the intrinsic and extrinsic mechanisms together,
we found no clear correlation between the PUFA-oxidised metabolites profile and the
microbiome profile. However, some specific bacteria groups (i.e., *Helicobacteraceae* and
Arcobacteraceae) increased during stressful periods, potentially linked to the lower oxygen
availability within the oysters due to the lack of gaping behaviour when emersed. These
microorganisms may provide important functions for the resilience and stability of the overall
microbial community, as well as for the health and survival of the oyster host in the thermally
challenging rocky intertidal environment. The overall microbiome stability can avoid dysbiosis
and contribute to the intrinsic adaptive adjustment of oyster physiology and biochemistry. To
our best knowledge, this is the first study to report the use of non-enzymatic PUFA-oxidised
metabolites as a proxy for oxidative stress levels in intertidal animals and profiling short-term

dynamics of the host-associated microbiome in an intertidal environment. Future investigations

should aim to explore the interplay between these biomolecules and microbial patterns when

stressful environmental conditions are unpredictable and beyond tolerance windows of the

hosts. Moreover, it is important to understand whether these functional links between intrinsic

and extrinsic mechanisms are modulated by local adaptation and can be shared among other

intertidal organisms.

Competing interests

725   The authors declare no competing or financial interests

Author Contributions

JDGE and BSA conceived the idea and designed the work. BSA carried out most of the work:

data collection, sample processing, statistical analyses, and data presentation. AWTC, HTY

GAW, and MG were involved in sample and environmental data collection. LKS performed

the PUFA-oxidised metabolites extraction and processing. TD supplied the in-house

synthesised oxidised lipids standards for PUFA metabolites quantification. JTY and TD were

involved in analysing the metabolite data. BSA led the manuscript writing with feedback and

input from all authors.

Acknowledgement

737   We would like to thank The Swire Group of Companies, especially Ms. Tina Chan and Cathay

738   Pacific staff for arranging air transportation of equipment to and from Thailand from Hong

739   Kong

Funding

742   JDGE was supported by the Research Grants Council (ECS 27124318) of Hong Kong.

Data availability

745   All raw data, and scripts will be made available once published.

References

Abele, D., Heise, K., Pörtner, H. O. and Puntarulo, S. (2002). Temperature-dependence of mitochondrial function and production of reactive oxygen species in the intertidal mud clam Mya arenaria. *J. Exp. Biol.* **205**, 1831–1841.

Ahmed, H. I., Herrera, M., Liew, Y. J. and Aranda, M. (2019). Long-term temperature stress in the Coral Model Aiptasia supports the "anna Karenina principle" for bacterial microbiomes. *Front. Microbiol.* **10**, 975.

Alberdi, A., Aizpurua, O., Bohmann, K., Zepeda-Mendoza, M. L. and Gilbert, M. T. P. (2016). Do Vertebrate Gut Metagenomes Confer Rapid Ecological Adaptation? *Trends Ecol. Evol.* **31**, 689–699.

Apprill, A. (2017). Marine animal microbiomes: Toward understanding host-microbiome interactions in a changing ocean. *Front. Mar. Sci.* **4**, 222.

Bäckhed, F., Roswall, J., Peng, Y., Feng, Q., Jia, H., Kovatcheva-Datchary, P., Li, Y., Xia, Y., Xie, H., Zhong, H., et al. (2015). Dynamics and stabilization of the human gut microbiome during the first year of life. *Cell Host Microbe* **17**, 690–703.

Bang, C., Dagan, T., Deines, P., Dubilier, N., Duschl, W. J., Fraune, S., Hentschel, U., Hirt, H., Hülter, N., Lachnit, T., et al. (2018). Metaorganisms in extreme environments: do microbes play a role in organismal adaptation? *Zoology* **127**, 1–19.

Basu, S. (2010). Fatty acid oxidation and isoprostanes: Oxidative strain and oxidative stress. *Prostaglandins Leukot. Essent. Fat. Acids* **82**, 219–225.

Birben, E., Sahiner, U. M., Sackesen, C., Erzurum, S. and Kalayci, O. (2012). Oxidative stress and antioxidant defense. *World Allergy Organ. J.* **5**, 9–19.

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., et al. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857.

Byrne, R. A., Gnaiger, E., McMahon, R. F. and Dietz, T. H. (1990). Behavioral and metabolic responses to emersion and subsequent reimmersion in the freshwater bivalve, Corbicula fluminea. *Biol. Bull.* **178**, 251–259.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A. and Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583.

Callahan, B. J., McMurdie, P. J. and Holmes, S. P. (2017). Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J.* **11**, 2639–2643.

Carey, H. V. and Duddleston, K. N. (2014). Animal-microbial symbioses in changing environments. *J. Therm. Biol.* **44**, 78–84.

Chapman, R. W., Mancia, A., Beal, M., Veloso, A., Rathbun, C., Blair, A., Holland, A. F., Warr, G. W., Didinato, G., Sokolova, I. M., et al. (2011). The transcriptomic responses of the eastern oyster, Crassostrea virginica, to environmental conditions. *Mol. Ecol.* **20**, 1431–1449.

Chow, C. Y. (2004). Foraging behaviour of Thais clavigera: The interplay of environmental variation and predator behaviour on sheltered rocky shores.

de Lorgeril, J., Lucasson, A., Petton, B., Toulza, E., Montagnani, C., Clerissi, C., Vidal-Dupiol, J., Chaparro, C., Galinier, R., Escoubas, J. M., et al. (2018). Immune-suppression by OsHV-1 viral infection causes fatal bacteraemia in Pacific oysters. *Nat. Commun.* **9**,.

Díaz-almeyda, E. M., Ryba, T., Ohdera, A. H., Collins, S. M., Shafer, N., Link, C., Prado-zapata, M., Ruhnke, C., Moore, M., Angel, A. M. G., et al. (2022). Thermal

Stress Has Minimal Effects on Bacterial Communities of Thermotolerant Symbiodinium
Cultures. *Front. Ecol. Evol.* **10**, 764086.

Dordević, D., Jančíková, S., Vítězová, M. and Kushkevych, I. (2021). Hydrogen sulfide
toxicity in the gut environment: Meta-analysis of sulfate-reducing and lactic acid
bacteria in inflammatory processes. *J. Adv. Res.* **27**, 55–69.

Douglas, G. M., Maffei, V. J., Zaneveld, J. R., Yurgel, S. N., Brown, J. R., Taylor, C. M.,
Huttenhower, C. and Langille, M. G. I. (2020). PICRUSt2 for prediction of
metagenome functions. *Nat. Biotechnol.* **38**, 669–673.

Dubousquet, V., Gros, E., Berteaux-Lecellier, V., Viguier, B., Raharivelomanana, P.,
Bertrand, C. and Lecellier, G. J. (2016). Changes in fatty acid composition in the giant
clam Tridacna maxima in response to thermal stress. *Biol. Open* **5**, 1400–1407.

Dupuy, A., Le Faouder, P., Vigor, C., Oger, C., Galano, J. M., Dray, C., Lee, J. C. Y.,
Valet, P., Gladine, C., Durand, T., et al. (2016). Simultaneous quantitative profiling of
20 isoprostanoids from omega-3 and omega-6 polyunsaturated fatty acids by LC-MS/MS
in various biological samples. *Anal. Chim. Acta* **921**, 46–58.

Durand, T., Bultel-Poncé, V., Guy, A., Berger, S., Mueller, M. J. and Galano, J. M.
(2009). New bioactive oxylipins formed by non-enzymatic free-radical-catalyzed
pathways: The phytoprostanes. *Lipids* **44**, 875–888.

Epstein, H. E., Torda, G. and van Oppen, M. J. H. (2019). Relative stability of the
Pocillopora acuta microbiome throughout a thermal stress event. *Coral Reefs* **38**, 373–
386.

Fadhlaoui, M. and Lavoie, I. (2021). Effects of temperature and glyphosate on fatty acid
composition, antioxidant capacity, and lipid peroxidation in the gastropod lymneae sp.
Water (Switzerland) **13**,.

Fan, L., Liu, M., Simister, R., Webster, N. S. and Thomas, T. (2013). Marine microbial
symbiosis heats up: the phylogenetic and functional response of a sponge holobiont to
thermal stress. *ISME J.* **7**, 991–1002.

Feng, J., Zhang, L., Tang, X., Xia, X., Hu, W. and Zhou, P. (2021). Season and geography
induced variation in sea cucumber (Stichopus japonicus) nutritional composition and gut
microbiota. *J. Food Compos. Anal.* **101**, 103838.

Flannigan, K. L., McCoy, K. D. and Wallace, J. L. (2011). Eukaryotic and prokaryotic
contributions to colonic hydrogen sulfide synthesis. *Am. J. Physiol. - Gastrointest. Liver*
Physiol. **301**, 188–193.

Folch, J., Lees, M. and Sloane Stanley, G. H. (1957). A simple method for the isolation and
purification of total lipides from animal tissues. *J. Biol. Chem.* **226**, 497–509.

Fontaine, S. S. and Kohl, K. D. (2022). The microbiome buffers tadpole hosts from heat
stress : a hologenomic approach to understand host-microbe interactions under warming.
J. Exp. Biol.

Fontaine, S. S., Mineo, P. M. and Kohl, K. D. (2022). Experimental manipulation of
microbiota reduces host thermal tolerance and fitness under heat stress in a vertebrate
ectotherm. *Nat. Ecol. Evol.* **6**, 405–417.

Foster, K. R., Schluter, J., Coyte, K. Z. and Rakoff-Nahoum, S. (2017). The evolution of
the host microbiome as an ecosystem on a leash. *Nature* **548**, 43–51.

Fox, J. and Weisberg, S. (2019). *An R Companion to Applied Regression.*

Freire, C. A., Welker, A. F., Storey, J. M., Storey, K. B. and Hermes-lima, M. (2011).
Oxidative Stress in Estuarine and Intertidal Environments (Temperate and Tropical). In
Oxidative Stress in Aquatic Ecosystems (ed. Abele, D.), Pablo, J.), Vázquez-Medina),
and Zenteno-Savín, T.), pp. 41–57. Blackwell Publishing Ltd.

Fu, Y., Wang, Y., Gao, H., Li, D., Jiang, R., Ge, L., Tong, C. and Xu, K. (2021).
Associations among Dietary Omega-3 Polyunsaturated Fatty Acids, the Gut Microbiota,

and Intestinal Immunity. *Mediators Inflamm.* **2021**,.

Galano, J. M., Lee, Y. Y., Oger, C., Vigor, C., Vercauteren, J., Durand, T., Giera, M.
and Lee, J. C. Y. (2017). Isoprostanes, neuroprostanes and phytoprostanes: An
overview of 25 years of research in chemistry and biology. *Prog. Lipid Res.* **68**, 83–108.

Garren, M., Son, K., Tout, J., Seymour, J. R. and Stocker, R. (2016). Temperature-
induced behavioral switches in a bacterial coral pathogen. *ISME J.* **10**, 1363–1372.

Garrity, S. D. (1984). Some Adaptations of Gastropods to Physical Stress on a Tropical
Rocky Shore Author (s): Stephen D . Garrity Published by : Wiley Stable URL :
http://www.jstor.org/stable/1941418 REFERENCES Linked references are available on
JSTOR for this article : You. *Ecology* **65**, 559–574.

Giomi, F., Mandaglio, C., Ganmanee, M., Han, G. D., Dong, Y., Williams, G. A. and
Sara, G. (2016). The importance of thermal history: Costs and benefits of heat exposure
in a tropical, rocky shore oyster. *J. Exp. Biol.* **219**, 686–694.

Green, T. J. and Barnes, A. C. (2010). Bacterial diversity of the digestive gland of Sydney
rock oysters, Saccostrea glomerata infected with the paramyxean parasite, Marteilia
sydneyi. *J. Appl. Microbiol.* **109**, 613–622.

Hand, S. C. and Hardewig, I. (1996). Downregulation of cellular metabolism during
environmental stress: Mechanisms and implications. *Annu. Rev. Physiol.* **58**, 539–563.

Harley, C. D. G. (2008). Tidal dynamics, topographic orientation, and temperature-mediated
mass mortalities on rocky shores. *Mar. Ecol. Prog. Ser.* **371**, 37–46.

Hartman, L. M., van Oppen, M. J. H. and Blackall, L. L. (2020). The effect of thermal
stress on the bacterial microbiome of exaiptasia diaphana. *Microorganisms* **8**, 1–18.

Helmuth, B. and Hofmann, G. E. (2001). Microhabitats, thermal heterogeneity, and patterns
of physiological stress in the rocky intertidal zone. *Biol. Bull.* **201**, 374–384.

Herlemann, D. P. R., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J. and
Andersson, A. F. (2011). Transitions in bacterial communities along the 2000 km
salinity gradient of the Baltic Sea. *ISME J.* **5**, 1571–1579.

Hermes-Lima, M. and Zenteno-Savín, T. (2002). Animal response to drastic changes in
oxygen availability and physiological oxidative stress. *Comp. Biochem. Physiol. - C*
Toxicol. Pharmacol. **133**, 537–556.

Hermes-Lima, M., Moreira, D. C., Rivera-Ingraham, G. A., Giraud-Billoud, M.,
Genaro-Mattos, T. C. and Campos, É. G. (2015). Preparation for oxidative stress
under hypoxia and metabolic depression: Revisiting the proposal two decades later. *Free*
Radic. Biol. Med. **89**, 1122–1143.

Herrera, M., Klein, S. G., Schmidt-Roach, S., Campana, S., Cziesielski, M. J., Chen, J.
E., Duarte, C. M. and Aranda, M. (2020). Unfamiliar partnerships limit cnidarian
holobiont acclimation to warming. *Glob. Chang. Biol.* **26**, 5539–5553.

Huang, Z., Yu, K., Fang, Y., Dai, H., Cai, H., Li, Z., Kan, B., Wei, Q. and Wang, D.
(2020). Comparative Genomics and Transcriptomics Analyses Reveal a Unique
Environmental Adaptability of Vibrio fujianensis. *Microorganisms* **8**, 555.

Hui, T. Y., Dong, Y., Han, G. D., Lau, S. L. Y., Cheng, M. C. F., Meepoka, C.,
Ganmanee, M. and Williams, G. A. (2020). Timing metabolic depression: Predicting
thermal stress in extreme intertidal environments. *Am. Nat.* **196**, 501–511.

Huus, K. E. and Ley, R. E. (2021). Blowing Hot and Cold: Body Temperature and the
Microbiome. *mSystems* **6**,.

Huyben, D., Roehe, B. K., Bekaert, M., Ruyter, B. and Glencross, B. (2020). Dietary
Lipid:Protein Ratio and n-3 Long-Chain Polyunsaturated Fatty Acids Alters the Gut
Microbiome of Atlantic Salmon Under Hypoxic and Normoxic Conditions. *Front.*
Microbiol. **11**, 589898.

Imbusch, R. and Mueller, M. J. (2000). Analysis of oxidative stress and wound-inducible

dinor isoprostanes F1 (Phytoprostanes F1) in plants. *Plant Physiol.* **124**, 1293–1303.

Jahn, U., Galano, J. M. and Durand, T. (2008). Beyond prostaglandins - Chemistry and biology of cyclic oxygenated metabolites formed by free-radical pathways from polyunsaturated fatty acids. *Angew. Chemie - Int. Ed.* **47**, 5894–5955.

Kandlikar, G. S., Gold, Z. J., Cowen, M. C., Meyer, R. S., Freise, A. C., Kraft, N. J. B., Moberg-Parker, J., Sprague, J., Kushner, D. J. and Curd, E. E. (2018). Ranacapa: An R package and shiny web app to explore environmental DNA data with exploratory statistics and interactive visualizations. *F1000Research* **7**, 1–19.

Klepac, C. N., Beal, J., Kenkel, C. D., Sproles, A., Polinski, J. M., Williams, M. A., Matz, M. V. and Voss, J. D. (2015). Seasonal stability of coral-Symbiodinium associations in the subtropical coral habitat of St. Lucie Reef, Florida. *Mar. Ecol. Prog. Ser.* **532**, 137–151.

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M. and Glöckner, F. O. (2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* **41**, 1–11.

Kodama, Y. and Watanabe, K. (2004). Sulfuricurvum kujiense gen. nov., sp. nov., a facultatively anaerobic, chemolithoautotrophic, sulfur-oxidizing bacterium isolated from an underground crude-oil storage cavity. *Int. J. Syst. Evol. Microbiol.* **54**, 2297–2300.

Kremer, N., Koch, E. J., El Filali, A., Zhou, L., Heath-Heckman, E. A. C., Ruby, E. G. and McFall-Ngai, M. J. (2018). Persistent Interactions with Bacterial Symbionts Direct Mature-Host Cell Morphology and Gene Expression in the Squid-Vibrio Symbiosis. *mSystems* **3**, 1–17.

Lathlean, J. A., Ayre, D. J. and Minchinton, T. E. (2014). Estimating latitudinal variability in extreme heat stress on rocky intertidal shores. *J. Biogeogr.* **41**, 1478–1491.

Lau, N. S., Ting, S. Y., Sam, K. K., Janaranjani, M., Wong, S. C., Wu, X., Waiho, K., Fazhan, H. and Shu-Chien, A. C. (2022). Comparative Analyses of Scylla olivacea Gut Microbiota Composition and Function Suggest the Capacity for Polyunsaturated Fatty Acid Biosynthesis. *Microb. Ecol.* 1–14.

Le Moullac, G., Cheize, M., Gastineau, O., Daniel, J. Y., Le Coz, J. R., Huvet, A., Moal, J., Pouvreau, S., Van Wormhoudt, A. and Samain, J. F. (2008). Ecophysiological and metabolic adaptations to sulphide exposure of the oyster Crassostrea gigas. *J. Shellfish Res.* **27**, 355–363.

Lee, Y. Y. and Lee, J. C.-Y. (2018). LC-MS/MS Analysis of Lipid Oxidation Products in Blood and Tissue Samples. In *Clinical Metabolomics: Methods and Protocols* (ed. Giera, M.), p. Springer Science+Business Media, LLC.

Lee, Y. Y., Galano, J. M., Leung, H. H., Balas, L., Oger, C., Durand, T. and Lee, J. C. Y. (2020). Nonenzymatic oxygenated metabolite of docosahexaenoic acid, 4(RS)-4-F4t-neuroprostane, acts as a bioactive lipid molecule in neuronal cells. *FEBS Lett.* **594**, 1797–1808.

Lent, C. M. (1969). Adaptations of the ribbed mussel, Modiolus demissus (Dillvvyn), to the intertidal habitat. *Integr. Comp. Biol.* **9**, 283–292.

Lenth, R. V. (2023). emmeans: Estimated Marginal Means, aka Least-Squares Means.

Letendre, J., Chouquet, B., Manduzio, H., Marin, M., Bultelle, F., Leboulenger, F. and Durand, F. (2009). Tidal height influences the levels of enzymatic antioxidant defences in Mytilus edulis. *Mar. Environ. Res.* **67**, 69–74.

Leung, K. S., Galano, J. M., Durand, T. and Lee, J. C. Y. (2015). Current development in non-enzymatic lipid peroxidation products, isoprostanoids and isofuranoids, in novel biological samples. *Free Radic. Res.* **49**, 816–826.

Li, Y. F., Yang, N., Liang, X., Yoshida, A., Osatomi, K., Power, D., Batista, F. M. and Yang, J. L. (2018). Elevated seawater temperatures decrease microbial diversity in the

gut of Mytilus coruscus. *Front. Physiol.* **9**, 839.

Li, J., Bates, K. A., Hoang, K. L., Hector, T. E., Knowles, S. C. L. and King, K. C.
(2022). Experimental temperatures shape host microbiome diversity and composition.
Glob. Chang. Biol. **00**, 1–16.

Lin, H. and Peddada, S. Das (2020). Analysis of compositions of microbiomes with bias
correction. *Nat. Commun.* **11**, 1–11.

Loeffler, C., Berger, S., Guy, A., Durand, T., Bringmann, G., Dreyer, M., Von Rad, U.,
Durner, J. and Mueller, M. J. (2005). B1-phytoprostanes trigger plant defense and
detoxification responses. *Plant Physiol.* **137**, 328–340.

Lokmer, A. and Mathias Wegner, K. (2015). Hemolymph microbiome of Pacific oysters in
response to temperature, temperature stress and infection. *ISME J.* **9**, 670–682.

Longford, S. R., Campbell, A. H., Nielsen, S., Case, R. J., Kjelleberg, S. and Steinberg,
P. D. (2019). Interactions within the microbiome alter microbial interactions with host
chemical defences and affect disease in a marine holobiont. *Sci. Rep.* **9**, 1–13.

Lund, I. and Steenfeldt, S. J. (2011). The effects of dietary long-chain essential fatty acids
on growth and stress tolerance in pikeperch larvae (Sander lucioperca L.). *Aquac. Nutr.*
17, 191–199.

Ma, K., Chen, S., Wu, Y., Ma, Y., Qiao, H., Fan, J. and Wu, H. (2022). Dietary
supplementation with microalgae enhances the zebrafish growth performance by
modulating immune status and gut microbiota. *Appl. Microbiol. Biotechnol.* **106**, 773–
788.

Maher, R. L., Schmeltzer, E. R., Meiling, S., McMinds, R., Ezzat, L., Shantz, A. A.,
Adam, T. C., Schmitt, R. J., Holbrook, S. J., Burkepile, D. E., et al. (2020). Coral
Microbiomes Demonstrate Flexibility and Resilience Through a Reduction in
Community Diversity Following a Thermal Stress Event. *Front. Ecol. Evol.* **8**, 555698.

Mallott, E. K., Skovmand, L. H., Garber, P. A. and Amato, K. R. (2022). *The fecal*
metabolome of black howler monkeys (Alouatta pigra) varies in response to seasonal
dietary changes .

Marangon, E., Laffy, P. W., Bourne, D. G. and Webster, N. S. (2021). Microbiome-
mediated mechanisms contributing to the environmental tolerance of reef invertebrate
species. *Mar. Biol.* **168**, 1–18.

Marshall, D. J. and McQuaid, C. D. (2011). Warming reduces metabolic rate in marine
snails: Adaptation to fluctuating high temperatures challenges the metabolic theory of
ecology. *Proc. R. Soc. B Biol. Sci.* **278**, 281–288.

Marshall, D. J. and McQuaid, C. D. (2020). Metabolic Regulation, Oxygen Limitation and
Heat Tolerance in a Subtidal Marine Gastropod Reveal the Complexity of Predicting
Climate Change Vulnerability. *Front. Physiol.* **11**, 1106.

Marshall, D. J., Dong, Y., McQuaid, C. D. and Williams, G. A. (2011). Thermal
adaptation in the intertidal snail Echinolittorina malaccana contradicts current theory by
revealing the crucial roles of resting metabolism. *J. Exp. Biol.* **214**, 3649–3657.

McDevitt-Irwin, J. M., Garren, M., McMinds, R., Vega Thurber, R. and Baum, J. K.
(2019). Variable interaction outcomes of local disturbance and El Niño-induced heat
stress on coral microbiome alpha and beta diversity. *Coral Reefs* **38**, 331–345.

McKnight, D. T., Huerlimann, R., Bower, D. S., Schwarzkopf, L., Alford, R. A. and
Zenger, K. R. (2019). Methods for normalizing microbiome data: An ecological
perspective. *Methods Ecol. Evol.* **10**, 389–400.

Mcmahon, B. R. (1988). Physiological responses to oxygen depletion in intertidal animals.
Integr. Comp. Biol. **28**, 39–53.

McMurdie, P. J. and Holmes, S. (2013). Phyloseq: An R Package for Reproducible
Interactive Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**,.

Miller, E., Morel, A., Saso, L. and Saluk, J. (2014). Isoprostanes and neuroprostanes as biomarkers of oxidative stress in neurodegenerative diseases. *Oxid. Med. Cell. Longev.* **2014**,.

Moghadam, N. N., Thorshauge, P. M., Kristensen, T. N., de Jonge, N., Bahrndorff, S., Kjeldal, H. and Nielsen, J. L. (2018). Strong responses of Drosophila melanogaster microbiota to developmental temperature. *Fly (Austin).* **12**, 1–12.

Muñoz, K., Flores-Herrera, P., Gonçalves, A. T., Rojas, C., Yáñez, C., Mercado, L., Brokordt, K. and Schmitt, P. (2019). The immune response of the scallop Argopecten purpuratus is associated with changes in the host microbiota structure and diversity. *Fish Shellfish Immunol.* **91**, 241–250.

Nakagawa, S., Saito, H., Tame, A., Hirai, M., Yamaguchi, H., Sunata, T., Aida, M., Muto, H., Sawayama, S. and Takaki, Y. (2017). Microbiota in the coelomic fluid of two common coastal starfish species and characterization of an abundant Helicobacter-related taxon. *Sci. Rep.* **7**, 1–10.

Narrowe, A. B., Albuthi-Lantz, M., Smith, E. P., Bower, K. J., Roane, T. M., Vajda, A. M. and Miller, C. S. (2015). Perturbation and restoration of the fathead minnow gut microbiome after low-level triclosan exposure. *Microbiome* **3**, 1–18.

Nearing, J. T., Douglas, G. M., Hayes, M., Macdonald, J., Desai, D., Allward, N., Jones, C. M. A., Wright, R., Dhanani, A., Comeau, A. M., et al. (2022). Microbiome differential abundance methods produce disturbingly different results across 38 datasets. *Nat. Commun.* **13**, 342.

Neu, A. T., Hughes, I. V., Allen, E. E. and Roy, K. (2021). Decade-scale stability and change in a marine bivalve microbiome. *Mol. Ecol.* **30**, 1237–1250.

Ng, T. P. T., Lau, S. L. Y., Seuront, L., Davies, M. S., Stafford, R., Marshall, D. J. and Williams, G. A. (2017). Linking behaviour and climate change in intertidal ectotherms: insights from littorinid snails. *J. Exp. Mar. Bio. Ecol.* **492**, 121–131.

Ng, T. P. T., Lau, S. L. Y., Davies, M. S., Stafford, R., Seuront, L., Hutchinson, N., Hui, T. T. Y. and Williams, G. A. (2021). Behavioral repertoire of high-shore littorinid snails reveals novel adaptations to an extreme environment. *Ecol. Evol.* **11**, 7114–7124.

Nguyen, Q. P., Karagas, M. R., Madan, J. C., Dade, E., Palys, T. J., Morrison, H. G., Pathmasiri, W. W., McRitche, S., Sumner, S. J., Frost, H. R., et al. (2021). Associations between the gut microbiome and metabolome in early life. *BMC Microbiol.* **21**, 1–19.

Nicastro, K. R., Zardi, G. I., McQuaid, C. D., Stephens, L., Radloff, S. and Blatch, G. L. (2010). The role of gaping behaviour in habitat partitioning between coexisting intertidal mussels. *BMC Ecol.* **10**,.

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Solymos, P., Stevens, M. H. H., Szoecs, E., et al. (2022). vegan: community ecology package. *R Packag. version 2.6-2*.

Paix, B., Othmani, A., Debroas, D., Culioli, G. and Briand, J. F. (2019). Temporal covariation of epibacterial community and surface metabolome in the Mediterranean seaweed holobiont Taonia atomaria. *Environ. Microbiol.* **21**, 3346–3363.

Pal, V. K., Bandyopadhyay, P. and Singh, A. (2018). Hydrogen sulfide in physiology and pathogenesis of bacteria and viruses. *IUBMB Life* **70**, 393–410.

Parris, D. J., Morgan, M. M. and Stewart, F. J. (2019). Feeding rapidly alters microbiome composition and gene transcription in the clownfish gut. *Appl. Environ. Microbiol.* **85**, 1–15.

Pernet, F., Tremblay, R., Redjah, I., Sévigny, J. M. and Gionet, C. (2008). Physiological and biochemical traits correlate with differences in growth rate and temperature adaptation among groups of the eastern oyster Crassostrea virginica. *J. Exp. Biol.* **211**,

1047   969–977.

Pierce, M. L. and Evan, J. (2019). Gut microbiomes of the eastern oyster (Crassostrea virginica) and the blue mussel (Mytilus edulis): temporal variation and the influence of marine aggregate-associated microbial communities. *mSphere* **4**, e00730-19.

Pimentel, Z. T., Dufault-Thompson, K., Russo, K. T., Scro, A. K., Smolowitz, R. M., Gomez-Chiarri, M. and Zhang, Y. (2021). Microbiome Analysis Reveals Diversity and Function of Mollicutes Associated with the Eastern Oyster, Crassostrea virginica. *mSphere* **6**, e00227-21.

Pita, L., Rix, L., Slaby, B. M., Franke, A. and Hentschel, U. (2018). The sponge holobiont in a changing ocean: from microbes to ecosystems. *Microbiome* **6**, 46.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J. and Glöckner, F. O. (2013). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Res.* **41**, D590–D596.

Rädecker, N., Pogoreutz, C., Gegner, H. M., Cárdenas, A., Perna, G., Geißler, L., Roth, F., Bougoure, J., Guagliardo, P., Struck, U., et al. (2022). Heat stress reduces the contribution of diazotrophs to coral holobiont nitrogen cycling. *ISME J.* **16**, 1110–1118.

Raffaelli, D. and Hawkins, S. (1999). *Intertidal ecology*. 2nd ed. Kluwer Academic Publishers.

Rivera-Ingraham, G. A., Rocchetta, I., Meyer, S. and Abele, D. (2013). Oxygen radical formation in anoxic transgression and anoxia-reoxygenation: Foe or phantom? Experiments with a hypoxia tolerant bivalve. *Mar. Environ. Res.* **92**, 110–119.

Rocca, J. D., Simonin, M., Blaszczak, J. R., Ernakovich, J. G., Gibbons, S. M., Midani, F. S. and Washburne, A. D. (2019). The Microbiome Stress Project: Toward a global meta-analysis of environmental stressors and their effects on microbial communities. *Front. Microbiol.* **9**, 3272.

Roy, J., Galano, J. M., Durand, T., Le Guennec, J. Y. and Lee, J. C. Y. (2017). Physiological role of reactive oxygen species as promoters of natural defenses. *FASEB J.* **31**, 3729–3745.

Samakraman, S., Williams, G. A. and Ganmanee, M. (2010). Spatial and Temporal Variability of Intertidal Rocky Shore Molluscs in Sichang Island, East Coast of Thailand. *Publ. Seto Mar. Biol. Lab.* **10**, 35–46.

Santoro, E. P., Borges, R. M., Espinoza, J. L., Freire, M., Messias, C. S. M. A., Villela, H. D. M., Pereira, L. M., Vilela, C. L. S., Rosado, J. G., Cardoso, P. M., et al. (2021). Coral microbiome manipulation elicits metabolic and genetic restructuring to mitigate heat stress and evade mortality. *Sci. Adv.* **7**, 19–21.

Sargent, J. R. (1976). The Structure, Metabolism and Function of Lipids in Marine organisms. In *Biochemical and biophysical perspectives in marine biology* (ed. Malins, D. C.) and Sargent, J. R.), pp. 149–212. New York: Academic Press.

Shankar, V., Homer, D., Rigsbee, L., Khamis, H. J., Michail, S., Raymer, M., Reo, N. V. and Paliy, O. (2015). The networks of human gut microbe-metabolite associations are different between health and irritable bowel syndrome. *ISME J.* **9**, 1899–1903.

Sharp, C. and Foster, K. R. (2022). Host control and the evolution of cooperation in host microbiomes. *Nat. Commun.* **13**, 1–15.

Sharp, K. H., Pratte, Z. A., Kerwin, A. H., Rotjan, R. D. and Stewart, F. J. (2017). Season, but not symbiont state, drives microbiome structure in the temperate coral Astrangia poculata. *Microbiome* **5**, 120.

Shaver, E. C., Shantz, A. A., McMinds, R., Burkepile, D. E., Thurber, R. L. V. and Silliman, B. R. (2017). Effects of predation and nutrient enrichment on the success and microbiome of a foundational coral. *Ecology* **98**, 830–839.

Shiu, J. H., Keshavmurthy, S., Chiang, P. W., Chen, H. J., Lou, S. P., Tseng, C. H.,

Justin Hsieh, H., Allen Chen, C. and Tang, S. L. (2017). Dynamics of coral-associated
bacterial communities acclimated to temperature stress based on recent thermal history.
Sci. Rep. **7**, 1–13.

Soen, Y. (2014). Environmental disruption of host-microbe co-adaptation as a potential
driving force in evolution. *Front. Genet.* **5**, 168.

Sokolova, I. M. and Pörtner, H. O. (2001). Physiological adaptations to high intertidal life
involve improved water conservation abilities and metabolic rate depression in littorina
saxatilis. *Mar. Ecol. Prog. Ser.* **224**, 171–186.

Somero, G. N. (2002). Thermal physiology and vertical zonation of intertidal animals:
Optima, limits, and costs of living. *Integr. Comp. Biol.* **42**, 780–789.

Sorek, M., Díaz-Almeyda, E. M., Medina, M. and Levy, O. (2014). Circadian clocks in
symbiotic corals: The duet between Symbiodinium algae and their coral host. *Mar.*
Genomics **14**, 47–57.

Stillman, J. H. (2002). Causes and consequences of thermal tolerance limits in rocky
intertidal porcelain crabs, genus Petrolisthes. *Integr. Comp. Biol.* **42**, 790–796.

Storey, K. B. and Storey, J. M. (2004). Oxygen Limitation and Metabolic Rate Depression.
In *Functional Metabolism: Regulation and Adaptation*, p. John Wiley & Sons, Inc.

Sussarellu, R., Fabioux, C., Camacho Sanchez, M., Le Goïc, N., Lambert, C., Soudant,
P. and Moraga, D. (2012). Molecular and cellular response to short-term oxygen
variations in the Pacific oyster Crassostrea gigas. *J. Exp. Mar. Bio. Ecol.* **412**, 87–95.

Thoma, I., Loeffler, C., Sinha, A. K., Gupta, M., Krischke, M., Steffan, B., Roitsch, T.
and Mueller, M. J. (2003). Cyclopentenone isoprostanes induced by reactive oxygen
species trigger defense gene activation and phytoalexin accumulation in plants. *Plant J.*
34, 363–375.

Tignat-Perrier, R., van de Water, J. A. J. M., Guillemain, D., Aurelle, D., Allemand, D.
and Ferrier-Pagès, C. (2022). The Effect of Thermal Stress on the Physiology and
Bacterial Communities of Two Key Mediterranean Gorgonians. *Appl. Environ.*
Microbiol. **88**,.

Vandamme, P. and De Ley, J. (1991). Proposal for a new family, Campylobacteraceae. *Int.*
J. Syst. Bacteriol. **41**, 451–455.

Voolstra, C. R. and Ziegler, M. (2020). Adapting with Microbial Help: Microbiome
Flexibility Facilitates Rapid Responses to Environmental Change. *BioEssays* **42**, 1–9.

Walke, J. B., Becker, M. H., Krinos, A., Chang, E. A. B., Santiago, C., Umile, T. P.,
Minbiole, K. P. C. and Belden, L. K. (2021). Seasonal changes and the unexpected
impact of environmental disturbance on skin bacteria of individual amphibians in a
natural habitat. *FEMS Microbiol. Ecol.* **97**, 1–14.

Wang, J., Ren, R. M. and Yao, C. L. (2018). Oxidative stress responses of Mytilus
galloprovincialis to acute cold and heat during air exposure. *J. Molluscan Stud.* **84**, 285–
292.

Wang, J., Ma, L. X. and Dong, Y. W. (2022). Coping with harsh heat environments:
molecular adaptation of metabolic depression in the intertidal snail Echinolittorina
radiata. *Cell Stress Chaperones*.

Webster, N. S., Cobb, R. E. and Negri, A. P. (2008). Temperature thresholds for bacterial
symbiosis with a sponge. *ISME J.* **2**, 830–842.

Weiss, S., Xu, Z. Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., Lozupone, C.,
Zaneveld, J. R., Vázquez-Baeza, Y., Birmingham, A., et al. (2017). Normalization
and microbial differential abundance strategies depend upon data characteristics.
Microbiome **5**, 1–18.

Welker, A. F., Moreira, D. C., Campos, É. G. and Hermes-Lima, M. (2013). Role of
redox metabolism for adaptation of aquatic animals to drastic changes in oxygen

availability. *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.* **165**, 384–404.

Wickham, H. (2008). *ggplot2: Elegant Graphics for Data Analysis*. New York, USA:
Springer Science+Business Media, LCC.

Williams, G. A. and Morritt, D. (1995). Habitat partitioning and thermal tolerance in a
tropical limpet, Cellana grata. *Mar. Ecol. Prog. Ser.* **124**, 89–103.

Yin, X., Chen, P., Chen, H., Jin, W. and Yan, X. (2017). Physiological performance of the
intertidal Manila clam (Ruditapes philippinarum) to long-term daily rhythms of air
exposure. *Sci. Rep.* **7**, 41648.

Yonny, M. E., Rodríguez Torresi, A., Cuyamendous, C., Réversat, G., Oger, C., Galano,
J. M., Durand, T., Vigor, C. and Nazareno, M. A. (2016). Thermal Stress in Melon
Plants: Phytoprostanes and Phytofurans as Oxidative Stress Biomarkers and the Effect of
Antioxidant Supplementation. *J. Agric. Food Chem.* **64**, 8296–8304.

Zaneveld, J. R., McMinds, R. and Thurber, R. V. (2017). Stress and stability: Applying the
Anna Karenina principle to animal microbiomes. *Nat. Microbiol.* **2**,.

Zhou, K., Aertsen, A. and Michiels, C. W. (2014). The role of variable DNA tandem
repeats in bacterial adaptation. *FEMS Microbiol. Rev.* **38**, 119–141.

Zhukova, N. V. (2019). Fatty acids of marine mollusks: Impact of diet, bacterial symbiosis
and biosynthetic potential. *Biomolecules* **9**, 1–25.

Ziegler, M., Seneca, F. O., Yum, L. K., Palumbi, S. R. and Voolstra, C. R. (2017).
Bacterial community dynamics are linked to patterns of coral heat tolerance. *Nat.*
Commun. **8**, 1–8.

1168

1169 Supplementary information

1170

1171

1172 **Fig. S1. Rarefaction curves illustrating the ASVs captured against the sampling depth**
1173 **which was generated for each sample using their corresponding quality-filtered V3-V4**
1174 **16SrRNA sequences**.

1175

1176

1177

Fig. S2. Variation in the alpha diversity indices of the *I. nucleus*-associated microbiome
can be explained by the host body temperature. Linear regression graph showing the
relationship between average body temperature (from 10 random independent replicates
collected at each timepoint) and (A) ASV richness (B) Shannon-Wiener index (H'). The
datapoints shown in each panel were the samples' replicates (n= 4-5) from their corresponding
timepoint. The shaded grey area indicates a 95% confidence interval for the predicted
regression line. Statistical result (least square regression analysis) is shown in the upper right
within each panel.

1186

1187 Table S1. Metadata describing the *Isognomon nucleus* samples' collected information

Date	Time	Timepoints label	Replicates	Sample name	Body (mm)
22.05.2019	0915	morning	1	ARM1	12.9
22.05.2019	0915	morning	2	ARM2	13.5
22.05.2019	0915	morning	3	ARM3	15.2
22.05.2019	0915	morning	4	ARM4	12.5
22.05.2019	0915	morning	5	ARM5	13.7
22.05.2019	1230	noon	1	BRM1	11.7
22.05.2019	1230	noon	2	BRM2	11.3
22.05.2019	1230	noon	3	BRM3	12.9
22.05.2019	1230	noon	4	BRM4	11.9
22.05.2019	1230	noon	5	BRM5	12.4
22.05.2019	1612	early-afternoon	1	CRM1	11.7
22.05.2019	1612	early-afternoon	2	CRM2	12.9
22.05.2019	1612	early-afternoon	3	CRM3	13.0
22.05.2019	1612	early-afternoon	4	CRM4	12.0
22.05.2019	1612	early-afternoon	5	CRM5	11.7
22.05.2019	1740	late-afternoon	1	DRM1	11.8
22.05.2019	1740	late-afternoon	2	DRM2	10.9
22.05.2019	1740	late-afternoon	3	DRM3	14.9
22.05.2019	1740	late-afternoon	4	DRM4	11.1
22.05.2019	1740	late-afternoon	5	DRM5	11.9

1188