Reconstructing cellular differentiation networks and identifying cell fate–determining features with CIBER

Longchen Xu[1,2,#], TingTing Cong[3,#], Hengyi Xu[4,#], Sijie Yang[1], Rahul Sinha, Ryo Yamamoto, Wei Zhang[1,2], Jianwei Wang[3,*] and Xun Lan[1,5,6,*]

[1]Tsinghua-Peking Joint Center for Life Sciences, Tsinghua University, Beijing 100084, China

[2]School of Life Sciences, Tsinghua University, Beijing 100084, China

[3]School of Pharmaceutical Sciences, Tsinghua University, Beijing 100084, China

[4]Eight-Year MD Program, Peking Union Medical College, Beijing 100084, China

[5]Department of Basic Medical Science, School of Medicine, Tsinghua University, Beijing 100084, China

[6]MOE Key Laboratory of Bioinformatics, Tsinghua University, Beijing 100084, China

[#]These authors contributed equally to this work

[*]Correspondence: jianweiwang@mail.tsinghua.edu.cn (J.W.), xlan@tsinghua.edu.cn (X.L.)

Abstract

Most existing trajectory inference methods are similarity-based and lack an understanding of the latent causality among differentiation processes. Here, we present CIBER, a <u>C</u>ausal <u>I</u>nference–<u>B</u>ased framework for the <u>E</u>valuation of feature effects and the <u>R</u>econstruction of cellular differentiation networks. CIBER is suitable for various types of omics data, including profilings of gene expression and chromatin accessibility. We show that CIBER can capture known cell-lineage structures and identify potential novo differentiation branches. By combining the CIBER-learned network with the structural causal model and applying *in silico* perturbation, we constructed an effect-matrix that quantifies the impacts of different features on each branch. We subsequently identified features important to the network structure through differentiation driver feature (DDF) analysis. We demonstrated that DDF analysis can identify features crucial to haematopoiesis but show no significant

29 difference in expression between lineages. We further experimentally verified the effect
30 of transcription factor *Bcl11b* on haematopoiesis, showing that it promotes lymphoid
31 lineage differentiation while suppressing myeloid lineage differentiation, which is
32 consistent with our predictions.

Introduction

Cellular differentiation is one of the most fundamental processes in biology. Single-cell
omics data have provided unprecedented opportunities for elucidating developmental
pathways and reconstructing cell lineages. Several computational trajectory inference
methods have been developed for studying cellular developmental dynamics. In most of
these methods, cells are ordered along deterministic[1–5] or probabilistic[6,7] trajectories
based on similarities in their expression patterns, and a numeric value known as the
pseudotime is assigned to each cell to indicate how far it has progressed along a dynamic
process of interest. These similarity-based trajectory inference methods tend to identify
correlations between different cell states; however, the process of cellular differentiation
is driven by its causal nature and thus may not match these methods properly.

Recently, a series of approaches[8–11] for cell fate prediction based on RNA velocity[12]
have emerged, which model the differentiation process with the spliced-to-unspliced
mRNA ratio in the form of differential equations, thereby restoring considerable cell
temporal information, and can then be utilized for causal gene regulatory network
reconstruction[13]. However, RNA velocity is limited to representing cell dynamics in the
transcriptome space[14] and is not suitable for other types of data. More general approaches
are required to identify causality between cellular states, not only through transcriptomic
profiling but also through different types of data, such as epigenomic profiling or
microarray assays.

Methods including Granger causality[15–17], Boolean networks[18–20] and Bayesian

networks[21–24] have been effectively applied for causal inference in studying biological

processes such as gene regulatory networks and protein-protein interactions. However,

the application of these causal inference methods is rare in scenarios such as the

reconstruction of cellular differentiation networks and the identification of cell fate–

determining features. Bayesian networks (BNs) provide a computationally efficient

approach for inferring causality among a set of variables and have been applied to infer

cellular networks[21], protein-protein interactions[22], protein-signalling networks[23], gene

regulatory networks[24,25], and long-range intrachromosomal interactions[26]. However, data-

based inferences of feasible BNs are challenging because the structure learning of BNs is

sensitive to parameters and requires considerable prior knowledge to obtain the expected

structure.

Here, we present CIBER, a novo causal inference–based framework, to reconstruct the

cellular differentiation networks and to identify cell fate–determining features. CIBER

provides a hybrid structure learning algorithm for detecting information flow among

different cell types and is suitable for both gene expression and chromatin accessibility

data. In addition to single-cell transcriptomic and epigenome sequencing data, which

assume contiguous cell states and are critical for typical trajectory inference methods,

CIBER can use microarray or bulk sequencing data in which the states of various cell

types are discrete. We then adopt the framework of the structural causal model[27,28] to the

directed acyclic graph (DAG) structure of the learned cellular differentiation network,

utilize *in silico* perturbation as interventions for causal inference, and build an effect

matrix that quantifies the impacts of different features on each branch of the cellular

differentiation network. Furthermore, differentiation driver feature (DDF) analysis is

defined to identify important features in the cellular differentiation process. In contrast to

regular differential feature analysis, in which the identified features are markers of each

cell type, DDFs capture the features that significantly influence the entire differentiation

process or a specific lineage. We demonstrated that DDF analysis can identify

differentiation-driving genes that are not differentially expressed among lineages and can

complement the regular differential feature analysis to help us understand the biological

processes comprehensively.

We applied CIBER framework to study haematopoiesis using various types of omics

or microarray data and obtained expected cell differentiation trajectories. The application

on a mouse microarray dataset with the most cell types showed that CIBER can identify

potential branches that disagreed with the conventional hierarchy while reported in

previous studies. In this study, we use regular differential feature (hereafter named RDF)

analysis to represent analysis identifying important features by directly comparing the

values in cell × feature matrices, such as differentially expressed gene analysis with

scRNA-seq datasets or differential deviation-score transcription factor (TF) analysis with

scATAC-seq datasets. In contrast, DDF analysis is performed by identifying important

driving features based on edge × feature matrices inferred by CIBER from the cell ×

feature matrices. DDF analysis enriches the techniques for identifying important features

related to cellular differentiation process and can complement RDF analysis to help us

understand biological processes more comprehensively. Through DDF analysis, CIBER

identifies cell fate–determining features that are not deemed statistically significant in

RDF analysis. We experimentally validated CIBER's prediction of the lymphoid-biased

effect of TF B-cell lymphoma/leukaemia 11B (*Bcl11b*).

Results

CIBER offers a robust Bayesian network structure learning strategy. Bayesian

networks are directed acyclic graphical models in which nodes represent variables and

edges represent probabilistic dependencies between nodes[29]. One major issue in Bayesian

network structure learning is that the obtained graph is extremely sensitive to parameters

and the observed data; thus, small changes in input may result in considerably different

outputs. To obtain a cellular differentiation network well-constructed for downstream

analysis, extensive expert knowledge is usually required as prior information to adjust the

learned structures. Here, we aim to develop a new framework that can generate robust

Bayesian network structures given minimal prior knowledge, such that the proposed

method can be generalized to investigate understudied biological processes.

The existing Bayesian network structure learning algorithms can be divided into three

categories: constraint-based algorithms, score-based algorithms and hybrid

algorithms[30,31]. Constraint-based algorithms construct a Bayesian network by first

generating an undirected skeleton via Markov blanket[32] or correlation and then orienting

the skeleton under acyclicity constraints to find the directed acyclic graph that best

satisfies the conditional independence test. The effectiveness and efficiency of constraint-

based algorithms are limited due to the lack of robustness and the high complexity of the

conditional independence test. Score-based algorithms evaluate how well a DAG fits the

observed data with a score such as the Akaike information criterion[33] or Bayesian

information criterion[34], with the goal of identifying the graph obtaining the highest score.

Major issues for score-based algorithms include the exponentially growing search space

as the number of variables increases and the search algorithm may lead to a local optimum.

One solution to the limitations of the abovementioned two types of algorithms is hybrid

algorithms, which first generate an undirected skeleton and then search for the DAG with

the highest score among the structures restricted to this skeleton. However, the result of

regular hybrid algorithms is sensitive to the parameters, and considerable prior knowledge

is required to obtain a proper structure.

To address the abovementioned issues and strengthen the robustness of Bayesian

network structure learning, we present in CIBER framework a new hybrid algorithm that

uses a mixed strategy of resampling over single cells and then combining the BNs learned

within each resampling (Fig. 1). The increasing size of single-cell omics data provides

hundreds or thousands of replicates of each cell type, allowing us to comprehensively

observe cell state transitions. CIBER takes an expression matrix as input and outputs a

feasible structure. A stratified resampling strategy is used to select random subsets of cells

to construct the network, resulting in a set of networks with various structures.

Specifically, for each subset of cells, we utilized the centroids of different cell types to

construct a Bayesian network, with each centroid representing the mean expression level

of its cell type. These centroids are connected to generate an undirected skeleton

according to Spearman's correlation or the mutual information between them, with each

edge representing the correlation between the two connected cell types. Several

algorithms for correlation estimation, such as CMI2NI[35] and SPACE[36], can be chosen

under the support of GeNeCK[37], an R framework for constructing gene regulatory

networks. Based on the undirected skeleton, we utilized the hill climbing algorithm

through the bnlearn[38] package to search for the DAG structure with the highest Bayesian

information criterion.

Fig. 1 | Simplified schematic of CIBER framework. The upper part shows the pipeline of the hybrid Bayesian network structure learning algorithm starting from cell resampling. Random subsets of cells are selected, and various DAGs are learned to represent the causation between the centroids of different cell types. Each DAG obtained within one resampling is referred to as a one-shot DAG for convenience in this study. Then, a combination strategy is applied over the one-shot DAGs to obtain the final Bayesian network. The bottom part shows the pipeline to generate effect matrix and to identify differentiation driving-genes. From the perspective of the structure causal model, a series of linear regressions are performed on the learned Bayesian network, with each child node taken as a dependent variable and its parent nodes taken as independent variables. A mixed strategy of resampling and interventions of *in silico* perturbation on a specific feature is utilized to obtain a pair of distributions for each edge, representing the reference and the intervened linear regression coefficients. The effect matrix is defined to quantify the difference between the two distributions and each element in the matrix reflects the effect of a feature on a differentiation branch. DDF analysis is performed on the effect matrix to identify cell fate–determining features.

The learned DAG is affected by the threshold used to evaluate correlations while constructing the undirected skeleton. We used a set of sequential correlation thresholds, such as from 0 to 1 in steps of 0.01, to obtain a series of DAGs varying from dense to

sparse. Approximate upper and lower boundaries for the number of edges in the network

can be assigned according to the expected complexity of the network. DAGs with the

number of edges within the specified boundaries are selected and combined to generate a

directed graph in which each edge is weighted according to the frequency at which it

appears. Each weight reveals the robustness of an edge, thereby reflecting the confidence

of the causation between the nodes it connects. However, the graph is not necessarily

acyclic and should be trimmed and binarized to produce an unweighted DAG that

preserves the most robust structure within this batch of cells. This DAG is named one-

shot DAG hereafter because it is obtained within one resampling. This process is then

repeated for multiple times to generate a set of one-shot DAGs, over which a second

combination is applied to generate a final DAG that captures the most consistent causation

between variables (see Methods for detailed description). In addition to single-cell omics

data, CIBER is suitable for microarray and bulk sequencing data, but without resampling

due to limited replicates.

CIBER captures known differentiation structures in various datasets. To test

whether CIBER can obtain well-established cell lineage structures, we applied it to study

haematopoiesis with various datasets, including mouse microarray data[39–41], mouse bone

marrow scRNA-seq data[42], human bone marrow scRNA-seq data[43] and human bone

marrow scATAC-seq data[44]. The mouse microarray dataset contains comprehensive

blood cell types, from which we selected a subset of representative types of progenitors

to reconstruct the early haematopoietic hierarchy. The subset includes 11 cell types —

haematopoietic stem cell (HSC), multipotent progenitor subset A (MPPa), multipotent

progenitor subset B (MPPb), common myeloid progenitor (CMP), common lymphoid

progenitor (CLP), granulocyte-macrophage progenitor (GMP), megakaryocyte-erythroid

progenitor (MEP), megakaryocyte progenitor (MkP), precolony-forming unit erythroid

(pCFU-E), B-cell-biased lymphoid progenitor (BLP) and double negative 1 thymocytes

(DN1) — each containing 3 or 4 biological replicates. In this microarray dataset, the

expression levels of 21,678 genes are detected by 45,101 probes[39–41]. If a gene is

simultaneously detected by several probes, we set its expression level to the median

among all the probes. Because of the stable expression levels over the limited size of

replicates, we omitted resampling and performed the analysis on a set of fixed points that

are the centroids obtained by averaging the expression level over biological replicates for

each cell type.

The results revealed that CIBER can extract robust lineage structures and provide

information on potential differentiation branches (Fig. 2b). Compared to the results

obtained with a typical hybrid algorithm consisting of mutual information and hill

climbing (hereafter named MI-hc), the networks derived from CIBER conformed more

to the conventional haematopoietic hierarchy and captured the bifurcation structure more

accurately (Fig. 2b, c). In these figures, the edges contradicting prior knowledge are

coloured in red. MI-hc is sensitive to parameters, and small changes in the correlation

threshold can result in considerably different structures (Extended Data Fig. 1), while the

structure obtained by CIBER is robust without the need to manually set the exact

correlation threshold. In addition to the well-known branches of the HSC→MPPa/MPPb

→lymphoid/myeloid lineages and those inside the lymphoid or myeloid lineage, CIBER

identified three unconventional interlineage edges that were reported to be possible in

previous studies, CLP→GMP[45] and CMP→BLP/DN1[46,47].

The application of CIBER to the single-cell omics datasets followed the entire pipeline,

including the resampling procedure. The structures learned by CIBER again identified

larger numbers and higher percentages of conventional edges compared to the results of

the typical hybrid algorithm MI-hc (Fig. 2d-f, Extended Data Fig. 2). Furthermore, the

cell types in CIBER-learned structures were in relative positions consistent with the low-

dimensional landscapes of diffusion maps[42], SPRING[43] or PCA[44] obtained in previous

studies.

Fig. 2 | CIBER captures conventional differentiation structures in various datasets. a, Conventional haematopoietic hierarchy. **b**, CIBER result of the mouse microarray dataset, with the blue/red-shaded part representing the lymphoid/myeloid lineage. **c**, Best-tuned structure learned with MI-hc, a typical hybrid algorithm consisting of mutual information and hill climbing, using the same dataset as **b**; the structures learned under other correlation thresholds are shown in Extended Data Fig. 1. The dashed edges represent unconventional interlineage branches that were reported to be possible in previous studies, while the red edges represent less credible branches. **b** and **c** show that the CIBER result is consistent with conventional knowledge with a larger number and a higher percentage of edges and captures the bifurcation structure more accurately than MI-hc. **d-f**, Low-dimensional landscapes[43], the CIBER and MI-hc results of the human scRNA-seq dataset. Results for more datasets are shown in Extended Data Fig. 2.

CIBER quantifies the effects of features on the cellular differentiation hierarchy.

The structural causal model (SCM) is a conventional causal inference framework that combines probabilistic graphical models and the notion of interventions[27,28]. Similar to Bayesian networks, SCM considers a set of associated observables in the form of a directed acyclic graph while viewing each node as a function of its parents in the graph. A natural restriction of SCM is the assumption of a linear additive noise model, in which the functions connecting each node to its parents are limited to the form of linear

regression (Methods). In the SCM framework, interventions can be performed by

changing the form of particular functions or the distribution of specific nodes to infer the

effect of causality.

Here, we introduced the concept of SCM into the directed acyclic graph learned by

CIBER and performed causal inference by applying *in silico* perturbation to the

expression of specific features as interventions (Fig. 1). Specifically, permutation of

features was performed on single-cell profiling data while deletion was performed on

microarray or bulk sequencing data. Then, we recorded the coefficient associated with

each edge in the linear additive SCM with/without interventions performed to a specific

feature and defined an effect matrix to measure the change in the coefficient due to the

interventions. Intuitively, the effect matrix quantifies the impacts of the features on each

edge in the cellular differentiation network, with a positive value implying a possible

promoting function on the corresponding differentiation branch and a negative value

implying a possible depressing function.

Differentiation driver feature (DDF) analysis identifies features determining cell

fate. Next, we developed DDF analysis, a novel method distinct from regular differential

feature (RDF) analysis, to identify important features crucial to the cellular differentiation

process via effect matrices obtained by *in silico* perturbation. Specifically, we defined

diffScores to assess the effect of individual features on the cellular differentiation network

or specific lineages. The diffScore of a gene on the entire network was defined by

summarizing the absolute values related to this gene in the effect matrix, while the

diffScore on a specific lineage was defined as the sum of the absolute values related to

the edges pointing to cell types in the lineage subtracting the original values of the edges

pointing from the lineage to other lineages (see Methods for detailed description).

DDF analysis provides a framework to systematically investigate a feature's impact on

the entire cellular differentiation hierarchy. Our study demonstrates that DDF analysis

can work effectively for scenarios in which cell types are limited. Additionally, DDF

analysis can be combined with RDF analysis to better identify important features due to

the independent information captured by the two methods.

To examine the effectiveness of DDF analysis, we first applied it to the mouse bone

marrow scRNA-seq dataset and compared the results with that of the RDF analysis

obtained by ranking genes according to the p value of Wilcoxon's signed-rank tests. To

compare the results of the two analyses, we selected the same number of top genes on the

two lists (Fig. 3a). The results showed that a few important cell fate–determining genes

identified by RDF analysis, such as *Gata2*[48], were not captured by DDF analysis; on the

other hand, DDF analysis identified important genes that were not differentially expressed

between the lymphoid and myeloid cell types, such as *Mpl*[49], *Ccl9*[50] and *Elane*[51];

nevertheless, some genes, such as *Flt3*[52], *Cd34*[53] and *Dntt*[54], were identified by both

analyses. We further conducted a literature search on the top-ranked genes and calculated

Spearman's correlation between the two gene lists (Fig. 3b-d). The results showed that

the RDF and DDF analyses both captured a part of the genes that have been proven to be

important in haematopoiesis.

Fig. 3 | DDF identifies cell fate–determining features. The plots show the results on RDF and DDF gene lists obtained with the mouse scRNA-seq dataset. **a**, Features are allocated by log fold change and log p value obtained in the RDF analysis, from which the top differentially expressed genes are coloured, as well as the same number of DDF-selected features important to the entire network. **b**, The number of shared top features between the two feature sets. **c**, Literature search result shows that the DDF and RDF analyses identify different important features, in which the red parts represent the reported features. **d**, Correlation between the ranked lists shows that the DDF analysis captures different information from the RDF analysis. The tiles represent the top 25%, 25~50%, 50~75% and bottom 25% features in RDF or DDF ranking lists. **e**, GSEA normalized enrichment scores (NES) with terms related to haematopoiesis or haematopoietic differentiation in the GO and KEGG databases. DDF list obtains higher NES than RDF on all the terms. Dots in the left part represent the NES of RDF and DDF ranking lists, while the distributions show the background NES given randomly ranked gene lists. (Hemo., Hemopoiesis; Diff., Differentiation; Lym., Lymphoid; Mye., Myeloid; Reg., Regulation; Pos., Positive; Neg., Negative.)

Then, we evaluated the behaviours of RDF and DDF lists on enriching important genes related to haematopoiesis or haematopoietic differentiation. Specifically, we calculated the normalized enrichment scores (NES) through GSEA for gene lists with respect to all

the GO and KEGG terms related to haemopoiesis or haematopoietic differentiation (Fig.

3e). GSEA results showed that DDF surpassed RDF on almost all these pathways on NES.

Note that the dataset did not contain any lymphoid-lineage cell types other than

lymphomyeloid-primed progenitor (LMPP), an early lymphoid-biased progenitor. The

results demonstrated that DDF analysis performs well on datasets with insufficient cell

types, due potentially to the fact that it examines the entire cellular differentiation network

rather than being restricted to the difference between subgroups of cells, as RDF analysis

did.

Next, we applied the same analyses to the human scRNA-seq dataset and mouse

microarray dataset. Similar results were obtained for the human scRNA-seq dataset to the

results of mouse scRNA-seq dataset described above (Extended Data Fig. 3). The

microarray dataset offered us an opportunity to study the behaviour of DDF analysis in

scenarios where most cell types are available and are therefore friendly to RDF analysis.

The result showed the RDF list performed slightly better than the DDF list in 12 out of

22 terms (Extended Data Fig. 4e). However, the literature search still showed that DDF

analysis identified important genes which are not differentially expressed, demonstrating

that DDFs can complement RDFs to better identify genes crucial to cell differentiation

(Extended Data Fig. 4a-c).

CIBER identifies driver TFs of biological processes. We then applied CIBER and DDF

analysis to a human bone marrow scATAC-seq dataset[44], and identified TFs crucial to

haematopoiesis based on deviation score matrix calculated with chromVAR[55]. RDF

analysis was also performed to identify the TFs with significant difference in deviation

score between lymphoid and myeloid cell types. Top TFs identified by DDF only includes,

IRF8, *BCL11A*, *SPI1*, *SPIB* and *CEBPB*; TFs, such as *TCF3*, *LYL1* and *EBF1*, were found

by the RDF analysis alone, while both analyses uncovered TFs such as *GATA1*, *GATA2*,

GATA3, *ID3* and *LMO2* (Fig. 4c). With lineage-specific diffScores, DDF analysis can

further quantify the potential effects of each TF on promoting or suppressing specific

lineages (Fig. 4d).

TF regulates cellular function by orchestrating the transcription of downstream target

genes. Next, we sought to test whether the target genes of the TFs identified in the

scATAC-seq dataset were also enriched in the DDF or RDF analysis of an independent

human bone marrow scRNA-seq dataset. we collected target genes for TFs uncovered by

both analysis from 5 databases (TRRUST[56], JASPAR[57], ChEA[58], Motifmap[59] and

ENCODE[60]) and ranked the target genes according to their significance in the DDF or

RDF analysis of the scRNA-seq dataset. Two types of metrics, normalized gene set

enrichment score (NES) and area under the receiver operating characteristic curve (AUC),

were calculated to measure the enrichment of targeted genes by each ranking algorithm

(Fig. 4e, Methods). When calculating AUC, the receiver operating characteristic (ROC)

curve was drawn by assigning TF-targeted genes as positive and all other genes as

negative samples. Background distributions of NES and AUC were obtained with 1,000

randomly shuffled ranking lists of all the genes. Results show that under both metrics,

DDF surpasses RDF on all gene sets targeted by different TFs (Fig. 4e, f, Extended Data

Fig. 5). For example, both DDF and RDF significantly surpass the random background,

and DDF scores higher for the *GATA1* target gene set; in the case of *BCL11B*, only the

DDF score is significantly higher than the random background (DDF NES p value=0.009,

RDF NES p value=0.103, DDF AUC p value<0.001, RDF p value=0.683). CIBER

quantifies the importance of TF target genes on cellular differentiation, thus provides

insight into the potential mechanism impacting the cellular differentiation process.

Fig. 4 | CIBER identifies driver TFs of biological processes. a, PCA reduction of a human bone marrow scATAC-seq dataset[44]. **b**, Differentiation hierarchy learned with CIBER. **c**, Top TFs identified by RDF and DDF analyses. Among them, several reported important TFs[44] are annotated. **d**, Lineage-specific diffScores quantify the effects of each TF on haematopoiesis under a higher resolution of promoting or suppressing specific lineages. **e**, Comparison of rankings obtained with DDF and RDF analyses using ROC or GSEA analyses applied with the set of genes regulated by each TF. AUC and GSEA NES results show DDF surpasses RDF on each gene set and significantly surpasses random background; background is set as the distribution of 1,000 random gene rankings. **f**, AUC and NES Results on *GATA1*-targeted and *BCL11B*-targeted gene sets respectively. For *GATA1*-targeted gene set, both DDF and RDF significantly surpass random background, and DDF scores higher; for *BCL11B*-targeted gene set, DDF significantly surpasses random background, while no statistical difference exists between RDF and the background (DDF NES p value=0.009, RDF NES p value=0.103, DDF AUC p value<0.001, RDF p value=0.683). Thus, compared with regular differential analysis, CIBER can better identify important TFs and enrich reported regulated genes, providing insight into potential regulated genes through which a TF influences the process of haematopoiesis.

In vivo experiments validate *BCL11B* as a crucial TF to fate determination in

haematopoiesis. In addition to the previously reported TFs, we noted that DDF analysis

of bone marrow scATAC-seq data identified *BCL11B* which showed no significant

difference in deviation scores between lymphoid and myeloid cell types (Fig. 4c).

Lineage-specific diffScores indicate that *IRF8*, *BCL11A* and *BCL11B* promote the

differentiation process of the lymphoid lineage while suppress the myeloid lineage,

consistent with previous report of *IRF8*[61] and *BCL11A*[62,63] (Fig. 4d). GSEA and ROC

analyses of bone marrow scRNA-seq demonstrated that *BCL11B* target genes were

enriched in DDF analysis, providing further evidence that the TF may influence the

process of haematopoiesis (Fig. 4e, f).

To explore the effect of *Bcl11b* on the differentiation of haematopoietic stem cells, we

constructed *Bcl11b*-deficient mice (*Bcl11b*[flox/flox], Vav-iCre[+]), hereafter named *Bcl11b*-

KO, to specifically knock out *Bcl11b* in the blood cells. Results showed that *Bcl11b*-KO

mice had a shorter survival time than WT mice in our cohort (Fig. 5b). Compared with

WT mice, the haematopoietic system of *Bcl11b*-KO mice after 5-fluorouracil (5-FU)

treatment had an increased proportion of reconstituted myeloid cells and a decreased

proportion of reconstituted lymphoid cells (Fig. 5d).

Fig. 5 | Experimental validation of CIBER predictions for *Bcl11b*. Survival experiments
(**a**) and results (**b**) of WT and *Bcl11b*-KO mice in their native state show the survival capacity
of *Bcl11b*-KO mice is hampered. Experiments (**c**) and percentages (**d**) of reconstituted B cells,
T cells and myeloid cells in peripheral blood of WT and *Bcl11b*-KO mice after 5-FU treatment.
e, Scheme of the transplantation strategy for *Bcl11b*-KO mice. Fifty HSCs freshly isolated
from 2-month-old WT or *Bcl11b*-KO mice were transplanted into lethally irradiated recipients
along with 3×10^5 competitor cells. Chimaera in peripheral blood was checked every month.
f, Lineage distribution of B cells, T cells, and myeloid cells among donor-derived cells in
peripheral blood four months after transplantation. (n=4-5 mice per group. Data are shown
as the mean ± SD.). **g**, Line plots display the contribution of donor cells to peripheral blood
overall (CD45.2+), B cells (B220+), T cells (CD3+) and myeloid cells (Mac-1+) every month
after transplantation. (n=4-5 mice per group. Data are shown as the mean ± SD.)

To evaluate the impact of *Bcl11b*-deficiency on the long-term haematopoietic
reconstitution capacity and differentiation of HSCs, we further performed competitive
transplantation assays. Fifty freshly isolated HSCs from WT or *Bcl11b*-KO mice were
injected into lethally irradiated recipient mice together with 3×10^5 total bone marrow cells
as a competitor (Fig. 5e). Chimaera in peripheral blood was evaluated every month until
4 months. Peripheral blood analysis of recipient mice every month calculated the
percentage of cells reconstructed by donor-drives HSCs to all these in recipient peripheral
blood and showed that the reconstitution capacity of *Bcl11b*-KO HSCs was severely
impaired in various cells, including B, T and myeloid lineages, compared to WT
transplanting the same number of HSCs (Fig. 5g). At the same time, by analysing the
composition of donor-derived cells in recipient mice, we found that *Bcl11b*-deficient
HSCs enhanced myeloid differentiation and decreased T-cell differentiation (Fig. 5f). In
summary, the knockout of *Bcl11b* resulted in a reduced reconstitution capacity of HSCs,
however, in the offspring cells, the proportion of the myeloid lineage is increased
compared to that of the lymphoid lineage.

Discussion

Large-scale single-cell omics data have provided unprecedented opportunities to study
developmental pathways and have enabled the reconstruction of lineage relationships
between cells. Multiple computational trajectory inference methods have been developed
to investigate cellular developmental dynamics based on snapshot measurements. Most
trajectory inference methods are based on expression pattern similarities and provide cell
fate probabilities. However, cellular differentiation processes are complex causation-
driven phenomena that are naturally more suited for analysis with causal inference
methods. The existing approaches for inferring causality among cellular states are mainly
based on RNA velocity, while other methods, such as those based on Granger causality,

Boolean networks and Bayesian networks, have been effectively used for causality

inference in other biological processes, such as gene regulatory networks.

In this study, we present CIBER framework based on causal inference to reconstruct

cellular differentiation networks and identify cell fate–determining features. CIBER

improved the robustness of existing hybrid Bayesian network structure learning

algorithms to obtain feasible structures with minimal prior knowledge. The applications

of CIBER on several datasets of haematopoietic cells show that it is suitable to dataset

obtained through various techniques on different species and that it may help in

identifying potential differentiation branches. CIBER built an effect matrix based on the

concept of the structural causal model and interventions of *in silico* perturbation on the

expression level of specific features. The effect matrix quantifies the impacts of features

on differentiation branches and can be used to identify the cell fate–determining features

through DDF analyses developed in this study. Unlike regular differential feature (RDF)

analysis, which identifies markers of specific cell types, DDF analysis captures distinctive

information on differentiation lineages. The applications on different datasets revealed

that DDF analysis was effective in scenarios in which cell types were limited for RDF

analysis, and that DDF analysis could complement RDF analysis to identify important

features comprehensively.

CIBER differs from existing trajectory inference methods in several aspects in terms

of reconstructing the cellular differentiation process. Most existing trajectory inference

methods are suitable only for single-cell omics data on the assumption that the static

snapshot of the cellular states is continuously distributed, while CIBER is applicable

under more conditions, such as when the number of cells or cell types is limited, and is

even applicable to datasets obtained by microarray or bulk sequencing techniques. Except

for the approaches based on RNA velocity, most existing trajectory inference methods

are similarity-based, while CIBER is a causation-based framework that provides a new

perspective on the cellular differentiation process.

Although CIBER can handle scenarios that existing trajectory inference methods

cannot, it has several limitations. To better perform the analysis after building the effect

matrix, at least two parents should exist for each child node in the obtained network. In

addition, CIBER may not be suitable for cases in which the network is too simple,

especially when the network is only a two-layer hierarchy, which prevents the

construction of the effect matrix for the same reason as the previous problem and even

impacts the robustness of BN structure learning. For extremely complex networks with

too many nodes and edges, CIBER may be time consuming. Typically, CIBER is most

effective for hierarchical structures with approximately 5 layers and less than 20 nodes.

Taken together, the CIBER framework provides a causal perspective for the

reconstruction of cellular differentiation networks and the evaluation of how features

influence differentiation branches, and DDF analysis enriches existing techniques for

identifying important features and can be combined with RDF analysis to help us

understand biological processes more comprehensively.

References

1. Haghverdi, L., Büttner, M., Wolf, F. A., Buettner, F. & Theis, F. J. Diffusion
pseudotime robustly reconstructs lineage branching. *Nat Methods* **13**, 845–
848 (2016).
2. Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell
trajectories. *Nature Methods* **14**, 979–982 (2017).
3. Setty, M. *et al.* Wishbone identifies bifurcating developmental trajectories
from single-cell data. *Nat Biotechnol* **34**, 637–645 (2016).
4. Welch, J. D., Hartemink, A. J. & Prins, J. F. SLICER: inferring branched,
nonlinear cellular trajectories from single cell RNA-seq data. *Genome*
Biology **17**, 106 (2016).
5. Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are
revealed by pseudotemporal ordering of single cells. *Nature Biotechnology*
32, 381–386 (2014).

6. Schiebinger, G. *et al.* Optimal-Transport Analysis of Single-Cell Gene
Expression Identifies Developmental Trajectories in Reprogramming. *Cell*
176, 928-943.e22 (2019).
7. Setty, M. *et al.* Characterization of cell fate probabilities in single-cell data
with Palantir. *Nature Biotechnology* **37**, 451–460 (2019).
8. Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing
RNA velocity to transient cell states through dynamical modeling. *Nat*
Biotechnol **38**, 1408–1414 (2020).
9. Lange, M. *et al.* CellRank for directed single-cell fate mapping. *Nat*
Methods **19**, 159–170 (2022).
10. Qiu, X. *et al.* Mapping transcriptomic vector fields of single cells. *Cell* **185**,
690-711.e45 (2022).
11. Li, C., Virgilio, M. C., Collins, K. L. & Welch, J. D. Multi-omic single-cell
velocity models epigenome–transcriptome interactions and improves cell
fate prediction. *Nat Biotechnol* 1–12 (2022) doi:10.1038/s41587-022-
01476-y.
12. La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498
(2018).
13. Qiu, X. *et al.* Inferring Causal Gene Regulatory Networks from Coupled
Single-Cell Expression Dynamics Using Scribe. *Cell Systems* **10**, 265-
274.e11 (2020).
14. Bergen, V., A. Soldatov, R., V. Kharchenko, P. & J. Theis, F. RNA
velocity—current challenges and future perspectives. *Molecular Systems*
Biology **17**, e10282 (2021).
15. Deshpande, A., Chu, L.-F., Stewart, R. & Gitter, A. Network inference with
Granger causality ensembles on single-cell transcriptomics. *Cell Reports*
38, 110333 (2022).
16. Finkle, J. D., Wu, J. J. & Bagheri, N. Windowed Granger causal inference
strategy improves discovery of gene regulatory networks. *Proceedings of*
the National Academy of Sciences **115**, 2252–2257 (2018).
17. Papili Gao, N., Ud-Dean, S. M. M., Gandrillon, O. & Gunawan, R.
SINCERITIES: inferring gene regulatory networks from time-stamped
single cell transcriptional expression profiles. *Bioinformatics* **34**, 258–266
(2018).
18. Font-Clos, F., Zapperi, S. & La Porta, C. A. M. Classification of triple-
negative breast cancers through a Boolean network model of the epithelial-
mesenchymal transition. *Cell Systems* **12**, 457-462.e4 (2021).
19. Hamey, F. K. *et al.* Reconstructing blood stem cell regulatory network
models from single-cell molecular profiles. *Proc Natl Acad Sci USA* **114**,
5822–5829 (2017).
20. Weidner, F. M. *et al.* Capturing dynamic relevance in Boolean networks
using graph theoretical measures. *Bioinformatics* **37**, 3530–3537 (2021).
21. Friedman, N. Inferring Cellular Networks Using Probabilistic Graphical
Models. *Science* **303**, 799–805 (2004).
22. Jansen, R. A Bayesian Networks Approach for Predicting Protein-Protein
Interactions from Genomic Data. *Science* **302**, 449–453 (2003).

23. Sachs, K., Perez, O., Pe'er, D., Lauffenburger, D. A. & Nolan, G. P. Causal Protein-Signaling Networks Derived from Multiparameter Single-Cell Data. *Science* **308**, 523–529 (2005).

24. Liu, F., Zhang, S.-W., Guo, W.-F., Wei, Z.-G. & Chen, L. Inference of Gene Regulatory Network Based on Local Bayesian Networks. *PLOS Computational Biology* **12**, e1005024 (2016).

25. Schütte, J. *et al.* An experimentally validated network of nine haematopoietic transcription factors reveals mechanisms of cell state stability. *eLife* **5**, e11469 (2016).

26. Zhang, X., Branciamore, S., Gogoshin, G., Rodin, A. S. & Riggs, A. D. Analysis of high-resolution 3D intrachromosomal interactions aided by Bayesian network modeling. *PNAS* **114**, E10359–E10368 (2017).

27. Pearl, J. Causal diagrams for empirical research. *Biometrika* **82**, 669–688 (1995).

28. Schölkopf, B. *et al.* Toward Causal Representation Learning. *Proceedings of the IEEE* **109**, 612–634 (2021).

29. Korb, K. & Nicholson, A. Bayesian Artificial Intelligence. in *Computer science and data analysis series* (2004). doi:10.5860/choice.41-5948.

30. Scutari, M. & Denis, J.-B. *Bayesian Networks: With Examples in R*. (Chapman and Hall/CRC, 2021). doi:10.1201/9780429347436.

31. Koller, D. & Friedman, N. *Probabilistic Graphical Models: Principles and Techniques*. (MIT Press, 2009).

32. Pearl, J. *Probabilistic reasoning in intelligent systems: networks of plausible inference*. (Morgan kaufmann, 1988).

33. Akaike, H. A new look at the statistical model identification. *IEEE transactions on automatic control* **19**, 716–723 (1974).

34. Gideon Schwarz. Estimating the Dimension of a Model. *The Annals of Statistics* **6**, 461–464 (1978).

35. Zhang, X., Zhao, J., Hao, J.-K., Zhao, X.-M. & Chen, L. Conditional mutual inclusive information enables accurate quantification of associations in gene regulatory networks. *Nucleic Acids Research* **43**, e31 (2015).

36. Peng, J., Wang, P., Zhou, N. & Zhu, J. Partial Correlation Estimation by Joint Sparse Regression Models. *Journal of the American Statistical Association* **104**, 735–746 (2009).

37. Zhang, M. *et al.* GeNeCK: a web server for gene network construction and visualization. *BMC Bioinformatics* **20**, 12 (2019).

38. Scutari, M. Learning Bayesian Networks with the bnlearn R Package. *Journal of Statistical Software* **35**, 1–22 (2010).

39. Seita, J. *et al.* Gene Expression Commons: an open platform for absolute gene expression profiling. *PLoS One* **7**, e40321 (2012).

40. Fathman, J. W. *et al.* Upregulation of CD11A on hematopoietic stem cells denotes the loss of long-term reconstitution potential. *Stem Cell Reports* **3**, 707–715 (2014).

41. Ambrosi, T. H. *et al.* Aged skeletal stem cells generate an inflammatory degenerative niche. *Nature* **597**, 256–262 (2021).

42. Nestorowa, S. *et al.* A single-cell resolution map of mouse hematopoietic stem and progenitor cell differentiation. *Blood* **128**, e20–e31 (2016).

43. Pellin, D. *et al.* A comprehensive single cell transcriptional landscape of
human hematopoietic progenitors. *Nature Communications* **10**, 2395
(2019).
44. Buenrostro, J. D. *et al.* Integrated Single-Cell Analysis Maps the
Continuous Regulatory Landscape of Human Hematopoietic Differentiation.
Cell **173**, 1535-1548.e16 (2018).
45. Graf, T. Differentiation plasticity of hematopoietic cells. *Blood* **99**, 3089–
3101 (2002).
46. Akashi, K., Traver, D., Miyamoto, T. & Weissman, I. L. A clonogenic
common myeloid progenitor that gives rise to all myeloid lineages. *Nature*
404, 193–197 (2000).
47. Prohaska, S. S., Scherer, D. C., Weissman, I. L. & Kondo, M.
Developmental plasticity of lymphoid progenitors. *Seminars in Immunology*
14, 377–384 (2002).
48. Single-cell ATAC-seq reveals GATA2-dependent priming defect in myeloid
and a maturation bottleneck in lymphoid lineages | Blood Advances |
American Society of Hematology.
https://ashpublications.org/bloodadvances/article/5/13/2673/475734/Single-
cell-ATAC-seq-reveals-GATA2-dependent.
49. The cytoplasmic domain of Mpl receptor transduces exclusive signals in
embryonic and fetal hematopoietic cells | Blood | American Society of
Hematology.
https://ashpublications.org/blood/article/100/6/2063/235068/The-
cytoplasmic-domain-of-Mpl-receptor-transduces.
50. ICSBP-mediated immune protection against BCR-ABL–induced leukemia
requires the CCL6 and CCL9 chemokines | Blood | American Society of
Hematology.
https://ashpublications.org/blood/article/113/16/3813/25078/ICSBP-
mediated-immune-protection-against-BCR-ABL.
51. CRISPR Mediated ELANE Single-Allele Knock-out Restores Proliferation
and Myeloid Differentiation of Neutropenia Patient Derived BM HSCs |
Blood | American Society of Hematology.
https://ashpublications.org/blood/article/136/Supplement%201/23/470437/C
RISPR-Mediated-ELANE-Single-Allele-Knock-out.
52. Sitnicka, E. *et al.* Key role of flt3 ligand in regulation of the common
lymphoid progenitor but not in maintenance of the hematopoietic stem cell
pool. *Immunity* **17**, 463–472 (2002).
53. Okuno, Y. *et al.* Differential regulation of the human and murine CD34
genes in hematopoietic stem cells. *Proceedings of the National Academy of*
Sciences **99**, 6246–6251 (2002).
54. Ng, S. Y.-M., Yoshida, T., Zhang, J. & Georgopoulos, K. Genome-wide
Lineage-Specific Transcriptional Networks Underscore Ikaros-Dependent
Lymphoid Priming in Hematopoietic Stem Cells. *Immunity* **30**, 493–507
(2009).
55. Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR:
inferring transcription-factor-associated accessibility from single-cell
epigenomic data. *Nat Methods* **14**, 975–978 (2017).

56. Han, H. *et al.* TRRUST v2: an expanded reference database of human and
mouse transcriptional regulatory interactions. *Nucleic Acids Research* **46**,
D380–D386 (2018).
57. Castro-Mondragon, J. A. *et al.* JASPAR 2022: the 9th release of the open-
access database of transcription factor binding profiles. *Nucleic Acids*
Research **50**, D165–D173 (2022).
58. Lachmann, A. *et al.* ChEA: transcription factor regulation inferred from
integrating genome-wide ChIP-X experiments. *Bioinformatics* **26**, 2438–
2444 (2010).
59. Xie, X., Rigor, P. & Baldi, P. MotifMap: a human genome-wide map of
candidate regulatory motif sites. *Bioinformatics* **25**, 167–174 (2009).
60. Consortium, T. E. P. A User's Guide to the Encyclopedia of DNA Elements
(ENCODE). *PLOS Biology* **9**, e1001046 (2011).
61. Differential IRF8 Transcription Factor Requirement Defines Two Pathways
of Dendritic Cell Development in Humans. *Immunity* **53**, 353-370.e8 (2020).
62. Luc, S. *et al.* Bcl11a Deficiency Leads to Hematopoietic Stem Cell Defects
with an Aging-like Phenotype. *Cell Reports* **16**, 3181–3194 (2016).
63. Tsang, J. C. H. *et al.* Single-cell transcriptomic reconstruction reveals cell
cycle and multi-lineage differentiation defects in Bcl11a-deficient
hematopoietic stem cells. *Genome Biol* **16**, 178 (2015).
638

Methods

Data preprocessing. Normalized mouse microarray, mouse bone marrow scRNA-seq,

human bone marrow scRNA-seq and human bone marrow scATAC-seq datasets were

obtained from the data sources. In the microarray dataset, each cell type had 3 or 4

biological replicates, and the expression levels of 21,678 genes were detected by 45,101

probes. For each gene detected by multiple probes, the median value was adopted as its

expression level. The top variable features were first selected for Bayesian network

structure learning; 3,000 features were selected in the mouse microarray dataset, 5,000 in

the mouse bone marrow scRNA-seq dataset, 4,000 in the human bone marrow scRNA-

seq dataset and 1,764 (all transcription factors) in the human bone marrow scATAC-seq

dataset.

Construction of cellular differentiation networks. In general, Bayesian network

structure learning is thought to be sensitive to both observations and parameters, and

small changes in input can lead to considerably different outputs. Hence, expert

experience is needed as prior knowledge to obtain a structure applicable for further study.

In CIBER framework, we adopted several strategies to ensure a well-structured hierarchy

with minimal prior knowledge of assigning credible upper and lower boundaries for the

number of edges in the network and setting stem cells as the root.

CIBER utilizes a multistep strategy to obtain robust BN structures — resampling,

undirected skeleton learning, DAG learning, combining DAGs and trimming branches.

Particularly, for datasets with limited biological replicates, such as microarray datasets,

the resampling step is omitted. By combining the DAGs obtained with different

resampled subsets of single cells and various parameters, we can extract the most robust

associations between variables, which have a higher probability of being the true

causation. The details of the steps are described below.

Resampling strategy. The resampling strategy maximizes the efficacy of large single-

cell omics datasets. A stratified resampling strategy was performed for the single-cell

omics datasets to generate a subset of cells, while this strategy was omitted for the

microarray dataset. For the microarray dataset and each subset of single cells after

resampling, we calculated the mean expression level of each cell type to obtain a mean

expression matrix, with each column representing a cell type and each row representing

a specific feature. The resampling ratio can be set to approximately 0.3 to prevent high

randomness or homogeneity between the obtained subsets. Similarly, the cell number of

each cell type was suggested to be greater than 50 to prevent random effects.

Undirected skeleton learning. With the mean expression matrix as an input, an

undirected skeleton was built to describe the correlation between each pair of cell types,

with each node representing a cell type and each edge representing the correlation

between the connected cell types. Various metrics can be utilized to evaluate the

correlation, such as Spearman's correlation or the mutual information. In addition to the

correlation, several algorithms developed for undirected skeleton construction can be
conveniently chosen under the support of GeNeCK[37]. The default algorithm used in
CIBER is CMI2NI[35], an approach that was developed for reverse-engineering gene
regulatory networks by quantifying the conditional mutual inclusive information between
two genes given a third gene by calculating the Kullback–Leibler divergence. The
undirected skeleton is determined by the threshold used for judging the correlation, and a
sparser skeleton will be obtained given a higher correlation threshold. A set of sequential
correlation thresholds was used to obtain a series of undirected skeletons for further DAG
learning.

DAG learning. Typically, two causally associated variables should also be correlated.
The undirected skeleton provides us with a reference that can be used to determine the
causal DAG. We excluded absent edges in the undirected skeleton in the alternative
DAGs and utilized the existing score-based structure learning algorithm offered in the
bnlearn[38] package to obtain the DAGs. The hill climbing algorithm was used to search
for the DAG with the highest Bayesian information criterion (BIC)[34]. BIC is commonly
used in Bayesian network structure learning to evaluate how well a graph G fits the
observed data D and can be formulated as follows:

695
$$\text{BIC}(G{:}D) = LL(G{:}D) - \|G\| \cdot \log|\text{D}|/2,$$

where $LL(G{:}D)$ describes the likelihood that G fits D, and $\|G\|$ and $\log|\text{D}|$ represent the
complexity of the graph structure and dataset, respectively. Thus, hill climbing seeks to
identify the least complex DAG that best explains the observed data.

Combination of DAGs and trimming branches. A DAG can be generated with each
correlation threshold under a resampled subset of cells. By changing the correlation
threshold, we can obtain a series of different DAGs. The DAGs with the number of edges

located between the prior-set upper and lower boundaries are selected and combined to

obtain a directed graph, with each edge weighted according to its frequency of occurrence.

According to the assumption that the true causation should be robust to different

parameters, a combined weighted DAG can be generated by extracting the highest

weighted edges and removing the lowest weighted edges in cycles.

Then, this DAG is binarized to obtain an unweighted DAG named one-shot DAG in

this study, which indicates the robust causation within this batch of cells. For microarray

or bulk sequencing datasets, the one-shot DAG is the final learned Bayesian network

structure. However, for single-cell omics datasets, the one-shot DAG reflects only the

possible causation within a specific subset of the whole dataset. Therefore, a similar

combination is applied over different one-shot DAGs to generate a quasi-final DAG.

Under some special conditions, when generating the quasi-final DAG, additive prior

knowledge may be needed to determine which edge should be severed if two edges with

the same weights occur in one cycle. The quasi-final DAG may contain some edges that

are not robust while being kept for the reason that their opposite edges occur even less.

An optional procedure can be performed on the quasi-final DAG to limit the maximum

number of parents for each node and to remove redundant low-weighted edges.

Specifically, under the structural causal model framework, we fit the quasi-final DAG to

the mean expression matrix of all the cells in the original dataset twice, once with the top

25% variable features and once with the top 25~50% variable features, and maintain the

edges that remain stable while removing edges with low-weighted coefficients.

Construction of the effect matrix. Based on the learned Bayesian network, the SCM

framework can be utilized to build an effect matrix that quantifies the effects of different

features on various differentiation branches. SCM is a conventional causal inference

framework that combines probabilistic graphical models and the notion of interventions.

This framework considers a set of associated observables with the nodes of a directed

acyclic graph while viewing each node X_i as a function of its parents $\mathbf{PA}_i$ in the graph:

729
$$X_i = h_i(\mathbf{PA}_i, U_i),$$

730   where U_i represents noise or exogenous variables. Under a linear additive noise

731   assumption, X_i can be viewed partially as a linear regression of $\mathbf{PA}_i$:

732
$$X_i = \sum_k \beta_{ik} \mathbf{PA}_{ik} + U_i.$$

Interventions are performed to infer causality based on the SCM equations. Typically, by

intervening in the form of a function or distribution of $\mathbf{PA}_i$, a latent expression of X_i can

be obtained. The difference between the observed and latent expressions of X_i is

measured to estimate the effect of causality.

In the construction of the effect matrix, we perform interventions by *in silico*

perturbation of permuting (or deleting) the expression of a specific feature in all cell types

and estimate the effect of causality by concentrating on the change in β_{ik} instead of the

change in X_i. The coefficient β_{ik} represents the association between two connected cell

types and reflects the information flowing from $\mathbf{PA}_{ik}$ to X_i and the information of the

corresponding differentiation branch. Thus, the change in the coefficients caused by

intervening in a specific feature quantifies the effect of that feature on the cellular

differentiation process.

For the single-cell omics dataset, we used a resampling strategy similar to that

performed in reconstructing networks to record the distribution of the coefficient

associated with a particular edge e_0 obtained without and with interventions on a specific

feature f_0, which we termed α_{ref} and α_{inter}, respectively, where α represents the

abovementioned β_{ik}. For one resampling and intervention, the matrix $\text{EffectMat}_{\text{sampling}}$

is defined as

751
$$\text{EffectMat}_{\text{samping}}[f_0, e_0] = \text{effect}(f_0, e_0) = \alpha_{\text{ref}} - \alpha_{\text{inter}},$$

where $\text{EffectMat}_{\text{samping}}[f_0, e_0]$ is the row f_0 and column e_0 element in the matrix, and

$\text{effect}(f_0, e_0)$ is the effect of feature f_0 on edge e_0. By resampling and permuting several

times, we obtain a series of $\text{EffectMat}_{\text{samping}}$ matrices, which can be horizontally merged

to generate a final effect matrix in which the rows consist of all features and the columns

consist of all edges multiplied by the resampling and permutation times. For microarray

or bulk sequencing datasets, the effect matrix is constructed with the same procedures,

except that the resampling step is omitted and the intervention is performed with deletion

instead of permutation.

DDF analysis. We developed DDF analysis to identify features important to specific

lineages or the entire cellular differentiation network according to the effect matrix. When

applying DDF analysis, diffScores are defined to evaluate the effects of different features

on the differentiation process or a specific lineage.

For the haematopoiesis network, we defined diffSum, $\text{diffScore}_{\text{lym}}$ and $\text{diffScore}_{\text{myl}}$ to

quantify the effects of the features on the entire haematopoietic hierarchy, the lymphoid-

lineage and the myeloid-lineage, respectively. Let e represent an edge; the effect of

feature f_0 on the entire haematopoietic cellular differentiation network G is defined as

768
$$\text{diffSum} = \sum_{e \in G} |\text{effect}(f_0, e)|.$$

769   The lymphoid-lineage and myeloid-lineage effects are defined as

770
$$\text{diffScore}_{\text{lym}} = \sum_{e \in \{S \to L\} \cup \{L \to L\}} |\text{effect}(f_0, e)| - \sum_{e \in \{L \to M\}} \text{effect}(f_0, e)$$

771　　and

$$\text{diffScore}_{\text{myl}} = \sum_{e \in \{S \to M\} \cup \{M \to M\}} |\text{effect}(f_0, e)| - \sum_{e \in \{M \to L\}} \text{effect}(f_0, e),$$

where S represents early cell types, including stem cells and multipotent cells, L

represents lymphoid-lineage cell types, M represents myeloid-lineage cell types, and

$\{S \to L\}$ represents the set of edges pointing from cell types in set S to cell types in set L.

Transcription factor–targeted gene set GSEA and ROC analyses. For each TF, we

collect the regulated genes from 5 databases (TRRUST, JASPAR, ChEA, Motifmap and

ENCODE) and apply GSEA and receiver operating characteristic (ROC) analyses on this

gene set. GSEA and ROC analyses are performed with the ranked gene lists obtained by

DDF and RDF analyses.

Specifically, normalized enrichment score (NES) is calculated in GSEA analysis.

When applying ROC analysis, we plot ROC curve by assigning the TF-targeted genes as

positive and the others as negative samples. In ROC curve, the true positive rate is referred

to as sensitivity and the true negative rate is referred to as specificity,

$$\text{Sensitivity} = \frac{\text{True Positive}}{\text{True Positive} + \text{False Negative}}$$

786　　and

$$\text{Specificity} = \frac{\text{True Negative}}{\text{True Negative} + \text{False Positive}}.$$

AUC score is calculated as the area under ROC curve.

Background distributions are obtained by calculating NES and AUC of 1,000 randomly

ordered lists of all the genes.

Competitive HSC transplantation and analysis. *Bcl11b*[flox/flox] mice were provided by

Dr. Yeguang Chen from Tsinghua University. In all experiments (unless otherwise

specified), mice were homozygous for the *Bcl11b* allele. *Bcl11b*-KO (C57BL/6, CD45.2)

mice, C57BL/6-SJL (CD45.1) (Stock No: 002014) and C57BL/6 WT (CD45.2) (Stock

No: 000664) mice were obtained from the Jackson Laboratory. CD45.1/2 mice were

heterozygotes from CD45.1 and CD45.2 mice. All mice were kept in specific-pathogen-

free (SPF), AAALAC-accredited animal care facilities at the Laboratory Animal

Research Center, Tsinghua University and all procedures were approved by the

Institutional Animal Care and Use Committee of Tsinghua University.

For the primary competitive HSC transplantation assay, 30 freshly isolated HSCs from

WT or *Bcl11b*-KO mice (CD45.2) were transplanted into lethally irradiated (10 Gy)

recipients (CD45.1/2) together with 3×10^5 competitor cells (CD45.1). Donor-derived

chimaerism (including B cells, T cells and myeloid cells) in the peripheral blood of

recipients was analysed at monthly intervals.

Two-tailed unpaired Student's *t* test was applied for analysing experiment data after

testing for normal distribution. All histogram and line graphs were plotted by GraphPad

Prism 8 software, and p value <0.05 was regarded as significant for all tests. All data are

depicted as the mean ± SD.

Data availability

The datasets used in this work were downloaded. Mouse microarray dataset[39–41] can be

found at GEO (GSE34723). Mouse bone marrow scRNA-seq data[42] can be found at GEO

(GSE81682). Human bone marrow scRNA-seq data[43] can be found at GEO (GSE117498).

Human bone marrow scATAC-seq data[44] can be found at GEO (GSE96772).

Code availability

CIBER toolkit is implemented in R and is available on Bioconductor and GitHub

(https://github.com/Lan-lab/CIBER). Tutorials of CIBER usage cases are also available

through GitHub (https://github.com/Lan-lab/CIBER/tutorials).

Acknowledgements

819 The authors are grateful to the High-Performance Computation platforms of Tsinghua

820 University for providing computational resources.

Author Contributions

Funding

824 This work was supported by National Natural Science Foundation of China (Grant No.

825 81972680 to X.L.), Beijing Natural Science Foundation (Grant No. 20201100463 to X.L.),

826 Tsinghua University-Peking University Jointed Center for Life Sciences (Grant No.

827 61020100119 to X.L.) and Alibaba innovation research programme.

Competing interests

829 The authors declare no competing interests.

Extended data

Extended Data Fig. 1 | Typical Bayesian network structure learning algorithm is sensitive to parameters. a-d, Networks learned by the typical Bayesian network structure learning strategy MI-hc given different correlation thresholds with the same microarray dataset. The dashed edges represent possible unconventional interlineage branches, while the red edges represent less credible branches.

CIBER   MI-hc

Extended Data Fig. 2 | CIBER-learned structures conform to prior knowledge and reduction landscapes better than the MI-hc algorithm. Plots show low-dimensional landscapes[42–44], CIBER and MI-hc results of (**a-c**) mouse scRNA-seq[42], (**d-f**) human scRNA-seq[43] and (**g-i**) human scATAC-seq[44] datasets respectively. The dashed edges represent unconventional interlineage edges that were reported to be possible in previous studies, while the red edges represent less credible edges. The results show that compared to the typical Bayesian network structure learning algorithm MI-hc, the CIBER-learned results were consistent with conventional knowledge with a larger number and a higher percentage of edges and captured the bifurcation structure more clearly.

Extended Data Fig. 3 | DDF identifies lineage driving features. The plots show the results
on RDF and DDF gene lists obtained with the human scRNA-seq dataset. **a**, Features are
allocated by log fold change and log p value obtained through RDF analysis; top RDF and
DDF features are coloured. **b**, The number of shared features between the two feature sets.
c, Literature search result shows that the DDF and RDF analyses identify different important
features, in which the red parts represent the reported features. **d**, Correlation between the
ranked lists. **e**, GSEA NES with terms related to haematopoiesis or haematopoietic
differentiation in the GO and KEGG databases. The terms for which DDF ranking list obtain
the higher NES are coloured in red. Dots in the left part represent the NES of RDF and DDF
ranking lists, while the distributions show the background NES given randomly ranked gene
lists.

Extended Data Fig. 4 | DDF results on the mouse microarray dataset. a, Features are allocated by log fold change and log p value obtained through RDF analysis; top RDF and DDF features are coloured. **b**, The number of shared top features between the two feature sets. **c**, Literature search result shows that the DDF and RDF analyses identify different important features, in which the red parts represent the reported features. **d**, Correlation between the ranked lists. **e**, GSEA NES with terms related to haematopoiesis or haematopoietic differentiation in the GO and KEGG databases. The terms for which DDF list obtain the higher NES are coloured in red. Dots in the left part represent the NES of RDF and DDF ranking lists, while the distributions show the background NES given randomly ranked gene lists. The RDF list performed slightly better than the DDF list in 12 out of 22 terms. However, the literature search still showed that DDF analysis identified some important genes which are not significantly expressed, demonstrating that DDFs can complement RDFs to better identify genes crucial to cell differentiation.

878 **Extended Data Fig. 5 | ROC curves of each TF-targeted gene set.**