A simple mechanism for integration of quorum sensing and cAMP signalling in *V. cholerae*

Lucas M. Walker[1], James R.J. Haycocks[1], Julia C. van Kessel[2], Triana N. Dalia[2], Ankur B. Dalia[2] and David C. Grainger[1*]

[1]School of Biosciences, University of Birmingham, Edgbaston B15 2TT, UK.

[2]Department of Biology, Indiana University, Bloomington, Indiana, USA.

*for correspondence

d.grainger@bham.ac.uk

Tel: +44 (0)121 414 5437

ABSTRACT

Many bacteria use quorum sensing to control changes in lifestyle. The process is regulated by microbially derived "autoinducer" signalling molecules, that accumulate in the local environment. Individual cells sense autoinducer abundance, to infer population density, and alter their behaviour accordingly. In *Vibrio cholerae*, quorum sensing signals are transduced by phosphorelay to the transcription factor LuxO. Unphosphorylated LuxO permits expression of HapR, which alters global gene expression patterns. In this work, we have mapped the genome-wide distribution of LuxO and HapR in *V. cholerae*. Whilst LuxO has a small regulon, HapR targets 32 loci. Many HapR targets coincide with sites for the cAMP receptor protein (CRP) that regulates the transcriptional response to carbon starvation. This overlap, also evident in other *Vibrio* species, results from similarities in the DNA sequence bound by each factor. At shared sites, HapR and CRP simultaneously contact the double helix and binding is stabilised by direct interaction of the two factors. Importantly, this involves a CRP surface that usually contacts RNA polymerase to stimulate transcription. As a result, HapR can block transcription activation by CRP. Thus, by interacting at shared sites, HapR and CRP integrate information from quorum sensing and cAMP signalling to control gene expression. This likely allows *V. cholerae* to regulate subsets of genes during the transition between aquatic environments and the human host.

INTRODUCTION

Vibrio cholerae is a Gram-negative bacterium responsible for the human disease cholera[1]. Estimates suggest 3 million annual infections, of which 100 thousand are fatal[2]. Most disease instances are attributed to the El Tor *V. cholerae* biotype, which is responsible for the ongoing 7th cholera pandemic[3]. Globally, over 1 billion people inhabit areas of endemicity and future climatic change is likely to exacerbate the risk of illness[2,4]. The success of *V. cholerae* as a pathogen is underpinned by an ability to colonise both aquatic ecosystems and the human intestinal tract[1]. In waterways, *V. cholerae* prospers by forming biofilms on arthropod exoskeletons. Degradation of these chitinous surfaces ultimately liberates *N*-acetylglucosamine (GlcNAc) for metabolism by the microbe[5]. Upon ingestion by a human host, *V. cholerae* express genetic determinants for acid tolerance, intestinal colonisation, and virulence. Diverse transcription factors regulate the transition and respond to signals including bile[6], temperature[7], nucleotide second messengers[8,9], and chitin availability[5]. Understanding these regulatory networks is important to determine how *V. cholerae* can switch between environments to cause disease outbreaks[3,10,11].

Quorum sensing is key for the transition of *V. cholerae* between ecological niches[12]. Briefly, *V. cholerae* produce at least 3 autoinducer (AI) signalling molecules: cholera AI-1 (CAI-1), AI-2, and 3,5-dimethylpyrazin-2-ol (DPO)[13]. In the environment, these compounds are detected by receptors in

neighbouring cells and indicate population density. Importantly, whilst AI-2 and DPO are produced

by multiple bacterial species, CAI-1 is only made by other members of the *Vibrio* genus[14]. Thus, *V.*

cholerae can determine the crude composition of bacterial populations. In the absence of their cognate

AIs, when population density is low, the receptors for CAI-I and AI-2 target the transcription factor

LuxO for phosphorylation via a phosphorelay system[13,15,16]. When phosphorylated, LuxO upregulates

the production of four small quorum regulatory RNAs (Qrrs)[17]. In turn, the Qrrs control expression of

two global transcription factors: AphA and HapR[17–19]. Importantly, whilst AphA production is

activated by Qrrs, synthesis of HapR is repressed. Hence, AphA and HapR control gene expression at

low and high cell density respectively[13,19]. A simplified outline of the LuxO dependent regulatory

pathway for HapR is illustrated in Figure 1a.

Identified as a regulator of *hapA*, required for *V. cholerae* migration through intestinal mucosa, HapR

is a TetR-family member that binds DNA as a homodimer via a N-terminal helix-turn-helix motif[20,21].

Many clinical isolates of pandemic *V. cholerae* have lost the ability to properly express HapR and this

may indicate adaptation to a more pathogenic lifestyle[3,10,22]. In *V. cholerae*, HapR regulates the

expression of ~100 genes to promote 'group behaviours' including natural competence, repression of

virulence genes, and escape from the host intestinal mucosa[23]. In other *Vibrio* spp., equivalent

regulons are larger. For example, LuxR in *Vibrio harveyi* regulates over 600 genes[24]. Expression of

HapR can be influenced by other factors. In particular, cAMP receptor protein (CRP), a regulator that

controls metabolism of alternative carbon sources, including chitin, upregulates HapR[25]. In this study,

we used chromatin immunoprecipitation and DNA sequencing (ChIP-seq) to identify direct DNA

binding targets of HapR and its upstream regulator, LuxO. We show that the degenerate DNA

consensus bound by HapR frequently overlaps targets for CRP. At such sites, HapR and CRP co-

operatively bind offset faces of the double helix. Strikingly, this occludes a key CRP surface required

to activate transcription. This simple mechanism allows *V. cholerae* species to integrate quorum

sensing, and cAMP signalling, in the control of gene expression.

RESULTS

Genome-wide DNA binding by HapR and LuxO in Vibrio cholerae

Whilst the impact of HapR on global gene expression in *V. cholerae* has been investigated, it is not

known which HapR responsive genes are directly controlled by the protein[23]. Similarly, the extent of

the direct LuxO regulon is unknown. Hence, we sought to map the binding of LuxO and HapR across

the *V. cholerae* genome. To facilitate this, *luxO* and *hapR* were cloned in plasmids pAMCF an

pAMNF respectively. The resulting constructs, encoding LuxO-3xFLAG or 3xFLAG-HapR, were

used to transform *V. cholerae* strain E7946. In subsequent ChIP-seq experiments, anti-FLAG

antibodies were used to select fragments of the *V. cholerae* genome bound with either LuxO or HapR.

The derived binding profiles are shown in Figure 1b. In each plot, genes are shown as blue lines

(outer two tracks) whilst the LuxO and HapR binding signals are red and green respectively (inner

two tracks). Examples of individual binding peaks for each factor are shown in Figure 1c. In total, we

identified 5 and 32 peaks for LuxO and HapR binding respectively (Table 1). Previous work

identified targets for LuxO adjacent to genes encoding the 4 Qrr sRNAs. We recovered all of these

known LuxO targets, and an additional binding site was identified between *VC1142* and *VC1143*.

These divergent genes encode cold shock-like protein CspD, and the Clp protease adaptor protein,

ClpS, respectively. Note that the LuxO binding signal at this locus is small, compared to the *qrr*1-4

targets, but may still be involved in transcription regulation (Figure 1-figure supplement 1). To

identify the sequence bound by LuxO, DNA regions overlapping LuxO binding peaks were inspected

using MEME. The motif identified matches the known consensus for LuxO binding and was found at

all LuxO targets (Table 1 and Figure 1d)[26]. The positions of LuxO binding sites with respect to genes,

and the functions encoded by these genes, are summarised in Figures 1e and 1f respectively.

Of the 32 peaks for HapR binding, 4 correspond to previously identified direct targets (*hapR*[27],

VC0241[28], *VC1851*[29] and *VCA0148*[28]). However, some known targets had weak or poorly

reproducible binding signals (Figure 1- figure supplement 2). A DNA motif common to all 32 HapR

ChIP-seq peaks matched prior descriptions of the DNA target for HapR or closely related

proteins[27,30,31] (Figure 1d). Occurrences of this HapR binging motif were most frequent in the 200 bp

preceding a gene start codon (Figure 1e). Most often, the genes adjacent to HapR binding peaks

encode protein functions related to metabolism, motility, and chemotaxis (Figure 1f). Overall, our

data suggest that LuxO primarily regulates gene expression via the 4 Qrr sRNA molecules.

Conversely, the genome-wide distribution of HapR is consistent with that of a global gene regulator

with many undefined regulatory roles.

HapR is a direct regulator of transcription at many target sites

We focused our attention on new HapR target promoters where adjacent coding sequence could be

used to predict encoded protein function. For these 24 targets, regulatory DNA was cloned upstream

of *lacZ* in plasmid pRW50T. Recombinants were then transferred to *V. cholerae* E7946, or the Δ*hapR*

derivative, by conjugation. Strains generated were cultured overnight before β-galactosidase activities

were determined. The results are shown in Figure 2a. Promoters were categorised as inactive,

unresponsive, repressed or activated by HapR. We identified 2 and 7 promoters subject to activation

and repression by HapR respectively. Of the remaining promoters, 6 were inactive and 9 unresponsive

to HapR in our conditions. Next, the 9 promoter DNA fragments responsive to HapR *in vivo* were

cloned upstream of the λ*oop* terminator in plasmid pSR. The resulting constructs were then provided

to housekeeping *V. cholerae* RNA polymerase, as templates for *in vitro* transcription, in the presence

and absence of HapR. The results are shown in Figure 2b where the expected size of transcripts

terminated by λ*oop* are marked with blue triangles[32]. Recall that the *VC1375* and *VC1403* promoters

were activated by HapR *in vivo* (Figure 2a). Consistent with this, HapR also activated the *VC1375*

promoter *in vitro* (Figure 3b, lanes 43-47). However, HapR did not activate *in vitro* transcription from

the *VC1403* promoter (Figure 3b, lanes 48-53). Indeed, interpretation of these data were hampered

because the location of the *VC1403* transcription start site (TSS) is not known[32]. Of the 7 promoters

repressed by HapR *in vivo*, we observed repression in 6 cases *in vitro* (*hapR*, *VC0585*, *VC2352*,

VCA0219, *VCA0663* and *VCA0960*) (lanes 7-42). Conversely, the *murQP* promoter (P*murQP*) subject

to repression by HapR *in vivo*, generated no transcript *in vitro* (lanes 1-6). Full gel images are shown

in Figure 2-figure supplement 1.

Transcription from the murQP *promoter requires CRP* in vivo *and* in vitro

The *murQP* operon (*VC0206-VC0207*) encodes functions important for recycling of peptidoglycan[33].

Briefly, cell wall derived *N*-acetylmuramic acid (MurNAc) is transported across the inner membrane,

and simultaneously phosphorylated, by the phosphotransferase system dependent permease MurP.

Resulting MurNAc-6P is hydrolysed by MurQ to generate *N*-acetylglucosamine 6-phosphate

(GlcNAc-6P). Intriguingly, GlcNAc-6P can also be derived from chitin break down and this coincides

with expression of HapR. Hence, we focused on understanding the role of HapR bound upstream of

murPQ. The HapR ChIP-seq binding signal at the *murQP* locus is shown in Figure 3a and the

associated regulatory region is shown in Figure 3b. The centre of the ChIP-seq peak for HapR is

marked by an asterisk and the predicted binding site is highlighted green. We reasoned that our

inability to detect transcription from P*murQP in vitro* was likely because an undefined transcriptional

activator is absent (Figure 2b). Inspection of the DNA sequence upstream of *murQP* identified a close

match to the consensus binding site for CRP (5'-TGTGA-N_6-TCACA-3'). Furthermore, this sequence

was located 41.5 bp upstream, of the *murQP* TSS (Figure 3b). This is a common scenario for CRP

dependent transcription activation[34]. To measure binding of CRP to the *murQP* regulatory region we

used electrophoretic mobility shift assays (EMSAs). Consistent with our prediction, CRP bound to

the *murQP* regulatory DNA (Figure 3c, lanes 1 and 2). To confirm that we had correctly identified the

binding site for CRP we made a series of P*murQP* derivatives. The Δ183 and Δ211 DNA fragments

have large upstream deletions (sites of truncation are shown by inverted triangles in Figure 3b, which

mark the 5' end of the remaining promoter DNA) but still bind CRP (Figure 3c, lanes 3-6).

Conversely, point mutations -35g and -49g, within the CRP site, prevent binding (Figure 3c, lanes 7-

8). To determine the impact of CRP on P*murQP* activity we first used *in vitro* transcription assays

(Figure 3d). Addition of CRP to reactions resulted in production of an RNA from P*murQP*. We

observed similar CRP dependence *in vivo* using β-galactosidase assays (Figure 3e, compare wild type

promoter activity with and without CRP). Furthermore, in wild type cells, the -35g and -49g

mutations reduced promotor activity whilst the Δ183 and Δ211 truncations did not (Figure 3e). We

note that the Δ211 derivative is much more active than the starting promoter DNA sequence, but
transcription remains totally dependent on CRP. Most likely, the truncation removes a repressive
DNA element upstream of the core promoter.

HapR and CRP bind a shared DNA site at the murQP *promoter*

At P*murQP*, the DNA site for CRP is completely embedded within the predicted HapR binding
sequence (Figure 3b). To better understand this unusual configuration, we used DNAseI footprinting.
The results are shown in Figure 4a. Lane 1 shows the pattern of DNAseI digestion in the absence of
bound protein. In the presence of CRP (lanes 2-4) a footprint was observed between positions -29 and
-59 bp relative to the *murQP* TSS. As is usual for CRP, and a consequence of DNA bending, the
footprint comprised protection from, and hypersensitivity to, DNAse I attack. Three distinct sites of
DNAseI hypersensitivity are marked by orange arrows alongside lane 4 in Figure 4a. The pattern of
DNAse I digestion in the presence of HapR is shown in lanes 5-8. The footprint due to HapR binding
exactly overlaps the region bound by CRP and results in complete protection of the DNA from
digestion between positions -29 and -58 (green bar adjacent to lane 8). We also observed changes in
the relative intensity of bands upstream of the HapR site between promoter positions -60 and -80. We
speculate that this may result from changes in DNA conformation. Importantly, there was one further
subtle difference between HapR and CRP induced banding patterns. Namely, in the presence of
HapR, a band was observed at position -58 (see green triangle adjacent to lane 8). With CRP, a band
was instead observed at position -59 (compare lanes 2-4 with 5-8). In a final set of assays, we
examined addition of CRP and HapR in unison. We reasoned that 3 outcomes were possible. First,
one of the two protein factors could outcompete the other. This should result in a DNAse I digestion
pattern identical to either the individual CRP or HapR footprint. Second, some DNA fragments in the
reaction could be bound by CRP and others by HapR. In this case, a mixed DNAse I digestion pattern,
containing all features of the individual footprints due to CRP and HapR, should occur. Third, CRP
and HapR could bind simultaneously. This might generate a DNAse I digestion pattern with
similarities to the CRP and HapR footprints. However, accessibility of the nucleic acid to DNAse I
would likely be altered in some way, with unpredictable outcomes. The result of the experiment was
analysed in lanes 9-12. The binding pattern matched only some aspects of the individual footprints for
CRP and HapR. Hence, we observed 2 of the 3 DNAse I hypersensitivity sites due to CRP binding.
Changes in the banding pattern upstream of the binding sequence, due to HapR, were also detected.
We did not observe the band at position -58 detected with HapR alone. Rather, we observed a band at
position -59. An additional band at position -26 (black triangle adjacent to lane 12) was unique to
these reactions. We conclude that HapR and CRP recognise the same section of the *murQP* regulatory
region and may bind in unison.

HapR and CRP bind the murQP *promoter co-operatively*

Fragments of the *murQP* regulatory DNA, simultaneously bound by CRP and HapR, are expected to
have distinct migratory properties during electrophoresis. Thus, we compared binding of CRP and/or
HapR using EMSAs. The results are shown in Figure 4b. As expected, addition of CRP to reactions
caused a distinct shift in electrophoretic mobility (lanes 1-5). Comparatively, at the concentration
used, HapR bound the DNA fragment poorly; we observed only smearing of the free DNA at the
highest HapR concentration tested (lanes 6-10). The binding pattern due to HapR was dramatically
different if DNA was pre-bound with CRP (lanes 11-15). In this scenario, even low concentrations of
added HapR were sufficient to generate a super-shifted nucleoprotein complex (lanes 11-15). These
data are consistent with HapR having a higher affinity for CRP-P*murQP* than P*murQP* alone. Hence,
HapR and CRP bind the *murQP* regulatory region co-operatively. A mundane explanation is that
increased molecular crowding, upon CRP addition, increases the effective concentration of HapR. To
exclude this possibility, we did two further sets of EMSA experiments. In the first set of assays, CRP
was added at a lower concentration. Thus, some DNA remained unbound (Figure 4c, lanes 1 and 2).
Hence, when added to such reactions, HapR could bind either the free DNA or the CRP-DNA
complex. Consistent with HapR preferentially binding the latter, all of the CRP-DNA complex was
super shifted upon HapR addition. Conversely, the free DNA remained unbound (compare lanes 2 and
4). In equivalent experiments, with point mutations -49g and -35g in the CRP site, neither CRP or
HapR were able to bind the DNA (lanes 5-8). In a second set of tests, we used the *hapR* regulatory
DNA that binds HapR but not CRP. If CRP addition increased the effective concentration of HapR,
this should result in much tighter HapR binding to the *hapR* promoter. However, this was not the case
(Figure 4-figure supplement 1). Taken together, our data are consistent with CRP and HapR co-
operatively binding the same DNA locus at the *murQP* promoter region.

HapR represses CRP dependent transcription from the murQP *promoter in vivo* and *in vitro*

Recall that, in the absence of CRP, P*murQP* is inactive *in vitro* (Figures 2b and 3d). Furthermore, the
promoter is subject to repression by HapR *in vivo* (Figure 2a). An explanation consistent with both
observations is that HapR directly counteracts CRP mediated activation. To test this, we used *in vitro*
transcription assays (Figure 4d). As expected, addition of CRP activated *murQP* transcription (lanes
1-4) and this was blocked by addition of HapR (lanes 5-8). We also repeated our prior *lacZ* fusion
experiments, using the Δ211 P*murQP* derivative, and *V. cholerae* E7946 lacking *crp* and/or *hapR*.
The result is shown in Figure 4e. Deletion of *hapR* caused increased transcription from P*murQP* only
when CRP was present. Hence, HapR also represses CRP dependent *murQP* transcription *in vivo*.

Binding sites for CRP and HapR overlap in a specific configuration genome-wide

Both CRP and HapR bind the same DNA region upstream of *murPQ*. This suggests similar nucleic
acid sequences are recognised by each factor. Figure 5a shows an alignment of DNA logos, derived
from CRP[9] (top) and HapR (bottom) ChIP-seq targets. The two motifs have features in common that

align best when the logo centres are offset by 1 base pair. This is consistent with the arrangement of

binding sites upstream of *murPQ* (Figure 3b). To understand the importance of this configuration we

first took a bioinformatic approach. The DNA sequences logos shown in Figure 5a were used to create

position weight matrices (PWMs) describing either the CRP or HapR binding site. We then searched

the *V. cholerae* genome, using each PWM, and calculated the distance between identified CRP and

HapR sites. The data for all sites within 100 bp of each other is shown in Figure 5b (top panel). In all

cases, the CRP and HapR targets were offset by 1 bp. We then repeated the analysis after randomising

the *V. cholerae* genome sequence (bottom panel). The number of overlapping targets was reduced

7-fold. An equivalent analysis of the *V. harveyi* genome produced similar results (Figure 5-figure

supplement1). Hence, sites for CRP and HapR have a propensity to coincide in a specific

configuration. That such sites are found more frequently in native genome sequences, compared to

those first randomised, suggests selection during genome evolution. We next sought to understand

how this arrangement might permit simultaneous and co-operative binding of CRP and HapR.

A structural model of the DNA-CRP-HapR ternary complex

To understand organisation of the DNA-CRP-HapR ternary complex we used structural modelling.

The *V. cholerae* CRP protein is 96 % identical to the equivalent factor in *Escherichia coli*. Similarly,

the *Staphylococcus aureus* factor QacR is 50 % similar to *V. cholerae* HapR. Previously, structural

biology tools were used to investigate *E. coli* CRP, and *S. aureus* QacR, bound with their cognate

DNA targets. We used this information to build a model for the DNA-CRP-HapR ternary complex.

Importantly, we ensured that the CRP and HapR binding centres were offset by 1 bp. When aligned in

this way, CRP and HapR recognise the same section of DNA via different surfaces of the double

helix. We examined the model in the context of our DNAse I footprinting data. Recall that CRP

binding upstream of *murPQ* induces three sites of DNAse I hypersensitivity (Figure 4a). These

correspond to positions -47, -38 and -34 with respect to the *murQP* TSS. Figure 5-figure supplement 2

shows these positions highlighted in the context of our model. In the presence of CRP alone, all sites

are surface exposed but position -34 is partially occluded by CRP (Figure 5-figure supplement 2a).

This likely explains why positions -47 and -38 are more readily cleaved by DNAse I (Figure 4a). With

both CRP and HapR, position -34 was completely protected from DNAse I attack (Figure 4a).

Consistent with the footprinting data, our model indicates that position -34 is almost completely

hidden upon binding of HapR (Figure 5-figure supplement 2b). Conversely, access to positions -47

and -38 is not altered (Compare Figure 5-figure supplement 2a and 2b).

Co-operative binding with HapR requires CRP residue E55

Co-operative DNA binding by transcription factors can result from their direct interaction[35–37]. In our

model, a negatively charged surface of CRP (including residue E55) is in close proximity to positively

charged HapR residue R123 (Figure 5c). In initial experiments, we mutated both protein surfaces to

remove the charged side chain, or replace the residue with an oppositely charged amino acid. We then
investigated consequences for HapR and CRP binding individually at P*murQP* using EMSAs (Figure
5-figure supplement 3). Whilst the CRP derivatives were able to bind the *murQP* regulatory region
normally, HapR variants were completely defective. This is likely because R123 sits at the HapR
dimerisation interface. Hence, we focused on understanding the contribution of CRP sidechain E55 to
co-operative DNA binding by HapR and CRP using EMSAs. The results are shown in Figure 5d.
Both wild type CRP, and CRP[E55A], were able to bind the *murQP* regulatory region similarly (lanes 1-4
and 10-13). As expected, HapR bound tightly to the wild type CRP:DNA complex (lanes 5-9).
Conversely, HapR had a lower affinity for DNA in complex with CRP[E55A] (lanes 14-18). This
suggests that the E55A mutation in CRP destabilises the interaction with HapR.
300
*Repression of P*murQP *by HapR requires CRP residue E55*
Residue E55 locates to a negatively charged surface of CRP called Activating Region 3 (AR3). This
determinant aids recruitment of RNA polymerase when CRP binds close to the promoter -35
element[38]. Hence, AR3 is likely to be important for activation of P*murQP* (Figure 3b). We inferred
that CRP lacking E55 should activate P*murQP* less efficiently but be less sensitive to negative effects
of HapR. To test these predictions, we used *in vitro* transcription assays. The results for CRP,
CRP[E55A] and CRP[E55R] are shown in Figure 5e. All CRP derivatives were able to activate transcription
from P*murQP*. However, consistent with an important role for AR3, the ability of the CRP[E55A] and
CRP[E55R] to activate transcription was impaired (compare lanes 1-5, 10-14 and 19-23). Crucially,
whilst HapR reduced transcription dependent on wild type CRP by 50-fold (compare lane 4 with lanes
6-9) only a 2-fold effect of HapR was observed with CRP[E55A] (compare lane 13 with lanes 15-18). In
the presence of CRP[E55R], HapR was even less effective (compare lane 22 with lanes 24-27).
313
High cell density locked V. cholerae are defective for growth on MurNAc
Phosphorylated LuxO activates expression of the Qrr sRNAs that inhibit *hapR* expression at low cell
density (Figure 1a). Consequently, deletion of *luxO* causes constitutive expression of HapR. Thus,
Δ*luxO V. cholerae* are "locked" in a high cell density state[29]. Our model predicts that such strains will
be defective for growth using MurNAc as the sole carbon source, as this requires expression of
murQP that is repressed by HapR. Furthermore, any such defect should be relieved upon deletion of
hapR. To test this, we constructed strains lacking different combinations of *luxO* and *hapR*. We also
tested a *V. cholerae* derivative lacking *murP*[39]. Figure 6a illustrates growth in M9 minimal media,
supplemented with MurNAc or glucose, and in LB-Miller medium. As expected, cells lacking *murP*
could not grow when MurNAc was the only carbon source but were not defective in other conditions
(compare grey data points in each panel). Cells lacking *hapR*, alone or in combination with *luxO*, had
a similar growth defect in all conditions. Strikingly, the *luxO* mutant (high cell density locked),
exhibited a growth defect only when MurNAc was the sole carbon source (compare red data points).

Specifically, these cells exhibited an extended lag phase in MurNAc. This extended lag phase was not

apparent when both *luxO* and *hapR* were deleted, consistent with the effect of *luxO* being mediated by

HapR-dependent repression of *murQP*.

Co-operative interactions between HapR and CRP are commonplace

In a final set of experiments, we turned our attention to other sites shared by CRP (prior work[9]) and

HapR (Table 1). We selected 5 such targets and examined binding of CRP and HapR using EMSAs.

At 1 target, adjacent to *VCA0218*, binding was not co-operative and free DNA remained when both

proteins were present (Figure 5-figure supplement 4). For 4 of the targets, we detected co-operative

binding of CRP and HapR, reminiscent of our experiments with P*murQP* DNA (compare Figure 5-

figure supplement 4 and Figure 4). At these loci (adjacent to *VC0102*, *VC1851*, *VCA0663* and

VCA0691) either HapR or CRP bound poorly to DNA in the absence of the other protein. However,

when both factors were added together, all DNA shifted into a distinct low mobility complex. We

conclude that co-operative binding of HapR and CRP to shared targets is common.

DISCUSSION

Previously, two studies have mapped DNA binding by HapR homologs in *Vibrio* species. For *V.*

harveyi, van Kessel and co-workers used ChIP-seq to identify 105 LuxR binding targets[30]. At 77 of

these sites, LuxR repressed transcription. Using ChIP-seq and global DNAse I footprinting, Zhang *et*

al. found 76 LuxR bound regions in *Vibrio alginolyticus*[31]. Regulatory effects were evident for 37

targeted genes, with 22 cases of LuxR mediated repression. In the present study, we identified 32

HapR bound sections of the *V. cholerae* genome. Consistent with prior work, repression of target

genes was the most common regulatory outcome. Furthermore, the DNA binding consensus derived

here for HapR is almost identical to motifs for LuxR binding in *V. harveyi* and *V. alginolyticus*.

Contrastingly, Tsou and colleagues used bioinformatic tools to predict HapR binding in *V. cholerae*[28].

Two different HapR binding motifs were proposed. Both partially match the HapR target sequence

proposed here. Most likely, the analysis of Tsou *et al.* was hampered by a paucity of targets from

which a full consensus could be derived. We note that our list of 32 HapR targets does not include all

known targets. However, on inspection, whilst insufficient to pass our stringent selection criteria,

weaker signals for HapR are evident at many such locations (Figure 1-figure supplement 2 and

Supplementary file 1). In particular, we note evidence for binding of HapR upstream of *hapA*, which

has previously been only inferred (Figure 1-figure supplement 2b). We note that previous relied on

computational predictions and *in vitro* DNA binding assays to identify potential HapR targets. That

not all such targets are bound *in vivo*, in the single growth condition tested here, is to be expected.

Recognition of shared DNA targets provides a simple mechanism for integration of quorum sensing

signals, relayed by HapR, and cAMP fluctuations, communicated by CRP. In the example presented

here, HapR acts to prevent transcription activation by co-binding the same DNA target with CRP

(Figure 4). Hence, at P*murQP*, the function of CRP switches from that of an activator to a co-

repressor with HapR (Figure 6b). This regulatory strategy is a logical consequence of *V. cholerae*

forming biofilms on chitinous surfaces. At low cell density, rapidly dividing cells must continually

remodel their cell wall. In these conditions, HapR is not expressed. Thus, MurQ and MurP are

produced and can convert cell wall derived MurNAc to GlcNAc-6P. Conversely, in high cell density

scenarios, usually involving adherence to chitin, cells divide infrequently, and remodelling of the cell

wall is reduced. In addition, GlcNAc-6P can be derived readily from chitin oligosaccharides. Hence,

cells locked in the high cell density state are defective for growth when supplied with MurNAc as the

sole carbon source (Figure 6a). We suggest that HapR and CRP are likely to coordinate the expression

of other metabolic enzymes in a similar way. Interestingly, AphA, another quorum sensing responsive

regulator, also acts alongside CRP at many *V. cholerae* promoters[40]. Indeed, AphA and CRP binding

sites can overlap but this results in competition between the factors[40]. Together with results presented

here, these observations highlight close integration of quorum sensing with gene control by cAMP in

V. cholerae.

MATERIALS AND METHODS

Strains, plasmids and oligonucleotides

Strains, plasmids and oligonucleotides used in this study are listed in Supplementary file 2. All *V.*

cholerae strains are derivatives of E7946[41]. Chromosomal deletions were made using the pKAS32

suicide plasmid for allelic exchange[42,43] or via splicing-by-overlap-extension PCR and chitin-induced

natural transformation[44]. The *E. coli* strain JCB387 was used for routine cloning[45]. Plasmids were

transferred into *V. cholerae* by either conjugation or transformation as described previously[9,40].

ChIP-seq and bioinformatics

Chromatin immunoprecipitation was done as in prior work[40] using strain E7946, carrying

plasmid pAMCF-*luxO* or pAMNF-*hapR*, and anti-FLAG antibodies . In both cases, control

experiments were done using the equivalent plasmid with no gene insert. Note that both

plasmids drive low level constitutive expression of 3xFLAG transcription factor

derivatives[46]. Lysates were prepared from LB-Lennox medium cultures, incubated with

shaking at 37 □ to an OD_{650} of ~1.1. Following sonication, the protein-DNA complexes were

immunoprecipitated with an anti-FLAG antibody (Sigma) and Protein A sepharose beads.

Immunoprecipitated DNA was blunt-ended, A- tailed, and ligated to barcoded adaptors

before elution and de-crosslinking. ChIP-seq libraries were then amplified by PCR and

purified. Library quality was assessed using an Agilent Tapestation 4200 instrument and

quantity determined by qPCR using an NEBnext library quantification kit (NEB). Libraries

were sequenced as described previously[46] and reads are available from ArrayExpress using accession code E-MTAB-11906. Single-end reads, from two independent ChIP-seq experiments for each strain, were mapped to the reference *V. cholerae* N16961 genome (chromosome I: NC_002505.1 and chromosome II: NC_002506.1) with Bowtie 2[47]. The read depth at each position of the genome was determined for each BAM file using multibamsummary. Each binding profile was then normalised to an average genome-wide read depth of 1 read per base. Following normalisation, the average read depth per base for each pair of replicates was calculated. The resulting files were used to generate the circular plots in Figure 1 using DNAplotter[48]. For peak selection, the files were viewed as graphs using the Artemis genome browser[49]. After visually identifying an appropriate cut-off, peaks were selected using the "create features from graph" tool. Note that our cut-off was selected to identify only completely unambiguous binding peaks. Hence, weak or less reproducible binding signals, even if representing known targets, were excluded (see Discussion for further details). For HapR, the window size, minimum feature size, and cut-off value were 100, 100 and 10 respectively. For LuxO, the equivalent values were 100, 100 and 4. The mid-point of features selected in this way was set as the peak centre. In each case, 300 bp of sequence from the peak centre was selected and the combined set of such sequences for each factor were analysed using MEME to generate DNA sequence logos[50].

β-galactosidase assays

Promoter DNA was fused to *lacZ* in plasmid pRW50T that can be transferred from *E. coli* to *V. cholerae* by conjugation[9]. Assays of β-galactosidase activity were done according to the Miller method[51]. Bacterial cultures were grown at 37 □ with shaking in LB-Lennox medium, supplemented with appropriate antibiotics, to an OD_{650} of ~1.1. Values shown are the mean of three independent experiments and error bars show the standard deviation.

Proteins

We purified *V. cholerae* CRP and RNA polymerase as described previously[9,40]. To generate HapR, *E. coli* T7 Express cells were transformed with plasmid pHis-tev-HapR, or derivatives, which encodes HapR with a His_6 tag and intervening site for the tobacco etch virus protease protease. Transformants were cultured in 40 ml LB-Lennox medium overnight, then sub-cultured in 1 L of fresh broth, with shaking at 37 ºC. When sub-cultures reached mid-log phase they were supplemented with 400 mM IPTG for 3 hours. Cells were then collected by centrifugation, resuspended in 40 ml of buffer 1 (40 ml 25 mM Tris-HCl pH 7.5, 1 mM EDTA and 1 M NaCl) and lysed by sonication. Inclusion bodies, recovered by centrifugation, were resuspended with 40 ml of buffer 2 (25 mM Tris-HCl pH 8.5 and 4

M urea) before the remaining solid material was again recovered and then solubilised using 40 ml of

buffer 3 (25 mM Tris-HCl pH 8.5 and 6 M guanidine hydrochloride). Cleared supernatant was applied

to a HisTrap HP column (GE healthcare) equilibrated with buffer A (25 mM Tris-HCl pH 8.5 and 1 M

NaCl). To elute His_6-HapR, a gradient of buffer B (25 mM Tris-HCl pH 8.5, 1 M NaCl and 1 M

imidazole) was used. Fractions containing His_6-HapR were pooled and the protein was transferred

into buffer X (50 mM HEPES, 1 M NaCl, 1 mM DTT, 5 mM EDTA and 0.1 mM Triton X-100) by

dialysis. Finally, we used Vivaspin ultrafiltration columns to reduce sample volume. The

concentration of His_6-HapR was then determined.

in vitro *transcription assays*

Experiments were done using our prior approach[40]. Plasmid templates were isolated from *E. coli* using

Qiagen Maxiprep kits. Each *in vitro* transcription assay contained 16 μg/ml DNA template in 40 mM

Tris pH 7.9, 5 mM MgCl2, 500 μM DTT, 50 mM KCl, 100 μg/ml BSA, 200 μM ATP/GTP/CTP, 10

μM UTP and 5 μCi α-P32-UTP. Purified HapR and CRP were added at the indicated concentrations

prior to the reaction start point. In experiments where CRP was used, the protein was incubated with

cAMP 37 ℃ prior to addition. Transcription was instigated by addition of RNA polymerase

holoenzyme prepared in advance by incubation of the core enzyme with a 4-fold excess of σ^{70} for 15

minutes at room temperature. After 10 minutes incubation at 37 ▢, reactions were stopped by the

addition of an equal volume of formamide containing stop buffer. Reactions were resolved on an 8%

(*w/v*) denaturing polyacrylamide gel, exposed on a Bio-Rad phosphor screen then visualised on a Bio-

Rad Personal Molecular Imager. The quantify transcript levels, we measured the intensity of bands

corresponding to RNAI and the RNA of interest using Quantity One software. After subtracting

background lane intensity, we calculated the RNA of interest to RNAI ratio. The maximum ratio was

set to 100 % activity with other ratios shown a percentage of this maximum. Experiments were

repeated at least twice with similar results.

Electrophoretic mobility shift assays and DNAse I footprinting

Promoter DNA fragments were excised from plasmid pSR and end-labelled with γ32-ATP using T4

PNK (NEB). EMSAs and DNase I footprints were done as previously described [40]. Full gel images are

shown in Figure 2-figure supplement 1. Experiments were repeated at least twice with similar results.

Structural modelling

The model of the ternary DNA-CRP-HapR complex was generated in PyMOL by aligning PDB

depositions 1jt0 (QacR-DNA complex) and 6pb6 (CRP-DNA complex). Alignments were done

manually and guided by the relative two-fold centres of symmetry for each complex. Each structure

was positioned so that their DNA base pairs overlapped and binding centres were offset by 1 base

pair. The Mutagenesis function of PyMOL was used to replace QacR sidechain K107, equivalent to

HapR R123[20], with an arginine residue. The double helix of the QacR DNA complex is hidden in the

final model.

REFERENCES

1. Nelson, E. J., Harris, J. B., Morris, J. G., Calderwood, S. B. & Camilli, A. Cholera

transmission: The host, pathogen and bacteriophage dynamic. *Nature Reviews Microbiology*

vol. 7 693–702 (2009).

2. Ali, M., Nelson, A. R., Lopez, A. L. & Sack, D. A. Updated Global Burden of Cholera in

Endemic Countries. *PLoS Negl. Trop. Dis.* **9**, (2015).

3. Domman, D. *et al.* Integrated view of Vibrio cholerae in the Americas. *Science* **358**, 789–793

(2017).

4. Asadgol, Z., Mohammadi, H., Kermani, M., Badirzadeh, A. & Gholami, M. The effect of

climate change on cholera disease: The road ahead using artificial neural network. *PLoS One*

14, e0224813 (2019).

5. Meiborn, K. L. *et al.* The Vibrio cholerae chitin utilization program. *PNAS* **101**, 2524–2529

(2004).

6. Hung, D. T. & Mekalanos, J. J. Bile acids induce cholera toxin expression in Vibrio cholerae

in a ToxT-independent manner. *PNAS* **102**, 3028–3033 (2005).

7. Weber, G. G., Kortmann, J., Narberhaus, F. & Klose, K. E. RNA thermometer controls

temperature-dependent virulence factor expression in Vibrio cholerae. *PNAS* **111**, 14241–

14246 (2014).

8. Krasteva, P. V. *et al.* Vibrio cholerae vpst regulates matrix production and motility by directly

sensing cyclic di-GMP. *Science* **327**, 866–868 (2010).

9. Manneh-Roussel, J. *et al.* Camp receptor protein controls Vibrio cholerae gene expression in

response to host colonization. *MBio* **9**, (2018).

10. Kamareddine, L. *et al.* Activation of Vibrio cholerae quorum sensing promotes survival of an

arthropod host. *Nat. Microbiol.* **3**, 243–252 (2018).

11. Weill, F.-X. *et al.* Genomic history of the seventh pandemic of cholera in Africa. *Science* **358**,

785–789 (2017).

12. Eickhoff, M. J. & Bassler, B. L. SnapShot: Bacterial Quorum Sensing. *Cell* vol. 174 1328-

1328.e1 (2018).

13. Mukherjee, S. & Bassler, B. L. Bacterial quorum sensing in complex and dynamically

changing environments. *Nature Reviews Microbiology* vol. 17 371–382 (2019).

14. Henke, J. M. & Bassler, B. L. Three parallel quorum-sensing systems regulate gene expression

in Vibrio harveyi. *J. Bacteriol.* **186**, 6902–6914 (2004).

15. Freeman, J. A. & Bassler, B. L. A genetic analysis of the function of LuxO, a two-component

response regulator involved in quorum sensing in Vibrio harveyi. *Mol. Microbiol.* **31**, 665–677

(1999).

16.   Freeman, J. A. & Bassler, B. L. Sequence and function of LuxU: A two-component

phosphorelay protein that regulates quorum sensing in Vibrio harveyi. *J. Bacteriol.* **181**, 899–

906 (1999).

17.   Lenz, D. H. *et al.* The small RNA chaperone Hfq and multiple small RNAs control quorum

sensing in Vibrio harveyi and Vibrio cholerae. *Cell* **118**, 69–82 (2004).

18.   Shao, Y. & Bassler, B. L. Quorum-sensing non-coding small RNAs use unique pairing regions

to differentially control mRNA targets. *Mol. Microbiol.* **83**, 599–611 (2012).

19.   Rutherford, S. T., Van Kessel, J. C., Shao, Y. & Bassler, B. L. AphA and LuxR/HapR

reciprocally control quorum sensing in vibrios. *Genes Dev.* **25**, 397–408 (2011).

20.   De Silva, R. S. *et al.* Crystal structure of the Vibrio cholerae quorum-sensing regulatory

protein HapR. *J. Bacteriol.* **189**, 5683–5691 (2007).

21.   Jobling, M. G. & Holmes, R. K. Characterization of hapR , a positive regulator of the Vibrio

cholerae HA/protease gene hap , and its identification as a functional homologue of the Vibrio

harveyi luxR gene. *Mol. Microbiol.* **26**, 1023–1034 (1997).

22.   Heidelberg, J. F. *et al.* DNA sequence of both chromosomes of the cholera pathogen Vibrio

cholerae. *Nature* **406**, 477–483 (2000).

23.   Nielsen, A. T. *et al.* RpoS controls the Vibrio cholerae mucosal escape response. *PLoS Pathog.*

2, 0933–0948 (2006).

24.   Van Kessel, J. C., Rutherford, S. T., Shao, Y., Utria, A. F. & Bassler, B. L. Individual and

combined roles of the master regulators apha and luxr in control of the Vibrio harveyi quorum-

sensing regulon. *J. Bacteriol.* **195**, 436–443 (2013).

25.   Silva, A. J. & Benitez, J. A. Transcriptional regulation of Vibrio cholerae

hemagglutinin/protease by the cyclic AMP receptor protein and RpoS. *J. Bacteriol.* **186**, 6374–

6382 (2004).

26.   Tu, K. C. & Bassler, B. L. Multiple small RNAs act additively to integrate sensory information

and control quorum sensing in Vibrio harveyi. *Genes Dev.* **21**, 221–233 (2007).

27.   Lin, W., Kovacikova, G. & Skorupski, K. Requirements for Vibrio cholerae HapR binding and

transcriptional repression at the hapR promoter are distinct from those at the aphA promoter. *J.*

Bacteriol. **187**, 3013–3019 (2005).

28.   Tsou, A. M., Cai, T., Liu, Z., Zhu, J. & Kulkarni, R. V. Regulatory targets of quorum sensing

in Vibrio cholerae: Evidence for two distinct HapR-binding motifs. *Nucleic Acids Res.* **37**,

2747–2756 (2009).

29.   Waters, C. M., Lu, W., Rabinowitz, J. D. & Bassler, B. L. Quorum sensing controls biofilm

formation in Vibrio cholerae through modulation of cyclic Di-GMP levels and repression of

vpsT. *J. Bacteriol.* **190**, 2527–2536 (2008).

30.   van Kessel, J. C., Ulrich, L. E., Zhulin, I. B. & Bassler, B. L. Analysis of activator and

repressor functions reveals the requirements for transcriptional control by LuxR, the master

regulator of quorum sensing in Vibrio harveyi. *MBio* **4**, (2013).

31. Zhang, J. *et al.* Binding site profiles and N-terminal minor groove interactions of the master

quorum-sensing regulator LuxR enable flexible control of gene activation and repression.

Nucleic Acids Res. **49**, 3274–3293 (2021).

32. Papenfort, K., Förstner, K. U., Cong, J. P., Sharma, C. M. & Bassler, B. L. Differential RNA-

seq of Vibrio cholerae identifies the VqmR small RNA as a regulator of biofilm formation.

PNAS **112**, E766–E775 (2015).

33. Borisova, M. *et al.* Peptidoglycan recycling in gram-positive bacteria is crucial for survival in

stationary phase. *MBio* **7**, (2016).

34. Savery, N. J. *et al.* Transcription activation at class II CRP-dependent promoters: Identification

of determinants in the C-terminal domain of the RNA polymerase α subunit. *EMBO J.* **17**,

3439–3447 (1998).

35. Wade, J. T., Belyaeva, T. A., Hyde, E. I. & Busby, S. J. W. A simple mechanism for co-

dependence on two activators at an Escherichia coli promoter. *EMBO J.* **20**, 7160–7167

(2002).

36. Kallipolitis, B. H., Nørregaard-Madsen, M. & Valentin-Hansen, P. Protein-protein

communication: Structural model of the repression complex formed by CytR and the global

regulator CRP. *Cell* **89**, 1101–1109 (1997).

37. Meibom, K. L., Kallipolitis, B. H., Ebright, R. H. & Valentin-Hansen, P. Identification of the

subunit of cAMP receptor protein (CRP) that functionally interacts with CytR in CRP-CytR-

mediated transcriptional repression. *J. Biol. Chem.* **275**, 11951–11956 (2000).

38. Rhodius, V. A. & Busby, S. J. W. Interactions between Activating Region 3 of the Escherichia

coli cyclic AMP receptor protein and region 4 of the RNA polymerase σ70 subunit:

Application of suppression genetics. *J. Mol. Biol.* **299**, 311–324 (2000).

39. Hayes, C. A., Dalia, T. N. & Dalia, A. B. Systematic genetic dissection of PTS in Vibrio

cholerae uncovers a novel glucose transporter and a limited role for PTS during infection of a

mammalian host. *Mol. Microbiol.* **104**, 568–579 (2017).

40. Haycocks, J. R. J. J. *et al.* The quorum sensing transcription factor AphA directly regulates

natural competence in Vibrio cholerae. *PLoS Genet.* **15**, e1008362 (2019).

41. Levine, M. M. *et al.* The pathogenicity of nonenterotoxigenic vibrio cholerae serogroup 01

biotype ei tor isolated from sewage water in brazil. *J. Infect. Dis.* **145**, 296–299 (1982).

42. Skorupski, K. & Taylor, R. K. Positive selection vectors for allelic exchange. *Gene* **169**, 47–52

(1996).

43. Dalia, A. B., McDonough, E. K. & Camilli, A. Multiplex genome editing by natural

transformation. *PNAS* **111**, 8937–8942 (2014).

44. Dalia, A. B. Natural cotransformation and multiplex genome editing by natural transformation

(MuGENT) of vibrio cholerae. in *Methods in Molecular Biology* vol. 1839 53–64 (2018).

45. Page, L., Griffiths, L. & Cole, J. A. Different physiological roles of two independent pathways

for nitrite reduction to ammonia by enteric bacteria. *Arch. Microbiol.* **154**, 349–354 (1990).

46. Sharma, P. *et al.* The multiple antibiotic resistance operon of enteric bacteria controls DNA

repair and outer membrane integrity. *Nat. Commun.* **8**, 1444 (2017).

47. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,

357–359 (2012).

48. Carver, T., Thomson, N., Bleasby, A., Berriman, M. & Parkhill, J. DNAPlotter: circular and

linear interactive genome visualization. *Bioinformatics* **25**, 119–120 (2008).

49. Carver, T., Harris, S. R., Berriman, M., Parkhill, J. & Mcquillan, J. A. Artemis: an integrated

platform for visualization and analysis of high-throughput sequence-based experimental data.

28, 464–469 (2012).

50. Bailey, T. L. *et al.* MEME Suite: Tools for motif discovery and searching. *Nucleic Acids Res.*

37, (2009).

51. Miller, J. *Experiments in molecular genetics. Experiments in molecular genetics* (Cold Spring

Harbor Laboratory, 1972).

ACKNOWLEDGEMENTS

This work was funded by BBSRC project grant BB/N005961/1 awarded to DCG, a BBSRC

MIBTP studentship (BB/M01116X/1, project reference 1898542) awarded to LMW, and

grant R35GM128674 from the National Institutes of Health (to ABD). We thank Kai

Papenfort, Jenny Ritchie and Joseph Wade, for commenting on the manuscript prior to

submission, and Melanie Blokesch for helpful discussions.

MATERIALS AVAILABILITY STATEMENT

Strains, plasmids and oligonucleotides are available on request.

FIGURE LEGENDS

Figure 1: Genome-wide distribution of HapR and LuxO in *Vibrio cholerae*.
610

a. **Simplified schematic overview of quorum sensing in *Vibrio cholerae*.** At low cell density,
expression of HapR is repressed by the Qrr sRNAs that depend on phosphorylated LuxO for
activation of their transcription. Arrows indicate activation and bar ended lines indicate
repression. For clarity, not all protein factors involved in the cascade have been included.
b. **Binding of LuxO and HapR across both *Vibrio cholerae* chromosomes.** In each plot the
outer two tracks (blue) are genes orientated in the forward or reverse direction. The LuxO and
HapR ChIP-seq binding signals are shown in red and green. LuxO binding peaks
corresponding to the *qrr1-4* loci are indicated. Tick marks are 0.25 Mbp apart.
c. **Example LuxO and HapR ChIP-seq binding peaks.** ChIP-seq coverage plots are shown for
individual experimental replicates. Data for LuxO and HapR are in green and red

respectively. Signals above or below the horizontal line correspond to reads mapping to the
top or bottom strand respectively. Gene are show as block arrows.
d. Sequence motifs derived from LuxO and HapR binding peaks using MEME.
e. Positions of LuxO and HapR binding peaks with respect to genes. The histograms show
the distribution of binding peak centres with respect to the start codon of the nearest gene.
f. Pie charts showing gene classes targeted by LuxO and HapR.
627
628 **Figure 2: HapR is a direct repressor of transcription at many target promoters.**
629
a. Activity of HapR targeted promoters in the presence and absence of HapR *in vivo*. The
promoter regions of HapR targeted genes were fused to *lacZ* in plasmid pRW50T and
constructs used to transform required bacterial strains. β-galactosidase activity was measured
in cell lysates taken from *Vibrio cholerae* E7946 (bars) or the Δ*hapR* derivative (open bars).
containing the *VC0857* promoter cloned upstream of *lacZ*. Standard deviation is shown for
three independent biological replicates. Cells were grown in LB-Lennox medium at 37 ºC to
an OD_{650} of ~1.1. Promoters were classified as inactive if, in both the presence and absence of
HapR, β-galactosidase activity was <2-fold higher than the equivalent no insert control. We
have labelled promoters with gene names or locus tags as most appropriate. Note that the
Table 1 footnote can be used to cross reference between locus tags and gene names where
relevant.
b. Activity of HapR targeted promoters in the presence and absence of HapR *in vitro*. The
gel images show results of *in vitro* transcription experiments. The DNA templates were
plasmid pSR derivatives containing the indicated regulatory regions. Experiments were done
with 0.4 µM RNA polymerase in the presence (0.25, 0.75, 1.0, 3.0, or 5.0 µM) and absence of
HapR. Except for the *VC1375* promoter, where the maximum HapR concentration was 3.0
µM. The RNAI transcript is plasmid-derived and acts as an internal control. Expected
transcript sizes, based on results from global transcription start site mapping experiments[32],
are indicated. Note that no *VC1403* transcript was detected in this prior study[32].
649
650 **Figure 3: Transcription from the *murQP* promoter requires CRP *in vivo* and *in vitro*.**
651
652 **a. HapR binding to the *murQP* regulatory region.** Genes are shown as block arrows. ChIP-
653 seq coverage plots are shown for individual experimental replicates. Signals above or below
654 the horizontal line correspond to reads mapping to the top or bottom strand respectively.
b. DNA sequence of the intergenic region upstream of *murQP*. For clarity, numbering is with
respect to the *murQP* transcription start site (TSS, +1). The TSS and promoter -10 element are
in bold. The *murQ* start codon is in blue. The HapR binding site, predicted by MEME analysis
of our ChIP-seq data for HapR, is in green. A potential CRP site is embedded within the
HapR binding sequence (orange). Sequences in red indicate point mutations used in this work.
Triangles show sites of truncation.
c. Binding of CRP to the *murQP* regulatory region and derivatives. Electrophoretic mobility
shift assays showing migration of the *murQP* regulatory region, or indicated derivatives, with
or without 0.1 µM CRP. The DNA fragment used is shown above each pair of lanes and
correspond to the truncations or point mutations indicated in panel b.
d. The *murQP* promoter is activated by CRP *in vitro*. The gel image shows the result of an *in
vitro* transcription assay. The DNA template was plasmid pSR carrying the *murQP* regulatory
region. Experiments were done with 0.4 µM RNA polymerase with or without 0.125, 0.25, or
0.5 µM CRP. The RNAI transcript is plasmid-derived and acts as an internal control.
e. The *murQP* promoter is activated by CRP *in vivo*. The bar chart shows results of
β-galactosidase activity assays. Cell lysates were obtained from wild type *V. cholerae* E7946
(solid green) or the Δ*crp* derivative, transformed with pRW50T derivatives containing the
indicated promoter derivatives fused to *lacZ*. Standard deviation is shown for three
independent biological replicates. Cells were grown in LB-Lennox medium at 37ºC to an
OD_{650} of ~1.1.

Figure 4: HapR and CRP co-operatively bind the same section of *murQP* regulatory DNA.

a. **Binding locations of HapR and CRP upstream of *murQP*.** The gel shows the result of DNase I footprinting experiment. The gel is calibrated with Sanger sequencing reactions. The pattern of DNase I cleavage in the absence of any proteins is in lane 1. Protection of DNA from DNase I cleavage in the presence of 0.11, 0.23 or 0.45 µM CRP is shown in lanes 2-4. Sites of DNAse I hypersensitivity due to CRP binding are indicated by orange triangles. Protection from DNase I cleavage in the presence of 0.5, 1.0, 2.0 or 3.0 µM HapR is shown in lanes 5-8. Protection from DNase I cleavage, dependent on HapR, is shown by a green bar. A DNAse I hypersensitive band, unique to reactions with HapR, is shown by a green triangle. In the presence of 0.45 µM CRP, increasing concentrations of HapR result in a different DNAse I cleavage pattern, including the appearance of a different site of hypersensitivity (black triangle).

b. **Binding of HapR and CRP upstream of *murQP* is co-operative.** Electrophoretic mobility shift assays showing migration of the *murQP* regulatory region with different combinations of CRP (0.025, 0.05, 0.1 or 0.2 µM) and HapR (0.5, 1.0, 2.0, 3.0 or 4.0 µM). For incubations with both factors, the same range of HapR concentrations was used with 0.2 µM CRP.

c. **Co-operative binding of CRP requires the shared HapR and CRP binding site.** Results of
an electrophoretic mobility shift assay, using the wild type *murQP* regulatory region or a
derivative with two point mutations in the shared recognition sequence, for HapR (4.0 µM)
and CRP (0.1 µM). Positions of mutations are shown in Figure 3b.

d. **HapR blocks CRP mediated activation of the *murQP* promoter *in vitro*.** The gel image
shows the result of an *in vitro* transcription assay. The DNA template was plasmid pSR
carrying the *murQP* regulatory region. Experiments were done with 0.4 µM RNA
polymerase, with or without 0.05, 0.1, 0.2 or 0.5 µM CRP 0.5, 1.0, 2.0 or 3.0 µM HapR, as
indicated. The RNAI transcript is plasmid-derived and acts as an internal control.

e. **HapR represses CRP mediated activation of the *murQP* promoter *in vivo*.** β-galactosidase
activity was measured in cell lysates taken from *Vibrio cholerae* E7946 (solid green bars),
Δ*hapR* derivative (open green bars), Δ*crp* variant (open orange bars), or cells lacking both
factors (orange outline with green patterned fill). Standard deviation is shown for three
independent biological replicates. Cells were grown in LB-Lennox medium to an OD_{650} of
~1.0 at 37 ºC.

Figure 5: HapR contacts Activation Region 3 of CRP.

a. **Binding sites for CRP and HapR are optimally aligned when offset by one base pair.** The
panel shows DNA sequences logos generated by aligning binding sites identified by ChIP-seq
analysis for CRP (top) and HapR (bottom). The centre of each motif is indicated by a dashed
line.

b. **Global overlap of CRP and HapR binding sites.** A position weight matrix (PWM),
corresponding to each DNA sequence logo shown in panel a, was created. The PWMs were
used to search the *V. cholerae* genome sequence using FIMO. Distances between the
identified CRP and HapR sites were calculated. Proximal sites were always overlapping and
offset by one base pair (top panel). Overlap was greatly reduced when the analysis was
applied to a randomised version of the same genome sequence (bottom panel).

c. **Model of the DNA-CRP-HapR complex**. The model was generated using PDB submissions
6pb6 (*E. coli* CRP in complex with a class II CRP dependent promoter) and 1jt0 (*S. aureus*
QacR bound to its DNA target). Note that QacR is closely related to *V. cholerae* HapR. The
structures were aligned so that the CRP and HapR binding centres were offset by one base
pair. Residue E55 of CRP (blue) is within Activating Region 3 of CRP that can interact with
the RNA polymerase sigma subunit at class II promoters. HapR residue R123 (red)
participates in HapR dimerisation and is proximal to E55 of CRP.

    d. Side chain E55 of CRP is required for stability of the DNA-CRP-HapR complex. Electrophoretic mobility shift assays showing migration of the *murQP* regulatory region with different combinations of CRP or CRP^{E55A} (0.15, 0.3 or 0.6 µM) and HapR (0.083, 0.125, 0.166 0.208 or 0.25 µM).

    e. HapR cannot repress transcription activated by CRP^{E55A}. Result of an *in vitro* transcription assay. The DNA template was plasmid pSR carrying the *murQP* regulatory region. Experiments were done with 0.4 µM RNA polymerase, with or without 0.05, 0.1, 0.2 or 0.5 µM CRP or CRP^{E55A} and 0.5, 1.0, 2.0 or 3.0 µM HapR, in the presence of 0.2 µM CRP, as indicated. The RNAI transcript is plasmid-derived and acts as an internal control.

Figure 6: Control of *murQP* expression by CRP and HapR at low and high cell density.

    a. *V. cholerae* locked at high cell density are defective for growth using MurNAc as the sole carbon source. Each panel illustrates the optical density of *V. cholerae* cultures at different timepoints after inoculation. Cells were grown at 32 ºC in M9 minimal media supplemented with the indicated carbon source (0.25 % *w/v*) or LB-Miller medium. Cells lacking *luxO*, but not *luxO* and *hapR*, mimic the high cell density state. Error bars show standard deviation from three separate experimental replicates.

    b. Model for coordination of MurNAc catabolism by CRP and HapR. In low *V. cholerae* population density conditions (left panel) cell division necessitates cell wall turnover. Expression of MurQP facilitates cell wall recycling and conversion of MurNAc to GlcNAc 6P for glycolysis (insert). At high cell density conditions (right panel) *V. cholerae* form biofilms on chitinous surfaces. Reduced cell division, and the availability of chitin derived GlcNAc 6P, reduces the need for MurQP.

SUPPLEMENTARY FIGURE LEGENDS

Figure 1-figure supplement 1: Binding of LuxO and the *qrr1* and *VC1142* loci.

    a. ChIP-seq coverage plots for individual experimental replicates. Signals above or below the horizontal line correspond to reads mapping to the top or bottom strand respectively.

    b. Sequence of the intergenic region between *VC1142* and *VC1143*. The LuxO target site is shown in red. Start codons for the divergent genes *VC1142* and *VC1143* are in green. Transcription start sites identified by Papenfort *et al*[32] are in uppercase with arrows indicating the direction of transcription. Underlined sequences are potential promoter -10 elements for the housekeeping RNA polymerase assuming a minimal requirement for the sequence 5'-TANNNT-3'.

Figure 1-figure supplement 2: Example HapR binding signals.

    a. Binding peaks for HapR that fall above our cut-off for peak selection. The HapR ChIP-seq binding profiles are shown in green and genes are shown as blue arrows.

    b. Binding peaks for HapR, at known targets, that fall below our cut-off for peak selection. Binding signals for HapR are shown at known target genes. These peaks for not selected by our analysis because the signal was too weak and/or insufficiently reproducible.

Figure 2-figure supplement 1: Original gel images.

Figure 4-figure supplement 1: Binding of HapR to the *hapR* promoter region in the presence and absence of CRP. Electrophoretic mobility shift assay showing migration of the *hapR* regulatory region with different combinations of CRP (0.0125, 0.025, 0.05 or 0.1 µM) and HapR (0.0625, 0.125, 0.25 or 0.5 µM). For incubations with both factors, the same range of HapR concentrations was used with 0.1 µM CRP.

Figure 5-figure supplement 1: Global overlap of CRP and HapR binding sites in *Vibrio harveyi*. A position weight matrix (PWM), corresponding to each DNA sequence logo shown in Figure 5a, was created. The PWMs were used to search the *V. harveyi* genome sequence (strain ATCC 33843) using FIMO. Distances between the identified CRP and HapR sites were calculated. Proximal sites were always overlapping and offset by one base pair (top panel). Overlap was greatly reduced when the analysis was applied to a randomised version of the same genome sequence (bottom panel).

Figure 5-figure supplement 2: Models of the DNA-CRP and DNA-CRP-HapR complexes. The models were generated using PDB submissions 6pb6 (*E. coli* CRP in complex with a class II CRP dependent promoter) and 1jt0 (S. aureus QacR bound to its DNA target). The DNA is shown in grey and positions hypersensitive to DNAse I cleave, in the context of the DNA-CRP complex, are highlighted red (Figure 4a). DNA position -34 is not cleaved by DNAse I in the context of the ternary DNA-CRP-HapR complex (Figure 4a). Consistent with this, position -34 is obscured by HapR binding.

Figure 5-figure supplement 3: Binding of CRP and HapR derivatives to P*murQP*. The figures shows results of electrophoretic mobility shift assays with CRP and derivatives (0.1, 0.2, 0.4 or 0.8 µM) or HapR and derivatives (0.25, 0.5, 1.0, 2.0 or 4.0 µM).

Figure 5-figure supplement 4: Co-operative DNA binding of HapR and CRP is common. Electrophoretic mobility shift assays showing migration of the indicated regulatory regions with different combinations of CRP (1 µM) and HapR (0.19 µM). For *VCA0691* the concentration of HapR was 0.57 µM.

SOURCE DATA LEGENDS

Figure 2b source data 1: Gel image TIFF file
Figure 2b source data 2: Gel image TIFF file
Figure 2b source data 3: Gel image TIFF file
Figure 2b source data 4: Gel image TIFF file
Figure 2b source data 5: Gel image TIFF file
Figure 2b source data 6: Gel image TIFF file
Figure 3c source data 1: Gel image TIFF file
Figure 3c source data 2: Gel image TIFF file
Figure 3d source data: Gel image TIFF file
Figure 4a source data: Gel image TIFF file
Figure 4b source data: Gel image TIFF file
Figure 4c source data 1: Gel image TIFF file
Figure 4c source data 2: Gel image TIFF file
Figure 4d source data: Gel image TIFF file
Figure 5c source data: Gel image TIFF file
Figure 5e source data: Gel image TIFF file
Figure 4-figure supplement 1: Gel image TIFF file
Figure 5-figure supplement 3 source data 1: Gel image TIFF file
Figure 5-figure supplement 3 source data 2: Gel image TIFF file
Figure 5-figure supplement 4 source data 1: Gel image TIFF file
Figure 5-figure supplement 4 source data 2: Gel image TIFF file
Figure 5-figure supplement 4 source data 3: Gel image TIFF file
Figure 5-figure supplement 4 source data 4: Gel image TIFF file

829 **Table 1: Locations of binding peaks from ChIP-seq experiments**

peak centre	gene(s)[4]	site location	site sequence	TSS[5]
		HapR ChIP-seq peaks		
		chromosome I		
99874	*VC0102<(VC0103)*	99863.5	aaattaataaaactgtcattta	99906 (+)
213457	*(VC0205)>VC0206*	213452.5	taattgtgattcttatcaccaa	213494 (+)
246366[1]	*VC0240<>VC0241*	246349.5	taattaagatggctataaacta	246430 (-)
463584	*VC0433*			
514422	*VC0484*	514430.5	ctactgaccttttcatcaataa	514427 (+)
516570	*(VC0486)*	516601.5	caactgagaaggcacacaatag	516545 (+)
534714	*(VC0502)*	534691.5	ctattataagctctatcagtgt	534805 (-)
547108	*VC0515*	547135.5	atagtaatattattgttaatag	549431 (-)
613328[3]	*VC0583[A]*	613357.5	ttattgagtgggtacataacaa	613427 (+)
716707	*VC0668*	716625.5	ctattgatgaggttatccacag	716537 (-)
735309	*VC0687<>VC0688*			
882854	*(VC0822)*	882825.5	taattatccactttatcaattg	883072 (-)
941187	*VC0880*	941164.5	cttttgacatttctgtcacaaa	941152 (+)
978577	*VC0916[R]*	978540.5	taattaatatccagctcaatta	978581 (+)
1356743	*VC1280<>VC1281[A]*	1356736.5	atattgatagaaataacaagtc	1356896 (+)
1379202	*VC1298<>VC1299*	1379180.5	ttcatgatagtttttgtaattat	1379189 (+)
1469384	*VC1375<>VC1376*	1469377.5	atattgatatatcacacatctt	1469374 (+)
1496023	*VC1403[A]<>VC1405*	1496025.5	tagttgatatttttataattgt	1495942 (+)
1533842	*(VC1437)*	1533854.5	tttgtgagtctcctgtcaataa	1533703 (-)
1990133[2]	*VC1851*	1990076.5	atattgagtaatcaattagtaa	1990031 (+)
2364721	*(VC2212)*	2364680.5	ctattaacagttttatttataa	2364774 (+)
2509878	*VC2352*	2509882.5	ttagtgacagatgcgtcattaa	2509790 (-)
2667349	*VC2486*	2667368.5	taattattaatttgaacaatag	2667206 (-)
		chromosome II		
163808[1]	*VCA0148*	163810.5	taattgattattgtgtaactat	163852 (-)
214589	*(VCA0198)*	214582.5	taattgataactttgacagtat	213494 (+)
237008	*VCA0218<>VCA0219[R]*	237019.5	taaataatatgaatatcagtaa	237053 (+)
247286	*VCA0224<>VCA0225*	247241.5	taaatgactaataagacaatta	247165 (-)
598444	*VCA0662<>VCA0663[R]*	598403.5	tttgtaataaatttgtcattaa	598413 (+)
630517	*VCA0691[A,]*	630559.5	ctattaacaggactgacattaa	631303 (+)
862737	*VCA0906*			
910196	*VCA0960[R]<>VCA0961*	910181.5	ctgattataaatttgtaaatat	910330 (+)
1021174	*VCA1070*	1021117.5	ctcctatccgattggtcactat	1021326 (+)
		LuxO ChIP-seq peaks		
		chromosome I		
1090129	*qrr1<> VC1021*	1090154	ttgcaaaatgcaa	1090182 (+)
1212442	*VC1142<>VC1143*	1212435	ttgcaaatcgcga	1212403 (-)
		chromosome II		

875	48415	*qrr2*	48347	`ttgcaatttgcaa`	48851 (-)
876	772208	*qrr3*	772149	`ttgcattttgcaa`	772227 (+)
877	908445	*qrr4*	908436	`ttgcaatttgcaa`	908475 (+)

878
[1]Identified by Tsou and co-worlers[28]
[2]Identified by Waters and co-workers[29]
[3]Identified by Lin and co-workers[27]
[4]Identified as activated (A) or repressed (R) by Nielsen and co-workers[23], VC0206, VC0240, VC0241, VC0583,
VC0668, VC0916, VC1021, VC1142, VC1143, VCA0219 correspond to *murQ*, *rfaD*, *rfbA*, *hapR*, *mutH*, *vpsU*, *luxO*,
cspD, clpS, hlyA respectively.
[5]Nearest transcription start site (TSS) identified by Papenfort *et al.*[32] with the symbol in parenthesis indicating the
direction of transcription. Note that these are not necessarily those TSSs subject to regulation by HarR or LuxO,
particularly if the regulators are acting as repressors, or if the gene subject to regulation is switched off for another
reason in the conditions of Papenfort *et al.*

a

b

a

b

```
      -270            -260            -250            -240            -230            -220            -210
       •               •               •               •               •               •               •
caccaatctggcggccactctcggaaagttgggtgcgacggctgacgatgcgttgtagtacggacattttttt

      -200            -190            -180            -170            -160            -150            -130
       •               •               •               •               •               •               •
acactcctcatgaataaattattccattgtaggaataatgcctgccacgcttcatgcttttcaatcaatgg

      -120            -110            -100             -90             -80             -70             -60
       •               •               •               •               •               •               •
tttcatggttgagcgtttattttttctcgatgctttttcacgattattccaaaaaacggctgcttaaagaataa
                                                                  ▲                                   ▲
                                                                 Δ183                                Δ211

      -50             -40              *              -30             -20             -10      +1      +10
       •               •               •               •               •               •               •
cct taattgtgattcttatcaccaa aatggaatatttttattcca **tagaat** cgcctc**A**attcgaaatttgccg
     g                     g
    CRP/HapR motif

      +20
       •
agaagcctt**atg**
```

c

d

e

a

markers

C T A G

[HapR]

[CRP]

-150
-140
-130
-120
-110
-100
-90
-80
-70
-60
-50
-40
-30
-20
-10
+1
+10
+20

Lane 1 2 3 4 5 6 7 8 9 10 11 12

b

[HapR]

[CRP]

HapR-CRP-DNA
CRP-DNA
P*murQP* DNA

Lane 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15

c

wild type   -49g -35g

HapR - - + +   - - + +
CRP   - + - +   - + - +

HapR-CRP-DNA
CRP-DNA
P*murQP* DNA

Lane 1 2 3 4   5 6 7 8

d

marker

[HapR]

[CRP]

120 nt
110 nt
100 nt

P*murQP* RNA
RNAI

Lane 1 2 3 4 5 6 7 8
% of max. activity 0 100 98 85 79 40 15 10

e

β-galactosidase activity (Miller units)

wild type   Δ*hapR*   Δ*crp*   Δ*crp*Δ*hapR*

V. cholerae strain

a Alignment of CRP (top) and HapR (bottom) binding motifs

binding motif centres offset by 1 bp

position

b *V. cholerae* N16961 genome sequence

HapR 55 7 CRP 82

V. cholerae N16961 scrambled sequence

HapR 61 1 CRP 88

number of binding sites

Distance between HapR and CRP sites

c CRP E55 R123 HapR

90°

d [HapR] [CRP]

HapR-CRP-DNA
CRP-DNA
P*murQ* DNA

Lane	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
% of CRP-DNA complex bound by HapR	N/A	N/A	N/A	N/A	52	69	100	100	100	N/A	N/A	N/A	N/A	36	50	53	79	91

wild type | E55A

CRP derivative

e marker [HapR] [CRP]

130 nt
120 nt — P*murQ* RNA
110 nt — RNAI
100 nt
90 nt

Lane	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27
% of max. activity for each CRP derivative	0	77	100	81	34	11	10	2	2	0	5	18	44	100	47	41	26	20	0	0	13	44	100	47	39	30	29

wild type | E55A | E55R

CRP derivative

a

b