A structured evaluation of genome-scale constraint-based modeling tools for microbial consortia

William T. Scott Jr.[1,2*‡], Sara Benito-Vaquerizo[1‡], Johannes Zimmerman[3], Djordje Bajić [4], Almut Heinken [5], Maria Suarez-Diez [1], Peter J. Schaap[1,2]

1 Laboratory of Systems and Synthetic Biology, Wageningen University & Research, Wageningen, the Netherlands. **2** UNLOCK, Wageningen University & Research and Delft University of Technology, the Netherlands. **3** Christian-Albrechts-University Kiel, Institute of Experimental Medicine, Research Group Medical Systems Biology, Michaelis-Str. 5, Kiel, 24105, Germany **4** Department of Biotechnology, Delft University of Technology, Van der Maasweg 9, 2629, HZ, Delft, the Netherlands **5** Inserm U1256 NGERE, Université de Lorraine, Nancy, France

‡These authors contributed equally to this work.
* william.scott@wur.nl

Abstract

Harnessing the power of microbial consortia is integral to a diverse range of sectors, from healthcare to biotechnology to environmental remediation. To fully realize this potential, it is critical to understand the mechanisms behind the interactions that structure microbial consortia and determine their functions. Constraint-based reconstruction and analysis (COBRA) approaches, employing genome-scale metabolic models (GEMs), have emerged as the state-of-the-art tool to simulate the behavior of microbial communities from their constituent genomes. In the last decade, many tools have been developed that use COBRA approaches to simulate multi-species consortia, under either steady-state, dynamic, or spatiotemporally varying scenarios. Yet, these tools have not been systematically evaluated regarding their software quality, most suitable application, and predictive power. Hence, it is uncertain which tools users should apply to their system and what are the most urgent directions that developers should take in the future to improve existing capacities.

This study conducted a systematic evaluation of COBRA-based tools for microbial communities using datasets from two-member communities as test cases. First, we performed a qualitative assessment in which we evaluated 24 published tools based on a list of FAIR (Findability, Accessibility, Interoperability, and Reusability) features essential for software quality. Next, we quantitatively tested the predictions in a subset of 14 of these tools against experimental data from three different case studies: a) syngas fermentation by *C. autoethanogenum* and *C. kluyveri* for the static tools, b) glucose/xylose fermentation with engineered *E. coli* and *S. cerevisiae* for the dynamic tools, and c) a Petri dish of *E. coli* and *S. enterica* for tools incorporating spatiotemporal variation. Our results show varying performance levels of the best qualitatively assessed tools when examining the different categories of tools. The differences in the mathematical formulation of the approaches and their relation to the results were also discussed. Ultimately, we provide recommendations for refining future GEM microbial modeling tools.

Author summary

Constraint-based modeling employing genome-scale reconstructions of microbial species has become one of the most successful approaches for studying, analyzing, and engineering microbial consortia. Over the past decade, many constraint-based modeling tools have been published to examine an immense variety of microbial consortia spanning from the application areas of bioremediation to food and health biotechnology. However, new potential users lack an overview of the quality and performance of existing metabolic modeling tools that would guide their choice. To tackle this issue, we examined 24 tools for genome-scale metabolic modeling of microbial consortia. After an initial qualitative screening, we quantitatively evaluated 14 adequate tools against published experimental data that included different organisms and conditions. We conducted simulations and evaluated model features such as predictive accuracy, computational time, and tractability in capturing critical physiological properties. We found that, generally, more up-to-date, accessible, and documented tools were superior in many important aspects of model quality and performance. Although, in some cases, we observed tradeoffs in older, less elaborate tools that can be more accurate or flexible. This work has broad implications to help researchers navigate the most suitable tools, and suggests to developers opportunities for improvement of the currently existing capabilities for metabolic modeling of multi-species microbial consortia.

Introduction

The goods and services provided by microbial consortia have long been harnessed in biotechnology. Recent years have seen a soar in the use of multi-species microbial consortia beyond their traditional application in the production of food and beverages [1, 2]. For example, microbial consortia are increasingly applied in the production of commodity chemicals [3–5], pharmaceuticals [6], or biofuels [7–9]; the valorization of waste and emissions [10–12], the improvement of sustainable agriculture systems [13–15], and applications in health [16] or bioremediation [17–19]. Besides the obvious advantage over established industrial processes in terms of sustainability, multi-species consortia offer a number of advantages compared to monocultures. These include the reduction of metabolic burden (a major issue) through division of labor [20–22], an enhanced substrate versatility, and an increased robustness to fluctuating environments [23, 24]. In consequence, synthetic ecology – the rational engineering of multi-species microbial consortia – is emerging as a new frontier in biotechnology and biomedicine. In order to advance this frontier, it is imperative to build predictive models that will allow us to design and control the composition and function of microbial communities [25, 26].

Constrained-based metabolic modeling (CBM) is a powerful computational approach that mechanistically predicts microbial metabolic traits from genomes. In the past decades, this method has proven invaluable as a guide for microbial experimental design and for elucidating metabolic engineering strategies [27]. Because metabolic traits are also a central determinant of ecological interactions in microbes (e.g. competition for resources or metabolite sharing), CBM also holds great promise as a predictive and an engineering tool in synthetic ecology [28]. In a nutshell, CBMs use genome-scale metabolic models (GEMs), a mathematical representation of the metabolic network encoded in an organism's genome, to simulate metabolic fluxes in a given environment [29]. In the case of single organisms, one of the most popular methods is Flux balance analysis (FBA) [30]. FBA optimizes a predefined objective function (e.g., biomass production) and assumes steady-state exponential growth (balanced growth). Dynamic-FBA (dFBA) -an extension of FBA- is applied to represent non-continuous

operations, such as batch or fed-batch reactors, by incorporating differential equations [31]
that describe the rate of change of the extracellular fluxes and the mass balances of the [32]
reactor [31, 32]. dFBA assumes a quasi-stationary state where internal dynamics are [33]
supposed to be much faster than external changes of the medium [33]. Going a step [34]
further, another general methodology has been developed to model microbial systems in [35]
which the extracellular environment varies spatially and temporally [34–36]. This [36]
approach is known as spatiotemporal FBA. A spatiotemporal FBA framework typically [37]
consists of partial differential equations (PDEs) conveyed in terms of time and spatial [38]
coordinates as independent variables [35]. The PDEs mainly characterize extracellular [39]
mass balance equations for biomass, metabolite, and potentially other chemical species [40]
concentrations. They account for the transport mechanisms that cause spatial [41]
disparities, including metabolite diffusion and liquid/gas phase convection. Some [42]
spatiotemporal FBA approaches choose to represent microbial biomass as individuals or [43]
agents (IBM) [37–39]. In contrast, others opt to represent microbial biomass at the [44]
population level (PLM) [34, 36]. [45]

With the continuous development of novel synthetic microbial consortia, [46]
considerable efforts have been made to extend constrained-based metabolic modeling to [47]
microbial communities [40–44]. In the past decades, many tools have become available [48]
with the aim of studying microbial interactions in the gut microbiome [45–47] or [49]
simulating the growth of microbial consortia in continuous and non-continuous [50]
environments. The increasing availability of these tools makes the selection process [51]
difficult for the user [48]. A key feature when selecting a tool is to verify whether it [52]
follows the FAIR (Findable, Accessible, Interoperable, and Reusable) principles [49]. In [53]
particular, adhering to the software FAIR guiding principles is best as they assure [54]
quality research maintenance and reproducibility [50, 51]. As such, the findability of a [55]
tool is based on the capacity of the metadata and software to be easily found by both [56]
humans and computers. Accessibility pertains to the ease of knowledge available at [57]
which a software tool can be accessed, possibly including authentication and [58]
authorization. Interoperability refers to the ability to communicate with other software [59]
via exchanged data (or metadata). The reusability of tools is related to how [60]
well-described (by metadata) and appropriately structured the software is so that [61]
outputs/results can be replicated, combined, reinterpreted, reimplemented, and/or used [62]
in different settings. Many studies have reviewed the state of the art of steady-state, [63]
dynamic, or spatiotemporal tools and followed qualitative assessments [40–44, 52–55]. [64]
However, no study has yet reviewed these tools quantitatively. Thus, a systematic [65]
evaluation of the latter element would be highly beneficial for the users and developers [66]
in the field. In this work, we have followed an extensive qualitative assessment to [67]
evaluate FAIR principles of available tools and a quantitative assessment to evaluate the [68]
performance of a subset of tools reproducing available experimental data of two-species [69]
communities. The following case studies were selected to quantitatively evaluate the [70]
tools: i) syngas fermentation by *Clostridium autoethanogenum* and *Clostridium kluyveri* [71]
for the static tools, b) xylose and glucose mixture fermentation with engineered [72]
Escherichia coli and *Saccharomyces cerevisiae* for the dynamic tools, and c) a Petri dish [73]
of *E. coli* and *Salmonella enterica* for the spatiotemporal tools. In particular, we tested [74]
tools that use constrained-based modeling approaches and GEM; based on steady-state, [75]
dynamic, or spatiotemporal conditions and suitable to model synthetic microbial [76]
communities of two species. Ultimately, we made recommendations for the best [77]
modeling tools to use based on qualitative and quantitative performance outcomes and [78]
for the future development and improvement of the tools. [79]

Overview of constrained-based modeling tools/approaches

Most of the community modeling tools available are based on steady-state, dynamic, and spatiotemporal conditions, describing continuous cultures, non-continuous cultures, and complex liquid and/or solid systems, for instance, a preferred environment of a Petri dish, respectively (Fig. 1). Steady-state approaches are suitable to describe

Fig 1. Overview of steady-state, dynamic, and spatiotemporal tools to model microbial communities. The division among steady-state, dynamic, and spatiotemporal tools is not firm since dynamic tools can be used to describe steady-state systems. Likewise, spatiotemporal tools can be used to describe dynamic and steady-state systems. However, most tools were specifically designed to be used for the highest dimensional cases.

growth in chemostats or continuous stir batch reactor (CSBR) systems. These tools require a single GEM of individual species and a community GEM, often generated by the tool. Extracellular metabolites and reactions of single species models must be defined by the user in the same namespace (unified identifier) upon constructing the community model. Most tools require the definition of medium composition, and some require either the relative abundance (microbial composition) or the growth rate of individual species. The community growth is often defined as the objective function,

and/or the species growth. The objective function is maximized under the specified constraints by computing the metabolic fluxes, microbial composition, or species growth rate, thus allowing a solution where microbial interactions can be inferred.

Tools based on dynamic approaches are suitable to describe non-continuous systems, such as batch serum bottles, batch or fed-batch reactors as well as some continuous systems, e.g., temporal dynamics of recovery from a perturbation in a CSBR. GEMs of species in the community are provided separately as inputs. All the tools require the medium composition in the form of initial concentration and substrate uptake rates and the kinetic parameters as inputs. The kinetic parameters are normally based on Michaelis-Menten-like kinetics, and thus, Michaelis-Menten constant (K_m) and maximum uptake rate of substrates ($q_{S_i,m}$) are required parameters. After optimization of individual species' growth rate, we can obtain information on cross-feeding metabolites, concentration dynamics of substrates, products and biomass, and metabolic fluxes.

Finally, the spatiotemporal tools aim to describe 2D dimensional surface environments such as mimicking simple solid-state Petri dish environments. In addition to the required inputs described for dynamic tools, these tools need information on the diffusion parameters. The diffusion parameters consist of diffusion coefficients for biomass and metabolites. The main output in spatiotemporal models is the spatial distribution of extracellular metabolites, biomass of the different species, as well as growth and uptake rates, at any given time point. This information can be then used to resolve space-dependent ecological interactions. For instance, we can observe how growth and competition for substrates occurs at the border of a colony, but not in the interior.

In this study, we evaluated a total of twenty-four tools/approaches based on steady-state (9) [43, 45–47, 56–61], dynamic (8) [62–69] and spatio-temporal (7) [34, 36–39, 70–72] methods according to their usability to model microbial communities using GEMs. A description of the tools/approaches is found in the Supplementary_File1.

Results

Both a qualitative and a quantitative assessment were performed to evaluate the modeling tools and approaches. The assessment workflow began by first constructing a list of 16 essential features for quality constraint-based GEM modeling of microbial communities. Each tool was rated according to performance on these qualitative metrics ranging from 1 for inadequate (Red) to 5 for excellent (Blue). The qualitative features are strongly related to FAIR Guiding Principles for research software which include aspects such as software availability, user support, traceability, interoperability, etc [51]. These 16 features were evaluated from the perspective of a fairly experienced user of COBRA methods and tools. A description of the criteria followed to evaluate each qualitative metric was described in a rubric (Supplementary File; S2 Table). Not all features could be examined since some tools were not readily available, or there was no tool developed for those approaches. These features were marked as 'Not applicable' (NA; grey square).

From the qualitative examination of the tools, a subset of tools/approaches were quantitatively evaluated for their potential to directly model microbial consortia under various conditions. For this subsequent evaluation, we needed data pertaining to substrate uptake rates, biomass composition and growth rates for static tools, substrate and product concentrations and biomass concentrations over time for dynamic tools, and media concentrations and spatial distribution of metabolites for spatiotemporal tools. However, these types of raw data are rarely available along with their metadata.

Therefore, only a few suitable candidate datasets were applicable to the surveyed modeling scenarios. We selected three datasets (Diender et al. [73], Hanly and Henson [74], and Harcombe et al. [34]), one dataset for each category of modeling tools, to use as case studies to validate the predictive capabilities of the tools. Every case study represents a consortium of two species. To have an unbiased comparison of the approaches, tools were not modified or augmented to revise their functionalities. Some tools were also available, but they were not quantitatively assessed since they were too specific to a particular application and not designed for general application.

In these evaluations, two main points were addressed: the quality of the tool for modeling microbial communities, and the predictability of the community behavior by the tool when presented with data from simple test cases.

Qualitative assessment - Static tools/approaches

Fig.2 shows the evaluation of the described features in every tool/approach based on the rubric (S2 Table). Joint FBA was not assessed since it is the oldest approach and did not include important parameters as the relative abundance of species. cFBA, RedCom, and NECom are approaches and therefore, some features were not evaluated (grey squares) since there is no tool developed specifically for those methods. The tools/approaches that best meet FAIR principles are MICOM, MMT, SteadyCom, and cFBA (see scores in S2 Figure). All tools/approaches are accessible for external use except CASINO and RedCom. OptCom is freely available for academic users upon request.

OptCom did not undergo updates throughout the years. SteadyCom is integrated as part of COBRA Toolbox [75] and has had some updates from the original publication. Small issues are fixed from time to time in SteadyCom. However, changes are not documented clearly. MMT and MICOM are routinely updated, and developers fix issues. Besides, all the modifications applied in every update of MICOM are described.

OptCom does not have community repository support, and developers only provide a contact person. SteadyCom and MMT offer good community support within COBRA Toolbox. MICOM has excellent user support since it has two channels available for discussions and support. cFBA only needs the installation of CBMPy [76]. SteadyCom and MMT are already available as part of the COBRA Toolbox, whose installation is fast and easy. MICOM is easy to install as it only needs a sentence of code and the installation of COBRApy [77]. OptCom was run using the 'OptCom' function from MICOM, and thus, they were equally evaluated.

OptCom requires GAMS and BARON as solvers also accessed through GAMS. GAMS is not a free programming language, and therefore, its use is more limited. All the dependencies of cFBA and MICOM are freely accessible. SteadyCom, RedCom, MMT, and NECom are written in Matlab, which is not free, but licenses are inexpensive for a wide group of users. Besides, SteadyCom and MMT require COBRA Toolbox, and RedCom requires *CellnetAnalyzer* [78] that are both freely available. Required solvers are either integrated by default or freely accessible to a wide group of users. CASINO uses the RAVEN Toolbox, a free software suited for Matlab [79].

The developers of OptCom provide a book with tutorials to run the tool. Some tutorials are included in COBRA Toolbox that explains case studies used with SteadyCom. However, they merely explain some aspects of the software. MMT has explanatory tutorials and includes README files for each main function of the tool, but it is not completely maintained to have the latest functionality. MICOM contains an extensive user manual explaining all aspects of the tool, and it is up to date (at the time of preparing this manuscript).

All the tools require certain expertise from the user to be run. OptCom is not considered a user-friendly tool as GAMS is used as the programming language, and it is

Fig 2. Qualitative assessment of the static tools/approaches. The colored squares indicate the evaluation of the specified feature in every tool/approach. The color scale (upper right) goes from excellent (blue) to inadequate (red). When a feature does not apply to the specified tool/approach or the feature was not evaluated, it is indicated as NA (Not applicable; grey). The metrics contained in the figure were inspired by [48] Colored squares with '*' indicate features of OptCom evaluated using the OptCom function from MICOM. Colored squares with '**' indicate that the evaluation is an assumption based on the given information. Scom: SteadyCom; MMT: Microbiome Modelling Toolbox. The tools/approaches are ordered by the year of the latest publication (e.g., MMT).

not an extended programming language. cFBA, MMT, and MICOM require some knowledge of programming and constrained-based modeling by the user. SteadyCom is somehow constrained to the case studies of the original publication, and thus, extending their use to new cases, and GEM requires more knowledge by the user.

Except for CASINO and RedCom where the source code is unavailable, the source code is accessible to all users.OptCom, MICOM, and MMT produce the same results over time. OptCom, cFBA, and MICOM produce consistent solutions without infeasibility or non-convergence issues. SteadyCom and MMT can lead, in some cases, to non-convergence solutions or infeasibilities.

NECom was applied to model a co-culture of two species. OptCom and cFBA have been used to model small microbial consortia (up to 3). CASINO, MMT, and MICOM can be used to model large communities (gut microbiota). SteadyCom and RedCom can be applied to model larger microbial communities than OptCom and cFBA (≈ 9).

OptCom toolbox used 'txt' files as input models that do not meet any community-level standard. cFBA, CASINO, SteadyCom, RedCom, MICOM, and

NECom meet SBML or COBRA community-level standards for model input and output files. Based on the original publication, we hypothesize that CASINO meets the SBML standard for model input and output since the RAVEN Toolbox is employed [79]. MMT meets both SBML and COBRA community-level standards for model inputs and outputs models in COBRA format. RedCom meets COBRA community-level standards. OptCom cannot be adjusted to standard GEM formats. cFBA uses CBMPy, which reads the standard SBML format. MICOM is used with COBRApy, which allows for different GEM formats. CASINO reads models imported in COBRA format. SteadyCom allows for the translation of GEM models from/to other formats. MMT allows for the import of SBML, XML, and COBRA models within the tool, and the user does not need to transform them before running the tool. NECom allows for the use of COBRA models.

Every tool/approach allows for adapting some parameters and constraints with more or less difficulty. MMT and MICOM provide a wide set of visualizations for different analyses. The rest of the tools do not produce visual outputs by default.

Quantitative assessment - Static tools/approaches

cFBA, SteadyCom, MICOM, and MMT scored higher than the other tools in the qualitative assessment (Figure in S1 Figure), and thus, they were also evaluated regarding their performance in reproducing an experimental case study. OptCom was also assessed because it was available as an additional function in MICOM, and it is highly cited ((Table in S1 Table). The case study consisted of syngas fermentation to medium-chain fatty acids by a co-culture of *C. autoethanogenum* and *C. kluyveri*. Syngas is a mixture of H_2, CO and CO_2. *C. autoethanogenum* can thrive on syngas by assimilating CO or CO_2 and H_2 and producing acetate and ethanol as byproducts. *C. kluyveri* does not grow on syngas. It needs acetate and ethanol to grow, and therefore, it depends on the direct cross-feeding of these metabolites by *C. autoethanogenum*. The study of Diender et al. [10] reported data on the steady-state consumption and production rates of CO, acetate, ethanol, butyrate, and caproate. The hydraulic retention time (HRT) of the reactor was fixed in the chemostat, suggesting that the species growth rates and the community growth rate were equal (Fig. 3A). This assumption is also considered in cFBA, SteadyCom, and MMT, as illustrated in Fig.3A. cFBA and MMT were run, constraining the growth rates, and thus were equal to the experiments. SteadyCom allows for the definition of the species growth rates and relative abundance. However, SteadyCom was not feasible in those conditions, and the unconstrained parameters resulted in higher growth rates than the experiments. OptCom and MICOM output lower growth rates than the other tools, being that OptCom's results were closer to the experimental values. MICOM can be run using the 'cooperative tradeoff' method, a two-step method that favors species growth rates. A minimum species growth rate can be enforced, which would likely result in a better fit with the experiments.

In the study of Diender et al., almost all the ethanol produced by *C. autoethanogenum* on CO was fully consumed by *C. kluyveri*, whereas the acetate produced by *C. autoethanogenum* was only partly consumed by *C. kluyveri* (Fig.3B) [80]. The fermentation of acetate and ethanol by *C. kluyveri* led to butyrate (≈ 0.11 mmol h^{-1}), caproate (≈ 0.035 h^{-1}), and H_2 (not showing here). Standard deviations for experimental values were not reported.

All tools predicted the exchange of acetate and ethanol between *C. autoethanogenum* and *C. kluyveri*, except MMT, which only predicted the exchange of ethanol (see fluxes in the online repository: https://doi.org/10.5281/zenodo.7573135). OptCom, SteadyCom, and MICOM predicted that the acetate produced by *C. autoethanogenum* was entirely consumed by *C. kluyveri* (see fluxes in the online repository:

Fig 3. Comparison of tool predictions to experimental data from the study of Diender et al. [10]. (A) Community and species growth rate and (B) Steady-state production rates of the main fermentation products obtained in the fermentation of CO by *C. autoethanogenum* and *C. kluyveri* with the assessed tools. cFBA shows the average fluxes and standard deviation of the samples used with Flux sampling.

https://doi.org/10.5281/zenodo.7573135). However, cFBA predicted acetate production by the co-culture (Fig.3B) and acetate consumption by *C. kluyveri*. Yet, acetate production was lower compared to the acetate measured in chemostat (≈ 0.73 mmol h^{-1}). MMT did not predict the production of acetate. Ethanol was difficult to detect in chemostat ($\prec 0.1$ mM), as most of it was consumed by *C. kluyveri*. Considering detection limits, OptCom and MICOM predicted a very similar rate of ethanol; cFBA predicted more ethanol than in chemostat; and in SteadyCom and MMT, ethanol was fully consumed by *C. kluyveri*. Butyrate production was predicted by cFBA and MMT. Caproate production was over-estimated by all the tools except cFBA, which predicted a value comparable to the experiments, and MMT, which did not predict caproate production.

Following the procedure of the modeling study of the same co-culture [80], flux sampling was used to compute the fluxes with cFBA. While the agreement of experimental data with model predictions is less accurate here than in the previous computational study for the lack of constraints, the new adaptation of cFBA can reproduce the experimental data better than the other tools [10]. MICOM, SteadyCom, and OptCom could also reproduce the cross-feeding of acetate and ethanol, which is a key feature in this co-culture. MMT only reproduced the exchange of ethanol, being infeasible for the CO uptake rate reported in the experiments. The adjustment of the coupling factor ('c') or the use of mgPipe pipeline [47] in MMT could lead to alternate outputs. Regardless, MMT was mainly designed to model the gut microbiome using AGORA models, and thus, it is more suitable to model large communities. The use of FVA as an alternative method to compute fluxes with the existing tools and approaches could also lead to additional information on cross-feeding metabolites

Qualitative assessment - Dynamic tools

From the qualitative assessment of all the dynamic tools, it was apparent that none were flawless (all features being good or excellent). There were, in fact, compromising aspects in many of the tools where tools were superior for some features while flawed for other features (see Fig.3 and supplementary file). For instance, when examining their availability, it was determined that dyMMM, DAPHNE, MMODES, and surfinFBA were good or excellent. Still, the tools were considered merely satisfactory when evaluating their interoperability, flexibility, and reproducibility, except MMODES. In addition, tools such as dMMM, DFBAlab, μbialSim, and MMODES were easy to install and contained accessible and tractable dependencies, but did not score above satisfactory levels for the software being traceable (Fig.4). Nevertheless, there was one tool, d-OptCom, that scored poorly compared with the other methods because the software was not readily available from the developers (available upon reasonable request). It was also constructed under an uncommonly utilized GAMS programming language.

Fig 4. Qualitative assessment of the dynamic tools. Colored squares indicate the evaluation of the specified feature in every tool. The color scale (upper right) goes from excellent (blue) to inadequate (red). When a feature does not apply to the specified tool or the feature was not evaluated, it is indicated as NA (Not applicable; grey). The metrics contained in the figure were inspired by [48]. The tools are ordered by year of publication.

Most of the tools received a good or excellent score for some features. For example, all the tools, except d-OptCom and DFBAlab, are open source (contain source code available to all users). DFBAlab also includes source code reachable to some users, but potential users first need to request the software to gain access as it is not available in

the public domain for all users. All tools were deemed at least satisfactory regarding their reproducibility and potential to visualize outputs. Furthermore, all the tools other than d-OptCom and DAPHNE adhere to COBRA community standards [81] and contain a user-friendly interface, at least to a satisfactory degree. This means most methods consist of relatively simple and direct ways of designating inputs, such as adding GEMs, setting up media compositions, and defining model constraints and kinetic parameters. Also, tools that were built incorporating existing COBRA frameworks, such as COBRAPy or COBRA Toolbox, typically scored well for being user-friendly and following current standards because they assimilate the current community standards and acquire greater functionality from the existing infrastructure.

All the dynamic tools received at least a satisfactory score for their interoperability except MCM, which got a poor score. This is due to MCM having difficulties handling some GEMs, especially those of eukaryotes with multiple compartments. Also, MCM utilizes an in-house convention when processing GEMs; thus, issues could arise when needing to employ the GEM in another tool or database. It is best that GEMs adhere to SBML standards [82]. Most software except MCM and d-OptCom earned at least a satisfactory score for dependencies. MCM earned a relatively low score because of the reliance on a not up-to-date micog.py extension (Python 2.7) for using GEMs. Overall, tools such as MMODES, μbialSim, and dyMMM received superior scores because they are readily available, accessible, and inoperable compared to the other tools.

Quantitative assessment - Dynamic tools

A case study comprising a dataset published by Hanly and Henson [74] was used to validate and quantitatively evaluate the dynamic tools. The experimental setup consists of a bioreactor operating in batch mode with a co-culture of *E. coli* and *S. cerevisiae* designed to study the efficient aerobic consumption of glucose/xylose mixtures. Each microbe utilizes a specific substrate. *S. cerevisiae* only consumes glucose, whereas the engineered *E. coli* strain ZSC113 only consumes xylose. This was done to prevent diauxic growth shown in monoculture for *S. cerevisiae*. The individual biomasses of the two microbial species as well as the concentration dynamics of the substrates glucose and xylose, were studied. Moreover, ethanol concentration dynamics were also measured because *S. cerevisiae* produces it during fermentation, thus, inhibiting the growth of *S. cerevisiae* and *E. coli*.

Furthermore, tools that scored 50 or above when summing up the qualitative scores were deemed of sufficient quality for a further quantitative examination. These tools were DFBAlab, dyMMM, μbialSim, and MMODES. We evaluated each tool's capability of simulating the observed kinetics using the same inputs across the methods. The error distribution for the different methods was determined (see Methods Section for a description of the normalized error calculation). *S. cerevisiae* and *E. coli* biomass concentration increases (5). DFBAlab was the only tool to predict the biomass formation of *E. coli* somewhat accurately though the simulation differed substantially from hours 5 to 12.5, showing no growth while growth was observed (5B). μbialSim and MMODES predicted growth for *E. coli*, but the biomass formations were at least an order of magnitude below measured biomass levels (5B). dyMMM predicted slight growth of *E. coli*. Overall, μbialSim and DFBAlab achieved the best predictions for *E. coli* based on the experimental data with R^2 values of 0.959 and 0.775, respectively. For the *S. cerevisiae* growth kinetics, the dyMMM tool most accurately simulated biomass formation (5D). While DFBAlab predicted growth for *S. cerevisiae*, the final biomass concentration was about 65% less than the experimental value (5D). μbialSim and MMODES simulated slight growth for *S. cerevisiae*.Overall, dyMMM and DFBAlab achieved the best predictions for *S. cerevisiae* based on the experimental data with R^2 values of 0.900 and 0.682, respectively (for all R^2 values, see supplementary materials).

Fig 5. Quantitative assessment of the dynamic tools. Experimental biomass concentration profiles of *E. coli* and *S. cerevisiae* (Panel A and C, respectively). Comparison of tool predictions of biomass concentration profiles with data for *E. coli* and *S. cerevisiae* (Panel B and D, respectively) from the study of Hanly and Henson. [74]

The measure and predicted kinetics for the consumption of glucose and xylose, as well as the formation of ethanol, are illustrated (Fig.6). Both sugars' concentration decrease over time, where glucose is completely consumed around 7.6 h while xylose remains in the medium longer and is used up entirely by around 13.2 h. This is due to a faster glucose uptake rate from *S. cerevisiae* compared to xylose's consumption rate by *E. coli*. There is a steady production of ethanol until the 7.5h duration. Then, there is a decrease in ethanol, perhaps because *S. cerevisiae* consumes ethanol under aerobic conditions. The co-culture featured a high increase in ethanol levels due to *S. cerevisiae* fermentation metabolism; ethanol is an inhibitor of *E. coli* growth. This ethanol inhibition from *S. cerevisiae* on *E. coli* growth was the only interspecies interaction identified in this experiment.

For the sugar substrates glucose and xylose, dyMMM and DFBAlab simulated the kinetics most precisely according to the experimental values (6A and 6B). However, DFBAlab predicted rapid xylose consumption at the end of fermentation. MMODES simulated a linear glucose consumption with a final concentration of about 52 mM, while xylose consumption was not predicted. μbialSim did not simulate glucose consumption, while xylose was consumed slightly with a final concentration of about 35mM. All the tools simulated ethanol production. Nevertheless, none of the tools could accurately estimate the dynamics of ethanol production. This is because the diauxic shift from glucose to ethanol by *S. cerevisiae* cannot be modeled by simple optimization as done by FBA. dFBAlab predicted the magnitude and kinetic characteristics most accurately, while μbialSim and MMODES gave an underestimation for the ethanol concentrations and reflected linear profiles (6C). Overall, DFBAlab achieved the best predictions for glucose, xylose, and ethanol kinetics based on the experimental data with R^2 values of 0.902, 0.978, and 0.440, respectively (for all R^2 values, see supplementary materials).

Fig 6. Quantitative assessment of the dynamic tools. Experimental extracellular metabolite concentration profiles of glucose, xylose, and ethanol (Panel A, C, and E, respectively). Comparison of tool predictions of biomass concentration profiles with data for glucose, xylose, and ethanol (Panel B, D, and F, respectively) from the study of Hanly and Henson. [74]

Qualitative assessment - Spatiotemporal tools

From the qualitative assessment of all the spatiotemporal tools, none of the tools received a perfect score. Many tools were highly rated in many categories, while several were also deficient in many categories (see Fig.7) and supplementary files). For example, COMETS, BacArena, CROMICS received excellent scores for software availability and being open-source because these tools and their source codes are readily available via a GitHub repository or a maintained website, whereas MatNet received an adequate score because neither the tool nor its source code can be easily accessed online. When evaluating the software maintenance, user support, and ease of installation of the software, two tools stood out as excellent: COMETS and BacArena. In contrast, many of the tools, including IndiMeSH, MiMoSA, ACBM, and CROMICS were merely rated as being satisfactory to poor for software maintenance and user support (Fig.7). However, except for MiMoSA, they were regarded as good for simple installation because of the accompanying instructions within the software download. Most of the tools received a good or excellent score for some features. For instance, for reproducibility, scalability, and visualization of results, all tools scored good to excellent, except ACBM and MatNet (for reproducibility), MiMoSA and MatNet (for scalability), and MiMoSA and MatNet (for visualization of results) (7). ACBM did score well for reproducibility because slight changes in the same GEM files caused the tool to be invalid or inoperable. In general, IBM methods scored lower in terms of scalability because of the additional computational costs compared to PLM methods. Most of the tools contained some functionality for visualizing results or rendered outputs which were easy to use by other software tools. Many of the tools have at least a satisfactory level of traceability, reliance on available software, and documentation. Only ACBM scored poorly for traceability because changes to the software cannot be detected, whereas most other tools contain mechanisms to trace various versions, and updates are documented.

All the spatiotemporal tools received at least satisfactory scores except MatNet for

Fig 7. Qualitative assessment of the spatiotemporal tools. Colored squares indicate the evaluation of the specified feature in every tool. The color scale (upper right) goes from excellent (blue) to inadequate (red). When a feature does not apply to the specified tool or the feature was not evaluated, it is indicated as NA (Not applicable; grey). The metrics contained in the figure were inspired by [48]. The tools are ordered by year of publication.

their user-friendly interfaces and numerical stability (7). These relative scores reflect how well the functionality, design, and incorporation of community-devised constraint-based modeling infrastructure into the methods. Also, some tools, such as COMETS and ACBM contain GUI options for novice users who may be less comfortable using command-line interfaces. Many of the tools were numerically stable, using examples implying users can start using the software assuming no infeasibilities occur under basic conditions. Overall, tools such as COMETS, BacArena, and CROMICS received superior scores because they are more readily available, up-to-date, open-source, and contain user-friendlier interfaces compared to the other tools.

Quantitative assessment - Spatiotemporal tools

A case study comprising of a dataset published by Harcombe et al. [34] was used to validate and quantitatively evaluate the spatiotemporal tools. The experimental setup consists of a two-member consortium of two mutant strains of *E. coli* K12 and *S. enterica* LT2 was designed to study the syntrophy between the two stains. *E. coli* K12 is deficient in methionine production, so it relies on methionine production from *S.*

enterica LT2. Conversely, *S. enterica* LT2 relies on the secretion of acetate by *E. coli* K12 because *S. enterica* LT2 cannot uptake lactose under microaerobic conditions. This experiment created the optimal condition to observe a mutualistic relationship where neither species can grow without the other being present. The two-species consortium was grown as a lawn on a Petri dish. Furthermore, experimentally, *E. coli* and *S. enterica* were grown overnight in permissive lactose Hypho minimal media (see Table S2 in [34] for more details) and then mixed at a ratio of 1:99 and 99:1. To examine the impact of time and space on the consortium, on LB, both *E. coli* and *S. enterica* can grow independently, and X-gal (5-bromo-4-chloro-3-indolyl-b-D-galactopyra- noside) was included in the plates so that blue *E. coli* colonies could be distinguished from white *S. enterica* colonies (see [34] for more details) The individual biomasses of the two microbial species were examined and measured over a 48-hour growth cycle by counting colonies. By the end of the cycle, the composition converged even when the inoculum frequencies varied by two orders of magnitude. Furthermore, the study also found a relationship concerning the spatial structure with the metabolite resources being allocated, which caused decreased growth between the species as they were moved further apart.

Fig 8. Quantitative assessment of the spatiotemporal tools. Relative abundance simulation profiles of *E. coli* and *S. enterica* from COMETS (Panel A) and BacArena (Panel B). Note biomasses used per individual organisms were 5 x 10^9 fg and 3 x 10^8 fg for *E. coli* and *S. enterica*, respectively. Experimental data used was from the study of Harcombe et al. [34]

From the qualitative assessment, there were four tools considered to be adequate for a further quantitative survey. Furthermore, from an enumeration of the qualitative scores, tools that scored overall 50 or above were deemed requisite quality (see supplementary material). These tools were COMETS, BacArena, IndiMeSH, and CROMICS. We evaluated each tool's capability of simulating the observed biomass dynamic as well as the spatial composition based on moving the species apart using similar inputs across the methods. The error distribution for the different methods was determined (see Methods Section for a description of the normalized error calculation). Over the 48-hour cycle, the community composition of species converged regardless of whether the initial frequency was 1% or 99% of *E. coli*. COMETS predicted species ratios for the two initial frequency conditions (8). COMETS predicted a composition of 79% ± 4% *E. coli*, which is not significantly different from the experimental frequency of 78% ± 6% (9A). However, BacArena predicted a composition of 63.2% *E. coli* (8B). The predictions of the community composition from IndiMeSH were determined not to be significantly different from both the experimental and COMETS simulation results (p-value of 0.66 and 0.33, respectively, two-tailed t-test) (see supplementary material).

The CROMICS results showed that regardless of the initial frequency, the system converged in a species ratio of 76.3% ± 0.1% for *E. coli* (see supplementary material).

The two-species consortium was used to examine the influence of spatial structure on resource allocation and growth within this mutualistic system. Here, we illustrate the evolution of the colony spatial distribution over time using a PBM (COMETS) and an IBM (BacArena) for their representations of biomass (9). The COMETS simulation shows a rapid rise in the population density of *E. coli* and a shift in the location within the grid from *E. coli* from 9h to 48h (9A). COMETS also predicted a moderate increase in population density in *S. enterica* as well as a slight shift in position from 9h to 48h (9A). BacArena takes a different approach and models each organism individually on a two-dimensional grid to simulate a spatial environment. For BacArena, the colony evolved first around the initial microbial positions of *E. coli* and *S. enterica* where *E. coli* reaches a near-final individual amount t = 9h (9B). The simulated *S. enterica* colony continued to grow more abruptly, however, from t = 9h to t=48h, filling up most of the remaining spatial elements(9B).

As a simple test, we attempted to simulate conditions where the growth of the species should be affected by increasing distance between them. Consistent with what was observed experimentally [34], the simulations using COMETS, IndiMeSH, and CROMICS showed similar degrees of decreased growth as they were initiated further apart (see supplementary material for output data for all the tools). However, the BacArena tool could not capture any variation in spatiotemporal growth as observed during the experiment.

Discussion

This study provides a survey of the chronicled expansion of GEM modeling methods for microbial communities throughout the past decade and recommends the best GEM modeling tools based on qualitative and quantitative assessments. In particular, we assessed tools that used constrained-based modeling methods and their performance to model synthetic microbial communities of two species. This study also illustrates the challenges and brings forth issues and recommendations that should be considered when developing future tools, so that overall serviceability and performance can best facilitate potential users.

The criteria to qualitatively evaluate the tools was based on the FAIR principles that we consider, should be applied for data, operating procedures, tools, and models [49, 51]. Generally, tools that scored the highest in the qualitative assessment were shown to perform best in the quantitative assessment, highlighting the importance of following the FAIR principles. On the contrary, tools that were less 'FAIR' could hardly be accessed nor used in this study, which negatively affects the future development of the tool. A tool needs to be FAIR to be used, to contribute to the field, and thus, to shed light on the continuous challenges that can be, later on, tackled in new tools. In fact, this study observed the latter, since the continual publication of GEM modeling methods for microbial consortia over the past decade, has generally improved in performance over time. The first tools have served as a proof of concept and have established the basis for developing the new tools. However, there were a few exceptions of tools that had high scores in the qualitative assessment and performed worse than the average in the quantitative assessment. In this study, the assessment was based on the modeling of two-species co-cultures, and therefore, the results could have varied when the tools were applied to larger communities, or to alternative systems. In addition, the limited availability of experimental data affected the selection of case studies, which might have an effect on the assessment of tools.

The current status of the array of static, dynamic, and spatiotemporal tools for

Fig 9. Quantitative assessment of the spatiotemporal tools. Colony spatial distribution over time simulation profiles of *E. coli* and *S. enterica* from COMETS (Panel A) and BacArena (Panel B). For Panel B, the red dots represent *E. coli* and the black dots represent *S. enterica*. Experimental data used was from the study of Harcombe et al. [34]

GEM modeling microbial consortia presents a substantial barrier to entry for researchers unfamiliar with specific requirements and idiosyncrasies that may arise within constraint-based modeling frameworks. Tools should be user-friendly enough for a novice user who may be familiar with metabolic modeling concepts such as flux balance analysis and has some basic programming knowledge. We believe that user-friendliness is also linked to the quality of available documentation and the existence of community support. Therefore, we recommend keeping manuals and related repositories up to date and including dependencies amenable to changing computational environments or, alternatively, describing which packages are compatible. When the software is open/free, a possible solution could be to provide a container with the tool and the dependencies. All changes should be equally specified and updated in all places where the tool can be found unless specified otherwise. It is crucial that the tool is accessible to the users. We recommend the tools can read several GEM formats and namespaces without the need to translate the input models using other packages (e.g., COBRA Toolbox, COBRApy). The translation of these models often leads to a loss of information and a lack of key attributes that might generate errors when running the tools. The community models built by some static tools could keep the original bounds of the transport reactions of a single GEM to avoid the user the re-definition of the bounds once the community model is built. This would extend the use of these tools to cases with unspecified mediums/diets. Dynamic tools should allow the possibility of including inhibitory effects to better predict community behavior.

Tools oriented for modeling small and big communities should incorporate sufficient ₅₂₄
series of examples featuring a range of application scenarios from small to big ₅₂₅
communities. Tools could incorporate case studies with GEMs of non-model organisms ₅₂₆
since many case studies are typically proven on communities of several *E. coli* species ₅₂₇
that use the same GEM. Other tools are meant to work with models retrieved from a ₅₂₈
specific database and defined namespace (e.g., AGORA [83]). The applicability of these ₅₂₉
tools to non-model organisms is critical for designing and optimizing synthetic ₅₃₀
communities. In the past decades, many synthetic communities have been established ₅₃₁
that do not depend on model organisms and have shown a high potential for producing ₅₃₂
commodity chemicals [5, 84–87]. On the other hand, GEMs require accurate genome ₅₃₃
annotation and curation to properly account for interactions mediated through ₅₃₄
pathways, but there is still difficulty identifying these components in microbial genomes. ₅₃₅
Therefore, most GEM modeling efforts have been restricted in application to ₅₃₆
high-quality GEMs (*E. coli* and *S. cerevisiae*). To extend the applicability to ₅₃₇
alternative GEMs, the quality of GEMs of single species and communities should follow ₅₃₈
the community standards and, thus, be verified using test suites such as MEMOTE [88]. ₅₃₉

Some tools are based on approaches that use FBA and, therefore, are limited to the ₅₄₀
maximization of the community growth rate or species growth rate. In those cases, FBA ₅₄₁
favors growth overproduction, and cross-feeding metabolites might be overlooked (3B). ₅₄₂
Objective functions can be improved by using experimental data, or frameworks can be ₅₄₃
refined in dynamic tools to use a bi-level objective routine [89]. Yet, these amendments ₅₄₄
may still not apply to all growth scenarios [90]. Flux sampling has been proven to be an ₅₄₅
effective tool to explore metabolism using GEM [32, 91, 92], and its use can mitigate the ₅₄₆
influence of the objective function. Flux sampling has also been successfully ₅₄₇
implemented to model microbial communities [80]. Therefore, we recommend integrating ₅₄₈
flux sampling methods in current and newly developed tools for modeling microbial ₅₄₉
communities in different environments, even though computational costs are high [93]. ₅₅₀

Future tools/approaches could consider using enzyme-constrained models to model ₅₅₁
microbial communities since they have shown great potential to better understand ₅₅₂
microbial phenotypes and phenomena such as overflow metabolism [32]. ₅₅₃

It is of the utmost importance that static tools/approaches incorporate the biomass ₅₅₄
composition of each species in their formulations beyond simply integrating the ratio by ₅₅₅
changing the stoichiometry of their biomass reaction. There is no clear consensus yet ₅₅₆
among static approaches on whether the species grow at balanced growth or whether ₅₅₇
the community and species growth rates differ. This challenges the selection of the ₅₅₈
objective function and the optimization approach of static tools. We believe that static ₅₅₉
tools best fit the simulation of continuous environments where the dilution rate is ₅₆₀
assumed to be constant, and thus, the community growth rate should be the same as ₅₆₁
single species [80]. Techniques that predict variable growth rates among microbes ₅₆₂
within a community should be incorporated into dynamic tools. The drawback, ₅₆₃
nonetheless, is the inability to account for longitudinal community composition. An ₅₆₄
original method for alleviating this limitation is integrating ecological models within ₅₆₅
existing constraint-based frameworks. Some researchers have already begun employing ₅₆₆
COBRA methods and evolutionary game theory [94]. However, in the future, it would ₅₆₇
be beneficial to bolster more tools in this direction. ₅₆₈

The results of this work suggest there is still much room for improving the overall ₅₆₉
quality of GEM modeling tools for microbial communities. Despite the immense number ₅₇₀
of proposed tools over the last decade, the availability, user-friendliness, and overall ₅₇₁
performance quality of the tools are disparate, and there is no perfect tool for all ₅₇₂
scenarios. In addition, although some of the tools which were available and accessible to ₅₇₃
a moderately novice-level user, there were still some flaws in the quantitative ₅₇₄
predictions of those tools. However, we determined some tools were exceptional and ₅₇₅

should be preferred starting points for researchers or developers. ₅₇₆

For instance, for static systems, cFBA is the approach that achieves the best outcome, but it largely requires manual adaptation. Moreover, it is not a tool but merely a generalizable method or algorithm. However, self-contained tools, including MICOM, OptCom, and SteadyCom, all produce reasonable results. None of them outperform the others, so they can be considered as a starting point, and given the usability characteristics, we recommend MICOM. For dynamic systems, the choice is more difficult to discern, but given our results, we suggest dFBAlab. MMODES is also a solid option, although discretion is advised when interpreting results from MMODES as predictions largely disagree with the measurements. Yet ethanol inhibition should be taken into account because the case we simulated contains ethanol inhibition. Therefore, an option or flexibility for augmenting a more complex Michaelis–Menten or Hill kinetic expression to reflect growth rate suppression at high ethanol concentrations would be desirable in a new tool. For spatiotemporal systems, we recommend either COMETS or BacArena because of their extensive development, accessible platforms, and documentation. COMETS performed better in simulating a simple coculture experiment. However, BacArena is an individual-based method that can be beneficial when studying heterogeneous cell populations.

Through this qualitative and quantitative survey, we have presented and analyzed a broad overview of many GEM modeling tools for microbial consortia. These tools have been successfully applied in various types of systems of microbial communities. In addition, while these constraint-based tools have significantly transformed our understanding of communities by incorporating mechanistic details and highlighting metabolic interactions, there are still many opportunities for improving modeling frameworks by making software FAIR, user-friendly, and improving the accuracy of simulations of various types of systems. The work presented here can guide researchers in selecting the proper modeling tools and help developers build upon suitable modeling frameworks for new software tools.

Materials and Methods

An extensive literature review was executed to identify the state of the art of tools for modeling microbial communities in steady-state, dynamic, and spatiotemporal environments that used constrained-based models. Every tool was rigorously studied to follow a qualitative assessment. For that, we created a rubric that described each of the evaluation levels for every feature (S2 Table). Tools were qualitatively assessed based on the latest version unless stated otherwise. In addition, we looked into studies that provided experimental data of synthetic microbial consortia of two species in every environment and assessed the performance of a set of each type of tool to reproduce them. The code used to produce our reported results can be downloaded from: https://doi.org/10.5281/zenodo.7573135

Evaluation of static tools

The case study chosen to evaluate the static tools was the production of medium-chain fatty acids from the fermentation of CO by a co-culture of *C. autoethanogenum* and *C. kluyveri*. The GEM of *C. autoethanogenum*, iCLAU786 [95], and the GEM of *C. kluyveri*, ickl708 [96], were downloaded from their original publications in SBML (xml) format and used as the input files in every tool. Extracellular metabolites (defined in compartment '_e') and exchange reactions ('EX_XX') common in both GEMs were modified to have the same namespace, as required to build the community model in all tools.

Experimental data was obtained from steady-state concentrations of fermentation products, total biomass, and hydraulic retention time (HRT) reported in the study of Diender et al. [10]. In particular, we simulated the condition defined in reactor run number three from the latter publication (Table 1 and Table 3; 116 mmol CO L^{-1} d^{-1} and 0 H$_2$). CO feed rate and steady-state concentrations of fermentation products were converted to mmol h^{-1}, and HRT (d) was converted to the growth rate (h^{-1}). The biomass species ratios were obtained from the computational study of the same co-culture [80], 0.4-0.6 *C. autoethanogenum*-*C. kluyveri*, respectively. CO feed rate was constrained to the experimental value in every tool unless stated otherwise. Additional constraints were applied and are specified per tool in the supplementary File (S3 and S4 Tables). Predicted steady-state fluxes of acetate, ethanol, butyrate, and caproate, as well as the community growth rate, were compared to the experimental measurements.

Below, we describe the specific methods applied to run every tool.

OptCom

OptCom was run using the OptCom function found within the MICOM module called 'micom.optcom', instead of using the code from the original publication. Therefore, we first installed MICOM using pip. The community model was created using the 'Community' function. The function requires a 'taxonomy' table as an input parameter, which contains information about the species, models, and species relative abundance (biomass fraction). In addition, we included the mass as the input parameter (0.22 g L^{-1}). OptCom function was run with the generated community model as input, selecting 'original' as the strategy parameter and setting a min_growth of 0.004 h^{-1} for each species. The community growth rate was maximized simultaneously with all individual growth rates, and fluxes were computed. The individual and community growth rates obtained were divided by the total biomass of the community (0.22 g L^{-1}), as MICOM assumes 1 g of biomass.

cFBA

cFBA was run using the methodology implemented in a previous computational study, where a community model of the same two species was built [80]. The community model was built manually, unifying the extracellular metabolites and extracellular reactions common between species in one single reaction or metabolite. We used the same community model without adding the extra reactions described in the latter publication. The species biomass reactions were constrained to the relative abundance multiplied by the growth rate (g L^{-1} h^{-1}), and the same as the community growth rate (0.021 h^{-1}). The solution space was computed using the 'sample' function in the flux_analysis submodule found in COBRApy. The results shown here are the average and standard deviation based on 10000 samples generated under the specified condition.

SteadyCom

SteadyCom was run using the 'SteadyCom' subroutine in COBRA Toolbox. The community model was built using 'createMultipleSpeciesModel.' The names and IDs for metabolites and exchange reactions in the shared compartment of the community [u] were added into the community model as 'infoCom' and 'indCom' fields using 'createMultipleSpeciesModel.' The bounds of the reactions defined in the [u] compartment (e.g.,'autoIEX_MG[u]tr,' EX_MG[u]') were imported from an excel file. SteadyCom was run using the built community model and the following options as input parameters: GRguess=0.5; algorithm=1; feasCrit=1 and BMweight=0.22. The community growth rate was maximized, and fluxes were computed.

Microbiome Modelling Toolbox (MMT)

MMT was run from COBRA Toolbox. The community model was created using the 'joinModelsPairwiseFromList' function. The coupling factor 'c' and threshold 'u' was maintained at their default values, 400 and 0, respectively. The bounds of the reactions defined in the [u] compartment of the generated community model were imported and defined from an excel file . The interactions and fluxes of the co-culture were next explored using the function called 'simulatePairwiseInteractions.' The species biomass reactions were constrained to the relative abundance multiplied by the growth rate (g L^{-1} h^{-1}). To run the latter function, the community model and the pairedModelInfo.mat file created by 'joinModelsPairwiseFromList' were used as the input parameters. Besides, 'saveSolutionsFlag' was selected to output the fluxes.

MICOM

MICOM was first installed using pip as described in the GitHub repository of MICOM. The community model was created using the 'Community' function following the procedure described in OptCom. The community growth rate was optimized using FBA by the MICOM function 'com.optimize', and fluxes and species growth rates were obtained. The individual and community growth rates obtained were divided by the total biomass of the community (0.22 g).

Evaluation of the dynamic tools

Similar to what was done by Hanly and coworkers [74], *Saccharomyces cerevisiae* S288C (iND750) [97] and *Escherichia coli* K-12 substr. MG1655's (iJR904) [98] GEMs, acquired from the BiGG database [99], were used to perform the model simulations with DyMMM, DFBAlab, MMODES, and μbialSim. Furthermore, the *E. coli* model(iJR904) was modified by constraining flux bounds for glucose exchange and glucose kinase at zero to mathematically reflect the associated gene deletions. The simulations were based on an aerobic xylose co-culture of *S. cerevisiae* and the engineered *E. coli* strain ZSC113 fermentation experiment. In this experiment, glucose and xylose concentrations are expected to decrease over time, while the ethanol concentration is expected to increase. Simultaneously, S. cerevisiae and E. coli biomass concentrations are expected to increase over time. For all simulations, any constraints and parameters given in the original tools were modified according to the experimental values from the respective dataset from Hanly and coworkers [74]. Please see the supplementary table for specific inputs. The simulated growth and metabolite concentration curves of *E. coli* and *S. cerevisiae*, as well as from glucose, xylose, and ethanol were compared to the experimental measurements.

The coefficient of determination, also known as R-squared (R^2) was calculated to quantitatively estimate the quality of the model fits and performance with the experimental data as in the work of Montgomery [100] for the dynamic and spatiotemporal cases.

The equation for R-squared is:
$R^2 = 1 - \frac{SSE}{SST}$
where: R^2 is the coefficient of determination, SSE is the sum of squared residuals, and SST is the total sum of squares, and $\frac{SSE}{SST}$ represents the ratio of the sum of squared residuals to the total sum of squares.

The equation for the sum of squared residuals:
$SSE = \sum_{i=1}^{n}(y_i - \hat{y}_i)^2$
where: SSE is the sum of squared residuals, y_i is the actual value of the dependent variable for the i^{th} observation, $\hat{y}_i$ is the predicted value of the dependent variable for

the i^{th} observation, and n is the number of observations.   719

The equation for the total sum of squares: 720

$SST = \sum_{i=1}^{n}(y_i - \bar{y})^2$ 721

where: SST is the total sum of squares, y_i is the actual value of the dependent   722
variable for the i^{th} observation, $\bar{y}$ is the mean of the dependent variable, and n is the   723
number of observations.   724

dyMMM   725

An implementation for this method is provided by the developers on sourceforge.net: 726
https://sourceforge.net/p/dymmm/wiki/Home/. This method requires a working   727
installation of COBRA Toolbox [101]. The provided formulation allows users to add   728
initial concentrations of limiting or important substrates as well as products of interest.   729
There is also a field to add initial biomass concentrations of microbes. We followed as   730
instructed in the source code by the developers to add the proper inputs regarding   731
initial concentration values of substrates and microbial biomasses, as well as GEMs,   732
where needed. These inputs we added as presented in the supplementary table.   733

DFBAlab   734

An implementation for this method is provided by the developers via the website   735
https://yoric.mit.edu/software. This method requires a working installation of IBM   736
CPLEX or Gurobi. A similar example is supplied in the tutorial to the user as the case   737
study used in this work. Although the original formulation of this method uses values   738
directly from the literature [63], in the implementation, the values are not exactly as   739
reported for our chosen case study. Therefore, the initial concentrations of glucose,   740
xylose, and microbial biomasses were adapted (see supplementary table), and the time   741
span of the simulation was modified from 10 to 14 hours.   742

μbialSim   743

An implementation for this method is supported via the website: 744
https://github.com/fcentler/microbialSim. This method requires a working installation   745
of COBRA Toolbox or CellNetAnalyzer. The provided example 2, based on a   746
batch-culture growth of a binary syntrophic community, served as a starting point for   747
employing our dynamic case study conditions. The reactor was defined according to   748
initial substrate concentrations (see supplementary table). The GEMs (iJR904 and   749
iND750) were loaded and adapted according to the COBRA Toolbox protocol within   750
μbialSim. In addition, reaction indices for the non-growth associate maintenance   751
(NGAM) and biomass reaction were provided. Exchange reactions were defined that   752
link compounds in the reactor to cellular uptake and secretion. The models were   753
parameterized according to the kinetic parameters provided in the literature (see   754
supplementary table). The simulation duration and time step size chosen were 14 and   755
0.02 h, respectively. dFBA was selected as the solver type between the options ODE   756
and dFBA. However, GPLK was used as the COBRA solver.   757

MMODES   758

An implementation for this method is supported via the website: 759
https://mmodes.readthedocs.io/en/latest/index.html. MMODES was installed via pip.   760
The example script and the documentation were used for instructions. First, the   761
medium file was defined according to the medium composition given and initial   762
conditions from the work of Hanly and Henson (see supplementary table). Next, the   763
kinetic parameters were defined and set according to values used by Hanly and Henson   764

(see supplementary table). Then, the GEMs (iJR904 and iND750) of the microbes, along with their respective initial biomasses and substrates they consume, were added to the model framework. The FBA optimization routine using the GPLK solver was selected. We chose a simulation time of 14 h with 0.02 h timesteps. Also, finite element analysis was used for the ordinary differential equation integrator.

Evaluation of the spatiotemporal tools

Similar to what was done by Harcombe et al. [34], simulations were performed using the GEMs for *E. coli* (iJO1366) [102] and *S. enterica* (iRR1083) [103]. We compared BacArena, IndiMeSH, and CROMICS, which were designed to study microbial communities using FBA and IBMs, with COMETS which was mainly designed to investigate the interrelationships of bacterial communities in space using FBA and PLMs. The goal was to compare the methods with one another for their potential in simulating trophic dependences of multispecies bacterial communities without making prior assumptions. Therefore, simulations mimicked the two-member consortium experiment studied in the original publication [34]. This consortium was composed of two mutant strains, *S. enterica* LT2 and *E. coli* K-12. *S. enterica* LT2 cannot metabolize and consume lactose, and *E. coli* K-12 cannot produce methionine. Hence, these two species participate in a mutualistic relationship because *S. enterica* relies on the secretion of acetate for a substrate by *E. coli* while *E. coli* needs *S. enterica* to produce methionine. Stoichiometric models of each species were modified to incorporate known genetic constraints according to the method of Harcombe and coworkers [34]. For instance, in the *E. coli* strain, the metB mutation was accounted for by constraining to zero the flux through the corresponding reaction (cystathionine γ-synthase). In *S. enterica*, methionine necessitated that we added a gain-of-function mutation in metA (homoserine transsuccinylase). This secretion was modeled as coupled to biomass where lactose is utilized by *E. coli*, so that as cells grew, they produced appropriate amino acid levels. A summary of boundary conditions, constraints, and parameters for all GEMS for spatiotemporal methods can be found in the supplementary material.

COMETS

The *in silico* experiment was set up as reported in the work of Harcombe et al. [34] (for more detail, see experimental procedures in the original work). The developers provide an implementation for this method via the website: https://www.runcomets.org/. This method requires a working installation of Java and Gurobi. The Python version of the COMETS Toolbox was installed using the Anaconda distribution.

BacArena

The developers provide a tutorial to help implement this method via https://bacarena.github.io/. This method requires a working installation of R and Sybil. Both GEMs of *E. coli* (iJO1366) and *S. enterica* (STM v1 multistrain) were retrieved from the BiGG database [99]. The *S. enterica* model was modified to ensure methionine production using the method of Harcombe et al. [34]. The biomass reaction was updated to include the production of 0.5 mmol gDW^{-1} of excreted extracellular methionine, which was balanced by an equal amount of intracellular methionine consumed. The methionine transporter (METtex, METabcpp) were set to export only (upper bounds set to 0). The *E. coli* model was modified to block flux through the corresponding reaction, cystathionine γ-synthase (CYSTL). To model metabolic exchanges between the microbes and compare the results of BacArena with the other methods, we performed the simulations with our method using a setup similar to

COMETS [34]. The simulations were carried out on a 50 times 50 grid environment for 48 hours. In the setups, a minimal medium stated in Harcombe et al. [34] was added to the environment with 1 mmol of lactose per grid position, oxygen, and several co-factors (calcium, chlorine, cobalt, potassium, iron, magnesium, ammonium, manganese, nickel, phosphate, zinc, and sulfate). To ensure the growth and mimic the preculture environment, an initial amount of 1.0 mmol methionine and acetate was added to each grid position. The amounts and biomasses of the initial species were also set according to initial frequencies used in Harcombe et al. [34]. The diffusion of metabolites was calibrated to the standard diffusion of glucose.

IndiMeSH

The developers provide an implementation for this method in the supporting materials of the original paper [39]. This method requires a working installation of MATLAB. The in silico experiment was performed to simulate microbial life on a two-dimensional surface in IndiMeSH, a rectangular, saturated pore network (coordination number: 4) was created with the same topology as in the COMETS simulations (for details, see S1A Fig in the supplementary material). Each pore has dimensions of 500x250x70 microns (LxWxH), as specified in the paper. Furthermore, we followed the setup as reported in the work of Borer et al. [39] (for details, see the Methods section in the original paper).

CROMICS

The developers provide an implementation for this method via a GitHub repository: https://github.com/EPFL-LCSB/cromics. This method requires a working installation of MATLAB and CPLEX solver. We used the setup detailed in the original paper [72], which mimics the setup of Harcombe and coworkers [34] for the two-species consortium of *S. enterica* and *E. coli* (for details, see the Methods section in the original paper [72]).

Supporting information

Acknowledgments

P.J.S. and W.S. acknowledge the Dutch Research Council (NWO), and Wageningen University and Research for their financial contribution to the Unlock initiative (NWO: 184.035.007). This work was also supported by the Netherlands Science Foundation (NWO) under the Programme 'Closed Cycles' (Project nr. ALWGK.2016.029) and the Netherlands Ministry of Education, Culture and Science under the Gravitation Grant nr. 024.002.002.

We thank Dr. William R. Harcombe and Dr. Jeremy Chacon from the Department of Biological Sciences at the University of Minnesota for assistance with the COMETS Toolbox for spatiotemporal modeling and interpretation of some experimental and model results. We thank Dr. Costas D. Maranas and Dr. Mohammad M. Islam from the Pennsylvania State University Department of Chemical Engineering for providing the d-OptCom tool and making suggestions for how to use the software. We thank Dr. Louca Stilianos from the Institute of Ecology and Evolution at the University of Oregon for correspondence and assistance employing the Microbial Community Modeler (MCM) tool.

Author Contributions

Authors' contributions: CRediT (Contribution Roles Taxonomy)- Conceptualization, W.T.S.J., S.B-V, M.S-D., P.J.S., and E.B-C.; Data Curation, W.T.S.J. and S.B-V., Formal Analysis, W.T.S.J., S.B-V., and J.Z.; Funding Acquisition, M.S-D, and P.J.S; Investigation, W.T.S.J., and S.B-V; Methodology, W.T.S.J., S.B-V., and M.S-D.; Resources, P.J.S, and M.S-D, Software, J.Z, D.B, and A.H..; Supervision, M.S-D, and P.J.S.; Validation, W.T.S.J. and S.B-V; Visualization, W.T.S.J., and S.B-V.; Writing—original draft, W.T.S.J., and S.B-V; Writing—review and editing, W.T.S.J., S.B-V., J.Z., D.B., A.H, M.S-D, and P.J.S. All authors have read and agreed to the published version of the manuscript.

References

1. Goers L, Freemont P, Polizzi KM. Co-culture systems and technologies: taking synthetic biology to the next level. Journal of the Royal Society Interface. 2014;11. doi:10.1098/RSIF.2014.0065.

2. Jia X, Liu C, Song H, Ding M, Du J, Ma Q, et al. Design, analysis and application of synthetic microbial consortia. Synthetic and Systems Biotechnology. 2016;1(2):109–117. doi:https://doi.org/10.1016/j.synbio.2016.02.001.

3. Cho JS, Kim GB, Eun H, Moon CW, Lee SY. Designing Microbial Cell Factories for the Production of Chemicals. Journal of the American Chemical Society. 2022;2:1781–1799. doi:10.1021/JACSAU.2C00344.

4. Duman-Özdamar ZE, dos Santos VAPM, Hugenholtz J, Suarez-Diez M. Tailoring and optimizing fatty acid production by oleaginous yeasts through the systematic exploration of their physiological fitness. Microbial Cell Factories. 2022;21:1–13. doi:10.1186/S12934-022-01956-5/FIGURES/3.

5. Benito-Vaquerizo S, Nouse N, Schaap PJ, Hugenholtz J, Brul S, Lopez-Contreras AM, et al. Model-driven approach for the production of butyrate from CO2/H2 by a novel co-culture of C. autoethanogenum and C. beijerinckii. Frontiers in Microbiology. 2022;doi:10.1101/2022.10.12.511882.

6. Wang EX, Ding MZ, Ma Q, Dong XT, Yuan YJ. Reorganization of a synthetic microbial consortium for one-step vitamin C fermentation. Microbial Cell Factories. 2016;15:1–11. doi:10.1186/S12934-016-0418-6/FIGURES/5.

7. Kleerebezem R, van Loosdrecht MC. Mixed culture biotechnology for bioenergy production. Current opinion in biotechnology. 2007;18(3):207–212.

8. Senne de Oliveira Lino F, Bajic D, Vila JCC, Sánchez A, Sommer MOA. Complex yeast–bacteria interactions affect the yield of industrial ethanol fermentation. Nature communications. 2021;12(1):1498.

9. Alper H, Stephanopoulos G. Engineering for biofuels: exploiting innate microbial capacity or importing biosynthetic potential? Nature Reviews Microbiology. 2009;7(10):715–723.

10. Diender M, Olm IP, Gelderloos M, Koehorst JJ, Schaap PJ, Stams AJM, et al. Metabolic shift induced by synthetic co-cultivation promotes high yield of chain elongated acids from syngas. Sci Rep. 2019;9:1–11. doi:10.1038/s41598-019-54445-y.

11. Hu J, Xue Y, Guo H, Gao Mt, Li J, Zhang S, et al. Design and composition of synthetic fungal-bacterial microbial consortia that improve lignocellulolytic enzyme activity. Bioresource Technology. 2017;227:247–255.

12. Cabau-Peinado O, Straathof AJ, Jourdin L. A general model for biofilm-driven microbial electrosynthesis of carboxylates from CO2. Frontiers in Microbiology. 2021;12:669218.

13. Baas P, Bell C, Mancini LM, Lee MN, Conant RT, Wallenstein MD. Phosphorus mobilizing consortium Mammoth P™ enhances plant growth. PeerJ. 2016;4:e2121.

14. Herrera Paredes S, Gao T, Law TF, Finkel OM, Mucyn T, Teixeira PJPL, et al. Design of synthetic bacterial communities for predictable plant phenotypes. PLoS biology. 2018;16(2):e2003962.

15. Hu J, Wei Z, Friman VP, Gu Sh, Wang Xf, Eisenhauer N, et al. Probiotic diversity enhances rhizosphere microbiome function and plant disease suppression. MBio. 2016;7(6):e01790–16.

16. Vrancken G, Gregory AC, Huys GR, Faust K, Raes J. Synthetic ecology of the human gut microbiota. Nature Reviews Microbiology. 2019;17(12):754–763.

17. Zhang C, Qi J, Cao Y. Synergistic Effect of Yeast-Bacterial Co-Culture on Bioremediation of Oil-Contaminated Soil. Bioremediation Journal. 2014;18. doi:10.1080/10889868.2013.847402.

18. Szulc A, Ambrożewicz D, Sydow M, Ławniczak Ł, Piotrowska-Cyplik A, Marecik R, et al. The influence of bioaugmentation and biosurfactant addition on bioremediation efficiency of diesel-oil contaminated soil: feasibility during field studies. Journal of environmental management. 2014;132:121–128.

19. Piccardi P, Vessman B, Mitri S. Toxicity drives facilitation between 4 bacterial species. Proceedings of the National Academy of Sciences. 2019;116(32):15979–15984.

20. Roell GW, Zha J, Carr RR, Koffas MA, Fong SS, Tang YJ. Engineering microbial consortia by division of labor. Microbial cell factories. 2019;18:1–11.

21. Thommes M, Wang T, Zhao Q, Paschalidis IC, Segrè D. Designing metabolic division of labor in microbial communities. MSystems. 2019;4(2):e00263–18.

22. Diender M, Parera Olm I, Sousa DZ. Synthetic co-cultures: novel avenues for bio-based processes. Current Opinion in Biotechnology. 2021;67:72–79. doi:https://doi.org/10.1016/j.copbio.2021.01.006.

23. Hays SG, Patrick WG, Ziesack M, Oxman N, Silver PA. Better together: engineering and application of microbial symbioses. Current opinion in biotechnology. 2015;36:40–49.

24. Erkus O, De Jager VC, Spus M, van Alen-Boerrigter IJ, Van Rijswijck IM, Hazelwood L, et al. Multifactorial diversity sustains microbial community stability. The ISME journal. 2013;7(11):2126–2136.

25. Sanchez A, Bajic D, Diaz-Colunga J, Skwara A, Vila JC, Kuehn S. The community-function landscape of microbial consortia. EcoEvoRxiv. 2022;.

26. Widder S, Allen RJ, Pfeiffer T, Curtis TP, Wiuf C, Sloan WT, et al. Challenges in microbial ecology: building predictive understanding of community function and dynamics. The ISME journal. 2016;10(11):2557–2568.

27. Burgard AP, Pharkya P, Maranas CD. Optknock: a bilevel programming framework for identifying gene knockout strategies for microbial strain optimization. Biotechnology and bioengineering. 2003;84(6):647–657.

28. Benito-Vaquerizo S, Nouse N, Schaap PJ, Hugenholtz J, Brul S, López-Contreras AM, et al. Model-driven approach for the production of butyrate from CO2/H2 by a novel co-culture of C. autoethanogenum and C. beijerinckii. Frontiers in Microbiology. 2022;13. doi:10.3389/fmicb.2022.1064013.

29. Passi A, Tibocha-Bonilla JD, Kumar M, Tec-Campos D, Zengler K, Zuniga C. Genome-scale metabolic modeling enables in-depth understanding of big data. Metabolites. 2022;12:14. doi:10.3390/METABO12010014.

30. Orth JD, Thiele I, Palsson BØ. What is flux balance analysis? Nature Biotechnology. 2010;28(3):245–248. doi:10.1038/nbt.1614.

31. Mahadevan R, Edwards JS, Doyle FJ. Dynamic Flux Balance Analysis of diauxic growth in Escherichia coli. Biophysical Journal. 2002;83:1331–1340. doi:10.1016/S0006-3495(02)73903-9.

32. Moreno-Paz S, Schmitz J, dos Santos VAPM, Suarez-Diez M. Enzyme-constrained models predict the dynamics of Saccharomyces cerevisiae growth in continuous, batch and fed-batch bioreactors. Microbial Biotechnology. 2022;15. doi:10.1111/1751-7915.13995.

33. Gottstein W, Olivier BG, Bruggeman FJ, Teusink B. Constraint-based stoichiometric modelling from single organisms to microbial communities. Journal of the Royal Society Interface. 2016;13(124):20160627.

34. Harcombe WR, Riehl WJ, Dukovski I, Granger BR, Betts A, Lang AH, et al. Metabolic resource allocation in individual microbes determines ecosystem interactions and spatial dynamics. Cell reports. 2014;7:1104. doi:10.1016/J.CELREP.2014.03.070.

35. Chen J, Gomez JA, Höffner K, Phalak P, Barton PI, Henson MA. Spatiotemporal modeling of microbial metabolism. BMC Systems Biology. 2016;doi:10.1186/s12918-016-0259-2.

36. Dukovski I, Bajić D, Chacón JM, Quintin M, Vila JCC, Sulheim S, et al. A metabolic modeling platform for the computation of microbial ecosystems in time and space (COMETS). Nature Protocols. 2021;16. doi:10.1038/s41596-021-00593-3.

37. Biggs MB, Papin JA. Novel Multiscale Modeling Tool Applied to Pseudomonas aeruginosa Biofilm Formation. PLOS ONE. 2013;8(10):null. doi:10.1371/journal.pone.0078011.

38. Bauer E, Zimmermann J, Baldini F, Thiele I, Kaleta C. BacArena: Individual-based metabolic modeling of heterogeneous microbes in complex communities. PLOS Computational Biology. 2017;13:e1005544. doi:10.1371/JOURNAL.PCBI.1005544.

39. Borer B, Ataman M, Hatzimanikatis V, Or D. Modeling metabolic networks of individual bacterial agents in heterogeneous and dynamic soil habitats (IndiMeSH). PLOS Computational Biology. 2019;15(6):1–21. doi:10.1371/journal.pcbi.1007127.

40. Heinken A, Basile A, Thiele I. Advances in constraint-based modelling of microbial communities. Current Opinion in Systems Biology. 2021;27:100346. doi:10.1016/J.COISB.2021.05.007.

41. Colarusso AV, Goodchild-Michelman I, Rayle M, Zomorrodi AR. Computational modeling of metabolism in microbial communities on a genome-scale. Current Opinion in Systems Biology. 2021;26:46–57. doi:10.1016/J.COISB.2021.04.001.

42. Ang KS, Lakshmanan M, Lee NR, Lee DY. Metabolic Modeling of Microbial Community Interactions for Health, Environmental and Biotechnological Applications. Current Genomics. 2018;doi:10.2174/1389202919666180911144055.

43. Shoaie S, Ghaffari P, Kovatcheva-Datchary P, Mardinoglu A, Sen P, Pujos-Guillot E, et al. Quantifying Diet-Induced Metabolic Changes of the Human Gut Microbiome. Cell metabolism. 2015;22:320–331. doi:10.1016/J.CMET.2015.07.001.

44. Gottstein W, Olivier BG, Bruggeman FJ, Teusink B. Constraint-based stoichiometric modelling from single organisms to microbial communities. Journal of The Royal Society Interface. 2016;13. doi:10.1098/RSIF.2016.0627.

45. Diener C, Gibbons SM, Resendis-Antonio O. MICOM: Metagenome-Scale Modeling To Infer Metabolic Interactions in the Gut Microbiota. mSystems. 2020;5. doi:10.1128/msystems.00606-19.

46. Baldini F, Heinken A, Heirendt L, Magnusdottir S, Fleming RMT, Thiele I. The Microbiome Modeling Toolbox: from microbial interactions to personalized microbial communities. Bioinformatics. 2018;35(13):2332–2334. doi:10.1093/bioinformatics/bty941.

47. Heinken A, Thiele I. Microbiome Modelling Toolbox 2.0: efficient, tractable modelling of microbiome communities. Bioinformatics. 2022;38(8):2367–2368. doi:10.1093/bioinformatics/btac082.

48. Mendoza SN, Olivier BG, Molenaar D, Teusink B. A systematic assessment of current genome-scale metabolic reconstruction tools. Genome Biology. 2019;20:1–20. doi:10.1186/S13059-019-1769-1/FIGURES/7.

49. Wilkinson MD, Dumontier M, Aalbersberg IJ, Appleton G, Axton M, Baak A, et al. The FAIR Guiding Principles for scientific data management and stewardship. Scientific Data 2016 3:1. 2016;3:1–9. doi:10.1038/sdata.2016.18.

50. Katz DS, Gruenpeter M, Honeyman T. Taking a fresh look at FAIR for research software. Patterns. 2021;2(3):100222. doi:https://doi.org/10.1016/j.patter.2021.100222.

51. Barker M, Chue Hong NP, Katz DS, Lamprecht AL, Martinez-Ortiz C, Psomopoulos F, et al. Introducing the FAIR Principles for research software. Scientific Data 2022 9:1. 2022;9(1):1–6. doi:10.1038/s41597-022-01710-x.

52. Bengtsson-Palme J. Microbial model communities: To understand complexity, harness the power of simplicity. Computational and Structural Biotechnology Journal. 2020;18:3987–4001. doi:10.1016/j.csbj.2020.11.043.

53. García-Jiménez B, Torres-Bacete J, Nogales J. Metabolic modelling approaches for describing and engineering microbial communities. Computational and Structural Biotechnology Journal. 2021;19:226. doi:10.1016/J.CSBJ.2020.12.003.

54. Eng A, Borenstein E. Microbial community design: methods, applications, and opportunities. Current Opinion in Biotechnology. 2019;58:117–128. doi:10.1016/j.copbio.2019.03.002.

55. Aguirre de Cárcer D. Experimental and computational approaches to unravel microbial community assembly. Computational and Structural Biotechnology Journal. 2020;18:4071–4081. doi:https://doi.org/10.1016/j.csbj.2020.11.031.

56. Stolyar S, Van Dien S, Hillesland KL, Pinel N, Lie TJ, Leigh JA, et al. Metabolic modeling of a mutualistic microbial community. Molecular Systems Biology. 2007;3(1):92. doi:https://doi.org/10.1038/msb4100131.

57. Zomorrodi AR, Maranas CD. OptCom: A multi-level optimization framework for the metabolic modeling and analysis of microbial communities. PLoS Computational Biology. 2012;doi:10.1371/journal.pcbi.1002363.

58. Khandelwal RA, Olivier BG, Röling WFM, Teusink B, Bruggeman FJ. Community Flux Balance Analysis for Microbial Consortia at Balanced Growth. PLoS ONE. 2013;doi:10.1371/journal.pone.0064567.

59. Hung S, Chan J, Simons MN, Maranas CD. SteadyCom: Predicting microbial abundances while ensuring community stability. PLOS Computational Biology. 2017;13:e1005539. doi:10.1371/JOURNAL.PCBI.1005539.

60. Koch S, Kohrs F, Lahmann P, Bissinger T, Wendschuh S, Benndorf D, et al. Redcom: A strategy for reduced metabolic modeling of complex microbial communities and its application for analyzing experimental datasets from anaerobic digestion. PLoS Computational Biology. 2019;doi:10.1371/journal.pcbi.1006759.

61. Cai J, Tan T, Chan SHJ. Predicting Nash equilibria for microbial metabolic interactions. Bioinformatics. 2021;36:5649–5655. doi:10.1093/BIOINFORMATICS/BTAA1014.

62. Zhuang K, Izallalen M, Mouser P, Richter H, Risso C, Mahadevan R, et al. Genome-scale dynamic modeling of the competition between Rhodoferax and Geobacter in anoxic subsurface environments. The ISME journal. 2011;5:305–316. doi:10.1038/ISMEJ.2010.117.

63. Gomez JA, Höffner K, Barton PI. DFBAlab: A fast and reliable MATLAB code for dynamic flux balance analysis. BMC Bioinformatics. 2014;doi:10.1186/s12859-014-0409-8.

64. Zomorrodi AR, Islam MM, Maranas CD. D-OptCom: Dynamic Multi-level and Multi-objective Metabolic Modeling of Microbial Communities. ACS Synthetic Biology. 2014;doi:10.1021/sb4001307.

65. Louca S, Doebeli M. Calibration and analysis of genome-based models for microbial ecology. eLife. 2015;4. doi:10.7554/ELIFE.08208.

66. Succurro A, Segrè D, Ebenhöh O. Emergent Subpopulation Behavior Uncovered with a Community Dynamic Metabolic Model of Escherichia coli Diauxic Growth. mSystems. 2019;4. doi:10.1128/msystems.00230-18.

67. Popp D, Centler F. μBialSim: Constraint-Based Dynamic Simulation of Complex Microbiomes. Frontiers in Bioengineering and Biotechnology. 2020;8:574. doi:10.3389/FBIOE.2020.00574/BIBTEX.

68. García-Jiménez B, Carrasco J, Medina J, Wilkinson MD. Dynamic simulations of microbial communities under perturbations: opportunities for microbiome engineering. Research Square Preprints. 2020;doi:10.21203/rs.2.24431/v1.

69. Brunner JD, Chia N. Minimizing the number of optimizations for efficient community dynamic flux balance analysis. PLOS Computational Biology. 2020;16(9):1–20. doi:10.1371/journal.pcbi.1007786.

70. Gardner JJ, Hodge BMS, Boyle NR. Multiscale Multiobjective Systems Analysis (MiMoSA): an advanced metabolic modeling framework for complex systems. Scientific Reports. 2019;9. doi:10.1038/s41598-019-53188-0.

71. Karimian E, Motamedian E. ACBM: An Integrated Agent and Constraint Based Modeling Framework for Simulation of Microbial Communities. Scientific Reports. 2020;10. doi:10.1038/s41598-020-65659-w.

72. Angeles-Martinez L, Hatzimanikatis V. Spatio-temporal modeling of the crowding conditions and metabolic variability in microbial communities. PLoS Computational Biology. 2021;17. doi:10.1371/journal.pcbi.1009140.

73. Diender M, Olm IP, Sousa DZ. Synthetic co-cultures: novel avenues for bio-based processes. Current Opinion in Biotechnology. 2021;67:72–79. doi:10.1016/J.COPBIO.2021.01.006.

74. Hanly TJ, Henson MA. Dynamic flux balance modeling of microbial co-cultures for efficient batch fermentation of glucose and xylose mixtures. Biotechnology and bioengineering. 2011;108(2):376–385.

75. Vlassis N, Pacheco MP, Sauter T. Fast Reconstruction of Compact Context-Specific Metabolic Network Models. PLOS Computational Biology. 2014;10(1):1–10. doi:10.1371/journal.pcbi.1003424.

76. Olivier B, Gottstein W, Molenaar D, Teusink B. CBMPy release 0.8.2; 2021. Available from: https://doi.org/10.5281/zenodo.5546608.

77. Ebrahim A, Lerman JA, Palsson BO, Hyduke DR. COBRApy: COnstraints-Based Reconstruction and Analysis for Python. BMC Systems Biology. 2013;7:74. doi:10.1186/1752-0509-7-74.

78. Klamt S, Saez-Rodriguez J, Gilles ED. Structural and functional analysis of cellular networks with CellNetAnalyzer. BMC systems biology. 2007;1(1):1–13.

79. Wang H, Marcišauskas S, Sánchez BJ, Domenzain I, Hermansson D, Agren R, et al. RAVEN 2.0: A versatile toolbox for metabolic network reconstruction and a case study on Streptomyces coelicolor. PLOS Computational Biology. 2018;14(10):1–17. doi:10.1371/journal.pcbi.1006541.

80. Benito-Vaquerizo S, Diender M, Parera Olm I, Martins dos Santos VAP, Schaap PJ, Sousa DZ, et al. Modeling a co-culture of *Clostridium autoethanogenum* and *Clostridium kluyveri* to increase syngas conversion to medium-chain fatty-acids. Computational and Structural Biotechnology Journal. 2020;doi:10.1016/j.csbj.2020.10.003.

81. Palsson B. Systems biology. Cambridge university press; 2015.

82. Keating SM, Waltemath D, König M, Zhang F, Dräger A, Chaouiya C, et al. SBML Level 3: an extensible format for the exchange and reuse of biological models. Molecular systems biology. 2020;16(8):e9110.

83. Heinken A, Acharya G, Ravcheev DA, Hertel J, Nyga M, Okpala OE, et al. AGORA2: Large scale reconstruction of the microbiome highlights wide-spread drug-metabolising capacities. bioRxiv. 2020;doi:10.1101/2020.11.09.375451.

84. Diender M, Stams AJM, Sousa DZ. Production of medium-chain fatty acids and higher alcohols by a synthetic co-culture grown on carbon monoxide or syngas. Biotechnol Biofuels. 2016;9:1–11. doi:10.1186/S13068-016-0495-0.

85. Charubin K, Papoutsakis ET. Direct cell-to-cell exchange of matter in a synthetic Clostridium syntrophy enables CO2 fixation, superior metabolite yields, and an expanded metabolic space. Metabolic Engineering. 2019;52:9–19. doi:10.1016/J.YMBEN.2018.10.006.

86. Moreira JPC, Diender M, Arantes AL, Boeren S, Stams AJM, Alves MM, et al. Propionate production from carbon monoxide by synthetic cocultures of *Acetobacterium wieringae* and propionigenic bacteria. Appl Environ Microbiol. 2021;87. doi:10.1128/AEM.02839-20.

87. Bäumler M, Schneider M, Ehrenreich A, Liebl W, Weuster-Botz D. Synthetic co-culture of autotrophic Clostridium carboxidivorans and chain elongating Clostridium kluyveri monitored by flow cytometry. Microbial Biotechnology. 2022;15. doi:10.1111/1751-7915.13941.

88. Lieven C, Beber ME, Olivier BG, Bergmann FT, Ataman M, Babaei P, et al. MEMOTE for standardized genome-scale metabolic model testing. Nature Biotechnology 2020 38:3. 2020;38(3):272–276. doi:10.1038/s41587-020-0446-y.

89. Lachance JC, Lloyd CJ, Monk JM, Yang L, Sastry AV, Seif Y, et al. BOFdat: Generating biomass objective functions for genome-scale metabolic models from experimental data. PLoS computational biology. 2019;15(4):e1006971.

90. Schuetz R, Kuepfer L, Sauer U. Systematic evaluation of objective functions for predicting intracellular fluxes in Escherichia coli. Molecular systems biology. 2007;3(1):119.

91. Herrmann HA, Dyson BC, Vass L, Johnson GN, Schwartz JM. Flux sampling is a powerful tool to study metabolism under changing environmental conditions. npj Systems Biology and Applications. 2019;5. doi:10.1038/s41540-019-0109-0.

92. Scott WT, Smid EJ, Block DE, Notebaart RA. Metabolic flux sampling predicts strain-dependent differences related to aroma production among commercial wine yeasts. Microbial cell factories. 2021;20(1):1–15.

93. Wiback SJ, Famili I, Greenberg HJ, Palsson BØ. Monte Carlo sampling can be used to determine the size and shape of the steady-state flux space. Journal of theoretical biology. 2004;228(4):437–447.

94. Zomorrodi AR, Segrè D. Synthetic ecology of microbes: mathematical models and applications. Journal of molecular biology. 2016;428(5):837–861.

95. Valgepea K, Loi KQ, Behrendorff JB, Lemgruber RdSP, Plan M, Hodson MP, et al. Arginine deiminase pathway provides ATP and boosts growth of the gas-fermenting acetogen Clostridium autoethanogenum. Metabolic Engineering. 2017;41:202–211. doi:10.1016/J.YMBEN.2017.04.007.

96. Zou W, Ye G, Zhang J, Zhao C, Zhao X, Zhang K. Genome-scale metabolic reconstruction and analysis for *Clostridium kluyveri*. Genome. 2018;61(8):605–613. doi:10.1139/gen-2017-0177.

97. Duarte NC, Herrgård MJ, Palsson BØ. Reconstruction and validation of Saccharomyces cerevisiae iND750, a fully compartmentalized genome-scale metabolic model. Genome research. 2004;14(7):1298–1309.

98. Reed JL, Vo TD, Schilling CH, Palsson BO. An expanded genome-scale model of Escherichia coli K-12 (iJR904 GSM/GPR). Genome biology. 2003;4(9):1–12.

99. King ZA, Lu J, Dräger A, Miller P, Federowicz S, Lerman JA, et al. BiGG Models: A platform for integrating, standardizing and sharing genome-scale models. Nucleic acids research. 2016;44(D1):D515–D522.

100. Montgomery DC. Design and analysis of experiments. John wiley & sons; 2017.

101. Heirendt L, Arreckx S, Pfau T, Mendoza SN, Richelle A, Heinken A, et al. Creation and analysis of biochemical constraint-based models using the COBRA Toolbox v. 3.0. Nature protocols. 2019;14(3):639–702.

102. Orth JD, Conrad TM, Na J, Lerman JA, Nam H, Feist AM, et al. A comprehensive genome-scale reconstruction of Escherichia coli metabolism—2011. Molecular systems biology. 2011;7(1):535.

103. Raghunathan A, Reed J, Shin S, Palsson B, Daefler S. Constraint-based analysis of metabolic capacity of Salmonella typhimurium during host-pathogen interaction. BMC systems biology. 2009;3(1):1–16.