1 **The origins and diversification of Holarctic brown bear populations inferred from**
2 **genomes of past and present populations**
3

4 **Keywords:** Brown bear, Demography, Evolution, Genome, Palaeogenome, Phylogenomics,
5

6 Takahiro Segawa[1], Alba Rey-Iglesia[2], Eline D Lorenzen[2], Michael V Westbury[2,*]
7
8 1. Center for Life Science Research, University of Yamanashi, Japan
9 2. Globe Institute, University of Copenhagen, Denmark
10
11 *Correspondence: m.westbury@sund.ku.dk

12
13
14 **Abstract**
15

The brown bear (*Ursus arctos*) is one of the survivors of the Late Quaternary
megafauna extinctions. However, despite being widely distributed across the Holarctic,
brown bears have experienced extensive range reductions, and even extirpations in some
geographic regions. Previous research efforts utilising genetic data have provided valuable
insights into their evolutionary history. However, most studies have been limited to
contemporary individuals or mitochondrial DNA, limiting insights into population processes
that preceded the present. Here we present genomic data from two Late Pleistocene brown
bears from Honshu, Japan, and eastern Siberia, and combine them with published
contemporary and ancient genomes from across the Holarctic range of brown bears to
investigate the evolutionary relationships among brown bear populations through time and
space. By including genomic data from Late Pleistocene and Holocene individuals sampled
outside the current distribution range we uncover diversity not present in the contemporary
populations. Notably, although contemporary individuals display geographically structured
populations most likely driven by isolation-by-distance, this pattern varies among the ancient
samples across different regions. The inclusion of ancient brown bears in our analysis
provides novel insights into the evolutionary history of brown bears and contributes to
understanding the populations and diversity lost during the Late Quaternary.

33
34 **Introduction**
35
The Pleistocene-Holocene transition was a period of major climatic upheaval,
culminating with a large number of megafaunal extinctions as a defining feature across the
globe [1]. However, despite this, many megafauna species still persist today, although they
may have experienced local extirpations, range reductions, and losses of genetic diversity
[2,3]. By comparing the genetic composition of populations inhabiting recently recolonised
regions to that of populations located in past refugia, genetic information from contemporary
individuals can be used to study the patterns of postglacial recolonisation after the Last
Glacial Maximum [4]. However, purely focussing on contemporary individuals means results
are restricted to surviving lineages, limiting our understanding of the population processes

that preceded the present. Genetic information from subfossil individuals (ancient DNA -
aDNA) enables the tracking of change in real time, the uncovering of potentially lost
lineages, and an overall better understanding of past responses to change [5].
48
The brown bear (*Ursus arctos*) is one of the survivors of the Late Pleistocene
megafauna extinctions. The species is one of the largest extant terrestrial carnivores, and is
distributed across the Holarctic [6]. The ability of brown bears to survive may be due to their
ecological flexibility and broad dietary range, observed in contemporary populations [7].
Brown bears have evolved a generalist omnivore strategy; although they possess
morphological traits characteristic of carnivores, their diets in many ecosystems comprise
primarily of plant matter [8]. Despite their current dietary flexibility, brown bears
experienced extensive range reductions during the Holocene, with extirpations in some
geographic regions driven by climate and environmental change, and more recently, humans
[9]. For example, based on fossil chronology, the relatively large steppe brown bear (*Ursus
arctos priscus*) of Eurasia disappeared ~25 kya. Although it was presumably well-adapted for
survival in Europe during the Late Pleistocene, its size may have inhibited adaptive flexibility
after the environment drastically changed at the onset of the Holocene [10].
62
Since its first utilisation in brown bears in 1994 [11], genetics has provided many
insights into the contemporary phylogeographic structuring of the species. However, the
majority of mitochondrial genetic clades used to form biogeographic hypotheses are
paraphyletic, and not unique to a single geographic region [12], making global inferences
difficult. A recent study using nuclear genomes from across the contemporary Holarctic
distribution showed phylogeographic patterns at the nuclear level do not mirror the
mitochondria, but instead found geographically structured populations following isolation by
distance. They suggested that current population structure may represent recent population
dynamics [13], as more ancient signals may have been broken down by gene flow and
recombination.
73
Ancient DNA from brown bears has provided valuable insights into their evolutionary
history, and provided valuable information regarding temporal changes in genetic diversity
and migration patterns [14–17]. However, studies of ancient brown bear specimens have thus
far been based on mitochondrial DNA, which represents just a single loci, is exclusively
maternally inherited, and may be heavily confounded by sex-biased migrations, gene flow,
and incomplete lineage sorting [13,18,19]. Nuclear data provides an opportunity to
circumvent the challenges associated with mitochondrial genomes [18]. Although ancient
nuclear genomic data is available for several brown bears, it has thus far only been used to
understand the evolutionary relationships between different Ursine bear species [20,21], not
to understand brown bear intraspecific relationships through time.
84
85   Here, we utilise a combination of contemporary and ancient nuclear genomes,
86 incorporating newly generated palaeogenomic data from two Late Pleistocene individuals
87 from Japan (0.11x; 32,464 ± 255 cal. BP [14]) and eastern Siberia (0.35x; molecularly dated

88 to 61,826 years BP [HPD 95%: 27,866–96,783 years BP][22]), to investigate the evolutionary
89 relationships among brown bear populations through time and space.
90
91 **Methods**
92
Data generation
We generated palaeogenomic data from a Late Pleistocene (32,464 ± 255 cal. BP
[14]) brown bear from the Japanese island of Honshu (JBB-32K [14]). Lab work was carried
out in a clean room for ancient DNA at the National Institute of Polar Research, Japan. We
extracted 100 mg of petrosal bone, and followed the DNA extraction procedure based on
silica pellets in solution as described in Orlando et al. [23]. Double-stranded DNA was
converted into an Illumina sequencing library using an NEBNext Quick DNA Library Prep
Master Mix Set for 454 (New England BioLabs) with 20 µl of DNA extract without DNA
fragmentation. Illumina sequencing adapters were added to the end-repair reaction. Indexing
PCR was prepared in the ancient DNA laboratory, carried out in a physically separate modern
laboratory. Each eluate (3 µl) was subjected to PCR amplification in 50-µl reactions
containing 1 KAPA HiFi HotStart Uracil+ReadyMix (Kapa Biosystems) and 0.3 µM Dual
Index primers of NEBNext Multiplex Oligos for Illumina (NEB). Three independent PCR
reactions from the same library were combined after PCR and purified using a NucleoSpin
Gel and PCR Clean-up kit (Macherey-Nagel). Sequencing was carried out on an Illumina
HiSeq XTen at Macrogen, Japan. Previously published data for this individual was
downloaded from DDBJ DRA IDs: DRR308533 and DRR308534 [14].
110
We generated palaeogenomic data from a Late Pleistocene (aged beyond radiocarbon
dating: >50 kya but molecularly dated to 61,826 years BP [HPD 95%: 27,866–96,783 years
BP][22]) brown bear from eastern Siberia (CGG_1_020006 [22]). Lab work was carried out
in a clean room for ancient DNA at Globe Institute, University of Copenhagen. We extracted
DNA from ~50 mg of bone powder using a modified version of the protocol from [24].
Modifications included overnight incubation with the extraction buffer at 42 °C instead of at
37 °C, the bone powder was pelleted out of suspension, and the supernatant concentrated
down to 150–200 µl for each sample using 30 kDa Amicon centrifugal filter unit
(Millipore). The remaining supernatant was mixed with 13x its volume of binding buffer and
DNA was purified with MinElute columns (Qiagen), following the manufacturer's
instructions with the exception of a 15-minute incubation at 37 °C during the elution step.
DNA extracts were transformed into sequencing libraries in 25 µl reactions following the
protocol from Meyer and Kircher [25] with the modification that the initial DNA
fragmentation was not performed, as ancient DNA is already highly fragmented, and
MinElute kit (Qiagen) was used for the purification steps. DNA libraries were indexed using
KAPA HiFi uracil + premix (KAPA Biosystems). The number of cycles for index PCRs
was determined from qPCR analysis. The resulting libraries were quantified on an Agilent
2100 Bioanalyzer, pooled at equimolar concentration and sequenced on an Illumina HiSeq
2500 at the University of Copenhagen using single-end 80 base pairs (bp) reads.
130

We additionally downloaded the raw reads from 137 contemporary and eight ancient
brown bear individuals, from across their present and past distributions [13,20,21,26–31] (Fig
1A and Supplementary Table 1). Seven of the ancient samples are from outside the current
distribution of brown bears, and include Ireland (n = 5), Honshu (Japan, n = 1), and Quebec
(Canada, n = 1).

Data processing and mapping

Contemporary brown bears
We processed the raw sequencing data from 35 contemporary brown bears (excluding
those from Endo et al and de Jong et al, detailed below) with the PALEOMIX v.1.2.12 [32]
pipeline. Adapters, stretches of Ns, and low-quality bases in reads were trimmed and filtered
with AdapterRemovalv2 [33] using default parameters. BWA-backtrack v0.7.15 [34] was
used to map the cleaned reads to the polar bear pseudo-chromosome genome [35,36] (based
on Genbank accession: GCA_000687225.1), which includes a mitochondrial genome
(Genbank accession: NC003428), with default parameters. Reads with mapping quality < 30
were filtered using SAMtools [37]. Duplicates were removed with Picard v2.6.0 [38].
Possible paralogs were filtered using SAMtools v1.6. Finally, local realignment around indels
was done using GATK (v 3.3) [39].
The individuals from Endo et al (n=6) and de Jong et al (n=96) followed a similar
protocol, but using Fastp v0.23.2 [40] for adapter trimming and duplicate removal using
SAMtools. To be used as an outgroup/ancestral sequence, the raw data for the spectacled bear
(*Tremarctos ornatus*) was downloaded (SRA accession code: SRR13602719) and mapped to
the same polar bear reference genome as above using the BWA mem algorithm.

Ancient brown bears
We removed illumina adapter sequences, low-quality reads (mean q <25), short reads
(<30bp), and merged overlapping read pairs with Fastp. We mapped the merged reads to the
polar bear reference genome using BWA-backtrack v0.7.15, with the seed disabled (-l 999),
and otherwise default parameters. Reads with mapping quality < 30 and duplicates were
filtered using SAMtools [37].

Population structure
We evaluated the relationships between the contemporary and ancient brown bears
included in this study by performing Principal Component Analyses (PCA). We performed
these using either pseudohaploid base calls or genotype likelihoods computed with
ANGSDv0.921 [41]. We computed genotype likelihoods in ANGSD for all individuals
specifying the parameters: minimum mapping and base qualities of 30 (-minmapQ 30 -minQ
30), calculate genotype likelihoods using the GATK algorithm (-GL 2), output a beagle
genotype likelihood file (-doGlf 2), calculate major and minor alleles based on genotype
likelihoods (-doMajorMinor 1), remove transitions (-rmtrans 1), only include SNPs with a p-
values <1e-6 (-SNP_pval 1e-6), only consider autosomal chromosomes (-rf), a minimum
minor allele frequency of 0.05 (-minmaf 0.05), skip triallelic sites (-skiptriallelic 1), only
consider reads mapping to one region uniquely (-uniqueonly 1), and only consider sites if at

least 132 individuals have coverage (-minind 132). We only included pseudochromosomes 1-
18 as previously recommended [35]. The pseudohaploid base call was computed using the
same parameters with the addition of -doIBS 2, -doCov 1, and only included SNPs if at least
two individuals had the minor allele (-minminor 2). To construct covariance matrices from
the genotype likelihoods datasets we used PCAngsd v0.98 [42]. We only included a single
Apennines mountain (Italy) brown bear in these analyses due to their high levels of
relatedness [29]. We repeated these steps with only individuals from Eurasia using the same
parameters but set the minimum individual parameter (MinInd) to 79. Due to its suitability
for high levels of missing data, we additionally performed principal component analyses for
the two datasets using EMU v0.9 [43]. As input, we generated consensus pseudohaploid base
calls in ANGSD using the same parameters as above but with the addition of the -dohaplocall
2. We converted the haploid calls into PLINK format using haploToPlink, which is part of
the ANGSD toolkit. We ran the PLINK file in EMU specifying to produce eigenvalues for
the first three principal components, and otherwise default parameters.
189
To investigate whether the low coverage, and ancient DNA damage may influence the
ancient individual's placement within a PCA, we simulated ancient DNA damage patterns,
short read lengths, and low coverage onto three contemporary individuals. The three
individuals were S235 (Central Asia) and Kirkenes (Europe), and DRR276778 (Japan). We
used Fastp to reduce the raw reads to 40bp then added ancient DNA patterns to the raw reads
of these three individuals using TAPAS v1.2 [44]. We mapped the artificially damaged and
short reads to the reference genome following the approaches specified above and we used
SAMtools to subsample the individuals. Ancient DNA damage patterns were taken from the
Sligo-14 mapdamage output for S235 and Kirkenes, and JBB-32K for DRR276778. We
downsampled S235 and Kirkenes to 0.1x and JBB-32K to 0.05x. We reran the genotype
likelihoods and pseudohaploid PCA with these three individuals replacing the original high-
quality data with the artificially damaged bam files.
202
Pairwise distances
To further estimate the relationships between ancient and contemporary Eurasian
individuals, we calculated pairwise distances between all Eurasian individuals (excluding
JBB-32K due to its clear placement within the Hokkaido brown bears in the PCA) using the
pseudohaploid baseball (-doIBS 2) mentioned above and converted these into a distance
matrix using the -makematrix 1 parameter in ANGSD. We extracted the columns
corresponding to the ancient individuals and rows corresponding to contemporary
individuals.
211
Ancestry proportions
We calculated the ancestry proportions of the eight ancient Eurasian individuals
(excluding JBB-32K) using admixfrog v0.7.2 [45]. For this analysis we considered three
ancestry states each represented by a single individual; East Europe - represented by
Kirkenes, Himalayas - represented by Himalaya1, and Central Asia - represented by S235. As
input for the reference panel we created a multi-individual variant call file with the three
individuals using BCFtools v1.15 [46], specifying chromosomes 1-18 (-f), minimum

mapping and base qualities of 20 (-q 20 -Q 20). We ran admixfrog specifying a minimum
read length of 30bp, a bin size of 50kb, and otherwise default parameters. When interpreting
the results, if a window was returned as having combined ancestry between any of the two
states, then we deemed this as ancestral.
223
D-statistics topology test
We used D-statistics to investigate whether our ancient Eurasian individuals were
more closely related to contemporary European or Asian individuals. D-statistics works by
calculating shared derived alleles between non-sister branches of a given topology [[[H1,
H2], H3], Outgroup]. An allele is considered as derived if it is different to the outgroup allele.
Although it is most commonly implemented to investigate differential levels of gene flow
between non-sister taxon, it can also be used to investigate shared polymorphisms and
population structure [47]. We subsampled the dataset to 42 Eurasian individuals
(Supplementary table S2). We calculated the D-statistics using a random base call approach
in ANGSD (-doabbababa 1), specifying only pseudochromosomes 1-18, the spectacled bear
as outgroup, and the following parameters: -minmapQ 30 -minQ 30 -blocksize 1000000 -
rmtrans 1 -uniqueonly 1. We summarised the results using a block jackknife approach with
the Rscript available in the ANGSD toolsuite (ANGSD_jackknife.R). As ANGSD computes
every possible ingroup triplet combination of D-statistics, we filtered out only the
combinations of interest. We filtered the output to only include the contemporary European
individuals in the H1 position, the contemporary Asian individuals in H2, and the ancient
individuals in the H3 position [[[Europe, Asia], Ancient], Outgroup]. This filtering allows us
to see if the ancient individual has more shared derived alleles with either the contemporary
European or contemporary Asian individual.
243
244 **Results**
245
Population structure
Regardless of the method used (genotype likelihood, pseudohaploid, EMU) we found
similar results in our PCAs, with clear differences between bears from different geographic
localities in a west-to-east cline (Fig 1B,C, Supplementary Fig S1). Further investigations
into Eurasia showed a very strong correlation between PC1 and longitude (R^2=0.81, Fig 1D).
The ~32kya Honshu Japanese individual (JBB-32K) clustered most closely with the
contemporary Hokkaido Japanese bears. The ~11kya Quebec sample clustered with the North
American bears, but not with any specific locality in the genotype likelihoods or
pseudohaploid data. However, it did cluster with the easternmost bears in the EMU-computed
PCA.
The remaining eight ancient individuals did not cluster where expected based on
geography. The Holocene Irish individuals (Leitrim-4, and Leitrim-5) clustered close to the
easternmost European individuals, whereas the Late Pleistocene Irish (Clare-12, Sligo-13,
Sligo-14), Austrian (Austria_LP), and Siberian (Denisova_LP, CGG_1_020006) cluster
closest to the Himalayan individual (seen as dark grey in Fig 1A) but in general do not cluster
within any contemporary individual.

Investigations into the role of ancient DNA damage in the placement of individuals
into the PCA showed that simulated ancient individuals clustered in the same location as they
did when using the high-quality version of the data, and that using genotype likelihoods may
be the most robust to aDNA damage (Supplementary Fig S2). This suggests that our
approaches are robust to aDNA damage and the results are biological rather than driven by
data bias.
When visualising PC3 in the Eurasian dataset, the differentiation between
contemporary European individuals became more apparent, yet retained the structuring
observed in PC1.
271
Pairwise distances
Overall, the Irish bears generally show the lowest pairwise distances (PWD) to
individuals from Europe, but this heavily overlaps with some Asian bears (Fig 2A).
Austria_LP and Denisova_LP show the smallest PWD to the Himalayan individual.
However, when excluding the Himalayas bear, Austria_LP is more or less similar to all bears,
and Denisova_LP is closer to Asian bears than European. PWD heavily overlapped between
CGG_1_020006 and Europe, Asia, and the Himalayas.
279
Ancestry proportions
We found relatively high proportions of most of the ancient Irish individual's ancestry
(to the exclusion of Sligo-14) to be eastern European, followed by central Asia, then
Himalayan (Fig 2B). There was slightly more European ancestry in the Holocene Irish
individuals (Leitrim-4 0.297 and Leitrim-5 0.318) relative to their Late Pleistocene
counterparts (Clare-12 0.246, Sligo-13 0.27, Sligo-14 0.046). The lowest level of ancestry in
all ancient individuals - when considering only the three single reference populations - was
Himalayan (<0.09). Large proportions of ancestry in all ancient individuals could not be
assigned to a single population, especially in Sligo-14 (0.91) and Denisova_LP (0.837), so
was denoted as ancestral. Austria_LP had similar values of eastern European and central
Asian ancestry (0.165 and 0.168 respectively), whereas CGG_1_020006 had slightly more
central Asian (0.205) relative to eastern European or Himalayan (0.149 and 0.059
respectively).
293
D-statistics topology test
All ancient Irish individuals, regardless of age, show similar D-statistics results (Fig
3). However, there is a slight shift towards more negative D-scores in the Holocene
individuals relative to the LP individuals, indicating more European ancestry in the Holocene
individuals. The mean D-scores of the Late Pleistocene individuals was -0.050, -0.056, -0.052
for Clare-12, Sligo-13, and Sligo-14 respectively and the mean D-scores of the Holocene
individuals was -0.077, -0.076 for Leitrim-4 and Leitrim-5, respectively. The ancient Asian
individuals, CGG_1_020006 and JBB-32K, showed more positive values when including
Asian individuals in the H2 position, suggesting a closer relationship to Asia than to Europe
(Fig 3). In contrast, Denisova_LP showed a clustering of values around zero, suggesting no
obvious differential affinity to either contemporary European or Asian individuals. However,
the mean D-score of 0.01 may indicate a slightly higher affinity to contemporary Asian

individuals. The ancient Austria_LP individual also showed similar D-scores between H2
individuals, with a clustering of values close to zero, but an overall tendency towards
negative D-scores (mean D-score -0.01) suggesting more European ancestry (Fig 3).
309
310   **Discussion**
311
Through the inclusion of genomic data from Late Pleistocene and Holocene brown
bear individuals sampled outside the current distribution range of the species (Fig 1A), we
uncover ancient diversity not present in the available contemporary data. In general, the
ancient individuals do not follow the clear phylogeographic subdivision found in
contemporary brown bear individuals, with geographically structured populations most likely
driven by isolation-by-distance (Fig 1). Therefore, although brown bears successfully made it
through the Late Quaternary megafauna extinction event, populations and diversity were lost
along with large areas of their past distribution.
320
A pattern of population continuity since the Late Pleistocene is not obvious in Europe,
where the ancient European individuals were widely distributed within our PCA analysis (Fig
1B,C) and not clustering nearest to their closest contemporary geographic counterparts. Most
notably, the ancient European (Irish and Austrian) individuals did not conform with the
hypothesis of the isolation-by-distance, derived from patterns of diversity and substructuring
in contemporary brown bear populations (Fig 1D). Our findings suggest brown bears
previously comprised disjunct populations. The two Holocene Irish individuals (aged 5-3.8
ky BP) appear to be part of the European cluster; they were both closer to European bears in
the PCA relative to their Late Pleistocene counterparts (aged 13.7-12.2 ky BP). However, the
low levels of data and filtering criteria may make individuals appear more closely related to
their closest cluster than they are. When considering the pairwise distance, ancestry, and D-
statistics analyses, the Holocene individuals only have slightly more European ancestry. The
phylogeographic structure of contemporary brown bears in Europe is considered consistent
with an "expansion/contraction" model of postglacial recolonisation, where populations
contracted into southern Europe during glacial periods and recolonised the rest of Europe
after the glacial period [12]. Therefore, the increased contemporary European ancestry in the
two Holocene Irish individuals may be due to genetic exchange with the encroaching
ancestors of the contemporary European populations. Lost diversity is in line with previous
findings of higher levels of mitochondrial diversity in Late Pleistocene European brown bears
relative to contemporary European bears [15].
341
In Late Pleistocene Europe, brown bears had higher δ^{15}N values, indicating more
carnivorous diets [22]. However, lower δ^{15}N values towards the end of the Late Pleistocene
and into the Holocene have been proposed as signs of dietary flexibility as the environment
changed, and brown bears shifted towards increasing herbivory [15]. Based on our findings,
we suggest the decline in δ15N may reflect a loss of the more carnivorous bears, which
carried their own unique genetic signal rather than the same bears shifting their diet.
Although available δ15N for the ancient Irish individuals do not show a clear difference

between time periods (Late Pleistocene; 6.53, 6.72; Holocene: 4.57, 8.83) [48], limited
sample size prevents us from exploring this hypothesis further.
351
In the PCAs, all Late Pleistocene European bears show a close relationship to the
Himalayan bear (Fig 1B,C), suggesting the Himalayan bear population may have been much
more widespread in the past. However, this finding may be due to an overrepresentation of
ancestrally shared alleles, and underrepresentation of population specific alleles; we only
have data from one Himalayan individual and filter rare variants (minimum minor allele
frequency 0.05 or minimum minor allele count 2) to remove the noise caused by low
coverage in our ancient samples. Admixfrog does not utilise the same filtering of rare alleles,
and provides support for a filtering-driven bias; we find the Late Pleistocene individuals have
relatively high levels of ancestral alleles and low levels of Himalayan-specific ancestry. Not
belonging to the same population as the Himalayan bear is further reinforced by the D-
statistics results that show a closer relationship of the Late Pleistocene European bears to
contemporary European individuals rather than to the Himalayan individual.
364
The Late Pleistocene Siberian individuals CGG_1_020006 (eastern Siberia) and
Denisova_LP (central Siberia) also do not clearly cluster with any contemporary individuals
in the PCA (Fig 1B,C). This suggests some lost diversity in Siberia, in accordance with our
observations in Europe, and . it is in agreement with previous work showing mitochondrial
lineages in Late Pleistocene Siberian individuals that have since been lost in contemporary
individuals [50]. However, Denisova_LP does conform to the isolation-by-distance
hypothesis when comparing PC1 and longitude (Fig 1D). This is also supported by the D-
statistics results, which place Denisova_LP as intermediate between contemporary European
and Asian individuals (Fig 3), as would be expected based on its intermediate geographic
locality.. The unique placement of CGG_1_020006 is in line with a previous study
investigating its mitochondrial genome, which found it belonged to a haplogroup not found in
living brown bears [22].
377
In Japan, all contemporary Hokkaido individuals cluster together in the PCAs and
appear to be one population (Fig 1B,C). This in contrast with previous hypotheses of multiple
migration events to Hokkaido based on mitochondrial genomes [14,51]he timing of
divergence suggested brown bears migrated from mainland Eurasia to Honshu, Japan's
largest island, at least twice [14], and to Hokkaido at least three times [51]. In our analyses,
the Late Pleistocene individual from Honshu (JBB-32K) falls within or very close to the
contemporary Hokkaido individuals (Fig 1B,C). The age of JBB-32K of 32,464 ± 255 cal. BP
suggests population continuity since at least this time. The mitochondrial genome is limited
in resolution, as it only represents a single loci and may be biased by gene flow and
incomplete lineage sorting. As gene flow has been reported in Japanese brown bears [30] and
both female and male brown bears are known to disperse [52], these hypothesised migration
events may represent gene flow events into an already established Japanese population,
mediated by a few individuals, and not large population movements. A proposed land bridge
in the Tsugaru Strait connected the islands of Hokkaido and Honshu around 140,000 years
ago during the glacial period of MIS6. At this time, sea levels were at their lowest and would

have facilitated connectivity between the two islands [53]. This relatively recent connectivity likely led to the shared recent ancestry between Honshu and Hokkaido bears we see in our results.

In contemporary North American brown bears, three genetically divergent mitochondrial lineages have been described, leading to the hypothesis of three independent colonisation events from Eurasia [12]. In contrast, the nuclear data are consistent with one single metapopulation, with several subpopulations likely driven by isolation-by-distance (Fig 1B). The ancient brown bear from Quebec, aged $11,279 \pm 30$ cal. BP, may represent lost diversity, as it does not cluster closely with any of the contemporary North American bears and brown bears are thought to have gone extinct in the region by the late 20th century [54]. The Quebec individual may have come from the most recent replacement/migrations towards the end of the LGM. Previous studies using specimens dated to before this have uncovered extinct and divergent mitochondrial lineages in Eastern Beringia, suggesting dispersals into North America at widely different points in time [16].

The utilisation of published palaeogenomic data for several European and one North American brown bear enabled the recovery of lost diversity in Europe and North America. However, by generating palaeogenomic data from an additional two Asian individuals, we were able to geographically expand our inferences across their Holarctic range. In doing so, we found that although Eurasian brown bears currently display patterns of isolation by distance across the two continents, this was not always the case. Taken together, our study shows that even with ultra-low coverage palaegenomes, important inferences can be made about how a species has changed through time which is not captured in the genomes of modern individuals alone.

Data accessibility
The raw Illumina sequence data for individual JBB-32K are available under DDBJ DRA ID: DRA015631. The raw Illumina sequence data for individual CGG_1_020006 are available under Bioproject PRJNA930648 on NCBI.

Funding
This study was supported by Grants-in-Aid for Scientific Research (no 22K18730 and 23KK0062) from the Japan Society for the Promotion of Science (JSPS), the Villum Fonden Young Investigator Programme, grant no. 37352.

Acknowledgements
We thank Dr. Naoki Kohno for his valuable discussions regarding sea level changes and periods of low sea levels in Japan.

References:

1.   Stuart AJ. 2015 Late Quaternary megafaunal extinctions on the continents: a short review. *Geol. J.* **50**, 338–363.

2.   Campos PF *et al.* 2010 Ancient DNA analyses exclude humans as the driving force
behind late Pleistocene musk ox (Ovibos moschatus) population dynamics. *Proc. Natl.*
Acad. Sci. U. S. A. **107**, 5675–5680.

3.   Campos PF *et al.* 2010 Ancient DNA sequences point to a large loss of mitochondrial
genetic diversity in the saiga antelope (Saiga tatarica) since the Pleistocene. *Mol. Ecol.*
19, 4863–4875.

4.   Provan J, Bennett KD. 2008 Phylogeographic insights into cryptic glacial refugia.
Trends Ecol. Evol. **23**, 564–571.

5.   Dalén L, Nyström V, Valdiosera C, Germonpré M, Sablin M, Turner E, Angerbjörn A,
Arsuaga JL, Götherström A. 2007 Ancient DNA reveals lack of postglacial habitat
tracking in the arctic fox. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 6726–6729.

6.   McLellan BN, Proctor MF, Huber D, Michel S. 2017 *Ursus arctos* (amended version of
2017 assessment). The IUCN Red List of Threatened Species 2017.

7.   Niedziałkowska M, Hayward MW, Borowik T, Jędrzejewski W, Jędrzejewska B. 2019
A meta-analysis of ungulate predation and prey selection by the brown bear Ursus arctos
in Eurasia. *Mammal Research* **64**, 1–9.

8.   Bojarska K, Selva N. 2012 Spatial patterns in brown bear Ursus arctos diet: the role of
geographical and environmental factors. *Mamm. Rev.* **42**, 120–143.

9.   Albrecht J, Bartoń KA, Selva N, Sommer RS, Swenson JE, Bischof R. 2017 Humans
and climate change drove the Holocene decline of the brown bear. *Sci. Rep.* **7**, 10399.

10.   Marciszak A, Schouwenburg C, Lipecki G, Talamo S, Shpansky A, Malikov D, Gornig
W. 2019 Steppe brown bear *Ursus arctos 'priscus'* from the Late Pleistocene of Europe.
Quat. Int. **534**, 158–170.

11.   Taberlet P, Bouvet J. 1994 Mitochondrial DNA polymorphism, phylogeography, and
conservation genetics of the brown bear Ursus arctos in Europe. *Proc. Biol. Sci.* **255**,
195–200.

12.   Davison J *et al.* 2011 Late-Quaternary biogeographic scenarios for the brown bear
(Ursus arctos), a wild mammal model species. *Quat. Sci. Rev.* **30**, 418–430.

13.   de Jong MJ *et al.* 2023 Range-wide whole-genome resequencing of the brown bear
reveals drivers of intraspecies divergence. *Commun Biol* **6**, 153.

14.   Segawa T *et al.* 2021 Ancient DNA reveals multiple origins and migration waves of
extinct Japanese brown bear lineages. *R Soc Open Sci* **8**, 210518.

15.   Ersmark E *et al.* 2019 Genetic turnovers and northern survival during the last glacial
maximum in European brown bears. *Ecol. Evol.* **9**, 5891–5905.

16.   Salis AT *et al.* 2022 Lions and brown bears colonized North America in multiple
synchronous waves of dispersal across the Bering Land Bridge. *Mol. Ecol.* **31**, 6407–
6421.

17. Barnes I, Matheus P, Shapiro B, Jensen D, Cooper A. 2002 Dynamics of Pleistocene
population extinctions in Beringian brown bears. *Science* **295**, 2267–2270.

18. Westbury MV *et al.* 2020 Hyena paleogenomes reveal a complex evolutionary history of
cross-continental gene flow between spotted and cave hyena. *Science Advances* **6**,
eaay0456.

19. Liu S *et al.* 2021 Ancient and modern genomes unravel the evolutionary history of the
rhinoceros family. *Cell* **184**, 4874–4885.e16.

20. Cahill JA *et al.* 2018 Genomic Evidence of Widespread Admixture from Polar Bears
into Brown Bears during the Last Ice Age. *Mol. Biol. Evol.* **35**, 1120–1129.

21. Barlow A *et al.* 2018 Partial genomic survival of cave bears in living brown bears. *Nat
Ecol Evol* **2**, 1563–1570.

22. Rey-Iglesia A, García-Vázquez A, Treadaway EC, van der Plicht J, Baryshnikov GF,
Szpak P, Bocherens H, Boeskorov GG, Lorenzen ED. 2019 Evolutionary history and
palaeoecology of brown bear in North-East Siberia re-examined using ancient DNA and
stable isotopes from skeletal remains. *Sci. Rep.* **9**, 4462.

23. Orlando L *et al.* 2013 Recalibrating Equus evolution using the genome sequence of an
early Middle Pleistocene horse. *Nature* **499**, 74–78.

24. Dabney J *et al.* 2013 Complete mitochondrial genome sequence of a Middle Pleistocene
cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci. U. S. A.*
110, 15758–15763.

25. Meyer M, Kircher M. 2010 Illumina sequencing library preparation for highly
multiplexed target capture and sequencing. *Cold Spring Harb. Protoc.* **2010**,
db.prot5448.

26. Glocke I, Meyer M. 2017 Extending the spectrum of DNA sequences retrieved from
ancient bones and teeth. *Genome Res.* **27**, 1230–1237.

27. Miller W *et al.* 2012 Polar and brown bear genomes reveal ancient admixture and
demographic footprints of past climate change. *Proc. Natl. Acad. Sci. U. S. A.* **109**,
E2382–90.

28. Cahill JA *et al.* 2013 Genomic evidence for island population conversion resolves
conflicting theories of polar bear evolution. *PLoS Genet.* **9**, e1003345.

29. Benazzo A *et al.* 2017 Survival and divergence in a small group: The extraordinary
genomic history of the endangered Apennine brown bear stragglers. *Proc. Natl. Acad.
Sci. U. S. A.* **114**, E9589–E9597.

30. Endo Y, Osada N, Mano T. 2021 Demographic History of the Brown Bear (Ursus
arctos) on Hokkaido Island, Japan, Based on Whole-Genomic Sequence Analysis.
Genome Biol.

31. Cahill JA, Stirling I, Kistler L, Salamzade R, Ersmark E, Fulton TL, Stiller M, Green
RE, Shapiro B. 2015 Genomic evidence of geographically widespread effect of gene

flow from polar bears into brown bears. *Mol. Ecol.* **24**, 1205–1217.

32.   Schubert M *et al.* 2014 Characterization of ancient and modern genomes by SNP
detection and phylogenomic and metagenomic analysis using PALEOMIX. *Nat. Protoc.*
9, 1056–1082.

33.   Schubert M, Lindgreen S, Orlando L. 2016 AdapterRemoval v2: rapid adapter trimming,
identification, and read merging. *BMC Res. Notes* **9**, 88.

34.   Li H, Durbin R. 2009 Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics* **25**, 1754–1760.

35.   Liu S *et al.* 2014 Population genomics reveal recent speciation and rapid evolutionary
adaptation in polar bears. *Cell* **157**, 785–794.

36.   Castruita JAS, Westbury MV, Lorenzen ED. 2020 Analyses of key genes involved in
Arctic adaptation in polar bears suggest selection on both standing variation and de novo
mutations played an important role. *BMC Genomics* **21**, 1–8.

37.   Li H *et al.* 2009 The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
25, 2078–2079.

38.   Broad institute. 2019 Picard Toolkit. See http://broadinstitute.github.io/picard.

39.   McKenna A *et al.* 2010 The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303.

40.   Chen S, Zhou Y, Chen Y, Gu J. 2018 fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* **34**, i884–i890.

41.   Korneliussen TS, Albrechtsen A, Nielsen R. 2014 ANGSD: Analysis of Next
Generation Sequencing Data. *BMC Bioinformatics* **15**, 356.

42.   Meisner J, Albrechtsen A. 2018 Inferring Population Structure and Admixture
Proportions in Low-Depth NGS Data. *Genetics* **210**, 719–731.

43.   Meisner J, Liu S, Huang M, Albrechtsen A. 2021 Large-scale inference of population
structure in presence of missingness using PCA. *Bioinformatics* **37**, 1868–1875.

44.   Taron UH, Lell M, Barlow A, Paijmans JLA. 2018 Testing of Alignment Parameters for
Ancient Samples: Evaluating and Optimizing Mapping Parameters for Ancient Samples
Using the TAPAS Tool. *Genes* **9**. (doi:10.3390/genes9030157)

45.   Peter BM. 2020 100,000 years of gene flow between Neandertals and Denisovans in the
Altai mountains. *bioRxiv*

46.   Narasimhan V, Danecek P, Scally A, Xue Y, Tyler-Smith C, Durbin R. 2016
BCFtools/RoH: a hidden Markov model approach for detecting autozygosity from next-
generation sequencing data. *Bioinformatics* **32**, 1749–1751.

47.   Westbury MV *et al.* 2018 Extended and Continuous Decline in Effective Population
Size Results in Low Genomic Diversity in the World's Rarest Hyena Species, the Brown

Hyena. *Mol. Biol. Evol.* **35**, 1225–1237.

48. Edwards CJ *et al.* 2011 Ancient hybridization and an Irish origin for the modern polar
bear matriline. *Curr. Biol.* **21**, 1251–1258.

49. Sommer RS, Benecke N. 2005 The recolonization of Europe by brown bears Ursus
arctos Linnaeus, 1758 after the Last Glacial Maximum. *Mamm. Rev.* **35**, 156–164.

50. Molodtseva AS *et al.* 2022 Phylogeography of ancient and modern brown bears from
eastern Eurasia. *Biol. J. Linn. Soc. Lond.* **135**, 722–733.

51. Hirata D *et al.* 2013 Molecular phylogeography of the brown bear (Ursus arctos) in
Northeastern Asia based on analyses of complete mitochondrial DNA sequences. *Mol.*
Biol. Evol. **30**, 1644–1652.

52. Støen O-G, Zedrosser A, Saebø S, Swenson JE. 2006 Inversely density-dependent natal
dispersal in brown bears Ursus arctos. *Oecologia* **148**, 356–364.

53. Lisiecki LE, Raymo ME. 2005 A Pliocene☐Pleistocene stack of 57 globally distributed
benthic δ18O records. *Paleoceanography* **20**. (doi:10.1029/2004pa001071)

54. Spiess A, Cox S. 1976 Discovery of the Skull of a Grizzly Bear in Labrador. *Arctic* **29**,
194–200.

563
564
565   **Figure legends:**
566
567
Figure 1: Brown bear sample localities and population structure. A) Map showing the
localities of the brown bear ancient and contemporary brown bear genomes analysed.
Calibrated radiocarbon ages of the ancient samples included are listed next to their locality.
'Late Pleistocene' next to Denisova_LP indicates a direct age is unknown, but it is listed as
Late Pleistocene in the original publication [26]. CGG_1_020006 was dated to beyond the
limit of radiocarbon dating (>50 kya) but molecularly dated to 61,826 years BP [HPD 95%:
27,866–96,783 years BP][23]. B) PCA generated from genotype likelihoods of all samples.
C) PCA generated from genotype likelihoods of the Eurasian samples only. Percentage
values in parentheses on the X and Y axes show the percentage of variation explained by the
principal component. Colours and symbols represent geography and age, respectively. D)
PC1 vs degrees longitude for the Eurasian individuals, excluding the Japanese individuals.
Dark grey circle shows the single Himalayan individual. The colour of CentralRussia1 was
changed to light red due to its geographic position in the Ural mountains and European-like
clustering in the PCA.
582
583   **Fig 2 - Relationship of ancient Eurasian individuals to contemporary individuals.** A)
584   Genome-wide pairwise distances to contemporary individuals. B) Ancestry proportions
585   calculated using three reference populations. Ancestral is defined as having combined
586   ancestry between any of the two reference populations. Coloured boxes above the names on

the X axis correspond to the geographic location (longitude) of the ancient sample as seen in Fig 1A.

Fig 3 - Test for population structure using D-statistics and the input topology [[[Europe, Asia], Ancient], Outgroup]. A negative D-score suggests a closer relationship of an ancient individual to western European individuals, whereas a positive D-score suggests a closer relationship to the Asian individual. Dot colours show the Asian individuals' geography (longitude) as seen in the map insert.. Bar colours next to their names represent the geographic location (longitude) of the ancient individuals based on the colours in Fig 1A.