Uncovering bacterial hosts of class 1 integrons in an urban coastal aquatic environment with a single-cell fusion-PCR technology

Qin Qi*‡, Timothy M Ghaly‡, Anahit Penesyan, Vaheesan Rajabal, Jeremy AC Stacey,

Sasha G Tetu, Michael R Gillings

* Correspondence: qin.qi@mq.edu.au

‡ These authors contributed equally

AUTHOR ADDRESS

School of Natural Sciences, Macquarie University, 14 Eastern Road, Sydney, NSW 2109,

Australia

Abstract

Horizontal gene transfer (HGT) is a key driver of bacterial evolution via transmission of genetic

materials across taxa. Class 1 integrons are genetic elements that correlate strongly with

anthropogenic pollution and contribute to the spread of antimicrobial resistance (AMR) genes

via HGT. Despite their significance to human health, there is a shortage of robust, culture-free

surveillance technologies for identifying uncultivated environmental taxa that harbour class 1

integrons. We developed a modified version of epicPCR (emulsion, paired isolation and

concatenation PCR) that links class 1 integrons amplified from single bacterial cells to

taxonomic markers from the same cells in emulsified aqueous droplets. Using this single-cell

genomic approach and Nanopore sequencing, we successfully assigned class 1 integron gene

cassette arrays containing mostly AMR genes to their hosts in coastal water samples that were

affected by pollution. Our work presents the first application of epicPCR for targeting variable,

multi-gene loci of interest. We also identified the *Rhizobacter* genus as novel hosts of class 1

integrons. These findings establish epicPCR as a powerful tool for linking taxa to class 1

integrons in environmental bacterial communities and offer the potential to direct mitigation

efforts towards hotspots of class 1 integron-mediated dissemination of AMR.

Synopsis

31　We present a novel single-cell genomic surveillance technology for identifying environmental

32　bacterial hosts of a class of mobile genetic elements that are linked to anthropogenic pollution

33　and contribute to the dissemination of antimicrobial resistance.

Keywords

36　Integrons, antimicrobial resistance (AMR), anthropogenic pollution, class 1 integron gene

37　cassettes, horizontal gene transfer (HGT), mobile genetic elements, environmental bacterial

38　hosts, single-cell genomics

Introduction

41　Horizontal gene transfer (HGT) is an important source of genomic variability upon which

42　selection can act [1-3]. Microbial adaptation via HGT most commonly occurs among closely

43　related taxa or between more distantly related taxa found in similar environments [4, 5]. A key

example is the dissemination of antimicrobial resistance (AMR) genes via conjugation, bacteriophage transduction, and natural transformation by extracellular DNA [6-9]. To understand the dynamics of HGT, we need to identify the sources of horizontally transferred genes and the range of taxa they subsequently inhabit [10-12]. This is challenging given the sheer diversity of microbial taxa and the difficulty of culturing many environmental microbes [13-16]. Metagenomic sequencing of bacterial communities does not require prior culturing of microbes [17, 18]. However, mobile genetic elements are often associated with repetitive DNA sequences and exhibit complex interactions with genome rearrangements, which limit our ability to attribute laterally transferred genes to host genomes due to assembly and/or taxonomic binning difficulties [19, 20].

epicPCR (emulsion, paired isolation and concatenation PCR) is a fusion PCR-based technique that overcomes these problems by co-amplifying taxonomic markers with functional genes of interest from single uncultured bacterial cells. Single cells are trapped in droplets generated by emulsifying an aqueous phase containing PCR reagents and environmental bacteria in an emulsion oil [21-26]. This compartmentalises the fusion PCR into parallel, single cell-level reactions. Within each droplet, chimeric amplicons that combine genes of interest with taxonomic markers corresponding to diverse phylogenetic groups are generated for high-throughput sequencing. To date, epicPCR has been successfully applied to target functional genes including dissimilatory sulfite reductase *dsrB* genes [21, 24], antibiotic resistance genes [22, 23, 26, 27], class 1 integron associated genes (*intI1*, *sul1*, Δ*qacE*) [23, 27, 28], and bacteriophage-borne ribonucleotide reductase *RNR* genes [25]. epicPCR amplicons typically comprise short fragments of genes of interest linked to the V4 hypervariable region of 16S rRNA markers. Illumina sequencing of recovered amplicons demonstrates the presence of the genes of interest in a variety of bacterial taxa. However, no studies have so far applied epicPCR to target variable, multi-gene loci of interest.

Class 1 integrons contribute to the dissemination of AMR genes via HGT in natural ecosystems

and clinical settings [29, 30]. Integrons are genetic elements that comprise an integron-integrase

gene (*intI*), an *attI* recombination site, and a promoter (P_C) that drives the expression of variable

gene cassettes [31-33]. IntI catalyses the integration and excision of gene cassettes via *attC* sites

that can be recombined with *attI* or other *attC* sites [33-36]. All class 1 integrons contain highly

conserved integron-integrase (*intI1*) and *attI1* sequences (Figure 1A). Clinical class 1 integrons

are strongly associated with AMR gene cassettes that confer resistance to antibiotics,

disinfectants, or heavy metals [29, 31, 37]. Mobile genetic elements such as broad host-range

plasmids and transposons have acquired class 1 integrons, resulting in the proliferation of class

1 integrons across diverse bacterial hosts [31, 38]. Recently, it has been reported that phage-

plasmids can also disseminate class 1 integrons [39].

Bacterial hosts of class 1 integrons have previously been isolated and characterised through

colony PCR screening and fosmid library preparation [29, 40-43]. The necessity to pre-isolate

environmental bacteria severely biases host identification towards the cultivated faction and

hinders our ability to attribute relative abundances of class 1 integrons to individual taxa. To

overcome these limitations, we developed a modified experimental and bioinformatic pipeline

for epicPCR that associates class 1 integrons with their bacterial hosts. Class 1 integron gene

cassette arrays are co-amplified with the V4 hypervariable region of the 16S rRNA marker

genes from individual bacterial cells (Figure 1B-C). The fused amplicons are sequenced using

Nanopore long-read technology. We applied stringent bioinformatic filtering to ensure that the

associations between the class 1 integrons and the taxonomic markers were *bona fide.* We

successfully tested our epicPCR pipeline on water samples from a coastal environment and

linked six unique class 1 integrons with variable gene cassettes to seven distinct environmental

bacterial taxa. While most of the gene cassettes encode AMR genes, we also linked a gene

cassette encoding a cupin-domain containing protein of unknown functions to two

Alphaproteobacteria hosts. Furthermore, our results demonstrated that *Rhizobacter* is a novel

host of class 1 integrons. Overall, our study shows that epicPCR is a powerful tool for

determining the environmental hosts of class 1 integrons with the potential to access previously

untapped carriers of these genetic elements. This paves the way for future studies that seek to

pinpoint the "sources" and "sinks" of class 1 integrons.

Materials and Methods

Isolation of bacteria from an urban coastal location

Coastal water samples were collected in three 50 mL biological replicates after rain in a rocky

intertidal zone downstream from a stormwater outlet at Shark Point (location coordinates: -

33.91, 151.27) near Sydney, NSW, Australia (Supplementary Figure 1). The samples were

filtered using EASYstrainer cell strainers with a mesh size of 40 μm (Greiner AG, Germany)

and centrifuged at 3220 *g* for 15 minutes. Cell pellets were resuspended in phosphate buffered

saline (PBS) solution at pH 7.4. Resuspended cells were stored as 25% glycerol stocks.

Bacterial cell count estimates by fluorescence microscopy

Approximately 10^5-10^6 cells were fixed and stained in PBS containing 4% formaldehyde and

2 μg/mL 4′,6-diamidino-2-phenylindole (DAPI) for 30 min in the dark with rotation at room

temperature. Re-suspended cells were diluted in PBS, transferred to a haemocytometer and

imaged under the x40 lens of an Olympus BX63 fluorescence microscope. Differential

interference contrast (DIC) and DAPI fluorescence channel images were overlaid and analysed

with ImageJ (National Institutes of Health, United States). Bacterial cell densities were

calculated from the average number of cells in five 0.2 mm x 0.2 mm square grids of the

counting chamber for each sample (Supplementary Figure 2).

Emulsion, paired-isolation and concatenation PCR (epicPCR)

Our hybrid epicPCR procedures were adapted from the protocols of Sakowski *et al* [25] and

Diebold *et al* [26]. Briefly, epicPCR was performed in three technical replicates for the three

biological replicates. Approximately 150,000 cells from the bacterial glycerol stocks were

pelleted and vortexed in a 75 µL PCR reagent mix containing GC buffer (1x), dNTP mix (0.4

mM), Phusion Hot-Start II polymerase (0.05 U/µL) (Thermo Scientific, United States), bovine

serum albumin solution (1 µg/µL) (Promega, United States), Lucigen Ready-Lyse lysozyme

(500 U/µL) (LGC Biosearch Technologies, United States) and oligonucleotide primers R926

(2 µM), *intI1*_outer (1 µM), and R519-*qacE* bridging primer (0.04 µM) (Supplementary Table

S2). To demonstrate the absence of false associations between 16S rRNA markers and class 1

integrons originating from different cells, we spiked one set of technical replicates with cells

of a class 1 integron-free *Escherichia coli* MG1655 strain at a total population frequency of

approximately 10%.

ABIL emulsion oil was prepared by supplementing mineral oil with 4% ABIL EM 90 (Redox,

Australia) and 0.05% Triton X-100 (Promega, United States). 425 µL ABIL oil was added to

the PCR mix, which was immediately emulsified at 4 ms^{-1} for 45 s using the FastPrep-24 bead

beating system (MP Biomedicals, United States). The water-in-oil emulsion was aliquoted into

8 portions, incubated at 37˚C for 10 min, and was subjected to the following PCR amplification

conditions: 98˚C for 5 minutes; 38 cycles of 98°C for 10 s, 59°C for 20 s, and 72°C for 2 min;

and finally 72°C for 5 min. The *intI1*_outer primer and the R519-*qacE* bridging primer bind to

intI1 in the 5'-conserved segment (CS) and *qacE* in the 3'-CS of class 1 integrons respectively

(Figure 1B). The 18 bp overhang introduced by the bridging primer to the *qacE* end allows the

intermediate PCR products to act as a long primer and amplify the V4-V5 hypervariable region

with the R926 reverse primer.

The emulsified PCR mix for each technical replicate was pooled, vortexed in 900 µL isobutanol

and 200 µL 5M NaCl and centrifuged for 1 min at ~20,000g with soft brake. The aqueous phase

at the bottom of the tubes was extracted, purified using the Monarch PCR & DNA Cleanup Kit

(New England Biolabs, United States), and eluted with nuclease-free water. Sera-Mag Select

magnetic beads (Cytiva, United States) were added to the eluted PCR products to deplete <300

bp DNA fragments. The size-selected DNA was used as template in 100 µL Phusion PCR mix

containing GC buffer (1x), dNTP mix (0.4 mM), Phusion Hot-Start II polymerase (0.04 U/µL),

AP27_short primer (0.8 µM), *intI1*_nested primer (0.4 µM), forward and reverse blocking

primers (0.32 µM each). The PCR mix subjected to the following thermocycling conditions in

8 aliquots: 98˚C for 30 s; 35 cycles of 98°C for 10 s, 59°C for 20 s and 72°C for 1 min 50 s;

followed by a final step of 72°C for 5 min. epicPCR products were visualised on 1% agarose

gel by electrophoresis and treated with Sera-Mag Select magnetic beads to deplete epicPCR

products containing cassette-less integrons.

To confirm that fusion between class 1 integron and 16S DNA fragments has occurred

(Supplementary Figure 3A), 2 ng of each purified epicPCR product was amplified using

*qacE*_F (0.4 µM) and AP28_short (0.4 µM) primers in GoTaq polymerase master mix

(Promega, United States) under the following thermocycling conditions: 95˚C for 30 s; 30

cycles of 95°C for 30 s, 55°C for 30 s and 72°C for 30 s; with a final step of 72°C for 5 min.

Successful amplification of the *qacE*-16S rRNA gene chimeric region should produce ~390 bp

gel bands.

Sanger sequencing of epicPCR products from the *Rhizobacter* genus

To confirm that class 1 integrons can be found in the *Rhizobacter* genus, we pooled approximately 150,000 cells from the three biological replicates and performed the epicPCR procedures in three technical replicates with the minor modifications shown in Supplementary Figure 3B: the R926 and AP27_short primers were replaced with *Rhizobacter*-specific primers *Rhizobacter*_16S_ R926 and *Rhizobacter*_16S_ R806 respectively, which were designed based on the 16S rRNA sequence of *R. gummiphilus* strain NS21 (NCBI accession no. CP015118) [44]. qacE_F was used as the nested forward primer. The resulting 389 bp epicPCR products were verified by Sanger sequencing (Marcogen Inc, South Korea).

Oxford Nanopore long-read sequencing

Purified epicPCR amplicons were treated with the NEBNext Ultra II End Repair/dA-Tailing Module (New England Biolabs, United States), purified with JetSeq Clean magnetic beads (Meridian Bioscience, United States), and eluted using nuclease-free water. End-repaired DNA was barcoded using the Native Barcoding Expansion kit (ONT, United Kingdom) and Blunt/TA Ligase Master Mix (New England Biolabs, United States). Barcoded samples were multiplexed and ligated to sequencing adaptor molecules in Adapter Mix II using the NEBNext Quick T4 DNA ligase (New England Biolabs, United States). The ligation products were washed twice with Short Fragmentation Buffer and eluted with Elution Buffer. The sequencing library was loaded into a FLO-MIN106D flow cell R9.4.1 and a MinION Mk1B sequencer (ONT, United Kingdom). The minimum threshold for read length was set to 1 kB on the MinKNOW operating software. Basecalling of FAST5 raw data was performed with the high accuracy option using Guppy v6.1.2 (ONT, United Kingdom). The native barcoded samples were demultiplexed using default parameters.

Analysis of Nanopore sequencing data of epicPCR amplicons

All filtering and processing steps were carried out using an in-house pipeline

(https://github.com/timghaly/Int1-epicPCR). First, the pipeline quality-filters reads using

NanoFilt v2.8.0 [45], removing those with an average read quality of <7 and read length <670 bp,

which represents the minimum length of an epicPCR product with a cassette-less integron.

Quality-filtered reads are then oriented and trimmed with the final nested forward and reverse

primer sequences using Pychopper v2.7.0 (https://github.com/epi2me-labs/pychopper).

Pychopper identifies both primers in each read using edlib v1.2.3 [46], orients the reads based on

the forward and reverse primer sequences, and discards reads that do not contain both primers

in the correct orientations. The pipeline then clusters the primer-oriented reads into amplicon-

specific clusters using isONclust v0.0.6.1 [47]. Error correction was performed on each cluster

using isONcorrect v0.0.8 [48], which can jointly use all cassette arrangements of the same

integron that occur in different clusters and allow efficient error correction even for amplicons

with low sequencing depths. A consensus sequence is then generated for each cluster using

spoa v4.0.7 [49]. All consensus sequences are then pooled, while removing any redundancies,

including reverse complement redundancies, using dedupe from BBTools v35

(https://github.com/kbaseapps/BBTools). Next, the consensus sequences are screened for the

R519-*qacE* bridging primer and the 157 bp region (covering the 5'-end of *intI1* and all of *attI1*)

using blastn from BLAST v2.2.31. Retained sequences must contain both regions. Any

sequences containing more than one hit to either of these sequences are considered unintended

chimeras and discarded. Finally, sequences are screened for the correctly fused 16S rRNA gene

fragment using Metaxa2 v2.2.3 [50]. The final output of the pipeline for each sample is a set of

full-length, primer-oriented amplicon consensus sequences that contain a complete *attI1*

sequence, R519-*qacE* bridging primer, and the V4 hypervariable region of the 16S rRNA gene.

Cd-hit v4.8.1 [51] was used to cluster chimeric epicPCR products comprising class 1 integron

gene cassette arrays and V4 hypervariable regions of 16S rRNA gene with ≥99% pairwise

identity in nucleotide sequences in at least 3 epicPCR replicates. For epicPCR products that

were found in fewer than 3 replicates, the NCBI Genome Workbench was used to perform

blastn searches of these sequences against a local database created using all the Nanopore reads

obtained in this study. Reads with ≥98% pairwise identity in nucleotide sequences and ≥98%

coverage were aligned using the Geneious bioinformatic software (Biomatters, New Zealand),

and the consensus sequences were generated from these alignments within each set of epicPCR

replicates. Consensus sequences that could be found in at least 3 epicPCR replicates and

showed ≥99% pairwise identity in nucleotide sequences and ≥99% identical sites were added

to the list of epicPCR products that have already been shortlisted by the Cd-hit algorithm.

IntegronFinder 2.0 [parameters: --local-max --gbk --promoter-attI --calin-threshold 1] was

used to predict the ORFs and *attC* sites [52].

Taxonomic classification for bacterial hosts of class 1 integrons

The sequences of V4 hypervariable regions of 16S rRNA gene sequences obtained in this study

were searched against the SILVA 16S rRNA gene database website [53] using the Alignment,

Classification and Tree (ACT) tool with a cut-off threshold of 0.80. The V4 region sequences

were aligned using the Geneious multiple alignment tool using default parameters and cost

matrix = 93% similarity. A phylogenetic tree was constructed with PhyML (v3.0) using the

maximum likelihood method based on the Hasegawa-Kishino-Yano (HKY85) model with

bootstrapping (number of replicates = 100) [54]. To verify genus-level taxonomic classifications,

the V4 hypervariable regions of 16S rRNA for the *Rhizobacter* and *Aquabacterium*-like hosts

found in this study were searched against the NCBI Nucleotide Collection by blastn with the

exclusion of "uncultured/environmental sample sequences". The 16S rRNA sequences of all

identical matches were searched against the SILVA 16S rRNA gene database. All the reference

genomes of *Rhizobacter* and *Aquabacterium* species were downloaded from the NCBI RefSeq

database (last accessed on 11 November 2022). Nucleotide alignment was performed against

the full 16S rRNA gene sequences from the RefSeq genomes using the Geneious bioinformatic

software.

Results and Discussion

Coastal water samples were collected in three biological replicates from the rockpools

downstream of a stormwater outlet at Shark Point, Sydney, New South Wales, Australia

(Supplementary Figure 1). We anticipated that class 1 integron-associated environmental

bacteria would be detected at this location, because surface runoff was expected to carry

pollutants from the surrounding urban areas. A preponderance of cells displayed unicellular

and planktonic morphologies (Supplementary Figure 2). We applied modified epicPCR

procedures to amplify class 1 integron gene cassette arrays from single bacterial cells

encapsulated by PCR reagent droplets homogenously dispersed in an oil emulsion and

successfully generated chimeric epicPCR products containing class 1 integrons and 16S rRNA

gene sequences (Figure 1). These epicPCR products were Nanopore sequenced. The median

number of reads per sample was 232,402. After stringent bioinformatic filtering, we identified

six class 1 integrons in seven distinct bacterial hosts (Figure 2). All seven combinations of

epicPCR products were observed in at least three independent replicates ($n{\geq}3$, $\geq$99% pairwise

nucleotide identity and $\geq$99% identical sites relative to the consensus sequence for each

epicPCR product). In addition, there were five combinations of class 1 integrons and hosts that

were detected in fewer than 3 replicates and were consequently excluded. Taxonomic

classifications were obtained using the SILVA ribosomal RNA gene database (Supplementary

Table S1) [53]. *Alphaproteobacteria* (*n*=2) and *Gammaproteobacteria* (*n*=5) were the two major

classes of hosts represented in our dataset, while *Burkholderiales* (*n*=4) was the most common

bacterial order. These observations are broadly consistent with previous findings that class 1

integrons are associated with *Gammaproteobacteria* [29, 42, 43]. The finding of integrons in two

Alphaproteobacteria is notable, because *Alphaproteobacteria* are rarely known to be

associated with integrons [55]. Currently, there are very few examples of *Alphaproteobacteria*

species that harbour class 1 integrons [56-58]. This underscores the potential of epicPCR

approaches to discover novel host-function associations.

Six different gene cassettes were detected, five of which encoded well-characterised AMR

genes (Table 1). The nucleotide sequences of all the gene cassettes reported in this study are

summarised in Supplementary Table S3. We compared their nucleotide sequences against

reference class 1 integron gene cassettes from the NCBI database and found that both the

coding and non-coding regions of these gene cassettes were highly conserved relative to the

reference sequences [34, 59]. In our dataset, there are six different *attC* recombination sites with

100% nucleotide sequence identity to those in the respective reference gene cassettes in Table

1. We also aligned all the nucleotide sequences in the 5'-CS and 3'-CS regions of all the class

1 integrons from this study and found that both the 157 bp *intI1/attI1* and 190 bp *attC/qacE*

regions in the chimeric amplicons were identical in all but one exception (Supplementary

Figure 4). Overall, these class 1 integron structures are typical of those recovered from clinical

bacterial isolates [29, 60, 61].

Class 1 gene cassettes detected by epicPCR

Cupins belong to a superfamily of proteins with a beta-barrel fold and were originally found as highly conserved domains in plant proteins [62]. We found a gene cassette encoding a cupin barrel domain of unknown functions in two *Alphaproteobacteria* hosts. One host was unclassified at the order level (Figure 2A), while the other belonged to the *Rhodobacteraceae* family (Figure 2B). This cupin-domain-encoding gene was previously observed in a class 1 integron gene cassette from an unknown host [59]. Based on identical nucleotide sequence matches from the NCBI nucleotide database, this gene cassette was detected in class 1 integrons, non-class 1 integrons, and CALINs (clusters of *attC*s lacking an associated integron-integrase) in the chromosomes or plasmids of several *Gammaproteobacteria* hosts (Supplementary Table S4). This observation suggests selective pressure for maintaining this gene cassette in both class 1 and non-class 1 integron genetic contexts. To the best of our knowledge, its association with an *Alphaproteobacteria* host has not been reported previously.

As expected, the majority of the class 1 integron gene cassettes we detected are AMR related. The dihydrofolate reductase encoded by *dfrA5* confers high levels of trimethoprim resistance in Gram-negative bacteria [63]. We found an association between the *dfrA5* gene cassette and a *Comamonadaceae*-family host (Figure 2C). The *blaOXA*$_{10}$ gene encodes a class D beta-lactamase that hydrolyses the semi-synthetic antibiotic oxacillin [64, 65]. In our dataset, the *blaOXA*$_{10}$ gene cassette was detected in a *Moraxellaceae*-family host (Figure 2D). Genes *qacF* and *qacG2* encode small multidrug resistance (SMR) efflux pumps that confer resistance to quaternary ammonium compounds (QACs) present in a wide range of disinfectants and detergents [31, 66]. We found a *qacG2* gene cassette in an *Aquabacterium*-like host (Figure 2E) and a *qacF* gene cassette at the proximal position in the gene cassette array in a *Rhizobacter* host (Figure 2F). Also found in the class 1 integron of the *Rhizobacter* host is an *aadA2* gene cassette. *aadA* genes confer resistance to the aminoglycoside antibiotics streptomycin and

310 spectinomycin [34, 67-70]. In the *Rhodocyclaceae*-family host, we found the same *aadA2* gene

311 cassette with a SNP in the non-coding region and a non-synonymous mutation (V257A) in the

312 coding region (Figure 2G and Supplementary Table S3). The phylogenetic relationship

313 between the seven hosts and the distribution of class 1 integron gene cassettes amongst them

314 are summarised in Figure 3.

316 Discovery of *Rhizobacter* as novel carriers of class 1 integrons

Rhizobacter is an environmental bacterial genus. Isolates of *Rhizobacter* species have been

successfully isolated from the rhizospheres of the *Panax* [71, 72] and *Bergenia* genera [73], as well

as from freshwater sediment [74]. There are still many questions regarding their potential

ecological roles. For example, the *R. gummiphilus* naturally degrades rubber [44, 75, 76], which

implies beneficial roles in promoting nutrient recycling and bioremediation in soil. On the other

hand, *R. daucus* is thought to be an agricultural plant pathogen that causes galls in carrot root

tissues [77]. Associations between the *Rhizobacter* genus and class 1 integrons have not been

reported previously. In all currently available NCBI RefSeq genomes for the *Rhizobacter* genus

($n = 11$), we detected various non-class 1 integrons but did not find any bioinformatic evidence

of class 1 integrons.

We found two *Rhizobacter* isolates with identical nucleotide sequence in the 16S rRNA V4

hypervariable region to that of the *Rhizobacter* host in our dataset (NCBI accession no.

AB835065.1 and LC734193.1) [78, 79]. To experimentally verify that the association between the

Rhizobacter and class 1 integron was *bona fide*, we designed *Rhizobacter*-specific 16S rRNA

gene primers that target the same primer binding sites in the 16S rRNA gene as R926 and R806.

By replacing the *intI1*_nested primer with *qacE*_F (the primer positions are indicated in

Supplementary Figure 3A) in our epicPCR experimental pipeline, we successfully obtained the

expected chimeric epicPCR product featuring a *qacE* fragment and the *Rhizobacter* phylogenetic marker. This provides the first experimental evidence of an association between class 1 integrons and the *Rhizobacter* genus.

A recent study applied a modelling approach to understand the emergence and spread of novel antimicrobial resistance genes [80]. The dispersal of antibiotic resistance genes from environmental bacteria to humans has been highlighted as a major knowledge gap that contributes to uncertainties in these models. Acquiring more experimental data for these gene mobilisation and HGT events can improve future quantitative risk assessment models for antibiotic resistance and our ability to curb transfer of antibiotic resistance genes between bacteria. epicPCR has the potential to contribute to this endeavour given its strengths in identifying environmental microbes that cannot be cultivated under laboratory conditions.

The seven bacterial hosts described in this study are expected to be a subset of all the class 1 integron hosts in our samples. Our bridging primer R519-*qacE* cannot bind to class 1 integrons that lack *qacE*/Δ*qacE*. These include many chromosomal class 1 integrons and atypical class 1 integrons with IS*26*-mediated deletions in their 3'-CS [29, 41]. To detect these class 1 integrons and their hosts requires new bridging primers that bind to the 3'-end of their gene cassette arrays. In addition, the presence of epicPCR products that were detected in fewer than 3 replicates in this study could be demonstrated by replacing the two *intI1*-targeting forward primers (*intI1*_outer and *intI1*_nested) with specific primers that target the 5'-end of specific gene cassettes of interest. The degenerate bases in the 16S rRNA gene-targeting primers also mean that certain hosts are less likely to be detected than others due to primer bias. Another inherent limitation is the variable resolution of taxonomic classifications when using 16S rRNA as phylogenetic marker.

We established epicPCR as a valuable surveillance platform for uncovering novel associations

between class 1 integrons with diverse cassette arrays and their bacterial hosts in environmental

water samples. Using this single-cell genomic approach, we successfully identified

Alphaproteobacteria as novel hosts of a gene cassette encoding a cupin-domain protein of

unknown functions, and the *Rhizobacter* genus as novel carriers of class 1 integrons. It is

envisaged that this technology can be deployed to analyse water samples from sewage

treatment plants, as well as effluents from industrial, clinical, and agricultural settings where

antimicrobial usage is high. These surveillance efforts will hopefully lead to a timelier

detection of hotspots of class 1 integron mediated AMR transmission and more targeted

environmental mitigation measures. epicPCR also paves the way for future studies that seek to

uncover the "sources" and "sinks" of integron gene cassettes in bacterial communities, as well

as possible barriers to integron-mediated HGT that may also exist [81]. With greater diversity in

combinations of gene cassette and bacterial hosts that are likely to be captured through future

epicPCR studies, we will be able to address outstanding questions and gain more insights into

possible origins of novel gene cassettes, which has so far remained the greatest mystery in the

field of integron research [32, 35, 82].

Figure Caption

Figure 1. Class 1 integrons and their co-amplification with 16S rRNA marker genes via

epicPCR.

(A) The class 1 integron carried by the IncW plasmid R388 has the canonical structure of

clinical class 1 integrons. *intI1* and *attI1* sequences are highly conserved in the 5'-conserved

segments (CS). The 3'-CS comprise a truncated quaternary ammonium compound (QAC)

resistance gene (Δ*qacE*) and a sulfonamide resistance gene (*sul1*). Between the 5'-CS and 3'-

CS are variable integron gene cassette arrays that contain *attC* recombination sites and gene

cassettes. Each gene cassette is typically associated with a single open-reading frame (ORF).

(B) During the first stage of epicPCR, the *intI1*_outer and R519_*qacE* bridging primers target

intI1 and *qacE* sequences in the 5'-CS and 3'-CS respectively and amplify the class 1 integron.

A short overhang introduced by the bridging primer to the *qacE* fragment end allows the

intermediate PCR product to act as a long primer and co-amplify the V4-V5 hypervariable

regions of the 16S rRNA marker gene in the presence of the R926 reverse primer.

(C) During the second stage of epicPCR, the *intI1*_nested and AP27_short nested primers

further amplify the chimeric amplicons pooled from the previous step. AP27_short introduces

a non-degenerate tail sequence (in blue) to the 3'-end of the amplicons. During standard PCR

verification of chimeric amplicon formation (Supplementary Figure 3A), this tail sequence is

targeted by the AP28_short non-degenerate primer, which is used in conjunction with the

*qacE*_F forward primer. The final chimeric epicPCR amplicons contain highly conserved

features of class 1 integrons, including a 157 bp *intI1-attI1* region in the 5'-CS and the

qacE/Δ*qacE* fragment in the 3'-CS.

Figure 2: Class 1 integrons identified through epicPCR alongside the taxonomic classification

of their respective bacterial hosts. The *intI1* gene fragment is displayed in blue, the *attI1* site in

purple, and all the gene cassette ORFs in green. The six different *attC* sites are shown in red.

Figure 3: Phylogenetic tree constructed using the maximum likelihood method for the V4

hypervariable regions of 16S rRNA genes for the seven hosts of class 1 integrons identified in

this study. The integron gene cassettes associated with *Alphaproteobacteria* (in blue) and

Gammaproteobacteria (in red) are grouped accordingly. Gene cassettes that were found in

Gammaproteobacteria hosts of the *Burkholderiales* order are shown in yellow.

Tables

408 **Table 1**. Comparison between the class 1 integron gene cassettes sequenced in this work and

409 annotated reference class 1 integron gene cassettes.

ORF in gene cassette	Predicted function of ORF gene product	NCBI reference gene cassette	% Nucleotide identity to NCBI reference gene cassette sequence
Cupin-domain-encoding gene	Unknown	FJ172388	99.8% nucleotide identity; 1 synonymous mutation in the coding region
dfrA5	Trimethoprim-resistant dihydrofolate reductase	X12868	100% nucleotide identity
blaOXA₁₀	Oxacillin-hydrolysing beta-lactamase	XXU37105	
qacG2	Quaternary ammonium compound efflux pump	AF327731	99.6% nucleotide identity; 2 non-synonymous mutations (S13F and A57G) in the coding region
qacF	Quaternary ammonium compound efflux pump	AF034958	99.6% nucleotide identity; 1 non-synonymous mutation (S10A) in the coding region
aadA2	Aminoglycoside (3'') adenylyltransferase conferring resistance to streptomycin and spectinomycin	X68227	In the *Rhizobacter* host: 100% nucleotide identity
			In the *Rhodocyclaceae* host: 99.6% nucleotide identity; 1 SNP in the non-coding region, 1 non-synonymous mutation (V257A) in the coding region

Associated Content

Supporting Information: Additional experimental results and methods on the assignment of taxonomy for bacterial hosts (Table S1); oligonucleotide primers (Table S2); nucleotide sequences of gene cassettes (Table S3); hosts of the cupin-domain-encoding gene cassette (Table S4); sampling location and photograph (Supplementary Figure 1); examples of fluorescence microscopy images for cell density estimation (Supplementary Figure 2); additional epicPCR amplicons (Supplementary Figure 3); nucleotide sequence alignments of conserved regions in sequenced class 1 integrons (Supplementary Figure 4) (PDF)

Author Contributions

Conceptualisation of study: TMG, QQ, SGT, MRG

Experimental investigation and validation: QQ, AP, JACS

Bioinformatic analysis: TMG, QQ, VR

Project management and supervision: MRG, SGT

Writing the original draft of the manuscript: QQ

Reviewing and editing the manuscript: All authors

Funding Sources

This work was funded by the Australian Research Council Discovery Project DP200101874 grant to MRG and SGT.

Acknowledgement

The authors thank Professor Marko Virta and his research group at the Department of Microbiology, University of Helsinki for sharing their expertise on epicPCR. We would like to acknowledge the Macquarie University Faculty of Science and Engineering Microscope Facility for access to its instrumentation and staff. We also thank Mr Richard Miller (IT Systems and HPC, Macquarie University) for his technical advice and support.

References

(1) Jain, R.; Rivera, M. C.; Moore, J. E.; Lake, J. A. Horizontal gene transfer accelerates genome innovation and evolution. Mol Biol Evol 2003, 20 (10), 1598-1602. DOI: 10.1093/molbev/msg154.

(2) Wiedenbeck, J.; Cohan, F. M. Origins of bacterial diversity through horizontal genetic transfer and adaptation to new ecological niches. FEMS Microbiol Rev 2011, 35 (5), 957-976. DOI: 10.1111/j.1574-6976.2011.00292.x.

(3) Boto, L. Horizontal gene transfer in evolution: Facts and challenges. Proc Biol Sci 2010, 277 (1683), 819-827. DOI: 10.1098/rspb.2009.1679.

(4) Beiko, R. G.; Harlow, T. J.; Ragan, M. A. Highways of gene sharing in prokaryotes. Proc Natl Acad Sci U S A 2005, 102 (40), 14332-14337. DOI: 10.1073/pnas.0504068102.

(5) Gillings, M. R.; Holley, M. P.; Stokes, H. W. Evidence for dynamic exchange of *qac* gene cassettes

between class 1 integrons and other integrons in freshwater biofilms. FEMS Microbiol Lett 2009, 296

(2), 282-288. DOI: 10.1111/j.1574-6968.2009.01646.x.

(6) Sørensen, S. J.; Bailey, M.; Hansen, L. H.; Kroer, N.; Wuertz, S. Studying plasmid horizontal

transfer *in situ*: A critical review. Nat Rev Microbiol 2005, 3 (9), 700-710. DOI: 10.1038/nrmicro1232.

(7) Gillings, M. R. Lateral gene transfer, bacterial genome evolution, and the Anthropocene. Ann N Y

Acad Sci 2017, 1389 (1), 20-36. DOI: 10.1111/nyas.13213.

(8) Brown-Jaque, M.; Calero-Cáceres, W.; Muniesa, M. Transfer of antibiotic-resistance genes via

phage-related mobile elements. Plasmid 2015, 79, 1-7. DOI: 10.1016/j.plasmid.2015.01.001.

(9) Lerminiaux, N. A.; Cameron, A. D. S. Horizontal transfer of antibiotic resistance genes in clinical

environments. Can J Microbiol 2019, 65 (1), 34-44. DOI: 10.1139/cjm-2018-0275.

(10) Hall, J. P.; Wood, A. J.; Harrison, E.; Brockhurst, M. A. Source-sink plasmid transfer dynamics

maintain gene mobility in soil bacterial communities. Proc Natl Acad Sci U S A 2016, 113 (29), 8260-

8265. DOI: 10.1073/pnas.1600974113.

(11) Zhou, H.; Beltrán, J. F.; Brito, I. L. Functions predict horizontal gene transfer and the emergence

of antibiotic resistance. Sci Adv 2021, 7 (43), eabj5056. DOI: 10.1126/sciadv.abj5056.

(12) Abe, K.; Nomura, N.; Suzuki, S. Biofilms: Hot spots of horizontal gene transfer (HGT) in aquatic

environments, with a focus on a new HGT mechanism. FEMS Microbiol Ecol 2020, 96 (5), fiaa031.

DOI: 10.1093/femsec/fiaa031.

(13) Cárdenas, J. P.; Quatrini, R.; Holmes, D. S. Genomic and metagenomic challenges and

opportunities for bioleaching: A mini-review. Res Microbiol 2016, 167 (7), 529-538. DOI:

10.1016/j.resmic.2016.06.007.

(14) Thompson, L. R.; Sanders, J. G.; McDonald, D.; Amir, A.; Ladau, J.; Locey, K. J.; Prill, R. J.;

Tripathi, A.; Gibbons, S. M.; Ackermann, G.; Navas-Molina, J. A.; Janssen, S.; Kopylova, E.; Vázquez-

Baeza, Y.; González, A.; Morton, J. T.; Mirarab, S.; Xu, Z. Z.; Jiang, L.; Haroon, M. F.; Kanbar, J.;

Zhu, Q.; Song, S. J.; Kosciolek, T.; Bokulich, N. A.; Lefler, J.; Brislawn, C. J.; Humphrey, G.; Owens, S. M.; Hampton-Marcell, J.; Berg-Lyons, D.; McKenzie, Fierer, V.; N.; Fuhrman, J. A.; Clauset, A.; Stevens, R. L.; Shade, A.; Pollard, K. S.; Goodwin, K. D.; Jansson, J. K.; Gilbert, J. A.; Knight, R.; The Earth Microbiome Project Consortium. A communal catalogue reveals Earth's multiscale microbial diversity. Nature 2017, 551 (7681), 457-463. DOI: 10.1038/nature24621.

(15) Bahram, M.; Hildebrand, F.; Forslund, S. K.; Anderson, J. L.; Soudzilovskaia, N. A.; Bodegom, P. M.; Bengtsson-Palme, J.; Anslan, S.; Coelho, L. P.; Harend, H.; Huerta-Cepas, J.; Medema, M. H.; Maltz, M. R.; Mundra, S.; Olsson, P. A.; Pent, M.; Põlme, S.; Sunagawa, S.; Ryberg, M.; Tedersoo, L.; Bork, P. Nature 2018, 560 (7717), 233-237. DOI: 10.1038/s41586-018-0386-6.

(16) Stewart, E. J. Growing unculturable bacteria. J Bacteriol 2012, 194 (16), 4151-4160. DOI: 10.1128/JB.00345-12.

(17) Douglas, G. M.; Langille, M. G. I. Current and promising approaches to identify horizontal gene transfer events in metagenomes. Genome Biol Evol 2019, 11 (10), 2750-2766. DOI: 10.1093/gbe/evz184.

(18) Seiler, E.; Trappe, K.; Renard, B. Y. Where did you come from, where did you go: Refining metagenomic analysis tools for horizontal gene transfer characterisation. PLoS Comput Biol 2019, 15 (7), e1007208. DOI: 10.1371/journal.pcbi.1007208.

(19) Ghurye, J. S.; Cepeda-Espinoza, V.; Pop, M. Metagenomic assembly: Overview, challenges and applications. Yale J Biol Med 2016, 89 (3), 353-362.

(20) Darmon, E.; Leach, D. R. Bacterial genome instability. Microbiology and molecular biology reviews: MMBR 2014, 78 (1), 1-39. DOI: 10.1128/MMBR.00035-13.

(21) Spencer, S. J.; Tamminen, M. V.; Preheim, S. P.; Guo, M. T.; Briggs, A. W.; Brito, I. L.; D, A. W.; Pitkanen, L. K.; Vigneault, F.; Virta, M. P.; Alm, E. J. Massively parallel sequencing of single cells by epicPCR links functional genes with phylogenetic markers. ISME J 2016, 10 (2), 427-436. DOI: 10.1038/ismej.2015.124.

(22) Roman, V. L.; Merlin, C.; Virta, M. P. J.; Bellanger, X. EpicPCR 2.0: Technical and
methodological improvement of a cutting-edge single-cell genomic approach. Microorganisms 2021, 9
(8), 1649. DOI: 10.3390/microorganisms9081649.

(23) Hultman, J.; Tamminen, M.; Parnanen, K.; Cairns, J.; Karkman, A.; Virta, M. Host range of
antibiotic resistance genes in wastewater treatment plant influent and effluent. FEMS Microbiol Ecol
2018, 94 (4), fiy038. DOI: 10.1093/femsec/fiy038.

(24) Qin, H.; Wang, S.; Feng, K.; He, Z.; Virta, M. P. J.; Hou, W.; Dong, H.; Deng, Y. Unraveling the
diversity of sedimentary sulfate-reducing prokaryotes (SRP) across Tibetan saline lakes using epicPCR.
Microbiome 2019, 7 (1), 71. DOI: 10.1186/s40168-019-0688-4.

(25) Sakowski, E. G.; Arora-Williams, K.; Tian, F.; Zayed, A. A.; Zablocki, O.; Sullivan, M. B.;
Preheim, S. P. Interaction dynamics and virus-host range for estuarine actinophages captured by
epicPCR. Nat Microbiol 2021, 6 (5), 630-642. DOI: 10.1038/s41564-021-00873-4.

(26) Diebold, P. J.; New, F. N.; Hovan, M.; Satlin, M. J.; Brito, I. L. Linking plasmid-based beta-
lactamases to their bacterial hosts using single-cell fusion PCR. eLife 2021, 10: e66834. DOI:
10.7554/eLife.66834.

(27) Sarekoski, A. K. The use of epicPCR to study antimicrobial resistance genes and their bacterial
hosts in human impacted environments in West Africa. University of Helsinki: 2020.

(28) Wei, Z.; Feng, K.; Wang, Z.; Zhang, Y.; Yang, M.; Zhu, Y. G.; Virta, M. P. J.; Deng, Y. High-
throughput single-cell technology reveals the contribution of horizontal gene transfer to typical
antibiotic resistance gene dissemination in wastewater treatment plants. Environ Sci Technol 2021, 55
(17), 11824-11834. DOI: 10.1021/acs.est.1c01250.

(29) Gillings, M.; Boucher, Y.; Labbate, M.; Holmes, A.; Krishnan, S.; Holley, M.; Stokes, H. W. The
evolution of class 1 integrons and the rise of antibiotic resistance. J Bacteriol 2008, 190 (14), 5095-
5100. DOI: 10.1128/JB.00152-08.

(30) Domingues, S.; Da Silva, G. J.; Nielsen, K. M. Global dissemination patterns of common gene

cassette arrays in class 1 integrons. Microbiology 2015, 161 (7), 1313-1337. DOI:

10.1099/mic.0.000099.

(31) Gillings, M. R.; Gaze, W. H.; Pruden, A.; Smalla, K.; Tiedje, J. M.; Zhu, Y. G. Using the class 1

integron-integrase gene as a proxy for anthropogenic pollution. ISME J 2015, 9 (6), 1269-1279. DOI:

10.1038/ismej.2014.226.

(32) Ghaly, T. M.; Geoghegan, J. L.; Tetu, S. G.; Gillings, M. R. The peril and promise of integrons:

beyond antibiotic resistance. Trends Microbiol 2020, 28 (6), 455-464. DOI: 10.1016/j.tim.2019.12.002.

(33) Escudero, J. A.; Loot, C.; Mazel, D. Integrons as adaptive devices. In Molecular Mechanisms of

Microbial Evolution, Rampelotto, P. H. Ed.; Springer International Publishing, 2018; 199-239.

(34) Partridge, S. R.; Tsafnat, G.; Coiera, E.; Iredell, J. R. Gene cassettes and cassette arrays in mobile

resistance integrons. FEMS Microbiol Rev 2009, 33 (4), 757-784. DOI: 10.1111/j.1574-

6976.2009.00175.x.

(35) Escudero, J. A.; Loot, C.; Nivina, A.; Mazel, D. The Integron: Adaptation on demand. Microbiol

Spectr 2015, 3 (2), MDNA3-0019-2014. DOI: 10.1128/microbiolspec.MDNA3-0019-2014.

(36) Recchia, G. D.; Stokes, H. W.; Hall, R. M. Characterisation of specific and secondary

recombination sites recognised by the integron DNA integrase. Nucleic Acids Res 1994, 22 (11), 2071-

2078. DOI: 10.1093/nar/22.11.2071.

(37) Zheng, W.; Huyan, J.; Tian, Z.; Zhang, Y.; Wen, X. Clinical class 1 integron-integrase gene - A

promising indicator to monitor the abundance and elimination of antibiotic resistance genes in an urban

wastewater treatment plant. Environ Int 2020, 135, 105372. DOI: 10.1016/j.envint.2019.105372.

(38) Jechalke, S.; Schreiter, S.; Wolters, B.; Dealtry, S.; Heuer, H.; Smalla, K. Widespread

dissemination of class 1 integron components in soils and related ecosystems as revealed by cultivation-

independent analysis. Front Microbiol 2013, 4, 420. DOI: 10.3389/fmicb.2013.00420.

(39) Pfeifer, E.; Bonnin, R. A.; Rocha, E. P. C. Phage-plasmids spread antibiotic resistance genes

through infection and lysogenic conversion. mBio 2022, 13 (5), e0185122. DOI: 10.1128/mbio.01851-

22.

(40) Waldron, L. S.; Gillings, M. R. Screening foodstuffs for class 1 integrons and gene cassettes. J Vis

Exp 2015, (100), e52889. DOI: 10.3791/52889.

(41) Dawes, F. E.; Kuzevski, A.; Bettelheim, K. A.; Hornitzky, M. A.; Djordjevic, S. P.; Walker, M. J.

Distribution of class 1 integrons with IS*26*-mediated deletions in their 3'-conserved segments in

Escherichia coli of human and animal origin. PLoS One 2010, 5 (9), e12754. DOI:

10.1371/journal.pone.0012754.

(42) Stokes, H. W.; Nesbø, C. L.; Holley, M.; Bahl, M. I.; Gillings, M. R.; Boucher, Y. Class 1 integrons

potentially predating the association with Tn*402*-like transposition genes are present in a sediment

microbial community. J Bacteriol 2006, 188 (16), 5722-5730. DOI: 10.1128/JB.01950-05.

(43) Rosewarne, C. P.; Pettigrove, V.; Stokes, H. W.; Parsons, Y. M. Class 1 integrons in benthic

bacterial communities: Abundance, association with Tn*402*-like transposition modules and evidence

for coselection with heavy-metal resistance. FEMS Microbiol Ecol 2010, 72 (1), 35-46. DOI:

10.1111/j.1574-6941.2009.00823.x.

(44) Kasai, D.; Imai, S.; Asano, S.; Tabata, M.; Iijima, S.; Kamimura, N.; Masai, E.; Fukuda, M.

Identification of natural rubber degradation gene in *Rhizobacter gummiphilus* NS21. Biosci Biotechnol

Biochem 2017, 81 (3), 614-620. DOI: 10.1080/09168451.2016.1263147.

(45) De Coster, W.; D'Hert, S.; Schultz, D. T.; Cruts, M.; Van Broeckhoven, C. NanoPack: Visualizing

and processing long-read sequencing data. Bioinformatics 2018, 34 (15), 2666-2669. DOI:

10.1093/bioinformatics/bty149.

(46) Šošic, M.; Šikic, M. Edlib: A C/C ++ library for fast, exact sequence alignment using edit distance.

Bioinformatics 2017, 33 (9), 1394-1395. DOI: 10.1093/bioinformatics/btw753.

(47) Sahlin, K.; Medvedev, P. *De novo* clustering of long-read transcriptome data using a greedy,

quality value-based algorithm. J Comput Biol 2020, 27 (4), 472-484. DOI: 10.1089/cmb.2019.0299.

(48) Sahlin, K.; Medvedev, P. Error correction enables use of Oxford Nanopore technology for

reference-free transcriptome analysis. Nat Commun 2021, 12 (1), 2. DOI: 10.1038/s41467-020-20340-

8.

(49) Vaser, R.; Sović, I.; Nagarajan, N.; Šikić, M. Fast and accurate *de novo* genome assembly from

long uncorrected reads. Genome Res 2017, 27 (5), 737-746. DOI: 10.1101/gr.214270.116.

(50) Bengtsson-Palme, J.; Hartmann, M.; Eriksson, K. M.; Pal, C.; Thorell, K.; Larsson, D. G.; Nilsson,

R. H. METAXA2: Improved identification and taxonomic classification of small and large subunit

rRNA in metagenomic data. Mol Ecol Resour 2015, 15 (6), 1403-1414. DOI: 10.1111/1755-

0998.12399.

(51) Li, W.; Godzik, A. Cd-hit: A fast program for clustering and comparing large sets of protein or

nucleotide sequences. Bioinformatics 2006, 22 (13), 1658-1659. DOI: 10.1093/bioinformatics/btl158.

(52) Neron, B.; Littner, E.; Haudiquet, M.; Perrin, A.; Cury, J.; Rocha, E. P. C. IntegronFinder 2.0:

Identification and analysis of integrons across bacteria, with a focus on antibiotic resistance in

Klebsiella. Microorganisms 2022, 10 (4), 700. DOI: 10.3390/microorganisms10040700.

(53) Quast, C.; Pruesse, E.; Yilmaz, P.; Gerken, J.; Schweer, T.; Yarza, P.; Peplies, J.; Glöckner, F. O.

The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools.

Nucleic Acids Res 2013, 41 (Database issue), D590-596. DOI: 10.1093/nar/gks1219.

(54) Guindon, S.; Dufayard, J. F.; Lefort, V.; Anisimova, M.; Hordijk, W.; Gascuel, O. New algorithms

and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0.

Syst Biol 2010, 59 (3), 307-321. DOI: 10.1093/sysbio/syq010.

(55) Cury, J.; Jove, T.; Touchon, M.; Neron, B.; Rocha, E. P. Identification and analysis of integrons

and cassette arrays in bacterial genomes. Nucleic Acids Res 2016, 44 (10), 4539-4550. DOI:

10.1093/nar/gkw319.

(56) Zhang, Y.; Lu, J.; Wu, J.; Wang, J.; Luo, Y. Potential risks of microplastics combined with superbugs: Enrichment of antibiotic resistant bacteria on the surface of microplastics in mariculture system. Ecotoxicol Environ Saf 2020, 187, 109852. DOI: 10.1016/j.ecoenv.2019.109852.

(57) Zhang, Y.; Yang, X.; Huang, L.; Zhang, R. Whole-genome analysis and description of an IMP-8-producing *Ochrobactrum anthropi*. J Glob Antimicrob Resist 2022, 29, 275-277. DOI: 10.1016/j.jgar.2022.03.016.

(58) Martini, M. C.; Quiroga, M. P.; Pistorio, M.; Lagares, A.; Centrón, D.; Del Papa, M. F. Novel environmental class 1 integrons and cassette arrays recovered from an on-farm bio-purification plant. FEMS Microbiol Ecol 2018, 94 (3), fix190. DOI: 10.1093/femsec/fix190.

(59) Gillings, M. R.; Xuejun, D.; Hardwick, S. A.; Holley, M. P.; Stokes, H. W. Gene cassettes encoding resistance to quaternary ammonium compounds: a role in the origin of clinical class 1 integrons? ISME J 2009, 3 (2), 209-215. DOI: 10.1038/ismej.2008.98.

(60) Martinez-Freijo, P.; Fluit, A. C.; Schmitz, F. J.; Grek, V. S.; Verhoef, J.; Jones, M. E. Class I integrons in Gram-negative isolates from different European hospitals and association with decreased susceptibility to multiple antibiotic compounds. J Antimicrob Chemother 1998, 42 (6), 689-696. DOI: 10.1093/jac/42.6.689.

(61) van Essen-Zandbergen, A.; Smith, H.; Veldman, K.; Mevius, D. Occurrence and characteristics of class 1, 2 and 3 integrons in *Escherichia coli*, *Salmonella* and *Campylobacter* spp. in the Netherlands. J Antimicrob Chemother 2007, 59 (4), 746-750. DOI: 10.1093/jac/dkl549.

(62) Dunwell, J. M.; Culham, A.; Carter, C. E.; Sosa-Aguirre, C. R.; Goodenough, P. W. Evolution of functional diversity in the cupin superfamily. Trends Biochem Sci 2001, 26 (12), 740-746. DOI: 10.1016/s0968-0004(01)01981-8.

(63) Krucinska, J.; Lombardo, M. N.; Erlandsen, H.; Estrada, A.; Si, D.; Viswanathan, K.; Wright, D. L. Structure-guided functional studies of plasmid-encoded dihydrofolate reductases reveal a common

mechanism of trimethoprim resistance in Gram-negative pathogens. Commun Biol 2022, 5 (1), 459.

DOI: 10.1038/s42003-022-03384-y.

(64) Hall, L. M.; Livermore, D. M.; Gur, D.; Akova, M.; Akalin, H. E. OXA-11, an extended-spectrum

variant of OXA-10 (PSE-2) beta-lactamase from *Pseudomonas aeruginosa*. Antimicrob Agents

Chemother 1993, 37 (8), 1637-1644. DOI: 10.1128/AAC.37.8.1637.

(65) Evans, B. A.; Amyes, S. G. OXA β-lactamases. Clin Microbiol Rev 2014, 27 (2), 241-263. DOI:

10.1128/CMR.00117-13.

(66) Gaze, W. H.; Zhang, L.; Abdouslam, N. A.; Hawkey, P. M.; Calvo-Bado, L.; Royle, J.; Brown, H.;

Davis, S.; Kay, P.; Boxall, A. B.; Wellington, E. M. H. Impacts of anthropogenic activity on the ecology

of class 1 integrons and integron-associated genes in the environment. ISME J 2011, 5 (8), 1253-1261.

DOI: 10.1038/ismej.2011.15.

(67) Antunes, P.; Machado, J.; Peixe, L. Dissemination of *sul3*-containing elements linked to class 1

integrons with an unusual 3' conserved sequence region among *Salmonella* isolates. Antimicrob Agents

Chemother 2007, 51 (4), 1545-1548. DOI: 10.1128/AAC.01275-06.

(68) Chang, C. Y.; Chang, L. L.; Chang, Y. H.; Lee, T. M.; Chang, S. F. Characterisation of drug

resistance gene cassettes associated with class 1 integrons in clinical isolates of *Escherichia coli* from

Taiwan, ROC. J Med Microbiol 2000, 49 (12), 1097-1102. DOI: 10.1099/0022-1317-49-12-1097.

(69) Clark, N. C.; Olsvik, O.; Swenson, J. M.; Spiegel, C. A.; Tenover, F. C. Detection of a

streptomycin/spectinomycin adenylyltransferase gene (*aadA*) in *Enterococcus faecalis*. Antimicrob

Agents Chemother 1999, 43 (1), 157-160. DOI: 10.1128/AAC.43.1.157.

(70) Binh, C. T.; Heuer, H.; Kaupenjohann, M.; Smalla, K. Diverse *aadA* gene cassettes on class 1

integrons introduced into soil *via* spread manure. Res Microbiol 2009, 160 (6), 427-433. DOI:

10.1016/j.resmic.2009.06.005.

(71) Yoon, M. H.; Ten, L. N.; Im, W. T.; Lee, S. T. *Methylibium fulvum* sp. nov., a member of the

Betaproteobacteria isolated from ginseng field soil, and emended description of the genus *Methylibium*.

Int J Syst Evol Microbiol 2007, 57 (Pt 9), 2062-2066. DOI: 10.1099/ijs.0.64909-0.

(72) Stackebrandt, E.; Verbarg, S.; Frühling, A.; Busse, H. J.; Tindall, B. J. Dissection of the genus

Methylibium: Reclassification of *Methylibium fulvum* as *Rhizobacter fulvus* comb. nov., *Methylibium*

aquaticum as *Piscinibacter aquaticus* gen. nov., comb. nov. and *Methylibium subsaxonicum* as

Rivibacter subsaxonicus gen. nov., comb. nov. and emended descriptions of the genera *Rhizobacter* and

Methylibium. Int J Syst Evol Microbiol 2009, 59 (Pt 10), 2552-2560. DOI: 10.1099/ijs.0.008383-0.

(73) Wei, L.; Si, M.; Long, M.; Zhu, L.; Li, C.; Shen, X.; Wang, Y.; Zhao, L.; Zhang, L. *Rhizobacter*

bergeniae sp. nov., isolated from the root of *Bergenia scopulosa*. Int J Syst Evol Microbiol 2015, 65

(Pt 2), 479-484. DOI: 10.1099/ijs.0.070318-0.

(74) Jin, L.; Ko, S. R.; Ahn, C. Y.; Lee, H. G.; Oh, H. M. *Rhizobacter profundi* sp. nov., isolated from

freshwater sediment. Int J Syst Evol Microbiol 2016, 66 (5), 1926-1931. DOI: 10.1099/ijsem.0.000962.

(75) Imai, S.; Yoshida, R.; Endo, Y.; Fukunaga, Y.; Yamazoe, A.; Kasai, D.; Masai, E.; Fukuda, M.

Rhizobacter gummiphilus sp. nov., a rubber-degrading bacterium isolated from the soil of a botanical

garden in Japan. J Gen Appl Microbiol 2013, 59 (3), 199-205. DOI: 10.2323/jgam.59.199.

(76) Linh, D. V.; Gibu, N.; Tabata, M.; Imai, S.; Hosoyama, A.; Yamazoe, A.; Kasai, D.; Fukuda, M.

Complete genome sequence of natural rubber-degrading, Gram-negative bacterium, *Rhizobacter*

gummiphilus strain NS21[T]. Biotechnol Rep (Amst) 2019, 22, e00332. DOI:

10.1016/j.btre.2019.e00332.

(77) Goto, M.; Kuwata, H. *Rhizobacter daucus* gen. nov., sp. nov., the causal agent of carrot bacterial

gall. Int J Syst Evol Microbiol 1988, 38 (3), 233-239. DOI: 10.1099/00207713-38-3-233.

(78) Watanabe, M.; Igarashi, K.; Kato, S.; Kamagata, Y.; Kitagawa, W. Critical effect of H_2O_2 in the

agar plate on the growth of laboratory and environmental strains. Microbiol Spectr 2022, e0333622.

DOI: 10.1128/spectrum.03336-22.

(79) Tanaka, T.; Kawasaki, K.; Daimon, S.; Kitagawa, W.; Yamamoto, K.; Tamaki, H.; Tanaka, M.;

Nakatsu, C. H.; Kamagata, Y. A hidden pitfall in the preparation of agar media undermines

microorganism cultivability. Appl Environ Microbiol 2014, 80 (24), 7659-7666. DOI:

10.1128/AEM.02741-14.

(80) Bengtsson-Palme, J.; Jonsson, V.; Hess, S. What is the role of the environment in the emergence

of novel antibiotic resistance genes? A modeling approach. Environ Sci Technol 2021, 55 (23), 15734-

15743. DOI: 10.1021/acs.est.1c02977.

(81) Kurland, C. G.; Canback, B.; Berg, O. G. Horizontal gene transfer: A critical view. Proc Natl Acad

Sci U S A 2003, 100 (17), 9658-9662. DOI: 10.1073/pnas.1632870100.

(82) Ghaly, T. M.; Gillings, M. R.; Penesyan, A.; Qi, Q.; Rajabal, V.; Tetu, S. G. The natural history of

integrons. Microorganisms 2021, 9 (11), 2212. DOI: 10.3390/microorganisms9112212.

A

intI1   *attI1*   ORF   *attC*   ORF   *attC*   Δ*qacE*   *sul1*   3'-CS

5'-CS

B

R519   16S rRNA gene fragment   R926

V4   V5

intI1   *attI1*   Variable integron gene cassette array   *qacE*

*intI1*_outer

Bridging primer

C

AP27_short

intI1   *attI1*   *qacE*   V4

*intI1*_nested