1 **Caloric restriction prevents inflammation and insulin dysfunction in middle-aged**

2 **ovariectomized mice**

Authors: Leticia Roberta Leme Sapatini[1†], Bruno Calsa[2†], Lais Jorge Marim[3], Júlia

Venturini Helaehil[1], Gabriela Bortolança Chiarotto[1] and Maria Esméria Corezola do

Amaral[1*]

[1]Graduate Program in Biomedical Sciences, University Center of Hermínio Ometto

Foundation, Araras, SP, Brazil

[2] Fetal Programming and Hydroelectrolyte Metabolism and Experimental Surgery,

Department of Internal Medicine, Faculty of Medical Sciences, State University of

Campinas, São Paulo, Brazil.

[3]Physiotherapy College, University Center of Hermínio Ometto Foundation, FHO,

Araras, SP, Brazil

[†]Co-first authors: Leticia Sapatini and Bruno Calsa

*Corresponding author:

Maria Esméria Corezola do Amaral

Graduate Program in Biomedical Science, University Center of Hermínio Ometto

Foundation, Av. Maximiliano Barutto nº 500, Jardim Universitário, 13607-339, Araras,

SP, Brazil

Tel: +55 (19) 3543 1474; Fax: +55 (19) 35 1412; E-mail: esmeria@fho.edu.br

Abstract

Loss of ovarian function is associated with increased visceral fat. In this study, we aimed to study the effects of caloric restriction (CR) on metabolism in ovariectomized mice. Female, 8–12-month-old mice were divided into three groups: OVX (ovariectomized mice), OVXR (40% CR) and Sham. CR increased insulin sensitivity and glucose tolerance. AMPK phosphorylation was observed in the liver of OVXR mice. CR also increased hepatic cholesterol and triglyceride levels. The reductions in the level of TBARS in the serum and liver and of H_2O_2 in the liver of OVXR mice suggested alterations in the redox state of the liver. Although expression of catalase protein was reduced by CR, expression of superoxide dismutase was not altered by CR. Although interleukin IL-6 and IL-10 levels in OVXR mice were similar to those in Sham mice, macrophage infiltration was reduced in OVXR mice. OVXR mice had increased sirtuin1 levels and decreased sirtuin3 levels in the liver. In conclusion, CR improved the condition of ovariectomized mice by reducing adiposity and increasing insulin sensitivity and glucose tolerance through a mechanism that may involve AMPK.

Keywords: caloric restriction, obesity, inflammation, menopause

Introduction

Numerous studies have shown that menopause is associated with unfavorable changes in body composition, such as increased abdominal fat deposition, and poor overall health [1, 2]. The deleterious effects of obesity are diverse and include various metabolic diseases, such as diabetes mellitus, and cardiovascular diseases, like coronary heart disease, heart attack, stroke, and hypertension, which adversely impact the quality of life and carry an increased risk of premature death [3-6]. However, the relationship

between menopause and obesity is complex. Sex hormones fluctuate with reproductive changes, such as menarche, pregnancy, and menopause, which may play an important role in the expansion of adipose tissue [1]. Hormones impact adipose tissue, and estrogens regulate glucose homeostasis and insulin sensitivity in several ways, by altering insulin production and secretion [7, 8], insulin-mediated glucose uptake in peripheral tissues, mitochondrial function [9], and the expression of inflammatory markers [10]. Given the apparent interaction between low estrogen concentrations and obesity in altering glucose and lipid homeostasis, increasing attention has been given to developing dietary strategies that preserve homeostasis during menopause.

Caloric restriction (CR) is one strategy that can reduce body weight, total body fat, and visceral fat, similar to exercise. CR is a non-pharmacological dietary intervention in which the average daily calorie intake is reduced without malnutrition or deprivation of essential nutrients. Animals on a calorie-restricted diet showed a significant (50%) increase in life expectancy [11]. Animal studies showed favorable effects of short-term CR on cardiometabolic changes, liver changes, and cellular senescence [12, 13]. In humans, experimental and observational studies have revealed that CR can reduce the incidence of cardiovascular disease, diabetes, dementia, frailty, and cancer [14, 15]. These observed benefits are related to CR-mediated activation of regulatory proteins in metabolic pathways, such as AMP-activated protein kinase (AMPK) and sirtuins (SIRTs). AMPK is a key sensor and regulator of glucose metabolism that activates SIRT1 deacetylase in response to changes in the redox state (NAD+/NADH and AMP/ATP ratios) of the cell [16]. Sirtuins are a group of highly conserved NAD+-dependent histones and protein deacetylases and/or ADP-ribosyl transferases that play important roles in various biological processes [17, 18, 8, 19]. SIRT1 functions as a metabolic/energy sensor that modulates the activity and/or gene

expression of several crucial transcription factors and transcription coactivators involved in metabolic homeostasis [20, 18, 21]. Reduced levels of NAD+ and SIRT1 activity have been detected under energy-overloaded metabolic profiles, such as a hyperlipidic diet [22, 23].

In contrast, CR increased NAD+ levels and induced activation of SIRT1 [22, 23]. SIRT1 functions as an essential regulator of hepatic lipid metabolism through SREBP-1c/ChREBP-dependent control of lipogenesis by inhibiting lipid synthesis and increasing PPARα/PGC-1α-dependent β-fatty acid oxidation [24]. SIRT1 regulates oxidative stress by increasing the transcriptional activity of antioxidant enzymes, such as superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPX) [24]. Activation of SIRT1 reduces nitric oxide synthase and lipid peroxidation products, which has an anti-oxidative stress effect in the liver [24]. SIRT3 is a mitochondrial protein that integrates cellular energy metabolism and plays an important role in preventing metabolic syndrome [25]. SIRT3 knockout mice showed characteristics of reduced mitochondrial metabolism, such as reduced fatty acid catabolism [25]. SIRT3 expression was reduced in male mice fed a high-fat diet [25, 26]. However, the consistency and reproducibility of results across investigations are varied, and key questions about the timing, duration, modality, and adverse effects of CR, as well as the influence of genotype and diet composition, remain [27]. Given the low adherence of humans to difficult-to-follow diets, research on dietary interventions should be conducted. This study examined the impact of 40% CR on the liver of middle-aged mice who underwent ovariectomy. CR was expected to have a positive effect on metabolism. The results showed that CR was associated with improvements in both metabolism, mainly glycemic metabolism, and inflammatory markers.

Materials and methods

Experimental animals and model establishment

The project was approved by the Committee on Ethics in Animal Use (CEUA) under protocol #009/2020. Eighteen C57BL/6J mice (8–12 months old, weighing ~20 g) were housed in individual cages at a constant temperature (22 °C) under a 12 h light/dark cycle with water and commercial chow *ad libitum*. For the ovariectomy, the mice were anesthetized by intraperitoneal injection with ketamine (50 mg/kg) and xylazine (5 mg/kg). After shaving the inguinal region, an abdominal incision was made, and the ovaries were located and removed. After surgery, the animals were placed in a heated chamber to avoid hypothermia. The animals in the control group (Sham) underwent the same procedure except the ovaries were not removed. At 16 weeks after surgery, the ovariectomy model mice were separated into two groups: the OVX (ovariectomized) and OVXR (ovariectomized plus 40% CR) groups. Sham and OVX mice were fed a commercial isocaloric diet *ad libitum*, and OVXR mice received 60% of the chow consumed by the OVX mice for 30 days, corresponding to 40% CR. All mice had free access to water throughout the experiment. The body weights of the animals were measured weekly throughout the experiment.

Intraperitoneal glucose tolerance test

The intraperitoneal glucose tolerance test (ipGTT) was administered 5 days before euthanasia after a 6 h fast. For the test, glucose (2 g/kg body weight) was injected intraperitoneally. Blood samples were collected before glucose overload (time zero), and at 15, 30, 60, and 120 min after glucose infusion. Blood glucose was determined using reagent strips and an Abbott® glucometer as previously described [28].

Intraperitoneal insulin tolerance test

The intraperitoneal insulin tolerance test (ipITT) was administered 3 days before

euthanasia after a 6 h fast. Insulin (regular crystalline, 0.75 U/kg body weight) was

administered intraperitoneally. Blood samples were collected before insulin injection

(time zero) and at 5, 10, 15, 20, 25, and 30 min after injection. Blood glucose was

determined using an Abbott glucometer and reagent strips. The rate constant for glucose

disappearance (Kitt) was calculated using the formula In2/t1/2. The half-life (t1/2) of

serum glucose was calculated from the slope of the minimum regression curve in the

linear phase of plasma glucose decline [29].

Homeostasis model assessments

Homeostasis model assessment of insulin resistance (HOMA-IR), insulin sensitivity

(HOMA%S), and β-cell function (HOMA%β) were calculated using the HOMA

calculator from the University of Oxford (www.dtu.ox.ac.uk/homacalculator).

Fat pad and Lee indexes

The perirenal adipose tissue weight and body weight were assessed, and the adipose

tissue index was calculated by dividing adipose tissue weight by body weight. The Lee

index was calculated using the following equation: $\sqrt[3]{}$body weight (g)÷nose to anus

length (cm). .

Biochemical assessments

Serum glucose, triglycerides, cholesterol, HDL, and total protein were determined using

commercial kits (Laborlab, Guarulhos, SP, Brazil). LDL was calculated using the

Friedewald equation, and VLDL was calculated by dividing triglycerides by 5. Insulin

serum was measured using ELISA (Rat/Mouse Insulin ELISA, #EZRMI-13K, Millipore

St. Charles, MO, USA)). Hepatic glycogen content was measured using a colorimetric

method [30]. Total liver lipids were extracted according to the method of Folch et al.

[31].

Preparation of liver homogenates for oxidative stress biomarker analysis

Liver samples (~50 mg) were homogenized in Krebs buffer (118 mM NaCl, 25 mM NaHCO$_3$, 1.2 mM KH$_2$PO$_4$, 4.7 mM KCl, 1.2 mM MgSO$_4$, 1.25 mM CaCl$_2$, 10 mM glucose, and 10 mM HEPES, pH 7.4). The obtained tissue homogenate was used for the determination of 2-thiobarbituric acid reactive substances (TBARS) and hydrogen peroxide (H$_2$O$_2$) levels. Total protein was quantified using the Biuret method, with bovine serum albumin as a standard, and measured at 540 nm.

Determination of hepatic H$_2$O$_2$

H$_2$O$_2$ levels in hepatic tissue were determined using the Amplex Red method. Briefly, 20 µL of homogenate was incubated with horseradish peroxidase (HRP, 1 U/mL) and Amplex Red (50 µM) at 37 °C for 30 min. Then, the absorbance was read at 560 nm. The H$_2$O$_2$ concentration was calculated by referring to a standard curve of known concentrations and normalized to the protein concentration.

Hepatic and systemic TBARS

Serum and liver TBARS contents were determined as described previously [32]. Briefly, 50 µL of serum or liver homogenate was mixed with 10 µL of 0.1 M Butylated Hydroxytoluene, 400 µL of 1% 2-thiobarbituric acid (TBA; Merck, St. Charles, MO, USA), and 200 µL of 20% phosphoric acid and heated at 100 °C for 15 min. After cooling, 1.5 mL of 1-butanol was added, and the optical density of the supernatant was determined at 532 nm.

Determination of hepatic n-acetylglucosaminidase

n-Acetylglucosaminidase (NAG) activity was measured to determine liver macrophagic infiltration. An aliquot (3.0 µL) of liver homogenate (0.08 M NaPO$_4$) was mixed with 30 µL of p-nitrophenyl-2-acetamide-β-D-glucopyranoside (Sigma-Aldrich) and diluted

in 50 µL of 50 mM citrate buffer. Finally, 50 µL of 0.2 M glycine was added, and the

absorbance was read at 405 nm.

Histological analysis of liver and adipose tissue

Liver and adipose tissues were fixed with 10% formalin and dehydrated. Then, small

fragments of the tissues were embedded in paraffin (Paraplast; Sigma, St. Louis, MO,

USA) using standard procedures. Slices (5 µm thick) were stained with hematoxylin-

eosin (HE), and images were acquired using a microscope (Leica DM 2000, Germany)

with LAS v.4.1 software. For each mouse, 40 images were obtained at 400×

magnification. The area (mm^2) containing adipocytes in each image was quantified

using the ImageJ software (National Institutes of Health, Bethesda, Maryland, USA).

Immunoblotting

Liver tissue fragments were homogenized in protein extraction buffer (10 mM EDTA,

100 mM Tris-HCL, 10 mM sodium pyrophosphate, 100 mM sodium fluoride, 100 mM

sodium orthovanadate, 10 mM phenylmethylsulfonyl fluoride, and 0.1 mg/mL

aprotinin). Protein content was determined according to the Biuret method. The samples

were mixed with Laemmli buffer (0.1% bromophenol blue, 1 M sodium phosphate,

50% glycerol, and 10% SDS). Total protein (50 µg) was separated by electrophoresis.

Then, the separated proteins were transferred to nitrocellulose membranes. After

transfer, the membranes were incubated at 4 °C, with shaking, overnight with the

following primary antibodies: IL-6, IL-10, TGF-β1, SIRT3, pIR, and IR (Santa Cruz,

Dallas, USA; 1:250), and SIRT1, pAMPK, and AMPK (Cell Signaling Danvers,

Massachusetts, USA; 1:1000). The membranes were then incubated with an HRP-

secondary antibody (Santa Cruz; Dallas, USA; 1:10,000), with shaking, followed by

development with chemiluminescent reagents (SuperSignal West Pico PLUS, Thermo

Scientific, Rockford, IL, USA) and bands intensities were determined by densitometry

197 scanning using the ImageJ software (National Institutes of Health, Bethesda, Maryland,

198 USA)).

Statistical analysis

200 Data are mean ± standard error of the mean. All data were analyzed using the Shapiro–

201 Wilk Normality test. For comparisons between experimental groups, one-way ANOVA

202 and Tukey's post-test were used. All statistical and graphical analyses were performed

203 using GraphPad Prism software (version 8.0). Significance was set at $p < 0.05$.

Results

Ovariectomy and caloric restriction characterization

The timeline of the experiment is shown in Figure 1A. After ovariectomy, the animals

were observed for 4 months. The animals in the OVXR group remained on a restricted

diet for the first month. The body weights of the animals were monitored throughout the

experiment and are presented in Figure 1B, which shows evident weight gain in the

OVX group compared to in the Sham group and marked weight loss during CR in the

OVXR group compared to in the Sham and OVX groups. The Lee index of the mice

was lower in the OVXR group than in the Sham and OVX groups (Figure 1C). Analysis

of the uterine weight of the mice showed evident uterine atrophy following ovariectomy

(Figure 1D).

Mice in the OVXR group had lower perirenal adipose tissue weight than mice in

the OVX group (Figure 1E); however, the difference was not statistically different.

OVX mice showed adipose tissue weight gain when compared to mice in the other

groups. HE staining of liver and adipose tissue (Figure 1F) showed slight steatosis in

OVXR group mice when compared to the Sham and OVX group mice. There was an

evident reduction in adipocyte diameter in OVXR mice when compared to that in OVX

mice.

Effects on glucose, insulin, and lipid metabolism

The GTT and ITT results (Figures 2A and 2B, respectively) showed a reduced area

under the curve for OVXR mice (Figure 2C) compared to that for OVX mice and an

increased Kitt for OVXR mice compared to that for OVX mice (Figure 2D). These

results suggest greater glucose tolerance and increased insulin sensitivity of OVXR

mice than of OVX mice. The HOMA-IR (Figure 2E), HOMA%S (Figure 2F), and

HOMA%B (Figure 2G) indexes indicated increased insulin sensitivity and beta cell

preservation in the OVXR mice compared to in the OVX mice. Consistent with these

results, liver glycogen levels were similar in OVXR and Sham mice but lower in OVX

than in OVXR mice (Table 1). No difference in the liver index was observed among the

groups. The reduced blood glucose levels (Figure 3F) and insulinemia (Figure 3G) in

the OVXR group compared to that in the Sham and OVX groups suggest improvement

in glycemic homeostasis under CR. Similar serum protein concentrations (Figure 3H) in

all groups indicate good nutritional status.

Serum cholesterol levels were lower in the OVXR group than in the OVX group

(Figure 3A). Serum triglycerides (Figure 3B) and VLDL (Figure 3E) were similar

among the groups. LDL-cholesterol (Figure 3D) was higher in the OVX group than in

the Sham group. Normalization of the HDL-cholesterol fraction was observed for

OVXR mice compared to Sham mice (Figure 3C), and HDL-cholesterol was lower in

OVX mice than in Sham mice. Hepatic cholesterol levels were similar among the

groups, but hepatic triglycerides were higher in OVXR mice than in Sham and OVX

mice (Table 1). This result is mirrored by the liver tissue histology, which showed lipid

vacuoles in the livers of OVXR mice (Figure 1F), suggesting metabolic adaptation to

the CR diet. Similar adipocyte diameter (Table 1) was observed in the OVXR and Sham

groups, whereas adipocyte diameter was increased in the OVX group when compared to

that in the OVXR group.

Oxidative stress balance

CAT protein expression was higher in OVX mice than in OVXR mice (Figure 4A), and

was similar in OVXR and Sham mice. SOD-2 protein expression (Figure 4B) was

similar among the groups. Representative bands of CAT, SOD-2, and beta-actin protein

are shown in Figure 4C. Analysis of TBARS content in the serum (Figure 4D) and liver

(Figure 4E) and H_2O_2 levels (Figure 4F) showed lower levels in the OVXR group than

in the OVX group, suggesting reduced oxidative stress in the OVXR group compared to

that in the OVX and Sham groups.

Hepatic inflammatory balance

Macrophage infiltration (NAG activity) in liver tissue was similar in the OVXR and

Sham groups (Table 1) but was higher in the OVX group than in the Sham and OVXR

groups. Levels of IL-6, a pro-inflammatory cytokine, were higher in OVX mice than in

Sham mice (Figure 5A), whereas levels of IL-10, anti-inflammatory cytokine, were

lower in OVX mice than in Sham mice (Figure 5B). Similar TGF protein expression

was observed in all groups (Figure 5C).

Hepatic metabolism

SIRT1 protein expression in the liver tissue was higher in OVXR mice than in Sham

mice (Figure 5D). In contrast, SIRT3 (Figure 5E) and IR (Figure 5G) protein levels

were lower in OVXR and OVX mice than in Sham mice. AMPK protein expression was

higher in OVXR mice than in OVX and Sham mice (Figure 5F). Representative protein

bands of AMPK, pAMPK, IL-6, IL-10, IR, pIR, SIRT1, SIRT3, TGF, and beta-actin are

shown in Figure 5H.

Discussion

CR is a well-accepted dietary intervention that can restore glucose homeostasis and

promote weight loss [33]. In this study, we expected that CR would ameliorate the

metabolic dysfunction and inflammation induced by ovariectomy. We observed weight

loss and lower adiposity in ovariectomized mice under CR. These mice also had higher

insulin sensitivity and normalized lipid profiles compared to the other mice. This can be

attributed due a CR-induced homeostatic effect on oxidative stress and inflammatory

balance in hepatic tissue.

Estrogen exerts protective effects on metabolism by regulating feeding behavior,

glucose and lipid uptake, and the insulin pathway [34]. The reduction in estrogen level

during menopause can lead to metabolism disruption, causing insulin resistance. In this

study, CR decreased fasting glucose and insulin levels. Corroborating these findings,

OVXR mice presented higher insulin sensitivity and glucose tolerance as evidenced by

the ipGTT and ipITT results and the HOMA index. These beneficial changes in

glycemic homeostasis can be attributed to increased expression of pAMPK/AMPK, but

not pIR/IR, in the liver. AMPK controls glycemia and gluconeogenesis and is an

important treatment target in diabetes [35]. Although the cholesterol fractions and

triglycerides were unchanged, CR reduced total serum cholesterol.

In previous studies, ovariectomized mice displayed a nonalcoholic fatty liver

disease (NAFLD) phenotype [36, 37]. Although we did not observe any changes in the

cholesterol and triglyceride contents in the livers of OVX mice, triglycerides were

increased in the liver of OVXR animals. Activation of AMPK inhibits hepatic

lipogenesis by inactivating acetyl CoA carboxylase [38], resulting in fat oxidation [39].

Interestingly, AMPK expression induced hepatic lipid accumulation and fatty acid

oxidation [35]. In this study, hepatic pAMPK/AMPK expression was significantly

increased in OVXR mice, suggesting its involvement in hepatic lipid metabolism. This

hepatic lipid storage in OVXR mice may be a transient metabolic process related to the

restricted diet. Under CR, the OVXR mice used fat as an energy source to maintain

cellular metabolism, which may explain the lipid accumulation in the liver. The

mechanism may involve rapid response of AMPK phosphorylation promoting uptake of

fatty acids stored in the liver [39, 35]. Our histological observations showed mild

hepatic steatosis and a reduced adipocyte area following CR, corroborating the proposed

mechanism. In a previous study, CR animals presented increased serum concentrations

of free fatty acids and SIRT1, which may indicate mobilization of triglycerides by

SIRT1 under CR [40].

Lipid accumulation increases oxidative stress and inflammation. This can

exacerbate release of reactive oxygen species and pro-inflammatory cytokines, such as

tumor-necrosis factor (TNF)-α and IL-6, in liver and adipose tissue. Previous studies

have reported an increase in IL-6 in menopause [41, 42]. After menopause, levels of the

anti-inflammatory cytokines IL-4, IL-10, and IL-12 are increased, which is related to an

increase in TNF-α, as a compensatory mechanism [43]. In our study, OVX mice showed

inflammatory effects, evidenced by an increase in IL-6 and a reduction in IL-10, due to

the reduction in estrogen levels. The levels of these cytokines were unchanged in

OVXR mice. Although TGF is an inflammatory factor that induces fibrosis, its

expression was unaffected by ovariectomy. NAG is an indicator of macrophage activity

and inflammation. The reduction in NAG level under CR suggests the anti-

inflammatory effect of CR. NAG concentrations have been linked to obesity and

diabetes [44].

In NAFLD, SIRT1 activation plays a beneficial role by decreasing oxidative

stress and inducing deacetylation of PGC-1α, leading to reduced pro-inflammatory

cytokine levels. Studies have shown that ovariectomy induced unfavorable

cardiovascular changes and that estrogen treatment activated SIRT1 and promoted

cardiovascular protection [45]. In mice, ovariectomy and SIRT3 depletion impaired the

antioxidative system response and increased lipid damage, body weight gain, and the

levels of oxidative stress-inducing genes [46]. Our findings showed that ovariectomy

increased SIRT1 and decreased SIRT3, suggesting these proteins function in a

regulatory mechanism to maintain normal metabolism.

The reductions in serum and liver TBARS contents and liver H_2O_2 levels in the

OVXR mice suggest changes in the redox state of the liver as a protective mechanism

against oxidative damage. The literature supports an increase in TBARS content

following ovariectomy, corroborating the findings of the present study [47]. The

reduction in TBARS contents in the OVXR group may explain the reduction in CAT

and the similar SOD expression levels in the present study, suggesting a balance of

mechanisms to combat oxidative stress under hypoestrogenism. The activity levels of

the antioxidant enzymes SOD, CAT, and GPX were significantly diminished in obesity

[48], and oxidative stress was enhanced as body weight increased in post-menopausal

women [49]. Here, we aimed to understand the specific functions of CR in the

"metabolic network" of tissues involved in whole-body glucose and lipid metabolism in

ovariectomized mice. Notably, while CR benefitted the ovariectomized mice by

reducing adiposity and increasing insulin sensitivity and glucose tolerance, suggesting a

balance in oxidative stress, it induced negative changes in lipid metabolism in the liver,

which was associated with changes in the protein expression of SIRT1 and SIRT3.

Finally, the results of our study suggest that ovariectomy can accelerate metabolic

changes from tissue damage to the body and that CR was able to restore, at least in part,

glycemic homeostasis.

Author statements

Acknowledgments

We thank Ana Cristina Pires Menegheti, Bruno Alves Cia, Lucas Orzari, Mateus

Eduardo Bortolanza da Silva, and Renata Barbieri for technical assistance.

Competing interests' statement

No conflicts of interest, financial or otherwise, are declared by the authors.

Author contribution statement

Sapatini LRL: Data curation, Investigation, Visualization, Writing – original draft.

Calsa B: Data curation, Formal analysis, Investigation, Validation, Visualization,

Writing – original draft, review and editing. **Marim LJ:** Investigation. **Helaehil JV:**

Data curation, Investigation. **Amaral MEC:** Conceptualization, Data curation, Formal

analysis, Funding acquisition, Investigation, Project administration, Resources,

Supervision, Validation, Visualization, Writing – review & editing.

Funding statement

This work was supported by the Herminio Ometto Foundation.

Data availability

Data generated or analyzed during this study are available form the corresponding

author upon reasonable request.

Preprint or repository availability

bioRxiv preprint doi: https://doi.org/10.1101/2023.02.09.527836

References

1. Davis SR, Castelo-Branco C, Chedraui P, Lumsden MA, Nappi RE, Shah D et al. (2012) Understanding weight gain at menopause. Climacteric 15:419-29.doi:10.3109/13697137.2012.707385.

2. Qian SW, Liu Y, Wang J, Nie JC, Wu MY, Tang Y et al. (2016) BMP4 Cross-talks With Estrogen/ERalpha Signaling to Regulate Adiposity and Glucose Metabolism in Females. EBioMedicine 11:91-100.doi:10.1016/j.ebiom.2016.07.034.

3. Camporez JP, Jornayvaz FR, Lee HY, Kanda S, Guigni BA, Kahn M et al. (2013) Cellular mechanism by which estradiol protects female ovariectomized mice from high-fat diet-induced hepatic and muscle insulin resistance. Endocrinology 154:1021-8.doi:10.1210/en.2012-1989.

4. Nappi RE, Kokot-Kierepa M (2012) Vaginal Health: Insights, Views & Attitudes (VIVA) - results from an international survey. Climacteric 15:36-44.doi:10.3109/13697137.2011.647840.

5. Neto NI, Rodrigues ME, Hachul AC, Moreno MF, Boldarine VT, Ribeiro EB et al. (2015) A Hyperlipidic Diet Combined with Short-Term Ovariectomy Increases Adiposity and Hyperleptinemia and Decreases Cytokine Content in Mesenteric Adipose Tissue. Mediators Inflamm 2015:923248.doi:10.1155/2015/923248.

6. Tominaga K, Yamauchi A, Egawa T, Tanaka R, Kawahara S, Shuto H et al. (2011) Vascular dysfunction and impaired insulin signaling in high-fat diet fed ovariectomized mice. Microvasc Res 82:171-6.doi:10.1016/j.mvr.2011.06.004.

7. Choi SB, Jang JS, Park S (2005) Estrogen and exercise may enhance beta-cell function and mass via insulin receptor substrate 2 induction in ovariectomized diabetic rats. Endocrinology 146:4786-94.doi:10.1210/en.2004-1653.

8. Santos L, Escande C, Denicola A (2016) Potential Modulation of Sirtuins by Oxidative Stress. Oxid Med Cell Longev 2016:9831825.doi:10.1155/2016/9831825.

9. Hamilton DJ, Minze LJ, Kumar T, Cao TN, Lyon CJ, Geiger PC et al. (2016) Estrogen receptor alpha activation enhances mitochondrial function and systemic metabolism in high-fat-fed ovariectomized mice. Physiol Rep 4.doi:10.14814/phy2.12913.

10. Straub RH (2007) The complex role of estrogens in inflammation. Endocr Rev
28:521-74.doi:10.1210/er.2007-0001.

11. Mattison JA, Colman RJ, Beasley TM, Allison DB, Kemnitz JW, Roth GS et al.
(2017) Caloric restriction improves health and survival of rhesus monkeys. Nat
Commun 8:14063.doi:10.1038/ncomms14063.

12. Li N, Guenancia C, Rigal E, Hachet O, Chollet P, Desmoulins L et al. (2016) Short-
term moderate diet restriction in adulthood can reverse oxidative, cardiovascular and
metabolic alterations induced by postnatal overfeeding in mice. Sci Rep
6:30817.doi:10.1038/srep30817.

13. Yzydorczyk C, Li N, Rigal E, Chehade H, Mosig D, Armengaud JB et al. (2019)
Calorie Restriction in Adulthood Reduces Hepatic Disorders Induced by Transient
Postnatal Overfeeding in Mice. Nutrients 11.doi:10.3390/nu11112796.

14. Al-Regaiey KA (2016) The effects of calorie restriction on aging: a brief review.
Eur Rev Med Pharmacol Sci 20:2468-73

15. Weiss EP, Fontana L (2011) Caloric restriction: powerful protection for the aging
heart and vasculature. Am J Physiol Heart Circ Physiol 301:H1205-
19.doi:10.1152/ajpheart.00685.2011.

16. Fulco M, Sartorelli V (2008) Comparing and contrasting the roles of AMPK and
SIRT1 in metabolic tissues. Cell Cycle 7:3669-79.doi:10.4161/cc.7.23.7164.

17. Finkel T, Deng CX, Mostoslavsky R (2009) Recent progress in the biology and
physiology of sirtuins. Nature 460:587-91.doi:10.1038/nature08197.

18. Li X (2013) SIRT1 and energy metabolism. Acta Biochim Biophys Sin (Shanghai)
45:51-60.doi:10.1093/abbs/gms108.

19. Vachharajani VT, Liu T, Wang X, Hoth JJ, Yoza BK, McCall CE (2016) Sirtuins
Link Inflammation and Metabolism. J Immunol Res
2016:8167273.doi:10.1155/2016/8167273.

20. Guclu A, Erdur FM, Turkmen K (2016) The Emerging Role of Sirtuin 1 in Cellular
Metabolism, Diabetes Mellitus, Diabetic Kidney Disease and Hypertension. Exp Clin
Endocrinol Diabetes 124:131-9.doi:10.1055/s-0035-1565067.

21. Xie J, Zhang X, Zhang L (2013) Negative regulation of inflammation by SIRT1.
Pharmacol Res 67:60-7.doi:10.1016/j.phrs.2012.10.010.

22. Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, Kalra A et al. (2006)
Resveratrol improves health and survival of mice on a high-calorie diet. Nature
444:337-42.doi:10.1038/nature05354.

23. Jiyun Ahn HL, Chang Hwa Jung, Tae Youl Ha, Young Jin Jang (2015) Effect of a
new dietary SIRT1 activator on high fat diet-induced obesity and the involvement of
microRNAs. FASEB 29

24. Ding RB, Bao J, Deng CX (2017) Emerging roles of SIRT1 in fatty liver diseases.
Int J Biol Sci 13:852-67.doi:10.7150/ijbs.19370.

25. Hirschey MD, Shimazu T, Jing E, Grueter CA, Collins AM, Aouizerat B et al.
(2011) SIRT3 deficiency and mitochondrial protein hyperacetylation accelerate the
development of the metabolic syndrome. Mol Cell 44:177-
90.doi:10.1016/j.molcel.2011.07.019.

26. Pinteric M, Podgorski, II, Hadzija MP, Bujak IT, Dekanic A, Bagaric R et al. (2020)
Role of Sirt3 in Differential Sex-Related Responses to a High-Fat Diet in Mice.
Antioxidants (Basel) 9.doi:10.3390/antiox9020174.

27. Ingram DK, de Cabo R (2017) Calorie restriction in rodents: Caveats to consider.
Ageing Res Rev 39:15-28.doi:10.1016/j.arr.2017.05.008.

28. Ayala JE, Samuel VT, Morton GJ, Obici S, Croniger CM, Shulman GI et al. (2010)
Standard operating procedures for describing and performing metabolic tests of glucose
homeostasis in mice. Dis Model Mech 3:525-34.doi:10.1242/dmm.006239.

29. Bonora E, Moghetti P, Zancanaro C, Cigolini M, Querena M, Cacciatori V et al.
(1989) Estimates of in vivo insulin action in man: comparison of insulin tolerance tests
with euglycemic and hyperglycemic glucose clamp studies. J Clin Endocrinol Metab
68:374-8.doi:10.1210/jcem-68-2-374.

30. Lo S, Russell JC, Taylor AW (1970) Determination of glycogen in small tissue
samples. J Appl Physiol 28:234-6.doi:10.1152/jappl.1970.28.2.234.

31. Folch J, Lees M, Sloane Stanley GH (1957) A simple method for the isolation and
purification of total lipides from animal tissues. J Biol Chem 226:497-509

32. Serra CA, Dos Reis AF, Calsa B, Bueno CS, Helaehil JV, de Souza SAR et al.
(2022) Quercetin prevents insulin dysfunction in hypertensive animals. J Diabetes
Metab Disord 21:407-17.doi:10.1007/s40200-022-00987-4.

33. Corezola do Amaral ME, Kravets V, Dwulet JM, Farnsworth NL, Piscopio R,
Schleicher WE et al. (2020) Caloric restriction recovers impaired beta-cell-beta-cell gap
junction coupling, calcium oscillation coordination, and insulin secretion in prediabetic
mice. Am J Physiol Endocrinol Metab 319:E709-E20.doi:10.1152/ajpendo.00132.2020.

34. Barros RP, Gustafsson JA (2011) Estrogen receptors and the metabolic network.
Cell Metab 14:289-99.doi:10.1016/j.cmet.2011.08.005.

35. Foretz M, Ancellin N, Andreelli F, Saintillan Y, Grondin P, Kahn A et al. (2005)
Short-term overexpression of a constitutively active form of AMP-activated protein
kinase in the liver leads to mild hypoglycemia and fatty liver. Diabetes 54:1331-
9.doi:10.2337/diabetes.54.5.1331.

36. Kim J, Lee H, Lim J, Lee H, Yoon S, Shin SS et al. (2017) The lemon balm extract
ALS-L1023 inhibits obesity and nonalcoholic fatty liver disease in female
ovariectomized mice. Food Chem Toxicol 106:292-305.doi:10.1016/j.fct.2017.05.059.

37. Lei Z, Wu H, Yang Y, Hu Q, Lei Y, Liu W et al. (2021) Ovariectomy Impaired
Hepatic Glucose and Lipid Homeostasis and Altered the Gut Microbiota in Mice With
Different Diets. Front Endocrinol (Lausanne)
12:708838.doi:10.3389/fendo.2021.708838.

38. Boudaba N, Marion A, Huet C, Pierre R, Viollet B, Foretz M (2018) AMPK Re-
Activation Suppresses Hepatic Steatosis but its Downregulation Does Not Promote
Fatty Liver Development. EBioMedicine 28:194-209.doi:10.1016/j.ebiom.2018.01.008.

39. Foretz M, Even PC, Viollet B (2018) AMPK Activation Reduces Hepatic Lipid
Content by Increasing Fat Oxidation In Vivo. Int J Mol Sci
19.doi:10.3390/ijms19092826.

40. Camargo TF, Zanesco AM, Pacher KAS, Andrade TAM, Alves AA, do Amaral
MEC (2020) Physiological profile regulation during weight gain and loss by
ovariectomized females: importance of SIRT1 and SIRT4. Am J Physiol Endocrinol
Metab 319:E769-E78.doi:10.1152/ajpendo.00465.2019.

41. Kim OY, Chae JS, Paik JK, Seo HS, Jang Y, Cavaillon JM et al. (2012) Effects of
aging and menopause on serum interleukin-6 levels and peripheral blood mononuclear
cell cytokine production in healthy nonobese women. Age (Dordr) 34:415-
25.doi:10.1007/s11357-011-9244-2.

42. Yasui T, Uemura H, Yamada M, Matsuzaki T, Tsuchiya N, Noguchi M et al. (2008)
Associations of interleukin-6 with interleukin-1beta, interleukin-8 and macrophage
inflammatory protein-1beta in midlife women. Cytokine 41:302-
6.doi:10.1016/j.cyto.2007.12.001.

43. Vural P, Canbaz M, Akgul C (2006) Effects of menopause and postmenopausal
tibolone treatment on plasma TNFalpha, IL-4, IL-10, IL-12 cytokine pattern and some
bone turnover markers. Pharmacol Res 53:367-71.doi:10.1016/j.phrs.2006.01.005.

44. Corral J, Miralles JM, Garcia-Pascual IJ, Corrales JJ, Garcia-Sastre A, Villar E
(1992) Increased serum N-acetyl-beta-D-glucosaminidase and alpha-D-mannosidase
activities in obese subjects. Clin Investig 70:880-4.doi:10.1007/BF00180432.

45. Sasaki Y, Ikeda Y, Miyauchi T, Uchikado Y, Akasaki Y, Ohishi M (2020)
Estrogen-SIRT1 Axis Plays a Pivotal Role in Protecting Arteries Against Menopause-

Induced Senescence and Atherosclerosis. J Atheroscler Thromb 27:47-
59.doi:10.5551/jat.47993.

46. Pinteric M, Podgorski, II, Popovic Hadzija M, Tartaro Bujak I, Tadijan A, Balog T
et al. (2021) Chronic High Fat Diet Intake Impairs Hepatic Metabolic Parameters in
Ovariectomized Sirt3 KO Mice. Int J Mol Sci 22.doi:10.3390/ijms22084277.

47. Ltaif M, Gargouri M, Soussi A (2021) Protective Effects of A. sativa against
Oxidative Stress-Induced Liver Damage in Ovariectomized Mice. Biomed Res Int
2021:5577498.doi:10.1155/2021/5577498.

48. Fernandez-Sanchez A, Madrigal-Santillan E, Bautista M, Esquivel-Soto J, Morales-
Gonzalez A, Esquivel-Chirino C et al. (2011) Inflammation, oxidative stress, and
obesity. Int J Mol Sci 12:3117-32.doi:10.3390/ijms12053117.

49. Mittal PC, Kant R (2009) Correlation of increased oxidative stress to body weight in
disease-free post menopausal women. Clin Biochem 42:1007-
11.doi:10.1016/j.clinbiochem.2009.03.019.

Figures legends

Fig. 1. Metabolic effects of caloric restriction on ovariectomized mice. (A) Experiment

timeline. (B) Body mass, (C) Lee index, (D) uterine weight, (E) Fat pad index, and (F)

histological examination of the liver and adipose tissue in sham operated (Sham),

ovariectomized (OVX), and ovariectomized + 40% caloric restriction (CR) (OVXR)

mouse groups. Data are mean ± SEM and were analyzed using one-way ANOVA

followed by post-hoc Tukey test. The numbers of mice (n) and p-values are indicated on

each graph.

Fig. 2. Biochemical analyses of glycemic homeostasis. (A) Intraperitoneal glucose

tolerance test (ipGTT); (B) intraperitoneal insulin tolerance test (ipITT); (C) area under

the curve (AUC) of the glucose tolerance test; (D) rate constant for glucose decay (Kitt)

during the ITT test; and (E) HOMA-IR, (F) HOMA%S, and (G) HOMA%B indexes.

Data are mean ± SEM and were analyzed by one-way ANOVA followed by post-hoc

Tukey test. The numbers of mice (n) and p-values are indicated on each graph.

Fig. 3. Biochemical analyses of lipid metabolism. Measurements of (A) serum

cholesterol, (B) serum triglycerides, (C) serum HDL, (D) serum LDL, (E) serum VLDL,

(F) fasting glucose, (G) fasting insulin, and (H) serum protein. Data are mean ± SEM

and were analyzed by one-way ANOVA followed by post-hoc Tukey test. The numbers

of mice (n) and p-values are indicated on each graph.

Fig. 4. Typical immunoblots of (A) catalase (CAT), (B) superoxide dismutase (SOD2),

and (C) representative blots. (D) Serum TBARS, (E) liver TBARS, and (F) liver H_2O_2

levels in the Sham, OVX, and OVXR groups. Data are mean ± SEM and were analyzed

by one-way ANOVA followed by post-hoc Tukey test. The numbers of mice (n) and p-

values are indicated on each graph.

Fig. 5. Typical immunoblots of (A) IL-6, (B) IL-10, (C) TGF-β1, (D) SIRT1, (E)

SIRT3, (F) pAMPK/AMPK, (G) pIR/IR, and (H) representative blots in the livers of

mice in the Sham, OVX, and OVXR groups. Data are mean ± SEM and were analyzed

by one-way ANOVA followed by post-hoc Tukey test. The numbers of mice (n) and p-

values are indicated on each graph.

Table 1: Hepatic and adipose characteristics

Parameters	Sham	OVX	OVXR	Statistical difference
Hepatic Chol (mg/100g tissue)	0.001435±0.0001	0.001637±0.0001	0.001796±0.0003	None
Hepatic TG (mg/100g tissue)	0.0005738±0.0001	0.0006981±0.0002	0.001502±0.0003[a,b]	a: p=0.003 vs Sham b: p=0.008 vs OVX
Hepatic Glycogen (mg/100g tissue)	0.8490±0.144	0.2902±0.059[a,c]	1.164±0.131	a: p=0.012 vs Sham c: p=0.0001 vs OVXR
Liver tissue index*	4.660±0.2420	4.454±0.1980	4.930±0.4340	None
Hepatic NAG (D.O/mg tissue)	0.03224±0.0017	0.04614±0.003[a,c]	0.03174±0.0053	a: p= 0.0421 vs Sham c: p= 0.0351 vs OVXR
Adipocytes area (mm²)	171.5±15.33	411.7±46.23 [a,c]	271.7±26.31	a: p<0.0001 vs Sham c: p=0.0054 vs OVXR

SHAM: control group; OVX, ovariectomized; OVXR (OVX + caloric restriction); *tissue weight (g)/ animal body weight X 100. Data express mean ± std error of mean (n=6/group) and values were compared by one-way ANOVA with Tukey post-test.