Caloric restriction prevents inflammation and insulin dysfunction in middle-aged

ovariectomized mice

Authors: Leticia Roberta Leme Sapatini[1†], Bruno Calsa[1†], Lais Jorge Marim[2], Júlia

Venturini Helaehil[1], and Maria Esméria Corezola do Amaral[1*]

[1]Graduate Program in Biomedical Sciences, University Center of Hermínio Ometto

Foundation, Araras, SP, Brazil

[2]Physiotherapy College, University Center of Hermínio Ometto Foundation, FHO,

Araras, SP, Brazil

[†]Co-first authors: Leticia Sapatini and Bruno Calsa

*Corresponding author:

Maria Esméria Corezola do Amaral

Graduate Program in Biomedical Science, Centro Universitário da Fundação Hermínio

Ometto, FHO, Av. Maximiliano Barutto nº 500, Jardim Universitário, 13607-339,

Araras, SP, Brazil

Tel: +55 (19) 3543 1474; Fax: +55 (19) 35 1412; E-mail: esmeria@fho.edu.br

Abstract

Loss of ovarian function is associated with increased visceral fat. In this study, we aimed to study the effects of caloric restriction (CR) on metabolism in ovariectomized mice. Female, 8–12-month-old mice were divided into three groups: OVX (ovariectomized mice), OVXR (ovariectomized mice on 40% CR) and Sham. CR increased insulin sensitivity and glucose tolerance as evidenced by the results of intraperitoneal glucose tolerance and intraperitoneal insulin tolerance tests, liver glycogen levels, and homeostasis model assessment indexes. Increased AMPK phosphorylation was observed in the liver of OVXR mice. CR also increased hepatic cholesterol and triglyceride levels. The reductions in the level of 2-thiobarbituric acid reactive substances in the serum and liver and of hydrogen peroxide in the liver of OVXR mice suggested alterations in the redox state of the liver. Although expression of catalase protein was reduced by CR, expression of superoxide dismutase was not altered by CR. Although interleukin (IL)-6 and IL-10 levels in OVXR mice were similar to those in Sham mice, macrophage infiltration was reduced in OVXR mice as evidenced by n-acetylglucosaminidase activity. OVXR mice had increased sirtuin (SIRT)1 levels and decreased SIRT3 levels in the liver. In conclusion, CR improved the condition of ovariectomized mice by reducing adiposity and increasing insulin sensitivity and glucose tolerance through a mechanism that may involve AMPK.

Keywords: caloric restriction, obesity, inflammation, menopause

Introduction

Numerous studies have shown that menopause is associated with unfavorable changes in body composition, such as increased abdominal fat deposition, and poor overall

health (Davis et al. 2012; Qian et al. 2016). The deleterious effects of obesity are

diverse and include various metabolic diseases, such as diabetes mellitus, and

cardiovascular diseases, like coronary heart disease, heart attack, stroke, and

hypertension, which adversely impact the quality of life and carry an increased risk of

premature death (Camporez et al. 2013; Nappi and Kokot-Kierepa 2012; Neto et al.

2015; Tominaga et al. 2011). However, the relationship between menopause and obesity

is complex. Sex hormones fluctuate with reproductive changes, such as menarche,

pregnancy, and menopause, which may play an important role in the expansion of

adipose tissue (Davis et al. 2012). Hormones impact adipose tissue, and estrogens

regulate glucose homeostasis and insulin sensitivity in several ways, by altering insulin

production and secretion (Choi et al. 2005; Santos et al. 2016), insulin-mediated glucose

uptake in peripheral tissues, mitochondrial function (Hamilton et al. 2016), and the

expression of inflammatory markers (Straub 2007). Given the apparent interaction

between low estrogen concentrations and obesity in altering glucose and lipid

homeostasis, increasing attention has been given to developing dietary strategies that

preserve homeostasis during menopause.

Caloric restriction (CR) is one strategy that can reduce body weight, total body

fat, and visceral fat, similar to exercise. CR is a non-pharmacological dietary

intervention in which the average daily calorie intake is reduced without malnutrition or

deprivation of essential nutrients. Animals on a calorie-restricted diet showed a

significant (50%) increase in life expectancy (Mattison et al. 2017). Animal studies

showed favorable effects of short-term CR on cardiometabolic changes, liver changes,

and cellular senescence (Li et al. 2016; Yzydorczyk et al. 2019). In humans,

experimental and observational studies have revealed that CR can reduce the incidence

of cardiovascular disease, diabetes, dementia, frailty, and cancer (Al-Regaiey 2016;

Weiss and Fontana 2011). These observed benefits are related to CR-mediated activation of regulatory proteins in metabolic pathways, such as AMP-activated protein kinase (AMPK) and sirtuins (SIRTs). AMPK is a key sensor and regulator of glucose metabolism that activates SIRT1 deacetylase in response to changes in the redox state (NAD+/NADH and AMP/ATP ratios) of the cell (Fulco and Sartorelli 2008). Sirtuins are a group of highly conserved NAD+-dependent histones and protein deacetylases and/or ADP-ribosyl transferases that play important roles in various biological processes (Finkel et al. 2009; Li 2013; Santos et al. 2016; Vachharajani et al. 2016). SIRT1 functions as a metabolic/energy sensor that modulates the activity and/or gene expression of several crucial transcription factors and transcription coactivators involved in metabolic homeostasis (Guclu et al. 2016; Li 2013; Xie et al. 2013). Reduced levels of NAD+ and SIRT1 activity have been detected under energy-overloaded metabolic profiles, such as a hyperlipidic diet (Baur et al. 2006; Jiyun Ahn 2015).

In contrast, CR increased NAD+ levels and induced activation of SIRT1 (Baur et al. 2006; Jiyun Ahn 2015). SIRT1 functions as an essential regulator of hepatic lipid metabolism through SREBP-1c/ChREBP-dependent control of lipogenesis by inhibiting lipid synthesis and increasing PPARα/PGC-1α-dependent β-fatty acid oxidation (Ding et al. 2017). SIRT1 regulates oxidative stress by increasing the transcriptional activity of antioxidant enzymes, such as superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPX) (Ding et al. 2017). Activation of SIRT1 reduces nitric oxide synthase and lipid peroxidation products, which has an anti-oxidative stress effect in the liver (Ding et al. 2017). SIRT3 is a mitochondrial protein that integrates cellular energy metabolism and plays an important role in preventing metabolic syndrome (Hirschey et al. 2011). SIRT3 knockout mice showed characteristics of reduced

mitochondrial metabolism, such as reduced fatty acid catabolism (Hirschey et al. 2011).

SIRT3 expression was reduced in male mice fed a high-fat diet (Hirschey et al. 2011; Pinteric et al. 2020). However, the consistency and reproducibility of results across investigations are varied, and key questions about the timing, duration, modality, and adverse effects of CR, as well as the influence of genotype and diet composition, remain (Ingram and de Cabo 2017). Given the low adherence of humans to difficult-to-follow diets, research on dietary interventions should be conducted. This study examined the impact of 40% CR on the liver of middle-aged mice who underwent ovariectomy. CR was expected to have a positive effect on metabolism. The results showed that CR was associated with improvements in both metabolism, mainly glycemic metabolism, and inflammatory markers.

Materials and methods

Experimental animals and model establishment

The project was approved by the Committee on Ethics in Animal Use (CEUA) under protocol #009/2020. Eighteen C57BL/6J mice (8–12 months old, weighing ~20 g) were housed in individual cages at a constant temperature (22 °C) under a 12 h light/dark cycle with water and commercial chow *ad libitum*. For the ovariectomy, the mice were anesthetized by intraperitoneal injection with ketamine (50 mg/kg) and xylazine (5 mg/kg). After shaving the inguinal region, an abdominal incision was made, and the ovaries were located and removed. After surgery, the animals were placed in a heated chamber to avoid hypothermia. The animals in the control group (Sham) underwent the same procedure except the ovaries were not removed. At 16 weeks after surgery, the ovariectomy model mice were separated into two groups: the OVX (ovariectomized) and OVXR (ovariectomized plus 40% CR) groups. Sham and OVX mice were fed a

commercial isocaloric diet *ad libitum*, and OVXR mice received 60% of the chow

consumed by the OVX mice for 30 days, corresponding to 40% CR. All mice had free

access to water throughout the experiment. The body weights of the animals were

measured weekly throughout the experiment.

Intraperitoneal glucose tolerance test

The intraperitoneal glucose tolerance test (ipGTT) was administered 5 days before

euthanasia after a 6 h fast. For the test, glucose (2 g/kg body weight) was injected

intraperitoneally. Blood samples were collected before glucose overload (time zero),

and at 15, 30, 60, and 120 min after glucose infusion. Blood glucose was determined

using reagent strips and an Abbott® glucometer as previously described (Ayala et al.

2010).

Intraperitoneal insulin tolerance test

The intraperitoneal insulin tolerance test (ipITT) was administered 3 days before

euthanasia after a 6 h fast. Insulin (regular crystalline, 0.75 U/kg body weight) was

administered intraperitoneally. Blood samples were collected before insulin injection

(time zero) and at 5, 10, 15, 20, 25, and 30 min after injection. Blood glucose was

determined using an Abbott glucometer and reagent strips. The rate constant for glucose

disappearance (Kitt) was calculated using the formula $In2/t1/2$. The half-life (t1/2) of

serum glucose was calculated from the slope of the minimum regression curve in the

linear phase of plasma glucose decline (Bonora et al. 1989).

Homeostasis model assessments

Homeostasis model assessment of insulin resistance (HOMA-IR), insulin sensitivity

(HOMA%S), and β-cell function (HOMA%β) were calculated using the HOMA

calculator from the University of Oxford (www.dtu.ox.ac.uk/homacalculator).

Fat pad and Lee indexes

The perirenal adipose tissue weight and body weight were assessed, and the adipose

tissue index was calculated by dividing adipose tissue weight by body weight. The Lee

index was calculated using the following equation: $\sqrt[3]{}$body weight (g)÷nose to anus

length (cm). .

Biochemical assessments

Serum glucose, triglycerides, cholesterol, HDL, and total protein were determined using

commercial kits (Laborlab, Guarulhos, SP, Brazil). LDL was calculated using the

Friedewald equation, and VLDL was calculated by dividing triglycerides by 5. Insulin

serum was measured using ELISA (Rat/Mouse Insulin ELISA, #EZRMI-13K, Millipore

St. Charles, MO, USA)). Hepatic glycogen content was measured using a colorimetric

method (Lo et al. 1970). Total liver lipids were extracted according to the method of

Folch et al. (1957).

Preparation of liver homogenates for oxidative stress biomarker analysis

Liver samples (~50 mg) were homogenized in Krebs buffer (118 mM NaCl, 25 mM

$NaHCO_3$, 1.2 mM KH_2PO_4, 4.7 mM KCl, 1.2 mM $MgSO_4$, 1.25 mM $CaCl_2$, 10 mM

glucose, and 10 mM HEPES, pH 7.4). The obtained tissue homogenate was used for the

determination of 2-thiobarbituric acid reactive substances (TBARS) and hydrogen

peroxide (H_2O_2) levels. Total protein was quantified using the Biuret method, with

bovine serum albumin as a standard, and measured at 540 nm.

Determination of hepatic H_2O_2

165 H_2O_2 levels in hepatic tissue were determined using the Amplex Red method. Briefly,

166 20 µL of homogenate was incubated with horseradish peroxidase (HRP, 1 U/mL) and

167 Amplex Red (50 µM) at 37 °C for 30 min. Then, the absorbance was read at 560 nm.

168 The H_2O_2 concentration was calculated by referring to a standard curve of known

169 concentrations and normalized to the protein concentration.

Hepatic and systemic TBARS

Serum and liver TBARS contents were determined as described previously (Serra et al. 2022). Briefly, 50 µL of serum or liver homogenate was mixed with 10 µL of 0.1 M Butylated Hydroxytoluene, 400 µL of 1% 2-thiobarbituric acid (TBA; Merck, St. Charles, MO, USA), and 200 µL of 20% phosphoric acid and heated at 100 °C for 15 min. After cooling, 1.5 mL of 1-butanol was added, and the optical density of the supernatant was determined at 532 nm.

Determination of hepatic n-acetylglucosaminidase

n-Acetylglucosaminidase (NAG) activity was measured to determine liver macrophagic infiltration. An aliquot (3.0 µL) of liver homogenate (0.08 M $NaPO_4$) was mixed with 30 µL of p-nitrophenyl-2-acetamide-β-D-glucopyranoside (Sigma-Aldrich) and diluted in 50 µL of 50 mM citrate buffer. Finally, 50 µL of 0.2 M glycine was added, and the absorbance was read at 405 nm.

Histological analysis of liver and adipose tissue

Liver and adipose tissues were fixed with 10% formalin and dehydrated. Then, small fragments of the tissues were embedded in paraffin (Paraplast; Sigma, St. Louis, MO, USA) using standard procedures. Slices (5 µm thick) were stained with hematoxylin-eosin (HE), and images were acquired using a microscope (Leica DM 2000, Germany) with LAS v.4.1 software. For each mouse, 40 images were obtained at 400× magnification. The area (mm^2) containing adipocytes in each image was quantified using the ImageJ software (National Institutes of Health, Bethesda, Maryland, USA).

Immunoblotting

Liver tissue fragments were homogenized in protein extraction buffer (10 mM EDTA, 100 mM Tris-HCL, 10 mM sodium pyrophosphate, 100 mM sodium fluoride, 100 mM sodium orthovanadate, 10 mM phenylmethylsulfonyl fluoride, and 0.1 mg/mL

aprotinin). Protein content was determined according to the Biuret method. The samples

were mixed with Laemmli buffer (0.1% bromophenol blue, 1 M sodium phosphate,

50% glycerol, and 10% SDS). Total protein (50 μg) was separated by electrophoresis.

Then, the separated proteins were transferred to nitrocellulose membranes. After

transfer, the membranes were incubated at 4 °C, with shaking, overnight with the

following primary antibodies: IL-6, IL-10, TGF-β1, SIRT3, pIR, and IR (Santa Cruz,

Dallas, USA; 1:250), and SIRT1, pAMPK, and AMPK (Cell Signaling Danvers,

Massachusetts, USA; 1:1000). The membranes were then incubated with an HRP-

secondary antibody (Santa Cruz; Dallas, USA; 1:10,000), with shaking, followed by

development with chemiluminescent reagents (SuperSignal West Pico PLUS, Thermo

Scientific, Rockford, IL, USA) and bands intensities were determined by densitometry

scanning using the ImageJ software (National Institutes of Health, Bethesda, Maryland,

USA)).

Statistical analysis

Data are mean ± standard error of the mean. All data were analyzed using the Shapiro–

Wilk Normality test. For comparisons between experimental groups, one-way ANOVA

and Tukey's post-test were used. All statistical and graphical analyses were performed

using GraphPad Prism software (version 8.0). Significance was set at $p < 0.05$.

Results

Ovariectomy and caloric restriction characterization

The timeline of the experiment is shown in Figure 1A. After ovariectomy, the animals

were observed for 4 months. The animals in the OVXR group remained on a restricted

diet for the first month. The body weights of the animals were monitored throughout the

experiment and are presented in Figure 1B, which shows evident weight gain in the

OVX group compared to in the Sham group and marked weight loss during CR in the

OVXR group compared to in the Sham and OVX groups. The Lee index of the mice

was lower in the OVXR group than in the Sham and OVX groups (Figure 1C). Analysis

of the uterine weight of the mice showed evident uterine atrophy following ovariectomy

(Figure 1D).

Mice in the OVXR group had lower perirenal adipose tissue weight than mice in

the OVX group (Figure 1E); however, the difference was not statistically different.

OVX mice showed adipose tissue weight gain when compared to mice in the other

groups. HE staining of liver and adipose tissue (Figure 1F) showed slight steatosis in

OVXR group mice when compared to the Sham and OVX group mice. There was an

evident reduction in adipocyte diameter in OVXR mice when compared to that in OVX

mice.

Effects on glucose, insulin, and lipid metabolism

The GTT and ITT results (Figures 2A and 2B, respectively) showed a reduced area

under the curve for OVXR mice (Figure 2C) compared to that for OVX mice and an

increased Kitt for OVXR mice compared to that for OVX mice (Figure 2D). These

results suggest greater glucose tolerance and increased insulin sensitivity of OVXR

mice than of OVX mice. The HOMA-IR (Figure 2E), HOMA%S (Figure 2F), and

HOMA%B (Figure 2G) indexes indicated increased insulin sensitivity and beta cell

preservation in the OVXR mice compared to in the OVX mice. Consistent with these

results, liver glycogen levels were similar in OVXR and Sham mice but lower in OVX

than in OVXR mice (Table 1). No difference in the liver index was observed among the

groups. The reduced blood glucose levels (Figure 3F) and insulinemia (Figure 3G) in

the OVXR group compared to that in the Sham and OVX groups suggest improvement

in glycemic homeostasis under CR. Similar serum protein concentrations (Figure 3H) in

all groups indicate good nutritional status.

Serum cholesterol levels were lower in the OVXR group than in the OVX group

(Figure 3A). Serum triglycerides (Figure 3B) and VLDL (Figure 3E) were similar

among the groups. LDL-cholesterol (Figure 3D) was higher in the OVX group than in

the Sham group. Normalization of the HDL-cholesterol fraction was observed for

OVXR mice compared to Sham mice (Figure 3C), and HDL-cholesterol was lower in

OVX mice than in Sham mice. Hepatic cholesterol levels were similar among the

groups, but hepatic triglycerides were higher in OVXR mice than in Sham and OVX

mice (Table 1). This result is mirrored by the liver tissue histology, which showed lipid

vacuoles in the livers of OVXR mice (Figure 1F), suggesting metabolic adaptation to

the CR diet. Similar adipocyte diameter (Table 1) was observed in the OVXR and Sham

groups, whereas adipocyte diameter was increased in the OVX group when compared to

that in the OVXR group.

Oxidative stress balance

CAT protein expression was higher in OVX mice than in OVXR mice (Figure 4A), and

was similar in OVXR and Sham mice. SOD-2 protein expression (Figure 4B) was

similar among the groups. Representative bands of CAT, SOD-2, and beta-actin protein

are shown in Figure 4C. Analysis of TBARS content in the serum (Figure 4D) and liver

(Figure 4E) and H_2O_2 levels (Figure 4F) showed lower levels in the OVXR group than

in the OVX group, suggesting reduced oxidative stress in the OVXR group compared to

that in the OVX and Sham groups.

Hepatic inflammatory balance

Macrophage infiltration (NAG activity) in liver tissue was similar in the OVXR and

Sham groups (Table 1) but was higher in the OVX group than in the Sham and OVXR

groups. Levels of IL-6, a pro-inflammatory cytokine, were higher in OVX mice than in

Sham mice (Figure 5A), whereas levels of IL-10, anti-inflammatory cytokine, were

lower in OVX mice than in Sham mice (Figure 5B). Similar TGF protein expression

was observed in all groups (Figure 5C).

Hepatic metabolism

SIRT1 protein expression in the liver tissue was higher in OVXR mice than in Sham

mice (Figure 5D). In contrast, SIRT3 (Figure 5E) and IR (Figure 5G) protein levels

were lower in OVXR and OVX mice than in Sham mice. AMPK protein expression was

higher in OVXR mice than in OVX and Sham mice (Figure 5F). Representative protein

bands of AMPK, pAMPK, IL-6, IL-10, IR, pIR, SIRT1, SIRT3, TGF, and beta-actin are

shown in Figure 5H.

Discussion

CR is a well-accepted dietary intervention that can restore glucose homeostasis and

promote weight loss (Corezola do Amaral et al. 2020). In this study, we expected that

CR would ameliorate the metabolic dysfunction and inflammation induced by

ovariectomy. We observed weight loss and lower adiposity in ovariectomized mice

under CR. These mice also had higher insulin sensitivity and normalized lipid profiles

compared to the other mice. This can be attributed due a CR-induced homeostatic effect

on oxidative stress and inflammatory balance in hepatic tissue.

Estrogen exerts protective effects on metabolism by regulating feeding behavior,

glucose and lipid uptake, and the insulin pathway (Barros and Gustafsson 2011). The

reduction in estrogen level during menopause can lead to metabolism disruption,

causing insulin resistance. In this study, CR decreased fasting glucose and insulin

levels. Corroborating these findings, OVXR mice presented higher insulin sensitivity

and glucose tolerance as evidenced by the ipGTT and ipITT results and the HOMA

index. These beneficial changes in glycemic homeostasis can be attributed to increased

expression of pAMPK/AMPK, but not pIR/IR, in the liver. AMPK controls glycemia

and gluconeogenesis and is an important treatment target in diabetes (Foretz et al.

2005). Although the cholesterol fractions and triglycerides were unchanged, CR reduced

total serum cholesterol.

In previous studies, ovariectomized mice displayed a nonalcoholic fatty liver

disease (NAFLD) phenotype (Kim et al. 2017; Lei et al. 2021). Although we did not

observe any changes in the cholesterol and triglyceride contents in the livers of OVX

mice, triglycerides were increased in the liver of OVXR animals. Activation of AMPK

inhibits hepatic lipogenesis by inactivating acetyl CoA carboxylase (Boudaba et al.

2018), resulting in fat oxidation (Foretz et al. 2018). Interestingly, AMPK expression

induced hepatic lipid accumulation and fatty acid oxidation (Foretz et al. 2005). In this

study, hepatic pAMPK/AMPK expression was significantly increased in OVXR mice,

suggesting its involvement in hepatic lipid metabolism. This hepatic lipid storage in

OVXR mice may be a transient metabolic process related to the restricted diet. Under

CR, the OVXR mice used fat as an energy source to maintain cellular metabolism,

which may explain the lipid accumulation in the liver. The mechanism may involve

rapid response of AMPK phosphorylation promoting uptake of fatty acids stored in the

liver (Foretz et al. 2018; Foretz et al. 2005). Our histological observations showed mild

hepatic steatosis and a reduced adipocyte area following CR, corroborating the proposed

mechanism. In a previous study, CR animals presented increased serum concentrations

of free fatty acids and SIRT1, which may indicate mobilization of triglycerides by

SIRT1 under CR (Camargo et al. 2020).

Lipid accumulation increases oxidative stress and inflammation. This can

exacerbate release of reactive oxygen species and pro-inflammatory cytokines, such as

tumor-necrosis factor (TNF)-α and IL-6, in liver and adipose tissue. Previous studies

have reported an increase in IL-6 in menopause (Kim et al. 2012; Yasui et al. 2008).

After menopause, levels of the anti-inflammatory cytokines IL-4, IL-10, and IL-12 are

increased, which is related to an increase in TNF-α, as a compensatory mechanism

(Vural et al. 2006). In our study, OVX mice showed inflammatory effects, evidenced by

an increase in IL-6 and a reduction in IL-10, due to the reduction in estrogen levels. The

levels of these cytokines were unchanged in OVXR mice. Although TGF is an

inflammatory factor that induces fibrosis, its expression was unaffected by ovariectomy.

NAG is an indicator of macrophage activity and inflammation. The reduction in NAG

level under CR suggests the anti-inflammatory effect of CR. NAG concentrations have

been linked to obesity and diabetes (Corral et al. 1992).

In NAFLD, SIRT1 activation plays a beneficial role by decreasing oxidative

stress and inducing deacetylation of PGC-1α, leading to reduced pro-inflammatory

cytokine levels. Studies have shown that ovariectomy induced unfavorable

cardiovascular changes and that estrogen treatment activated SIRT1 and promoted

cardiovascular protection (Sasaki et al. 2020). In mice, ovariectomy and SIRT3

depletion impaired the antioxidative system response and increased lipid damage, body

weight gain, and the levels of oxidative stress-inducing genes (Pinteric et al. 2021). Our

findings showed that ovariectomy increased SIRT1 and decreased SIRT3, suggesting

these proteins function in a regulatory mechanism to maintain normal metabolism.

The reductions in serum and liver TBARS contents and liver H_2O_2 levels in the

OVXR mice suggest changes in the redox state of the liver as a protective mechanism

against oxidative damage. The literature supports an increase in TBARS content

following ovariectomy, corroborating the findings of the present study (Ltaif et al.

2021). The reduction in TBARS contents in the OVXR group may explain the reduction

in CAT and the similar SOD expression levels in the present study, suggesting a balance

of mechanisms to combat oxidative stress under hypoestrogenism. The activity levels of

the antioxidant enzymes SOD, CAT, and GPX were significantly diminished in obesity

(Fernandez-Sanchez et al. 2011), and oxidative stress was enhanced as body weight

increased in post-menopausal women (Mittal and Kant 2009). Here, we aimed to

understand the specific functions of CR in the "metabolic network" of tissues involved

in whole-body glucose and lipid metabolism in ovariectomized mice. Notably, while

CR benefitted the ovariectomized mice by reducing adiposity and increasing insulin

sensitivity and glucose tolerance, suggesting a balance in oxidative stress, it induced

negative changes in lipid metabolism in the liver, which was associated with changes in

the protein expression of SIRT1 and SIRT3. Finally, the results of our study suggest

that ovariectomy can accelerate metabolic changes from tissue damage to the body and

that CR was able to restore, at least in part, glycemic homeostasis.

Author statements

Acknowledgments

We thank Ana Cristina Pires Menegheti, Bruno Alves Cia, Lucas Orzari, Mateus

Eduardo Bortolanza da Silva, and Renata Barbieri for technical assistance.

Competing interests' statement

No conflicts of interest, financial or otherwise, are declared by the authors.

Author contribution statement

Sapatini LRL: Data curation, Investigation, Visualization, Writing – original draft.

Calsa B: Data curation, Formal analysis, Investigation, Validation, Visualization,

Writing – original draft, review and editing. **Marim LJ:** Investigation. **Helaehil JV:**

Data curation, Investigation. **Amaral MEC:** Conceptualization, Data curation, Formal

analysis, Funding acquisition, Investigation, Project administration, Resources,

Supervision, Validation, Visualization, Writing – review & editing.

Funding statement

This work was supported by the Herminio Ometto Foundation.

Data availability

Data generated or analyzed during this study are available form the corresponding

author upon reasonable request.

References

Al-Regaiey, K.A. (2016). The effects of calorie restriction on aging: a brief review. *Eur Rev Med Pharmacol Sci,* 20(11), 2468-2473.

Ayala, J.E., Samuel, V.T., Morton, G.J., Obici, S., Croniger, C.M., Shulman, G.I., Wasserman, D.H., McGuinness, O.P., & Consortium, N.I.H.M.M.P.C. (2010). Standard operating procedures for describing and performing metabolic tests of glucose homeostasis in mice. *Dis Model Mech,* 3(9-10), 525-534. https://doi.org/10.1242/dmm.006239

Barros, R.P., & Gustafsson, J.A. (2011). Estrogen receptors and the metabolic network. *Cell Metab,* 14(3), 289-299. https://doi.org/10.1016/j.cmet.2011.08.005

Baur, J.A., Pearson, K.J., Price, N.L., Jamieson, H.A., Lerin, C., Kalra, A., Prabhu, V.V., Allard, J.S., Lopez-Lluch, G., Lewis, K., Pistell, P.J., Poosala, S., Becker, K.G., Boss, O., Gwinn, D., Wang, M., Ramaswamy, S., Fishbein, K.W., Spencer, R.G., Lakatta, E.G., Le Couteur, D., Shaw, R.J., Navas, P., Puigserver, P., Ingram, D.K., de Cabo, R., & Sinclair, D.A. (2006). Resveratrol improves the health and survival of mice on a high-calorie diet. *Nature,* 444(7117), 337-342. https://doi.org/10.1038/nature05354

Bonora, E., Moghetti, P., Zancanaro, C., Cigolini, M., Querena, M., Cacciatori, V., Corgnati, A., & Muggeo, M. (1989). Estimates of in vivo insulin action in man: comparison of insulin tolerance tests with euglycemic and hyperglycemic glucose clamp studies. *J Clin Endocrinol Metab,* 68(2), 374-378. https://doi.org/10.1210/jcem-68-2-374

Boudaba, N., Marion, A., Huet, C., Pierre, R., Viollet, B., & Foretz, M. (2018). AMPK re-activation suppresses hepatic steatosis but its downregulation does not promote fatty

liver development. *EBioMedicine,* 28194-209. https://doi.org/10.1016/j.ebiom.2018.01.008

Camargo, T.F., Zanesco, A.M., Pacher, K.A.S., Andrade, T.A.M., Alves, A.A., & do Amaral, M.E.C. (2020). Physiological profile regulation during weight gain and loss by ovariectomized females: the importance of SIRT1 and SIRT4. *Am J Physiol Endocrinol Metab,* 319(4), E769-E778. https://doi.org/10.1152/ajpendo.00465.2019

Camporez, J.P., Jornayvaz, F.R., Lee, H.Y., Kanda, S., Guigni, B.A., Kahn, M., Samuel, V.T., Carvalho, C.R., Petersen, K.F., Jurczak, M.J., & Shulman, G.I. (2013). Cellular mechanism by which estradiol protects female ovariectomized mice from high-fat diet-induced hepatic and muscle insulin resistance. *Endocrinology,* 154(3), 1021-1028. https://doi.org/10.1210/en.2012-1989

Choi, S.B., Jang, J.S., & Park, S. (2005). Estrogen and exercise may enhance beta-cell function and mass via insulin receptor substrate 2 induction in ovariectomized diabetic rats. *Endocrinology,* 146(11), 4786-4794. https://doi.org/10.1210/en.2004-1653

Corezola do Amaral, M.E., Kravets, V., Dwulet, J.M., Farnsworth, N.L., Piscopio, R., Schleicher, W.E., Miranda, J.G., & Benninger, R.K.P. (2020). Caloric restriction recovers impaired beta-cell-beta-cell gap junction coupling, calcium oscillation coordination, and insulin secretion in prediabetic mice. *Am J Physiol Endocrinol Metab,* 319(4), E709-E720. https://doi.org/10.1152/ajpendo.00132.2020

Corral, J., Miralles, J.M., Garcia-Pascual, I.J., Corrales, J.J., Garcia-Sastre, A., & Villar, E. (1992). Increased serum N-acetyl-beta-D-glucosaminidase and alpha-D-mannosidase activities in obese subjects. *Clin Investig,* 70(10), 880-884. https://doi.org/10.1007/BF00180432

Davis, S.R., Castelo-Branco, C., Chedraui, P., Lumsden, M.A., Nappi, R.E., Shah, D.,
Villaseca, P., & Writing Group of the International Menopause Society for World
Menopause, D. (2012). Understanding weight gain at menopause. *Climacteric,* 15(5),
419-429. https://doi.org/10.3109/13697137.2012.707385

Ding, R.B., Bao, J., & Deng, C.X. (2017). Emerging roles of SIRT1 in fatty liver
diseases. *Int J Biol Sci,* 13(7), 852-867. https://doi.org/10.7150/ijbs.19370

Fernandez-Sanchez, A., Madrigal-Santillan, E., Bautista, M., Esquivel-Soto, J.,
Morales-Gonzalez, A., Esquivel-Chirino, C., Durante-Montiel, I., Sanchez-Rivera, G.,
Valadez-Vega, C., & Morales-Gonzalez, J.A. (2011). Inflammation, oxidative stress,
and obesity. *Int J Mol Sci,* 12(5), 3117-3132. https://doi.org/10.3390/ijms12053117

Finkel, T., Deng, C.X., & Mostoslavsky, R. (2009). Recent progress in the biology and
physiology   of   sirtuins.   *Nature,*   460(7255),   587-591.
https://doi.org/10.1038/nature08197

Folch, J., Lees, M., & Sloane Stanley, G.H. (1957). A simple method for the isolation
and purification of total lipids from animal tissues. *J Biol Chem,* 226(1), 497-509.

Foretz, M., Even, P.C., & Viollet, B. (2018). AMPK activation reduces hepatic lipid
content by increasing fat oxidation in vivo. *Int J Mol Sci,* 19(9).
https://doi.org/10.3390/ijms19092826

Foretz, M., Ancellin, N., Andreelli, F., Saintillan, Y., Grondin, P., Kahn, A., Thorens,
B., Vaulont, S., & Viollet, B. (2005). Short-term overexpression of a constitutively
active form of AMP-activated protein kinase in the liver leads to mild hypoglycemia
and fatty liver. *Diabetes,* 54(5), 1331-1339. https://doi.org/10.2337/diabetes.54.5.1331

Fulco, M., & Sartorelli, V. (2008). Comparing and contrasting the roles of AMPK and

SIRT1 in metabolic tissues. *Cell Cycle,* 7(23), 3669-3679.

https://doi.org/10.4161/cc.7.23.7164

Guclu, A., Erdur, F.M., & Turkmen, K. (2016). The emerging role of sirtuin 1 in

cellular metabolism, diabetes mellitus, diabetic kidney disease and hypertension. *Exp*

Clin Endocrinol Diabetes, 124(3), 131-139. https://doi.org/10.1055/s-0035-1565067

Hamilton, D.J., Minze, L.J., Kumar, T., Cao, T.N., Lyon, C.J., Geiger, P.C., Hsueh,

W.A., & Gupte, A.A. (2016). Estrogen receptor alpha activation enhances

mitochondrial function and systemic metabolism in high-fat-fed ovariectomized mice.

Physiol Rep, 4(17). https://doi.org/10.14814/phy2.12913

Hirschey, M.D., Shimazu, T., Jing, E., Grueter, C.A., Collins, A.M., Aouizerat, B.,

Stancakova, A., Goetzman, E., Lam, M.M., Schwer, B., Stevens, R.D., Muehlbauer,

M.J., Kakar, S., Bass, N.M., Kuusisto, J., Laakso, M., Alt, F.W., Newgard, C.B., Farese,

R.V., Jr., Kahn, C.R., & Verdin, E. (2011). SIRT3 deficiency and mitochondrial protein

hyperacetylation accelerate the development of metabolic syndrome. *Mol Cell,* 44(2),

177-190. https://doi.org/10.1016/j.molcel.2011.07.019

Ingram, D.K., & de Cabo, R. (2017). Calorie restriction in rodents: Caveats to consider.

Ageing Res Rev, 3915-3928. https://doi.org/10.1016/j.arr.2017.05.008

Jiyun Ahn, H.L., Jung, C.H., Ha, T.Y., Jang, Y.J. (2015). Effect of a new dietary SIRT1

activator on high fat diet-induced obesity and the involvement of microRNAs. *FASEB*

29(S1). https://doi.org/10.1096/fasebj.29.1_supplement.405.8

Kim, J., Lee, H., Lim, J., Lee, H., Yoon, S., Shin, S.S., & Yoon, M. (2017). The lemon

balm extract ALS-L1023 inhibits obesity and nonalcoholic fatty liver disease in female

ovariectomized mice. *Food Chem Toxicol,* 106(Pt A), 292-305. https://doi.org/10.1016/j.fct.2017.05.059

Kim, O.Y., Chae, J.S., Paik, J.K., Seo, H.S., Jang, Y., Cavaillon, J.M., & Lee, J.H. (2012). Effects of aging and menopause on serum interleukin-6 levels and peripheral blood mononuclear cell cytokine production in healthy nonobese women. *Age (Dordr),* 34(2), 415-425. https://doi.org/10.1007/s11357-011-9244-2

Lei, Z., Wu, H., Yang, Y., Hu, Q., Lei, Y., Liu, W., Nie, Y., Yang, L., Zhang, X., Yang, C., Lin, T., Tong, F., Zhu, J., & Guo, J. (2021). Ovariectomy impaired hepatic glucose and lipid homeostasis and altered the gut microbiota in mice with different diets. *Front Endocrinol (Lausanne),* 12708838. https://doi.org/10.3389/fendo.2021.708838

Li, N., Guenancia, C., Rigal, E., Hachet, O., Chollet, P., Desmoulins, L., Leloup, C., Rochette, L., & Vergely, C. (2016). Short-term moderate diet restriction in adulthood can reverse oxidative, cardiovascular and metabolic alterations induced by postnatal overfeeding in mice. *Sci Rep,* 630817. https://doi.org/10.1038/srep30817

Li, X. (2013). SIRT1 and energy metabolism. *Acta Biochim Biophys Sin (Shanghai),* 45(1), 51-60. https://doi.org/10.1093/abbs/gms108

Lo, S., Russell, J.C., & Taylor, A.W. (1970). Determination of glycogen in small tissue samples. *J Appl Physiol,* 28(2), 234-236. https://doi.org/10.1152/jappl.1970.28.2.234

Ltaif, M., Gargouri, M., & Soussi, A. (2021). Protective effects of *A. sativa* against oxidative stress-induced liver damage in ovariectomized mice. *Biomed Res Int,* 20215577498. https://doi.org/10.1155/2021/5577498

Mattison, J.A., Colman, R.J., Beasley, T.M., Allison, D.B., Kemnitz, J.W., Roth, G.S., Ingram, D.K., Weindruch, R., de Cabo, R., & Anderson, R.M. (2017). Caloric

restriction improves the health and survival of rhesus monkeys. *Nat Commun,* 814063.

https://doi.org/10.1038/ncomms14063

Mittal, P.C., & Kant, R. (2009). Correlation of increased oxidative stress to body weight

in disease-free postmenopausal women. *Clin Biochem,* 42(10-11), 1007-1011.

https://doi.org/10.1016/j.clinbiochem.2009.03.019

Nappi, R.E., & Kokot-Kierepa, M. (2012). Vaginal health: Insights, views & attitudes

(VIVA) - results from an international survey. *Climacteric,* 15(1), 36-44.

https://doi.org/10.3109/13697137.2011.647840

Neto, N.I., Rodrigues, M.E., Hachul, A.C., Moreno, M.F., Boldarine, V.T., Ribeiro,

E.B., Oyama, L.M., & Oller do Nascimento, C.M. (2015). A hyperlipidic diet combined

with short-term ovariectomy increases adiposity and hyperleptinemia and decreases

cytokine content in mesenteric adipose tissue. *Mediators Inflamm,* 2015923248.

https://doi.org/10.1155/2015/923248

Pinteric, M., Podgorski, II, Popovic Hadzija, M., Tartaro Bujak, I., Tadijan, A., Balog,

T., & Sobocanec, S. (2021). Chronic high fat diet intake impairs hepatic metabolic

parameters in ovariectomized Sirt3 KO mice. *Int J Mol Sci,* 22(8).

https://doi.org/10.3390/ijms22084277

Pinteric, M., Podgorski, II, Hadzija, M.P., Bujak, I.T., Dekanic, A., Bagaric, R., Farkas,

V., Sobocanec, S., & Balog, T. (2020). Role of Sirt3 in differential sex-related

responses to a high-fat diet in mice. *Antioxidants (Basel),* 9(2).

https://doi.org/10.3390/antiox9020174

Qian, S.W., Liu, Y., Wang, J., Nie, J.C., Wu, M.Y., Tang, Y., Zhao, Y.X., Li, X.,

Huang, H.Y., Guo, L., Liu, X.S., Xu, C.J., & Tang, Q.Q. (2016). BMP4 Cross-talks

with estrogen/eralpha signaling to regulate adiposity and glucose metabolism in females. *EBioMedicine,* 1191-100. https://doi.org/10.1016/j.ebiom.2016.07.034

Santos, L., Escande, C., & Denicola, A. (2016). Potential modulation of sirtuins by oxidative stress. *Oxid Med Cell Longev,* 20169831825. https://doi.org/10.1155/2016/9831825

Sasaki, Y., Ikeda, Y., Miyauchi, T., Uchikado, Y., Akasaki, Y., & Ohishi, M. (2020). Estrogen-SIRT1 axis plays a pivotal role in protecting arteries against menopause-induced senescence and atherosclerosis. *J Atheroscler Thromb,* 27(1), 47-59. https://doi.org/10.5551/jat.47993

Serra, C.A., Dos Reis, A.F., Calsa, B., Bueno, C.S., Helaehil, J.V., de Souza, S.A.R., de Oliveira, C.A., Vanzella, E.C., & do Amaral, M.E.C. (2022). Quercetin prevents insulin dysfunction in hypertensive animals. *J Diabetes Metab Disord,* 21(1), 407-417. https://doi.org/10.1007/s40200-022-00987-4

Straub, R.H. (2007). The complex role of estrogens in inflammation. *Endocr Rev,* 28(5), 521-574. https://doi.org/10.1210/er.2007-0001

Tominaga, K., Yamauchi, A., Egawa, T., Tanaka, R., Kawahara, S., Shuto, H., & Kataoka, Y. (2011). Vascular dysfunction and impaired insulin signaling in high-fat diet-fed ovariectomized mice. *Microvasc Res,* 82(2), 171-176. https://doi.org/10.1016/j.mvr.2011.06.004

Vachharajani, V.T., Liu, T., Wang, X., Hoth, J.J., Yoza, B.K., & McCall, C.E. (2016). Sirtuins link inflammation and metabolism. *J Immunol Res,* 20168167273. https://doi.org/10.1155/2016/8167273

Vural, P., Canbaz, M., & Akgul, C. (2006). Effects of menopause and postmenopausal

tibolone treatment on plasma TNFalpha, IL-4, IL-10, IL-12 cytokine pattern and some

bone   turnover   markers.   *Pharmacol   Res,*   53(4),   367-371.

https://doi.org/10.1016/j.phrs.2006.01.005

Weiss, E.P., & Fontana, L. (2011). Caloric restriction: powerful protection for the aging

heart and vasculature. *Am J Physiol Heart Circ Physiol,* 301(4), H1205-1219.

https://doi.org/10.1152/ajpheart.00685.2011

Xie, J., Zhang, X., & Zhang, L. (2013). Negative regulation of inflammation by SIRT1.

Pharmacol Res, 67(1), 60-67. https://doi.org/10.1016/j.phrs.2012.10.010

Yasui, T., Uemura, H., Yamada, M., Matsuzaki, T., Tsuchiya, N., Noguchi, M.,

Yuzurihara, M., Kase, Y., & Irahara, M. (2008). Associations of interleukin-6 with

interleukin-1beta, interleukin-8 and macrophage inflammatory protein-1beta in midlife

women. *Cytokine,* 41(3), 302-306. https://doi.org/10.1016/j.cyto.2007.12.001

Yzydorczyk, C., Li, N., Rigal, E., Chehade, H., Mosig, D., Armengaud, J.B., Rolle, T.,

Krishnasamy, A., Orozco, E., Siddeek, B., Juvet, C., Vergely, C., & Simeoni, U. (2019).

Calorie restriction in adulthood reduces hepatic disorders induced by transient postnatal

overfeeding in mice. *Nutrients,* 11(11). https://doi.org/10.3390/nu11112796

Figures legends

Fig. 1. Metabolic effects of caloric restriction on ovariectomized mice. (A) Experiment timeline. (B) Body mass, (C) Lee index, (D) uterine weight, (E) Fat pad index, and (F) histological examination of the liver and adipose tissue in sham operated (Sham), ovariectomized (OVX), and ovariectomized + 40% caloric restriction (CR) (OVXR) mouse groups. Data are mean ± SEM and were analyzed using one-way ANOVA followed by post-hoc Tukey test. The numbers of mice (n) and p-values are indicated on each graph.

Fig. 2. Biochemical analyses of glycemic homeostasis. (A) Intraperitoneal glucose tolerance test (ipGTT); (B) intraperitoneal insulin tolerance test (ipITT); (C) area under the curve (AUC) of the glucose tolerance test; (D) rate constant for glucose decay (Kitt) during the ITT test; and (E) HOMA-IR, (F) HOMA%S, and (G) HOMA%B indexes. Data are mean ± SEM and were analyzed by one-way ANOVA followed by post-hoc Tukey test. The numbers of mice (n) and p-values are indicated on each graph.

Fig. 3. Biochemical analyses of lipid metabolism. Measurements of (A) serum cholesterol, (B) serum triglycerides, (C) serum HDL, (D) serum LDL, (E) serum VLDL, (F) fasting glucose, (G) fasting insulin, and (H) serum protein. Data are mean ± SEM and were analyzed by one-way ANOVA followed by post-hoc Tukey test. The numbers of mice (n) and p-values are indicated on each graph.

Fig. 4. Typical immunoblots of (A) catalase (CAT), (B) superoxide dismutase (SOD2), and (C) representative blots. (D) Serum TBARS, (E) liver TBARS, and (F) liver H_2O_2 levels in the Sham, OVX, and OVXR groups. Data are mean ± SEM and were analyzed by one-way ANOVA followed by post-hoc Tukey test. The numbers of mice (n) and p-values are indicated on each graph.

580 **Fig. 5.** Typical immunoblots of (A) IL-6, (B) IL-10, (C) TGF-β1, (D) SIRT1, (E)

581 SIRT3, (F) pAMPK/AMPK, (G) pIR/IR, and (H) representative blots in the livers of

582 mice in the Sham, OVX, and OVXR groups. Data are mean ± SEM and were analyzed

583 by one-way ANOVA followed by post-hoc Tukey test. The numbers of mice (n) and p-

584 values are indicated on each graph.

Table 1: Hepatic and adipose characteristics

Parameters	Sham	OVX	OVXR	Statistical difference
Hepatic Chol (mg/100g tissue)	0.001435±0.0001	0.001637±0.0001	0.001796±0.0003	None
Hepatic TG (mg/100g tissue)	0.0005738±0.0001	0.0006981±0.0002	0.001502±0.0003[a,b]	a: p=0.003 vs Sham b: p=0.008 vs OVX
Hepatic Glycogen (mg/100g tissue)	0.8490±0.144	0.2902±0.059[a,c]	1.164±0.131	a: p=0.012 vs Sham c: p=0.0001 vs OVXR
Liver tissue index*	4.660±0.2420	4.454±0.1980	4.930±0.4340	None
Hepatic NAG (D.O/mg tissue)	0.03224±0.0017	0.04614±0.003[a,c]	0.03174±0.0053	a: p= 0.0421 vs Sham c: p= 0.0351 vs OVXR
Adipocytes area (mm^2)	171.5±15.33	411.7±46.23 [a,c]	271.7±26.31	a: p<0.0001 vs Sham c: p=0.0054 vs OVXR

SHAM: control group; OVX, ovariectomized; OVXR (OVX + caloric restriction); *tissue weight (g)/ animal body weight X 100. Data express mean ± std error of mean (n=6/group) and values were compared by one-way ANOVA with Tukey post-test.