The iron-dopamine D1 coupling modulates neural signatures of
working memory across adult lifespan

Jonatan Gustavsson[1], Jarkko Johansson[2], Farshad Falahati[1], Micael Andersson[4], Goran Papenberg[1],

Bárbara Avelar-Pereira[1], Lars Bäckman[1], Grégoria Kalpouzos*[1], Alireza Salami*[1,3,4]

[1]Aging Research Center, Karolinska Institutet and Stockholm University, Sweden. [2]Umeå University,

Faculty of Medicine, Department of Radiation Sciences, Sweden. [3]Wallenberg Center for Molecular

Medicine at Umeå University [4] Department of Integrative Medical Biology, Umeå University, Umeå,

Sweden

*Authors share the position as last authors.

Abstract

Brain iron overload and decreased integrity of the dopaminergic system have been independently

reported as brain substrates of cognitive decline in aging. Dopamine (DA), and iron are co-localized in

high concentrations in the striatum and prefrontal cortex (PFC), but follow opposing age-related

trajectories across the lifespan. DA contributes to cellular iron homeostasis and the activation of D1-

like DA receptors (D1DR) alleviates oxidative stress-induced inflammatory responses, suggesting a

mutual interaction between these two fundamental components. Still, a direct in-vivo study testing

the iron-D1DR relationship and their interactions on brain function and cognition across the lifespan

is rare. Using PET and MRI data from the DyNAMiC study (n=180, age=20-79, %50 female), we

showed that elevated iron content was related to lower D1DRs in DLPFC, but not in striatum,

suggesting that dopamine-rich regions are less susceptible to elevated iron. Critically, older

individuals with elevated iron and lower D1DR exhibited less frontoparietal activations during the

most demanding task, which in turn was related to poorer working-memory performance. Together,

our findings suggest that the combination of elevated iron load and reduced D1DR contribute to

disturbed PFC-related circuits in older age, and thus may be targeted as two modifiable factors for

future intervention.

Keywords: Dopamine, Iron, Working memory, Age, BOLD

Highlights

•   First study demonstrating the association between regional iron and dopamine D1DR in adult

humans.

•   The interplay between age-related elevated iron and diminished D1DR explained lower task-

related brain activity, which in turn was related to poorer task performance.

•   Our findings iron-DA coupling can help progress the understanding of the mechanisms

behind DA-related neurodegeneration.

1 Introduction

Aging is associated with cognitive decline (Gorbach et al., 2017; Salthouse, 2012) and concomitant alterations in structural, functional, and molecular properties of the brain (Andrews-Hanna et al., 2007; Bäckman et al., 2000; Gorbach et al., 2017; Zimmerman et al., 2006). Elevated iron content and decreased integrity of the dopaminergic system are typically reported as independent brain substrates of age-related cognitive decline (Bäckman et al., 2006, 2011; Cools & D'Esposito, 2011; Daugherty et al., 2015; Gustavsson et al., 2022; Hallgren & Sourander, 1958; Kalpouzos, 2018; Landau et al., 2009; Nyberg, Andersson, et al., 2009; Salami et al., 2019). Intracellular non-heme iron is involved in several fundamental neurobiological processes, including dopamine (DA) metabolism (Hare & Double, 2016; Mills et al., 2010). More specifically, iron is a cofactor of tyrosine hydroxylase during DA synthesis, indicating a critical role of iron in DA metabolism as well as in the development of the dopaminergic system (Erikson et al., 2001; Ortega et al., 2007; Unger et al., 2014; Zucca et al., 2017).

Iron is stored in the ferritin protein which keeps it from harming brain cells and is released upon metabolic demand. However, unbound iron accumulates with advancing aging and exerts detrimental effects on cellular integrity. Elevated iron content contributes to poorer myelin integrity (Steiger et al., 2016) and grey-matter atrophy (Daugherty & Raz, 2015, 2016), as well as altered frontostriatal activity (Kalpouzos et al., 2017; Rodrigue et al., 2020; Salami et al., 2021) along with disrupted functional connectivity in aging (Salami et al., 2018). Similar to consequences of elevated iron in aging, disruption of the DA system contributes to age-related neurocognitive impairment (Bäckman et al., 2006, 2011; Cools & D'Esposito, 2011; Landau et al., 2009; Nyberg, Dahlin, et al., 2009; Rieckmann et al., 2011; Salami et al., 2019). Given DA and iron are co-localized in high concentrations in the striatum, age-related iron dyshomeostasis combined with disturbance of the DA system may become harmful for brain integrity and cognition. Direct in-vivo studies testing the

synergistic effects between the dopaminergic system and iron on brain function and cognition in

aging are rare.

Past studies suggest that iron may cause neurotoxicity through DA oxidation, which may in turn

contribute to loss of dopaminergic neurons (Hald & Lotharius, 2005; Hare & Double, 2016; Youdim et

al., 1993; Zucca et al., 2017). Animal studies have demonstrated that iron-induced damage causing

DA depletion (Poetini et al., 2018) and deterioration of dopaminergic cells (Kaur et al., 2003) could be

restored after iron chelation. In contrast, a longitudinal study showed that dopaminergic cell death

precedes iron elevation in parkinsonian monkey (He et al., 2003). A study with young, middle-aged,

and older rhesus monkeys reported an association between iron and stimulus-evoked levels of DA

(Cass et al., 2007), with older monkeys exhibiting more iron accumulation and less DA release.

Although it remains unclear whether elevated iron is the cause or consequence of DA degeneration,

it is relatively well acknowledged in animal studies that these two key chemical components of the

brain interact with each other (Hare & Double, 2016).

As opposed to ample animal studies on iron-DA coupling, evidence for such an association across the

adult lifespan in humans is scarce. A recent longitudinal study showed that developmental changes in

pre-synaptic, but not post-synaptic, DA receptors were positively associated with iron accumulation

through adolescence and young adulthood (Larsen et al., 2020). Still, iron and DA play ambivalent

roles during early and later adulthood (Johansson et al., 2022; Kalpouzos et al., 2017; Salami et al.,

2021). Recent work showed that older individuals with genetic predisposition for lower DA (by proxy

of the COMT Val158Met genetic polymorphism) accumulated more iron in the dorsolateral

prefrontal cortex (DLPFC) and striatum over 3 years (Gustavsson et al., 2022). Postsynaptic DA

markers, particularly DA D1-like receptor (D1DRs) which are the most abundantly expressed receptor

subtypes, are among the most age-sensitive DA systems (Karrer et al., 2019). Moreover, the

activation of D1 and D2 receptors alleviates oxidative stress-induced inflammatory responses (Shao

et al., 2013; Yan et al., 2015; Zhu et al., 2018). Hence, with diminished DADRs, the capacity of the

protective response counteracting oxidative stress and neuroinflammation due to iron overload is

lessened, and may lead to increased damage and ferroptosis (Hald & Lotharius, 2005).

Motivated by the lack of human studies, we investigated the relationship between dopaminergic

receptors and iron content, and their interactions on brain function and cognition. Using the largest

D1DR dataset across the world, we studied 180 healthy individuals (20-79 years old) who underwent

D1DR Positron Emission Tomography (PET) assessment using [^{11}C]SCH23390, iron approximation

made with magnetic resonance imaging (MRI) based quantitative susceptibility mapping (QSM;

(Langkammer et al., 2012)), and functional magnetic resonance imaging (fMRI) while performing a

working-memory task. We first examined regional variation in the link between D1DR and iron

content with the hypothesis that greater iron content is related to lower D1DR, with a possible

interaction with age (C.f. Salami et al., 2021; Kalpouzos et al.,2018). We targeted DLPFC and striatum,

because of abundantly expressed D1DR (Shohamy & Adcock, 2010), pronounced age-related D1DR

differences (Karrer et al., 2017), and association between *COMT* polymorphism (i.e., a dopaminergic

gene) and iron accumulation reported in these regions (Gustavsson et al., 2022). Given the

association of iron and DA to neural circuits of working memory (c.f., Salami et al., 2021; Salami et al.,

2019), we predicted an interactive effect of iron and D1DR on load-dependent BOLD modulation and

working memory processing (Gustavsson et al., 2022; Spence et al., 2020). To this end, we applied

multivariate partial-least squares (PLS; ((McIntosh & Lobaugh, 2004)) which enables a simultaneous

analysis of iron content, D1DR, age, on BOLD associations across all task conditions in a data-driven

fashion. If these variables are simultaneously related to neural circuits of WM, PLS should reveal a

single network showing that older individuals with elevated iron and decreased D1DR (i.e., toxic iron-

DA coupling) exhibit lower task-related BOLD-response, particularly within the frontoparietal

network. However, if D1DR and iron differentially modulate BOLD response (c.f., Salami et al., 2019;

Salami et al., 2021), PLS should reveal different networks.

2 Materials and methods

This study uses data from the DopamiNe Age Connectome Cognition (DyNAMiC) project approved by

the Umeå University Regional Ethical Board. All participants signed informed consent prior to data

collection (for details about DyNAMiC, see Nordin et al., 2022).

2.1 Participants

One-hundred and eighty adults (mean age 49.8 ± 17.4 years; range 20-79; 90 females) from Umeå,

Sweden, were recruited to participate in the project and underwent the full protocol, including

cognitive testing, MRI, and PET assessments. All participants were native Swedish speakers, right-

handed, and had no history of neurological illnesses. Of the sample of 180 participants, 3 dropped

out of PET scanning, 4 were excluded due to incidental brain abnormalities, 3 due to failed

reconstruction of QSM images. Additionally, brain and behavioural data were screened for univariate

and multivariate outliers. Univariate outliers were defined as values greater or less than 3.29 SD from

the mean (Tabachnick & Fidell, 2013) and excluded as pairwise deletions per ROIs and modality.

Multivariate outliers were identified according to Mahalanobis distance ($p < 0.001$; Tabachnick &

Fidell, 2013). As a result, five participants were identified as univariate outliers on their iron (n=3) or

D1DR (n=2) values, and three were multivariate outliers. Thus, data from 162 individuals were used

in the analyses involving iron content and D1D1R. Furthermore, three outliers were identified based

on their online WM performance and excluded from analyses.

2.2 Neuroimaging acquisition and pre-processing

Participants underwent positron emission tomography on a Discovery 690 PET/CT scanner (General

Electric) and MRI on a Discovery MR750 3.0 T scanner (General Electric) equipped with a 32-channel

phased-array head coil at two separate occasions at Umeå University Hospital.

2.2.1 PET acquisition

PET was conducted to assess whole-brain D1DR using radioligand [^{11}C]SCH23390 at rest (for details

see Nordin et al., 2022). The scanning session started with a low-dose CT image, followed by an

intravenous bolus injection of the radioligand. Participants were instructed to lay still and stay awake

with eyes open. An individually fitted thermoplastic mask was attached to the bed surface during

scanning to minimize head movement.

2.2.2 Structural MR acquisition

High-resolution anatomical T1-weighted images were acquired using 3D fast-spoiled gradient echo

sequence with the following parameters: 176 sagittal slices, slice thickness = 1 mm, voxel size 0.49 x

0.49 x 1 mm^3, repetition time (TR) = 8.2 ms, echo time (TE) = 3.2 ms, flip angle = 12°, field of view

(FOV) = 250 x 250 mm, no spacing.

2.2.3 Iron acquisition

Images for iron estimation was obtained using a 3D multi-echo gradient-recalled echo sequence

(meGRE). The parameters were as follows: 124 axial slices, voxel size 1 x 1 x 1.2 mm^3, TR = 31 ms, flip

angle = 17°, FOV = 256 x 256 mm, no spacing. The first TE was 1.78 ms followed by seven additional

TEs with a 2.87 ms interval.

2.2.4 Functional MR acquisition

The functional images were sampled using single-shot multiband EPI sequence with 37 axial slices,

voxel size 1.95 x 1.95 x 3.9 mm^3, 0.5 mm spacing, TR = 2.000 ms, TE = 30 ms, flip angle = 80°, and FOV

= 250 × 250 mm. Ten dummy scans were collected at the start of the sequence.

2.2.5 In-scanner working memory task

A numerical N-back task was administered in the scanner (Salami et al., 2018). A sequence of single

numbers appeared on the screen for 1.5s, with an interstimulus interval of 0.5s. During every item

presentation, subjects indicated whether the digit on the screen was the same as the one shown 1, 2,

or 3 digits back by pressing one of the two adjacent buttons with the index (Yes response) or middle

finger (No response). Each working-memory load condition (1-, 2-, and 3-back) was presented over

nine blocks in random order (interblock interval, 22 s) with each block consisting of 10 trials. For

every block, 10 trials were performed with four matches (requiring a "yes" response) and six

nonmatches (requiring a "no" response). The N-back blocks were counterbalanced and trial sequence

was the same for all participants. The maximum score for each condition was 81, 72, 63, respectively.

Performance was calculated as an error-adjusted discrimination score by subtracting the proportion

of false alarms (i.e., a wrong answer judged to be correct, or, in other terms, answering Yes, when

the correct answer is No) from the proportion of correct hits (i.e., answering Yes when the correct

answer is Yes).

2.2.6 PET processing

PET data obtained with $[^{11}C]SCH23390$ was processed with the following steps: To estimate receptor

availability (i.e., D1DR) in targeted regions, binding potential relative to non-displaceable binding in a

reference region (BP_{ND}; Innis et al., 2007) was used with cerebellum as reference. The processing of

the PET data included correction for head movement by using frame-to-frame image co-registration,

and co-registered with T1-weighted MRI data with re-slicing to MRI voxel size using Statistical

Parametric   Mapping   (SPM12:   Wellcome   Trust   Centre   for   Neuroimaging,

http://www.fil.ion.ucl.ac.uk/spm/). To model the regional time-activity course (TAC) data we used

simplified reference tissue model (SRTM; Lammertsma & Hume, 1996).

2.2.7 MRI processing

Quantitative susceptibility mapping. Processing of QSM is referred from Gustavsson et al. (2022).

Approximation of iron content was inferred from susceptibility values derived from QSM images.

Morphology-enabled dipole inversion (MEDI; T. Liu et al., 2011) is a method for QSM estimation that

selects the solution that minimizes the discrepancy in the number of voxels belonging to edges

between the susceptibility image and the magnitude image. Here, we used the recommended

nonlinear variant of MEDI proposed by Liu et al. (2013). Initially, the total field map was estimated

from the complex meGRE images by performing a nonlinear least square fitting on a voxel-by-voxel

basis. The resulting frequency map was then spatially unwrapped using a guided region-growing

unwrapping algorithm (Xu & Cumming, 1999). The background fields, the superimposed field

contributions that are not caused by the sources inside the brain and mainly generated by air-tissue

interference, were eliminated using a nonparametric technique based on Projection onto Dipole

Fields (PDF: Liu et al., 2011). Finally, the corrected frequency map was used as input for the field-to-

source inverse problem to calculate susceptibility maps. The MEDI Toolbox

(http://weill.cornell.edu/mri/pages/qsm.html) was used to calculate QSM images.

Due to the singularity of dipole kernel at the centre of k-space, the generated QSM images contain

relative susceptibility values, which may not necessarily be comparable across subjects. To address

this issue, a region of the corticospinal tract was selected as a zero-reference region due to its

resilience to age-related degeneration (de Groot et al., 2015) and stable susceptibility across

adulthood (Li et al., 2014). Using white-matter areas as reference regions has previously been

recommended due to their low standard deviations of susceptibility, indicating low inter-subject

variation similar (Straub et al., 2017) or even lower than CSF (Deistung et al., 2013). The process of

zero-referencing is described in detail by Garzón and colleagues (2017).

Automated segmentation of cortical and deep gray-matter structures was performed with the

Freesurfer image analysis suite — version 6 (http://surfer.nmr.mgh.harva rd.edu/) using T1-weighted

images (Fischl et al., 2002; Fischl, Salat, et al., 2004; Fischl, Van Der Kouwe, et al., 2004).

Next, QSM and the segmentation results were resampled to the native structural space. Then,

statistics including average and standard deviation were computed on the QSM maps. We merged

segmented rostral and caudal middle frontal regions from the left and right hemispheres to form

DLPFC (Fig. 1A), and the left and right caudate and putamen to form striatum (supplementary

materials Fig. 1A).

Prior to computing statistics on the QSM maps, the boundary of segmentations was eroded by one

voxel, and a fraction (15%) of the most extreme values was removed to avoid the influence of high

signal from neighbouring vessels and obtain more robust estimates (Garzón et al., 2017).

2.2.8 Functional MRI processing

Pre-processing of the fMRI data, performed in SPM12 software, included slice-timing correction and

motion correction by unwarping and re-alignment to the first image of each volume. The fMRI

volumes were then normalized to a sample-specific template generated using Diffeomorphic

Anatomical Registration using Exponentiated Lie algebra (DARTEL: Ashburner, 2007), affine

alignment to MNI standard space, and spatial smoothing with a 6-mm full width at half maximum

(FWHM) Gaussian kernel (voxel size = $2 \times 2 \times 2$ mm^3).

The pre-processed fMRI data were analysed with spatiotemporal PLS (McIntosh et al., 2004;

McIntosh & Lobaugh, 2004) to assess the BOLD association with iron, D1DR, and age across the three

experimental WM conditions (1-, 2-, and 3-back). PLS determines time-varying distributed patterns of

neural activity as a function of experimental variables (1-, 2-, and 3-back) and regional iron and

D1DR. This technique allows for the identification of patterns/networks, which reflect association

changes across all regions of the brain simultaneously, rather than assemblies of independent

regions, thus ruling out the need for multiple-comparison correction. A detailed description of

spatiotemporal PLS analysis for fMRI data has been given in previous reports (Garrett et al., 2010;

Grady & Garrett, 2014; Salami et al., 2010, 2012, 2014).

The onset of each stimulus within each block of images (1-, 2-, and 3-back) was averaged across

blocks for each condition. A cross-block correlation matrix was computed as the correlation between

neural activity across experimental conditions (1-, 2-, and 3-back) and variables of interest (age,

D1DR, iron) across different regions. Then, the correlation matrix was decomposed using singular

value decomposition (SVD), to identify a set of orthogonal latent variables (LVs) representing linear

combinations of the original variables:

SVD $_{CORR}$ = USV ⍰

This decomposition produces a left singular vector of regional susceptibility weights (U), a right

singular vector of BOLD weights (V), and a diagonal matrix of singular values (S). This analysis

produces orthogonal LVs that optimally represent relations between the variables of interest (age,

D1DR, iron) and BOLD. Note that PLS is not mathematically susceptible to collinearity similar to the

multiple regression approach (Leibovitch et al., 1999). Each LV contains a spatial pattern exhibiting

the brain regions whose activity shows the strongest simultaneous relations to the input variables. To

obtain a summary measure of each participant's expression of a particular LV pattern, subject-

specific "brain scores" are computed by multiplying each voxel's (i) weight (V) from each LV (j) by the

BOLD value in that voxel for person (m), and summing over all (n) brain voxels:

$$\sum_{i=l}^{n} \mathrm{v}_{ij} \mathrm{BOLD}_{im}$$

Taken together, a brain score represents the degree to which each subject contributes to the

multivariate spatial pattern captured by a latent variable.

The statistical significance of each LV was assessed using permutation testing. This procedure

involved reshuffling the rows of the data matrix and recalculating the LVs of the reshuffled matrix

using the same SVD approach. The number of times a permuted singular value exceeds the original

singular value yields the probability of significance of the original LV (McIntosh et al., 1996). In the

present study, 1000 permutations were performed. In addition, the stability of voxel saliencies

contributing to each LV was determined with bootstrap estimation of standard errors (SEs), using

1000 bootstrap samples (Efron & Tibshirani, 1986). The Bootstrap Ratio (BSR: the ratio between

voxel saliences and estimated SEs) was computed and voxels with BSR > 3.29 (similar to a Z-score of

3.29, corresponding to p = 0.001) were considered reliable. All reliable clusters comprised contiguous

voxels, with a distance between clusters of at least 10 mm. Moreover, the upper and lower

percentiles of the bootstrap distribution were used to generate 95% confidence intervals (CIs)

around the correlation scores to facilitate interpretation (McIntosh & Lobaugh, 2004). For instance, a

significant difference between correlation scores in different conditions is indicated by non-

overlapping CIs. Similarly, brain or correlation scores were considered unreliable when CIs

overlapped with zero.

PLS uses all conditions of an experiment and variables of interest at once, thus offering an additional

dimension by simultaneously considering both similarities and differences across the experimental

variables. If the variables of interest (i.e., age, iron content, D1DR) are similarly related to brain

regions, PLS reveals a pattern with reliable loadings (with/without quantitative differences) for all

variables. If a single variable dominates the pattern, PLS should reveal a reliable loading for that

variable only. Alternatively, if different variables (e.g., D1DR and iron) differentially modulate BOLD

response (e.g., Load-dependent effects of dopamine on BOLD as shown in Salami et al., (2019) versus

load-independent effect of iron on BOLD shown in Salami et al., 2021), PLS may reveal two distinctive

networks.

2.3 Additional statistical analyses

To assess age-effects on cognitive performance, a multivariate analysis of covariance (MANCOVA)

was conducted with WM load-conditions as dependent variables and age-groups (younger (age 20-

39) vs. middle-aged (age 40-59) vs. older (age 60-79)) as between-subjects factors. Follow-up

independent t-tests were conducted to assess significant differences between age groups. To assess

the relationship between iron content and D1DR, we conducted partial correlation analyses for each

region of interest with iron content and D1DR as dependent variables, controlling for age. As control

analyses, we performed the same analyses but included sex, education, and regional grey-matter

volume as covariates.

3. Results

3.1 Demographics

Demographic information along with data on body mass index (BMI), brain volumes, D1DR, and N-

back performance are presented in Table 1.

Table 1. Participant characteristics			
		Mean	**SD**
N (Women)	162 (79)		
Age		50	17
Age range		20-78	
MMSE		28.8	1.1
BMI		25.2	3.9
Education		15	3.4
Volume[a]			
DLPFC		20478	2696
Striatum		11473	1282
Iron[b]			
DLPFC		0.0430	0.0140
Striatum		0.0998	0.0286
D1DR			
DLPFC		0.3582	0.0690
Striatum		1.6130	0.2182
WM performance			
1-back accuracy[c]		75.18	6.65
1-back (H-FA)[d]		0.86	
2-back accuracy		60.99	8.77
2-back (H-FA)		0.72	
3-back accuracy		48.10	9.57
3-back (H-FA)		0.59	

Note. MMSE = Mini Mental State Examination,
DLPFC = Dorsolateral prefrontal cortex, SD = Standard deviation.
[a] Volume (mL) adjusted for total intracranial volume
[b] Approximation of iron, based on susceptibility in parts per million
[c] Accuracy calculated as the sum of correct responses
[d] Proportion of false alarms (FA) subtracted from proportion of hits (H)

The MANCOVA conducted on N-Back performance showed a significant main effect on load conditions ($F_{2,155}$ = 228.24, $p < 0.001$, Wilk's Λ = 0.253, partial η^2 = 0.747), and an age-group effect on load conditions ($F_{4,310}$ = 9.43, $p < 0.001$, Wilk's Λ = 0.795, partial η^2 = 0.108). Post-hoc analysis for the lowest WM load (1-back) showed that the older participants performed less accurately compared to both middle-aged (t_{89} = -3.974, $p < 0.001$) and younger participants (t_{71} = -5.707, $p < 0.001$). There was no significant difference between younger and middle-aged participants ($p = 0.07$). For 2-back, older participants performed significantly poorer compared to both middle-aged (t_{105} = -4.015, $p < 0.001$)

and younger participants (t_{85} = -8.905, p < 0.001). Middle-aged participants performed significantly

poorer compared to younger participants (t_{98} = -4.176, p < 0.001). Lastly, for highest WM load (3-

back), older participants performed significantly poorer compared to both middle-aged (t_{105} = -5.300

p < 0.001) and younger participants (t_{102} = -7.950, p < 0.001). Middle-aged participants performed

significantly poorer compared to younger participants (t_{105} = -2.921, p = 0.004).

3.2 Iron content in DLPFC, but not in striatum, was negatively associated with D1DR

Both striatal and DLPFC iron increased with advancing age (striatum: r = 0.551, p < 0.001; DLPFC: r =

0.244, p = 0.002). D1DRs in both regions decreased with advancing age (striatum: r = -0.62, p < 0.001;

DLPFC: r = -0.565, p = < 0.001).

The partial correlation analysis for iron content and D1DR in DLPFC showed a significant negative

association across the whole sample (r = -0.272, p < 0.001), suggesting that greater iron content was

linked to lower D1DR (Fig. 1B). Further group level analyses revealed that this correlation was

similarly expressed across different age groups (Younger: r = -0.10, p = 0.48; Middle-aged: r = -0.30, p

= 0.026; Older: r = -0.37, p = 0.006). However, striatal iron content was unrelated to D1DR across the

whole sample (r = -0.101, p = 0.20; supplementary materials, Fig. 1B). No significant association was

observed within each age groups (Ps > 0.05), except for a trend-level link in the older individuals (r = -

0.24, p = 0.08).

Figure 1. The relationship between iron content and D1DR in dorsolateral prefrontal cortex. **(A)**
Surface map representing distribution of cortical iron (susceptibility in parts per million; left column)
and D1DR ([^{11}C]SCH23390 BP$_{ND}$; right column) with dorsolateral prefrontal cortex (DLPFC) outlined.
(B) Scatterplot depicting the association between greater iron content and lower D1DR in DLPFC. All
values are z-transformed residuals adjusting for age.

3.3 Iron-D1DR couplings in DLPFC modulates WM-related BOLD in a load-dependent fashion

Given the iron-D1DR link in DLPFC, we next investigated whether iron-D1DR coupling modulated
neural correlates of PFC-related function across the adult lifespan. We used behavioural PLS to assess
the presence of multivariate spatial patterns of task-related BOLD response dependent on age, iron
content, and D1DR across load conditions during N-back working memory task. The analysis
identified two significant latent variables (LVs). LV1 represented a brain-wide network with increased
activation in older individuals in a load-independent fashion. This network was largely unrelated to
D1DR or iron in DLPFC (supplementary materials 5.1.2, Table 1, Fig 2).

The second LV represented the canonical WM network (c.f. Nagel et al., 2009; Salami et al., 2018).
This LV (permuted p = 0.02, 17.97% of cross-block covariance: Fig. 3A) demonstrated that older
individuals with elevated iron content and lower D1DR in DLPFC exhibited lower BOLD response in
the frontoparietal network during 3-back (Fig. 3B). In contrast, higher BOLD response in the

frontoparietal network during 1-back was associated with older age and lower D1DR in DLPFC, but

not with iron content. For 2-back, no associations were considered reliable as the CI:s overlapped

with zero (Fig. 3B). As opposed to the frontoparietal WM network, the default-mode network (Fig. 3A

(Blue areas); supplementary Table 2) was less deactivated during 3-back in older individuals with

higher iron content and lower D1DR in DLPFC. Taken together, these results revealed that iron-D1DR

coupling modulates neural correlates of the working memory in a load-dependent fashion.

Figure 3. Multivariate relationship of BOLD response patterns during working-memory task identified

by task PLS. **A)** The regions depicted in a yellow colour mainly correspond to the frontoparietal

network, whereas the blue colour mainly represents the default-mode network. The frontoparietal

network showed greater activation in correspondence to the behavioural measures across different

task conditions, whereas the default-mode network showed greater deactivation (i.e., less

activation). **B)** BOLD association across age, iron content (QSM), and D1DR. Interpretation of the

figure should be as reliable positive and negative associations of activation (BOLD response) when

the confidence intervals (CI:s) do not overlap with zero. As the frontoparietal network displayed

increased activation, a positive association is interpreted as greater activation, whereas a negative

association is interpreted as less activation. For the default-mode network the opposite applies. **C)**

Greater activation was associated with poorer performance during 1-back, but greater performance

during 3-back. All values are z-transformed residuals adjusting for age.

3.4 Load-dependent BOLD response is differentially associated with task performance.

We have shown that increasing age was concomitant with increased and decreased activations

within the frontoparietal network during 1-back and 3-back, respectively. We next tested the

relationship of brain activation in relation to task performance. The brain score during 1-back was

negatively associated with 1-back performance ($r = -0.324$, $p < 0.001$; Fig. 3C), but positively with 3-

back ($r = 0.216$, $p = 0.006$; Fig. 3C). No significant association was observed during 2-back ($r = 0.05$,

$p=0.4$). Taken together, our results suggest that less frontoparietal activity during 3-back as well as

greater frontoparietal activity during 1-back are associated with less efficient working memory

function.

3.5 Control analyses

To confirm that the results obtained from the partial correlation analyses were not due to

confounding variables, we performed additional analyses in which we controlled for sex, education,

and regional grey-matter volume. The inclusion of the additional variables did not result in a

noticeable change of the outcome and all the patterns remained consistent. A full description of the

results is given in the supplementary results.

4. Discussion

In this study, we provide first in-vivo evidence for an association between D1DR and brain iron across

the adult lifespan, and how iron-D1DR synergy may contribute to disturbed brain responses during

WM task performance across the lifespan. The main findings were that greater iron content was

associated with lower D1DR in DLPFC, but not in striatum, across the adult lifespan. We also found

that older individuals with elevated iron content and lower D1DR in DLPFC – a presumably

deleterious synergy – exhibited lower frontoparietal activity along with less deactivation of the DMN

during high demand condition and in turn poorer WM performance. Although the associations

revealed by these data are cross-sectional, a plausible interpretation is that elevated iron and lower

D1DR together form a toxic couple (c.f. Hare & Double, 2016), which would contribute to reduced

network dynamics and impaired WM processing.

A key finding of the present study is the association between higher iron content and lower D1DR in

DLPFC across the adult lifespan. The exact mechanism underlying the association between iron

content and DA receptors is poorly understood. The dopaminergic system is important to cellular

iron homeostasis, as documented in *in vitro* studies (Dichtl et al., 2018; Liu et al., 2021). Supporting

this idea, we recently demonstrated that older adults with genetic predisposition for lower DA levels

accumulated more iron in DLPFC and striatum over time compared to individuals with presumably

higher DA levels (Gustavsson et al., 2022). Age-related differences in pre- and postsynaptic DA

markers (Bäckman et al., 2006, 2010) may also contribute to iron dyshomeostasis in aging, resulting

in disruption of transportation of iron via transferrin and storage in the ferritin protein (Ward et al.,

2014). Disruption of iron handling may lead to increase of ferrous iron, which triggers oxidative

stress, neuroinflammation, and cell loss due to ferroptosis (Daugherty et al., 2015; Mazhar et al.,

2021; Salami et al., 2021). The activation of D1 and D2/3 receptors alleviates oxidative stress-induced

inflammatory responses (Shao et al., 2013; Yan et al., 2015; Zhu et al., 2018). However, whether age-

related loss of DA receptors leads to an iron-related increase of oxidative stress and ferroptosis, or

whether an age-related increase of iron causes loss of DA receptor cells can only be determined in a

longitudinal setting.

We did not observe any significant association between iron and D1DR in striatum, except for a

trend-level negative association in older age. The reason for these regional variations is unclear. One

possibility is that, as D1DR are more expressed in striatum compared to cortex (Shohamy & Adcock,

2010), striatal regions are less vulnerable or better at attenuating the damage from higher iron

content. If too many receptors diminishes, or the threshold of the capacity for dealing with oxidative

stress is exceeded, the iron-induced damage to the cells may lead to ferroptosis (Lillig et al., 2008).

This concords well with Parkinson's Disease studies reporting regional variations in synuclein

pathology despite brain-wide iron accumulation (McCann et al., 2016), suggesting that additional

factors may contribute to iron serving a pathological role.

The multivariate PLS analysis revealed that in older age the combination of higher iron and lower

D1DR within DLPFC was related to high frontoparietal recruitment during low-demanding tasks along

with weak frontoparietal upregulation at higher task demand, which in turn was associated with

poorer working memory performance. These results concord with the Compensation-Related

Utilization of Neural Circuits Hypothesis (CRUNCH; (Cappell et al., 2010; Mattay et al., 2006; Nyberg

et al., 2014; Reuter-Lorenz & Cappell, 2008; Schneider-Garces et al., 2010)), which postulates that

neural activity increases at low demanding task-levels in older adults compared to younger adults,

but is reduced at more demanding levels.

The unique and shared contributions of iron content and neuroinflammation to astrocytic

dysfunction in the neurovascular coupling has been proposed to account for the reduced BOLD

response (Kalpouzos et al., 2017; Salami et al., 2018, 2021). Higher iron content has been related to

inflammation (Haider, 2015; Salami et al., 2021), which can be detrimental to cells during prolonged

periods (Hald & Lotharius, 2005; Zecca et al., 2004), thus contributing to astrocytic dysfunction. In

vitro studies have demonstrated that DA receptors activation may alleviate and supress

neuroinflammation (Liu et al., 2021; Shao et al., 2013; Yan et al., 2015). A more efficient

dopaminergic system (e.g., manifested by greater receptor availability) protect against negative

effects of iron and neuroinflammation on brain function. In support of this notion, a past study

showed that age-related D1DR alteration alone may contribute to less efficient engagement of

working memory circuits (Bäckman et al., 2011). Relatedly, the importance of DA for the

frontoparietal BOLD response has been further demonstrated through pharmacological

administration of a dopaminergic antagonist, leading to poorer working-memory performance

(Fischer et al., 2010). Our results add novel contribution to the previous work, by showing that the

combination of elevated iron and D1DR reduction in DLPFC in aging, possibly reflecting synergistic

iron-D1DR effects, exerts a deleterious effect in neural circuits of WM. Furthermore, during the most

demanding WM condition (3-back), lower activation in the frontoparietal network was related to

worse performance, in concordance with previous reports (Salami et al., 2021). Longitudinal data are

needed to identify the primary mechanism of disturbed working memory circuit in older age.

Our study is the first to link regional iron content to DA receptor availability, by showing that greater

iron content is related to lower D1DR. Critically, an interplay between age-related elevated iron

content and diminished D1 receptor availability may provide a mechanistic understanding of how

iron-DA coupling may exert deleterious effects on neural function and ultimately cognition. Elevated

brain iron has been implicated in several neurodegenerative disorders, including Parkinson's disease,

which is characterized by loss of dopaminergic cells (Ward et al., 2014). Thus, the observed findings

have implications for better understanding the mechanisms behind DA-related neurodegeneration.

Author credit statement

Conceptualisation: A.S. and G.K.; Software, M.A.; Methodology, J.G., G.K., G.P., A.S.; Formal Analysis

J.G., G.K., A.S., J.J., F.F., G.P., M.A., and B.A.P.; Visualisation, J.G. Writing – original draft, J.G., G.K.,

A.S., F.F.; Writing – Review & Editing G.P., J.J., L.A., B.A.P.

Acknowledgement

We thank the staff of the DyNAMiC project, Frida Magnusson and Emma Simonsson, staff at MRI and

PET labs at Umeå University Hospital, and all our participants. We also thank Robin Pedersen for his

contribution with the figures.

Conflict of interest

The authors declare no conflict of interest.

Ethics approval

This study was approved by the Regional Ethical board and the local Radiation Safety Committee in

Umeå, Sweden.

Data availability statement

Data from the DyNAMiC project cannot be made publicly available due to ethical and legal

restrictions. However, access to these original data may be available upon request from the

corresponding author.

5. References

Andrews-Hanna, J. R., Snyder, A. Z., Vincent, J. L., Lustig, C., Head, D., Raichle, M. E., & Buckner, R. L.
(2007). Disruption of Large-Scale Brain Systems in Advanced Aging. *Neuron*, *56*(5), 924–935.
https://doi.org/10.1016/j.neuron.2007.10.038

Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *NeuroImage*, *38*(1), 95–113.
https://doi.org/10.1016/j.neuroimage.2007.07.007

Bäckman, L., Ginovart, N., Dixon, R. A., Wahlin, T. B. R., Wahlin, Å., Halldin, C., & Farde, L. (2000).
Age-related cognitive deficits mediated by changes in the striatal dopamine system. *American*
Journal of Psychiatry, *157*(4), 635–637. https://doi.org/10.1176/appi.ajp.157.4.635

Bäckman, L., Karlsson, S., Fischer, H., Karlsson, P., Brehmer, Y., Rieckmann, A., MacDonald, S. W. S.,
Farde, L., & Nyberg, L. (2011). Dopamine D1 receptors and age differences in brain activation
during working memory. *Neurobiology of Aging*, *32*(10), 1849–1856.
https://doi.org/10.1016/j.neurobiolaging.2009.10.018

Bäckman, L., Nyberg, L., Lindenberger, U., Li, S. C., & Farde, L. (2006). The correlative triad among
aging, dopamine, and cognition: Current status and future prospects. *Neuroscience and*
Biobehavioral Reviews, *30*(6), 791–807. https://doi.org/10.1016/j.neubiorev.2006.06.005

Cappell, K. A., Gmeindl, L., & Reuter-Lorenz, P. A. (2010). Age differences in prefontal recruitment
during verbal working memory maintenance depend on memory load. *Cortex*, *46*(4), 462–473.
https://doi.org/10.1016/j.cortex.2009.11.009

Cass, W. A., Grondin, R., Andersen, A. H., Zhang, Z., Hardy, P. A., Hussey-Andersen, L. K., Rayens, W.
S., Gerhardt, G. A., & Gash, D. M. (2007). Iron accumulation in the striatum predicts aging-
related decline in motor function in rhesus monkeys. *Neurobiology of Aging*, *28*(2), 258–271.
https://doi.org/10.1016/j.neurobiolaging.2005.12.010

Cools, R., & D'Esposito, M. (2011). Inverted-U–Shaped Dopamine Actions on Human Working
Memory and Cognitive Control. *Biological Psychiatry*, *69*(12), e113–e125.
https://doi.org/https://doi.org/10.1016/j.biopsych.2011.03.028

Daugherty, A. M., Haacke, E. M., & Raz, N. (2015). Striatal iron content predicts its shrinkage and

changes in verbal working memory after two years in healthy adults. *Journal of Neuroscience*,
35(17), 6731–6743. https://doi.org/10.1523/JNEUROSCI.4717-14.2015

Daugherty, A. M., & Raz, N. (2015). Appraising the Role of Iron in Brain Aging and Cognition: Promises
and Limitations of MRI Methods. *Neuropsychology Review*, *25*(3), 272–287.
https://doi.org/10.1007/s11065-015-9292-y

Daugherty, A. M., & Raz, N. (2016). Accumulation of iron in the putamen predicts its shrinkage in
healthy older adults: A multi-occasion longitudinal study. *NeuroImage*, *128*(1), 11–20.
https://doi.org/10.1016/j.neuroimage.2015.12.045

de Groot, M., Ikram, M. A., Akoudad, S., Krestin, G. P., Hofman, A., Lugt, A., Niessen, W. J., &
Vernooij, M. W. (2015). Tract-specific white matter degeneration in aging: The Rotterdam
Study. *Alzheimer's & Dementia*, *11*(3), 321–330. https://doi.org/10.1016/j.jalz.2014.06.011

Deistung, A., Schäfer, A., Schweser, F., Biedermann, U., Turner, R., & Reichenbach, J. R. (2013).
Toward in vivo histology: A comparison of quantitative susceptibility mapping (QSM) with
magnitude-, phase-, and R2*-imaging at ultra-high magnetic field strength. *NeuroImage*, *65*,
299–314. https://doi.org/10.1016/j.neuroimage.2012.09.055

Dichtl, S., Haschka, D., Nairz, M., Seifert, M., Volani, C., Lutz, O., & Weiss, G. (2018). Dopamine
promotes cellular iron accumulation and oxidative stress responses in macrophages.
Biochemical Pharmacology, *148*, 193–201. https://doi.org/10.1016/j.bcp.2017.12.001

Efron, B., & Tibshirani, R. (1986). Bootstrap Methods for Standard Errors, Confidence Intervals, and
Other Measures of Statistical Accuracy. *Statistical Science*, *1*(1), 15–51.
https://doi.org/10.1214/ss/1177013815

Erikson, K. M., Jones, B. C., Hess, E. J., Zhang, Q., & Beard, J. L. (2001). Iron deficiency decreases
dopamine D1 and D2 receptors in rat brain. *Pharmacology Biochemistry and Behavior*, *69*(3–4),
409–418. https://doi.org/10.1016/S0091-3057(01)00563-9

Fischer, H., Nyberg, L., Karlsson, S., Karlsson, P., Brehmer, Y., Rieckmann, A., MacDonald, S. W. S.,
Farde, L., & Bäckman, L. (2010). Simulating Neurocognitive Aging: Effects of a Dopaminergic
Antagonist on Brain Activity During Working Memory. *Biological Psychiatry*, *67*(6), 575–580.
https://doi.org/10.1016/j.biopsych.2009.12.013

Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., Van Der Kouwe, A., Killiany,
R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., & Dale, A. M. (2002). Whole
brain segmentation: Automated labeling of neuroanatomical structures in the human brain.
Neuron, *33*(3), 341–355. https://doi.org/10.1016/S0896-6273(02)00569-X

Fischl, B., Salat, D. H., Van Der Kouwe, A. J. W., Makris, N., Ségonne, F., Quinn, B. T., & Dale, A. M.
(2004). Sequence-independent segmentation of magnetic resonance images. *NeuroImage*,
23(SUPPL. 1), 69–84. https://doi.org/10.1016/j.neuroimage.2004.07.016

Fischl, B., Van Der Kouwe, A., Destrieux, C., Halgren, E., Ségonne, F., Salat, D. H., Busa, E., Seidman, L.
J., Goldstein, J., Kennedy, D., Caviness, V., Makris, N., Rosen, B., & Dale, A. M. (2004).
Automatically Parcellating the Human Cerebral Cortex. *Cerebral Cortex*, *14*(1), 11–22.
https://doi.org/10.1093/cercor/bhg087

Garrett, D. D., Kovacevic, N., McIntosh, A. R., & Grady, C. L. (2010). Blood oxygen level-dependent
signal variability is more than just noise. *Journal of Neuroscience*, *30*(14), 4914–4921.
https://doi.org/10.1523/JNEUROSCI.5166-09.2010

Garzón, B., Sitnikov, R., Bäckman, L., & Kalpouzos, G. (2017). Can transverse relaxation rates in deep
gray matter be approximated from functional and T2-weighted FLAIR scans for relative brain

iron quantification? *Magnetic Resonance Imaging*, *40*, 75–82.
https://doi.org/10.1016/j.mri.2017.04.005

Gorbach, T., Pudas, S., Lundquist, A., Orädd, G., Josefsson, M., Salami, A., de Luna, X., & Nyberg, L.
(2017). Longitudinal association between hippocampus atrophy and episodic-memory decline.
Neurobiology of Aging, *51*, 167–176. https://doi.org/10.1016/j.neurobiolaging.2016.12.002

Grady, C. L., & Garrett, D. D. (2014). Understanding variability in the BOLD signal and why it matters
for aging. *Brain Imaging and Behavior*, *8*(2), 274–283. https://doi.org/10.1007/s11682-013-
9253-0

Gustavsson, J., Papenberg, G., Falahati, F., Laukka, E. J., & Kalpouzos, G. (2022). Contributions of the
Catechol-O-Methyltransferase Val158Met Polymorphism to Changes in Brain Iron Across
Adulthood and Their Relationships to Working Memory. *Frontiers in Human Neuroscience*, *16*.
https://doi.org/10.3389/fnhum.2022.838228

Haider, L. (2015). Inflammation, Iron, Energy Failure, and Oxidative Stress in the Pathogenesis of
Multiple Sclerosis. *Oxidative Medicine and Cellular Longevity*, *2015*(3), 1–10.
https://doi.org/10.1155/2015/725370

Hald, A., & Lotharius, J. (2005). Oxidative stress and inflammation in Parkinson's disease: Is there a
causal link? *Experimental Neurology*, *193*(2), 279–290.
https://doi.org/10.1016/j.expneurol.2005.01.013

Hallgren, B., & Sourander, P. (1958). The Effect of Age on the Non-Haemin Iron in the Human Brain.
Journal of Neurochemistry, *3*(1), 41–51. https://doi.org/10.1111/j.1471-4159.1958.tb12607.x

Hare, D. J., & Double, K. L. (2016). Iron and dopamine: A toxic couple. *Brain*, *139*(4), 1026–1035.
https://doi.org/10.1093/brain/aww022

He, Y., Thong, P. S., Lee, T., Leong, S. ., Mao, B. Y., Dong, F., & Watt, F. (2003). Dopaminergic cell
death precedes iron elevation in MPTP-injected monkeys. *Free Radical Biology and Medicine*,
35(5), 540–547. https://doi.org/doi.org/10.1016/j.neurobiolaging.2005.12.010

Innis, R. B., Cunningham, V. J., Delforge, J., Fujita, M., Gjedde, A., Gunn, R. N., Holden, J., Houle, S.,
Huang, S.-C., Ichise, M., Iida, H., Ito, H., Kimura, Y., Koeppe, R. A., Knudsen, G. M., Knuuti, J.,
Lammertsma, A. A., Laruelle, M., Logan, J., … Carson, R. E. (2007). Consensus Nomenclature for
in vivo Imaging of Reversibly Binding Radioligands. *Journal of Cerebral Blood Flow &*
Metabolism, *27*(9), 1533–1539. https://doi.org/10.1038/sj.jcbfm.9600493

Johansson, J., Nordin, K., Pedersen, R., Karalija, N., Papenberg, G., Andersson, M., Korkki, S. M.,
Riklund, K., Guitart-Masip, M., Rieckmann, A., Bäckman, L., Nyberg, L., & Salami, A. (2022). Bi-
phasic patterns of age-related differences in dopamine D1 receptors across the adult lifespan.
BioRxiv, 1–35. https://doi.org/10.1101/2022.05.24.493225

Kalpouzos, G. (2018). Brain iron accumulation, and motor and cognitive decline in normal aging.
Revue de Neuropsychologie, *10*(3), 205. https://doi.org/10.3917/rne.103.0205

Kalpouzos, G., Garzón, B., Sitnikov, R., Heiland, C., Salami, A., Persson, J., & Bäckman, L. (2017).
Higher Striatal Iron Concentration is Linked to Frontostriatal Underactivation and Poorer
Memory in Normal Aging. *Cerebral Cortex*, *27*(6), 3427–3436.
https://doi.org/10.1093/cercor/bhx045

Karrer, T. M., Josef, A. K., Mata, R., Morris, E. D., & Samanez-Larkin, G. R. (2017). Reduced dopamine
receptors and transporters but not synthesis capacity in normal aging adults: a meta-analysis.
Neurobiology of Aging, *57*, 36–46. https://doi.org/10.1016/j.neurobiolaging.2017.05.006

Kaur, D., Yantiri, F., Rajagopalan, S., Kumar, J., Mo, J. Q., Boonplueang, R., Viswanath, V., Jacobs, R.,
Yang, L., Beal, M. F., DiMonte, D., Volitaskis, I., Ellerby, L., Cherny, R. A., Bush, A. I., & Andersen,
J. K. (2003). Genetic or Pharmacological Iron Chelation Prevents MPTP-Induced Neurotoxicity In
Vivo. *Neuron*, *37*(6), 899–909. https://doi.org/10.1016/S0896-6273(03)00126-0

Lammertsma, A. A., & Hume, S. P. (1996). Simplified Reference Tissue Model for PET Receptor
Studies. *NeuroImage*, *4*(3), 153–158. https://doi.org/10.1006/nimg.1996.0066

Landau, S. M., Lal, R., O'Neil, J. P., Baker, S., & Jagust, W. J. (2009). Striatal dopamine and working
memory. *Cerebral Cortex*, *19*(2), 445–454. https://doi.org/10.1093/cercor/bhn095

Langkammer, C., Schweser, F., Krebs, N., Deistung, A., Goessler, W., Scheurer, E., Sommer, K.,
Reishofer, G., Yen, K., Fazekas, F., Ropele, S., & Reichenbach, J. R. (2012). Quantitative
susceptibility mapping (QSM) as a means to measure brain iron? A post mortem validation
study. *NeuroImage*, *62*(3), 1593–1599. https://doi.org/10.1016/j.neuroimage.2012.05.049

Larsen, B., Olafsson, V., Calabro, F., Laymon, C., Tervo-Clemmens, B., Campbell, E., Minhas, D.,
Montez, D., Price, J., & Luna, B. (2020). Maturation of the human striatal dopamine system
revealed by PET and quantitative MRI. *Nature Communications*, *11*(1), 1–10.
https://doi.org/10.1038/s41467-020-14693-3

Leibovitch, F. S., Black, S. E., Caldwell, C. B., McIntosh, A. R., Ehrlich, L. E., & Szalai, J. P. (1999). Brain
SPECT imaging and left hemispatial neglect covaried using partial least squares: The sunnybrook
stroke study. *Human Brain Mapping*, *7*(4), 244–253. https://doi.org/10.1002/(SICI)1097-
0193(1999)7:4<244::AID-HBM3>3.0.CO;2-K

Li, W., Wu, B., Batrachenko, A., Bancroft-Wu, V., Morey, R. A., Shashi, V., Langkammer, C., De Bellis,
M. D., Ropele, S., Song, A. W., & Liu, C. (2014). Differential developmental trajectories of
magnetic susceptibility in human brain gray and white matter over the lifespan. *Human Brain
Mapping*, *35*(6), 2698–2713. https://doi.org/10.1002/hbm.22360

Lillig, C. H., Berndt, C., & Holmgren, A. (2008). Glutaredoxin systems. *Biochimica et Biophysica Acta -
General Subjects*, *1780*(11), 1304–1317. https://doi.org/10.1016/j.bbagen.2008.06.003

Liu, T., Liu, J., De Rochefort, L., Spincemaille, P., Khalidov, I., Ledoux, J. R., & Wang, Y. (2011).
Morphology enabled dipole inversion (MEDI) from a single-angle acquisition: Comparison with
COSMOS in human brain imaging. *Magnetic Resonance in Medicine*, *66*(3), 777–783.
https://doi.org/10.1002/mrm.22816

Liu, T., Wisnieff, C., Lou, M., Chen, W., Spincemaille, P., & Wang, Y. (2013). Nonlinear formulation of
the magnetic field to source relationship for robust quantitative susceptibility mapping.
Magnetic Resonance in Medicine, *69*(2), 467–476. https://doi.org/10.1002/mrm.24272

Liu, Z., Zhai, X. R., Du, Z. S., Xu, F. F., Huang, Y., Wang, X. Q., Qiu, Y. H., & Peng, Y. P. (2021). Dopamine
receptor D2 on CD4+ T cells is protective against neuroinflammation and neurodegeneration in
a mouse model of Parkinson's disease. *Brain, Behavior, and Immunity*, *98*(April), 110–121.
https://doi.org/10.1016/j.bbi.2021.08.220

Mattay, V. S., Fera, F., Tessitore, A., Hariri, A. R., Berman, K. F., Das, S., Meyer-Lindenberg, A.,
Goldberg, T. E., Callicott, J. H., & Weinberger, D. R. (2006). Neurophysiological correlates of age-
related changes in working memory capacity. *Neuroscience Letters*, *392*(1–2), 32–37.
https://doi.org/10.1016/j.neulet.2005.09.025

Mazhar, M., Din, A. U., Ali, H., Yang, G., Ren, W., Wang, L., Fan, X., & Yang, S. (2021). Implication of
ferroptosis in aging. *Cell Death Discovery*, *7*(1), 1–9. https://doi.org/10.1038/s41420-021-
00553-6

McCann, H., Cartwright, H., & Halliday, G. M. (2016). Neuropathology of α-synuclein propagation and
braak hypothesis. *Movement Disorders*, *31*(2), 152–160. https://doi.org/10.1002/mds.26421

McIntosh, A. R., Bookstein, F. L., Haxby, J. V., & Grady, C. L. (1996). Spatial pattern analysis of
functional brain images using partial least squares. *NeuroImage*, *3*(3 I), 143–157.
https://doi.org/10.1006/nimg.1996.0016

McIntosh, A. R., Chau, W. K., & Protzner, A. B. (2004). Spatiotemporal analysis of event-related fMRI
data using partial least squares. *NeuroImage*, *23*(2), 764–775.
https://doi.org/10.1016/j.neuroimage.2004.05.018

McIntosh, A. R., & Lobaugh, N. J. (2004). Partial least squares analysis of neuroimaging data:
Applications and advances. *NeuroImage*, *23*(SUPPL. 1), 250–263.
https://doi.org/10.1016/j.neuroimage.2004.07.020

Mills, E., Dong, X. P., Wang, F., & Xu, H. (2010). Mechanisms of brain iron transport: Insight into
neurodegeneration and CNS disorders. *Future Medicinal Chemistry*, *2*(1), 51–64.
https://doi.org/10.4155/fmc.09.140

Nordin, K., Gorbach, T., Pedersen, R., Lundmark, V. P., Johansson, J., Andersson, M., Mcnulty, C.,
Riklund, K., Wåhlin, A., Papenberg, G., Kalpouzos, G., Bäckman, L., & Salami, A. (2022).
DyNAMiC: A prospective longitudinal study of dopamine and brain connectomes: A new
window into cognitive aging. *Journal of Neuroscience Research*, *January*, 1–25.
https://doi.org/10.1002/jnr.25039

Nyberg, L., Andersson, M., Forsgren, L., Jakobsson-Mo, S., Larsson, A., Marklund, P., Nilsson, L. G.,
Riklund, K., & Bäckman, L. (2009). Striatal dopamine D2 binding is related to frontal BOLD
response during updating of long-term memory representations. *NeuroImage*, *46*(4), 1194–
1199. https://doi.org/10.1016/j.neuroimage.2009.03.035

Nyberg, L., Andersson, M., Kauppi, K., Lundquist, A., Persson, J., Pudas, S., & Nilsson, L.-G. (2014).
Age-related and Genetic Modulation of Frontal Cortex Efficiency. *Journal of Cognitive
Neuroscience*, *26*(4), 746–754. https://doi.org/10.1162/jocn_a_00521

Nyberg, L., Dahlin, E., Stigsdotter Neely, A., & BÄckman, L. (2009). Neural correlates of variable
working memory load across adult age and skill: Dissociative patterns within the fronto-parietal
network: Cognition and Neurosciences. *Scandinavian Journal of Psychology*, *50*(1), 41–46.
https://doi.org/10.1111/j.1467-9450.2008.00678.x

Ortega, R., Cloetens, P., Devès, G., Carmona, A., & Bohic, S. (2007). Iron storage within dopamine
neurovesicles revealed by chemical nano-imaging. *PLoS ONE*, *2*(9).
https://doi.org/10.1371/journal.pone.0000925

Poetini, M. R., Araujo, S. M., Trindade de Paula, M., Bortolotto, V. C., Meichtry, L. B., Polet de
Almeida, F., Jesse, C. R., Kunz, S. N., & Prigol, M. (2018). Hesperidin attenuates iron-induced
oxidative damage and dopamine depletion in Drosophila melanogaster model of Parkinson's
disease. *Chemico-Biological Interactions*, *279*(September 2017), 177–186.
https://doi.org/10.1016/j.cbi.2017.11.018

Reuter-Lorenz, P. A., & Cappell, K. A. (2008). Neurocognitive aging and the compensation hypothesis.
Current Directions in Psychological Science, *17*(3), 177–182. https://doi.org/10.1111/j.1467-
8721.2008.00570.x

Rieckmann, A., Karlsson, S., Karlsson, P., Brehmer, Y., Fischer, H., Farde, L., Nyberg, L., & Bäckman, L.
(2011). Dopamine D1 receptor associations within and between dopaminergic pathways in
younger and elderly adults: Links to cognitive performance. *Cerebral Cortex*, *21*(9), 2023–2032.

https://doi.org/10.1093/cercor/bhq266

Rodrigue, K. M., Daugherty, A. M., Foster, C. M., & Kennedy, K. M. (2020). Striatal iron content is
linked to reduced fronto-striatal brain function under working memory load. *NeuroImage*,
210(October 2019), 116544. https://doi.org/10.1016/j.neuroimage.2020.116544

Salami, A., Avelar-Pereira, B., Garzón, B., Sitnikov, R., & Kalpouzos, G. (2018). Functional coherence of
striatal resting-state networks is modulated by striatal iron content. *NeuroImage*, *183*(April),
495–503. https://doi.org/10.1016/j.neuroimage.2018.08.036

Salami, A., Eriksson, J., Kompus, K., Habib, R., Kauppi, K., & Nyberg, L. (2010). Characterizing the
neural correlates of modality-specific and modality-independent accessibility and availability
signals in memory using partial-least squares. *NeuroImage*, *52*(2), 686–698.
https://doi.org/10.1016/j.neuroimage.2010.04.195

Salami, A., Eriksson, J., & Nyberg, L. (2012). Opposing effects of aging on large-scale Brain systems for
memory encoding and cognitive control. *Journal of Neuroscience*, *32*(31), 10749–10757.
https://doi.org/10.1523/JNEUROSCI.0278-12.2012

Salami, A., Garrett, D. D., Wåhlin, A., Rieckmann, A., Papenberg, G., Karalija, N., Jonasson, L.,
Andersson, M., Axelsson, J., Johansson, J., Riklund, K., Lövdén, M., Lindenberger, U., Bäckman,
L., & Nyberg, L. (2019). Dopamine D 2/3 binding potential modulates neural signatures of
working memory in a load-dependent fashion. *Journal of Neuroscience*, *39*(3), 537–547.
https://doi.org/10.1523/JNEUROSCI.1493-18.2018

Salami, A., Papenberg, G., Sitnikov, R., Laukka, E. J., Persson, J., & Kalpouzos, G. (2021). Elevated
Neuroinflammation Contributes to the Deleterious Impact of Iron Overload on Brain Function in
Aging. *NeuroImage*, *230*(August 2020), 117792.
https://doi.org/10.1016/j.neuroimage.2021.117792

Salami, A., Rieckmann, A., Fischer, H., & Bäckman, L. (2014). A multivariate analysis of age-related
differences in functional networks supporting conflict resolution. *NeuroImage*, *86*, 150–163.
https://doi.org/10.1016/j.neuroimage.2013.08.002

Salthouse, T. (2012). Consequences of age-related cognitive declines. Annual review of psychology.
Annu Rev Psychol., *63*, 201-226. https://doi.org/10.1146/annurev-psych-120710-
100328.Consequences

Schneider-Garces, N. J., Gordon, B. A., Brumback-Peltz, C. R., Shin, E., Lee, Y., Sutton, B. P., Maclin, E.
L., Gratton, G., & Fabiani, M. (2010). Span, CRUNCH, and beyond: Working memory capacity
and the aging brain. *Journal of Cognitive Neuroscience*, *22*(4), 655–669.
https://doi.org/10.1162/jocn.2009.21230

Shao, W., Zhang, S., Tang, M., Zhang, X., Zhou, Z., Yin, Y., Zhou, Q., Huang, Y., Liu, Y., Wawrousek, E.,
Chen, T., Li, S., Xu, M., Zhou, J., Hu, G., & Zhou, J. (2013). Suppression of neuroinflammation by
astrocytic dopamine D2 receptors via αB-crystallin. *Nature*, *494*, 90–94.
https://doi.org/https://doi.org/10.1038/nature11748

Shohamy, D., & Adcock, R. A. (2010). Dopamine and adaptive memory. *Trends in Cognitive Sciences*,
14(10), 464–472. https://doi.org/10.1016/j.tics.2010.08.002

Spence, H., McNeil, C. J., & Waiter, G. D. (2020). The impact of brain iron accumulation on cognition:
A systematic review. *PLoS ONE*, *15*(10 October). https://doi.org/10.1371/journal.pone.0240697

Steiger, T. K., Weiskopf, N., & Bunzeck, N. (2016). Iron level and myelin content in the ventral
striatum predict memory performance in the aging brain. *Journal of Neuroscience*, *36*(12),
3552–3558. https://doi.org/10.1523/JNEUROSCI.3617-15.2016

Straub, S., Schneider, T. M., Emmerich, J., Freitag, M. T., Ziener, C. H., Schlemmer, H. P., Ladd, M. E.,
& Laun, F. B. (2017). Suitable reference tissues for quantitative susceptibility mapping of the
brain. *Magnetic Resonance in Medicine*, *78*(1), 204–214. https://doi.org/10.1002/mrm.26369

Tabachnick, B., & Fidell, L. (2013). Using Multivariate Statistics. In *Singular Spectrum Analysis* (6th
ed.). Pearson. https://doi.org/10.1007/978-1-4757-2514-8_3

Unger, E. L., Bianco, L. E., Jones, B. C., Allen, R. P., & Earley, C. J. (2014). Low brain iron effects and
reversibility on striatal dopamine dynamics. *Experimental Neurology*, *261*, 462–468.
https://doi.org/10.1016/j.expneurol.2014.06.023

Ward, R. J., Zucca, F. A., Duyn, J. H., Crichton, R. R., & Zecca, L. (2014). The role of iron in brain ageing
and neurodegenerative disorders. *The Lancet Neurology*, *13*(10), 1045–1060.
https://doi.org/10.1016/S1474-4422(14)70117-6

Xu, W., & Cumming, I. (1999). A region-growing algorithm for InSAR phase unwrapping. *IEEE*
Transactions on Geoscience and Remote Sensing, *37*(1), 124–134.
https://doi.org/10.1109/36.739143

Yan, Y., Jiang, W., Liu, L., Wang, X., Ding, C., Tian, Z., & Zhou, R. (2015). Dopamine Controls Systemic
Inflammation through Inhibition of NLRP3 Inflammasome. *Cell*, *160*(1–2), 62–73.
https://doi.org/10.1016/j.cell.2014.11.047

Youdim, M. B. H., Ben-Shachar, D., & Riederer, P. (1993). The possible role of iron in the
etiopathology of parkinson's disease. *Movement Disorders*, *8*(1), 1–12.
https://doi.org/10.1002/mds.870080102

Zecca, L., Youdim, M., Riederer, P., Connor, J. R., & Crichton, R. R. (2004). Iron, brain ageing and
neurodegenerative disorders. *Nature Reviews Neuroscience*, *5*(11), 863–873.
https://doi.org/10.1038/nrn1537

Zhu, J., Hu, Z., Han, X., Wang, D., Jiang, Q., Ding, J., Xiao, M., Wang, C., Lu, M., & Hu, G. (2018).
Dopamine D2 receptor restricts astrocytic NLRP3 inflammasome activation via enhancing the
interaction of β -arrestin2 and NLRP3. *Cell Death & Differentiation*, *25*, 2037–2049.
https://doi.org/10.1038/s41418-018-0127-2

Zimmerman, M. E., Brickman, A. M., Paul, R. H., Grieve, S. M., Tate, D. F., Gunstad, J., Cohen, R. A.,
Aloia, M. S., Williams, L. M., Clark, C. R., Whitford, T. J., & Gordon, E. (2006). The relationship
between frontal gray matter volume and cognition varies across the healthy adult lifespan.
American Journal of Geriatric Psychiatry, *14*(10), 823–833.
https://doi.org/10.1097/01.JGP.0000238502.40963.ac

Zucca, F. A., Segura-Aguilar, J., Ferrari, E., Muñoz, P., Paris, I., Sulzer, D., Sarna, T., Casella, L., & Zecca,
L. (2017). Interactions of iron, dopamine and neuromelanin pathways in brain aging and
Parkinson's disease. *Progress in Neurobiology*, *155*(3), 96–119.
https://doi.org/10.1016/j.pneurobio.2015.09.012