1
2

DRomics, a workflow to model and make sense of dose-response (multi-)omics data in (eco)toxicology

Marie Laure Delignette-Muller*[1], Aurélie Siberchicot[1], Floriane Larras[1], Elise Billoir[2]

[1] Université de Lyon, Université Lyon 1, CNRS, VetAgro Sup, UMR 5558, Laboratoire de Biométrie et Biologie Evolutive, 69622 Villeurbanne, France
[2] Université de Lorraine, CNRS, LIEC, F-57000 Metz, France

*Corresponding author
Correspondence: marielaure.delignettemuller@vetagro-sup.fr or dromics@univ-lyon1.fr

16
17

ABSTRACT
Omics technologies has opened new possibilities to assess environmental risks and to understand the mode(s) of action of pollutants. Coupled to dose-response experimental designs, they allow a non-targeted assessment of organism responses at the molecular level along an exposure gradient. However, describing the dose-response relationships on such high-throughput data is no easy task. In a first part, we review the software available for this purpose, and their main features. We set out arguments on some statistical and modeling choices we have made while developing DRomics (an R package released in its first version in 2019) and its positioning compared to others tools. The DRomics main analysis workflow is made available through a web interface, namely a shiny app named DRomics-shiny. Next, we present the new functionalities recently implemented. DRomics has been augmented especially to be able to handle varied omics data considering the nature of the measured signal (e.g. counts of reads in RNAseq) and the way data were collected (e.g. experimental versus *in situ* data). Another important upgrade is the development of tools to ease the biological interpretation of results. Various functions are proposed to visualize, summarize and compare the responses, for different biological groups (defined from biological annotation), optionally at different experimental levels (e.g. measurements at several omics level or in different experimental conditions). A new shiny app named DRomicsInterpreter-shiny is dedicated to the biological interpretation of results.

Keywords: Dose-response modeling, BenchMark Dose (BMD), Adverse Outcome Pathway (AOP), Mode of Action (MoA), environmental risk assessment, toxicogenomics, transcriptomics, proteomics, metabolomics, multi-omics

40
41

Introduction

Dose-response (DR) modeling belongs to the toolkit of ecotoxicologists. The latter are used to this approach when working on apical endpoints (e.g. reproduction, photosynthesis, …). The derived sensitivity thresholds, e.g. Effective Concentrations (ECx), BenchMark Dose (BMD), are at the basis of regulatory risk assessment.

The recent years have seen the emergence of works using DR omics data (transcriptomic, proteomic, metabolomic, …) in ecotoxicology (Zhang *et al.*, 2018). Some of those works aim at a mechanistic understanding of adverse effects (Adverse Outcome Pathway perspective - AOP). They can identify potential Modes of Action of pollutants (MoAs) at the molecular level, that generally need to be validated in a second step using targeted experiments (Andersen *et al.* 2018). Typical DR designs, with many doses (>6), ensure a good description of the DR relationship and a robust and precise estimation of a sensitivity threshold (such as the benchmark dose – BMD) that is useful to fix regulatory thresholds (Ewald *et al.*, 2022). However, such designs are less commonly used in omics studies, as tools classically used to analyse omics data are dedicated to differential expression analysis between few conditions (limma, Ritchie *et al.*, 2025, DESeq2, Love *et al.*, 2014, EdgeR, Robinson *et al.*, 2010). The analysis of such data typically starts with a pairwise differential analysis to the control followed by an enrichment procedure to identify GO (Gene Ontology) terms or KEGG (Kyoto Encyclopedia of Genes and Genomes) biological pathways of differentially expressed items (contigs, proteins, …) (Dubois *et al.*, 2019, Murat El Houdigui *et al.*, 2019, Meier *et al.*, 2020, Zhan *et al.*, 2021). In a second time, some perform a DR modeling on differentially expressed items to estimate a BMD per item and summarize the sensitivity of each pathway for example by the median of BMDs of corresponding pathways (Meier *et al.*, 2020, Zhan *et al.*, 2021).

The purpose of this paper is to present the new version of the DRomics ("Dose Response for Omics") R package and its two companion interactive web applications, DRomics-shiny and DRomicsInterpreter-shiny. We first review the software available to tackle a DR analysis of high-throughput omic data and explain how DRomics distinguishes from other tools. Then, we present the various functionalities we added to DRomics from its first version published in 2018 and especially explain the way it can be used to make sense of DR (multi-)omics data in environmental risk assessment.

DRomics original features compared to other tools dedicated to dose-response omics data

Based on literature review in toxicology and ecotoxicology, we identified five tools available for DR analysis of high-throughput omics data (Table 1). Their workflows consist of successive steps: first, the selection of items (e.g. contigs) significantly regulated along the gradient of exposure. Then, for the selected items, DR relationships are modeled and BMD are derived from these DR models. The BMD-zSD, the most often used and recommended version of the BMD, is defined as the dose that leads to a response (BMR – benchmark response) pointing a difference from the response in controls of more than z times (e.g. with z=1, EFSA, 2017) the residual standard deviation of the DR model.

The first piece of software developed to analyze high dimensional DR data, in particular gene expression data, was BMDExpress, released in its first version in 2007 (Yang *et al.*, 2007). It was since updated and augmented in BMDExpress-2 (Philips *et al.*, 2019). FastBMD (Ewald *et al.*, 2021) implements the same methods, as framed by the US National Toxicology Program (NTP, 2018), with the sake of being faster and friendlier to end-users than BMDExpress, and through a web-based interface. Again in the context of toxicogenomics, BBMD (Ji *et al.*, 2022) has the specificity to use model averaging to account for BMD uncertainty related to the underlying DR model, and BMDx (Serra *et al.*, 2020) allows a comparison of BMD values of a transcriptomics experiment at different time points or from different experiments. We developed DRomics in 2018 (Larras *et al.*, 2018) for DR analysis of any omics data (e.g. transcriptomics, metabolomics), including non-sequenced organisms or communities (meta-omics), biological models commonly used in the field of ecotoxicology. The main characteristics and functionalities of those five tools are summarized in Table 1.

Table 1 - Main characteristics of the tools available for DR analysis of high-throughput omics data

	Software	DRomics	BMDExpress 2	FastBMD	BBMD	BMDx
General	References	Larras *et al*., 2018	Yang *et al*., 2007 Philips *et al*., 2019	Ewald *et al*., 2020	Ji *et al*., 2022	Serra *et al*., 2020
	Main application area	Ecotoxicology	Toxicology	Toxicology	Toxicology	Toxicology
	Platform	R package, Web app. (R Shiny) free without registration	free standalone application that must be locally installed	Web app. free without registration	Web app. with registration (free or premium accounts)	Web app. (R Shiny) to be launched locally (no server - source on GitHub)
	Open source	yes (CRAN, GitHub)	yes (GitHub)	no	no	yes (GitHub)
	Programming languages	R	Java, C, Fortran	R, JavaServer Faces (JSF)	Python, Javascript	R
	Date of the first version	2019	2007	2021	2022	2019
	Date of last revision (as on 02/02/2023)	2023	2020 BMDExpress 3 in prep. https://github.com/auerbachs/BMDExpress-3	2022	2022	2022
	Dependencies to other statistical R packages	limma, DESeq2 (Bioconductor packages)	not mentionned	limma, vsn, edgeR, DESeq2, genefilter and preprocessCore (Bioconductor packages)	not mentionned	drc, bmd, alr3, jtools (alr3 and bmd no more available on CRAN)
Import/ Visualisation of data	Types of data for which the tool is dedicated	RNAseq, microarray, continuous omics data, continuous anchoring data	Transcriptomics data (does not differentiate microarray and RNAseq data) or other continuous data	RNAseq, microarray	RNAseq, microarray, continuous anchoring data, binary anchoring data	Transcriptomics data (does not differentiate microarray and RNAseq data)
	Preprocessing of transcriptomic data (microarray/RNAseq) supported by the software	yes	no	yes	no	no
	Proposed plots to check imported data	PCA + Boxplot	PCA + Density	PCA + Boxplot + Density	PCA + Density	none
Selection of significant responses	Selection of significant responses to the dose gradient	yes	yes	yes	yes	yes
	Filter on the fold change for selection	no	optional (by default yes, > 2)	optional (by default yes > 1)	yes mandatory (> 1.5 at least)	no
	Control of the FDR for this selection	yes	optional (by default no)	optional (by default no)	yes	optional
	Proposed methods for selection	ANOVA, linear or quadratic trend test (for selection of monotonic and biphasic responses)	ANOVA, William's trend test, Oriogen	ANOVA	ANOVA or adaptation of William's trend test and Oriogen for selection of monotonic responses	ANOVA or trend test (monotonic)

Dose-response modeling	Shapes of proposed models	monotonic and biphasic	monotonic, biphasic, multiphasic (3rd and 4th order polynomials)	monotonic, biphasic, multiphasic (3rd and 4th order polynomials)	monotonic (Linear, Hill, Pow, Exp2, Exp3, Exp4, Exp5)	monotonic, biphasic, multiphasic (3rd order polynomials)
	User selection of models	no	yes	yes	yes	yes
	Proposed criterion for best fit model choice	AICc (default), AIC, BIC	AIC and comparison of nested models for polynomial models	AIC	Model averaging	AIC
	Characterization of the response	increasing (up-regulated), decreasing (down-regulated), U-shape, bell-shape	none	none	increasing, decreasing	none
	Toxicity threshold provided	BMD (-zSD and -xfold)	BMD-zSD	BMD-zSD	BMD (-zSD or -xfold)	BMD-zSD
	Method for computing confidence intervals on the toxicity threshold	bootstrap	based on the likelihood but not precisely described	profile likelihood method	model averaging	not found
Biological interpretation	Includes the annotation step	no	yes (6 species)	yes (14 species)	yes (6 species)	yes (3 species)
	Available organisms for annotation	none	*H. sapiens, M. musculus, C. lupus familiaris, R. norvegicus, D. melanogaster, D. rerio*	*H. sapiens, M. musculus, R. norvegicus, C. elegans, D. melanogaster, D. rerio, S. cerevisiae, A. thalian, B. taurus, G. gallus, C. japonica, X laevis, P. promelas, O. mykiss,* + annotation-free pipeline, Seq2Fun Ortholog	*H. sapiens, M. musculus, R. norvegicus, , D. melanogaster, D. rerio, B. taurus*	*H. sapiens, M. musculus, R. norvegicus*
	Annotation available databases	none	GO, reactome	KEGG, GO (BP, MF, CC), reactome	GeneID, GO, reactome, KEGG	KEGG, reactome, GO
	Functions to compare the responses at different experimental levels (multi-omics, diff. conditions, …)	yes	not found	not found	not found	yes (multiple time points, multiple experiments)

DRomics-shiny
Main workflow

DRomicsInterpreter-shiny
to help biological
interpretation

95   **Figure 1** - Diagram of the DRomics functionalities and of the perimeter of associated tools. Bold font indicates new functions added to the DRomics R package since its first version.

New features in the DRomics DR modeling workflow

Figure 1 maps the DRomics workflow and the functionalities offered to explore the results. First
developed for microarray data, DRomics is now able to handle RNAseq data, metabolomic or other
continuous omic data (e.g. proteomics data) or even continuous non omic data (e.g. growth data) that
could be used for phenotypical anchoring (respectively imported using RNAseqdata(),
continuousomicdata() and continuousanchoringdata() functions). The same modeling workflow, choosing
the best fit models among our complete family of models (as described previously), was declined for each
type of data (e.g. intensities, counts) to ensure the comparability of results (e.g. transcriptomics vs.
metabolomics, transcriptomics vs. anchoring). Writing in the R language ensures the interoperability with
functions of the Bioconductor packages. Thus, functions of the DESeq2 and limma packages are internally
called within the package to normalize and/or transform omic data and to implement the trend tests for
selecting significantly responsive items. The call to additional R functions can be added for example to
correct omics data for a potential batch effect (see an example using ComBat-seq in the package vignette:
https://cran.r-project.org/web/packages/DRomics/vignettes/DRomics_vignette.html#insitudata).
Various functions were also added to the package (highlighted in bold type in Figure 1) to respond to
user requests. Among them we can cite PCAdataplot() for a visualization of data after the importation step
and detection of potential outlier samples or batch effect, targetplot() to visualize the response of targeted
items whatever they are selected or not in the DRomics workflow, bmdboot() and bmdplot() to compute
and visualize confidence intervals on BMD values using bootstrap. Moreover, we performed modifications
in the modeling workflow to ensure a better robustness of results on *in situ* data with no replicate due to
the uncontrolled dose, or on data with a low number of doses (4 or 5). For example, we changed the default
information criterion used for best model selection from the AIC to the AICc, as recommended by Burnham
and Anderson (2004), and limited the set of models to those with a maximum of 4 parameters for designs
with less than 5 doses.

New functions and the new Shiny application to help biological interpretation

In toxicology, while working on sequenced and well-annotated organisms, items (e.g. genes) can be
functionally annotated prior to DR analysis. BMDExpress integrates this annotation step for 6 model
species on the basis of GO or reactome databases, and FastBMD for 14 model species on the basis of
GeneID, GO, Reactome or KEGG databases (see Table 1). Such an annotation of all the items whatever they
respond or not to the dose gradient exposition, enables the classical enrichment analysis (Wu *et al.*, 2021).
This analysis consists in highlighting gene sets/pathways which are the most overrepresented among the
responding ones.
In ecotoxicology, one often works on non-model organisms or even on samples from environmental
communities (freshwater biofilm for example – Creusot *et al.*, 2021). This implies the need for the user to
retrieve and manage its own annotation, which is a challenging task, especially for RNAseq experiments
for which the number of measured contigs may be huge (several millions of contigs). We thus considered
that this annotation step could be done after the selection/modeling workflow, to reduce the number of
items to annotate, and so the difficulty of this task. Due to the great diversity of annotation pipelines that
can be developed for such non-model organisms, we did not include an annotation step in DRomics.
However, and to support the interpretation of the workflow results in view of a biological annotation
provided by the user, we recently developed new functions and a second shiny application (named
DRomicsInterpreter-shiny) (Figure 1).
After augmenting the DRomics output with information about the functional role of items, the
graphical representations offered by DRomics give a new insight into the results. Further, provided a
common annotation system is used at different biological scales under study, DRomics can be used to
compare the response at various levels. Figures 2 and 3 give some illustrations on an example with two
molecular levels from Larras *et al.* (2020): transcriptomics and metabolomics responses of *Scenedesmus*
vacuolatus to triclosan exposure. Various functions are proposed to visualize/summarize/compare the

responses, for the different biological groups (defined from biological annotation), optionally at the
different experimental levels.

- the trends (increasing, decreasing, U-shape or bell-shape) of the DR curves (using the trendplot() function, see a one-level example on Figure 1),
- the group/pathway-level sensitivity calculated as a quantile of BMD values of the group (using the sensitivityplot() function, see an one-level example on Figure 1 and a multilevel-level example on Figure 2),
- for selected groups/pathways, all the BMD values with their confidence intervals (using the bmdplot () function, see a multilevel-level example on Figure 1, right part)
- for selected groups/pathways, all the BMD values with the corresponding DR curve signal coded as a color gradient (using the bmdplotwithgradient() function, see a multi-level example on Figure 3, left part).
- for selected groups/pathways, all the DR curves represented as curves (using the curvesplot() function, see a multi-level example on Figure 3, right part),

Those functions can also be used to compare the response at one molecular level but measured under
different experimental conditions (different time points, different pre-exposure scenarios, in vitro/in vivo,
etc.). The selectgroups() function was also developed to help the user to focus its interpretation on the
most represented and/or the most sensitive biological groups (see an example of use in the package
vignette: https://cran.r-
project.org/web/packages/DRomics/vignettes/DRomics_vignette.html#selectgroups).

Figure 2 - Illustration of the use of the function sensitivityplot() here to summarize the sensitivity of the responding
KEGG pathways at two molecular levels using transcriptomic and metabolomic data published in Larras *et al*. (2020) on
Scenedesmus vacuolatus exposed to triclosan

Figure 3 - Summary of the DR curves of all contigs and metabolites corresponding to two specific KEGG pathways: «membrane transport» and «lipid metabolism» (from data published in Larras *et al.* (2020) on *Scenedesmus vacuolatus* exposed to triclosan) on the left using the bmdplotwithgradient() function and on the right using the curvesplot() fuction. The signal was shifted by putting the initial theoretical value at 0 and scaled so as to make it vary in the same amplitude to focus on the shape of the responses and not on the amplitude of the signal (it is an option of both functions).

Perspectives

In the report of the National Toxicology Program (NTP, 2018) model averaging is mentioned as an interesting feature to make the BMD estimate less dependent of the choice of the best DR model. BBMD (Ji *et al.*, 2022) and the next version of BMDExpress in preparation (BMDExpress-3 – https://github.com/auerbachs/BMDExpress-3/) include a Bayesian model averaging procedure. However, we do not plan to implement model averaging in DRomics because the BMD estimation is not our only purpose. We also want to characterize each response by its trend, which is itself dependent of the model choice and not averageable. Instead, we plan to add an alternative to our current bootstrap procedure, enabling the fit of a different model at each bootstrap iteration, to be able to pass the uncertainty due to the model choice both on the BMD uncertainty and on the trend uncertainty.

Concerning the modeling workflow, so far, we added specific functionalities to analyse continuous anchoring data, essentially to prevent comparisons of BMD values at different biological scales but with BMD obtained from different analysis workflows, which could induce a bias. As anchoring data may be non-continuous, such as dichotomous survival data, or reproduction data reported as number of offspring per individual-day (Delignette-Muller *et al.*, 2014), we plan new developments in DRomics to be able to analyse those data properly taking into account their nature.

Concerning our recent development of functions to help the biological interpretation of DRomics results, we plan to enlarge the range of the methods proposed, by imaging new plots and summaries to explore and characterize the responses and their diversity within a biological group/pathway.

Conclusion

The development of DRomics has been driven by the demands of ecotoxicologists to help make full sense of their dose-response (multi-)omics studies. Hence, DRomics was augmented to provide a common workflow to handle (meta-)transcriptomics, proteomics, metabolomics and/or anchoring data. This creates the foundations for a proper comparison of responses at different omics levels (and anchoring endpoints) and a mechanistic understanding in an AOP perspective. Along with functional annotations, DRomics outputs (response trends, BMD, *etc.*) can now be processed using a series of graphical functions thought to help their biological interpretation at the metabolic pathway level. The comparison is made easy (i) of different measurements, for instance transcripto- and metabolomics, (ii) of different biological materials, for instance communities with/without pre-exposure history, or (iii) of experimental settings, for instance

204 successive timepoints or different temperatures. Moreover, two special cases have been addressed:
205 experiments with a batch effect and *in situ* sampling with no replicate. DRomics future direction and
206 evolutions depend on upcoming challenges and needs brought by (eco)toxicologists.

Acknowledgements

208 The authors thank colleagues and students whose requests and/or discussions contributed to the
209 recent evolution of DRomics, especially Olivier Armant, Ellis Franklin, Sandrine Frelon, Sophie Prud'homme,
210 Mechthild Schmitt-Jansen and Philippe Veber.

Data, scripts, code, and supplementary information availability

Scripts and code are available online: DRomics is written in R, freely available from CRAN and
encompasses two Shiny applications (https://cran.r-project.org/web/packages/DRomics/). Those Shiny
applications are also freely available online. The homepage https://lbbe.univ-lyon1.fr/fr/dromics gathers
links to all resources related to DRomics, e.g. the two shiny applications, a cheat sheet and a complete
vignette to start with DRomics. The github https://aursiber.github.io/DRomics/ hosts DRomics stable as
well as development versions.
Data used in this paper are embedded in the DRomics package.

Conflict of interest disclosure

220 The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
221 relation to the content of the article.

Funding

This work has been financially supported by the Graduate School H2O'Lyon (ANR-17-EURE-0018) of
Université de Lyon (UdL), within the program "Investissements d'Avenir" operated by the French National
Research Agency (ANR)", by the "Ecosphère continentale et côtière" (EC2CO) interdisciplinary program
from the Centre National de la Recherche Scientifique (CNRS, France), as parts of the DROMADERE and the
ECORA projects, and by the "Agence Nationale de la Recherche" as a part of the Chroco project (ANR-21-
CE34-0003).

References

Andersen, M. E., Pendse, S. N., Black, M. B., & McMullen, P. D. (2018). Application of transcriptomic data,
visualization tools and bioinformatics resources for informing mode of action. Current Opinion in
Toxicology, 9, 21-27. https://doi.org/10.1016/j.cotox.2018.05.003
Burnham, KP, Anderson DR (2004). Multimodel inference: understanding AIC and BIC in model selection.
Sociological methods & research, 33(2), 261-304. https://doi.org/10.1177/0049124104268644
Creusot, N., Chaumet, B., Eon, M., Mazzella, N., Moreira, A., & Morin, S. (2021). Metabolomics insight into
the influence of environmental factors in responses of freshwater biofilms to the model herbicide
diuron. Environmental Science and Pollution Research, 29(20), 29332-29347.
https://doi.org/10.1007/s11356-021-17072-7
Delignette-Muller, M. L., Lopes, C., Veber, P., & Charles, S. (2014). Statistical handling of reproduction data
for exposure-response modeling. Environmental science & technology, 48(13), 7544-7551.
https://doi.org/10.1021/es502009r
Desforges, J. P., Legrand, E., Boulager, E., Liu, P., Xia, J., Butler, H., ... & Crump, D. (2021). Using
Transcriptomics and Metabolomics to Understand Species Differences in Sensitivity to Chlorpyrifos in
Japanese Quail and Double-Crested Cormorant Embryos. Environmental Toxicology and Chemistry,
40(11), 3019-3033. https://doi.org/10.1002/etc.5174

Dubois, C., Pophillat, M., Audebert, S., Fourquet, P., Lecomte, C., Dubourg, N., ... & Frelon, S. (2019). Differential modification of the C. elegans proteome in response to acute and chronic gamma radiation: Link with reproduction decline. Science of the total environment, 676, 767-781. https://doi.org/10.1016/j.scitotenv.2019.04.039

EFSA Scientific Committee; Hardy, A., Benford, D., Halldorsson, T., Jeger, M. J., Knutsen, K. H., More, S., Mortensen, A., Naegeli, H., Noteborn, H., Ockleford, C., Ricci, A., Rychen, G., Silano, V., Solecki, R., Turck, D., Aerts, M., Bodin, L., Davis, A., Edler, L., Gundert-Remy, U., Sand, S., Slob, W., Bottex, B., Abrahantes, J. C., Marques, D. C., Kass, G., Schlatter, J. R. (2017) Update: Use of the Benchmark Dose Approach in Risk Assessment. EFSA J., 15 (1). https://efsa.onlinelibrary.wiley.com/doi/full/10.2903/j.efsa.2017.4658

Ewald, J. D., Soufan, O., Crump, D., Hecker, M., Xia, J., & Basu, N. (2020). EcoToxModules: custom gene sets to organize and analyze toxicogenomics data from ecological species. Environmental Science & Technology, 54(7), 4376-4387. https://doi.org/10.1021/acs.est.9b06607

Ewald, J., Soufan, O., Xia, J., Basu, N. (2021). FastBMD: an online tool for rapid benchmark dose–response analysis of transcriptomics data. Bioinformatics, 37(7), 1035-1036. https://doi.org/10.1093/bioinformatics/btaa700

Ewald, J. D., Basu, N., Crump, D., Boulanger, E., & Head, J. (2022). Characterizing Variability and Uncertainty Associated with Transcriptomic Dose–Response Modeling. Environmental Science & Technology, 56(22), 15960-15968. https://doi.org/10.1021/acs.est.2c04665

Gündel, U., Kalkhof, S., Zitzkat, D., von Bergen, M., Altenburger, R., & Küster, E. (2012). Concentration–response concept in ecotoxicoproteomics: Effects of different phenanthrene concentrations to the zebrafish (Danio rerio) embryo proteome. Ecotoxicology and environmental safety, 76, 11-22. https://doi.org/10.1016/j.ecoenv.2011.10.010

Ji, C., Weissmann, A., Shao, K. (2022). A computational system for Bayesian benchmark dose estimation of genomic data in BBMD. Environment International, 161, 107135. https://doi.org/10.1016/j.envint.2022.107135

Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation network analysis. BMC bioinformatics, 9(1), 1-13. https://doi.org/10.1186/1471-2105-9-559

Larras, F., Billoir, E., ... & Delignette-Muller, M. L. (2018). DRomics: A turnkey tool to support the use of the dose–response framework for omics data in ecological risk assessment. Environmental science & technology, 52(24), 14461-14468. https://doi.org/10.1021/acs.est.8b04752

Larras, F., Billoir, E., Scholz, S., Tarkka, M., Wubet, T., Delignette-Muller, M. L., Schmitt-Jansen, M. (2020). A multi-omics concentration-response framework uncovers novel understanding of triclosan effects in the chlorophyte Scenedesmus vacuolatus. Journal of Hazardous Materials. https://doi.org/10.1016/j.jhazmat.2020.122727

Love, M. I., Huber, W., Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology, 15, 550. https://doi.org/10.1186/s13059-014-0550-8

Martínez, R., Codina, A. E., Barata, C., Tauler, R., Piña, B., & Navarro-Martín, L. (2020). Transcriptomic effects of tributyltin (TBT) in zebrafish eleutheroembryos. A functional benchmark dose analysis. Journal of Hazardous Materials, 122881. https://doi.org/10.1016/j.jhazmat.2020.122881

Meier, M. J., Dodge, A. E., Samarajeewa, A. D., & Beaudette, L. A. (2020). Soil exposed to silver nanoparticles reveals significant changes in community structure and altered microbial transcriptional profiles. Environmental Pollution, 258, 113816. https://doi.org/10.1016/j.envpol.2019.113816

Murat El Houdigui, S., Adam-Guillermin, C., Loro, G., Arcanjo, C., Frelon, S., Floriani, M., ... & Armant, O. (2019). A systems biology approach reveals neuronal and muscle developmental defects after chronic exposure to ionising radiation in zebrafish. Scientific reports, 9(1), 1-15. https://doi.org/10.1038/s41598-019-56590-w

National Toxicology Program. (2018). NTP Research Report on National Toxicology Program Approach to Genomic Dose-Response Modeling: Research Report 5. https://ntp.niehs.nih.gov/ntp/results/pubs/rr/reports/rr05_508.pdf

Peddada, S. D., Lobenhofer, E. K., Li, L., Afshari, C. A., Weinberg, C. R., & Umbach, D. M. (2003). Gene selection and clustering for time-course and dose–response microarray experiments using order-restricted inference. Bioinformatics, 19(7), 834-841. https://doi.org/10.1093/bioinformatics/btg093

Phillips, J. R., Svoboda, D. L., Tandon, A., Patel, S., Sedykh, A., Mav, D., ... & Gift, J. S. (2019). BMDExpress
2: enhanced transcriptomic dose-response analysis workflow. Bioinformatics, 35(10), 1780-1782.
https://doi.org/10.1093/bioinformatics/bty878
R Core Team (2021). R: A language and environment for statistical computing R Foundation for Statistical
Computing, Vienna, Austria. URL https://www.R-project.org/
Robinson, M. D., McCarthy, D. J., Smyth, G. K. (2010). edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics, 26(1), 139-140.
https://doi.org/10.1093/bioinformatics/btp616
Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., Smyth, G. K. (2015). limma powers differential
expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research, 43(7), e47.
https://doi.org/10.1093/nar/gkv007
Serra, A., Saarimäki, L. A., Fratello, M., Marwah, V. S., Greco, D. (2020). BMDx: a graphical Shiny application
to perform Benchmark Dose analysis for transcriptomics data. Bioinformatics, 36(9), 2932-2933.
https://doi.org/10.1093/bioinformatics/btaa030

313