**IDSL.CSA: Composite Spectra Analysis for Chemical Annotation of Untargeted Metabolomics Datasets**

Sadjad Fakouri Baygi[1], Yashwant Kumar[2], Dinesh Kumar Barupal[1]*

[1] Department of Environmental Medicine and Public Health, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA

[2] Non-communicable Diseases Division, Translational Health Science and Technology Institute, Faridabad, Haryana, 121001, India

* Corresponding author: Address: CAM Building, 3rd floor, 17 E 102nd St, New York, NY 10029 Email: dinesh.kumar@mssm.edu

**Abstract:**

Poor chemical annotation of high-resolution mass spectrometry data limit applications of untargeted metabolomics datasets. Our new software, the Integrated Data Science Laboratory for Metabolomics and Exposomics – Composite Spectra Analysis (IDSL.CSA) R package, generates composite mass spectra libraries from MS1-only data, enabling the chemical annotation of LC/HRMS peaks regardless of the availability of MS2 fragmentation spectra. We demonstrate comparable annotation rates for commonly detected endogenous metabolites in human blood samples using IDSL.CSA libraries versus data dependent acquisition (DDA) MS2 libraries in validation tests. IDSL.CSA can create and search composite spectra libraries from any untargeted metabolomics dataset generated using high-resolution mass spectrometry coupled to liquid or gas chromatography. The cross-applicability of these libraries across independent studies can improve overall annotation rates in metabolomics and exposomics projects, providing access to new biological insights that may be missed due to the lack of MS2 fragmentation data. The IDSL.CSA package is available in the R CRAN repository (https://cran.r-project.org/package=IDSL.CSA. Detailed documentation and tutorials are provided at https://github.com/idslme/IDSL.CSA .

**Keywords: Metabolomics, Peak Annotation, High resolution mass spectrometry, Spectra Search**

**Introduction**

Metabolomics has a great potential to uncover biological mechanisms and biomarkers that can be translated to innovative products to improve human health, food security and the environment. One major approach of metabolomics is the untargeted profiling of the small molecule chemical space in a biospecimen to create large databases of chemical measurements under different phenotype[1] and genotype backgrounds[1, 2]. High-resolution mass spectrometry coupled with liquid chromatography (LC/HRMS) is the most commonly used technique for untargeted metabolomics assays. However, not all peaks in an LC/HRMS dataset are annotated with chemical information, mainly for two reasons 1) MS2 fragmentation data are not available for every detected LC/HRMS peak and 2) Available MS2 libraries poorly cover the expected chemical space. As a result, a large volume of untargeted metabolomics data remains to be underutilized, leading to missed discovery opportunities.

A single chemical compound can generate several ions species due to chemical reactions
including fragmentation and adduct formations in the electrospray ionization (ESI) source,
collectively known as the MS1 data. Sporadically, selected ions (precursors) can be fragmented
further to generate mass spectra (MS2) that can be searched against data-dependent acquisition
(DDA) mass spectral libraries such as NIST 2020 for chemical annotation[3, 4] along with several
advances in annotating and interpreting DDA MS2 data[5-7]. Existing approaches[8-11] for MS1 data
annotation are limited to group the correlating ions in order to flag LC/HRMS peaks as potential
isotopologues, adducts or in-source fragments or assigning molecular formula using isotope
patterns[12]. Promisingly, these approaches including RAMClust[8], eISA[11], CorrDec[13], IIMN[10], MS-
FLO[14], and MetaboAnnotatoR[15] also suggests that co-eluting ions at MS1 level can be exported
as a spectrum which may match an entry in a DDA MS/MS library, enabling the chemical
annotation of untargeted metabolomics data using only MS1 data. Consistent ion annotation for
common metabolites across multiple studies suggests that related ESI signals for a compound
are trackable and can be utilized to annotate LC/HRMS peaks[12, 16-19]. There is a need to advance
the computational resources to annotate compounds using these grouped peaks in untargeted
LC/HRMS datasets. Converting these correlating ions on MS1 level to re-usable mass spectra
libraries can be useful for annotating peaks in independent studies that may or may not have MS2
data.

59
Here, we propose to create composite spectra libraries of correlating LC/HRMS peaks in the MS1
data from untargeted metabolomics assays and to use these libraries for annotating peaks by
mass spectral library searches. For that, we have developed a new R package IDSL.CSA
(https://cran.r-project.org/package=IDSL.CSA) for creating CSA libraries and a companion R
package IDSL.FSA (https://cran.r-project.org/package=IDSL.FSA ) for mass spectral similarity
searches.

**Results:**

First, we analyzed the elution profile for authentic reference compounds in LC/HRMS data. There
were 255 chromatographic peaks detected for a test file (pool_P1_A1-A12.mzML) by the
IDSL.IPA software[19] ( https://zenodo.org/record/7530009 ). Out of those, 153(60%) had at least
one neighboring peak in a retention time window of 0.01 minutes. Many of these co-occurring
ions may have originated from a single compound eluting from the LC column. For example, a
cluster of 209.292, 192.065 and 146.060 m/z ions related to Kynurenine compound standard
followed an almost identical elution profile (Figure S.1) at retention time 4.18 minutes. To
automatically test if those neighboring peaks are related to a single compound, we have
developed a new spectra deconvolution algorithm to group them by measuring the elution profile
similarity. Extracted ion chromatograms (EICs) for individual peaks were extracted from the raw
data file. Then, EICs were smoothed using the Local Polynomial Regression Fitting (LOESS)
regression to provide a robust estimate of the elution profile similarity. At a Pearson correlation
coefficient threshold of 0.98 between the EICs, 76.9% peaks were grouped into 42 EIC groups or
CSA clusters ( https://zenodo.org/record/7530028 ) for the test file. Clusters were selected only if
they had at least two peaks from the IDSL.IPA peak list (https://zenodo.org/record/7530009 ) and
a difference between minimum and maximum m/z values was greater than eight. These two
criteria ensured that the CSA clusters had m/z values outside the isotopologue ranges which
obviously have highly similar elution profiles. Observations of these clusters motivated us explore
their utility in annotating compounds in untargeted LC/HRMS datasets.

We checked if a CSA cluster can represent a deconvoluted mass spectrum for a compound. For
each CSA cluster, we exported all m/z values within a cluster, the retention time at the apex, the
peak height values at the apex of for each ion's EIC as the intensity values, the correlation
statistics and the additional metadata to a mass spectra file in the NIST MSP format (
https://zenodo.org/record/7530023). The file had 42 spectra. They were named as 'CSA Spectra'.
Searching these spectra against the NIST 2020 library suggested high confidence annotations
for 5 clusters (https://zenodo.org/record/7530063 ). See the method section for spectral search
parameters. Top hits for these clusters were Purine, L-Kynurenine, N-Acetyl histidine,
Homoserine, N-Acetylneuraminic acid (Figure S.2). The annotations were also confirmed by the
data dependent acquisition (DDA) MS2 spectra (Figure S.3, https://zenodo.org/record/7530151 ).
These results suggested that CSA Spectra may be used for annotating compounds in LC-HRMS
data with limited or no availability of DDA data which consistent with previous reports (cite
RAMClust[8] and MetaboAnnotatorR[15]).

We developed a workflow for automatically creating a CSA spectra library for authentic standards.
A total of 359 unique chemical standards had confirmed reverse phase retention time and
precursor m/z values across 54 LC/HRMS data in the ESI positive mode (
https://zenodo.org/record/7530170 ). For these standards, 214(59.6%) were associated with a
CSA spectrum by the workflow. Searching these CSA spectra against mass spectral databases
suggested that only up to 42(19.6%) were similar to DDA spectra, indicating that a majority of
CSA spectra contained different in-source ESI adducts. The spectra library is provided at
(https://zenodo.org/record/7530184 ).

Next, we extended the CSA workflow to cover known annotations that have been reported for
LC/HRMS data of biological specimens. In a publicly available untargeted metabolomics study
(ST000923) of human stool samples, 177 annotations were targeted in 600 LC/HRMS data files
(https://zenodo.org/record/7530227 ) for creating a CSA spectra library. Only MS1 data were
available for this study. Up to 105(59.3%) compounds had an associated CSA spectra with a
median frequency of 204, min of 1, max of 589 and a total of 25,351 CSA spectra were
deconvoluted for the target compounds from 600 samples. Then, we identified variants of CSA
spectra that are detected across multiple LC/HRMS data files. By using the entropy similarity
approach[20] (See method), we found that the average number of CSA spectra was 2.5 per
compound. Searching these CSA spectra against mass spectral databases suggested that only
46(26%) were similar to DDA spectra. The CSA spectra library for the annotated compounds for
the study has been provided at (https://zenodo.org/record/7530237 ).

Next, we scaled and applied the CSA deconvolution algorithm to the ST001000 (n=200) study to
detect all possible CSA spectra in LC/HRMS data files. For this study, a total of 139,018 CSA
spectra were deconvoluted with a median of 631 per file. A total of 16,659 unique CSA spectra
with at least 5% detection frequency were detected for this study
(https://zenodo.org/record/7530245 ).

To annotate the deconvoluted CSA spectra, we have first searched them against MS/MS
databases. For the ST001000 study, 802(4.8%) CSA spectra had a high confidence spectral
similarity match (https://zenodo.org/record/7530263 ). Because a compound can have multiple
CSA variants across the entire study, unique base peaks by chemical name and InChiKeys were
selected to subset the aligned peak table for the study. A total of 377 unique peaks were selected
and annotated with 321 unique InChiKeys and 343 unique chemical names

(https://zenodo.org/record/7530275 ). This subset aligned table represents a data matrix for
annotated compounds by the CSA workflow.

Next, we tested if a CSA spectra library created for one study can enable annotations of
compounds in a different study. We searched the unique CSA spectra variants for ST001000
study against the CSA library that we have generated earlier for the study ST000923. The spectral
search results (https://zenodo.org/record/7530302 ) suggested that 34 additional compounds
(InChiKeys) can be annotated for study ST001000 that were not covered by the DDA library
search (https://zenodo.org/record/7530310 ).

To verify the accuracy of the annotations obtained using only the spectral searches for CSA
spectra, we have matched their retention time against the published data dictionaries for
ST001000 and ST000923 studies (See methods). Data for both studies were collected by the
same laboratory using identical analytical conditions, but ST000923 had 174 annotated
compounds (InChiKeys), of which 89 were not previously annotated for the ST001000 study. Of
the 488 annotations (InChiKeys) by CSA spectra, 60 were confirmed by matching their retention
time in the published data dictionaries for ST000923 and ST000100 studies in the Metabolomics
WorkBench repository. Only 26(26.4%) of the hits were probable false positives or in-source
fragments. The IDSL.CSA workflow suggested 301 new annotations (unique InChiKeys) for the
ST001000 study (https://zenodo.org/record/7530359 ).

To demonstrate the biological significance of the annotated compounds for the ST001000 study,
we conducted a chemical set enrichment analysis using ChemRICH software[21]. ChemRICH
identified the chemical classes that were found to be significantly different between individuals
with Crohn's disease and healthy controls (https://zenodo.org/record/7530364 ). Our analysis
revealed that several chemical classes, including cholic acids, amino acids, aminosalicylic acids,
biogenic amines, hexosamines, and vitamins, were significantly different between these two
groups (https://zenodo.org/record/7530366 ). The ChemRICH result suggests that the altered gut
microbiota ecology[22] in Crohn's disease patients may also be connected to metabolic pathways
involving these chemical classes.

Finally, we have incorporated the workflows for creating CSA libraries and annotating those using
mass spectral similarity searches into two standard R package called 'IDSL.CSA' and 'IDSL.FSA'
available on CRAN repository at (https://cran.r-project.org/package=IDSL.CSA) and .
(https://cran.r-project.org/package=IDSL.FSA ). We have also added workflows for processing
data-dependent (DDA) and data-independent (DIA) acquisitions. The package includes a user-
friendly parameter file in Microsoft Excel format that allows users to run different workflows and
ensure reproducible data processing. Additionally, we also extended the CSA deconvolution to
cover nominal mass data for GC/MS data using a secondary 'IDSL.NPA' R package
(https://cran.r-project.org/package=IDSL.NPA). The parameter tables are extensive and cover
commonly used settings as well as several new parameters to optimize spectra deconvolution
and spectral searches (Table S1-S4). Documentation, tutorials, and code for the software are
available for IDSL.CSA and IDSL.FSA R packages in the GitHub repository at
https://github.com/idslme/IDSL.CSA and https://github.com/idslme/IDSL.FSA, respectively.

**Discussion:**

We have developed a simple and easy to use integrated workflow of IDSL.CSA and IDSL.FSA R
packages that can improve the chemical annotation rates of LC/HRMS peaks in untargeted
metabolomics datasets. The integrated approach using IDSL.IPA[19], IDSL.CSA and IDSL.FSA R

packages (Figure 1) contains easy to use steps for 1) creating CSA, DDA and DIA spectral
libraries 2) performing mass spectral similarity searches using spectral entropy[20], cosine
similarity, normalized Euclidean mass error[12] 3) prefiltering library spectra for faster searches
(Section S.1) 4) refining deconvolution results aligned table (Section S.2) and 5) ranking
annotations using spectra search results from all the samples within a study.

We have developed a new workflow to create composite spectra libraries from LC-HRMS
datasets. A CSA spectrum may be similar to a counter DDA spectrum of the compound but may
often include additional ionization reactions (Figure S.4). As recommended by earlier reports[12, 16-
19], IDSL.CSA workflows also utilize both individual file level geometry alignment of LC/HRMS
peaks and the co-detection frequency across all samples within a study to capture different
variants of CSA spectra for a compound. For example, two variants of CSA spectra for kynurenine
can contain different ESI adducts such as $[M+Na]^+$ and $[M-H_2O+H]^+$ (Figure S.5). The composition
of CSA variant spectra for a chemical depends on the instrument type, analytical method, sample
matrix, gradient additives and other factors. These variants can increase the specificity of library
searches. We argue that these CSA spectra should be catalogued in a mass spectral library and
in a community driven MS database such as GNPS[10]. A CSA mass spectral library is a collection
of unique composite spectra variants of different chemical compounds which can be created using
MS1 only data for annotated peaks and authentic standards. To create CSA libraries, raw
LC/HRMS data for known annotations or authentic standards are needed, which are readily
available for over 2000 publicly available metabolomics datasets in EBI Metabolights[23], GNPS
Massive[24] and Metabolomics Workbench[25] repositories.

The IDSL.CSA workflows incorporate several essential and novel features for processing
untargeted metabolomics datasets. It encapsulates a full workflow of steps including peak
detection[19], alignment, DDA/DIA deconvolution, library generation, spectra search and annotation
ranking, into a single line of R command that need all parameters in an input Microsoft Excel file.
The workflow applies a critical step of LOESS smoothing followed by the cubic spline smoothing
method to minimize chromatogram jaggedness while computing correlation among ion intensities.
CSA variants and consensus CSA spectra both were created for a compound to enable cross-
instrument searches. It also interprets a CSA spectrum by identifying ion species of commonly
observed ESI adducts. The optimized pre-filtering methods using precursor, spectra markers,
spectra entropy enabled faster searching of larger libraries with millions of spectra. A key unique
strength of our approach is to identify recurring CSA spectra across multiple samples which is
helpful in generating high confidence CSA libraries. It also standardizes publicly available mass
spectra data with inconsistent fields to a standard storage format in R, making IDSL.CSA and
IDSL.FSA packages fully compatible with existing public MS/MS libraries and the NIST MS/MS
database. With these features, IDSL.CSA workflow can streamline generation of high-quality data
matrices with improved chemical annotation rates from untargeted metabolomics datasets.
IDSL.CSA is a useful addition to the growing pool of software to improve the annotation of MS1
data in untargeted metabolomics studies.

It is a major challenge in metabolomics that 2/3 of untargeted studies in the Metabolomics
Workbench data repository (https://www.metabolomicsworkbench.org) have only unannotated
peaks. To overcome this hurdle to some extent, IDSL.CSA workflows may be able to annotate a
substantial number of peaks in these studies using available reference DDA and newly created
CSA libraries. By utilizing MS1 data in an exhaustive way, our workflow can minimize the

underutilization of untargeted metabolomics datasets in studying basic metabolic processes and
biomonitoring of environmental chemical exposure.

**Material and methods:**

**Publicly available LC/HRMS test datasets:** Raw mass spectra data and known annotations
were accessed for MSV000088661, ST000923 and ST001000 from Massive UCSD
(https://massive.ucsd.edu ), Metabolomics WorkBench
(https://www.metabolomicsworkbench.org) data repositories. Raw data were converted to the
centroid mzML format using the MSConvert utility.

**R packages:** IDSL.CSA R package has been provided via the R-CRAN repository (https://cran.r-
project.org/package=IDSL.CSA). IDSL.MXP R package[19] (https://CRAN.R-
project.org/package=IDSL.MXP) was used to read LC/HRMS files in mzML/mzXML/netCDF data
formats. IDSL.IPA R package (version 2.6)[19] (https://CRAN.R-project.org/package=IDSL.IPA)
was used for generating chromatographic peak lists and aligned peak tables for each study.
IDSL.FSA R package (https://cran.r-project.org/package=IDSL.FSA) was used for mass spectral
similarity searches.

**Generation of mass spectral libraries:** IDSL.CSA workflow can generate three types of mass
spectral libraries – 1) composite spectra (CSA) 2) data-dependent acquisition (DDA) and 3) data
independent acquisition (DIA). **CSA:** LC/HRMS peaks from IDSL.IPA peak list for each sample
were grouped by estimating the Pearson correlation among the chromatographic peaks after
LOESS smoothing followed by a cubic spline interpolation. First, peaks were sorted by intensity
and the most abundant peak was selected as a seed peak for creating a CSA spectrum. Next, a
retention time threshold was applied to narrow down candidate fragmentation peaks. Then,
Pearson correlation was estimated among the candidate peaks. One peak was assigned to only
a single CSA spectrum to avoid redundancies. For true positive annotations, the seed peak was
reported m/z value. **DDA:** DDA spectra with precursor details were extracted. When more than
one DDA spectra for a IDSL.IPA peak was observed, three options were provided 1) integrated
all DDA spectra for a peak 2) select the most abundant 3) de-noise spectra by correlation
statistics. **DIA:** DIA spectra were created by the same approach used for CSA, except that the
fragment ions were obtained from the $MS^E$ (MS level = 2) data channels[16]. Spectra from these
three methods were exported to the MSP format. Parameter files are provided at
https://zenodo.org/record/7530387.

**Mass spectral similarity search:** Spectral similarity search parameters were provided in a
Microsoft excel format (https://zenodo.org/record/7530387) to the IDSL.FSA R package. If mass
spectral libraries for standards were from different sources and in different formats such as mgf,
they were converted to a standard msp file and separated by ionization mode (+/-). To enable
faster searches in R, msp files for known annotations and authentic standards were converted
into a fragmentation spectra database (FSDB) that stored pre-calculated spectral entropy for each
spectrum in the MSP file (https://zenodo.org/record/7530387). FSDBs for public databases
including the Global Natural Product Social Molecular Networking (GNPS) and Mass bank of
North America (MoNA) for positive and negative modes are provided at
(https://zenodo.org/record/7530397). Similar to Li *et. al.,*[20] adjacent fragments within the
instrument resolution were resolved followed by a noise removal (Default = 1%). A threshold
above a baseline (%) was used to determine characteristic spectra markers, and then a minimum

threshold (%) of matched characteristic spectra markers was used to allow partial symmetry matches in case of presence of instrumental noises. Cosine and entropy similarity scores were computed for spectra matching. The cosine similarity was computed using equation (1) only using the library fragments.

$$CosineSimilarity = \sum_{i=1}^{NP} \frac{I_i^{lib} I_i^{exptl}}{\sqrt{\sum_{i=1}^{NP}(I_i^{lib})^2} \sqrt{\sum_{i=1}^{NP}(I_i^{exptl})^2}} \tag{1}$$

where $I_i$ and $NP$ represent the intensity of the fragment, and number of matched fragment peaks in the fragmentation spectra, respectively. Superscripts of *lib* and *exptl* represent library and experimental fragmentation spectra, respectively. Entropy similarity was calculated from spectral entropy (*S*) values described by Li *et. al.*[20]

$$S = -\sum_p I_p \ln I_p \tag{2}$$

where $I_p$ is normalized intensity to the summation of intensities ($\Sigma I_p = 1$). A merged mass spectrum *lib:exptl* is generated by 1:1 mixing the normalized *lib* and *exptl* spectra. Then, the entropy similarity was calculated using equation (3).

$$EntropySimilarity = 1 - \frac{2S^{lib:exptl} - S^{lib} - S^{exptl}}{\ln 4} \tag{3}$$

An additional weight transformation was proposed to adjust spectral entropy using equation (4).

$$I_p^{new} = I_p^w \begin{cases} w = 1 & , \quad S \geq 3 \\ w = 0.25(1 + S), & \quad S < 3 \end{cases} \tag{4}$$

The transformed $I_p^{new}$ should also be re-normalized ($\Sigma I_p^{new} = 1$). Normalized Euclidean mass error (*NEME*) was calculated using the equation (5) to further asses the fragmentation spectra. An entropy similarity cutoff (Default ≥ 0.75) may be used to filter out candidate hits.

$$NEME = \sqrt{\frac{\sum_{i=1}^{NP}(M_i^{lib} - M_i^{exptl})^2}{NP}} \tag{5}$$

where $M_i$ represents the mass of the fragmentation peaks. A maximum threshold for *NEME* may be used to cut off hits with higher mass errors.

When the entire candidate hits were matched for an experimental fragmentation spectra, the candidate hits are sorted using the following equation.

$$Score = CosineSimilarity * EntropySimilarity^2 \tag{6}$$

For each sample in a study, spectra search result table was generated. An example table is provided at (https://zenodo.org/record/7530263 ). If the aligned table generated by the IDSL.IPA R package was available, spectra search results were summarized for annotation frequency and overall median ranks.

**Parameter files:** IDSL.IPA and IDSL.CSA parameter files for MSV000088661, ST000923 and ST001000 studies are provided at (https://zenodo.org/record/7530409). These files include – 1) creating reference CSA spectra for authentic standard 2) creating CSA spectra for true positive annotation ST001000 study and 3) creating CSA spectra in a full deconvolution mode for

ST001000 and 4) IDSL.IPA data processing settings. After filling up parameters in these files, the excel sheets were used directly as an input for running the workflow function in the IDSL.CSA R package.

**Statistical analysis:** Student *t*-test and ChemRICH analysis was performed in R to find significantly different chemical and chemical sets in the Crohn's disease patients in comparison to the healthy control group.

**Funding:** The research is in part supported by NIH grants U2CES026561, R01ES032831, R01ES033688, U2CES026555 P30ES023515, T32HD049311, K12ES033594, U2CES030859, UL1TR004419 and UL1TR001433.

**Conflict of interest:** DKB has been a consultant for the Brightseed Inc, California, USA.

**Author's contribution:** SFB and DKB planned the study, prepared the results and drafted the manuscript. SFB and DKB coded the IDSL.CSA package. YK provided test LC/HRMS data for authentic standards of metabolites. All authors have reviewed the manuscript content.

**Supporting information:**

Supporting spreadsheet tables, data sources, workflow and computational method details and additional benchmarks.

# References:

(1) Cheng, Y.; Schlosser, P.; Hertel, J.; Sekula, P.; Oefner, P. J.; Spiekerkoetter, U.; Mielke, J.; Freitag, D. F.; Schmidts, M.; Investigators, G.; et al. Rare genetic variants affecting urine metabolite levels link population variation to inborn errors of metabolism. *Nat Commun* **2021**, *12* (1), 964. DOI: 10.1038/s41467-020-20877-8

(2) Hu, J.; Yao, J.; Deng, S.; Balasubramanian, R.; Jimenez, M. C.; Li, J.; Guo, X.; Cruz, D. E.; Gao, Y.; Huang, T.; et al. Differences in Metabolomic Profiles Between Black and White Women and Risk of Coronary Heart Disease: an Observational Study of Women From Four US Cohorts. *Circ Res* **2022**, *131* (7), 601-615. DOI: 10.1161/CIRCRESAHA.121.320134

(3) Yang, X.; Neta, P.; Stein, S. E. Quality control for building libraries from electrospray ionization tandem mass spectra. *Anal Chem* **2014**, *86* (13), 6393-6400. DOI: 10.1021/ac500711m

(4) Yang, X.; Neta, P.; Stein, S. E. Extending a Tandem Mass Spectral Library to Include MS(2) Spectra of Fragment Ions Produced In-Source and MS(n) Spectra. *J Am Soc Mass Spectrom* **2017**, *28* (11), 2280-2287. DOI: 10.1007/s13361-017-1748-2

(5) Domingo-Almenara, X.; Montenegro-Burke, J. R.; Benton, H. P.; Siuzdak, G. Annotation: A Computational Solution for Streamlining Metabolomics Analysis. *Anal Chem* **2018**, *90* (1), 480-489. DOI: 10.1021/acs.analchem.7b03929

(6) Duhrkop, K.; Fleischauer, M.; Ludwig, M.; Aksenov, A. A.; Melnik, A. V.; Meusel, M.; Dorrestein, P. C.; Rousu, J.; Bocker, S. SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure information. *Nat Methods* **2019**, *16* (4), 299-302. DOI: 10.1038/s41592-019-0344-8

(7) Guijas, C.; Montenegro-Burke, J. R.; Domingo-Almenara, X.; Palermo, A.; Warth, B.; Hermann, G.; Koellensperger, G.; Huan, T.; Uritboonthai, W.; Aisporna, A. E.; et al. METLIN: A Technology Platform for Identifying Knowns and Unknowns. *Anal Chem* **2018**, *90* (5), 3156-3164. DOI: 10.1021/acs.analchem.7b04424

(8) Broeckling, C. D.; Afsar, F. A.; Neumann, S.; Ben-Hur, A.; Prenni, J. E. RAMClust: a novel feature
clustering method enables spectral-matching-based annotation for metabolomics data. *Anal Chem*
**2014**, *86* (14), 6812-6817. DOI: 10.1021/ac501530d
(9) Kuhl, C.; Tautenhahn, R.; Bottcher, C.; Larson, T. R.; Neumann, S. CAMERA: an integrated strategy for
compound spectra extraction and annotation of liquid chromatography/mass spectrometry data sets.
*Anal Chem* **2012**, *84* (1), 283-289. DOI: 10.1021/ac202450g
(10) Schmid, R.; Petras, D.; Nothias, L. F.; Wang, M.; Aron, A. T.; Jagels, A.; Tsugawa, H.; Rainer, J.; Garcia-
Aloy, M.; Duhrkop, K.; et al. Ion identity molecular networking for mass spectrometry-based
metabolomics in the GNPS environment. *Nat Commun* **2021**, *12* (1), 3832. DOI: 10.1038/s41467-021-
23953-9
(11) Xue, J.; Domingo-Almenara, X.; Guijas, C.; Palermo, A.; Rinschen, M. M.; Isbell, J.; Benton, H. P.;
Siuzdak, G. Enhanced in-Source Fragmentation Annotation Enables Novel Data Independent Acquisition
and Autonomous METLIN Molecular Identification. *Anal Chem* **2020**, *92* (8), 6051-6059. DOI:
10.1021/acs.analchem.0c00409
(12) Baygi, S. F.; Banerjee, S. K.; Chakraborty, P.; Kumar, Y.; Barupal, D. K. IDSL.UFA Assigns High-
Confidence Molecular Formula Annotations for Untargeted LC/HRMS Data Sets in Metabolomics and
Exposomics. *Anal Chem* **2022**, *94* (39), 13315-13322. DOI: 10.1021/acs.analchem.2c00563
(13) Tada, I.; Chaleckis, R.; Tsugawa, H.; Meister, I.; Zhang, P.; Lazarinis, N.; Dahlen, B.; Wheelock, C. E.;
Arita, M. Correlation-Based Deconvolution (CorrDec) To Generate High-Quality MS2 Spectra from Data-
Independent Acquisition in Multisample Studies. *Anal Chem* **2020**, *92* (16), 11310-11317. DOI:
10.1021/acs.analchem.0c01980
(14) DeFelice, B. C.; Mehta, S. S.; Samra, S.; Cajka, T.; Wancewicz, B.; Fahrmann, J. F.; Fiehn, O. Mass
Spectral Feature List Optimizer (MS-FLO): A Tool To Minimize False Positive Peak Reports in Untargeted
Liquid Chromatography-Mass Spectroscopy (LC-MS) Data Processing. *Anal Chem* **2017**, *89* (6), 3250-
3255. DOI: 10.1021/acs.analchem.6b04372
(15) Graca, G.; Cai, Y.; Lau, C. E.; Vorkas, P. A.; Lewis, M. R.; Want, E. J.; Herrington, D.; Ebbels, T. M. D.
Automated Annotation of Untargeted All-Ion Fragmentation LC-MS Metabolomics Data with
MetaboAnnotatoR. *Anal Chem* **2022**, *94* (8), 3446-3455. DOI: 10.1021/acs.analchem.1c03032
(16) Baygi, S. F.; Fernando, S.; Hopke, P. K.; Holsen, T. M.; Crimmins, B. S. Nontargeted Discovery of
Novel Contaminants in the Great Lakes Region: A Comparison of Fish Fillets and Fish Consumers. *Environ*
*Sci Technol* **2021**, *55* (6), 3765-3774. DOI: 10.1021/acs.est.0c08507
(17) Fakouri Baygi, S.; Fernando, S.; Hopke, P. K.; Holsen, T. M.; Crimmins, B. S. Automated Isotopic
Profile Deconvolution for High Resolution Mass Spectrometric Data (APGC-QToF) from Biological
Matrices. *Anal Chem* **2019**, *91* (24), 15509-15517. DOI: 10.1021/acs.analchem.9b03335
(18) Fakouri Baygi, S.; Fernando, S.; Hopke, P. K.; Holsen, T. M.; Crimmins, B. S. Decadal Differences in
Emerging Halogenated Contaminant Profiles in Great Lakes Top Predator Fish. *Environ Sci Technol* **2020**,
*54* (22), 14352-14360. DOI: 10.1021/acs.est.0c03825
(19) Fakouri Baygi, S.; Kumar, Y.; Barupal, D. K. IDSL.IPA Characterizes the Organic Chemical Space in
Untargeted LC/HRMS Data Sets. *J Proteome Res* **2022**, *21* (6), 1485-1494. DOI:
10.1021/acs.jproteome.2c00120
(20) Li, Y.; Kind, T.; Folz, J.; Vaniya, A.; Mehta, S. S.; Fiehn, O. Spectral entropy outperforms MS/MS dot
product similarity for small-molecule compound identification. *Nat Methods* **2021**, *18* (12), 1524-1531.
DOI: 10.1038/s41592-021-01331-z
(21) Barupal, D. K.; Fiehn, O. Chemical Similarity Enrichment Analysis (ChemRICH) as alternative to
biochemical pathway mapping for metabolomic datasets. *Sci Rep* **2017**, *7* (1), 14567. DOI:
10.1038/s41598-017-15231-w

(22) Franzosa, E. A.; Sirota-Madi, A.; Avila-Pacheco, J.; Fornelos, N.; Haiser, H. J.; Reinker, S.; Vatanen, T.;
Hall, A. B.; Mallick, H.; McIver, L. J.; et al. Gut microbiome structure and metabolic activity in
inflammatory bowel disease. *Nat Microbiol* **2019**, *4* (2), 293-305. DOI: 10.1038/s41564-018-0306-4
(23) Haug, K.; Cochrane, K.; Nainala, V. C.; Williams, M.; Chang, J.; Jayaseelan, K. V.; O'Donovan, C.
MetaboLights: a resource evolving in response to the needs of its scientific community. *Nucleic Acids Res*
**2020**, *48* (D1), D440-D444. DOI: 10.1093/nar/gkz1019
(24) Wang, M.; Carver, J. J.; Phelan, V. V.; Sanchez, L. M.; Garg, N.; Peng, Y.; Nguyen, D. D.; Watrous, J.;
Kapono, C. A.; Luzzatto-Knaan, T.; et al. Sharing and community curation of mass spectrometry data with
Global Natural Products Social Molecular Networking. *Nat Biotechnol* **2016**, *34* (8), 828-837. DOI:
10.1038/nbt.3597
(25) Sud, M.; Fahy, E.; Cotter, D.; Azam, K.; Vadivelu, I.; Burant, C.; Edison, A.; Fiehn, O.; Higashi, R.; Nair,
K. S.; et al. Metabolomics Workbench: An international repository for metabolomics data and metadata,
metabolite standards, protocols, tutorials and training, and analysis tools. *Nucleic Acids Res* **2016**, *44*
(D1), D463-470. DOI: 10.1093/nar/gkv1042

**Figure 1.** Integrated workflow of the IDSL.IPA, IDS.CSA and IDSL.FSA packages to deconvolute and annotate composite spectra.

**For Table of Contents Only**