1

An INDEL genomic approach to explore population diversity of phytoplankton

Martine Devic*[1], Louis Dennu[1], Jean-Claude Lozano[1,] Cédric Mariac[2], Valérie Vergé[1], Philippe Schatt[1], François-Yves Bouget*[1], François Sabot*[2]

*Corresponding authors :

martine.devic@obs-banyuls.fr

francois-yves.bouget@obs-banyuls.fr

francois.sabot@ird.fr

Author affiliations :

[1] Laboratoire d'Océanographie Microbienne (LOMIC), CNRS/Sorbonne University, UMR 7621, Observatoire Océanologique, 66650 Banyuls/mer, France.

[2] Diversité, adaptation et développement des plantes (DIADE), University of Montpellier, Cirad, IRD, UMR 232, Centre IRD de Montpellier, 911 avenue Agropolis, BP 604501, 34394, Montpellier Cedex 5, France.

Keywords : Phytoplankton, *Bathycoccus*, Intraspecific Diversity, Genotyping, Bloom

Abstract (258 words)

Background: Although metabarcoding and metagenomic approaches have generated large datasets on worldwide phytoplankton species diversity, the intraspecific genetic diversity underlying the genetic adaptation of marine phytoplankton to specific environmental niches remains largely unexplored. This is mainly due to the lack of biological resources and tools for monitoring the dynamics of this diversity in space and time. **Results:** To gain insight into population diversity, a novel method based on INDEL markers was developed on *Bathycoccus prasinos* (Mamiellophyceae), an abundant and cosmopolitan species with strong seasonal patterns. Long read sequencing was first used to characterise structural variants among the genomes of six *B. prasinos* strains sampled from geographically distinct regions in the world ocean. Markers derived from identified insertions/deletions were validated by PCR then used to genotype 55 *B. prasinos* strains isolated during the winter bloom 2018-2019 in the bay of Banyuls-sur-Mer (Mediterranean Sea, France). This led to their classification into eight multi-loci genotypes and the sequencing of strains representative of local diversity, further improving the available genetic diversity of *B. prasinos*. Finally, selected markers were directly tracked on environmental DNA sampled during 3 successive blooms from 2018 to 2021, showcasing a fast and cost-effective approach to follow local population dynamics.

Conclusions: This method, which involves (i) pre-identifying the genetic diversity of *B. prasinos* in environmental samples by PCR, (ii) isolating cells from selected environmental samples and (iii) identifying genotypes representative of *B. prasinos* diversity for sequencing, can be used to comprehensively describe the diversity and population dynamics not only in *B. prasinos* but also potentially in other generalist phytoplankton species.

Background

Marine phytoplankton, including picoeukaryotic algae, is responsible for a large fraction of

primary production [1]. In temperate regions, the abundance and diversity of the phytoplankton

is often seasonal and occurs in bursts, as algal blooms. *Per se*, blooms have a large impact on

global primary production and therefore the understanding of the genetic basis of

phytoplankton adaptation to seasonal niches and the effects of ocean warming on

phytoplankton blooms are of the utmost importance.

Meta-ribosomal barcoding on the nuclear or plastidial 18/16S rRNA gene has opened the

access to massive data in time and space and has accelerated the study of phytoplankton

species and genera diversity in natural communities. However, metabarcoding approaches do

not provide information on intraspecific genetic variations. Assessing only interspecific

diversity leads to the underestimation of the diversity of the populations. Equally, a single

isolate cannot represent the diversity of a population. Since natural selection acts on variation

among individuals within populations, it is essential to incorporate both intra and interspecific

trait variability into community ecology [2]. Indeed, Raffard et al. [3] demonstrated that

intraspecific variation has significant ecological effects across a large set of species,

confirming a previous estimate based on a more restricted species set [4]. Furthermore, it has

been shown that diversity within species is rapidly decreasing, making them more

homogeneous and highlighting the need to preserve intraspecific variations since this

intraspecific diversity reinforces the overall population stability in the face of environmental

change [4].

In diatoms, intraspecific variations play a key role in the response of species to several

important environmental factors such as light, salinity, temperature and nutrients [5]. Modelling

approaches suggest that intraspecific variation extends bloom periods by providing variability

in competitive interactions between species under changing conditions, and contributes to

fitness in temporary microhabitats. Intraspecific genetic diversity allows local adaptation by

optimising physiological responses, allowing local populations to be exceptionally fit and

competitive in their respective habitat.

Since an assembly of genotypically diverse individuals constitutes a population,

methodological approaches have been developed in order to determine the genetic variation

among individuals [6]. One of the main challenges is the difficulty of isolating a sufficient

number of individuals for classical diversity analyses while ensuring representativity of diverse

subpopulations. Historically, sequence-based intraspecific markers correspond to small

nucleotide repeats such as microsatellites [7], chloroplastic [8] and mitochondrial [9] genes

and a few nuclear genes that were applied to several hundreds of isolated individuals. The

development of Randomly Amplified Polymorphic DNA (RAPD) markers [10] and more

recently of Restriction-site-Associated DNA sequencing techniques (RADseq) [11] allowed the

discovery and genotyping of thousands of genetic markers for any given species at relatively

low-cost. Some of these approaches have been used to analyse the diversity of populations

during algal blooms [12]. Recently, the dramatic increase of the number of sequenced

genomes led to large-scale diversity studies with large sets of nuclear genes or whole genome

comparisons. However, most of these approaches require the isolation of a large number of

individuals. As a consequence, the intraspecific diversity remains poorly documented in

marine phytoplankton.

Widely distributed from the equator to arctic and antarctic poles with a marked seasonality in

temperate and polar regions, picoeukaryotes belonging to the *Mamiellales* order

(*Bathycoccus*, *Ostreococcus* and *Micromonas*) are abundant and have a cosmopolitan

presence, illustrating a high capacity for adaptation to a wide range of contrasting

environments [13–16]. Novel, rapid and cheap sequencing technologies have given access to

Mamiellales interspecific diversity by metagenomic [16,17] and metatranscriptomic

approaches [18]. However, to date, very little information is available on intraspecific diversity

of *Bathycoccaceae*, with the exception of *Ostreococcus tauri* from mediterranean lagoons [19]

. Unlike *O. tauri*, which is usually not detectable in publicly available metagenomes,

Bathycoccus sp. is the most cosmopolitan *Mamiellophyceae* genus. Although *Bathycoccus*

sp. is abundant in the world Ocean, Metabarcoding of 18S ribosomal RNA (V9 region) does

not allow the identification of *Bathycoccus* species [20,21]. Through metagenomic analysis,

Bathycoccus genus was divided into two species, the polar and temperate *Bathycoccus*

prasinos type B1 genome [13,22] and the tropical *Bathycoccus calidus* type B2 genome

[21,23,24]. Thus the cosmopolitan nature of *Bathycoccus* from poles to equator is due to the

combination of both B1 and B2 species.

In the bay of Banyuls, *Bathycoccus* and *Micromonas* bloom yearly from November to April, *B.*

prasinos being one of the most abundant species [15,25]. The highly reproducible yearly

occurrence of *B. prasinos* in the Banyuls bay during the last decade [15] raises the question of

the persistence of a *B. prasinos* population adapted to the bay or of a variation of the

population structure each year. In addition, since outside of the bloom period *Mamiellales* are

virtually absent from the bay, is the *B. prasinos* bloom initiated by an uptake of resident "resting

cells" in the sediment or by a fresh input carried by North western Mediterranean currents

along the Gulf of Lion? At present no resting stages that can act as inoculum of subsequent

blooms have been described for *B. prasinos.*

To assess the intraspecific diversity of *B. prasinos* worldwide and locally in the bay of Banyuls,

we developed an efficient method to isolate strains together with whole genome sequencing by

Oxford Nanopore Technology (ONT) in order to identify Structural Variations (SV) in *B.*

prasinos genomes. Diversity markers designed from INDEL (insertion or deletion of bases in

the genome of an organism) were used to genotype *B. prasinos* strains and populations from

environmental samples.

Methods

Algal strains and culture conditions

World-wide *B. prasinos* strains were obtained from the Roscoff Culture Collection (RCC) centre (See Supplementary Table 1, Additional File 1). Strain RCC1105 from which the current *B. prasinos* reference genome originated [22] was lost and replaced by its clonal culture RCC4222. The strains were cultivated in 100 mL flasks in filtered artificial seawater (1×10^{-3} M TrisHCl pH 7.2, 24.55 g/L NaCl, 0.75 g/L KCl, 4.07 g/L MgCl2 6H2O, 1.47 g/L CaCl2 2H2O, 6.04 g/L MgSO4 7H2O, 0.21 g/L NaHCO3, 0.00138 g/L NaH2PO4, 0.075 g/L NaNO3, 1.28×10^{-6} g/L H2SeO3) supplemented with trace metals and vitamins B1 (300 nM) and B12 (1 nM). Cultures were maintained under constant gentle agitation in an orbital platform shaker (Heidoph shaker and mixer unimax 1010). Sunlight irradiation curves recreating realistic light regimes at a chosen latitude and period of the year were applied in temperature-controlled incubators (Panasonic MIR-154-PE).

Cell isolation

Surface water was collected at 3 metre depth at SOLA buoy in Banyuls bay, North Western Mediterranean Sea, France (42°31′N, 03°11′E) approximately every week from December, 2018 to March, 2019 ; November, 2019 to March, 2020 and October, 2020 to April, 2021. Two ml aliquots were used to determine the quantity and size of phytoplankton by flow cytometry. For the 2018/2019 bloom, 50 ml were filtered through a 1.2-µm pore-size acrodisc (FP 30/1.2 CA-S cat N° 10462260 Whatman GE Healthcare Sciences) and used to inoculate 4 culture flasks with 10 ml of filtrate each. The sea water was supplemented by vitamins B1 and B12, NaH2PO4, NaNO3 and metal traces at the same final concentration as artificial sea water (ASW), antibiotics (Streptomycin sulfate and Penicillin at 50 µg/ml) were added to half of the cultures. The cultures were incubated under light and temperature conditions similar to those during the sampling date for 3-4 weeks. The presence of picophytoplankton was analysed by a

BD Accuri C6 flow cytometer. Generally, superior results were obtained without antibiotics.

Cultures containing at least 90% of picophytoplankton with only residual nanophytoplankton

were used for plating on agarose (0.21%). Colonies appearing after 10 days were hand-picked

and further cultured in 2 ml ASW in deepwell plates (Nunc, Perkin Elmer, Hessen, Germany)

for 10 days. Cells were cryopreserved at this stage. (See Supplementary Figure 1, Additional

File 2) Circa 500 clones were cryopreserved. At the same time, DNA extraction and PCR were

performed in order to identify *B. prasinos* strains.

157 DNA extraction, genome sequencing, assembly and PCR
158 amplification

For PCR analysis, total DNA was extracted from 4 ml *B. prasinos* cell cultures according to the

Plant DNAeasy Qiagen protocol. For whole genome sequencing by Oxford Nanopore

Technology (ONT), DNA was extracted by a CTAB method from 100 ml culture principally

based on Debladis et al. [26]. ONT libraries were prepared using the Rapid Barcoding

Sequencing kit (SQK-RBK004) and deposited on R9.4 flow cells for sequencing run. For

environmental samples, 5 litres of seawater at SOLA 3 metre depth were passed through 3

microns and 0.8 micron pore filters. DNA from cells collected on the 0.8 micron filters were

extracted using the Plant DNAeasy Qiagen protocol with the addition of a proteinase K

treatment in the AP1 buffer. PCR was performed using the Red Taq polymerase Master mix

(VWR) with the required primers (See Supplementary Appendix 1, Additional File 3) and

corresponding DNA. For sequencing, the PCR products were purified using the NucleoSpin

Gel and PCR Clean-up kit (Macherey-Nagel reference 740609.50) and sent to GENEWIZ for

Sanger sequencing.

Raw ONT Fast5 data were basecalled using Guppy 4.0.15 (https://nanoporetech.com) and the

HAC model, and QC performed using NanoPlot 1.38.1 [27]. All reads with a QPHRED higher

than 8 were retained and subjected to genome assembly using Flye 2.8 [28] under standard

options for ONT data. Raw assemblies were then polished with 3 turns of Racon 1.4.3 with

standard options [29] after mapping of raw reads on the previous round sequence using

minimap2 2.24 (-ax map-ont mode) [30]. Final scaffolding was performed using Ragoo 1.1 [31]

upon the original *B. prasinos* reference genome (GCA_002220235.1) [22]. Final QC of

assemblies was performed using QUAST 5.0 [32].

181   Variant calling and size control

Draft assemblies were aligned to *B. prasinos* reference genome using MUMmer 4.0.0 [33]

nucmer (options: –maxmatch -c 500 -b 500 -l 100). Alignments were then filtered for identity

(>90%) and length (>100bp) using delta-filter from MUMmer 4.0.0 (options: -m -i 90 -l 100).

Variant calling on the resulting delta files was performed with SVanalyzer 0.36

(https://github.com/nhansen/SVanalyzer) SVrefine subcommand, and resulting VCFs were

merged with bcftools merge 1.18 [34].

Size control on VCF was performed using a set of home-made Python scripts available at

https://forge.ird.fr/diade/genomecodes under GPLv3 licence.

190   Determination of Growth Rates

Cells isolated during December, 2018, January and February, 2019 in Banyuls bay were used

in this experiment. For each culture condition the cell number was determined by flow

cytometry daily, for 9 days. The growth rate was determined as Ln(N)/dT, where N is the cell

concentration per ml and T the time (days). The determination of the maximal growth rate

(μmax) and subsequent statistical analysis were conducted in accordance with the

methodology described in Guyon et al. [35].

Results

INDELs diversity of selected *Bathycoccus prasinos* strains

Figure 1 : Structural variant identification in collection strains of B. prasinos. (A) Summary of the sequencing and analysis strategy for the identification of INDELs structural variations and the design of PCR diversity markers. (B) Distribution frequency of identified SV along a 20Kb sliding window on nuclear chromosomes. Each inner circle corresponds to the distribution of SV within a sequenced strain. The outer circle corresponds to the cumulative frequency of all sequenced strains.

With the aim of creating a genetic resource of natural variants for *B. prasinos*, we first undertook a search for genetic markers of diversity that could be used to differentiate isolates from natural communities (Fig. 1A). From the few strains available in the Roscoff culture collection, we examined the genetic diversity of *B. prasinos* and selected geographically dispersed strains, so that the markers could potentially be used locally as well as worldwide. Oxford Nanopore Technology (ONT) was used to sequence the genomes of 6 selected strains spread along a latitudinal gradient from the Baffin bay (67° N) to the Mediterranean Sea (40° N) (See Supplementary Table 1, Additional File 1). After individual *de novo* assembly, each genome was compared to the reference genome of the RCC1105 strain from the Banyuls bay

[22] to identify 200 to 2,000 bp long INDELs. Selective amplification of these variable size

INDELs by PCR generates different size DNA fragments that can be used to genotype new

isolated strains as well as to track population diversity in environmental samples. Once

sequenced, new genotypes further enrich the marker resource that could be used

subsequently for genotyping *B. prasinos* isolates (Fig. 1A).

A total of 1,346 unique SV were identified, the contribution of each strain being highly variable

(Table 1). Strain RCC5417 from the Baffin bay was the most geographically distant from the

reference strain, and also the most diverse with 536 SV of size > 200 bp identified. Conversely,

RCC1615, with only a 4X sequencing depth, made the smallest contribution to SV

identification up to 10,000 bp but it was the main contributor to larger SVs (>10,000 bp), with 73

SV detected. This may be due to the overall lower assembly contiguity and quality of this strain

genome (Table 1; See Supplementary Table 1, Additional File 1). Despite these disparities, SV

were evenly distributed along *B. prasinos* nuclear genome, with the exception of the outlier

chromosome 19 [22], for which few variations were identified, most likely because of the

hypervariable structure of this chromosome (Fig. 1B). In the case of the big outlier

chromosome 14, no contrast in variant identification could be detected with other nuclear

chromosomes.

Table 1: Number of structural variations identified in selected strains of *B. prasinos.*
Individual loci corresponds to the cumulated number of variable genomic positions without duplicates.
238

Size range

	200 - 2000 bp	2000 - 10000 bp	> 10000 bp	Total
RCC5417	371	125	40	**536**
RCC685	210	131	28	**369**
RCC4752	146	77	28	**251**
RCC1613	147	66	29	**242**
RCC1868	85	66	38	**189**
RCC1615	18	16	73	**107**
Individual loci	765	385	196	**1346**

Search for intraspecific diversity markers

Figure 2 : Distribution of structural variations shared by three or more strains. The vertical barplot represents the number of structural variations between 200 and 2000 bp detected only within the corresponding group of strains. The horizontal barplot represents the number of structural variations detected within the corresponding strain.

To select putative PCR markers of *B. prasinos* diversity several criteria were considered. The

size of the amplified fragment had to be between 0.2 and 2 Kb and sufficiently different among

genomes to be unambiguously visualised on agarose gel after amplification with a single set of

primers. In addition, selected INDEL SV had to be found in the genome of at least three strains.

The aim of this selection was to identify the most divergent markers among the largest

available genetic diversity of *B. prasinos*, with the expectation that some of these variations

could potentially be found and tracked in local communities (Fig.1 A). In total, 765 unique SV

with sizes ranging between 0.2 and 2 Kb were identified (Table 1). However, only 44 of them

were found in at least three strains, thus meeting our criteria as putative markers (Fig. 2).

Manual analysis of INDEL flanking sequences was performed to identify conserved

sequences that could be used to design primers for PCR amplification of INDELS. Four

markers were selected for experimental validation of INDELs by PCR (Fig. 3) (See

Supplementary Appendix 2, Additional File 3).

Figure 3 : Diversity markers in collection strains. PCR amplification of diversity markers (A) TOC1 promoter, (B) Flavodoxin-like, (C) yrdC promoter, (D) Zinc Finger, (E) TIMa. Variations in amplification efficiency are due to mismatches between primer sets and genomes.

The "TOC1 promoter" marker, located on chromosome 17 (RCC1105: NC_023992.1:308225..308959) corresponds to an INDEL variation in the intergenic region upstream of gene Bathy17g01510, a homolog of circadian clock gene TOC1. This diversity marker is bi-allelic with amplified fragments of 700 or 2,200 bp in size (Fig. 3A).

The "Flavodoxin-like" marker, located on chromosome 3 (RCC1105: NC_024006.1:379834..380504) corresponds to an INDEL variation in the gene Bathy03g02080 which encodes a protein containing amino stretches repeats of various lengths. This marker shows 5 alleles producing amplified fragment lengths of 600, 800, 1000, 1,200 and 1,400 bp (Fig. 3B).

The "yrdC promoter" marker, located on chromosome 1 (RCC1105: NC_024008.1:800915..801320) corresponds to an INDEL variation in the promoter region of a yrdC domain-containing protein of unknown function (Bathy01g04300). In the tested strains, this diversity marker is tri-allelic and shows amplified fragments of 200, 400 or 1,400 bp.

Amplification efficiency was lower in strains RCC1615, RCC685 and RCC4752 and no

amplification was obtained for strain RCC5417 (Fig. 3C).

Finally, the "Zinc finger" marker, located on chromosome 15 (RCC1105:

NC_023994.1:446402..447132), corresponds to variation in the number of Zinc finger repeats

encoded in the gene Bathy15g02320. This marker is biallelic and shows amplified fragments of

730 or 810 bp. No amplification was obtained for strain RCC5417 and RCC1868 (Fig.3 D).

The best sets of primers designed for markers "yrdC promoter" and "Zinc finger" show

differences in amplification efficiencies between strains due to the lack of conservation in PCR

primer sites in these strain genomes. However, the sequences of all amplified INDELs were

verified by sequencing and correspond to the expected amplification.

A fifth additional marker, named "TIMa", was designed to amplify a 530 bp fragment. This

marker was selected because it is located in the BOC region of chromosome 14 at the border

between a conserved region and an outlier region putatively involved in the definition of mating

types in other *Bathycocacceae* (RCC1105: NC_023995.1:564926..565456) [19]. As expected

this marker was amplified in the 6 genomes which would correspond to the same putative

mating type (Fig 3E).

Isolation and identification of *Bathycoccus prasinos* Multi Loci

Genotypes in the Banyuls bay

Surface water was collected weekly at the SOLA buoy in Banyuls bay from December the 3rd,

2018 to March the 19th, 2019. During this period, the sea temperature ranged between 15.87°C

in December and 10.68°C in February. We implemented a protocol to isolate *B. prasinos* cells

based on filtration through 1.2 μm pore-size filters to remove larger cells of nanoplankton. This

pore size allows the passage of *B. prasinos* cells (which size is estimated to be of 1.5 μm) and

most importantly eliminates the potential larger predators. After a period of acclimation of two

weeks in natural sea water supplemented with nitrate, phosphate and vitamins B1 and B12,

cells were isolated by plating in low melting agarose supplemented sea water. Out of more

than 400 colonies, only light green-yellow coloured colonies were picked and sub-cultured

subsequently. In order to identify putative *B. prasinos* strains, amplification of a fragment of the

LOV-HK gene (Bathy10g02360) was performed. These primers did not amplify LOV-HK

sequences in other *Mamiellales* (*Ostreococcus* or *Micromonas*). The identity of these clones

was further confirmed by ribotyping (amplification of a 2kb ribosomal DNA fragment followed

by sequencing) (See Supplementary Appendix 1, Additional File 3). In total, 55 *B. prasinos*

isolates were recovered for nine sampling dates (See Supplementary Table 2, Additional File

1).

The five markers, including the four diversity markers and the TIMa marker were used in

combination to distinguish the different isolates of *B. prasinos* into Multi Loci Genotypes

(MLG). Eight MLG were identified in Banyuls' strains, MLG1 and MLG2 being dominant with 30

and 15 isolated strains respectively (Table 2). These two dominant MLG were distinguished

only by the TIMa marker, which was amplified only in MLG1. For yrdC promoter marker, the

200 bp allele was dominant with an amplification in 54 isolates. The 1,400 bp allele, identified

in worldwide strain, was not detected in the Banyuls bay strain. For the TOC1 promoter

marker, the 1,200 bp allele was dominant with an amplification in 51 isolates. For the

Flavodoxin-like marker, the 1,200 bp allele was dominant with an amplification in 49 isolates.

From the 6 remaining isolates, 3 alleles could be identified, including a new 1,600 bp allele;

however the 1,000 bp allele was not detected. Finally, for the Zinc finger marker, the dominant

allele was the 730 bp one, amplified in 52 isolates.

Table 2: Marker characterisation of multi-loci genotypes in *B. prasinos* strains isolated from
Banyuls bay. Marker values correspond to amplified sizes in base pairs.
333

Marker

	yrdC	TOC1	Flavodoxin	Zinc finger	TIMa	Number of isolates
MLG1	200	2200	1200	730	530	30
MLG2	200	2200	1200	730	n.d.	15
MLG3	200	2200	800	730	530	4
MLG4	200	700	1200	730	530	2
MLG5	200	2200	1200	820	n.d.	1
MLG6	200	700	1200	820	n.d.	1
MLG7	400	700	1400	820	530	1
MLG8	200	2200	1600	730	530	1

n.d. : Not Detected

The genotypic diversity observed by the MLG classification was also correlated with biological

and physiological diversities of the strains. Indeed, by monitoring the growth response of

representative strains under different temperature and light conditions corresponding to those

observed at different parts of the bloom in the Banyuls bay, we showed clear differences in

growth rate between MLG and conditions. Specifically, there were notable variations between

strains from MLG1/MLG2 (isolated in December/January) and other tested MLG (isolated in

February) (See Supplementary Figure 2, Additional File 2).

Diversity of INDEL markers

Seven isolates corresponding to 6 MLG were selected for whole genome sequencing and

assembly following the methodology described in Figure 1A. The genome assemblies

revealed that the two MLG, which failed to amplify the TIMa marker, have a different haplotype

of the chromosome 14 outlier region. A complementary marker called "TIMb" was designed

from these genome assemblies, corresponding to a 1,300 bp amplified fragment. TIMb was

detected in isolates for which TIMa was not amplified (See Supplementary Figure 3, Additional

File 2).

Figure 4 : Worldwide and local diversity in marker alleles and variant loci
(A) Sizes and frequencies in marker alleles within worldwide strains (n=6) and local strains from the
Banyuls Bay (n=55). (B) Specific and shared variant loci (200 < size < 2000 bp) between at least one
worldwide sequenced strains (n=6) and one sequenced strains from the Banyuls Bay(n=7). (C)
Distribution frequency of the 79 shared variant loci (200 < size < 2000 bp) in the 13 sequenced
strains.

Comparative analysis of the polymorphism in our markers between worldwide strains and

newly isolated strains revealed that all five markers were at least biallelic in the Banyuls local

population (Fig. 4A). Furthermore, new alleles were identified through the genotyping of

Banyuls bay isolates for Flavodoxin-like and TIMa/b markers, thus extending the diversity

observed of our markers. The pipeline of INDEL identification was applied to the seven strains

sequenced from the Banyuls bay (Fig. 1A). This led to the detection of 211 unique INDELs of

sizes ranging between 0.2 and 2 Kb (Fig. 4B). Within these loci, 79 corresponding to 37% of

366 loci diversity in the Banyuls bay, were shared with worldwide strains (Fig. 4B). On average,

367 each variant locus was found in 5.6 strains across the two populations (Fig. 4C).

Determination of allelic variants in environmental samples

The identification of dominant MLG during the 2018/2019 bloom raises the question of their

yearly or occasional prevalence in the Banyuls bay. However, isolating strains is highly time

consuming and not appropriate to follow population dynamics at higher frequency. To

overcome this, our diversity markers were directly tracked on environmental DNA samples

during 3 successive blooms from 2018 to 2021. Five litres of seawater were sampled once a

week and filtered between 3 and 0.8 μm. DNA extracted from 0.8 μm filters was used for PCR

analysis of presence/absence variations in alleles of yrdC promoter, TOC1 promoter and

TIMa/TIMb markers (Table 3). Flavodoxin-like and Zinc finger markers could not be used on

complex environmental DNA samples due to the presence of high background.

The initiation of the 2019/2020 bloom was defined by the exclusive presence of the 200 bp

allele of yrdC promoter marker from the third week of November to the third week of

December. This was followed from January to March by the unambiguous detection of both the

200 bp and the 400 bp alleles. During this bloom, TIMa and TIMb markers were both amplified

in all samples, with the exception of the second week of March, during which only the TIMa

marker was detected.

Table 3: Marker alleles detection in environmental DNA samples from successives *B. prasinos* b
looms in the Banyuls bay. YrdC promoter (A : 200bp, B : 400bp), TOC1 promoter (A : 2200bp, B:
700bp), TIM a/b (a : 530 bp, b : 1300bp). Missing datapoints are represented by empty cells.
387

	Oct				Nov				Dec				Jan				Feb				Mar				Apr	
	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2
yrdC promoter																										
2018 - 2019																A,B	A,B	A,B	A,B	A,B	A	A,B	A,B	A,B		
2019 - 2020							A	A	A	A	A		A,B	A,B		A,B	A,B	A,B	A,B	A,B	A,B	A,B				
2020 - 2021	A,B	A,B		A,B	n.d.		n.d.		A,B	A,B	A,B		A,B		A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B
TOC1 promoter																										
2018 - 2019																A,B	A,B	A,B	A,B	A,B		A,B	A,B	A,B		
2019 - 2020								A,B	A,B	A,B	A,B		A,B	A,B		A,B	B	A,B	A,B	A,B	A,B	A,B				
2020 - 2021	A,B	A,B		n.d.	n.d.		n.d.		A,B	A,B	A,B		A,B		A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B	A,B
TIM a/b																										
2018 - 2019																a,b	a,b	a,b	a,b	a,b	a,b	a,b	a,b	a,b		
2019 - 2020							a,b	a,b	a,b	a,b	a,b		a,b	a,b		a,b	a,b	a,b	a,b	a,b	a,b	a				
2020 - 2021	a	a		a	n.d.		n.d.		a	a,b	a		a,b		a,b	a,b	a,b	a,b	a,b	a,b	a,b	a,b	a,b	a,b	a,b	a,b

n.d. : Not Detected

Several distinctions can be done with the 2020/2021 bloom, the first one being that no

amplification was detected in November samples, an observation consistent with the overall

decrease in picophytoplankton abundance (See Supplementary Figure 4, Additional File 2).

For yrdC promoter marker, the 400 bp allele was observed in October and December, hence

between one and three months earlier than in the 2019/2020 bloom. In addition, only TIMa was

amplified in October and December 2020. Both alleles of the TOC1 promoter marker were

amplified in nearly all samples where *B. prasinos* could be detected and were therefore not

informative in this context (Table 3). From the study of these successive blooms, we were able

to conclude that their onset differed in both chronology and population diversity. Furthermore,

we demonstrated that population diversity changes both within and between blooms at a given

site and that diversity markers, designed from the worldwide diversity, could be used to

successfully follow these local population dynamics.

Discussion

We aimed to develop a time and cost effective method to produce a biological resource and

characterise the intraspecific diversity of the marine microalga *B. prasinos*. In recent years, the

cost of high-throughput sequencing has fallen sharply, favouring blind sequencing approaches

without prior genotyping [19]. However, the isolation of new strains from complex microbial

communities is time consuming, requiring multiple steps from sample isolation in remote areas

of the ocean, to incubations, shipment and isolation by subculturing on agarose plates.

Moreover, the incubation step can lead to the amplification of strains that, without genotyping,

could be sequenced multiple times, while missing rare genotypes that contribute to the fitness

of a species under changing environmental conditions. Given the bottleneck steps of strain

isolation and whole genome sequencing in complex microbial communities, we turned to PCR

amplification as a potentially effective approach for (i) preliminary screening of environmental

samples to retain and incubate only those containing *B. prasinos* cells, (ii) identifying incubates

containing *B. prasinos* for subsequent isolation of strains, (iii) genotyping and identifying the

main MLGs prior to sequencing. These first steps of screening by PCR could be potentially

performed during oceanographic cruise to guide and optimise sample selection and culturing

on boats. Remarkably, our protocol proved efficient and selective for *B. prasinos* isolation

since *Ostreococcus* and *Micromonas* strains were not obtained even though *Micromonas* is as

abundant as *B. prasinos* in the Banyuls bay in winter [15]. This is all the more important as flow

cytometry cannot distinguish the *Bathycoccus* genus from other *Mamiellales*.

Design of diversity markers from INDELs structural variations

While PCR, combined with DNA fragment size estimation by gel electrophoresis, has already

been used to follow intraspecific diversity [12], this approach requires the specific amplification

of size-varying loci. This leads to challenges such as identifying putative marker loci that vary

in size, are sufficiently conserved to be representative of subpopulations and can be found in

conserved regions for which PCR primers can be designed. To address this, we explored the

utility of INDEL variations within coding sequences, which offer higher stability in clonal

populations compared to SNP and microsatellite markers, and are therefore good candidates

for the design of size markers to track intraspecific diversity [36,37]. Furthermore, information

on such INDELs variation and on repeated/low complexity sequences are now becoming

easily available through the leveraging of the recent ONT and PacBio long read sequencing

technologies, providing access to structural variations and marker designs previously

overlooked [31,38,39].

By long read sequencing a selection of only six geographically distinct strains available in

culture collection, we were able to obtain a first picture of the genetic diversity of *B. prasinos.* T

he implemented genome assembly pipeline provided the draft genomes required for

comparison and identification of structural variations. This resulted in the initial identification of

683 INDEL variation candidates, including 44 meeting our criteria for the design of diversity

markers. While each candidate marker must be validated experimentally, this demonstrates

the effectiveness of long reads whole genome sequencing for the identification of diversity

marker loci using a limited biological resource. Four loci were selected for PCR validation,

resulting in the diversity markers Flavodoxin-like, TOC1 promoter, yrdC promoter and Zinc

finger. All markers were successfully amplified in a majority of strains even though a lack of

conservation in PCR primer sites resulted in uneven amplification signals for yrdC promoter

and Zinc finger markers. As such, nucleotide diversity remains a limiting factor for the design of

universal markers when considering phylogenetically distant strains. Nonetheless, the pipeline

developed in this study, by systematically mapping all INDEL and iteratively integrating them in

the database, provides an extensive resource to identify novel markers (Fig. 1a; Fig. 4).

As expected, fewer alleles were detected in the local Mediterranean population for the

designed markers, but genetic diversity was still observed for all of them. Since all markers

correspond to structural variations either within coding sequences or regulatory sequences,

this raises the question of whether they play a role in local adaptation. For instance, in the case

of the central circadian clock TOC1, the variation was found in the promoter which contains a

key *cis*-regulatory Evening element that is essential for the circadian regulation of TOC1 [40].

However, no clear pattern of allele distribution was identified neither in the worldwide strains

nor in the sampled Banyuls population. The potential role of these INDELs in adaptation would

therefore require additional experimental investigation.

Genotyping and phenotyping in local populations of *B. prasinos*

The five diversity markers developed in this study were tested in combination on 55 freshly

isolated Mediterranean strains of *B. prasinos*. This resulted in their classification into eight

multi-loci genotypes. This highlights, firstly, the efficiency of our protocol for isolating *B.*

prasinos strains from complex seawater samples and its effectiveness in preserving genotypic

diversity. This also includes the TIMa/b markers which were manually designed from

assembled genomes to detect different haplotypes of the big outlier chromosome 14 (BOC) as

described by Blanc-Mathieu et al. [19] in *Ostreococcus tauri*. This marker combination allowed

us to confirm the existence of at least two distinct haplotypes of the BOC in the *B. prasinos*

subpopulation from the Banyuls bay, and thus can be used to distinguish these putative mating

types in newly isolated strains and environmental samples.

A high level of intraspecific diversity has been reported in the few studies describing the

genetic diversity of blooming phytoplankton populations [41–44]. For example, in diatoms

(although not directly comparable as microsatellite markers have a higher mutation rate than

other regions of the genome), more than 600 individuals were genotyped using microsatellite

markers, and it was estimated that the blooming population was comprised of at least 2400

different genotypes [41]. Therefore, the observation of eight MLG in *B. prasinos* mediterranean

population is most likely an underestimation due to the low number of markers used and their

limited polymorphism. However, further improvement in the accuracy of our marker set can be

achieved in an iterative way: through the identification of new size alleles (as exemplified by

the 1,600 bp allele of the flavodoxin-like marker identified in the Banyuls population), or by the

sequencing and integration of newly isolated strains distinguished by our original set of primer,

further increasing the number of candidate loci. In this study, the integration of 7 newly

sequenced genomes from the Banyuls bay nearly doubled the number of putative universal

markers and provided population specific variants that could be used in local subpopulations

tracking.

The first physiological characterization of selected strains revealed distinct growth responses,

the strains isolated in December and January showing reduced growth under lower

temperature (MLG1 and MLG2) and short photoperiod (MLG1) which suggest that these

strains may correspond to seasonal ecotypes. Before concluding whether the different MLG

classes correspond to different ecotypes, it will be necessary to characterise the physiological

response of more strains, in particular those belonging to each of the dominant MLG1 and

MLG2 classes.

Tracking intraspecific diversity directly in environmental samples

Isolation of *B. prasinos* strains is too laborious and costly to be performed systematically on a large scale. For this reason, we attempted to assess intraspecific diversity in environmental samples during three consecutives annual blooms of *B. prasinos* in the Banyuls bay. Although flavodoxin-like was the most resolutive marker in isolated strains, PCR on environmental DNA yielded a smear on electrophoresis gel, possibly due to high polymorphism. Three diversity markers, TIMa/b, yrdC and TOC1 promoter, were suitable for amplifying environmental DNA, highlighting local population dynamics within and between bloom events. While this approach based on PCR amplification provides only qualitative data at this stage, approaches based on quantitative PCR could be developed to track the dynamics of genetic diversity in *B. prasinos* blooms directly from environmental samples [45]. As such, by accessing a level of structural diversity hardly obtainable through metagenomic short read mapping, it is complementary with approaches of metagenomic characterisation through SNP, as proposed by Da silva et al. [46] .

Variations in interannual presence/absence of alleles raise the question of the nature of the highly reproducible yearly occurrence of *B. prasinos* in the Banyuls bay. Seasonal blooms may result either from re-activation of "dormant/survivor" cells from the water column (whose genetic fingerprint will determine the genetic profile of the next bloom) or by yearly *de novo* seeding by cells carried by the north Mediterranean current along the Gulf of Lion. At first glance, our preliminary qualitative results are in favour of the introduction of a new population rather than "resurrection" of cells from the previous bloom, since the allelic composition is distinct between the end of a bloom and the onset of the next (Table 3). By monitoring the temporal population structures of the dinoflagellate *Alexandrium minutum* in two estuaries in France, Dia et al. [44] showed that interannual genetic differentiation was greater than intra-bloom differentiation. Alternation of genotypes/populations has also been observed with diatoms in the dominance of one of the two sympatric populations of *Pseudonitzschia*

multistriata, which could be due either to environmental factors favouring one population over

the other or intrinsic factors coupled to the obligate sexual life cycle of *P. multistriata* [47]. Thus

the observed fluctuations in allele frequencies could equally be the result of new inoculum from

currents or sexual reproduction. Even though sexual reproduction has not been demonstrated

in *B. prasinos*, there is genomic evidence that it may occur [48], and potential mating types

were identified in our study (TIMa/TIMb markers). Sexual recombination generates new

combinations of alleles, whereas clonality favours the spread of the fittest genotype through

the entire population [44]. Erdner et al. [49] propose for *A. fundyense* that mitosis is the primary

mode of multiplication during blooms, whereas mating is triggered presumably in response to

unfavourable conditions at the end of blooms, with vegetative cells not overwintering in the

water column. Knowing that (i) *B. prasinos* blooms are followed by severe bottlenecks between

one bloom and the next [15], (ii) allele occurrences were different between the end of one

bloom and the onset of the next, and (iii) structural markers are very stable in mitotic dividing

cells, the hypothesis of rare vegetative cells remaining in the water column between the

blooms is unlikely, except if those remaining cells were produced by sexual reproduction. The

alternative hypothesis of new strains brought by current is however still equally probable. To

test these hypotheses, more quantitative approaches based on real time quantitative PCR, or

direct mapping of INDEL markers on the 7 year metagenomic time series in the Banyuls bay

could potentially be used [50].

541 **Conclusions**

In this paper, we describe the development of a new type of diversity markers based on INDEL

structural variants obtained by long read sequencing. After validation on freshly isolated

individuals, markers were used *in situ* on environmental samples. Whole genome sequencing

of the new MLG refined a marker resource that could be further used to guide strain isolation

and genotype new strains from the world ocean. In addition, the assessment of the

physiological performances of genotyped strains suggest the presence of seasonal ecotypes

in the Banyuls bay. Finally, PCR on environmental samples provided insight into the genetic

diversity of *B. prasinos* seasonal blooms.

This INDEL marker resource represents a new tool for grasping the maximum diversity of

newly isolated *B. prasinos* with the long term goal of identifying putative molecular

mechanisms involved in adaptation to environmental niches in this cosmopolitan genus. The

marker resource could also potentially be used to check culture purity or to follow specific

genotypes in competition experiments in culture or in microcosms. The developed pipeline

aimed at optimising sequencing efforts and building a collection of sequenced genomes

representative of *B. prasinos* intraspecific diversity. This pipeline could easily be applied to

other cosmopolitan and cultivable planktonic species that are available in collection, whether

they are related to *B. prasinos* (e.g. *Ostreococcus* RCC809, *Micromonas commoda*,

Micromonas pusilla) or share similar genome features.

List of abbreviations

ASW : Artificial Sea Water

BOC : Big Outlier Chromosome

INDEL : Insertion/Deletion

ONT : Oxford Nanopore Tech

PCR : Polymerase Chain Reaction

RCC : Roscoff Culture Collection

SV : Structural Variation

Declarations

Availability of data and materials

Whole genome sequences and basecalled reads of the strains sequenced in this study as well as variant calling outputs are available at Zenodo https://doi.org/10.5281/zenodo.11395682. Strains isolated in Banyuls are available on request. All codes are available at https://forge.ird.fr/diade/genomecodes under the GPLv3 licence.

Funding

The work was financed by an internal LOMIC Microproject to MD and the ANR Clima-Clock, ANR-20-CE20-0024 to FYB.

Author contribution

MD, FYB and FS conceived the work and acquired funding. MD extracted high molecular weight genomic DNA. CM and MD performed the ONT sequencing. LD and FS performed bioinformatic analysis. MD designed and validated the diversity markers. PS analysed the seawater samples by flow cytometry. MD and VV isolated Banyuls strains during the winter bloom, genotyped by MD and JCL. MD and JCL determined the diversity of seawater samples. MD, LD and FYB wrote the article. JCL and FS took part in the critical reviews. All authors approved the final version for submission.

Acknowledgments

We are grateful to the captain and the crew of the RV 'Nereis II' for their help in acquiring the samples. We thank the Roscoff Culture Collection for providing access to collected strains. Additional ONT were performed with the help of Christel Llauro and Marie Mirouze LGDP. The authors acknowledge the ISO 9001 certified IRD itrop HPC (member of the South Green Platform) at IRD Montpellier for providing HPC resources that have contributed to the research

591 results reported within this paper (URL: https://bioinfo.ird.fr/ - http://www.southgreen.fr). We

592 thank Thomas Roscoe for critically reading the manuscript.

593 References

594 [1] Li WKW, Rao DVS, Harrison WG, Smith JC, Cullen JJ, Irwin B, et al. Autotrophic Picoplankton
595   in the Tropical Ocean. Science 1983;219:292–5. https://doi.org/10.1126/science.219.4582.292.
596
597 [2] Violle C, Enquist BJ, McGill BJ, Jiang L, Albert CH, Hulshof C, et al. The return of the variance:
598   intraspecific variability in community ecology. Trends in Ecology & Evolution 2012;27:244–52.
599   https://doi.org/10.1016/j.tree.2011.11.014.
600
601 [3] Raffard A, Santoul F, Cucherousset J, Blanchet S. The community and ecosystem
602   consequences of intraspecific diversity: a meta-analysis. Biological Reviews 2019;94:648–61.
603   https://doi.org/10.1111/brv.12472.
604
605 [4] Des Roches S, Post DM, Turley NE, Bailey JK, Hendry AP, Kinnison MT, et al. The ecological
606   importance of intraspecific variation. Nat Ecol Evol 2018;2:57–64.
607   https://doi.org/10.1038/s41559-017-0402-5.
608
609 [5] Godhe A, Rynearson T. The role of intraspecific variation in the ecological and evolutionary
610   success of diatoms in changing environments. Phil Trans R Soc B 2017;372:20160399.
611   https://doi.org/10.1098/rstb.2016.0399.
612
613 [6] Rynearson TA, Bishop IW, Collins S. The Population Genetics and Evolutionary Potential of
614   Diatoms. In: Falciatore A, Mock T, editors. The Molecular Life of Diatoms, Cham: Springer
615   International Publishing; 2022, p. 29–57. https://doi.org/10.1007/978-3-030-92499-7_2.
616
617 [7] Srivastava S, Avvaru AK, Sowpati DT, Mishra RK. Patterns of microsatellite distribution across
618   eukaryotic genomes. BMC Genomics 2019;20:153. https://doi.org/10.1186/s12864-019-5516-5.
619
620 [8] Wheeler GL, Dorman HE, Buchanan A, Challagundla L, Wallace LE. A review of the prevalence,
621   utility, and caveats of using chloroplast simple sequence repeats for studies of plant biology.
622   Applications in Plant Sciences 2014;2:1400059. https://doi.org/10.3732/apps.1400059.
623
624 [9] Galtier N, Nabholz B, Glémin S, Hurst GDD. Mitochondrial DNA as a marker of molecular
625   diversity: a reappraisal. Mol Ecol 2009;18:4541–50. https://doi.org/10.1111/j.1365-
626   294X.2009.04380.x.
627
628 [10] Lewis RJ, Jensen SI, DeNicola DM, Miller VI, Hoagland KD, Ernst SG. Genetic variation in the
629   diatom Fragilaria capucina (Fragilariaceae) along a latitudinal gradient across North America. Pl
630   Syst Evol 1997;204:99–108. https://doi.org/10.1007/BF00982534.
631
632 [11] Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA. Harnessing the power of RADseq
633   for ecological and evolutionary genomics. Nat Rev Genet 2016;17:81–92.
634   https://doi.org/10.1038/nrg.2015.28.
635
636 [12] Rengefors K, Kremp A, Reusch TBH, Wood AM. Genetic diversity and evolution in eukaryotic
637   phytoplankton: revelations from population genetic studies. Journal of Plankton Research
638   2017;39:165–79. https://doi.org/10.1093/plankt/fbw098.
639
640 [13] Joli N, Monier A, Logares R, Lovejoy C. Seasonal patterns in Arctic prasinophytes and inferred
641   ecology of Bathycoccus unveiled in an Arctic winter metagenome. ISME J 2017;11:1372–85.
642   https://doi.org/10.1038/ismej.2017.7.
643

644 [14] Tragin M, Vaulot D. Novel diversity within marine Mamiellophyceae (Chlorophyta) unveiled by
645   metabarcoding. Sci Rep 2019;9:5190. https://doi.org/10.1038/s41598-019-41680-6.
646
647 [15] Lambert S, Tragin M, Lozano J-C, Ghiglione J-F, Vaulot D, Bouget F-Y, et al. Rhythmicity of
648   coastal marine picoeukaryotes, bacteria and archaea despite irregular environmental
649   perturbations. ISME J 2019;13:388–401. https://doi.org/10.1038/s41396-018-0281-z.
650
651 [16] Leconte J, Benites LF, Vannier T, Wincker P, Piganeau G, Jaillon O. Genome Resolved
652   Biogeography of Mamiellales. Genes 2020;11:66. https://doi.org/10.3390/genes11010066.
653
654 [17] Richter DJ, Watteaux R, Vannier T, Leconte J, Frémont P, Reygondeau G, et al. Genomic
655   evidence for global ocean plankton biogeography shaped by large-scale current systems. eLife
656   2022;11:e78129. https://doi.org/10.7554/eLife.78129.
657
658 [18] Simmons MP, Sudek S, Monier A, Limardo AJ, Jimenez V, Perle CR, et al. Abundance and
659   Biogeography of Picoprasinophyte Ecotypes and Other Phytoplankton in the Eastern North
660   Pacific Ocean. Appl Environ Microbiol 2016;82:1693–705. https://doi.org/10.1128/AEM.02730-
661   15.
662
663 [19] Blanc-Mathieu R, Krasovec M, Hebrard M, Yau S, Desgranges E, Martin J, et al. Population
664   genomics of picophytoplankton unveils novel chromosome hypervariability. Sci Adv
665   2017;3:e1700239. https://doi.org/10.1126/sciadv.1700239.
666
667 [20] De Vargas C, Audic S, Henry N, Decelle J, Mahé F, Logares R, et al. Eukaryotic plankton
668   diversity in the sunlit ocean. Science 2015;348:1261605.
669   https://doi.org/10.1126/science.1261605.
670
671 [21] Vannier T, Leconte J, Seeleuthner Y, Mondy S, Pelletier E, Aury J-M, et al. Survey of the green
672   picoalga Bathycoccus genomes in the global ocean. Scientific Reports 2016;6:37900–37900.
673   https://doi.org/10.1038/srep37900.
674
675 [22] Moreau H, Verhelst B, Couloux A, Derelle E, Rombauts S, Grimsley N, et al. Gene
676   functionalities and genome structure in Bathycoccus prasinos reflect cellular specializations at
677   the base of the green lineage. Genome Biol 2012;13:R74. https://doi.org/10.1186/gb-2012-13-8-
678   r74.
679
680 [23] Limardo AJ, Sudek S, Choi CJ, Poirier C, Rii YM, Blum M, et al. Quantitative biogeography of
681   picoprasinophytes establishes ecotype distributions and significant contributions to marine
682   phytoplankton. Environmental Microbiology 2017;19:3219–34. https://doi.org/10.1111/1462-
683   2920.13812.
684
685 [24] Bachy C, Yung CCM, Needham DM, Gazitúa MC, Roux S, Limardo AJ, et al. Viruses infecting a
686   warm water picoeukaryote shed light on spatial co-occurrence dynamics of marine viruses and
687   their hosts. ISME J 2021;15:3129–47. https://doi.org/10.1038/s41396-021-00989-9.
688
689 [25] Lambert S, Lozano J-C, Bouget F-Y, Galand PE. Seasonal marine microorganisms change
690   neighbours under contrasting environmental conditions. Environmental Microbiology
691   2021;23:2592–604. https://doi.org/10.1111/1462-2920.15482.
692
693 [26] Debladis E, Llauro C, Carpentier M-C, Mirouze M, Panaud O. Detection of active transposable
694   elements in Arabidopsis thaliana using Oxford Nanopore Sequencing technology. BMC
695   Genomics 2017;18:537. https://doi.org/10.1186/s12864-017-3753-z.
696
697 [27] De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. NanoPack: visualizing and
698   processing long-read sequencing data. Bioinformatics 2018;34:2666–9.
699   https://doi.org/10.1093/bioinformatics/bty149.
700
701 [28] Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat
702   graphs. Nat Biotechnol 2019;37:540–6. https://doi.org/10.1038/s41587-019-0072-8.
703

[29] Vaser R, Sović I, Nagarajan N, Šikić M. Fast and accurate de novo genome assembly from long uncorrected reads. Genome Res 2017;27:737–46. https://doi.org/10.1101/gr.214270.116.

[30] Li H. New strategies to improve minimap2 alignment accuracy. Bioinformatics 2021;37:4572–4. https://doi.org/10.1093/bioinformatics/btab705.

[31] Alonge M, Soyk S, Ramakrishnan S, Wang X, Goodwin S, Sedlazeck FJ, et al. RaGOO: fast and accurate reference-guided scaffolding of draft genomes. Genome Biol 2019;20:224. https://doi.org/10.1186/s13059-019-1829-6.

[32] Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. Versatile genome assembly evaluation with QUAST-LG. Bioinformatics 2018;34:i142–50. https://doi.org/10.1093/bioinformatics/bty266.

[33] Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. MUMmer4: A fast and versatile genome alignment system. PLoS Comput Biol 2018;14:e1005944. https://doi.org/10.1371/journal.pcbi.1005944.

[34] Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of SAMtools and BCFtools. GigaScience 2021;10:giab008. https://doi.org/10.1093/gigascience/giab008.

[35] Guyon J-B, Vergé V, Schatt P, Lozano J-C, Liennard M, Bouget F-Y. Comparative Analysis of Culture Conditions for the Optimization of Carotenoid Production in Several Strains of the Picoeukaryote Ostreococcus. Marine Drugs 2018;16:76. https://doi.org/10.3390/md16030076.

[36] Ruggiero MV, D'Alelio D, Ferrante MI, Santoro M, Vitale L, Procaccini G, et al. Clonal expansion behind a marine diatom bloom. The ISME Journal 2018;12:463–72. https://doi.org/10.1038/ismej.2017.181.

[37] Mérot C, Oomen RA, Tigano A, Wellenreuther M. A Roadmap for Understanding the Evolutionary Significance of Structural Genomic Variation. Trends in Ecology & Evolution 2020;35:561–72. https://doi.org/10.1016/j.tree.2020.03.002.

[38] Mantere T, Kersten S, Hoischen A. Long-Read Sequencing Emerging in Medical Genetics. Front Genet 2019;10:426. https://doi.org/10.3389/fgene.2019.00426.

[39] Wellenreuther M, Mérot C, Berdan E, Bernatchez L. Going beyond SNPs: The role of structural genomic variants in adaptive evolution and species diversification. Mol Ecol 2019;28:1203–9. https://doi.org/10.1111/mec.15066.

[40] Corellou F, Schwartz C, Motta J-P, Djouani-Tahri EB, Sanchez F, Bouget F-Y. Clocks in the Green Lineage: Comparative Functional Analysis of the Circadian Architecture of the Picoeukaryote Ostreococcus. The Plant Cell 2009;21:3436–49. https://doi.org/10.1105/tpc.109.068825.

[41] Rynearson TA, Armbrust EV. Maintenance of clonal diversity during a spring bloom of the centric diatom Ditylum brightwellii. Molecular Ecology 2005;14:1631–40. https://doi.org/10.1111/j.1365-294X.2005.02526.x.

[42] Alpermann TJ, Beszteri B, John U, Tillmann U, Cembella AD. Implications of life-history transitions on the population genetic structure of the toxigenic marine dinoflagellate Alexandrium tamarense. Molecular Ecology 2009;18:2122–33. https://doi.org/10.1111/j.1365-294X.2009.04165.x.

[43] Lebret K, Kritzberg ES, Figueroa R, Rengefors K. Genetic diversity within and genetic differentiation between blooms of a microalgal species. Environmental Microbiology 2012;14:2395–404. https://doi.org/10.1111/j.1462-2920.2012.02769.x.

[44] Dia A, Guillou L, Mauger S, Bigeard E, Marie D, Valero M, et al. Spatiotemporal changes in the

764 genetic diversity of harmful algal blooms caused by the toxic dinoflagellate Alexandrium
765 minutum. Molecular Ecology 2014;23:549–60. https://doi.org/10.1111/mec.12617.
766
767 [45] Demir-Hilton E, Sudek S, Cuvelier ML, Gentemann CL, Zehr JP, Worden AZ. Global distribution
768 patterns of distinct clades of the photosynthetic picoeukaryote Ostreococcus. ISME J
769 2011;5:1095–107. https://doi.org/10.1038/ismej.2010.209.
770
771 [46] Da Silva O, Ayata S, Ser-Giacomi E, Leconte J, Pelletier E, Fauvelot C, et al. Genomic
772 differentiation of three pico-phytoplankton species in the Mediterranean Sea. Environmental
773 Microbiology 2022:1462-2920.16171. https://doi.org/10.1111/1462-2920.16171.
774
775 [47] D'Alelio D, d'Alcalà MR, Dubroca L, Diana Sarn, Zingone A, Montresor M. The time for sex: A
776 biennial life cycle in a marine planktonic diatom. Limnology and Oceanography 2010;55:106–14.
777 https://doi.org/10.4319/lo.2010.55.1.0106.
778
779 [48] Benites LF, Bucchini F, Sanchez-Brosseau S, Grimsley N, Vandepoele K, Piganeau G.
780 Evolutionary Genomics of Sex-Related Chromosomes at the Base of the Green Lineage.
781 Genome Biology and Evolution 2021;13:evab216. https://doi.org/10.1093/gbe/evab216.
782
783 [49] Erdner DL, Richlen M, McCauley LAR, Anderson DM. Diversity and Dynamics of a Widespread
784 Bloom of the Toxic Dinoflagellate Alexandrium fundyense. PLOS ONE 2011;6:e22965.
785 https://doi.org/10.1371/journal.pone.0022965.
786
787 [50] Beauvais M, Schatt P, Montiel L, Logares R, Galand PE, Bouget F-Y. Functional redundancy of
788 seasonal vitamin B12 biosynthesis pathways in coastal marine microbial communities.
789 Environmental Microbiology 2023;25:3753–70. https://doi.org/10.1111/1462-2920.16545.