Proteins with anti-apoptotic action in the hemolymph of caterpillars of the Megalopygidae family acts by maintaining the structure of the cellular cytoskeleton

Nathalia Delazeri de Carvalho[1], **Henrique Krambeck Rofatto**[1], **Karina de Senna Villar**[1], **Roberta Fiusa Magnelli**[2], **Ronaldo Zucatelli Mendonça**[1]

1 Laboratório de Parasitologia, 2 Centro Bioindustrial do Instituto Butantan, São Paulo SP.
Correspondence to Ronaldo Zucatelli Mendonça *
Laboratório de Parasitologia, Instituto Butantan
Av. Vital Brasil 1500, CEP 05503-900, São Paulo, Brazil.
Phone 55 11 2627 9420 ; Fax: +55 011 2627 9581   e-mail: zucatelli@uol.com.br

ABSTRACT

Brazil has a very large biological variety, which is an almost inexhaustible source of substances of pharmacological and biotechnological interest. Several studies have demonstrated the presence of bioactive peptides in insect hemolymph and their potential use as therapeutic agents. However, few data are available regarding molecules extracted from insects with anti-apoptotic action. The objective of this work was to identify and isolate proteins from the hemolymph of caterpillars of the *Megalopygidae* family with pharmacological and biotechnological interest. Two species of this family were studied**,** *Podalia sp and Megalopyge albicolis.* Cytotoxicity tests on Vero and Sf-9 cells revealed that the hemolymph of both caterpillars were cytotoxic only at concentrations greater than 5%v/v. In the anti-apoptotic activity assays, it was verified that the supplementation of cell cultures with only 1% of hemolymph v/v is sufficient to inhibit cell death by apoptosis induced by different inducers such as terbutyl, actinomycin D, hydrogen peroxide or even by nutrient depletion. For this study, cells were staining with trypan blue, crystal violet and fluorescent markers to cytoskeleton (actin and tubulin), mitochondria membrane electric potential (JC-1) and apoptosis marker (acridine orange and ethidium). The protein responsible for anti-apoptotic action was isolated through gel filtration chromatography, using an AKTA purifier high-resolution liquid chromatography system. The hemolymph was fractionated into 3 pools for *Podalia sp* and 6 pools for *M. abicolis*. In the antiapoptotic tests, semi purified hemolymph from both caterpillars showed anti-apoptotic effect in VERO and SF-9 cells, pre-treated with only 1% v/v of hemolymph and induced to death by different and apoptotic inductors. Was observed that the molecule with anti-apoptotic effect are present in pool 3 in both hemolymphs. This protector effect blocked and attenuated the disruption of the cytoskeleton (actin filaments), being that the protective effect also was observed on the integrity of the mitochondrial membrane of SF-9 cells pre-treated with both hemolymphs and treated with the apoptosis inducer Terbutil at concentrations of 25 to 100µM.

Key words: *Podalia sp*, *M. albicolis*, antiapoptotic action, hemolymph, bio-active peptide; mitochondrial membrane permeabilization; apoptosis; bio-prospection, Vero cells, SF-9 cells, cytoskeleton, actin filaments, tubulin.

Abbreviations: HB: Hemolymph; t-BPH: Tert-Butylhydroperoxide; PI: Propidium Iodide;   AO: acridine orange; EB: ethidium bromide; **(**DiOC6 (3), 3,3'Dihexyloxacarbocyanine Iodide;   JC-1: 3,3'E-Tetraethylbenzimidazolylcar-bocyanine Iodide;   MMP: Mitochondrial Membrane Permeabilization.

1. INTRODUCTION

Insects are believed to have emerged in the Devonian period, over 350 million years ago. During this period, insects lived and survived in almost all ecological niches on the planet, from deserts, polar regions, contaminated environments or extremely harmful to the development of life, developing protection and defense mechanisms for each adverse situation. Several studies have been carried out demonstrating the presence of pharmacologically active substances in the hemolymph of insects with antiviral action (Mendonça et al, 2023; Carvalho et al, 2022, Carmona et al, 2020; Coelho et al, 2018, Greco et al, 2009, Carmo et al, 2012) antimicrobial and antifungal action (Cantero et al, 2022, (Hayashida et al 2019), Zhu et al., 2000, Gross et al., 1998; Johns et al., 1998; Lanz-Mendoza et al., 1996; Lowenberger et al., 1999) antifreeze (Kristiansen et al, 2005) hormonal (Huberman et al., 2000; Jones et al., 1993; Kurata et al., 1994; Lin et al., 1998), enzymatic effects, among others. (Guerrero et al., 1999; Hamdaoui et al., 1998; Jiang et al., 1999; Rosenfeld and Vanderberg, 1998; Shiotsuki et al., 2000; Yamamoto et al., 1999). Some works have been published demonstrating the positive effect of hemolymph in promoting growth (Kanaya and Kobayashi, 2000; Rhee et al., 1999; Rhee and Park, 2000), production of recombinant proteins (Vieira et al, 2010, Mendonca et al., 2008; Woo et al., 1997). Kanaya and Kobayashi (2000) isolated and characterized a protein present in the hemolymph of the silkworm (Bombix mori) capable of increasing the activity of a recombinant protein (luciferase) by approximately 10,000 times. Some studies have been carried out seeking to isolate and characterize the factors involved in these effects (Nussbaumer et al., 2000; Ochanda et al., 1992; Shishikura et al., 1996; Shishikura et al., 1997) and only some studies have demonstrated the site of action of these proteins. We have verified the occurrence of cell death by apoptosis in cell cultures depleted of nutrients (Mendonca et al., 2002; Meneses-Acosta et al., 2001) or by induction of chemical agents (Souza et al., 2005). We have shown that supplementation of insect and mammalian cell cultures with *Lonomia obliqua* hemolymph can extend culture viability by preventing death by apoptosis (Maranga et al., 2003; Raffoul et al., 2005; Souza et al., 2005). Recently we have observed that this antiapoptotic effect of hemolymph is associated with the mitochondria patway, acting in the mitochondria membrane permeabilization (MMP) and with the maintenance of electrochemical potential of the mitochondria (Mendonça and Martins, 2022). These informations makes of hemolymph an interesting source of additives for cell cultures. However, studies with caterpillars of the *Megalopygidae* family are very scarce. Megalopygids (hairy caterpillars) are generally solitary, non-aggressive, from 1 cm to 8 cm in length. Its dorsal "hairs" are long bristles, fragile, silky and harmless, of varied colors that camouflage the true pointed and stinging bristles. The base of the bristle is a portion implanted in the caterpillar's skin, consisting of a single gland, filled with toxic substance. When pressed upon contact, the gland releases the poison that travels through a channel, being injected into the human skin causing injury. Some species of caterpillar are especially beautiful and attractive, which makes people, especially children, run their hand over the caterpillar. Species of Megalopygids can vary in color from gray to red, pink, yellow, black brown. Some popular names are given to these caterpillars such as caterpillar-dog, flannel caterpillar, caterpillar-kitten, hat-armor and bug-hairy.

In Brazil there are approximately 50,000 species of lepidopterans, being about 220 species in the Megalopygidae family. These caterpillar are found mostly in the neotropics regions (Cardoso and Junior 2005). Megalopygidae and the Saturniidae family *(Lonomia obliqua),* are responsible for the majority of accidents by caterpillar in Brazil. Reports of irritating properties of lepidopterans are described by Pliny the Elder and Galen during the Roman Empire (Fonseca, F, 1950).The first reports in the American continent were made by priest José de Anchieta in his Letters from São Vicente" (1569) and Marcgrave and Piso, the fathers of Natural History in Brazil, recorded cases of accidents with caterpillars in the Northeastern region, in 1658 (Rotberg A, 1990). Megalopygid venom can cause an extreme allergic reaction, resulting in rashes, inflammation, blisters, and breathing difficulties. The principal species in this family are *Podalia sp* and *Megalopyge albicolis.*   Studies aiming at the search for substances with pharmacological or biotechnological action are practically non-existent. We recently carried out a study with these two caterpillars in order to identify substances with antiviral action. In this work we identified substances with potent antiviral action against Measles virus,   Influenza   virus (H1N1), Herpes Simplex Virus and picornavirus (Carvalho et al, 2022). Studies on cell death are of great importance. When the cell dies, it guarantees the regulation of the body's homeostasis and important functions, such as organ remodeling and control of the immunity system. The entire process of controlling cell death, known as apoptosis, is regulated by a network of proteins. Knowledge of cell death mechanisms can lead to important strategies in the control of various physiological processes. We previously identified substances with potent anti-apototic action in the hemolymph of caterpillars of the Saturnidade family (Souza et al., 2005) and *Megalopyge*   (Mazzoni et al, 2014). So, due to the importance of studies on cell death,   the objective of this research is the identification, isolation and purification of substances with anti-apoptotic action in the hemolymph of caterpillars of the Megalopygidae family.

3. MATERIALS AND METHODS

3.1 Cell culture

3.1.1 Cell line, culture medium and cell culture

VERO cells (Cercophitecus aethiops African green monkey kidney fibroblast cells), were cultured at 37ºC in T-flasks containing 5 ml of Leibovitz medium (L-15) supplemented with 10% fetal bovine serum. Cell cultures were also performed in 6, 12, 24 or 96 well microplates.

Sf-9 cells (spodoptera frugiperda pupal ovary strain) were cultured at 29°C and 120 rpm in Schott flasks with a working volume of 13 mL in SF-900 medium or in microplates.

Cell growth and morphology were monitored using an Olympus CK2 inverted optical microscope at 100, 200 or 400x magnification.

3.2 Isolation and purification of proteins

3.2.1 Hemolymph

The hemolymph used in this study was obtained from caterpillars of the Megalopygidae family (*Podalia sp and M. albicolis*), by hemolymph leakage after cutting the caterpillar's pseudofeet. The hemolymph thus collected was stored at –20°C. Before use, the hemolymph was thawed, centrifuged at 1,000 g for 10 minutes and filtered through a 0.22 μm sterilizing membrane. The material was kept at -4°C and used as a supplement to the cultures at a concentration of 1% to 2% (v/v).

3.2.2 Hemolymph fractionation by chromatography

For the fractionation of the hemolymph of *Podalia sp and M. albicolis*, a high-resolution liquid chromatography system AKTA purifier (Amersham Pharmacia Biotech) was used, consisting of two pumps, ultraviolet light reader, programmer – Unicorn 4.0 software and fraction collector Frac 900 (Amersham Pharmacia Biotech) coupled. 0.5 mL sample of hemolymph was subjected to gel filtration chromatography on "Superdex 75" columns in Tris buffer (20mM) and Nacl (150mM) with a flow rate of 0.5 mL/min. The fractions obtained were pooled.

3.2.3 Determination of protein concentration

The protein concentration present in the hemolymph and in its fractions was determined by the NanoDrop 2000. The absorbance reading was performed at 280 nm.

3.2.4 SDS-PAGE polyacrylamide gel electrophoresis

Electrophoresis of polyacrylamide gels, for visualization of the protein bands of the total hemolymph and its fractions, were carried out according to (Laemmli, 1970), under denaturing conditions, in the presence of sodium dodecyl sulfate (SDS).

3.3 Cytotoxicity e Genotoxicity of hemolymph

3.3.1 Cytotoxicity of hemolymph.

To evaluate cytotoxicity, 0.5; 1; 2 or 5% of total hemolymph or its fractions obtained by chromatography was applied to cell culture. The morphology of the cultures was monitored through daily observation in an inverted optical microscope at 100x and 200x magnification. At the end of the cultivation, the wells were stained with crystal violet (0.25%) and photographed, determining the morphological changes that occurred in the cells.

The cytotoxicity on Vero cells of both hemolymph by apoptosis inducers cells were also evaluated by the MTT method, as described by Mosmann, 1983). For this, Vero cells were treated with 2% v/v of hemolymph or Tert-butyl at concentrations 0,5 and 1mM for 18 hours. The medium was removed and 100μl of a 0.5 mg/ml MTT solution was added and the plate was again incubated at 37°C for 4 hours. Next, the supernatant was removed and 100μl of DMSO was added. The formazan crystals were then solubilized by hand stirring. Absorbance was determined in a microplate reader at 540nm. The ability of cells to reduce MTT provides an indication of mitochondrial activity and integrity, which are interpreted as measures of cell viability. The technique was first described to evaluate compounds with antitumor activity, which is closely related to cytotoxicity.

3.3.2 Genotoxicity of hemolymph

The genotoxicity of raw hemolymphs and its fractions obtained by gel filtration chromatography was evaluated by the comet method as described by Magneli et al, (2021).

3.4 Chemical inducers of apoptosis

Tert-butyl hydrogen peroxide (t-BPH) (25 to 2000 μM), Hydrogen Peroxide (H_2O_2) (25 to 2000 μM) and Actinomycin D (200 and 800ng/mL) were used as apoptosis chemical inducers.

3.5 Characterization of the biological activity of hemolymphs and its fractions

3.5.1 Effect of *Podalia sp* hemolymph in promoting cell growth

The total hemolymph of ***Podalia sp*** was tested in **Sf-9 cell** cultures, seeking to identify its ability to promote cell growth. For this, 1% of the total hemolymph v/v was added, at the time of cell inoculum to the culture of Sf-9 cells. Culture samples were obtained daily and the number of viable cells was determined by the trypan blue staining method.

3.5.2 Anti-apoptotic action of hemolymphs

3.5.2.1 Protector effect of *Podalia sp* hemolymph in death induced by nutrient depletion

The effect of total hemolymph and its fractions on **VERO cell** growth in culture with depleted medium was analyzed. To this, after the cells confluence, the medium culture was changed by PBS. At the time of nutrient depletion, 1% v/v of total hemolymph or its fraction 3 obtained from gel filtration column chromatography was added to the and analysed for 96 hous.

3.5.2.2 Protector effect of *Podalia sp* and M. albicolis hemolymph in death induced by t-BPH and H_2O_2

The apoptotic effect of ***Podalia sp*** and ***M. albicolis*** hemolymph on protection against cell death induced by chemical inducers of apoptosis. **VERO** cells were pre-treated or not with 1 or 2% (v/v) crude hemolymph for 1 hour. After this period, cell death was induced with 0,5mM or 1 mM of **t-BPH** or **H_2O_2** (25, 50, 100 or 400µM), for 18 hours. Cells were stained with crystal violet and observed under an inverted optical microscope (400x).

3.5.3 Determination of cell death by epifluorescence microscopy, confocal microscopy and optical microscopy

Total hemolymph of ***Podalia sp or M. albicolis*** and its fractions, obtained by chromatography, were tested in cell cultures, seeking to verify factors present in the hemolymph that promote cellular protection against death by apoptosis. For this, total hemolymph (1% v/v) or its fractions (2% v/v) were added to the cultures of **Vero and Sf-9** cells and after 1 hour the cell death inducers were added. After 2 days, for Vero cells or 5 days for Sf-9 cells, of cell contact with the inducer, the cultures were stained and analyzed under epifluorescence optical microscopy.To this, 5 µl volume of a solution containing 100 µg/ml of acridine orange **(AO)** and 100 µg/ml of ethidium bromide **(EB)** was added to 100 µl of cell culture (**Vero or Sf-9**) at a concentration between 0.5 and 1 x 10^6 cells/mL. This mixture was examined in epifluorescence microscopy with ultraviolet light and blue filter and confocal microscopy. Apoptosis was also visualized in light microscopy after staining with trypan blue.

3.5.4 Determination of cell death by flow cytometry

Vero cells were treated or not with total hemolymph from *Podalia sp or M. albicolis* (1% v/v) and after 1 hour t-BPH or Actinomycin D was added. After 18 hours, samples containing 10^6 cells were obtained. These were centrifuged at 800g for 5 minutes, the pellet was fixed in ice-cold 70% ethanol and stored at -20°C. On the day of analysis, ethanol was removed by centrifugation (800g for 5 minutes) and cells were incubated with 1 mL of Vindelev lysis buffer (1 mg/ml citrate, 50 µg/mL propidium iodide, 1 mL NP -40 (0.1%), 700 U/ml RNase, 0.01 M NaCl). After 10 minutes, the cells were centrifuged (10000g for 5 minutes) and the pellet resuspended in 1 mL of FACS buffer. The samples were processed in a FACS Calibur (BD Biosciences) with a wavelength of 488 nm for excitation and 620 nm for emission, determining the different phases of the cell cycle.

3.5. 5 Determination of alteration of the cytoskeleton of Vero cells by phalloidin-FITC labeling

One of the first changes observed in apoptotic cells is the loss of cell adhesion due to alterations in the cytoskeleton by microfilament retraction. To visualize the actin filaments, label the Vero cells with FITC-conjugated phalloidin. Furthermore, a co-labelling of the nuclei with propidium iodide was performed. For this, the cells were seeded in plates with coverslips at a maximum density of 10^5 cells/mL and after 24 hours they were pre-treated or not with the total hemolymph of *Podalia or M. albicolis* (1% v/v) for 1 hour and then induced with different doses of Terbutyl (25, 50 and 100µM) for 4 hours. After this time, the cell medium was removed and the cells were fixed with 4% paraformaldehyde in PBS for 20 minutes at room temperature. Cells were washed 3 times/5 minutes with PBS. Next, the cells were permeabilized with 0.1% Triton X-100 in PBS at room temperature for 15 minutes. After this time, a block was performed with PBS+BSA 1%+ RNase (100µg/mL), 40 minutes at 37 ºC. These cells were then incubated with FITC-conjugated phalloidin (20µg/ml) diluted in PBS+BSA1%, 3 hours at 37°C. Cells were washed 3 times, 1 minute each with PBS and incubated with propidium iodide (20 µg/ml) diluted in PBS+BS1%, for 1.5 h at 37°C. Cells were again washed 3 times/5 minutes with PBS. The slides were passed under a confocal microscope (LSM 510 META, ZEISS). The labeling with phalloidin-FITC was followed in laser starting at 488 nm with the emission filter the BP500-550 nm, while for the iodide it was used to study the laser 543 nm with the emission filter LP560 nm.

3.5.6 Labeling of tubulin in Vero cells

For the visualization of tubulin, **Vero cells** were labeled with anti-tubulin antibodies. Additionally, a co-labelling of the nuclei with propidium iodide was carried out. For this, the cells were seeded in microplates containing coverslips at a maximum density of 10^5 cells/mL. After 24 hours, the cell medium was removed and the cells were fixed with 4% paraformaldehyde in PBS for 20 minutes at 37°C. Cells were washed 3 times/5 minutes with PBS. After, the cells were permeabilized with 0.1% Triton X-100 in PBS at room temperature for 10 minutes. After this time, blocking was performed with PBS+ Triton X-100 0.05% + FBS 5% + RNAse (100µg/mL), 40 minutes at 37 °C. Cells were washed 3 times, 1 min each with PBS + 5% FBS. Cells were again washed 3 times/5 minutes with PBS. Cells were incubated with primary antibody, anti-beta-tubulin produced in mice (Sigma), (1 µg/mL + PBS + 0.05% Triton X-100 + 5% FBS) at 4°C overnight. After this time the cells were washed 3 times/5 minutes each with PBS. Then, the cells were incubated with blocking solution for 2 hours at room temperature with the secondary antibody, anti-mouse IgG conjugated to **ALexa Fluor 488** (Life Technologies), (10 µg/mL + PBS + Triton X-100 0.05% + 5% FBS at room temperature for 2 hours. Cells were then washed 2 times/5 minutes each with PBS. Propidium iodide (10µg/mL) was placed for 1 hour and after this time the cells were again washed 3 times/5 minutes each with PBS. Slides were then mounted with VectaShield and sealed with enamel. The slides were analyzed under a confocal microscope (LSM 510 META, ZEISS). The labeling for Tubulin was analyzed with a 488 nm laser for excitation and with a BP500-550 nm emission filter, while for iodide a 543 nm laser was used for excitation with an LP560 nm emission filter.

3.5.7 Determination of mitochondrial membrane potential ($\Delta\Psi$m)

For the determination of the effect of hemolymph in protecting tert-butyl-induced apoptosis **(t-BPH), Sf-9 cells** were cultured in 12-well microplates. After semi-confluence the cells were treated with 1% (v/v) crude hemolymph. After 1 hour of contact, the cells were treated with 600 µM of t-BPH. The cultures were kept in an oven for 4 hours and after this period the cells were washed with PBS and treated with the dye JC-1 (iodide of 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolylcarbocyanine) (2µM). The cells were analyzed and photographed in confocal microscopy.

4. RESULTS AND DISCUSSION

4.1 Determination of cell death by epifluorescence microscopy, and optical microscopy

Apoptosis in **Vero and Sf-9 cells** was observed by epifluorescence microscopy with ultraviolet light and blue filter after stained with acridine orange and ethidium bromide **(figure 1)**. Apoptosis was also visualized in light microscopy after staining with trypan blue, as can be seen in **Figure 2.** The photos in Figures 1 and 2 are representative of all the experiments carried out to determine apoptosis by microscope.

Figura 1: Photomicrograph of **Vero and Sf-9 cells**. Cells at different stages of death were stained with acridine orange and ethidium bromide. Viable cells (V) are stained in green. In red are observed cells in process of early apoptosis (AP) or late APt apoptosis). Also are observed cells in necrosis process (NC). Cells were observed under an epifluorescence microscope (100x (or 400x)

Figura 2: Photomicrograph of Sf-9 cells showing viable (v) or at different stages of death apoptosis (AP). The formation of apoptotic bodies (CA) and cells with later apoptosis (APt) is also visible. A sample of a Sf-9 cell culture was obtained on day 5, stained with trypan blue and visualized under an optical microscope. (400x).

4.2 Hemolymph fractionation by chromatography

To determine which fraction is responsible for the observed effects, total hemolymph from **Podalia sp and M. albicolis** were semi-purified in a gel filtration column (superdex 75). The obtained fractions were pooled in 3 pools for *Podalia sp* and 6 pools to *M. albicolis* and tested for their anti-apoptotic activity. In this first step of purification in gel filtration columns, the molecular weight profile of the different proteins that make up the hemolymph was determined. The chromatographic profile is shown in **figure 3 and 4** and the polyacrylamide gel of fractions in **figure 5**

Figura 3: Purification of *Podalia sp* hemolymph by gel filtration chromatography on a "Superdex 75" column. Elution was performed with Tris (20mM) and Nacl (150mM) solution. The flow used was 0.5 ml/minute and fractions of 1 ml were collected. The fractions were separated into 3 fractions as indicated by the bars

Figura 4: Purification of **M. albicolis** hemolymph by gel filtration chromatography on a "Superdex 75" column. Elution was performed with Tris (20mM) and Nacl (150mM) solution. The flow used was 0.5 ml/minute and fractions of 1 ml were collected. Fractions were separated into 6 groups as indicated

Figure 5: 12% SDS-PAGE electrophoresis gel. Crude hemolymph from **Podalia sp** (Pd), **M. albicolis** and its fractions obtained by gel filtration chromatography: Column 1 (PM) corresponds to the calibration molecular weight. Pd (*Podalia sp* crude). Ma (*M. albicolis* crude). The gel was stained with Coomassie Blue dye.

4.2.1 Determination of protein concentration

The protein concentration present in the hemolymphs and in its fractions obtained by gel filtration chromatography was determined by the NanoDrop 2000. The absorbance reading was performed at 280 nm. The results are presented in the **table 1**.

Table 1. Measurement of protein concentration in the crude hemolymph of *Podalia sp, M. albicolis* and the fractions obtained from the purification of the hemolymph by gel filtration chromatography on a "Superdex 75" column.

Podalia SP	mg/mL	M. albicolis	mg/mL
Hemolymph crude	41,012	Hemolymph crude	42,591
Pool 1	1,060	Pool 1	1,085
Pool 2	1,943	Pool 2	0,259
Pool 3	0,828	Pool 3	0,383
		Pool 4	0,046
		Pool 5	0,129
		Pool 6	0,100

4.2.3 Cytotoxicity

The cytotoxicity of total hemolymph of **Podalia sp** and **M. albicolis** was determined in Vero and Sf-9, at concentrations of 0.5, 1, 2, 5 and 10% v/v. Hemolymphs were cytotoxic only at concentrations greater than 5% of the final culture volume, demonstrating that its application in smaller amounts in cultures is feasible. At lower concentrations, no significant cell death was observed in any of the tested cells. From this result, the use of 1% and 2% (v/v) was determined in all tests (date not showed).

4.2.4 Genotoxicity of *Podalia sp and M. Albicolis* hemolymph

The genotoxicity of crude hemolymph from *Podalia sp and M. Albicolis* and its fractions obtained by gel filtration chromatography was evaluated by the comet method. In our study, according to the classification obtained, 90% of the comets were class 0 (not damaged) or class 1 (little damage), showing that the tested material is safe and non-genotoxic (date not showed).

4.3 Effect of *Podalia sp* hemolymph in promoting cell growth

The effect of total hemolymph on **SF-9 cell** growth was analyzed. As can be seen in **figure 6,** the culture treated with hemolymph had no effect on the number of cells, however we can observe a prolongation of cell viability, an indication of a protective action.

Figure 6: Cultures of **SF-9 cells** in Schott flasks containing 13 ml of Sf-900 medium, maintained at 29°C and 120 rpm. Cultivation was started with 3×10^5 cells/ml. Samples were obtained daily and the number of viable cells determined by trypan blue staining.

4.3.1 Protector effect of *Podalia sp* hemolymph in death induced by nutrient depletion

The effect of total hemolymph and its fractions on **VERO cell** growth in culture with depleted medium was analyzed. To this, the medium culture was changed by PBS e analysed for 96 hous. As can be seen in **figure 7,** the withdrawal of nutrients leads to cell death. However, when added *Podalia* **sp** hemolymph or fraction 3 obtained from gel filtration chromatography, the morphology of the culture is maintained.

Figure 7: Effect of *Podalia sp* hemolymph on protection from cell death induced by nutrient depletion. At the time of nutrient depletion, 1% v/v of total hemolymph or its fraction 3 obtained from gel filtration column chromatography was added to the culture. After 96 hours the cells were observed in an inverted optical microscope (400x).

We also tested the ability of total hemolymph of *M. Albicolis* to protect against apoptosis induced by nutrient depletion. To this, **VERO cells** were maintained in PBS for 96 hours plus 1% v/v total **M. albicolis** hemolymph. As can be seen in **Figure 8**, the removal of nutrients leads to the death of the cell culture. However, when *M. albicolis* hemolymph was added to the culture depleted of culture medium, the culture morphology was maintained

Figure 8: Effect of *M. albicolis* hemolymph on protection from cell death induced by nutrient depletion. At the time of nutrient depletion, 1% v/v of total hemolymph was added to the culture. As a control, wells from the same culture plate were maintained in L15 medium with 10% fetal bovine serum. After 96 hours the cells were observed in an inverted optical microscope (400x).

4.4 Effect of apoptosis inducers on Sf-9

The action of the chemical inducers of apoptosis, **t-BPH** (2mM) **Actinomycin D** (200 ng/ml) and **Hydrogen Peroxide** (H_2O_2) (2mM), was analyzed in **Sf-9 cells**. The cultures were kept in contact with the inducer for 18 hours and cell viability was determined by trypan blue. As can be seen in **Figure 9**, all 3 inducers were able to induce cell death at the tested concentrations, 200ng/ml for actinomycin and 2 mM for t-BPH 2 H_2O_2.

Figure 9: SF-9 cells treated or not with the apoptosis inducers t-BPH (2mM), Actinomycin D (200ng/ml) or H2O2 (2mM). After 18 hours of exposure, cells were stained with trypan blue and counted in a Newbauer chamber. *$p<0.05$ compared to control (ANOVA and Newman-Keuls multiple comparison).

4.5 Anti-apoptotic action of crude hemolymph from *Podalia sp*

The protection of hemolymph from ***Podalia sp*** (2% v/v) on **Vero cells,** treated with t-BPH (0,5 or 1 mM) after 18 hours was determined. The cells and cell viability was determined by MTT. As can be seen, the hemolymph of *Podalia sp* reduce the induction of apoptosis death by **t-BPH** (**figure 10, 11**).

Figure 10: Viability of **Vero cells** (by MTT) after 18 hours of exposure with **t-BPH** (1mM) and treated with crude hemolymph from ***Podalia sp*** (2% v/v)

Figure 11: Effect of *Podalia sp* hemolymph on protection against cell death induced by chemical inducers of apoptosis. VERO cells were pre-treated or not with 2% (v/v) crude hemolymph for 1 hour. After this period, cell death was induced with 0,5mM of t-BPH for 18 hours. Cells were stained with crystal violet and observed under an inverted optical microscope (400x).

To verify if there hemolymph of **_Podalia sp_** promotes cellular protection against **H_2O_2**, VERO cells were treated with 1% v/v of the hemolymph of _Podalia sp_ 1 hour before the addition of H_2O_2 (25, 50 or 100µM), (**figure 12**). As can be seen, total hemolymph from _Podalia sp_ was able to protect VERO cells from death induced by H_2O_2

Figure 12: Effect of hemolymph in protecting cell death induced by Hydrogen Peroxide (H_2O_2). VERO cells were pre-treated or not with 1% (v/v) of total hemolymph from _Podalia sp_ and after 1 hour cell death was induced with H_2O_2 at concentrations of 25, 50 or 100µM for 18 hours. Cells were stained with crystal violet and observed under an inverted optical microscope at 400x magnification.

4.6 Anti-apoptotic action of crude hemolymph from *M. albicolis*

To study the anti-apoptotic effect of **M. albicolis** hemolymph, **VERO** cells were pre-treated or not with total hemolymph (1% v/v) for 1 hour. After this period, cell death was induced with **H_2O_2** (0,4mM) (**figure 13a**) or **t-BPH** (0,5mM) (**figure 13b**) for 18 hours. Total hemolymph from *M. albicolis* was able to neutralize the effects of apoptosis inducers

Figure 13 a: Effect of *M. albicolis* hemolymph on protection against cell death induced by chemical inducers of apoptosis. VERO cells were pre-treated or not with 1% (v/v) crude hemolymph for 1 hour. After this period, cell death was induced with 0,4 mM of H_2O_2. for 18 hours. The cells were observed in an inverted optical microscope (400x).

Figure 13 b: Effect of *M. albicolis* hemolymph on protection against cell death induced by chemical inducers of apoptosis (t-BPH). VERO cells were pre-treated or not with 1% (v/v) crude hemolymph for 1 hour. After this period, cell death was induced with 0,5 mM of t-BPH for 18 hours. Cells were stained with crystal violet and observed under an inverted optical microscope (400x).

4.7 Analysis of the anti-apoptotic effect of hemolymph from *Podalia* and M. albicolis by staining acridine orange (AO) and ethidium bromide (BE) with AO/BE

Cell death by apoptosis in **SF-9 cell** cultures was also determined by confocal microscopy after staining with acridine orange (AO) and ethidium bromide (BE) fluorophores. For this, SF-9 cells were treated with 1% v/v of hemolymph or its fractions 1 hour before the addition of **(t-BPH)** at a concentration of 1mM (**Figure 14 a**) or **Actinomycin D** at a concentration of 800ng/mL (**Figure 14 b**). The total hemolymph of ***Podalia sp and M. albicolis*** showed anti-apoptotic effect with both inducers. Viable cells stain uniformly green while apoptotic cells stain red

Figure 14 a: Morphology of SF-9 cells stained with Acridine Orange and Ethidium Bromide. A) cells treated with **1mM t-BPH**; B) cells treated with 1mM t-BPH and *M. albicolis* hemolymph; C) cells treated with 1mM t-BPH and Podalia sp hemolymph (arrow: apoptotic cells).

Figure 14 b : Morphology of SF-9 cells stained with Acridine Orange and Ethidium Bromide. A) cells treated with **Actinomycin D 800ng/ml;** B) cells treated with Actinomycin D 800ng/mL and *M. albicolis* hemolymph; C) cells treated with Actinomycin D 800ng/mL and *Podalia sp* hemolymph (arrow: apoptotic cells).

VERO cells were maintained in PBS for 24 hours or in PBS plus 1% v/v of total hemolymph from *Podalia sp* as described above. Hemolymph was able to protect VERO cells from death induced by nutrient depletion **(Figure 15).**

VERO IN PBS

VERO IN PBS + 1% Hb Podalia sp

Figure 15: Effect of hemolymph on protection from cell death induced by nutrient depletion. VERO cells were kept in PBS for 24 hours and treated or not with 1% (v/v) total hemolymph from ***Podalia sp.*** Cells were stained with acridine orange and ethidium bromide and observed under a confocal microscope.

4.8 Analysis of the anti-apoptotic effect of Podalia hemolymph in flow cytometry

Determination of DNA content by flow cytometry has been widely used to identify apoptosis in cell culture. For this, VERO cells were treated with 1% v/v of **Podalia sp** hemolymph 1 hour before adding **t-BPH** at a concentration of 0,5mM. **Figure 16** shows a histogram of the control culture, with a typical DNA profile of these cells, where the G1, S and G2 phases are visible and the sub-G1 population is low. In cultures where the cells were treated only with t-BPH an increase in the sub-G1 population can be observed, indicating a process of death by apoptosis. In the culture where cell death was induced, but with the previous addition of hemolymph, there was an increase in populations in other phases of the cycle, indicating that hemolymph acts by stimulating cell proliferation and preventing cell death by apoptosis

Figure 16: Histogram of the DNA of VERO cells treated with 1% v/v of *Podalia sp* hemolymph 1 hour before the addition of t-BPH at a concentration of 0,5 mM

4.9 Determination of alteration of the cytoskeleton of VERO cells by labeling with phalloidin-FITC

Studies have provided important information about how the actin cytoskeleton contributes to the control of growth, both in transformed cells and in non-transformed cells, indicating its participation in the regulation of several cellular processes related to transformation, such as proliferation, anchorage-independent growth and contact inhibition of growth (Pawlak and Helfman, 2001). Cellular transformation into fibroblasts, as well as the process of apoptosis, is characterized by several morphological changes, with the rounded phenotype and the decrease in actin bundles of the cytoskeleton and stress fibers being striking features (Janmey and Chaponnier, 1995). In control cells, labeled with phalloidin-FITC, a network of highly organized actin filaments is observed, with abundant stress fibers, most of which have an orientation parallel to the major cell axis. With the induction of cell death by **t-BPH** there was a significant reduction in the number of stress fibers and a shortening of those that prevailed. In order to study the possible effect of hemolymph in maintaining the cytoskeleton of cells induced to apoptosis, VERO cells were pre-treated with **Podalia sp** hemolymph and treated with $100\mu M$ of t-BPH for 4 hours. When VERO cells were pre-treated with total hemolymph from **Podalia sp,** before apoptosis induction, the cytoskeleton was maintained, showing the stress fibers clearly evident. Therefore, the hemolymph of **Podalia sp** showed a protective effect, blocking or mitigating the disruption of the cytoskeleton.

Figure 17: Actin filaments in Vero cells (phalloidin labeling). (A) Control VERO cells showing abundant fibers parallel to the major cell axis (arrows). The nucleus was stained with propidium iodide. (B) VERO cells treated with t-BPH (100µM) for 4 hours. (C) Cells pretreated with total Podalia hemolymph (1% v/v) for 1 hour. After this period, they were treated with t-BPH at a concentration of 100µM for 4 hours. In cells pre-treated with hemolymph, the cytoskeleton was maintained, showing clearly evident stress fibers (arrow). The nucleus was not marked. Cells were analyzed by confocal microscopy.

Likewise, VERO cells were pre-treated with hemolymph of *M. albicolis* total by 1 hour before apoptosis induction with 100µM t-BPH. As can be seen, the cytoskeleton and nucleus were also maintained in the culture treated with *M. albicolis* hemolymph, demonstrating the protective effect, blocking or mitigating the disruption of the cytoskeleton. (**figure 18**).

Figure 18: Actin filaments in Vero cells (phalloidin labeling). (A) Cells treated with t-BPH at a concentration of 100µM for 4 hours. (B) VERO cells pre-treated with *M. albicolis* total hemolymph (1% v/v) for 1 hour. After this period, the cells were induced to death with t-BPH at a concentration of 100µM, for 4 hours. In cells pre-treated with hemolymph, the cytoskeleton was maintained, showing clearly evident stress fibers (arrow). The core was labeled with propidium iodide. Cells were analyzed by confocal microscopy.

4.10 Labeling of tubulin in VERO cells

The addition of hemolymph also maintained the tubulin structure in VERO cells even after the addition of 100μM t-BPH. Microtubules were evidenced by a primary anti-β tubulin antibody, followed by a secondary anti-mouse IgG antibody conjugated to ALexa Fluor 488 (**Figure 19**). Microtubules appear distributed throughout the cytoplasm of cells similar to the organization of normal cells. Mitotic cells with microtubule organization characteristic of normal mitotic spindles can also be seen.

Figure 19: Microtubules in Vero cells after induction of apoptosis with 100μM t-BPH but pre-treated with M. albicolis total hemolymph (1% v/v) for 1 hour. Microtubules were evidenced by primary anti-β tubulin antibody, followed by secondary anti-mouse IgG antibody conjugated to ALexa Fluor 488. The nucleus was labeled with propidium iodide. Chromosomes are highlighted in red and microtubules in green. The blue arrow points to polar microtubules and the white arrow points to astral microtubules. Cells were analyzed by confocal microscopy.

4.11 Determination of mitochondrial membrane potential ($\Delta\Psi$m) in SF-9 cells

Oxidative stress is related to the loss of mitochondrial integrity and the triggering of the intrinsic pathway of apoptosis. We used the dye JC-1 (5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolylcarbocyanine iodide) to evaluate the integrity of the mitochondrial membrane of **Sf-9 cells** pre-treated or not with the hemolymph of *Podalia* and induced apoptosis with **t-BPH** (600μM). Red staining indicates healthy mitochondria with preserved membrane potential (A, C). Green staining indicates loss of mitochondrial membrane electrical potential (B). (**Figure 20**). The following figures show representative images of the analysis with JC-1 in confocal microscopy.

Figure 20: Sf-9 cells labeled with JC-1. (A) control cells; (B) cells treated with t-BPH (600µM) for 4 hours; (C) Cells pre-treated with hemolymph OF *Podalia's* total (1 hr) and treated with t-BPH (600µM) for 4 hours.

Mitochondria are involved in many processes essential for cell survival, including energy production, redox control, calcium homeostasis, and certain metabolic and biosynthetic pathways. In addition, mitochondria play an essential role in the cell death process known as apoptosis. This mitochondrial pathway of death by apoptosis can suffer dysfunctions causing various diseases such as cancer, diabetes, ischemia and neurodegenerative disorders such as Parkinson's and Alzheimer's (Bouchier-Hayes et al., 2005). Due to the numerous activation mechanisms of death by apoptosis, the determination of the mechanism of action of the different anti-apoptotic proteins, especially those with a broad spectrum of action, whether inhibiting death by different inducers or preventing this death in different cell types, can be of great importance. high medical and biotechnological interest. Studies with an anti-apoptotic protein isolated from the hemolymph of *Lonomia obliqua* showed that its likely action would be on the mitochondrial membrane, keeping the electrical potential high and preventing membrane permeability (Mendonça & Martins, 2022).

The hemolymph of *Podalia and M. albicolis* showed a very similar action with the anti-apoptotic protein of *Lonomia obliqua*. It is also suggested that the site of action of this protein could be similar to that of the Bcl-(x) family proteins since, in all experiments, a block or decrease in death by apoptosis was observed, with evident protection of the potential of the mitochondrial membrane . Due to the broad spectrum of action of these hemolymphs, which act both in insect and mammalian cells, either by inducing apoptosis by chemical or nutritional agents, there is a molecule present in the hemolymph capable of acting as a general suppressor of the apoptosis process, for example a conserved pathway in many organisms.

The addition of hemolymph to cultures not only neutralized or reduced the chemically induced death, but also the physiological death that normally occurs due to culture stress. Cells treated with hemolymph remained alive longer, indicating that this protein could also be used as a supplement to culture media. By blocking cell death, the cell remains functional, and this functionality is a fundamental indication when thinking about using a molecule with these characteristics as a drug.

The hemolymph of the caterpillars *Podalia sp and M. albicolis* showed an anti-apoptotic effect both in insect cells and in mammalian cells, showing that eventually this protein is acting in some basic mechanism of cell death.

Silkworm hemolymph has been reported to inhibit apoptosis in later stages of culture, both in insect and human cells (Choi et al., 2002). Studies have shown the occurrence of cell death by apoptosis in cultures of mammalian and insect cells with nutrient depletion (Mendonca et al., 2002; Meneses-Acosta et al., 2001) or by induction of chemical agents (Souza et al. , 2005), The cell death is a very important factor in production processes that limited the industrial production of some proteins of economic interest., One the way to increase the cellular productivity is to inhibit or attenuate the cellular death. Earlier, we have demonstrated the presence of a potent anti-apoptotic protein in Lonomia obliqua hemolymph which extends the cell culture viability through apoptosis prevention (Maranga et al., 2003; Raffoul et al., 2005; Souza et al., 2005). Has been reported also that mitochondria has one important action in the apoptosis control process, being that Mitochondria Membrane Permeabilization (MMP) an important stage in this process. We have showed also that the addition of *Lonomia obliqua* hemolymph in the culture lead to a prolongation of the cellular, leading a high electrochemical potential of the mitochondria, avoiding the loss of the membrane permeability and the Cytochrome-C release. Thus, even though they originate from different species, we believe that the hemolymph of *Podalia sp* e *M. albicolis* can also act through similar mechanisms to the hemolymph of *lonomia obliqua.* By the other hand, the hemolymph from *Podalia sp* and *M. albicolis* caterpillars did not prove to be cytotoxic or genotoxic to cell cultures at concentrations below 5%, what may be useful for use as a cell cultivation additive. Even though the electrophoretic profiles of both caterpillars are different, the effects were similar, both in mammalian and insect cell, which may indicate the presence of components with functions and mechanism of action similar. Crude hemolymph from both caterpillars showed anti-apoptotic effect in VERO and SF-9 cells pre-treated with only 1% v/v of hemolymph and induced to death by different, and potent apoptotic inductors as terbutyl, actinomycin D, hydrogen peroxide and by depletion of nutrients.This protector effect blocked and attenuated the disruption of the cytoskeleton (actin filaments), being that the protective effect also was observed on the integrity of the mitochondrial membrane of SF-9 cells pre-treated with both hemolymphs and treated with the apoptosis inducer Terbutil at concentrations of 25 to 100µM.

5. CONCLUSION

The hemolymph of *Podalia sp* and *M. albicolis* caterpillars did not prove to be cytotoxic or genotoxic to cell cultures at concentrations below 5%. The electrophoretic profile of the hemolymph of *Podalia sp* was different from the electrophoretic profile of *M. Albicolis.* Was observed in the hemolymphs of the caterpillars *Podalia sp.*, and *M. albicolis* molecules of pharmacological and biotechnological interest. Crude hemolymph from both caterpillars showed anti-apoptotic effect in both, VERO and SF-9 cells, against different inducers as terbutyl, actinomycin D and hydrogen peroxide and by depletion of nutrients. The hemolymph of both caterpillars showed a protective effect, blocking or attenuating the disruption of the cytoskeleton (actin filaments) of VERO cells treated with the apoptosis inducer Terbutil at concentrations of 25 to 100μM. Protective effect was observed on the integrity of the mitochondrial membrane of SF-9 cells pre-treated with both hemolymphs and treated with the apoptosis inducer Terbutil at concentrations of 25 to 100μM

Acknowledgments

The authors acknowledge the financial support received from FAPESP:2010/52434-6 and a doctorate fellowship to Nathalia Delazeri de Carvalho from FAPESP-2011/50095-2

Conflict of interest. The authors of the article "Proteins with anti-apoptotic action in the hemolymph of caterpillars of the Megalopygidae family acts by maintaining the structure of the cellular cytoskeleton " declare that there are no known conflicts of interest associated with this publication and that there was no significant financial support for this work that could have influenced its outcome.

7. REFERENCES

Bouchier-Hayes, L., Lartigue, L., Newmeyer, D.D., 2005. Mitochondria: pharmacological manipulation of cell death. J Clin Invest. 115**,** 2640-7.

Cantero TM, Silva Junior PI , Negri G , Nascimento RM Mendonça RZ . 2022 Antimicrobial activity of flavonoids glycosides and pyrrolizidine alkaloids from propolis of Scaptotrigona aff. Postica. Toxin Reviews 41:4,

Carmona, G.S., Sousa, I.L.M., Laurentino, R.V., Junior, D.L., Pantoja, A.V., Mendonça, R.Z., Nascimento, R.M. and Valentin, F.N. 2020 Propolis and its components can help reduce the physiopathological consequences of covid-19 infection. International Journal of Development Research 10(10): 41067-41069

Carvalho, N.D; Curt, S.P; Oliveira, M. I.; Rofatto, H.K; Figueiredo, C.A; Senna Villar, K; Magnelli, R.F.; Mendonça, RZ. 2022. Antiviral Activity of the Hemolymph of Podalia sp and M. Albicolis (Lepidoptera: Megalopigydae Journal of Infectious Diseases & Therapy v.10, p.1 - 7

Choi, S.S., Rhee, W.J., Park, T.H., 2002. Inhibition of human cell apoptosis by silkworm hemolymph. Biotechnol Prog. 18, 874-8.

Cardoso, A.E.C and Junior V.H. 2005 Accidents caused by lepidopterans (moth larvae and adult): study on the epidemiological, clinical and therapeutic aspects. An. Bras. Dermatol. 80 (6)

Fonseca F. Acidentes por lagartas. In: Animais peçonhentos. São Paulo: Instituto Butantã; 1950. p. 284-96

Coelho, G.R.,Figueiredo, C.A., Negri, G., Fernandes-Silva, C.C.,Senna Villar,K. Badari, J.C., Oliveira,M.I., Barbosa, T.F., Taniwaki, N.N., Namiyama G.M., and Mendonça R.Z. 2018. Antiviral Activity of Geopropolis Extract from Scaptotrigona Aff. Postica against Rubella Virus Journal of Food Research; Vol. 7, No. 6; pag 91-106

Greco, K.N., Mendonça, R.M.Z., Moraes, R.H.P., Mancini, D.A.P., Mendonça, R.Z., 2009. Antiviral activity of the hemolymph of Lonomia obliqua (Lepidoptera: Saturniidae). Antiviral Research. 84, 84-90.

Gross, J., Muller, C., Vilcinskas, A., Hilker, M., 1998. Antimicrobial activity of exocrine glandular secretions, hemolymph, and larval regurgitate of the mustard leaf beetle phaedon cochleariae. J Invertebr Pathol. 72, 296-303.

Guerrero, B., Perales, J., Gil, A., Arocha-Pinango, C.L., 1999. Effect on platelet FXIII and partial characterization of Lonomin V, a proteolytic enzyme from Lonomia achelous caterpillars. Thromb Res. 93, 243-52.

Hamdaoui, A., Wataleb, S., Devreese, B., Chiou, S.J., Vanden Broeck, J., Van Beeumen, J., De Loof, A., Schoofs, L., 1998. Purification and characterization of a group of five novel peptide serine protease inhibitors from ovaries of the desert locust, Schistocerca gregaria. FEBS Lett. 422, 74-8.

Hayashida, Patricia Yumi, Silva Junior Pedro Ismael, Mendonça, Ronaldo Zucatelli. 2019. *Lonomia obliqua*'s hemolymph: new horizons to antibiotics. Toxicon 121, 1-16

Huberman, A., Aguilar, M.B., Navarro-Quiroga, I., Ramos, L., Fernandez, I., White, F.M., Hunt, D.F., Shabanowitz, J., 2000. A hyperglycemic peptide hormone from the Caribbean shrimp Penaeus (litopenaeus) schmitti. Peptides. 21, 331-8.

Janmey, P.A., Chaponnier, C., 1995. Medical aspects of the actin cytoskeleton. Curr Opin Cell Biol. 7, 111-7.

Jiang, H., Wang, Y., Kanost, M.R., 1999. Four serine proteinases expressed in Manduca sexta haemocytes. Insect Mol Biol. 8, 39-53.

Johns, R., Sonenshine, D.E., Hynes, W.L., 1998. Control of bacterial infections in the hard tick Dermacentor variabilis (Acari: Ixodidae): evidence for the existence of antimicrobial proteins in tick hemolymph. J Med Entomol. 35, 458-64.

Jones, G., Manczak, M., Horn, M., 1993. Hormonal regulation and properties of a new group of basic hemolymph proteins expressed during insect metamorphosis. J Biol Chem. 268, 1284-91.

Kanaya, T., Kobayashi, J., 2000. Purification and characterization of an insect haemolymph protein promoting in vitro replication of the Bombyx mori nucleopolyhedrovirus. J Gen Virol. 81, 1135-41.

Kristiansen, E., Ramløv, H., Hagen, L., Pedersen, S.A., Andersen, R.A., Zachariassen, K.E. 2005. Isolation and characterization of hemolymph antifreeze proteins from larvae of the longhorn beetle Rhagium inquisitor (L.) Comp Biochem Physiol B Biochem Mol Biol Sep;142(1):90-7.

Kurata, K., Nakamura, M., Okuda, T., Hirano, H., Shinbo, H., 1994. Purification and characterization of a juvenile hormone binding protein from hemolymph of the silkworm, Bombyx mori. Comp Biochem Physiol B Biochem Mol Biol. 109, 105-14.

Laemmli, U.K., 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature. 227, 680-5.

Lanz-Mendoza, H., Bettencourt, R., Fabbri, M., Faye, I., 1996. Regulation of the insect immune response: the effect of hemolin on cellular immune mechanisms. Cell Immunol. 169, 47-54.

Lin, C.Y., Chen, S.H., Kou, G.H., Kuo, C.M., 1998. Identification and characterization of a hyperglycemic hormone from freshwater giant prawn, Macrobrachium rosenbergii. Comp Biochem Physiol A Mol Integr Physiol. 121, 315-21.

Lowenberger, C., Charlet, M., Vizioli, J., Kamal, S., Richman, A., Christensen, B.M., Bulet, P., 1999. Antimicrobial activity spectrum, cDNA cloning, and mRNA expression of a newly isolated member of the cecropin family from the mosquito vector Aedes aegypti. J Biol Chem. 274, 20092-7.

Maranga, L., Mendonca, R.Z., Bengala, A., Peixoto, C.C., Moraes, R.H., Pereira, C.A., Carrondo, M.J., 2003. Enhancement of Sf-9 cell growth and longevity through supplementation of culture medium with hemolymph. Biotechnol Prog. 19, 58-63.

Mazzoni, M.K.F., de Carvalho, N.D., Rofatto, H.K., Moraes, R.E.P., and Mendonça, R.Z. 2014. Study of antiapoptotic effect of a protein isolated from Megalopyge albicolis (Lepidoptera: Megalopygidae) hemolymph. BMC Proc. 2014; 8 (Suppl 4): P148. Published online Oct1. doi: 10.1186/1753-6561-8-S4-P148

Mendonca, R.Z., Arrozio, S.J., Antoniazzi, M.M., Ferreira, J.M., Jr., Pereira, C.A., 2002. Metabolic active-high density VERO cell cultures on microcarriers following apoptosis prevention by galactose/glutamine feeding. J Biotechnol. 97, 13-22.

Mendonca, R.Z., Greco, K.N., Sousa, A.P., Moraes, R.H., Astray, R.M., Pereira, C.A., 2008. Enhancing effect of a protein from Lonomia obliqua hemolymph on recombinant protein production. Cytotechnology. 57, 83-91.

Mendonça RZ and Martins, LM. 2022. The antiapoptotic effect of Lonomia obliqua Hemolymph is associated with the mitochondria patway. Journal of Membrane Science & Technology Vol. 12 Iss. 9 No: 1000298

Mendonça, R.Z., Fernandez, A.C., Nascimento RM Silva Junior PI. 2022 Antiviral action of aqueous extracts of propolis from Scaptotrigona aff. Postica against Zica, Chikungunya and Mayaro Virus. (bioRxiv 2022 doi: https://doi.org/10.1101/2022.11.03.515030)

Meneses-Acosta, A., Mendonca, R., Merchant, H., Covarrubias, L., Ramirez, O., 2001. Comparative characterization of cell death between Sf9 insect cells and hybridoma cultures. Biotechnol Bioeng. 72, 441-57.

Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. J Immunol Methods. 65, 55-63.

Nardone, R.M., 1977. Toxicity testing in vitro. . In Growth, nutrition and metabolism of cells in culture. Vol., Academic. New York, pp. 471-495.

Nussbaumer, C., Hinton, A.C., Schopf, A., Stradner, A., Hammock, B.D., 2000. Isolation and characterization of juvenile hormone esterase from hemolymph of Lymantria dispar by affinity- and by anion-exchange chromatography. Insect Biochem Mol Biol. 30, 307-14.

Ochanda, J.O., Osir, E.O., Nguu, E.K., Olembo, N.K., 1992. Isolation and properties of 600-kDa and 23-kDa haemolymph proteins from the tsetse fly, Glossina morsitans: their possible role as biological insecticides. Scand J Immunol Suppl. 11, 41-7.

Pan, J., Kurosky, A., Xu, B., Chopra, A.K., Coppenhaver, D.H., Singh, I.P., Baron, S., 2000. Broad antiviral activity in tissues of crustaceans. Antiviral Res. 48, 39-47.

Pawlak, G., Helfman, D.M., 2001. Cytoskeletal changes in cell transformation and tumorigenesis. Curr Opin Genet Dev. 11, 41-7.

Raffoul, T., Swiech, K., Arantes, M.K., Sousa, Á.P.B.d., Mendonça, R.Z., Pereira, C.A., Suazo, C.A.T., 2005. Performance evaluation of CHO-K1 cell in culture medium supplemented with hemolymph. Brazilian Archives of Biology and Technology. 48, 85-95.

Rhee, W.J., Park, T.H., 2000. Silkworm hemolymph inhibits baculovirus-induced insect cell apoptosis. Biochem Biophys Res Commun. 271, 186-90.

Rosenfeld, A., Vanderberg, J.P., 1998. Identification of electrophoretically separated proteases from midgut and hemolymph of adult Anopheles stephensi mosquitoes. J Parasitol. 84, 361-5.

Rotberg A. Taturanas e mariposas. Folha de São Paulo. (Acess in: 27 jan 1990

Shiotsuki, T., Bonning, B.C., Hirai, M., Kikuchi, K., Hammock, B.D., 2000. Characterization and affinity purification of juvenile hormone esterase from Bombyx mori. Biosci Biotechnol Biochem. 64, 1681-7.

Shishikura, F., Abe, T., Ohtake, S., Tanaka, K., 1996. Purification and characterization of a 58,000-Da proteinase inhibitor from the hemolymph of a solitary ascidian, Halocynthia roretzi. Comp Biochem Physiol B Biochem Mol Biol. 114, 1-9.

Shishikura, F., Abe, T., Ohtake, S., Tanaka, K., 1997. Purification and characterization of a 39,000-Da serine proteinase from the hemolymph of a solitary ascidian, Halocynthia roretzi. Comp Biochem Physiol B Biochem Mol Biol. 118**,** 131-41.

Souza, A.P., Peixoto, C.C., Maranga, L., Carvalhal, A.V., Moraes, R.H., Mendonca, R.M., Pereira, C.A., Carrondo, M.J., Mendonca, R.Z., 2005. Purification and characterization of an anti-apoptotic protein isolated from Lonomia obliqua hemolymph. Biotechnol Prog. 21**,** 99-105.

Vieira, H.L., Pereira, A.C., Peixoto, C.C., Moraes, R.H., Alves, P.M., Mendonca, R.Z., 2010. Improvement of recombinant protein production by an anti-apoptotic protein from hemolymph of Lonomia obliqua. Cytotechnology. 62**,** 547-55.

Woo, S.D., Kim, W.J., Kim, H.S., Choi, J.Y., Jin, B.R., Kang, S.K., 1997. Effect of silkworm hemolymph on the expression of E. coli beta-galactosidase in insect cell lines infected with recombinant baculoviruses. Mol Cells. 7**,** 572-4.

Yamamoto, Y., Watabe, S., Kageyama, T., Takahashi, S.Y., 1999. Purification and characterization of Bombyx cysteine proteinase specific inhibitors from the hemolymph of Bombyx mori. Arch Insect Biochem Physiol. 42**,** 119-29.

Zhu, S., Li, W., Jiang, D., Zeng, X., 2000. Evidence for the existence of insect defensin-like peptide in scorpion venom. IUBMB Life. 50**,** 57-61.