SARS-COV-2 NSP5 ANTAGONIZES MHC II EXPRESION BY SUBVERTING HISTONE DEACETYLASE 2

Nima Taefehshokr[1*], Alex Lac[1*], Angela M Vrieze[1], Brandon H Dickson[1], Peter N Guo[1], Catherine Jung[1], Eoin N Blythe[1,2], Corby Fink[1,2], Jimmy D Dikeakos[1], Gregory A Dekaban[1,2], Bryan Heit[1,2,†]

1. Department of Microbiology and Immunology, and the Western Infection, Immunity and Inflammation Centre, The University of Western Ontario, London, Ontario, Canada, N6A 5C1; 2. Robarts Research Institute, London, Ontario, Canada, N6A 3K7. †Corresponding Author: bheit@uwo.ca. *These authors contributed equally to this article.

SARS-CoV-2 interferes with antigen presentation by downregulating MHC II on antigen presenting cells, but the mechanism mediating this process is unelucidated. Herein, analysis of protein and gene expression in human antigen presenting cells reveals that MHC II is downregulated by the SARS-CoV-2 main protease, NSP5. This suppression of MHC II expression occurs via decreased expression of the MHC II regulatory protein CIITA. This downregulation of CIITA is independent of NSP5's proteolytic activity, and rather, NSP5 delivers HDAC2 to the CIITA promoter via an IRF3-dependent mechanism. Here, HDAC2 deacetylates and inactivates the CIITA promoter. This loss of CIITA expression prevents further expression of MHC II, with this suppression alleviated by ectopic expression of CIITA or knockdown of HDAC2. These results identify a mechanism by which SARS-CoV-2 can limit MHC II expression, thereby delaying or weakening the subsequent adaptive immune response.

SARS-CoV-2 | NSP5 | Major Histocompatibility Complex II | CIITA | Antigen Presentation | IRF3

Introduction

First identified in December 2019, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has rapidly become a leading global cause of morbidity and mortality. This high virulence is due, in part, to the presence of multiple mechanisms which allow SARS-CoV-2 to evade and alter host immune responses, thereby delaying viral clearance and prolonging the period in which patients remain infected [1,2]. Of note, humoral immunity against SARS-CoV-2 wanes twice as quickly compared to other human-infecting coronaviruses [3]. Even though vaccination has greatly reduced morbidity and mortality, breakthrough infections still occur and are becoming more common due to the emergence of new, more immunoevasive, SARS-CoV-2 variants [4,5]. Therefore, understanding the immunoevasion mechanisms of SARS-CoV-2 is important to better understand this deadly disease and to develop better targeted treatments and vaccines.

Although professional antigen presenting cells (pAPCs) such as macrophages and dendritic cells (DCs) do not express the SARS-CoV-2 receptor ACE2, they can be infected via Fc-receptor dependent phagocytosis of antibody-opsonized virions, and potentially through the efferocytosis of SARS-CoV-2 infected apoptotic cells [6–12]. While SARS-CoV-2 is unable to establish a productive infection in macrophages or DCs, viral early genes are expressed and drive a multi-pronged immunoevasion response. Firstly, the expression of pro-inflammatory cytokines is induced and contributes to the cytokine storm [12]. This cytokine response is typified by high circulating levels of IL-2, IL-6, IL-7, IL-8, IL-10, IP-10, G-CSF, MCP-1, MIP1-α, and TNF-α [13–15]. While this mechanism drives a potent inflammatory response, the cytokine profile is more typical of bacterial infections, and promotes both NK cell exhaustion and reduced NK cell cytotoxicity, thereby producing a non-productive innate immune response that can exacerbate tissue damage [16,17]. Secondly, SARS-CoV-2 suppresses the anti-viral interferon (IFN) pathway, reducing the production of type I and type II IFNs. This suppression is driven by ORF6,

which sequesters inactive signal transducer and activator of transcription 1 (STAT1) and STAT2 in the cytosol, thereby preventing their nuclear translocation and blocking the primary signalling pathway that initiates IFN responses [18]. Moreover, membrane protein and non-structural protein 13 further inhibits IFN-I production by degrading TANK-binding kinase 1 [19,20]. Thirdly, infected macrophages and dendritic cells often die, resulting in long-term depletion of some subsets [21]. Fourthly, SARS-CoV-2 directly suppresses antigen presentation on Major Histocompatibility Complex class I (MHC I) via ORF8, which directs the trafficking of MHC I to the lysosome where it is degraded [22]. MHC I is further suppressed by ORF6, which inactivates CITA/NLRC5—the master regulator of MHC I expression [23]. Finally, infection of alveolar DCs reduces their ability to migrate to draining lymph nodes and suppresses expression of MHC class II and the class II transcriptional activator (CIITA) required for MHC II expression [24]. This suppression of MHC II occurs across a range of pAPCs in SARS-CoV-2 infected patients and in cells infected *in-vitro* [25–27]. Moreover, some non-professional antigen presenting cells express MHC II in response to infection – including type II alveolar epithelial cells which are one of the primary targets of SARS-CoV-2 infection in the alveolus [28,29]. However, the cellular mechanism which downregulates MHC II in these cells has remained elusive.

MHC II expression is driven by type II interferon signaling [30]. Activation of the IFN-γ receptor complex leads to the structural rearrangements in the receptor complex and activation of Janus kinase 1 and 2 (JAK1/2) tyrosine kinases and phosphorylation and homodimerization of STAT1, which translocates to the nucleus to induce transcription of IFN-γ-inducible genes [31]. Here, STAT1 activates interferon regulatory factor -1 and -3 (IRF-1/3), with STAT1 and IRF-1/3 then cooperatively inducing expression of CIITA [32,33]. CIITA, via its intrinsic acetyltransferase activity, can then acetylate histones at the MHC II promoter, which decondenses the chromatin to allow access for transcription factors that regulate MHC II expression [34]. Once the chromatin is opened, CIITA and regulatory factor X (RFX) form an enhanceosome complex on the MHC II promoter which recruits and activates additional transcription factors that induce the transcription of MHC II [35]. While inhibition of STAT1 nuclear import by SARS-CoV-2 ORF6 may account for some inhibition of MHC II expression [18], tissue-resident DCs natively express significant amounts of CIITA [36], whereas ORF6 expression is limited until 12-16 hours post-infection [37] – more than sufficient time to induce presentation of SARS-CoV-2 antigens following infection. Moreover, IFN-independent mechanisms can drive some MHC II expression in macrophages and DCs [38–40]. Therefore, a more direct form of MHC II suppression is likely invoked by SARS-CoV-2. A critical regulator of MHC II expression is histone deacetylase 2 (HDAC2), which can suppress the expression of both CIITA and MHC II through deacetylation of histones within their promoters [41]. Gordon *et al*. recently mapped the SARS-CoV-2 protein interactome and identified non-structural protein 5 (NSP5) as an HDAC2 interactor, and identified a putative NSP5 cleavage site near the nuclear localization signal (NLS) of HDAC2 [42]. NSP5 – also known as main protease and 3-chymotrypsin like protease – is translated as part of the polyprotein expressed early after viral entry, and cleaves this polyprotein into 11 individual proteins which form the complex that translates full viral RNA and allows for reproduction of the viral genome and the production of mature virions [43]. Through interactions with HDAC2, NSP5 may mediate the epigenetic reprogramming of infected cells, which in pAPCs may include suppression of MHC II expression. Indeed, epigenomic changes are required for SARS-CoV-2 reproduction [44,45], and similar epigenetic reprogramming is known to suppress MHC II expression in Middle East respiratory syndrome–related coronavirus (MERS-CoV) [46]. Therefore, in this study we tested the hypothesis that SARS-CoV-2 NSP5 inhibits MHC II expression through interactions with HDAC2.

Results

SARS-CoV-2 NSP5 Downregulates MHC II in Professional Antigen Presenting Cells
SARS-CoV-2 NSP5 is the main viral protease and plays an essential role in viral infection and pathogenesis [47,48]. As main proteases are required for the processing of coronavirus polyproteins [49], deletion of NSP5 or its inactivation abrogates productive infection by SARS-CoV-2 [43,49,50] To assess the effects of NSP5 on the MHC II antigen presentation system, primary human moDCs were transduced with either empty or NSP5-

expressing lentiviral vectors bearing a zsGreen marker. Flow cytometry was used to quantify surface expression of total MHC II on transduced (zsGreen+) moDCs (**Figure 1A-B, S1**), with NSP5 expression reducing the cell surface expression of MHC II to an extent similar to that observed in SARS-CoV-2 patients (30-50%, **Figure 1C-D**) [25–27]. This downregulation was not a general suppression of the MHC II presentation system, as expression of the co-stimulatory molecule CD86 was not affected by NSP5 expression (**Figure 1E**). Some pathogens such as Human cytomegalovirus reduce their immunogenicity by diverting intracellular trafficking such that MHC II molecules fail to reach the cell surface [51]. To test this possibility, we transduced J774.2 macrophages with NSP5-expressing or empty vectors, labeled the plasma membrane with wheat-germ agglutinin, followed by labeling for total cellular MHC II. Three-dimensional reconstructions of these cells were used to differentiate between cytosolic/vesicular MHC II and cell-surface MHC II, comparing both non-transfected (zsGreen-negative) to transfected (zsGreen-positive) cells in the same samples (**Figure 1F**). Quantitation of these micrographs revealed no changes in the portion of MHC II localized to the cell surface versus intracellular vacuoles (**Figure 1G**) but did identify the same decrease in MHC II expression that was observed with flow cytometry (**Figure 1D,H**), indicating that NSP5 does not affect the trafficking of MHC II to the cell surface, but rather decreases its overall expression. Moreover, this effect was only observed in NSP5-expressing cells, indicating that this effect is cell-intrinsic and not due to NSP5-induced changes in the expression of cytokines or other secreted factors.

Figure 1: NSP5 Suppresses MHC II at the Protein Level in Primary Monocyte-Derived Human Dendritic Cells. moDCs were transduced with lentiviral vectors lacking a transgene (empty vector) or bearing NSP5, with both vectors containing an IRES-zsGreen marker. **A-B)** Flow cytometry dot plots showing cell surface MHC II expression and transduction (zsGreen+) of moDCs transduced with an empty vector control (A) or NSP5-expressing lentiviral vector (B). **C)** Histogram of cell surface MHC II expression levels on moDCs transduced with an empty vector (red) or NSP5-expressing vector (cyan). **D-E)** Quantification of cell surface MHC II (D) and CD86 (E) in moDCs transduced with either an empty (Empty) or NSP5-expressing (NSP5) lentiviral vector. MFI is normalized to the MFI of the untransduced cells in the Empty-vector sample. **F)** Z-slice through a macrophage stained for the plasma membrane and MHC II, showing the segmentation of vesicular/cytosolic versus surface (membrane) MHC II. Scale bar is 10 μm. **G-H)** Quantification of the fraction of MHC II on the plasma membrane (G) and total cellular MHC II (H) in macrophages that have been transfected either with an empty or NSP5-expressing vector, comparing non-transfected (zsGreen-negative) to transfected (zsGreen-positive) cells in both conditions. Data is representative of, or quantifies a minimum of 3 independent experiments, * = p < 0.05; n.s. = p > 0.05 compared to non-transfected empty vector, Kruskal-Wallace test with Dunn Correction.

NSP5 Suppresses CIITA and MHC II Transcription

Next, the subcellular localization of NSP5 was determined to identify potential mechanisms accounting for the downregulation of MHC II. Quantitative microscopy of HeLa cells expressing NSP5-FLAG, the ER marker KDEL-GFP, the Golgi marker GalT-mCherry, and with the nuclei stained with Hoechst, determined that approximately half of the cellular NSP5 was localized to the nucleus, with the remainder associated with the ER

(**Figure 2A-B**). The nuclear localization of NSP5 was confirmed by blocking importin-mediated nuclear transport (**Figure 2C-D**). While localization to the ER is consistent with the known role of NSP5 in forming the viral replication complex [52], the role for nuclear NSP5 remains unclear. Unexpectedly, bioinformatic analysis of NSP5 failed to identify either a classical or bipartite nuclear localization signal, or a nuclear export signal. This suggests that NSP5 may be carried into the nucleus through interactions with other cellular proteins, similar to the hepatitis delta antigen [53].

Figure 2: NSP5 Suppresses CIITA and MHC II expression. A-B) Fluorescent z-projection (A) and Manders colocalization analysis (B) quantifying the portion of NSP5 colocalized with the nucleus of HeLa cells co-transfected with NSP5-FLAG (yellow), GalT-mCherry (Golgi/magenta), KDEL-eGFP (ER/cyan), and stained with Hoechst (DNA/grey). Manders colocalization analysis compares the fraction of NSP5 colocalized with the nucleus, ER and Golgi (Manders) to the Manders ratio from the same image when the NSP5 image was randomized (Random). **C-D)** Fluorescent z-projections (C) and quantification (D) of the fraction of NSP5 in the nucleus of vehicle-treated (DMSO) versus ivermectin-treated HeLa cells expressing NSP5-FLAG (yellow) and stained for DNA with Hoechst (cyan). **E)** RT-qPCR quantification of RFX5, CIITA, and MHC II mRNA levels in moDCs transduced with empty or NSP5-expressing lentiviral vectors. **F)** Quantification of the promoter activity of the CIITA pI, CIITA pIII/IV, and MHC II promoters using a dual-luciferase assay in RAW 264.7 macrophages co-transfected with empty or NSP5-expressing lentiviral vectors. **G)** Quantification of MHC II promoter activity in RAW 264.7 macrophages co-transfected with CIITA and NSP5. (-) indicates the sample was transfected with empty vector rather than CIITA or NSP5. Scale bars are 10 μm. Images are representative of a minimum of 30 cells captured across 3 independent experiments. n = minimum of 3. * = $p < 0.05$; n.s. = $p > 0.05$, compared to Random (B), DMSO (D), or Empty Vector (E-F), paired t-test (B) or Mann-Whitney test (D, E, F).

The nuclear localization of NSP5 suggests that downregulation of MHC II occurs via a transcriptional mechanism. To test this hypothesis, we used RT-qPCR to measure the mRNA levels of MHC II, as well as RFX5 and CIITA—the two transcription factors which act as the master regulators of MHC II expression. Interestingly, while RFX5 expression was unchanged, NSP5 significantly downregulated the expression of CIITA and MHC II relative to the levels normally found on resting moDCs (**Figure 2E**). In humans, CIITA is transcribed from three separate promoters: pI—which drives expression in myeloid cells, pIII—which drives expression in lymphoid cells, and pIV—which drives IFN-γ-induced expression in non-immune cells such as epithelia [54,55]. To assay the activity of the MHC II and CIITA promoters, we constructed dual-luciferase reporters of the MHC II promoter, the CIITA pI promoter, and the CIITA pIII/pIV promoters, followed by quantification of NSP5's effect on these promoters' activity in macrophages. NSP5 expression strongly suppressed both the MHC II promoter and the CIITA pI promoter, whereas the CIITA pIII/IV promoters were minimally active, producing insufficient signal to observe any effect of NSP5 (**Figure 2F**). MHC II transcription is dependent on CIITA, therefore this suppression of MHC II expression may be due to NSP5-dependent repression of CIITA expression, or alternatively, may be a product of NSP5 suppression of both the CIITA and MHC II promoters. To differentiate between these possibilities, we quantified MHC II promoter activity in cells ectopically expressing CIITA and NSP5 (**Figure 2G**). CIITA expression greatly increased MHC II promoter activity, with co-expression of NSP5 having no effect on MHC II promoter activity in the presence of ectopically expressed CIITA. These data indicate that NSP5 suppresses the transcription of CIITA, and the resulting absence of CIITA then limits MHC II expression.

NSP5-Mediated Suppression of MHC II Expression is Dependent on HDAC2

MHC II expression is regulated by CIITA through several mechanisms. Induction of MHC II expression begins with CIITA binding to distal enhancers located several kilobases 5′ to the MHC II promoter [56], followed by acetylation of histones and transcription factors within the core MHC II promoter protein complex by CIITA. This acetylation induces the formation of an enhanceosome complex comprised of CIITA, RFX5, CREB and NF-Y, wherein CIITA activates TAF family transcription factors via its intrinsic protein acetylation and kinase activity, thus initiating MHC II transcription [57,58]. CIITA expression is induced by IFNγ through activation of the transcription factor IRF1, and via toll-like receptor (TLR) and IL-1 family cytokines via activation of the transcription factor IRF3 [32,59,60], and is negatively regulated by HDAC2-mediated promoter deacetylation [61]. Critically, SARS-CoV-2 NSP5 can interact with HDAC2, suggesting that NSP5 utilizes HDAC2 to silence the CIITA promoter [42]. Consistent with this model, HDAC2 co-precipitated with FLAG-tagged NSP5 in an anti-FLAG immunoprecipitation (**Figure 3A**). Knockdown of HDAC2 (**Figure 3B**) had a profound restorative effect on CIITA and MHC II mRNA levels, with HDAC2 knockdown not only reversing – but increasing above baseline – expression of both genes (**Figure 3C**). NSP5 is comprised of a proteolytic domain formed by the interface of the globular A and B domains, which positions two key catalytic residues (H41 and C145) in a binding cleft formed between the two domains. A C-terminal B' chain folds over this cleft, coordinating both the substrate and three water molecules within the active site [62]. We inactivated the catalytic site by generating NSP5^{H41A} and NSP5^{C145S} point mutants, and deleted the proteolytic (NSP5$^{\Delta1-192}$) and B' domains (NSP5$^{\Delta199-306}$, **Figure 3D**). While our ectopically expressed NSP5 displayed proteolytic activity against a FRET cleavage reporter containing the NSP4/NSP5 cleavage site—known to be a target of NSP5 [63]—both point mutants lost this proteolytic activity (**Figure S2A-G**).The deletion mutants were unstable and could not be expressed at levels amenable for immunoprecipitation, but immunoprecipitation of the point mutants revealed that both retained some binding capacity for HDAC2 (**Figure 3E**), and retained their ability to suppress CIITA and MHC II promoter activity (**Figure 3F**).

Figure 3: NSP5 Requires HDAC2 to Suppress MHC II Expression. A) Co-immunoprecipitation of HDAC2 with NSP5-FLAG from primary human moDCs transduced with empty or NSP5-FLAG-expressing lentiviral vectors. **B)** Expression of HDAC2, as quantified by immunoblotting, in moDCs that are untreated (UT) or treated with either scrambled (siScrm) or HDAC2-targeting (siHDAC2) siRNAs. HDAC2 protein level is quantified relative to GAPDH protein levels in the same lysates. **C)** RT-qPCR quantification of the impact of scrambled (siScrm) or HDAC2-targeting (siHDAC2) siRNA on CIITA and MHC II mRNA levels in primary human moDCs transduced with empty or NSP5-expressing lentiviral vectors. **D)** Protease-inactivating (H41A and C145S) point-mutants, and deletion of the catalytic (A/B, Δ1-192) and ligand-stabilizing (B', Δ199-306) domains were generated to assay the roles of these sites in NSP5 activity. **E)** Immunoprecipitation of HDAC2 with NSP5[H41A] and NSP5[C145S] mutants. **F)** Dual-luciferase assay quantification of CIITA pI and MHC II promoter activity in RAW264.7 macrophages that were either untransfected (Untrans.), or co-transfected with the CIITA or MHC II luciferase constructs plus either the empty vector (Empty), or with one of wild-type NSP5 (WT), NSP5[H41A], or NSP5[C145S] vectors. Data is normalized to Empty. n = 5, * = p < 0.05; n.s. = p > 0.05, Kruskal-Wallis test with Dunn correction.

NSP5 Induces Deacetylation of the CIITA and MHC II Promoters

Given the dependence of NSP5 on HDAC2 for its suppressive effect on MHC II transcription, it is likely that NSP5 is modulating protein acetylation at the CIITA or MHC II promoters. Because lentiviral transduction of primary human macrophages is inefficient (30-50% transduction), FRET microscopy was used in lieu of conventional ChIP to quantify total acetylation at the MHC II and CIITA promoters. This assay measures the transfer of excitation energy from a Cy3-labeled anti-acetyl-lysine antibody to ATTO647-N labeled FISH probes specific to the MHC II promoter plus 3 kb of 5′ enhancer regions, the CIITA-pI promoter plus 1.5 kb of 5′ sequence, and the region containing the CIITA pIII and pIV promoters (CIITA pIII/IV), with the zsGreen marker of the lentiviral vector used to restrict analysis to transduced cells. This approach has the added advantage that both histone acetylation and acetylation of promoter-bound transcription factors are measured, with acetylation of transcription factors such as CIITA known to enhance their activity [64]. Primary human macrophages were transduced with empty or NSP5-expressing vectors and treated with either scrambled or HDAC2-targeting siRNA. Acetylated-lysine staining was concentrated in the nucleus, with weaker staining present in the cytosol, with one or two FISH probes in-focus within each nucleus (**Figure 4A**). Neither NSP5 expression, nor HDAC2 depletion, altered the quantity or distribution of cellular acetyl-lysine staining (**Figure 4B-C**), indicating that neither NSP5

expression nor HDAC2 knockdown globally affected lysine acetylation. In untransduced cells, and in cells transduced with an empty vector, a significant FRET signal could be observed at the CIITA pI and MHC II promoters, while a much weaker FRET signal was detected at the CIITA pIII/IV promoter (**Figure 4D-F**), consistent with the pI promoter driving CIITA expression in myeloid cells. Critically, ectopic expression of NSP5 significantly reduced acetylation at all three promoters, with HDAC2 knockdown restoring promoter acetylation in NSP5-expressing cells (**Figure 4D-F**).

It seemed unusual that a virus whose closest known wild ancestor is found in bats (BANAL-20-236) would have activity against human MHC II and CIITA promoters [65,66]. This activity may suggest that NSP5 has undergone significant evolution following the zoonotic transmission of SARS-CoV-2 to humans, or alternatively, that HDAC2 is highly conserved between bats and humans. Phylogenetic and amino acid conservation analysis revealed that the latter is the case – HDAC2 is strikingly conserved across vertebrates, with most residues completely conserved between humans and a range of vertebrates known to be infected by SARS-CoV-2 or to frequently contact humans (**Figure 4G**). A similar degree of conservation is also observed across all major clades of vertebrates (**Figure S3**). In contrast, NSP5 is modestly conserved across the major coronavirus clades, but shows high conservation of its catalytic site, as well as complete conservation between SARS-CoV, SARS-CoV-2, and bat coronavirus BANAL-20-236 (**Figure 4H**).

NSP5 Targets the CIITA Promoter via Interactions with IRF3

As NSP5 is a protease, and proteolytic cleavage of some HDACs are known to promote their activity [67], we next assessed whether NSP5 was capable of cleaving HDAC2. Immunoblot analyses were performed to compare differences in HDAC2 protein levels in cells transfected with either NSP5-expressing or empty vectors. Despite a predicted NSP5 cleavage site [42], we observed no change in HDAC2 protein levels or mass in NSP5-expressing cells (**Figure 5A**). While this predicted cleavage site should produce an 11 kDa decrease in the mass of HDAC2, HDACs are often cleaved closer to the C-terminus, resulting in cleavage products barely distinguishable from the uncleaved protein. As such, we constructed an intramolecular FRET cleavage probe comprised of an mVenus-HDAC2-mTurquoise2 fusion protein. Intact, this probe has high FRET efficiency between the mVenus and mTurquoise2 fluorophores, with cleavage resulting in separation of the fluorophores and loss of the FRET signal (**Figure S2H-J**). Consistent with the immunoblotting results, no change in FRET efficiency was observed in NSP5-transfected versus empty-vector transfected cells in either the cytosol or nucleus (**Figure 5B**), confirming that NSP5 does not modulate HDAC2 activity through proteolytic cleavage. We next quantified the subcellular distribution of HDAC2 and the NSP5 mutants generated in Figure 3 and found that inactivation of the NSP5 catalytic site had no impact on the distribution of either protein (**Figure 5C-D**). Although poorly expressed, the rare cells expressing NSP5 deletion mutants lacking the proteolytic (NSP5$^{\Delta 1-192}$) or B′ (NSP5$^{\Delta 199-306}$) domains displayed a hyper-nuclear localization compared to wild-type, but also had no impact on the distribution of HDAC2, indicating that HDAC2 unlikely to be the protein used by NSP5 to gain access to the nucleus (**Figure 5C-D**). Naik *et al* demonstrated that NSP5 can interact with IRF3, and IRF3 is known to bind to the CIITA PI promoter, suggesting that NSP5 may deliver HDAC2 to the CIITA promoter via interactions with IRF3 [68–70]. Using FISH-FRET analysis of A549 cells—a type II human alveolar lung epithelial cell line—we found that IFNγ stimulation increased CIITA PI, PIII/IV and MHC II promoter acetylation, and as expected, NSP5 expression inhibited promoter acetylation (**Figures 5E-G, S4)**. Critically, this suppression of promoter acetylation by NSP5 was largely reversed by an IRF3-targeting siRNA, consistent with a model in which NSP5 delivers HDAC2 to the CIITA promoter via interactions with IRF3 (**Figures 5E-G, S5**).

Figure 4: NSP5 Modulates Acetylation of the CIITA and MHC II promoters. A) Representative micrograph of primary human macrophages transduced with empty or NSP5-expressing lentiviral vectors. Cells were stained for acetyl-lysine (Lys^{ac}, cyan), ATTO647N-labeled MHC II FISH probes (yellow), with the FRET signal between the Lys^{ac} and FISH probes within the insert shown in the FRET panel. Scale bars are 10 μm. **B-C)** Quantification of the nuclear:cytosolic acetyl-lysine distribution (B) and total cellular acetyl-lysine content (C) in primary human macrophages transduced with empty or NSP5-expressing lentiviral vectors and treated with scrambled (siScrm) or HDAC2-targeting (siHDAC2) siRNAs. "Untransduced" quantifies macrophages which did not take up the empty or NSP5-expressing

lentiviral vector. **D-F)** Quantification of the acetylation of the CIITA pI (D), CIITA pIII/pIV (E) and MHC II (F) promoters, as quantified by the FRET efficiency between Cy3-labeled anti-acetyl-lysine staining and ATTO647N-stained FISH probes in the same cells used for quantification of panels B and C. **G-H)** Domain structure, amino acid conservation (Shannon Entropy, H (X)), and phylogenetic trees for (G) HDAC2 across a range of vertebrate species which humans frequently contact or which are possible vectors of SARS-CoV2, and (H) representative species of the four genera of coronaviruses. HDAC2 consists of four major domains, a homodimerization (HD) domain, deacetylase domain, an IAC (E/D)E motif, and a coil-coil domain which mediate interactions with other transcription factors. NSP5 contains a bipartite protease domain (A/B) with two critical catalytic residues (H41 and C145), and a B' C-terminal domain which stabilizes ligands in the protease domain. Data are representative of (A) or quantifies (B-F) three independent experiments. * = p < 0.05; n.s. = p > 0.05, Kruskal-Wallis test with Dunn correction. Flat bars indicate the statistical significance for all groups beneath the bar; legged bars indicate the statistical significance between the groups below the legs.

Discussion

In this study we demonstrate that SARS-CoV-2 NSP5 suppresses expression of CIITA across a range of professional and non-professional antigen presenting cells, thereby suppressing MHC II expression. This suppression of CIITA expression occurs via a pathway in which NSP5 delivers HDAC2 to the CIITA promoter via interactions between NSP5 and promoter-bound IRF3. HDAC2 then deacetylates histones at the CIITA promoter, decreasing CIITA expression, thereby blocking MHC II expression (**Figure S5**). These findings explain the decreased MHC II expression observed COVID-19 patients and in *in-vitro* infection models [25–27]. This suppressive mechanism occurred across multiple types of professional APCs (human moDCs, human macrophages, and a mouse macrophage cell line) and in human non-professional APCs (type II alveolar epithelial cells). As SARS-CoV-2 infects pAPCs including DCs, macrophages, and B cells—as well as non-professional APCs in the lung—this suppression of MHC II has the potential to dramatically decrease the activation of CD4$^+$ T cells in COVID-19 patients [6–8]. Such a loss of CD4$^+$ T cell activity likely contributes to the shorter-lived humoral immunity and propensity for reinfection of COVID-19, and may contribute to the aberrant formation of memory and effector CD4$^+$ T cell subsets in individuals with severe disease [24,71–74].

The suppression of MHC I and MHC II expression, or removal of these proteins from the cell surface, is a commonly employed viral immunoevasion strategy. For MHC I, mis-directing or re-internalizing MHC I after its translation are the predominant mechanisms used to limit immunogenicity of infected cells [75]. For example, HIV-1 Nef mediates the AP-1-dependent endocytosis of MHC I and its sequestration in a Golgi-proximal compartment, thus limiting cytotoxic T cell activity against infected cells [76,77]. Similarly, SARS-CoV-2 utilizes ORF8 to direct MHC I into lysosomes for degradation, and uses ORF6 to suppress MHC I expression by downregulating expression of the transcription factor CITA/NLRC5 [18,22]. Influenza A and B also downregulate surface MHC I via proteasomal degradation or via endocytosis and sequestration in cytosolic vacuoles, respectively [78]. Downregulation of MHC II is rarer, as only viruses that infect APCs have the potential to exhibit this activity. HSV-1 reduces cell surface levels of MHC II by directing MHC II into multivesicular bodies [79,80]. HIV-1 Nef, through accumulation of αβIi complexes in intracellular vesicles, can also suppress trafficking of peptide-loaded MHC II to the cell surface [51]. Human cytomegalovirus protein US2 degrades HLA-DR-α and DM-α chains, while the US3 protein competes with the invariant chain for binding to MHC II α/β complexes, thereby restricting MHC II intracellular trafficking [81,82]. Transcriptional downregulation of MHC II is also observed, often through targeting of CIITA. Similar to what we have reported in this study, Kaposi sarcoma-associated herpesvirus also targets the IRF3 binding site in the CIITA promoter—but unlike SARS-CoV-2—this virus expresses a viral IRF3 which is inhibitory to CIITA expression [69]. Epstein-Barr virus impairs CIITA expression in B cells by suppressing the activity of E47 and PU.1, thereby preventing these transcription factors from binding the pIII promoter [83,84]. The coronavirus MERS-CoV was shown to transcriptionally downregulate both MHC I and MHC II through an undefined epigenetic mechanism [46]. Given the epigenetic basis of SARS-CoV-2's suppression of CIITA and MHC II, it is possible that the downregulation of MHC II by MERS-CoV is mediated by the same NSP5-mediated mechanism that we identified in this study, and that this activity may be conserved across pAPC-infecting coronaviruses.

Figure 5: NSP5 Suppresses CIITA Promoter Activity via IRF3. A-B) No evidence of HDAC2 cleavage by NSP5 could be detected by (A) HDAC2 immunoblotting of primary human moDCs transduced with empty or NSP5-expressing lentiviral vectors (UT = untransduced cells), or by (B) changes in intermolecular FRET of an mVenus-HDAC2-mTouquise2 fusion protein in HeLa cells co-transfected with empty or NSP5-expressing lentiviral vectors. **C-D)** Fluorescent micrographs (C) and quantification (D) of NSP5 (Yellow) and HDAC2 (Magenta) nuclear localization in HeLa cells transfected with wild-type (WT) or mutant NSP5. The nucleus has been stained with DAPI (Cyan). Scale bar is 10 μm. **E-G)** Quantification of FISH-FRET at the CIITA pI (E), CIITA pII/IV (F) and MHC II (G) promoters in NSP5-transfected A549 cells treated with IFNγ and a non-targeting (Scrm) or IRF3-depleating siRNA. Images are representative of a minimum of 30 cells imaged over 3 independent experiments. Data is presented as mean ± SEM (B) or quartiles, n = 3, * = p < 0.05; n.s. = p > 0.05 compared to Empty (B), WT (D), or the indicated groups (E-G), Kruskal-Wallis test with Dunn correction.

The proteolytic activity of coronaviral NSP5 is known to be important for its immunosuppressive activity. Porcine deltacoronavirus NSP5 suppresses host antiviral IFN signaling by proteolysis of the host antiviral proteins NF-kappa-B essential modulator [85], STAT2 [86], and mRNA decapping protein 1a [87]. Previous computational studies also predicted a SARS-CoV-2 NSP5 cleavage site (^{380}VQMQ|AIPE^{387}) in HDAC2, with cleavage removing an 11.6 kDa C-terminal fragment containing the HDAC2 NLS [42]. This cleavage was proposed to reduce HDAC2 localization to the nucleus and thus limit its ability to attenuate inflammatory responses. Surprisingly, we did not observe evidence that NSP5 cleaves or inactivates HDAC2, with no cleaved HDAC2 (~44 kDa) appearing in our immunoblotting experiments, nor with any cleavage detected in our intramolecular FRET assay of either cytosolic or nuclear-localized HDAC2. This is consistent with the work of Naik *et al.* who observed a similar non-proteolytic interaction of NSP5 with HDAC2 and IRF3 [68]. Further demonstrating that cleavage is not required for this

interaction, the catalytically inactive NSP5^{H41A} and NSP5^{C145S} point mutants maintained their interaction with HDAC2 and had the same suppressive effect on CIITA pI and MHC II promoter activity as wild-type NSP5. Why HDAC2 is not cleaved by NSP5, despite containing an accessible SARS-CoV-2 NSP5 consensus sequence, is unclear. However, non-proteolytic functions of SARS-CoV-2 NSP5 have been reported, including inducing SUMOylation of MAVS to promote inflammation [88] and suppressing activation of RIG-I by preventing formation of antiviral stress granules via interactions with G3BP1 [89]. Additionally, coronavirus-mediated epigenetic reprogramming has been reported previously, with both SARS-CoV and MERS-CoV altering inhibitory and activating histone modifications across a range of genes [90]. This regulation is targeted, with these viruses able to selectively activate or repress expression of specific genes. Consistent with this observation, we found that SARS-CoV-2 NSP5 selectively inhibited expression of MHC II and CIITA without affecting expression of CD86 and RFX5, or globally suppressing acetylation.

HDAC2 is a promising target for treatment of SARS-CoV-2 infection. Indeed, epigenetic changes consistent with HDAC2 activity drives the cytokine storm that is responsible for much of the pathology seen in SARS-CoV-2 patients, with the degree of epigenetic change correlated to disease severity [45,91–93]. NSP5 is thought to drive much of the cytokine response in the infected lung via HDAC2, and indeed, HDAC2 is required for the upregulation of many pro-inflammatory molecules in endothelial and myeloid cells undergoing SARS-CoV-2 infection [94–96]. HDAC inhibitors exhibit potent anti-inflammatory effects and therefore may antagonize many of the inflammatory pathways activated by SARS-CoV-2 [97,98]. Outside of inflammation, HDACs positively regulate ACE2 expression, and consequentially, HDAC2 inhibitors decrease SARS-CoV-2 entry and replication through reduced ACE2 expression [97,99,100]. Moreover, HDAC inhibitors downregulate pro-inflammatory cytokines, reduce lung fibrosis, prevent viral entry into the central nervous system, and decrease neurological damage [98,101,102]. Thus, HDAC2 inhibition may improve patient outcomes through multiple mechanisms in addition to restoration of MHC II expression. Critically, targeting a host protein would reduce the likelihood of SARS-CoV-2 evolving resistance to this treatment.

Understanding how SARS-CoV-2 modulates the MHC II antigen presentation pathway provides an important insight into the immunoevasion tactics used by this virus and may help to provide directions for the design of future COVID-19 vaccines or therapeutics. This study identifies one mechanism through which SARS-CoV-2 suppresses MHC II expression. Furthermore, our data indicate that SARS-CoV-2 may utilize NSP5 to modulate a broad array of immune responses via targeting HDAC2, which in addition to its effects on MHC II expression identified herein, has also been identified as a positive regulator of the cytokine storm that underlies much of the pathology of COVID-19. Indeed, HDAC2 inhibition has been proposed as a therapeutic approach for COVID-19, with the findings from this study further validating HDAC2 inhibitors as potentially valuable treatments for this disease [101].

Materials & Methods

Materials

The TGN46-GFP and KDEL-mRFP plasmids were gifts from Dr. Sergio Grinstein (Hospital for Sick Children, Toronto, Canada) The pMD2.G (plasmid 12259) and pDR8.2 (plasmid 12263) packaging vectors and pcDNA3-myc-CIITA (plasmid 14650) were purchased from AddGene. All DNA primers and synthesized genes were from IDT (Coralville, Iowa), and the sequences for all primers used in this study can be found in Supplemental Table S1. Tissue culture medium, fetal bovine serum (FBS), and trypsin were from Wisent (St. Bruno, Canada). Recombinant cytokines were from Peprotech (Cranbury, NJ). All cell lines and Lympholyte-poly were from Cedarlane labs (Burlington, Canada). Polybrene, ivermectin, and 100K Amicon centrifugal filters were purchased from EMD Millipore Corp (USA). CD14 Positive Cell Selection Kit,

FcBlock, and anti-DYKDDDDK Tag (L5) were from BioLegend (San Diego, California). The #1.5 thickness coverslips and 16% paraformaldehyde (PFA) were from Electron Microscopy Sciences (Hatfield, PA). FDA-traceable PLA filament was purchased from Filaments.ca (Mississauga, Canada). RNeasy Mini Kit was from Qiagen (Germantown, MD). Permaflor, WGA-Alexa Fluor 647, DAPI, Hoescht, HALT protease/phosphatase inhibitors, Dithiobis[succinimidyl propionate] and dithio-bismaleimidoethane, were purchased from ThermoFisher Canada (Mississauga, Canada). The suppliers and all antibodies and labeling reagents used in this study can be found in Supplemental Table S2. Atto647N NT Labeling Kit was from Jena Biotech (Jena, Germany). Accell cell-penetrating SMARTpool scrambled and HDAC2-targeting siRNAs were from Horizon Discovery (Cambridge,

UK). Instagene, iScript Select cDNA Synthesis Kit, 4%-20% SDS-PAGE gels, SsoFast EvaGreen Supermix, and all protein blotting reagents/gels were from BioRad Canada (Mississauga, Canada). Renilla luciferase internal control vector pRL-TK and the dual luciferase reporter assay kit was from Promega (Madison, WI). Phusion PCR enzyme, all restriction enzymes, HiFi Gibson Assembly Kit, and T4 DNA ligase were from NEB Canada (Whitby, Canada). The pLVX-zsGreen lentiviral vector and Retro-X Universal Packaging System was purchased from Takara Bio (San Jose, California). All lab plasticware, PolyJet and GenJet transfection reagent, and DNA isolation kits were from FroggaBio (Concord, Canada), and all laboratory chemicals were from Bioshop Canada (Burlington, Canada).

NSP5 Cloning and Retroviral Packaging

The RNA sequence NSP5 from the Wuhan SARS-CoV-2 strain [103] was synthesized such that a start and stop codon were added to the 5' and 3' end of the NSP5 sequence, along with 20 bp of homology to the pLVX-zsGreen vector at the EcoRI restriction site. The resulting NSP5 sequence was cloned into EcoRI-digested pLVX-zsGreen by Gibson assembly. Point mutants were generated by amplifying the entirety of this original vector with phosphorylated primers that incorporate the point mutation in the first base pair of the forward primer, while deletion mutants were generated by amplifying the vector from either side of the desired deletion with phosphorylated primers. After amplification, the parental plasmid was removed by DpnI digestion and the amplicons circularized with T4 DNA ligase. All primers used for RT-qPCR can be found in Supplemental Table S1. To produce pseudotyped lentivirus containing empty vector or NSP5, 3×10^6 HEK293T cells were grown in 75 cm2 tissue culture flasks, then transfected with PolyJet transfection reagent (500 μL complex containing 4 μg pMD2.G and 10 μg pDR8.2 packaging vectors, and 10 μg of pLVX expression vector). Following transfection, cells were incubated at 37°C/5% CO_2 for 18 hr at which point the media was exchanged for 8 mL of DMEM supplemented with 10% FBS and returned to the incubator for 48 hr. The media was transferred to a sterile 50 mL conical centrifuge tube and topped up to 20% FBS, then centrifuged at 4000×g for 5 minutes, and the supernatant filtered with a 0.2 μm syringe filter into a new 50 mL conical tube. The pseudotyped lentivirus was then concentrated using a 100 kDa centrifugal filter unit at 4000×g at 4°C for 45 min per 15 mL of filtrate. Concentrated pseudotyped lentivirus was aliquoted and stored at -80 °C and thawed at room temperature prior to use.

Human Macrophage and Dendritic Cell Culture, Transduction, and siRNA Treatment

The collection of blood and cells from healthy donors was approved by the Health Science Research Ethics Board of the University of Western Ontario and was performed in accordance with the guidelines of the Tri-Council policy statement on human research. Blood was drawn into heparinized vacuum collection tubes, layered on an equal volume of Lympholyte-poly and centrifuged at 300×g for 35 min at 20 °C. The top band of peripheral blood mononuclear cells was collected and washed once (300×g, 6 min, 20 °C) with phosphate-buffered saline. For dendritic cell differentiation, a CD14 selection kit was used to isolate monocytes according to manufacturer's instruction. The

selected CD14+ cells were cultured in RPMI-1640+10% FBS and 1% antibiotic–antimycotic solution with GM-CSF (100 ng/ml) and IL-4 (100 ng/ml) for 4 days yielding immature monocyte-derived DCs (moDCs). To produce macrophages, selected CD14+ cells were cultured in the presence of M-CSF (10 ng/ml) for 6 days. ~1×10^6 moDCs or macrophages were centrifuged with 20 transducing units of lentiviral vectors per cell at 800×g at 32°C for 90 min with Polybrene (10 μg/ml). After centrifugation, the cells were incubated at 37°C with 5% CO_2, and 8 hr later fresh media plus cytokines were added and the cells incubated for 72 hr. For siRNA knockdown, 1 μM of Accell cell-permeant siRNA was added to the cells and incubated at 37°C with 5% CO_2 for 72-96 hr.

Cell Line Culture and Transfection

HeLa, RAW264.7, and J774.2 were cultured in DMEM supplemented with 10% FBS, A549 cells were cultured in Ham's F12 supplemented with 10% FBS and were grown at 37°C/5% CO_2 incubator. Cells were split 1:10 upon reaching >80% confluency by either scraping cells into suspension (RAW and J774) or by trypsinization, diluting in fresh medium, and replating in a new tissue culture flask. Genjet DNA transfection reagent was used to transfect plasmids into HeLa cells, as per the manufacturer's instructions. Briefly, for each well in a 6-well plate, 1 μg of DNA was diluted into 50 μl of serum-free DMEM, followed by 3 μl of Genjet reagent. The resulting mixture was incubated for 10 min at room temperature, and then added dropwise to the HeLa cells. Cells were incubated for at least 18 hr at 37 °C in a 5% CO_2 before collection. A549 cells were transfected using Lipofectamine 3000, transfecting 1 μg of DNA per well of a 12-well plate, using 2 μL of P2000 and 3 μL Lipofectamine per transfection. A Neon transfection system (Thermo Fisher Scientific AG) was used to transfect plasmids into RAW264.7 cells according to the manufacturer's instructions. Briefly, 1×10^6 cells were resuspended in 10 μL of buffer R containing 5 μg of plasmid DNA and electroporated using a single 20 ms pulse at 1680V. Lentiviral transductions were used for transducing plasmids into J774.2 cells as described above.

Flow Cytometry

HLA-DR and CD86 expression on the surface of moDCs was measured following 72 hrs transduction with NSP5-ZsGreen or empty-ZsGreen pseudotyped lentivirus and subsequent 24 hr stimulation with 100 ng/uL IFN-γ and incubated at 37°C/5% CO_2. After stimulation 3×10^5 cells per condition were washed with PBS and blocked for 30 min on ice with FcBlock. Cells were stained on ice for 30 min using eFluor670-FVD and conjugated primary antibodies as indicated in Supplemental Table S2. Cells were fixed with 4% PFA in PBS for 15 min then washed with PBS. Expression levels were measured using a FACSCanto (BD), live moDCs were identified based on FVD-eFluor780 viability dye staining and forward scatter and side scatter profiles. Singlets were gated on the forward area scatter and forward height scatter profiles (**Figure S1**). For cell sorting, singlets were gated on the forward area scatter and forward height scatter profiles, then transduced cells were identified by a positive zsGreen signal and this population sorted into the receiving tube. Flow cytometry data were analyzed using FlowJo (v10.8). All antibodies, dyes, and dilutions used for flow cytometry can be found in Supplemental Table S2.

Immunoprecipitation and Immunoblotting

Prior to lysis, cells were washed 3× with cold PBS. For immunoprecipitations, proteins were reversibly cross-linked using the ReCLIP method [104]. Briefly, cells were incubated for 1 hr at room temperature in PBS + 0.5 mM dithiobis[succinimidyl propionate] and 0.5 mM dithio-bismaleimidoethane). This medium was aspirated, and crosslinking quenched by the addition of 5 mM L-cysteine in 20 mM Tris-Cl, pH 7.4 for 10 min at room temperature. Cells were suspended in 300 µL of RIPA lysis buffer, and 50 µL pre-washed anti-DYKDDDDK Tag (L5) beads, rotating for 1 hr. Beads were washed with PBS and immunoprecipitated protein eluted using 0.1 M glycine at pH 2.8. For immunoblotting, cells were lysed with 300 µL RIPA buffer supplemented with 1 mM PMSF and Halt protease and phosphatase inhibitor cocktail at the manufacturer's recommended concentration. Proteins were loaded on a 4–15% gradient SDS-PAGE gels and transferred onto PVDF membrane. The membrane was blocked for 5 minutes with EveryBlot Blocking Buffer (BioRad), incubated overnight at 4°C with the desired primary antibodies (Supplemental Table 2), washed 3 × 5 min with TBS-T, incubated with appropriate IR700 or IR800 secondary antibodies, 1:2,500 dilution, for 1 hr at room temperature in TBS-T. The blots were washed 3 × 15 min washes in TBS-T and visualized with an Odyssey CLx (LI-COR Biosciences, Lincoln, Nebraska). Densitometry was performed in ImageJ/FIJI [105,106].

RT-qPCR

Total RNA was isolated from FACS sorted cells transduced with either empty or NSP5-expressing lentiviral vectors using RNeasy Mini Kit as per manufacturer's instructions. Samples were eluted in 30-50 µL of RNAse-free water. RNA concentration and quality were measured using a NanoDrop 1000 Spectrophotometer. cDNA was obtained from total RNA using the iScript Select cDNA Synthesis Kit according to manufacturer's instructions using an equal amount of starting RNA and equal mix of the oligo $(dT)_{20}$ primer mixes. RT-qPCR was performed using SsoFast EvaGreen Supermix with an equal amount of starting cDNA. Reactions were run on a QuantStudio 3 Real-Time PCR System for 40 amplification cycles. Relative expression of genes of interest was calculated using the ΔΔCt method, with GAPDH serving as the reference gene.

Dual-Luciferase Promoter Activity Assay

RAW264.7 cells were seeded at 1×10^6 cells/well in a 12-well plate and transfected as indicated with Luc-HLA-DRA, Luc-CIITA pI/, Luc-CIITA pIII/IV, Renilla luciferase internal control vector pRL-TK, and NSP5-FLAG or pLVX-IRES-ZsGreen using the Neon electroporation system according to the manufacturer's protocol. 72 hr post-transfection, cells were lysed with 1x Passive lysis buffer supplemented with EDTA-free HALT protease inhibitor at the manufacturer's recommended concentration. Dual luciferase assays were performed using a Dual-Luciferase Reporter Assay Kit according to the manufacturer's instructions, with measurements performed on a Cytation 5 luminescence microplate reader. Firefly luciferase readings were relative to Renilla luciferase readings to account for differences in transfection efficiencies and cell count between samples.

Immunofluorescence Microscopy

Cells of interest were seeded at a density of 1000 cells/mm^2 into either 18 mm circular coverslips placed into the wells of a 12-well plate, or into the wells of a custom-printed 15-well imaging chamber [107]. Cells were fixed in 4% paraformaldehyde in PBS for 20 min at room temperature. For plasma membrane staining, cells were stained with 5 µg/mL Alexa Fluor 647-conjugated wheat germ agglutinin for 10 min at 10 °C, then fixed in 4% paraformaldehyde in PEM buffer (80 mM PIPES, 1 mM EGTA, 1 mM $MgCl_2$) for 10 min at 37 °C. If permeabilization was required, fixed cells were treated with permeabilization buffer (PBS + 0.1% triton X-100 + 2.5% BSA); otherwise, cells were blocked with antibody buffer (2.5% BSA in PBS). Anti-FLAG, -MHC II, or -acetyl-lysine were diluted to the concentration indicated in Supplemental Table S1 in antibody buffer and incubated with the cells for 1 hr. Cells were then washed 3 × 15 min with PBS, and then an appropriate secondary antibody added at a 1:1,000 to 1:2,500 dilution in antibody buffer for 1 hr, followed by washing 3 × 15 min with PBS. Samples were either imaged immediately or mounted on a slide using Permafluor before imaging. All incubations and washes were performed at room temperature.

All samples were imaged on a Leica DMI6000B equipped with a Hamamatsu ORCA-Flash4 CMOS camera, fast filter wheels equipped with a Chroma Sedat Quad and custom Fluorescent Resonance Energy Transfer (FRET) filter wheels, operated using Leica LAS-X software. Unless otherwise noted, all cells were imaged using a 100×/1.40 NA objective lens, with Z-stacks acquired with 0.4 µm between slices. Z-stacks were deconvolved in LAS-X using a 10-iteration blinded deconvolution. Images were exported to ImageJ/FIJI for analysis [105,106]. For co-localization studies, the JaCoP plugin was used to calculate the Manders Ratio of NSP5-FLAG and nuclear DAPI or Hoechst staining, or transgene-delineated ER or Golgi markers [108]. To calculate the fraction of NSP5 in the nucleus, a manual region of interest (ROI) was drawn around the nucleus and whole cell and the integrated intensity of each was measured. The fraction of NSP5 in the nucleus was calculated as $\frac{Nuclear\ NSP5}{Total\ NSP}$.

FRET and FISH-FRET were quantified as described below. To calculate the mean fluorescence intensity of MHC-II in transduced macrophages, the background subtracted channel was thresholded to create a binary mask using the default setting in ImageJ, then a sum slices Z-projection was created. A manual ROI was drawn around the whole cell and the integrated intensity was measured. To determine the surface to cytosol ratio of MHC II in transduced macrophages, the background subtracted channels for wheat germ agglutinin and MHC II were thresholded as described above. Then, the image calculator was used to display all overlapping and non-overlapping pixels, representing MHC II on the cell surface and cytosol, respectively. A summed Z-projection was created, and a manual ROI was drawn around the whole cell to measure the integrated density for both surface- and cytosolic-MHC II. The surface-to-cytosol ratio was calculated as $\frac{Surface\ MHC\ II}{Cytosolic\ MHC\ II}$ and the fraction of total MHC II on the membrane was calculated as $\frac{Surface\ MHC\ II}{Surface\ MHC\ II\ +\ Cytosolic\ MHC\ II}$.

Intramolecular FRET

DNA comprised of the human HDAC2 gene with flanking BglII and BamHI restriction sites was synthesized and cloned into pmVenus (L68V)-mTurquoise2 (AddGene #60493) such that a fusion protein of mVenus-HDAC2-mTurqoise2 was produced. To measure NSP5 proteolysis, a codon-optimized DNA sequence encoding the 20 amino acids of the region of the SARS-CoV-2 ORF1ab transcript that codes for the terminal 10 amino acids of NSP4 and initial 10 amino acids of NSP5 (QTSITSAVLQSGFRKMAFPS) was cloned into the BamHI site of the pmVenus (L68V)-mTurquoise2 vector. This site is a known substrate for NSP5 [63]. HeLa cells were then transfected with these constructs with or without NSP5, with mTurquoise2 alone (donor-only sample), with mVenus alone (acceptor-only sample), or with the pmVenus (L68V)-mTurquoise2 vector (positive control). After 24 hours of expression, protein translation was inhibited with 100 μM cycloheximide and the cells incubated for 6 hours to allow NSP5 proteolytic activity to occur in the absence of new protein synthesis. Tiled images of the each well were collected, acquiring the donor (mTurqoise2), acceptor (mVenus) and FRET channels at 40× magnification, using the same excitation and camera settings across all samples. FRET efficiency was then calculated using an implementation of the approach of van Rheenen et al. [109] using a custom-written script in FIJI. In each repeat, the correction values for donor cross-talk (β, donor-only Ida/Idd), donor cross-excitation (α, acceptor-only Idd/Iaa), acceptor cross-excitation (γ, acceptor-only Ida/Iaa), and FRET cross-talk (δ, acceptor-only Idd/Ida), were calculated using donor-only or acceptor-only images and custom-written scripts in FIJI. FRET efficiency (E_A) was then calculated in background subtracted images using the formula:

$$E_A(i) = \frac{\left(I_{dd} - \beta I_{dd} - \left(I_{aa} \times (\gamma - \alpha\beta)\right)\right)}{I_{aa}(1 - \beta\delta)}$$

In ImageJ/FIJI the acceptor-only image was then thresholded, and the "Analyze particles" feature used to generate separate ROIs for each cell in each image, and these ROIs were used to quantify the FRET signal of each cell. The maximum theoretical FRET efficiency for the mTurquise2/mVenus FRET pair is 0.3744, while the maximum theoretical FRET efficiency of the Cy3/ATTO647N FRET pair is 0.3063 [110].

FISH-FRET

To measure levels of acetyl-lysine at the MHC II and CIITA promoters, human DNA was purified from a cheek swab using Instagene as per the manufacturer's instructions. 3500 bp amplicons starting before the promoter and ending at the end of the first exon were amplified with Phusion DNA polymerase as per the manufacturer's instructions using the primers from Supplemental Table S1. Amplicons were gel purified and cloned into EcoRV-digested pBluescript II using a HiFi Assembly Kit as per the manufacturer's instructions. The resulting plasmids were labeled with ATTO647N and fragmented using a ATTO647N NT Labeling Kit, producing fragments averaging 200 nucleotides. Primary human macrophages or A549 cells were plated into the 7.5 mm wells of a customized imaging chamber [107], transduced with lentiviral vectors (macrophages) or transfected (A549 cells) with NSP5 or an empty vector, and treated with siRNA as described above. These cells were stained for immune-FISH as per the protocol of Ye et al. [111], including wells which were left unstained, or stained only with the donor (acetyl-lysine-Cy3) or acceptor (FISH probes). Briefly, cells were fixed, permeabilized and immunostained for acetyl-lysine as described above. After labeling a secondary fixation was performed for 10 min with 2% PFA. FISH probes diluted 1:2,500 in hybridization solution (50% formamide, 10% dextran sulfate, 0.3 M NaCl, 30 mM sodium citrate) and denatured at 75°C for 10 min and then cooled to 37°C. Simultaneously, the cells were incubated at 70°C for 2 min in 70% formamide, 0.3 M NaCl, and 30 mM sodium citrate. The cells were dehydrated by immersing in 75%, 90% and 100% ethanol, 2 min/immersion, then air-dried. The cells were incubated with the denatured FISH probes overnight at 37°C, washed 3 × 5 min with 50% formamide, 0.3 M NaCl, and 30 mM sodium citrate at 42°C, then washed 3 × 5 min with 0.05% Tween 20 in 0.6 M NaCl, and 60 mM sodium citrate. The cells were then washed 3 × 5 min in PBS and immediately imaged.

Tiled images of each well were collected, acquiring the zsGreen, donor, acceptor, and FRET channels at 40× magnification, using the same excitation and camera settings across all samples. FRET efficiency was then calculated as described above. A trained algorithm in Ilastik [112] was used to identify cells based on the acetyl-lysine straining and to classify each cell as zsGreen[+] (transduced) or zsGreen[-] (untransduced). The resulting classifications were exported to FIJI where they were used to assign each FISH probe in the image and the corresponding FRET signal to transduced or untransduced groups. The FRET signal in each sample was then normalized to that observed in the scrambled siRNA-treated, zsGreen[-] nuclei.

NSP5 NLS Analysis

The protein sequence of NSP5 was analyzed for the presence of monotonic and bipartite nuclear localization signals using the default settings on four different prediction algorithms: cNLS Mapper (https://nls-mapper.iab.keio.ac.jp/cgi-bin/NLS_Mapper_form.cgi), 4 state HMM on NLStradamus (http://www.moseslab.csb.utoronto.ca/NLStradamus/), seqNLS (http://mleg.cse.sc.edu/seqNLS/), and in InterProScan (https://www.ebi.ac.uk/interpro/search/sequence/).

Phylogenetic Analysis

Using the protein sequence of human HDAC2 and the NCBI BLASTp tool (https://blast.ncbi.nlm.nih.gov), the protein sequences of HDAC2 from a range of species representing the major vertebrate clades were identified. The same approach, using the protein sequence of NSP5 from SARS-CoV-2 was used to identify NSP5 protein sequences across the four coronavirus genera. These sequences were imported into MEGA XI, and a MUSCLE alignment of the protein sequences was generated [113]. Pairwise distances were then calculated using a Poisson model assuming uniform rates across sites, and maximum likelihood trees were generated using a 500-iteration bootstrapping approach. Per-residue conservation was quantified using the Shannon Entropy calculator on the Protein Residue Conservation Prediction server (https://compbio.cs.princeton.edu/conservation/) [114].

Statistical Analysis

Using GraphPad Prism, a Shapiro-Wilk test was used to determine whether data was parametrically or non-parametrically distributed, and data was then analyzed using an appropriate 2-tailed statistical test, as indicated in the figure legends. Parametric data is presented as mean ± SEM, while non-parametric data is presented as box-and-whisker or violin plots with median and quartiles.

Acknowledgments

We would like to thank Kristin Chadwick from the London Regional Flow Cytometry Facility for her assistance in the design and optimization of the flow cytometry assay. This work was funded by a Canadian Institutes of Health Research (CIHR) Project Grant (PJT-162203) to BH. AL, BHD, and ENB were funded by Canada Graduate Scholarships – Master's (CGSM) from the CIHR. The funding agencies had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interest Statement

The authors have no competing interests.

Preprint Notice

This publication is a scientific preprint and has not undergone peer review.

References

1.   Zhu, N.; Zhang, D.; Wang, W.; Li, X.; Yang, B.; Song, J.; Zhao, X.; Huang, B.; Shi, W.; Lu, R.; et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* **2020**, *382*, 727–733, doi:10.1056/NEJMoa2001017.

2.   Taefehshokr, N.; Taefehshokr, S.; Hemmat, N.; Heit, B. Covid-19: Perspectives on Innate Immune Evasion. *Frontiers in Immunology* **2020**, *11*, 2549, doi:10.3389/fimmu.2020.580641.

3.   Townsend, J.P.; Hassler, H.B.; Wang, Z.; Miura, S.; Singh, J.; Kumar, S.; Ruddle, N.H.; Galvani, A.P.; Dornburg, A. The Durability of Immunity against Reinfection by SARS-CoV-2: A Comparative Evolutionary Study. *The Lancet Microbe* **2021**, *2*, e666–e675, doi:10.1016/S2666-5247(21)00219-6.

4.   Bergwerk, M.; Gonen, T.; Lustig, Y.; Amit, S.; Lipsitch, M.; Cohen, C.; Mandelboim, M.; Levin, E.G.; Rubin, C.; Indenbaum, V.; et al. Covid-19 Breakthrough Infections in Vaccinated Health Care Workers. *N Engl J Med* **2021**, *385*, 1474–1484, doi:10.1056/NEJMoa2109072.

5.   Kuhlmann, C.; Mayer, C.K.; Claassen, M.; Maponga, T.; Burgers, W.A.; Keeton, R.; Riou, C.; Sutherland, A.D.; Suliman, T.; Shaw, M.L.; et al. Breakthrough Infections with SARS-CoV-2 Omicron despite MRNA Vaccine Booster Dose. *The Lancet* **2022**, *399*, 625–626, doi:10.1016/S0140-6736(22)00090-3.

6.   Zhou, R.; To, K.K.-W.; Wong, Y.-C.; Liu, L.; Zhou, B.; Li, X.; Huang, H.; Mo, Y.; Luk, T.-Y.; Lau, T.T.-K.; et al. Acute SARS-CoV-2 Infection Impairs Dendritic Cell and T Cell Responses. *Immunity* **2020**, *53*, 864-877.e5, doi:10.1016/j.immuni.2020.07.026.

7.   Grant, R.A.; Morales-Nebreda, L.; Markov, N.S.; Swaminathan, S.; Querrey, M.; Guzman, E.R.; Abbott, D.A.; Donnelly, H.K.; Donayre, A.; Goldberg, I.A.; et al. Circuits between Infected Macrophages and T Cells in SARS-CoV-2 Pneumonia. *Nature* **2021**, *590*, 635–641, doi:10.1038/s41586-020-03148-w.

8.   Pontelli, M.C.; Castro, I.A.; Martins, R.B.; Veras, F.P.; Serra, L.L.; Nascimento, D.C.; Cardoso, R.S.; Rosales, R.; Lima, T.M.; Souza, J.P.; et al. Infection of Human Lymphomononuclear Cells by SARS-CoV-2. *bioRxiv* **2020**, 2020.07.28.225912, doi:10.1101/2020.07.28.225912.

9.   Salina, A.C.G.; dos Santos, D.; Rodrigues, T.S.; Fortes-Rocha, M.; Freitas-Filho, E.G.; Alzamora-Terrel, D.L.; Castro, I.M.; Fraga-Silva, T.F.; de Lima, M.H.; Nascimento, D.C.; et al. Efferocytosis of SARS-CoV-2-Infected Dying Cells Impairs Macrophage Anti-Inflammatory Functions and Clearance of Apoptotic Cells. *eLife* **2022**, *11*, e74443, doi:10.7554/eLife.74443.

10.   Junqueira, C.; Crespo, Â.; Ranjbar, S.; de Lacerda, L.B.; Lewandrowski, M.; Ingber, J.; Parry, B.; Ravid, S.; Clark, S.; Schrimpf, M.R.; et al. FcγR-Mediated SARS-CoV-2 Infection of Monocytes

Activates Inflammation. *Nature* **2022**, *606*, 576–584, doi:10.1038/s41586-022-04702-4.

11. Wendisch, D.; Dietrich, O.; Mari, T.; von Stillfried, S.; Ibarra, I.L.; Mittermaier, M.; Mache, C.; Chua, R.L.; Knoll, R.; Timm, S.; et al. SARS-CoV-2 Infection Triggers Profibrotic Macrophage Responses and Lung Fibrosis. *Cell* **2021**, *184*, 6243-6261.e27, doi:10.1016/j.cell.2021.11.033.

12. Sefik, E.; Qu, R.; Junqueira, C.; Kaffe, E.; Mirza, H.; Zhao, J.; Brewer, J.R.; Han, A.; Steach, H.R.; Israelow, B.; et al. Inflammasome Activation in Infected Macrophages Drives COVID-19 Pathology. *Nature* **2022**, *606*, 585–593, doi:10.1038/s41586-022-04802-1.

13. Huang, C.; Wang, Y.; Li, X.; Ren, L.; Zhao, J.; Hu, Y.; Zhang, L.; Fan, G.; Xu, J.; Gu, X.; et al. Clinical Features of Patients Infected with 2019 Novel Coronavirus in Wuhan, China. *The Lancet* **2020**, *395*, 497–506, doi:10.1016/S0140-6736(20)30183-5.

14. Del Valle, D.M.; Kim-Schulze, S.; Huang, H.-H.; Beckmann, N.D.; Nirenberg, S.; Wang, B.; Lavin, Y.; Swartz, T.H.; Madduri, D.; Stock, A.; et al. An Inflammatory Cytokine Signature Predicts COVID-19 Severity and Survival. *Nature Medicine* **2020**, *26*, 1636–1643, doi:10.1038/s41591-020-1051-9.

15. Song, P.; Li, W.; Xie, J.; Hou, Y.; You, C. Cytokine Storm Induced by SARS-CoV-2. *Clinica Chimica Acta* **2020**, *509*, 280–287, doi:10.1016/j.cca.2020.06.017.

16. Lassen, M.G.; Lukens, J.R.; Dolina, J.S.; Brown, M.G.; Hahn, Y.S. Intrahepatic IL-10 Maintains NKG2A^{+}Ly49^{-} Liver NK Cells in a Functionally Hyporesponsive State. *J. Immunol.* **2010**, *184*, 2693, doi:10.4049/jimmunol.0901362.

17. Mazzoni, A.; Salvati, L.; Maggi, L.; Capone, M.; Vanni, A.; Spinicci, M.; Mencarini, J.; Caporale, R.; Peruzzi, B.; Antonelli, A.; et al. Impaired Immune Cell Cytotoxicity in Severe COVID-19 Is IL-6 Dependent. *J Clin Invest* **2020**, *130*, 4694–4703, doi:10.1172/JCI138554.

18. Miorin Lisa; Kehrer Thomas; Sanchez-Aparicio Maria Teresa; Zhang Ke; Cohen Phillip; Patel Roosheel S.; Cupic Anastasija; Makio Tadashi; Mei Menghan; Moreno Elena; et al. SARS-CoV-2 Orf6 Hijacks Nup98 to Block STAT Nuclear Import and Antagonize Interferon Signaling. *Proceedings of the National Academy of Sciences* **2020**, *117*, 28344–28354, doi:10.1073/pnas.2016650117.

19. Sui, L.; Zhao, Y.; Wang, W.; Wu, P.; Wang, Z.; Yu, Y.; Hou, Z.; Tan, G.; Liu, Q. SARS-CoV-2 Membrane Protein Inhibits Type I Interferon Production Through Ubiquitin-Mediated Degradation of TBK1. *Front Immunol* **2021**, *12*, 662989, doi:10.3389/fimmu.2021.662989.

20. Sui, C.; Xiao, T.; Zhang, S.; Zeng, H.; Zheng, Y.; Liu, B.; Xu, G.; Gao, C.; Zhang, Z. SARS-CoV-2 NSP13 Inhibits Type I IFN Production by Degradation of TBK1 via P62-Dependent Selective Autophagy. *J Immunol* **2022**, *208*, 753–761, doi:10.4049/jimmunol.2100684.

21. Chang, T.; Yang, J.; Deng, H.; Chen, D.; Yang, X.; Tang, Z.-H. Depletion and Dysfunction of Dendritic Cells: Understanding SARS-CoV-2 Infection. *Frontiers in Immunology* **2022**, *13*.

22. Zhang Yiwen; Chen Yingshi; Li Yuzhuang; Huang Feng; Luo Baohong; Yuan Yaochang; Xia Baijin; Ma Xiancai; Yang Tao; Yu Fei; et al. The ORF8 Protein of SARS-CoV-2 Mediates Immune Evasion through down-Regulating MHC-I. *Proceedings of the National Academy of Sciences* **2021**, *118*, e2024202118, doi:10.1073/pnas.2024202118.

23. Yoo, J.-S.; Sasaki, M.; Cho, S.X.; Kasuga, Y.; Zhu, B.; Ouda, R.; Orba, Y.; de Figueiredo, P.; Sawa, H.; Kobayashi, K.S. SARS-CoV-2 Inhibits Induction of the MHC Class I Pathway by Targeting the STAT1-IRF1-NLRC5 Axis. *Nat Commun* **2021**, *12*, 6602, doi:10.1038/s41467-021-26910-8.

24. Saichi, M.; Ladjemi, M.Z.; Korniotis, S.; Rousseau, C.; Ait Hamou, Z.; Massenet-Regad, L.; Amblard, E.; Noel, F.; Marie, Y.; Bouteiller, D.; et al. Single-Cell RNA Sequencing of Blood Antigen-Presenting Cells in Severe COVID-19 Reveals Multi-Process Defects in Antiviral Immunity. *Nature Cell Biology* **2021**, *23*, 538–551, doi:10.1038/s41556-021-00681-2.

25. Wilk, A.J.; Rustagi, A.; Zhao, N.Q.; Roque, J.; Martínez-Colón, G.J.; McKechnie, J.L.; Ivison, G.T.; Ranganath, T.; Vergara, R.; Hollis, T.; et al. A Single-Cell Atlas of the Peripheral Immune Response in Patients with Severe COVID-19. *Nat Med* **2020**, *26*, 1070–1076, doi:10.1038/s41591-020-0944-y.

26. Giamarellos-Bourboulis, E.J.; Netea, M.G.; Rovina, N.; Akinosoglou, K.; Antoniadou, A.; Antonakos, N.; Damoraki, G.; Gkavogianni, T.; Adami, M.-E.; Katsaounou, P.; et al. Complex Immune Dysregulation in COVID-19 Patients with Severe Respiratory Failure. *Cell Host & Microbe* **2020**, *27*, 992-1000.e3, doi:10.1016/j.chom.2020.04.009.

27. Munnur, D.; Teo, Q.; Eggermont, D.; Lee, H.H.Y.; Thery, F.; Ho, J.; van Leur, S.W.; Ng, W.W.S.; Siu, L.Y.L.; Beling, A.; et al. Altered ISGylation Drives Aberrant Macrophage-Dependent Immune Responses during SARS-CoV-2 Infection. *Nat Immunol* **2021**, *22*, 1416–1427, doi:10.1038/s41590-021-01035-8.

28. Munster, V.J.; Feldmann, F.; Williamson, B.N.; van Doremalen, N.; Pérez-Pérez, L.; Schulz, J.; Meade-White, K.; Okumura, A.; Callison, J.; Brumbaugh, B.; et al. Respiratory Disease in Rhesus Macaques Inoculated with SARS-CoV-2. *Nature* **2020**, *585*, 268–272, doi:10.1038/s41586-020-2324-7.

29. Ziegler, C.G.K.; Allon, S.J.; Nyquist, S.K.; Mbano, I.M.; Miao, V.N.; Tzouanas, C.N.; Cao, Y.; Yousif, A.S.; Bals, J.; Hauser, B.M.; et al. SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues. *Cell* **2020**, *181*, 1016-1035.e19, doi:10.1016/j.cell.2020.04.035.

30. Steimle Viktor; Siegrist Claire-Anne; Mottet Annick; Lisowska-Grospierre Barbara; Mach Bernard Regulation of MHC Class II Expression by Interferon-γ Mediated by the Transactivator Gene CIITA. *Science* **1994**, *265*, 106–109, doi:10.1126/science.8016643.

31. Hu, X.; Ivashkiv, L.B. Cross-Regulation of Signaling Pathways by Interferon-Gamma: Implications for Immune Responses and Autoimmune Diseases. *Immunity* **2009**, *31*, 539–550, doi:10.1016/j.immuni.2009.09.002.

32. Muhlethaler-Mottet, A.; Di Berardino, W.; Otten, L.A.; Mach, B. Activation of the MHC Class II Transactivator CIITA by Interferon-Gamma Requires Cooperative Interaction between Stat1 and USF-1. *Immunity* **1998**, *8*, 157–166, doi:10.1016/s1074-7613(00)80468-9.

33. Zhang, G.; Zhang, Z.; Liu, Z. Interferon Regulation Factor-3 Is a Critical Regulator of the Mature of Dendritic Cells from Mice. *Scandinavian Journal of Immunology* **2013**, *77*, 13–20, doi:10.1111/sji.12005.

34. Beresford, G.W.; Boss, J.M. CIITA Coordinates Multiple Histone Acetylation Modifications at the HLA-DRA Promoter. *Nat Immunol* **2001**, *2*, 652–657, doi:10.1038/89810.

35. Masternak, K.; Muhlethaler-Mottet, A.; Villard, J.; Zufferey, M.; Steimle, V.; Reith, W. CIITA Is a Transcriptional Coactivator That Is Recruited to MHC Class II Promoters by Multiple Synergistic Interactions with an Enhanceosome Complex. *Genes Dev.* **2000**, *14*, 1156–1166, doi:10.1101/gad.14.9.1156.

36. Landmann, S.; Mühlethaler-Mottet, A.; Bernasconi, L.; Suter, T.; Waldburger, J.-M.; Masternak, K.; Arrighi, J.-F.; Hauser, C.; Fontana, A.; Reith, W. Maturation of Dendritic Cells Is Accompanied by Rapid Transcriptional Silencing of Class II Transactivator (Ciita) Expression. *Journal of Experimental Medicine* **2001**, *194*, 379–392, doi:10.1084/jem.194.4.379.

37. Kim, D.; Kim, S.; Park, J.; Chang, H.R.; Chang, J.; Ahn, J.; Park, H.; Park, J.; Son, N.; Kang, G.; et al. A High-Resolution Temporal Atlas of the SARS-CoV-2 Translatome and Transcriptome. *Nat Commun* **2021**, *12*, 5120, doi:10.1038/s41467-021-25361-5.

38. Zhou, H.; Su, H.S.; Zhang, X.; Douhan, J.; Glimcher, L.H. CIITA-Dependent and -Independent Class II MHC Expression Revealed by a Dominant Negative Mutant. *J. Immunol.* **1997**, *158*, 4741.

39. Williams, G.S.; Malin, M.; Vremec, D.; Chang, C.H.; Boyd, R.; Benoist, C.; Mathis, D. Mice Lacking the Transcription Factor CIITA--a Second Look. *International Immunology* **1998**, *10*, 1957–1967, doi:10.1093/intimm/10.12.1957.

40. Lee, K.-W.; Lee, Y.; Kim, D.-S.; Kwon, H.-J. Direct Role of NF-KB Activation in Toll-like Receptor-Triggered HLA-DRA Expression. *European Journal of Immunology* **2006**, *36*, 1254–1266, doi:10.1002/eji.200535577.

41. Gialitakis, M.; Kretsovali, A.; Spilianakis, C.; Kravariti, L.; Mages, J.; Hoffmann, R.; Hatzopoulos, A.K.; Papamatheakis, J. Coordinated Changes of Histone Modifications and HDAC Mobilization Regulate the Induction of MHC Class II Genes by Trichostatin A. *Nucleic Acids Research* **2006**, *34*, 765–772, doi:10.1093/nar/gkj462.

42. Gordon, D.E.; Jang, G.M.; Bouhaddou, M.; Xu, J.; Obernier, K.; White, K.M.; O'Meara, M.J.; Rezelj, V.V.; Guo, J.Z.; Swaney, D.L.; et al. A SARS-CoV-2 Protein Interaction Map Reveals Targets for Drug Repurposing. *Nature* **2020**, doi:10.1038/s41586-020-2286-9.

43. Jin, Z.; Du, X.; Xu, Y.; Deng, Y.; Liu, M.; Zhao, Y.; Zhang, B.; Li, X.; Zhang, L.; Peng, C.; et al. Structure of Mpro from SARS-CoV-2 and Discovery of Its Inhibitors. *Nature* **2020**, *582*, 289–293, doi:10.1038/s41586-020-2223-y.

44. Salgado-Albarrán, M.; Navarro-Delgado, E.I.; Del Moral-Morales, A.; Alcaraz, N.; Baumbach, J.; González-Barrios, R.; Soto-Reyes, E. Comparative Transcriptome Analysis Reveals Key Epigenetic Targets in SARS-CoV-2 Infection. *npj Syst Biol Appl* **2021**, *7*, 21, doi:10.1038/s41540-021-00181-x.

45. Chlamydas, S.; Papavassiliou, A.G.; Piperi, C. Epigenetic Mechanisms Regulating COVID-19 Infection. *Epigenetics* **2021**, *16*, 263–270, doi:10.1080/15592294.2020.1796896.

46. Josset, L.; Menachery, V.D.; Gralinski, L.E.; Agnihothram, S.; Sova, P.; Carter, V.S.; Yount, B.L.; Graham, R.L.; Baric, R.S.; Katze, M.G. Cell Host Response to Infection with Novel Human Coronavirus EMC Predicts Potential Antivirals and Important Differences with SARS Coronavirus. *mBio* **2013**, *4*, e00165, doi:10.1128/mBio.00165-13.

47. Yashvardhini, N.; Kumar, A.; Jha, D.K. Analysis of SARS-CoV-2 mutations in the main viral protease (NSP5) and its implications on the vaccine designing strategies. *Vacunas* **2021**, 10.1016/j.vacun.2021.10.002, doi:10.1016/j.vacun.2021.10.002.

48. Zhang Linlin; Lin Daizong; Sun Xinyuanyuan; Curth Ute; Drosten Christian; Sauerhering Lucie; Becker Stephan; Rox Katharina; Hilgenfeld Rolf Crystal Structure of SARS-CoV-2 Main Protease Provides a Basis for Design of Improved α-Ketoamide Inhibitors. *Science* **2020**, *368*, 409–412, doi:10.1126/science.abb3405.

49. Hegyi, A.; Ziebuhr, J. 2002 Conservation of Substrate Specificities among Coronavirus Main Proteases. *Journal of General Virology 83*, 595–599, doi:10.1099/0022-1317-83-3-595.

50. Lee, J.; Kenward, C.; Worrall, L.J.; Vuckovic, M.; Gentile, F.; Ton, A.-T.; Ng, M.; Cherkasov, A.; Strynadka, N.C.J.; Paetzel, M. X-Ray Crystallographic Characterization of the SARS-CoV-2 Main Protease Polyprotein Cleavage Sites Essential for Viral Processing and Maturation. *Nat Commun* **2022**, *13*, 5196, doi:10.1038/s41467-022-32854-4.

51. Stumptner-Cuvelette, P.; Morchoisne, S.; Dugast, M.; Le Gall, S.; Raposo, G.; Schwartz, O.; Benaroch, P. HIV-1 Nef Impairs MHC Class II Antigen Presentation and Surface Expression. *Proc Natl Acad Sci U S A* **2001**, *98*, 12144–12149, doi:10.1073/pnas.221256498.

52. Santerre, M.; Arjona, S.P.; Allen, C.N.; Shcherbik, N.; Sawaya, B.E. Why Do SARS-CoV-2 NSPs Rush to the ER? *J Neurol* **2021**, *268*, 2013–2022, doi:10.1007/s00415-020-10197-8.

53. Xia, Y.P.; Yeh, C.T.; Ou, J.H.; Lai, M.M. Characterization of Nuclear Targeting Signal of Hepatitis Delta Antigen: Nuclear Transport as a Protein Complex. *Journal of Virology* **1992**, *66*, 914–921, doi:10.1128/jvi.66.2.914-921.1992.

54. Morris, A.C.; Beresford, G.W.; Mooney, M.R.; Boss, J.M. Kinetics of a Gamma Interferon Response: Expression and Assembly of CIITA Promoter IV and Inhibition by Methylation. *Mol Cell Biol* **2002**, *22*, 4781–4791, doi:10.1128/MCB.22.13.4781-4791.2002.

55. Zinzow-Kramer, W.M.; Long, A.B.; Youngblood, B.A.; Rosenthal, K.M.; Butler, R.; Mohammed, A.-U.-R.; Skountzou, I.; Ahmed, R.; Evavold, B.D.; Boss, J.M. CIITA Promoter I CARD-Deficient Mice Express Functional MHC Class II Genes in Myeloid and Lymphoid Compartments. *Genes Immun* **2012**, *13*, 299–310, doi:10.1038/gene.2011.86.

56. Krawczyk, M.; Peyraud, N.; Rybtsova, N.; Masternak, K.; Bucher, P.; Barras, E.; Reith, W. Long Distance Control of MHC Class II Expression by Multiple Distal Enhancers Regulated by Regulatory Factor X Complex and CIITA. *The Journal of Immunology* **2004**, *173*, 6200–6210, doi:10.4049/jimmunol.173.10.6200.

57. Soe, K.C.; Devaiah, B.N.; Singer, D.S. Transcriptional Coactivator CIITA, a Functional Homolog of TAF1, Has Kinase Activity. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* **2013**, *1829*, 1184–1190, doi:10.1016/j.bbagrm.2013.09.001.

58. Raval, A.; Howcroft, T.K.; Weissman, J.D.; Kirshner, S.; Zhu, X.-S.; Yokoyama, K.; Ting, J.; Singer, D.S. Transcriptional Coactivator, CIITA, Is an Acetyltransferase That Bypasses a Promoter Requirement for TAFII250. *Molecular Cell* **2001**, *7*, 105–115, doi:10.1016/S1097-2765(01)00159-9.

59. Mantegazza, A.R.; Guttentag, S.H.; El-Benna, J.; Sasai, M.; Iwasaki, A.; Shen, H.; Laufer, T.M.; Marks, M.S. Adaptor Protein-3 in Dendritic Cells Facilitates Phagosomal Toll-like Receptor Signaling and Antigen Presentation to CD4(+) T Cells. *Immunity* **2012**, *36*, 782–794, doi:10.1016/j.immuni.2012.02.018.

60.   Mantegazza, A.R.; Zajac, A.L.; Twelvetrees, A.; Holzbaur, E.L.F.; Amigorena, S.; Marks, M.S. TLR-Dependent Phagosome Tubulation in Dendritic Cells Promotes Phagosome Cross-Talk to Optimize MHC-II Antigen Presentation. *Proceedings of the National Academy of Sciences of the United States of America* **2014**, *111*, 15508–15513, doi:10.1073/pnas.1412998111.

61.   Kong, X.; Fang, M.; Li, P.; Fang, F.; Xu, Y. HDAC2 Deacetylates Class II Transactivator and Suppresses Its Activity in Macrophages and Smooth Muscle Cells. *J. Mol. Cell. Cardiol.* **2009**, *46*, 292–299, doi:10.1016/j.yjmcc.2008.10.023.

62.   Lee, J.; Worrall, L.J.; Vuckovic, M.; Rosell, F.I.; Gentile, F.; Ton, A.-T.; Caveney, N.A.; Ban, F.; Cherkasov, A.; Paetzel, M.; et al. Crystallographic Structure of Wild-Type SARS-CoV-2 Main Protease Acyl-Enzyme Intermediate with Physiological C-Terminal Autoprocessing Site. *Nat Commun* **2020**, *11*, 5877, doi:10.1038/s41467-020-19662-4.

63.   Zhao, Y.; Zhu, Y.; Liu, X.; Jin, Z.; Duan, Y.; Zhang, Q.; Wu, C.; Feng, L.; Du, X.; Zhao, J.; et al. Structural Basis for Replicase Polyprotein Cleavage and Substrate Specificity of Main Protease from SARS-CoV-2. *Proceedings of the National Academy of Sciences* **2022**, *119*, e2117142119, doi:10.1073/pnas.2117142119.

64.   Spilianakis, C.; Papamatheakis, J.; Kretsovali, A. Acetylation by PCAF Enhances CIITA Nuclear Accumulation and Transactivation of Major Histocompatibility Complex Class II Genes. *Mol Cell Biol* **2000**, *20*, 8489–8498, doi:10.1128/MCB.20.22.8489-8498.2000.

65.   Zhou, H.; Ji, J.; Chen, X.; Bi, Y.; Li, J.; Wang, Q.; Hu, T.; Song, H.; Zhao, R.; Chen, Y.; et al. Identification of Novel Bat Coronaviruses Sheds Light on the Evolutionary Origins of SARS-CoV-2 and Related Viruses. *Cell* **2021**, *184*, 4380-4391.e14, doi:10.1016/j.cell.2021.06.008.

66.   Delaune, D.; Hul, V.; Karlsson, E.A.; Hassanin, A.; Ou, T.P.; Baidaliuk, A.; Gámbaro, F.; Prot, M.; Tu, V.T.; Chea, S.; et al. A Novel SARS-CoV-2 Related Coronavirus in Bats from Cambodia. *Nat Commun* **2021**, *12*, 6563, doi:10.1038/s41467-021-26809-4.

67.   Paroni, G.; Mizzau, M.; Henderson, C.; Del Sal, G.; Schneider, C.; Brancolini, C. Caspase-Dependent Regulation of Histone Deacetylase 4 Nuclear-Cytoplasmic Shuttling Promotes Apoptosis. *MBoC* **2004**, *15*, 2804–2818, doi:10.1091/mbc.e03-08-0624.

68.   Naik, N.G.; Lee, S.-C.; Veronese, B.H.S.; Ma, Z.; Toth, Z. Interaction of HDAC2 with SARS-CoV-2 NSP5 and IRF3 Is Not Required for NSP5-Mediated Inhibition of Type I Interferon Signaling Pathway. *Microbiol Spectr* 10, e02322-22, doi:10.1128/spectrum.02322-22.

69.   Zuo, J.; Hislop, A.D.; Leung, C.S.; Sabbah, S.; Rowe, M. Kaposi's Sarcoma-Associated Herpesvirus-Encoded Viral IRF3 Modulates Major Histocompatibility Complex Class II (MHC-II) Antigen Presentation through MHC-II Transactivator-Dependent and -Independent Mechanisms: Implications for Oncogenesis. *J Virol* **2013**, *87*, 5340–5350, doi:10.1128/JVI.00250-13.

70.   ENCODE Project Consortium An Integrated Encyclopedia of DNA Elements in the Human Genome. *Nature* **2012**, *489*, 57–74, doi:10.1038/nature11247.

71.   Seow, J.; Graham, C.; Merrick, B.; Acors, S.; Pickering, S.; Steel, K.J.A.; Hemmings, O.; O'Byrne, A.; Kouphou, N.; Galao, R.P.; et al. Longitudinal Observation and Decline of Neutralizing Antibody Responses in the Three Months Following SARS-CoV-2 Infection in Humans. *Nat Microbiol* **2020**, *5*, 1598–1607, doi:10.1038/s41564-020-00813-8.

72.   Ibarrondo, F.J.; Fulcher, J.A.; Goodman-Meza, D.; Elliott, J.; Hofmann, C.; Hausner, M.A.; Ferbas, K.G.; Tobin, N.H.; Aldrovandi, G.M.; Yang, O.O. Rapid Decay of Anti-SARS-CoV-2 Antibodies in Persons with Mild Covid-19. *N Engl J Med* **2020**, *383*, 1085–1087, doi:10.1056/NEJMc2025179.

73.   Kaneko, N.; Kuo, H.-H.; Boucau, J.; Farmer, J.R.; Allard-Chamard, H.; Mahajan, V.S.; Piechocka-Trocha, A.; Lefteri, K.; Osborn, M.; Bals, J.; et al. Loss of Bcl-6-Expressing T Follicular Helper Cells and Germinal Centers in COVID-19. *Cell* **2020**, *183*, 143-157.e13, doi:10.1016/j.cell.2020.08.025.

74.   Wiech, M.; Chroscicki, P.; Swatler, J.; Stepnik, D.; De Biasi, S.; Hampel, M.; Brewinska-Olchowik, M.; Maliszewska, A.; Sklinda, K.; Durlik, M.; et al. Remodeling of T Cell Dynamics During Long COVID Is Dependent on Severity of SARS-CoV-2 Infection. *Frontiers in Immunology* **2022**, *13*.

75.   van de Weijer, M.L.; Luteijn, R.D.; Wiertz, E.J.H.J. Viral Immune Evasion: Lessons in MHC Class I Antigen Presentation. *Semin Immunol* **2015**, *27*, 125–137, doi:10.1016/j.smim.2015.03.010.

76.   Dikeakos, J.D.; Thomas, L.; Kwon, G.; Elferich, J.; Shinde, U.; Thomas, G. An Interdomain Binding Site on HIV-1 Nef Interacts with PACS-1 and PACS-2 on Endosomes to down-Regulate MHC-I. *Molecular biology of the cell* **2012**, *23*, 2184–2197, doi:10.1091/mbc.E11-11-0928.

77.   Dirk, B.S.; Pawlak, E.N.; Johnson, A.L.; Van Nynatten, L.R.; Jacob, R.A.; Heit, B.; Dikeakos, J.D. HIV-1 Nef Sequesters MHC-I Intracellularly by Targeting Early Stages of Endocytosis and Recycling. *Scientific reports* **2016**, *6*, 37021, doi:10.1038/srep37021.

78.   Koutsakos, M.; McWilliam, H.E.G.; Aktepe, T.E.; Fritzlar, S.; Illing, P.T.; Mifsud, N.A.; Purcell, A.W.; Rockman, S.; Reading, P.C.; Vivian, J.P.; et al. Downregulation of MHC Class I Expression by Influenza A and B Viruses. *Frontiers in Immunology* **2019**, *10*.

79.   Temme, S.; Eis-Hübinger, A.M.; McLellan, A.D.; Koch, N. The Herpes Simplex Virus-1 Encoded Glycoprotein B Diverts HLA-DR into the Exosome Pathway. *The Journal of Immunology* **2010**, *184*, 236–243, doi:10.4049/jimmunol.0902192.

80.   Trgovcich, J.; Johnson, D.; Roizman, B. Cell Surface Major Histocompatibility Complex Class II Proteins Are Regulated by the Products of the Gamma(1)34.5 and U(L)41 Genes of Herpes Simplex Virus 1. *J Virol* **2002**, *76*, 6974–6986, doi:10.1128/jvi.76.14.6974-6986.2002.

81.   Hegde, N.R.; Tomazin, R.A.; Wisner, T.W.; Dunn, C.; Boname, J.M.; Lewinsohn, D.M.; Johnson, D.C. Inhibition of HLA-DR Assembly, Transport, and Loading by Human Cytomegalovirus Glycoprotein US3: A Novel Mechanism for Evading Major Histocompatibility Complex Class II Antigen Presentation. *J Virol* **2002**, *76*, 10929–10941, doi:10.1128/jvi.76.21.10929-10941.2002.

82.   Tomazin, R.; Boname, J.; Hegde, N.R.; Lewinsohn, D.M.; Altschuler, Y.; Jones, T.R.; Cresswell, P.; Nelson, J.A.; Riddell, S.R.; Johnson, D.C. Cytomegalovirus US2 Destroys Two Components of the MHC Class II Pathway, Preventing Recognition by CD4+ T Cells. *Nat Med* **1999**, *5*, 1039–1043, doi:10.1038/12478.

83.   Lin, J.-H.; Lin, J.-Y.; Chou, Y.-C.; Chen, M.-R.; Yeh, T.-H.; Lin, C.-W.; Lin, S.-J.; Tsai, C.-H. Epstein-Barr Virus LMP2A Suppresses MHC Class II Expression by Regulating the B-Cell Transcription Factors E47 and PU.1. *Blood* **2015**, *125*, 2228–2238, doi:10.1182/blood-2014-08-594689.

84.   van der Stoep, N.; Quinten, E.; Rezende, M.M.; van den Elsen, P.J. E47, IRF-4, and PU.1 Synergize to Induce B-Cell-Specific

Activation of the Class II Transactivator Promoter III (CIITA-PIII). *Blood* **2004**, *104*, 2849–2857, doi:10.1182/blood-2004-03-0790.

85. Zhu, X.; Fang, L.; Wang, D.; Yang, Y.; Chen, J.; Ye, X.; Foda, M.F.; Xiao, S. Porcine Deltacoronavirus Nsp5 Inhibits Interferon-β Production through the Cleavage of NEMO. *Virology* **2017**, *502*, 33–38, doi:10.1016/j.virol.2016.12.005.

86. Zhu, X.; Wang, D.; Zhou, J.; Pan, T.; Chen, J.; Yang, Y.; Lv, M.; Ye, X.; Peng, G.; Fang, L.; et al. Porcine Deltacoronavirus Nsp5 Antagonizes Type I Interferon Signaling by Cleaving STAT2. *J Virol* **2017**, *91*, e00003-17, doi:10.1128/JVI.00003-17.

87. Zhu, X.; Chen, J.; Tian, L.; Zhou, Y.; Xu, S.; Long, S.; Wang, D.; Fang, L.; Xiao, S. Porcine Deltacoronavirus Nsp5 Cleaves DCP1A To Decrease Its Antiviral Activity. *J Virol* **2020**, *94*, e02162-19, doi:10.1128/JVI.02162-19.

88. Li, W.; Qiao, J.; You, Q.; Zong, S.; Peng, Q.; Liu, Y.; Hu, S.; Liu, W.; Li, S.; Shu, X.; et al. SARS-CoV-2 Nsp5 Activates NF-KB Pathway by Upregulating SUMOylation of MAVS. *Frontiers in Immunology* **2021**, *12*.

89. Zheng, Y.; Deng, J.; Han, L.; Zhuang, M.-W.; Xu, Y.; Zhang, J.; Nan, M.-L.; Xiao, Y.; Zhan, P.; Liu, X.; et al. SARS-CoV-2 NSP5 and N Protein Counteract the RIG-I Signaling Pathway by Suppressing the Formation of Stress Granules. *Signal Transduct Target Ther* **2022**, *7*, 22, doi:10.1038/s41392-022-00878-3.

90. Menachery, V.D.; Eisfeld, A.J.; Schäfer, A.; Josset, L.; Sims, A.C.; Proll, S.; Fan, S.; Li, C.; Neumann, G.; Tilton, S.C.; et al. Pathogenic Influenza Viruses and Coronaviruses Utilize Similar and Contrasting Approaches To Control Interferon-Stimulated Gene Responses. *mBio* **2014**, *5*, e01174-14, doi:10.1128/mBio.01174-14.

91. Khokhar, M.; Tomo, S.; Purohit, P. MicroRNAs Based Regulation of Cytokine Regulating Immune Expressed Genes and Their Transcription Factors in COVID-19. *Meta Gene* **2022**, *31*, 100990, doi:10.1016/j.mgene.2021.100990.

92. Balnis, J.; Madrid, A.; Hogan, K.J.; Drake, L.A.; Chieng, H.C.; Tiwari, A.; Vincent, C.E.; Chopra, A.; Vincent, P.A.; Robek, M.D.; et al. Blood DNA Methylation and COVID-19 Outcomes. *Clinical Epigenetics* **2021**, *13*, 118, doi:10.1186/s13148-021-01102-9.

93. Moura, M.C. de; Davalos, V.; Planas-Serra, L.; Alvarez-Errico, D.; Arribas, C.; Ruiz, M.; Aguilera-Albesa, S.; Troya, J.; Valencia-Ramos, J.; Vélez-Santamaria, V.; et al. Epigenome-Wide Association Study of COVID-19 Severity with Respiratory Failure. *eBioMedicine* **2021**, *66*, doi:10.1016/j.ebiom.2021.103339.

94. Bensussen, A.; Álvarez-Buylla, E.R.; Díaz, J. SARS-CoV-2 Nsp5 Protein Causes Acute Lung Inflammation, A Dynamical Mathematical Model. *Frontiers in Systems Biology* **2021**, *1*.

95. Shweta, S.; Krishna K, S. Valproic Acid in Prevention and Treatment of COVID-19. *Int J Respir Pulm Med* **2020**, *7*, 1–7, doi:10.23937/2378-3516/1410138.

96. Teodori, L.; Sestili, P.; Madiai, V.; Coppari, S.; Fraternale, D.; Rocchi, M.B.L.; Ramakrishna, S.; Albertini, M.C. MicroRNAs Bioinformatics Analyses Identifying HDAC Pathway as a Putative Target for Existing Anti-COVID-19 Therapeutics. *Frontiers in Pharmacology* **2020**, *11*.

97. Liu, K.; Zou, R.; Cui, W.; Li, M.; Wang, X.; Dong, J.; Li, H.; Li, H.; Wang, P.; Shao, X.; et al. Clinical HDAC Inhibitors Are Effective Drugs to Prevent the Entry of SARS-CoV2. *ACS Pharmacol. Transl. Sci.* **2020**, *3*, 1361–1370, doi:10.1021/acsptsci.0c00163.

98. Sixto-López, Y.; Correa-Basurto, J. HDAC Inhibition as Neuroprotection in COVID-19 Infection. *Current Topics in Medicinal Chemistry* **2022**, *22*, 1–10, doi:10.2174/1568026622666220303113445.

99. Saiz, M.L.; DeDiego, M.L.; López-García, D.; Corte-Iglesias, V.; Baragaño Raneros, A.; Astola, I.; Asensi, V.; López-Larrea, C.; Suarez-Alvarez, B. Epigenetic Targeting of the ACE2 and NRP1 Viral Receptors Limits SARS-CoV-2 Infectivity. *Clinical Epigenetics* **2021**, *13*, 187, doi:10.1186/s13148-021-01168-5.

100. Temmam, S.; Vongphayloth, K.; Baquero, E.; Munier, S.; Bonomi, M.; Regnault, B.; Douangboubpha, B.; Karami, Y.; Chrétien, D.; Sanamxay, D.; et al. Bat Coronaviruses Related to SARS-CoV-2 and Infectious for Human Cells. *Nature* **2022**, *604*, 330–336, doi:10.1038/s41586-022-04532-4.

101. P, K.M.; Sivashanmugam, K.; Kandasamy, M.; Subbiah, R.; Ravikumar, V. Repurposing of Histone Deacetylase Inhibitors: A Promising Strategy to Combat Pulmonary Fibrosis Promoted by TGF-β Signalling in COVID-19 Survivors. *Life Sciences* **2021**, *266*, 118883, doi:10.1016/j.lfs.2020.118883.

102. Ripamonti, C.; Spadotto, V.; Pozzi, P.; Stevenazzi, A.; Vergani, B.; Marchini, M.; Sandrone, G.; Bonetti, E.; Mazzarella, L.; Minucci, S.; et al. HDAC Inhibition as Potential Therapeutic Strategy to Restore the Deregulated Immune Response in Severe COVID-19. *Frontiers in Immunology* **2022**, *13*.

103. Wu, A.; Peng, Y.; Huang, B.; Ding, X.; Wang, X.; Niu, P.; Meng, J.; Zhu, Z.; Zhang, Z.; Wang, J.; et al. Genome Composition and Divergence of the Novel Coronavirus (2019-NCoV) Originating in China. *Cell Host & Microbe* **2020**, *27*, 325–328, doi:10.1016/j.chom.2020.02.001.

104. Smith, A.L.; Friedman, D.B.; Yu, H.; Carnahan, R.H.; Reynolds, A.B. ReCLIP (Reversible Cross-Link Immuno-Precipitation): An Efficient Method for Interrogation of Labile Protein Complexes. *PloS one* **2011**, *6*, e16206, doi:10.1371/journal.pone.0016206.

105. Schindelin, J.; Arganda-Carreras, I.; Frise, E.; Kaynig, V.; Longair, M.; Pietzsch, T.; Preibisch, S.; Rueden, C.; Saalfeld, S.; Schmid, B.; et al. Fiji: An Open-Source Platform for Biological-Image Analysis. *Nature methods* **2012**, *9*, 676–682, doi:10.1038/nmeth.2019.

106. Schneider, C.A.; Rasband, W.S.; Eliceiri, K.W. NIH Image to ImageJ: 25 Years of Image Analysis. *Nat Methods* **2012**, *9*, 671–675, doi:10.1038/nmeth.2089.

107. Tepperman, A.; Zheng, D.J.; Taka, M.A.; Vrieze, A.; Le Lam, A.; Heit, B. Customizable Live-Cell Imaging Chambers for Multimodal and Multiplex Fluorescence Microscopy. *Biochemistry and cell biology = Biochimie et biologie cellulaire* **2020**, *98*, 612–623, doi:10.1139/bcb-2020-0064.

108. Bolte, S.; Cordelières, F.P. A Guided Tour into Subcellular Colocalization Analysis in Light Microscopy. *Journal of microscopy* **2006**, *224*, 213–232, doi:10.1111/j.1365-2818.2006.01706.x.

109. van Rheenen, J.; Langeslag, M.; Jalink, K. Correcting Confocal Acquisition to Optimize Imaging of Fluorescence Resonance Energy Transfer by Sensitized Emission. *Biophysical Journal* **2004**, *86*, 2517–2529, doi:10.1016/S0006-3495(04)74307-6.

110. Lambert, T.J. FPbase: A Community-Editable Fluorescent Protein Database. *Nature Methods* **2019**, *16*, 277–278, doi:10.1038/s41592-019-0352-8.

111.   Ye, C.J.; Lawrenson, L.; Liu, G.; Stevens, J.; Bremer, S.; Ye, K.J.; Heng, H.H.Q. Simultaneous Fluorescence Immunostaining and FISH. 24.

112.   Berg, S.; Kutra, D.; Kroeger, T.; Straehle, C.N.; Kausler, B.X.; Haubold, C.; Schiegg, M.; Ales, J.; Beier, T.; Rudy, M.; et al. Ilastik: Interactive Machine Learning for (Bio)Image Analysis. *Nat Methods* **2019**, *16*, 1226–1232, doi:10.1038/s41592-019-0582-9.

113.   Tamura, K.; Stecher, G.; Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution* **2021**, *38*, 3022–3027, doi:10.1093/molbev/msab120.

114.   Capra, J.A.; Singh, M. Predicting Functionally Important Residues from Sequence Conservation. *Bioinformatics* **2007**, *23*, 1875–1882, doi:10.1093/bioinformatics/btm270.